

6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

Commercial Plans

2020 Prior Authorization Criteria

Health First Commercial Plans, Inc. is doing business under the name of Health First Health Plans. Health First Health Plans does not discriminate on the basis of race, color, national origin, disability, age, sex, gender identity, sexual orientation, or health status in the administration of the plan, including enrollment and benefit determinations.

36194_MPINFO7673(10/19)

Drugs

acitretin

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy.

Required Medical Information

Documented diagnosis of severe psoriasis

Age Restriction 18 years of age or older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Will not be approved for the treatment of acne.

ACTIMMUNE (interferon gamma-1B)

Drugs ACTIMMUNE

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Hypersensitivity to interferon gamma, E. coli derived proteins, or any component of the formulation.

Required Medical Information

Documentation of diagnosis of chronic granulomatous disease or severe malignant osteoporosis.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

ADCIRCA (tadalafil (Pulmonary Hypertension))

Drugs

tadalafil (pulm. hypertension), tadalafil oral tablet 20 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Receiving nitrate therapy (includes intermittent use).

Required Medical Information

Age Restriction 18 years or older

Prescriber Restriction Cardiologist or Pulmonologist

Coverage Duration Plan year

-

Other Criteria Sildenafil citrate (generic Revatio indicated for Pulmonary Hypertension) must be tried first. **Drugs** adefovir

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Optimal treatment duration is unknown.

Drugs ADEMPAS

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Pregnancy. Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form. Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline).

Required Medical Information

Age Restriction 18 years of age or older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For renewal, medication was effective (i.e. improved 6 minute walk distance, oxygen saturation, etc.)

Drugs AFINITOR ORAL TABLET 10 MG, 2.5 MG, 5 MG, 7.5 MG, everolimus (antineoplastic)

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with Sutent or Nexavar (Afinitor only), or B) Progressive pancreatic, nonfunctional GI or lung neuroendocrine tumors (NET) that are unresectable, locally advanced or metastatic (Afinitor only), or C) Renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery (Afinitor only), or D) Advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin (Afinitor only), or E) Subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection (Afinitor or Afinitor Disperz only).

Age Restriction

18 years of age or older for RCC, pNET, and renal angiomyolipoma with TSC. 1 year of age or older for SEGA.

Prescriber Restriction

Coverage Duration Plan year

Drugs ALINIA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation indicating treatment needed for diarrhea caused by Giardia lamblia or Cryptosporidium parvum.

Age Restriction

Age 1 year or older (Suspension) Age 12 years or older (Tablets)

Prescriber Restriction

Coverage Duration 30 days

Drugs ALIQOPA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented diagnosis of relapsed follicular lymphoma (FL) who have received at least two prior systemic therapies.

Age Restriction

Prescriber Restriction Oncologist or Hematologist

Coverage Duration Through end of benefit year

Drugs

alosetron

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Initial therapy for Irritable Bowel Syndrome (IBS) in the male gender.

Required Medical Information

Initial Therapy for Irritable Bowel Syndrome (IBS): 1. Confirmed diagnosis of IBS with diarrhea predominant symptoms for at least 6 months Reauthorization for Irritable Bowel Syndrome (IBS): 1. Recurrence of diarrhea predominant IBS, AND 2. documentation of positive clinical response while on Lotronex.

Age Restriction

Patient must be at least 18 years of age or older

Prescriber Restriction

Prescriber must be specially trained gastrointestinal physician

Coverage Duration

IBS Initial Therapy: 12 weeks Reauthorization: 6 months

Other Criteria

Initial Therapy for Irritable Bowel Syndrome (IBS): 1. Failure to both: a. An antispasmodic (e.g. dicyclomine) AND b. An anti-diarrhea agent (e.g. loperamide, diphenoxylate-atropine)

AMPYRA (dalfampridine)

Drugs

dalfampridine

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

History of seizure. Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute). Patient currently using any other forms of 4-aminopyridine.

Required Medical Information

Diagnosis of multiple sclerosis AND patient is ambulatory AND patient has walking impairment.

Age Restriction

Prescriber Restriction Neurologist

Coverage Duration

Initial: 3 months. Renewal: Plan year

Other Criteria

For renewal, documentation that walking speed has improved from baseline must be provided.

Drugs APTIOM

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs APTIVUS, APTIVUS (WITH VITAMIN E)

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 2 years of age and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Do not use Aptivus in treatment-naïve patients.

Drugs ARCALYST

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Use in combination with other IL-1 inhibitors (e.g. Ilaris, Kineret) or tumor necrosis factor (TNF) inhibitors (e.g. Enbrel, Humira, Remicade, etc). Individual is receiving live vaccines. Exhibiting evidence of active or chronic infection(s), including tuberculosis, or a history of recurrent infections. Individual has not had a tuberculin skin test (TST) or Centers for Disease Control (CDC)-recommended equivalent to evaluate for latent tuberculosis prior to initiating treatmentwith rilonacept.

Required Medical Information

Documented diagnosis of cryopyrin-associated period syndromes (CAPS), including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS). Member's diagnosis of CAPS must be confirmed by either NRLP=3 gene mutation OR overproduction of interleukin-1.

Age Restriction

12 years of age and older

Prescriber Restriction

Prescribed by or in consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist, or other medical specialist.

Coverage Duration Plan year

Other Criteria Approve doses based on FDA labeling.

Armodafinil (NUVIGIL)

Drugs

armodafinil

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis and treatment history.

Age Restriction Patient must be at least 17 years or older

Prescriber Restriction

Idiopathic hypersomnia-- approve if the diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)

Coverage Duration Plan year

Drugs ATRIPLA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older

Prescriber Restriction

Coverage Duration Plan year

Drugs AUBAGIO

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Concurrent use of Aubagio with other disease-modifying agents used for multiple sclerosis (MS) [eg, Avonex, Rebif, Betaseron, Extavia, Copaxone, Tysabri, Tecfidera, or Gilenya].

Required Medical Information

Documented diagnosis of relapsing form of MS (RRMS, SPMS with relapses, or PRMS) and previous MS therapies tried.

Age Restriction

Prescriber Restriction Prescribed by or in consultation with a neurologist or MS specialist.

Coverage Duration

Plan year

Other Criteria

For use in a relapsing form of MS, approve if: 1) Patient is currently taking teriflunomide (Aubagio), OR 2) Patient has tried dimethyl fumarate (Tecfidera), interferon beta-1a intramuscular (Avonex) and glatiramer acetate (Copaxone).

Drugs AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Drugs AVONEX INTRAMUSCULAR PEN INJECTOR KIT, AVONEX INTRAMUSCULAR SYRINGE KIT

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

Required Medical Information

Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis

Age Restriction

Prescriber Restriction

Prescribed by or in consultation with a neurologist or MS specialist.

Coverage Duration Plan year

Drugs AYVAKIT

Covered Uses

Exclusion Criteria

Required Medical Information

documentation of unresectable or metastatic GIST harboring a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs BALVERSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented diagnosis of locally advanced or metastatic urothelial carcinoma AND member has susceptible FGFR3 or FGFR2 genetic alteration as detected by an FDA-approved companion diagnostic AND disease has progressed during or following at least one line of prior platinum-containing chemotherapy including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.

Age Restriction 8 years or older

Prescriber Restriction

Coverage Duration 3 Months

Drugs BANZEL ORAL SUSPENSION, BANZEL ORAL TABLET 200 MG, 400 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Banzel is not covered for members with the diagnosis of Familial Short QT syndrome

Required Medical Information

Documentation of diagnosis. Documentation of previous therapies and that the current medication regimen is inadequate to control disease.

Age Restriction Must be 1 years of age or older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Patient must be refractory to at least 2 of the following: Felbamate (Felbatol), Lamotrigine (Lamictal), Topiramate (Topamax), Valproic acid (Depakene), Divalproex sodium (Depakote)

Drugs BARACLUDE ORAL SOLUTION, entecavir

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs BAVENCIO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 Months

Drugs BELSOMRA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented trial and failure of two formulary alternatives AND documented medication review to rule out medication induced insomnia.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

BENLYSTA

Drugs BENLYSTA SUBCUTANEOUS AUTO-INJECTOR

Covered Uses All FDA-approved indications not otherwise exluded for part D

Exclusion Criteria

Required Medical Information Documentation from the medical record of diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Benznidazole

Drugs

benznidazole

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Patients who have used disulfiram within two weeks of initiation of benznidazole

Required Medical Information

Documentation of a consultation with an infectious disease specialist. Reviewer will verify patient claim history to confirm that patient has not used disulfiram within two weeks prior to benznidazole initiation

Age Restriction

Prescriber Restriction

Coverage Duration 60 days

Benzodiazepines

Drugs DIAZEPAM INTENSOL, diazepam oral concentrate

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs BETASERON SUBCUTANEOUS KIT

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

Required Medical Information

Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis.

Age Restriction

Prescriber Restriction

Prescribed by or in consultation with a neurologist or MS specialist.

Coverage Duration

Plan year

Other Criteria

Approve if: 1) Patient is currently taking Betaseron, OR 2) Patient has tried dimethyl fumarate (Tecfidera), interferon beta-1a intramuscular (Avonex) and glatiramer acetate (Copaxone)

Drugs BRAFTOVI ORAL CAPSULE 50 MG, 75 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDAapproved test and the medication will be used in combination with binimetinib.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs CAPLYTA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs

raloxifene, tamoxifen

Covered Uses

This criteria is a copay review process. The medications tamoxifen or raloxifene may be eligible for \$0 cost-share for women 35 years of age or older who: 1) do not have a history of breast cancer, and 2) are being prescribed tamoxifen or raloxifene for the purpose of primary prevention of invasive breast cancer because the member is deemed high risk, and 3) are post-menopausal, if prescribed raloxifene (this requirement does not apply to tamoxifen)

Exclusion Criteria

Women under 35 years of age, history of breast cancer

Required Medical Information

A 5-year predicted risk of breast cancer greater than or equal to 1.66%, as calculated by the Gail model.

Age Restriction 35 years and older

Prescriber Restriction

Coverage Duration 5 years

Drugs BROVANA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Must have documented failure, intolerance or contraindication to a long-acting beta agonist formulary product OR be unable to use a hand-actuated device.

Drugs BRUKINSA

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation of mantle cell lymphoma (MCL) in patients who have received at least one prior therapy.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 Months

Drugs BUPHENYL ORAL TABLET, sodium phenylbutyrate oral tablet

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis confirmed by enzymatic, biochemical or genetic testing. Buphenyl will be used for chronic management of urea cycle disorders (UCD).

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Buprenorphine Products

Drugs

buprenorphine HCl sublingual, buprenorphine-naloxone sublingual film 12-3 mg, 2-0.5 mg, 4-1 mg, 8-2 mg

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Being used for the treatment of pain OR patient is using short or long acting narcotics concurrently with Suboxone/Subutex.

Required Medical Information

The indicated diagnosis and medication usage must be supported by documentation from the patient's medical records.

Age Restriction

Must be 16 years of age or older.

Prescriber Restriction

Prescribing provider must have a DEA number starting with the letter X, AND physician must be listed on the Buprenorphine Physician Locator maintained by the Substance Abuse and Mental Health Services Administration (SAMSHA).

Coverage Duration 12 months

Drugs BUSULFEX

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs BYNFEZIA

Covered Uses

Exclusion Criteria

Required Medical Information

To be used to treat of the following •Treatment of severe diarrhea/flushing episodes associated with metastatic carcinoid tumors in adult patients•Treatment of profuse watery diarrhea associated with vasoactive intestinal peptide tumors (VIPomas) in adult patients•To reduce the amount of growth hormone and insulin like growth factor 1 in adult patients with acromegaly who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation, and bromocriptine mesylate at maximally tolerated doses

Age Restriction

Prescriber Restriction

Coverage Duration

Through end of benefit year

Drugs AIMOVIG AUTOINJECTOR, AJOVY AUTOINJECTOR, AJOVY SYRINGE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinical documentation of migraines and member has tried and failed two formulary alternatives for migraine prophylaxis with two different mechanism of action.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs CALQUENCE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented diagnosis of mantle cell lymphoma (MCL) who have received at least one prior therapy.

Age Restriction

Prescriber Restriction Oncologist or hematologist

Coverage Duration 3 Months

Drugs capecitabine

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction Oncologist

Coverage Duration Plan year

CAPRELSA (vandetanib)

Drugs CAPRELSA ORAL TABLET 100 MG, 300 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria Congenital long QT syndrome

Required Medical Information

Diagnosis of symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease.

Age Restriction 18 years or older

Prescriber Restriction Oncologist or endocrinologist

Coverage Duration 3 Months

CASPOFUNGIN

Drugs

caspofungin

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A)Empirical therapy for presumed fungal infections in febrile, neutropenic patients, or B)Treatment of candidemia and other Candida infections (intraabdominal abscesses, peritonitis and pleural space infections), or C)Treatment of esophageal candidiasis, or D)Treatment of invasive aspergillosis in patients who are refractory to or intolerant of other therapies (amphotericin B, itraconazole).Age Restrictions: 3 months of age or older

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Drugs CAYSTON

Covered Uses

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs

Age Restriction 7 years of age and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations)

Drugs CESAMET

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation that nabilone is being used for the treatment of the nausea and vomiting associated with cancer chemotherapy in patients who have a failed to respond adequately to conventional antiemetic treatments (e.g. Zofran and Emend).

Age Restriction

Older than 10 months

Prescriber Restriction Oncologist

Coverage Duration

Through the duration of chemotherapy

Other Criteria

A substantial proportion of any group of patients treated with nabilone can be expected to experience disturbing psychotomimetic reactions not observed with other antiemetic agents. Because of its potential to alter the mental state, nabilone is intended for use under circumstances that permit close supervision of the patient by a responsible individual, particularly during the initial use of nabilone and during dose adjustments. Nabilone is not intended for use on an asneeded basis or as the first antiemetic product prescribed for a patient.

cinacalcet

Drugs

cinacalcet

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of secondary hyperparathyroidism due to chronic kidney disease on dialysis, Or Hypercalcemia due to parathyroid carcinoma, Or severe hypercalcemia in patients with primary hyperparathyroidism who are unable to undergo parathyroidectomy.

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Drugs CLOVIQUE

Covered Uses

Exclusion Criteria

Not recommended in cystinuria or rheumatoid arthritis. Not indicated for biliary cirrhosis.

Required Medical Information

For use in patients with diagnosis of Wilson disease who are intolerant to penicillamine.

Age Restriction

Geriatric patients: Use with caution. Initiate at lower end of the dosing range.Pediatric dosage: Children and adolescents: Oral: Initial: 20 mg/kg/day (round dose to the nearest 250 mg) in 2 to 3 divided doses. Maximum initial daily dose: 1,000 mg/day. titrate dose based on clinical response and free serum copper (non-ceruloplasmin bound copper) concentrations and/or 24-hour urinary copper excretion. usual maintenance dose: 900 to 1,500 mg/day in 2 to 3 divided doses

Prescriber Restriction

Coverage Duration Through end of benefit year.

Drugs

clozapine oral tablet, disintegrating

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

If the patient has any of the following contraindications: agranulocytosis, bone marrow suppression, coma, ileus, leukopenia, myocarditis or neutropenia, OR if the patient has CNS depression, dementia-related psychosis or uncontrolled epilepsy.

Required Medical Information

A statement showing the patient is unwilling or unable to take tablets or capsules orally or at high risk for non-compliance AND is not receiving other tablets or capsules indicating that the patient can take non-dissolvable tablets.

Age Restriction

Prescriber Restriction Part of a clozapine registry.

Coverage Duration Plan Year

Drugs COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

Required Medical Information

Diagnosis of relapsing-remitting multiple sclerosis OR diagnosis of first clinical episode with MRI features consistent with multiple sclerosis.

Age Restriction

Prescriber Restriction By or in consultation with a Neurologist or a Certified MS Specialist

Coverage Duration Plan year

Drugs COPIKTRA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of the following: A) relapsed or refractory CLL or SLL after at least two prior therapies OR B)relapsed or refractory FL after at least two prior systemic therapies.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs CORLANOR ORAL SOLUTION

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Must be clinically diagnosed with A) stable, symptomatic chronic heart failure in adults with left ventricular ejection fraction less than or equal to 35% supported by documentation from the patient's medical records, AND have sinus rhythm with resting heart rate greater than or equal to 70 beats per minute, AND be on maximally tolerated doses of beta blockers unless contraindicated, AND be on optimal therapy with standard treatment of ACEI or ARB unless intolerant or contraindicated, AND be on optimal therapy with standard treatment of an aldosterone antagonist unless intolerant or contraindicated, AND documentation of trial and failure of Entresto OR B)stable symptomatic heart failure due to dilated cardiomyopathy (DCM) in pediatric patients aged 6 months and older AND have sinus rhythm with an elevated heart rate.

Age Restriction

Prescriber Restriction Cardiologist

Coverage Duration Through end of benefit year

Drugs COSENTYX, COSENTYX (2 SYRINGES), COSENTYX PEN, COSENTYX PEN (2 PENS)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented trial and failure of preferred TNF inhibitors (Enbrel and Humira)AND negative Tuberculin test prior to therapy AND patient is free of any clinically important active infections.

Age Restriction

Prescriber Restriction

Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs CRIXIVAN ORAL CAPSULE 200 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older (off-label dosing for pediatrics)

Prescriber Restriction

Coverage Duration Plan year

Drugs CYCLOSET

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information Documented diagnosis of type 2 diabetes mellitus

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs CYSTAGON

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Cysteamine is contraindicated in patients who have demonstrated hypersensitivity to cysteamine or penicillamine hypersensitivity.

Required Medical Information

Documentation of diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Other Criteria

Do not administer intact cysteamine capsules to children less than 6 years old because of aspiration risk. Capsules may be administered by sprinkling contents over food.

Drugs DAURISMO ORAL TABLET 100 MG, 25 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used in combination with low-dose cytarabine for the treatment of newlydiagnosed acute myeloid leukemia (AML), and the patient is A) 75 years of age or older old OR B)has comorbidities that preclude use of intensive induction chemotherapy.

Age Restriction

Prescriber Restriction oncology

Coverage Duration 3 months

Drugs DESCOVY

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs DIACOMIT ORAL CAPSULE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication is prescribed for the treatment of seizures associated with confirmed diagnosis of Dravet syndrome AND member has been inadequately controlled on clobazam and valproate (unless contraindicated) despite optimized therapy AND the member will be receiving concurrent clobazam therapy.

Age Restriction

Member is 2 years of age or older

Prescriber Restriction

Medication is prescribed by a neurologist

Coverage Duration

3 months

Other Criteria

Reauthorization requires documentation of significant decrease in the frequency of seizures

Drugs diazoxide

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used for the management of hypoglycemia due to hyperinsulinism associated with one of the following conditions:A) Adults: Inoperable islet cell adenoma or carcinoma, or extrapancreatic malignanc OR B) Infants and Children: Leucine sensitivity, islet cell hyperplasia, nesidioblastosis, extrapancreatic malignancy, islet cell adenoma, or adenomatosis.

Age Restriction

Prescriber Restriction

Coverage Duration 12 months

Drugs

diclofenac sodium topical gel 3 %

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis and treatment history

Age Restriction DO NOT use Solaraze in children.

Prescriber Restriction

Coverage Duration 3 months

Other Criteria Must have failed topical 5-FU cream

Didanosine (VIDEX)

Drugs

didanosine oral capsule, delayed release (DR/EC) 250 mg, 400 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 2 weeks and older.

Prescriber Restriction

Coverage Duration Plan year

Drugs DIFICID

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of Clostridium difficile associated diarrhea (CDAD) with one of the following: A) Patient has mild to moderate CDAD and failure, contraindication or intolerance to oral Flagyl (metronidazole) and oral Vancocin (vancomycin), or B) Patient has severe CDAD.

Age Restriction 18 years or older

Prescriber Restriction

Coverage Duration 10 Days

Drugs

doxercalciferol oral

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Drugs

dronabinol oral capsule 10 mg, 2.5 mg, 5 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

For treatment of chemotherapy-induced nausea or vomiting refractory to conventional antiemetic agents: 1. Patient is receiving cancer chemotherapy, AND 2. Failure to preferred 5HT-3 receptor antagonist. preferred agents include ondansetron or granisetron, AND 3. Failure to one of the following agents: a. Antihistamine b. Corticosteroid c. Prokinetic agent d. Antipsychotic. For treatment of anorexia associated with weight loss in patients with HIV: documentation of trial and failure, contraindication, or intolerance to megestrol.

Age Restriction

18 years old and greater for the treatment of anorexia associated with weight loss in patients with HIV

Prescriber Restriction

Coverage Duration 3 Months

Drugs DUPIXENT PEN, DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Patient must have the following: A) Moderate-to-severe atopic dermatitis (eczema)AND submission of medical records (e.g. chart notes, laboratory values) documenting the following: Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories:a. Topical calcineurin inhibitor b. High potency topical corticosteroid. OR B) Moderate-to-severe asthma AND submission of medical records documenting the following: 1. Patient has ONE of the following: a. Asthma with eosinophilic phenotype with eosinophil count greater than or equal to 300 cells/mcL in the past 12 months, or b. Oral corticosteroid dependent asthma with at least 1 month of daily oral corticosteroid use within the last 3 months AND 2. Inadequate control of asthma symptoms after a minimum of 3 months of compliant use of one of the following: a. Inhaled corticosteroids & long acting beta2 agonist, or b. Inhaled corticosteroids & long acting muscarinic antagonist. OR C)Chronic rhinosinusitis with nasal polyposis (CRSwNP)AND submission of medical records (e.g. chart notes, laboratory values) documenting the following: Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories: 1. Inadequate response, intolerance or contraindication to ONE medication in EACH of the following categories: a. Nasal corticosteroid spray and b. Oral corticosteroid.

Age Restriction

18 or older

Prescriber Restriction

Dermatologist or allergist/immunologist

Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Other Criteria

Renewals require submission of medical records (e.g. chart notes, laboratory values) documenting improvement of the condition.

EDECRIN (Ethacrynic Acid)

Drugs EDECRIN

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria Use for pediatrics 12 months and younger

Required Medical Information

Documentation of diagnosis.

Age Restriction 13 months and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Must have documented failure, intolerance or contraindication to at least 2 other loop diuretics.

Drugs EDURANT

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older.

Prescriber Restriction

Coverage Duration Plan year

Drugs ELIGARD, ELIGARD (3 MONTH), ELIGARD (4 MONTH), ELIGARD (6 MONTH)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinical documentation of advanced prostate cancer diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration 12 months

Drugs ELZONRIS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Patient must have a definitive diagnosis of Blastic Plasmacytoid Dendritic Cell Neoplasm (BPDCN) AND Patient has CD123 positive expressing disease AND Patient has a baseline serum albumin level of at least 3.2 g per dL

Age Restriction Patient is 2 years of age or older

Prescriber Restriction

Coverage Duration 6 months

Drugs EMCYT

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documented diagnosis of metastatic and/or progressive prostate cancer.

Age Restriction 18 years or older

Prescriber Restriction Oncologist prescriber

Coverage Duration 6 months

Drugs EMEND ORAL CAPSULE 40 MG, 80 MG, EMEND ORAL CAPSULE, DOSE PACK

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration 3 Months

Drugs EMSAM

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Pheochromocytoma. Patient is taking or will take any of the following: SSRIs, SNRIs, tricyclic antidepressants (TCAs), bupropion, buspirone, meperidine, tramadol, methadone, pentazocine, dextromethorphan, St. John's wort, mirtazapine, cyclobenzaprine, oral selegiline, other MAOIs, oxcarbazepine, carbamazepine, and/or sympathomimetic amines.

Required Medical Information

Diagnosis of major depressive disorder, AND 1) Failure of at least two generic oral antidepressants from different classes(at least one should be from the following list: selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, mirtazapine, or bupropion unless contraindicated), AND 2) Patient had an adequate washout period (for patients previously on agents requiring a washout period)

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs EMTRIVA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs ENBREL SUBCUTANEOUS SOLUTION

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Active serious infection (including tuberculosis).

Required Medical Information

Diagnosis of one of the following: A) Moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to at least one non-biologic disease modifying anti-rheumatic drugs (DMARD) and one NSAID for at least 3 months, or B) Moderate to severe polyarticular juvenile idiopathic arthritis and patient had an inadequate response, intolerance or contraindication to at least one DMARD and one NSAID for at least 3 months, OR C) Psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate, or D) Ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to at least two NSAIDs, or E) Moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had an inadequate response, intolerance or contraindication to conventional therapy with to at least two of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (i.e. methotrexate, cyclosporine, acitretin, sulfasalazine) for at least 3 months.

Age Restriction

2 years of age or older for JIA. 4 years of age or older for plaque psoriasis. 18 years of age or older for all other indications.

Prescriber Restriction

Coverage Duration

Through end of benefit year

Other Criteria

Preferred NSAIDs include: ibuprofen, naproxen, ketoprofen, meloxicam. Preferred DMARDs include: methotrexate, sulfasalazine, hydroxychloroquine, cuprimine, azathioprine, leflunomide. Dosing as per FDA approved labeling.

Drugs

ENBREL SUBCUTANEOUS RECON SOLN, ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML), ENBREL SURECLICK

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Tuberculosis (TB), or invasive fungal infections or other active seriousinfections, or history if recurrent infections. Individuals who have not had a tuberculin skin test or Center for Disease Control recommended equivalent to evaluate for latent tuberculosis prior to initiating therapy. Using in combination with other TNF antagonists, IL-1R antagonists, janus kinase inhibitor, anti-cd20 monoclonalantibodies or selective co-stimulation modulators.

Required Medical Information

Documentation of diagnosis, treatment history and TB evaluation. FDA-approved indications include: Ankylosing Spondylitis (AS), Polyarticular Juvenile Idiopathic Arthritis (PJIA), Plaque Psoriasis, Psoriatic Arthritis (PsA), Rheumatoid Arthritis (RA).

Age Restriction

2 years of age or older for JIA. 18 years of age or older for all other indications.

Prescriber Restriction

Rheumatologist (RA, PJIA, PsA, AS), Dermatologist for Plaque Psoriasis

Coverage Duration Plan year

Other Criteria

APPROVE for AS if patient has had an inadequate response, intolerance or contraindication to one or more NSAIDs (e.g. ibuprofen, naproxen, meloxicam, celecoxib). APPROVE for PJIA if patient has has an inadequate response, intolerance or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) (e.g., hydroxychloroquine [HCQ], sulfasalazine, methotrexate [MTX], leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months. APPROVE for Plaque Psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) if patient has had an inadequate response, intolerance or contraindication to conventional therapy with to at least two of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least 3 consecutive months. APPROVE for PsA if patient has had an inadequate response, intolerance, or contraindication to MTX. APPROVE for RA if patient has had an inadequate response to, intolerance to, or contraindication to at least one non-biologic disease modifying anti-rheumatic drugs (DMARD) Patient has been tested for TB and latent TB has been ruled out or is being treated. Dosing as per FDA approved labeling.(e.g., HCQ, sulfasalazine, MTX, leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months.

Drugs ENDARI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented diagnosis of sickle cell disease.

Age Restriction 5 years and older

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs ENTRESTO

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Use of Entresto with an Angiotensin Converting Enzyme (ACE) Inhibitor or an ACE Inhibitor-Containing Product. Use of Entresto with an Angiotensin II Receptor Blocker (ARB) or an ARB-Containing Product. Use of Entresto with Tekturna® (aliskiren tablets) or a Tekturna-Containing Product in patients with diabetes.

Required Medical Information

Clinical documentation of FDA-approved indication for treatment AND the patient has a left ventricular ejection fraction (LVEF) less than or equal to 40% prior to initiation of Entresto.

Age Restriction 18 years and older.

Prescriber Restriction Cardiologist

Coverage Duration Through end of benefit year

Drugs EPCLUSA ORAL TABLET 400-100 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Current alcohol, drug use or reinfection after 3 months of successful treatment. Alcohol urine metabolite and drug screen required.

Required Medical Information

Provider must submit medical records documenting the diagnosis of chronic hepatitis C with genotype and subtype, if applicable (i.e., genotypes 1, 2, 3, 4, 5, or 6) AND submit medical records documenting viral load taken within 6 months of beginning therapy, AND submit medical records documenting F2-F4 fibrosis with a fibrosis score of 0.48 and up or be a documented health Care worker in direct patient care setting.

Age Restriction

18 years of age and older

Prescriber Restriction

Prescribed by, or in consultation with, a gastroenterologist, hepatologist, or infectious disease physician.

Coverage Duration

12 weeks or as defined by current AASLD/IDSA guidance.

Other Criteria

Mavyret, generic Ledipasvir/Sofosbuvir, or generic Sofosbuvir/Velpatasvir must be tried first in patients with chronic hepatitis C. Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance. Must have contraindication to or be unable to tolerate Mavyret, generic Ledipasvir/Sofosbuvir, and generic Sofosbuvir/Velpatasvir.

Drugs EPIDIOLEX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of the following A) diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS)or seizures associated with Dravet syndrome (DS): AND B) normal serum transaminases (ALT and AST) and total bilirubin levels: AND C) inadequate treatment response, intolerance, or contraindication to TWO generic antiepileptic medications (i.e. clobazam, Valproic acid, Lamotrigine, Levetiracetam, Topiramate, etc.)

Age Restriction

2 years of age or older

Prescriber Restriction

Coverage Duration Through end of benefit year

Other Criteria

Dosing is within the FDA labeled dose of up to 20mg/kg/day.

Drugs EPIVIR HBV ORAL SOLUTION

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 3 months of age or older

Prescriber Restriction

Coverage Duration Plan year

EPZICOM (Abacavir/Lamivudine)

Drugs

abacavir-lamivudine, EPZICOM

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented negative HLA-B*5701 screening. Drugs ERLEADA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented diagnosis of non-metastatic, castration-resistant prostate cancer (NM-CRPC).

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Drugs ERTACZO

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 12 years and older.

Prescriber Restriction

Coverage Duration 4 weeks

Other Criteria Failure to generic topical antifungal medications.

Drugs ESBRIET

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

The patient has a diagnosis of idiopathic pulmonary fibrosis confirmed by a high resolution CT scan or biopsy AND the patient does not have evidence or suspicion of an alternative interstitial lung disease diagnosis AND liver function tests have been performed prior to start of therapy and base line PFTs provided.

Age Restriction

Prescriber Restriction

Prescribed by or in consultation with a pulmonologist

Coverage Duration

Through benefit year

Other Criteria

For renewal, patient experienced stabilization from baseline or a less than 10 percent decline in force vital capacity AND the patient has not experienced AST or ALT elevations greater than 5 times the upper limit of normal or greater than 3 times the upper limit of normal with signs or symptoms of severe liver damage.

Drugs EVENITY

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Diagnosis of Osteoporosis in a postmenopausal female AND one or more of the following: 1) History of osteoporotic fracture, or 2) Documented trial and failure of bisphosphonate or 3) Documented contraindication or intolerance to bisphosphonate therapy. Patient has not received more than 1 year of therapy with Evenity.

Age Restriction

Prescriber Restriction

Coverage Duration 12 months

Drugs

AKÝNZEO (FOSNETUPITANT) INTRAVENOUS RECON SOLN, AKYNZEO (NETUPITANT), ALECENSA, ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG, ALUNBRIG ORAL TABLETS, DOSE PACK, BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG, CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG, COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY), COTELLIC, ERIVEDGE, FARYDAK, GILOTRIF, GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG, HYCAMTIN, ICLUSIG ORAL TABLET 15 MG, 45 MG, IDAMYCIN PFS, INLYTA, IRESSA, LONSURF, MEKINIST ORAL TABLET 0.5 MG, 2 MG, NINLARO, ODOMZO, ROZLYTREK ORAL CAPSULE 100 MG, 200 MG, RUBRACA, RYDAPT, SANCUSO, STIVARGA, TAGRISSO ORAL TABLET 40 MG, 80 MG, TOPOSAR, VALCHLOR, VARUBI, VENCLEXTA, VENCLEXTA STARTING PACK, XERMELO, ZEJULA, ZELBORAF, ZYKADIA ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication would be used for an FDA approved indication.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 12 months

Other Criteria For authorization, please submit to EviCore at evicore.com or call at 877-825-7722. Drugs EVOTAZ

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs EXJADE

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Creatinine clearance less than 40 mL/minute. Platelet count less than 50 x 109/L. Poor performance status. Severe (Child-Pugh class C) hepatic impairment. High-risk myelodysplastic syndromes. Advanced malignancies. Gastrointestinal ulceration or hemorrhage.

Required Medical Information

Medical documentation of FDA approved diagnosis, serum ferritin levels and serum creatinine.

Age Restriction

2 years of age or older for chronic iron overload due to transfusions. 10 years of age or older for chronic iron overload due to NTDT.

Prescriber Restriction

Coverage Duration 3 months

Drugs FANAPT ORAL TABLET, FANAPT ORAL TABLETS, DOSE PACK

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and treatment history.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Approve if member has tried two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

Drugs FASENRA PEN

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation that the patient has asthma with an eosinophilic phenotype defined as blood eosinophils greater than or equal to 300 cells/µL within previous 12 months or greater than or equal to 150 cells/µL within 6 weeks of dosing AND medication will be used in combination with a corticosteroid inhaler and long acting beta2-agonist AND Patient must have experienced two or more exacerbations in the previous year OR require daily oral corticosteroids

Age Restriction

Prescriber Restriction

Coverage Duration 6 months

Drugs

fentanyl citrate buccal lozenge on a handle

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Coverage not provided in the management of acute or postoperative pain (including headache/migraines), opiod non-tolerant patients, patients with known intolerance or hypersensitivity to the drug or fentanyl.

Required Medical Information

Diagnosis of cancer AND 1. Use is for breakthrough cancer pain, AND 2. Patient is opioid tolerant and taking at least 60 mg morphine/day, at least 25 mcg transdermal fentanyl/hour, at least 30 mg of oxycodone daily, at least 8 mg oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer, AND 3. Other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, AND 4. Prescriber is registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program.

Age Restriction

16 years or older

Prescriber Restriction

Prescribed by an oncologist or pain specialist.

Coverage Duration 3 months

Other Criteria

Patient must have tried and failed or not responded to the following formulary short-acting narcotics, Oxycodone and morphine. Available only to those enrolled in the Transmucosal Immediate Release Fentanyl (TIRF) EMS Program.

Drugs FERRIPROX ORAL TABLET 500 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Diagnosis of transfusional iron overload due to thalassemia syndromes AND patient has failed prior chelation therapy with Desferal or Exjade (failure is defined as a serum ferritin level greater than 2,500 mcg/L) or patient has a contraindication or intolerance to Desferal or Exjade AND Patient has an absolute neutrophil count greater than 1.5 x 109/L.

Age Restriction

Prescriber Restriction Prescribed by a hematologist/oncologist or hepatologist

Coverage Duration Per treatment

Other Criteria

For renewal, patient has experienced at least a 20% reduction in serum ferritin levels and has an absolute neutrophil count greater than 0.5 x 109/L

Drugs FINTEPLA

Covered Uses

Exclusion Criteria

Required Medical Information

Patients had a clinical diagnosis of Dravet syndrome and seizures that were inadequately controlled on at least 1 antiepileptic drug (AED) OR treatment including vagal nerve stimulation OR ketogenic diet.

Age Restriction Children 2 and older

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs ICATIBANT

Covered Uses All FDA-approved indications not otherwise exluded for part D

Exclusion Criteria

Required Medical Information

Documentation of clinical diagnosis of hereditary angioedema or C1 inhibitor deficiency and having angioedema attacks.

Age Restriction

Prescriber Restriction

Must be prescribed by an allergist, immunologist, hematologist, or a physician that specializes in the treatment of HAE or related disorders.

Coverage Duration 3 months

Drugs FIRDAPSE, RUZURGI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Member has a history of seizures

Required Medical Information

The diagnosis has been confirmed by one of the following: A) Presence of anti-P/Q-type voltage-gated calcium channel (VGCC) antibodies OR B) Characteristic electromyography (EMG).

Age Restriction

Prescriber Restriction Neurologist

Coverage Duration Through the benefit year

Fondaparinux (ARIXTRA)

Drugs

fondaparinux subcutaneous syringe 10 mg/0.8 mL, 2.5 mg/0.5 mL, 5 mg/0.4 mL, 7.5 mg/0.6 mL

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

CrCl (EGFR) less than 30mL/min. Patient's weight less than 50kg.

Required Medical Information

Clinical documentation of FDA approved indication for treatment. Patient's weight and creatinine clearance (CrCl).

Age Restriction

18 years and older.

Prescriber Restriction

Coverage Duration

Per treatment. Post-op DVT prophylaxis 1. hip/knee replacement max of 35 days. 2. abdominal surgery

Drugs FORTEO

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Because of an increased incidence of osteosarcoma, Forteo should not be prescribed for patients who are at increased baseline risk for osteosarcoma (including those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, open epiphyses, or prior radiation therapy involving the skeleton).

Required Medical Information

Approve the following: 1. Treatment of postmenopausal women with osteoporosis at high risk for fracture OR 2. Increase of bone mass in men with primary or hypogonadal osteoporosis at high risk for fracture, AND member meets the following criteria: a. fracture OR b. BMD screening results of -2.5 or below OR c. previously failure / contraindication / intolerance of an oral bisphosphonate AND Prolia 3. Treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy at high risk for fracture, AND member meets the following criteria: a. previous fracture, OR b. multiple risk factors for fracture, OR c. previous failure/contraindication/intolerance of the following: oral bisphosphonate AND Prolia. Patient has not received more than 2 years of therapy with Forteo.

Age Restriction

Prescriber Restriction

Coverage Duration

For initial therapy up to 1 year. Continuation up to 1 year not to exceed 2 years of total therapy.

Other Criteria

Treatment failure is defined as documented continued bone loss after at leaast three months despite treatment with a bisphosphonate or Prolia. Note: Since the effects of long-term treatment with teriparatide are not known at this time, therapy for more than 2 years duration is considered experimental and investigational.

Drugs FUZEON SUBCUTANEOUS RECON SOLN

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria For treatment-experienced patients.

Drugs FYCOMPA ORAL SUSPENSION, FYCOMPA ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs GENVOYA

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs GILENYA ORAL CAPSULE 0.5 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure. History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker. Baseline QTc interval greater than or equal to 500 ms. Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol). Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

Required Medical Information

Diagnosis of a relapsing form of multiple sclerosis or diagnosis of first clinical episode with MRI features consistent with MS AND Patient will be observed for signs and symptoms of bradycardia in a controlled setting for at least 6 hours after the first dose.

Age Restriction

Prescriber Restriction

Prescribed by or in consultation with a neurologist or MS specialist.

Coverage Duration Plan year

Other Criteria

In patients with relapsing forms of MS, Avonex, Copaxone, and Tecfidera must be tried before any other formulary agent will be approved.

Drugs OMNITROPE

Covered Uses

All FDA approved indications not otherwise excluded by Health Plan. Additional off-label coverage is provided for (note - some growth hormone drugs may be labeled for 1 or more of these indications): adult growth hormone deficiency, growth failure in children small for gestational age or with intrauterine growth retardation, idiopathic short stature, GH deficiency associated with Turner Syndrome, growth failure secondary to chronic renal failure/insufficiency in children who have not received a renal transplant, short stature associated with Noonan Syndrome, short bowel syndrome, and for the treatment of Prader-Willi Syndrome.

Exclusion Criteria

Coverage is not provided for constitutional delayed growth

Required Medical Information

Pediatric GHD: epiphyses must be confirmed open in patients 10 years of age and older, AND 1. diagnosis confirmed by any 2 provocative tests or by both low IGF-1 and IGFBP-3 levels in patients who meet the height related conditions of coverage, 2. diagnosis confirmed by 2 provocative tests and both low IGF-1 and IGF-BP3 in patients not meeting height related coverage conditions, or 3. 3 pituitary hormone deficiencies in pt with irreversible hypothalamic-pituitary structural lesions or panhypopituitarism. Growth failure from CRF: PGHD criteria must be met without the provocative tests or IGF-1 and IGF-BP3 related conditions. Idiopathic Short Stature: epiphyses must be confirmed as open in patients greater than or equal 10 years of age, height must be less than or equal - 2.25 sds from the mean. Small for Gestational Age: failure to manifest catch up growth by age 2 defined as birth weight, birth length, or both that are more than 2 sds mean normal values following adjustment for age and gender. Turner's syndrome and Noonan Syndrome: epiphyses must be confirmed as open and when on therapy. Adult GHD: requires either 1. a negative GH provocative test when the AGHD is due to childhood onset GHD, pituitary or hypothalamic disease, surgery or radiation therapy, or trauma, OR 2. 3 pituitary hormone deficiencies and baseline serum IGF-I levels below the age- and sex-appropriate reference range when the AGHD is due to irreversible hypothalamic-pituitary structural lesions are from childhood. Short bowel syndrome: when receiving specialized nutritional support.

Age Restriction

7 years of age or older for Idiopathic short stature

Prescriber Restriction

Pediatric endocrinologist for ISS

Coverage Duration

1 month for short bowel syndrome, 12 months for other indications

Other Criteria

Height related conditions of coverage: 1. height below the third percentile for their age and gender related height, 2. growth velocity subnormal greater than or equal 2 standard deviations (sds) from the age related mean, 3. delayed skeletal maturation greater than or equal 2 sds below the age/gender related mean. Renewals for PGHD, CFR, SGA, Turner's and Noonan Syndromes require growth response of greater than or equal 4.5 cm/yr (pre-pubertal) or greater than or equal 2.5 cm/yr (post-pubertal) AND open epiphyses. For pediatric patients with irreversible hypothalamic-pituitary structural lesions or panhypopituitarism coverage is renewable if the patient has had 3 pituitary hormone deficiencies. Renewals for short bowel syndrome is provided in the presence of clinical benefit (such as, decreasing the patient?s intravenous nutritional requirements). Renewals for Prader-Willi syndrome is provided if pt has increase in lean body mass or decrease in fat mass. Renewals for AGHD is provided in the presence of a growth response of greater than or equal 1.5 cm/yr AND open epiphyses. Renewals for AGHD is provided in the presence of clinical benefit (e.g., increase in total lean body mass, increase in IGF-1 and IGFBP-3 levels, or increase in exercise capacity).

Drugs HARVONI ORAL TABLET 90-400 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

1. Active IV drug users, 2. Active alcohol users, 3. Reinfection after 6 months cure

Required Medical Information

For initial authorization (12 weeks maximum), provider must submit completed HCV Tx form, medical records documenting the diagnosis of chronic hepatitis C with genotype and subtype, if applicable, medical records documenting viral load taken within 6 months of beginning therapy AND submit medical records documenting advanced fibrosis as corresponding to a FibroSure or a Liver Biopsy proven. Other fibrosis scores, physical findings, or clinical evidence consistent with cirrhosis as attested by the prescribing physician may be also considered. For any retreatment or extension of PA, 100% compliance will be required (Claim hx). Urine alcohol metabolite and drug screen required.

Age Restriction

Patient must be 18 years of age or older

Prescriber Restriction

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

Coverage Duration

Based on the AASLD treatment guidelines

Other Criteria

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance. Documentation of F2-F4 fibrosis (fibrosis score of 0.48 and greater) or patient is Health Care worker in direct patient care setting. Must have contraindication to or be unable to tolerate Mavyret, generic Ledipasvir/Sofosbuvir, and generic Sofosbuvir/Velpatasvir.

Drugs HERCEPTIN HYLECTA

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of 1) HER2 overexpressing node positive or node negative breast cancer as part of a treatment regimen consisting of a) doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel or b)docetaxel and carboplatin or c) as a single agent following multi-modality anthracycline based therapy OR 2)HER2-overexpressing metastatic breast cancer either a) in combination with paclitaxel for first-line treatment or b) as a single agent for patients who have received one or more chemotherapy regimens for metastatic disease.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs HORIZANT

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Documented diagnosis of postherpetic neuralgia (PHN) and trial of gabapentin, or diagnosis of moderate to severe primary restless legs syndrome (RLS) in adults and trial of ropinirole and pramipexole.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs

HUMIRA PEN, HUMIRA PEN CROHNS-UC-HS START, HUMIRA PEN PSOR-UVEITS-ADOL HS, HUMIRA SUBCUTANEOUS SYRINGE KIT 10 MG/0.2 ML, 20 MG/0.4 ML, 40 MG/0.8 ML, HUMIRA(CF) PEDI CROHNS STARTER, HUMIRA(CF) PEN CROHNS-UC-HS, HUMIRA(CF) PEN PSOR-UV-ADOL HS, HUMIRA(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 80 MG/0.8 ML, HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Tuberculosis (TB), or invasive fungal infections or other active serious infections, or history if recurrent infections. Individuals who have not had a tuberculin skin test or Center for Disease Control recommended equivalent to evaluate for latent tuberculosis prior to initiating treatmeth. Using in combination with other TNF antagonists. IL-1R antagonists, janus kinase inhibitor, anti-cd20 monoclonalantibodies or selective co-stimulation modulators.

Required Medical Information

Documentation of diagnosis, treatment history and TB evaluation. FDA-approved indications include Ankylosing Spondylitis (AS), Moderate to Severe Crohn's Disease (CD), Hidradenitis Suppurativa (HS), Polyarticular Juvenile Idiopathic Arthritis (PJIA), Plaque Psoriasis, Psoriatic Arthritis (PsA), Rheumatoid Arthritis (RA), Moderate to Severe Ulcerative Colitis (UC), Uveitis

Age Restriction

2 years of age or older for JIA. 6 years of age and older for pediatric Crohn's disease. 18 years of age or older for all other indications.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

APPROVE for AS if patient has had an inadequate response, intolerance or contraindication to one or more NSAIDs (e.g. ibuprofen, naproxen, meloxicam, celecoxib). APPROVE for HS if patient has had an inadequate response, intolerance or contraindication to one or more of the following: intralesional or oral corticosteroids, systemic antibiotics, isotretinoin. APPROVE for PJIA if patient has had an inadequate response, intolerance or contraindication to one or more nonbiologic DMARDs (e.g., hydroxychloroquine [HCQ], sulfasalazine, MTX, leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months. APPROVE for Plague Psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) if patient has had an inadequate response, intolerance or contraindication to conventional therapy with to at least two of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (i.e. MTX, cyclosporine, acitretin, sulfasalazine) for at least 3 consecutive months. APPROVE for PsA if patient has had an inadequate response, intolerance, or contraindication to MTX. APPROVE for RA if patient has had inadequate response to, intolerance to, or contraindication to at least one non-biologic DMARD (e.g., HCQ, sulfasalazine, MTX, leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months. APPROVE for Uveitis if patient has tried one of the following therapies: periocular, intraocular, or systemic corticosteroids, immunosuppressants (azathioprine, MTX, mycophenolate mofetil, cyclophosphamide, cyclosporine). Patient has been tested for TB and latent TB has been ruled out or is being treated. Dosing as per FDA approved labeling.

Drugs IBRANCE

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs IDHIFA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

1. Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation AND 2. Prescriber agrees to monitor for signs and symptoms of differentiation syndrome.

Age Restriction

Prescriber Restriction Oncologist or Hematologist

Coverage Duration Through the end of benefit year

imatinib oral tablet 100 mg, 400 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), or B) Ph+ acute lymphoblastic leukemia (ALL), or C) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, or D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, or E) Hypereosinophilic syndrome or chronic eosinophilic leukemia, or F) Myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutation or with c-KIT mutational status unknown.

Age Restriction

1 year of age or older - newly diagnosed CML in the chronic phase or newly diagnosed Ph+ ALL. 18 years of age or older for other indications.

Prescriber Restriction

Coverage Duration Plan year

Drugs IMBRUVICA ORAL CAPSULE 140 MG, 70 MG, IMBRUVICA ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Drugs INBRIJA

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

patient is not currently taking or has recently (within 2 weeks) taken a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine and tranylcypromine)

Required Medical Information

Patient is currently treated with carbidopa/levodopa AND is experiencing intermittent OFF episodes secondary to Parkinson's disease

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Other Criteria

Reauthorization requires physician attestation of medications efficacy

Drugs INCRELEX

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Insulin-like growth factor therapy is considered NOT medically necessary when any of the following criteria are met: Final adult height has been reached as determined by the 5th percentile of adult height OR the bone epiphyses are closed OR the patient is older than 18 years of age. Contraindicated in neonates, patients with closed epiphyses, and suspected neoplasia.

Required Medical Information

1. All of the following: a. Diagnosis of severe primary IGF-1 deficiency. b. Height standard deviation score of -3.0 or less. c. Basal IGF-1 standard deviation score of -3.0 or less. d. Normal or elevated growth hormone. e. Open finger epiphyses on last boneradiograph GH gene deletion: a. Diagnosis of growth hormone gene deletion who have developed neutralizing antibodies to GH, AND b. Have open finger epiphyses on last bone radiograph.

Age Restriction

The patient is between 2 years -18 years old for Increlex therapy

Prescriber Restriction

Must be endocrinologist to prescribe

Coverage Duration 6 months to 1 year

Other Criteria

Not a substitute for GH treatment. For renewal, Patient had a minimum growth rate of at least 2 cm/year.

Drugs INREBIC

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Patients on treatment with ruxolitinib before initiation must taper and discontinue according to ruxolitinib prescribing information

Required Medical Information

Must provide labs showing patient is not thiamine deficient before starting drug

Age Restriction 18 years of age and older

Prescriber Restriction Oncologist or Hematologist

Coverage Duration 3 months

Drugs INTELENCE ORAL TABLET 100 MG, 200 MG, 25 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria For treatment-experienced patients.

Drugs INTRON A INJECTION

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Type B viral Hepatitis (HBeAg positive): Serum HBsAg positive for at least six months, AND elevated serum ALT 2 times ULN or moderate to severe hepatitis or fibrosis on biopsy. Type B Viral Hepatitis (HBeAg negative) HBsAG positive for at least 6 months AND BHV DNA level of 2000 IU/ml or more than 11,200 copies/ml AND One of the following, persistent ALT 2 times UNL or moderate to severe hepatitis or fibrosis on biopsy. Documentation must be provided showing trial and failure to our preferred agent Peg-Intron. Chronic Hepatitis C: Positive HCV antibody and HCV RNA. Documentation must be provided showing trial and failure to our preferred agent Peg-Intron. Condyloma Acuminatum or Perianal Warts: Must have documentation of trial and failure to preferred alternative or intolerance/contraindication to preferred alternatives. For external perianal warts, condylox gel, for external genital warts, podofilox, or imiquimod. Hairy Cell Leukemia: Medical documentation indicating diagnosis. Malignant Melanoma: Indicated as adjuvant to surgical treatment with malignant melanoma who are free of disease but at high risk for systemic recurrence, within 56 days of surgery. Follicular Lymphoma: Indicated for the initial treatment of clinically aggressive follicular Non-Hodgkins Lymphoma in conjunction with anthracycline-containing combination chemotherapy. Efficacy in patients with low-grade, low-tumor burden follicular Non-Hodgkins Lymphoma has not been demonstrated. AIDS-Related Kaposis Sarcoma: Indicated for the treatment of selected patients. The likelihood of response to therapy is greater in patients who are without systemic symptoms, who have limited lymphadenopathy and who have a relatively intact immune system as indicated by total CD4 count.

Age Restriction

For Hepatitis B- age 1 or older, For Hepatitis C - age 3 or older, All other diagnoses- 18 years or older.

Prescriber Restriction

Coverage Duration Plan year

Drugs INVEGA SUSTENNA, paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 6 mg, 9 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of Schizophrenia AND documented treatment failure or intolerable side effects from treatment with two formulary antipsychotic medications such as risperidone, ziprasidone,

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs INVIRASE ORAL TABLET

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Pretreatment EKG should be done and therapy is NOT to be initiated if QT interval exceeds 450 msec. Must be used in combination with ritonavir.

Drugs ISENTRESS HD, ISENTRESS ORAL POWDER IN PACKET, ISENTRESS ORAL TABLET, ISENTRESS ORAL TABLET, CHEWABLE

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs ISTURISA ORAL TABLET 1 MG, 10 MG, 5 MG

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosis of Cushing disease: Treatment of Cushing disease in adults for whom pituitary surgery is not an option or has not been curative.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Itraconazole (Sporanox)

Drugs

itraconazole oral capsule

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Other Criteria

For onychomycosis - must have documented failure, intolerance or contraindication to terbinafine.

Drugs GAMMAGARD LIQUID

Covered Uses

All medically accepted indications not otherwise excluded by Health Plan, chronic inflammatory demyelinating polyneuropathy, multifocal motor neuropathy, dermatomyositis, polymyositis, Guillain-Barre syndrome (GBS), relapsing-remitting multiple sclerosis (RRMS), myasthenia gravis, Lambert-Eaton myasthenic syndrome, Kawasaki syndrome, idiopathic thrombocytopenic purpura, pure red cell aplasia (PRCA), fetal/neonatal alloimmune thrombocytopenia, and prophylaxis of bacterial infections in B-cell chronic lymphocytic leukemia (CLL), bone marrow/hematopoietic stem cell transplant (BMT/HSCT) recipients, and pediatric HIV infection.

Exclusion Criteria

IgA deficiency with antibodies to IgA and a history of hypersensitivity. History of anaphylaxis or severe systemic reaction to human immune globulin or product components.

Required Medical Information

Documentation of diagnosis and previous treatment. For dermatomyositis and polymyositis: standard 1st line treatments (corticosteroids or immunosuppressants) have been tried but were unsuccessful or not tolerated OR patient is unable to receive standard therapy because of a contraindication or other clinical reason. For GBS: physical mobility must be severely affected such that the patient requires an aid to walk AND IVIG therapy must be initiated within 2 weeks of symptom onset. For RRMS: standard 1st line treatments (e.g. interferon, glatiramer, dimethyl fumarate) have been tried but were unsuccessful or not tolerated OR patient is unable to receive standard therapy because of a contraindication or other clinical reason. For CLL: serum IgG less than 500 mg/dL OR a history of recurrent bacterial infections. For BMT/HSCT: serum IgG less than 400 mg/dL. For pediatric HIV infection: serum IgG less than 400 mg/dL OR a history of recurrent bacterial infections. PRCA is secondary to parvovirus B19 infection. For all indications: patients with any of the following risk factors for renal dysfunction must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: pre-existing renal insufficiency, diabetes mellitus, age over 65 years, volume depletion, sepsis, paraproteinemia, or receiving concomitant nephrotoxic drugs. For all indications: patients with any of the following risk factors for thrombosis must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: pre-existing renal insufficiency, diabetes mellitus, age over 65 years, volume depletion, sepsis, paraproteinemia, or receiving concomitant nephrotoxic drugs. For all indications: patients with any of the following risk factors for thrombosis must receive the minimum dose or concentration available of IVIG and the minimum infusion rate practicable: age 45 years or older, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs JAKAFI ORAL TABLET 10 MG, 15 MG, 20 MG, 25 MG, 5 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Statement of diagnosis for treatment of patients with intermediate or highrisk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis and lab work indicating a complete blood count and platelet count before initiating therapy and recent lab work indicating complete blood count and platelet count for a dosage adjustment. Lab work must indicate platelets are more than 50 x 109/L and dose must be less than 50 mg per day. No dose increases will be approved within 4 weeks of therapy and not more frequently than every 2 weeks. If no spleen reduction or symptom improvement after 6 months then discontinue the drug.

Age Restriction

18 years or older

Prescriber Restriction

Myelofibrosis: Prescribed by a Hematologist/Oncologist

Coverage Duration 3 months

Drugs JYNARQUE ORAL TABLETS, SEQUENTIAL 45 MG (AM)/ 15 MG (PM), 60 MG (AM)/ 30 MG (PM), 90 MG (AM)/ 30 MG (PM)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Medication requested is being used to slow kidney function decline AND Liver function laboratory values (ALT, AST and bilirubin) have been reviewed and are appropriate before initiation.

Age Restriction

Patient is 18 years of age or older

Prescriber Restriction

Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs KALETRA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs KALYDECO ORAL GRANULES IN PACKET 50 MG, 75 MG, KALYDECO ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Medical documentation of cystic fibrosis AND member has a G551D, G1244E, G1349D, G178R, G551S, R117H, S1251N, S1255P, S549N, or S549R mutation in the CFTR gene AND member does not have a Homozygous F508del mutation in CFTR gene.

Age Restriction

Ivacaftor oral granules are approved in patients 2 years of age and older. Ivacaftor oral tablets are approved in patients 6 years of age and older.

Prescriber Restriction

Endocrinologist or Pulmonologist

Coverage Duration Plan year

Drugs KESIMPTA PEN

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondaryprogresive MS)

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs KISQALI, KISQALI FEMARA CO-PACK

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documentation that the member has tried and failed or has a contraindication to Ibrance.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs KOSELUGO ORAL CAPSULE 10 MG, 25 MG

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation that the member has neurofibromatosis type 1 (NF1) who have symptomatic, inoperable plexiform neurofibromas (PN).

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 months

Drugs KUVAN ORAL TABLET,SOLUBLE

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction 1 month and older

Prescriber Restriction

Coverage Duration

Initial: 2 months. Renewal: through plan year

Other Criteria

For initial approval, Patient will have phenylalanine levels measured one week after starting therapy and periodically for up to two months of therapy to determine response. For renewal, patient has been determined to be a responder to therapy (i.e. phenylalanine levels have decreased by at least 30% from baseline) and phenylalanine levels will be measured periodically during therapy.

Lamivudine (EPIVIR)

Drugs

lamivudine oral solution, lamivudine oral tablet 100 mg, 150 mg, 300 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs LEDIPASVIR-SOFOSBUVIR

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy, AND 3) fibrosis score to confirm appropriate duration of treatment, AND 4) documentation of previous HCV therapies to confirm appropriate duration of treatment.

Age Restriction

Patient must be 12 years of age or older

Prescriber Restriction

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

Coverage Duration

12 to 24 weeks based on the AASLD treatment guidelines

Other Criteria

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

Drugs LENVIMA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Through benefit year

ambrisentan, LETAIRIS

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Known or suspected pregnancy. Treat women of child-bearing potential only after a negative pregnancy test and treat only women who are using two reliable methods of contraception OR have had a tubal sterilization OR a Copper T 380A IUD or LNg 20 IUD inserted.

Required Medical Information

Diagnosis of Pulmonary Arterial Hypertension (PAH) AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential. Trial and failure of Revatio or Adcirca.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration 3 months

Drugs LEUKINE INJECTION RECON SOLN

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Chemotherapy or radiotherapy within 24 hours or concomitantly, excess leukemic myeloid blasts in the bone marrow or blood (10% or greater), hypersensitivity to granulocyte-macrophage colony-stimulating factor (GM-CSF) or yeast-derived products, allergic or anaphylactoid reactions to the medication in the past.

Required Medical Information

Medical statement indicating diagnosis AND trial and failure of preferred agent neupogen AND Absolute Neutrophil Count less than 10,000/mm3 and CBC with differential.

Age Restriction

Patients requiring prophylaxis of febirle neutropenia in acute myelogenous leukemia following induction chemotherapy must be at least 55 years of age, other diagnoses do not specify an age restriction

Prescriber Restriction Oncologist or Hematologist

Coverage Duration Plan year

leuprolide

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.

Required Medical Information

Diagnosis of one of the following: A) Advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), or B) Central precocious puberty (Lupron Depot-Ped) AND submission of pubertal gonadal sex steroid levels (testosterone greater than 30 ng/dL, estradiol greater than 20 pg/mL AND a pubertal LH increase upon native GnRH stimulation AND pelvic ultrasound assessment (girls) is required for approval along with notes indicating premature development of secondary sexual characteristics at or before the age of 8 yrs in girls and 9 yrs in boys and significant advancement of bone age and/or a poor adult height prediction AND other causes of sexual precocity must be excluded.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

fosamprenavir, LEXIVA

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

lidocaine topical adhesive patch, medicated 5 %, lidocaine topical ointment

Covered Uses

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded by Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

linezolid

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Patients that are currently myelosuppressed due to any cause

Required Medical Information

Culture and sensitivity reports verifying: 1. VRE infections within past 30 days. 2. Nosocomial pneumonia (MRSA) within past 30 days. 3. Nosocomial or CAP (MSSA or S. pneumoniae) within past 30 days and failure/resistance to 2 preferred antibiotics 4. Complicated SSI without osteomyelitis (MRSA) within past 30 days. 5. Uncomplicated SSI (MRSA) within past 30 days or empirical treatment of uncomplicated or community-acquired complicated SSI without osteomyelitis (MRSA likely) and failure/resistance to 2 preferred antibiotics. 6. Uncomplicated or complicated SSI without osteomyelitis (MSSA, S. pyogenes, or S. agalactiae (complicated SSI only)) within past 30 days and failure/resistance to 2 preferred antibiotics.

Age Restriction

Prescriber Restriction

Presribing physician must be an infectious disease specialist

Coverage Duration

Non-MRSA nosocomial or community acquired pneumonia, SSI: 14 days Other uses: 28 days

Other Criteria

Nosocomial or community acquired pneumonia (MSSA or S. pneumoniae) preferred antibiotics: Amoxicilin/Clavulanate, Azithromycin, Cephalexin, Clarithromycin, Levaquin. Uncomplicated SSI (MRSA) or empirical treatment of patients with uncomplicated or community-acquired complicated SSI without osteomyelitis (MRSA likely) preferred antibiotics: Trimethoprim/sulfamethoxazole, Tetracycline, Doxycycline, Minocycline, Clindamycin. Uncomplicated or complicated SSI without osteomyelitis (MSSA, S. pyogenes, or S. agalactiae (complicated SSI only)) preferred antibiotics: Amoxicillin/clavulanate, Cephalexin, Ciprofloxacin, Clindamycin, Levaquin, Trimethoprim/Sulfamethoxazole, Dicloxacillin. Drugs LOKELMA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of elevated serum potassium (greater than 5.0 mEq/L) and the beneficiary has failure, contraindication or intolerance to sodium polystyrene sulfonate oral suspension.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration 6 months

Drugs LORBRENA ORAL TABLET 100 MG, 25 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) whose disease has progressed on one of the following: 1. crizotinib and at least one other ALK inhibitor for metastatic disease: or 2. alectinib as the first ALK inhibitor therapy for metastatic disease: or 3. ceritinib as the first ALK inhibitor therapy for metastatic disease.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

LUPRON DEPOT, LUPRON DEPOT (3 MONTH), LUPRON DEPOT (4 MONTH), LUPRON DEPOT (6 MONTH), LUPRON DEPOT-PED, LUPRON DEPOT-PED (3 MONTH)

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.

Required Medical Information

Diagnosis of one of the following: A) Advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), or B) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only) AND 1. For initial authorization, patient has had an inadequate pain control response or has an intolerance or contraindication to one of the following: Danazol or combination [estrogen/progesterone] oral contraceptives or progestins, or 2. For retreatment course, patient is experiencing recurrence of symptoms after an initial course of therapy with leuprolide acetate and norethindrone acetate 5 mg daily will be co-administered, or C) Anemia due to uterine Leiomyomata (Fibroids) (3.75 mg 1-month &11.25 mg 3-month depots only) AND patient is preoperative AND has tried and had an inadequate response to monotherapy with iron, or D) Central precocious puberty (Lupron Depot-Ped) AND submission of pubertal gonadal sex steroid levels (testosterone greater than 30 ng/dL, estradiol greater than 20 pg/mL AND a pubertal LH increase upon native GnRH stimulation AND pelvic ultrasound assessment (girls) is required for approval along with notes indicating premature development of secondary sexual characteristics at or before the age of 8 yrs in girls and 9 yrs in boys and significant advancement of bone age and/or a poor adult height prediction AND other causes of sexual precocity must be excluded.

Age Restriction

Prescriber Restriction

Oncologist, Endocrinologist, or Gynecologist to prescribe

Coverage Duration

Plan year

Other Criteria

For endometriosis and uterine fibroids, patient will be using nonhormonal contraception during and for 12 weeks after therapy.

Drugs LYNPARZA ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

1. Documentation of deleterious germline BRCA mutated ovarian cancer AND 2. Documentation of at least 3 prior chemotherapy regimens that have been ineffective or not tolerated AND 3. Lynparza will be used as monotherapy.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs MAKENA (PF), MAKENA INTRAMUSCULAR OIL 250 MG/ML (1 ML)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinical documentation of singleton pregnancy (i.e. one fetus) AND a history of singleton spontaneous preterm birth defined as delivery prior to 37 weeks gestation AND the pregnancy is between 16 weeks, 0 days and 20 weeks, 6 days gestation AND the requested dose and frequency is in accordance with FDA-approved labeling.

Age Restriction

16 years of age or older

Prescriber Restriction

Coverage Duration

coverage is provided until week 37 (through 36 weeks, 6 days) of gestation

MAVENCLAD (10 TABLET PACK), MAVENCLAD (4 TABLET PACK), MAVENCLAD (5 TABLET PACK), MAVENCLAD (6 TABLET PACK), MAVENCLAD (7 TABLET PACK), MAVENCLAD (8 TABLET PACK), MAVENCLAD (9 TABLET PACK)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Individual with current malignancy OR human immunodeficiency virus (HIV) infection OR an active chronic infection (e.g., hepatitis or tuberculosis)

Required Medical Information

Documentation of diagnosis of relapsing multiple sclerosis, including relapsing-remitting disease or active secondary progressive disease AND the patient has had a trial and inadequate response or intolerance to at least one alternative drug indicated for the treatment of multiple sclerosis.

Age Restriction

Prescriber Restriction Coverage Duration

1 year

Other Criteria Will only be approved for 2 treatment cycles.

Drugs MAVYRET

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

For retreatment, patient was non-adherent to initial regimen as evidenced by medical record and/or pharmacy claims OR patient continues to engage in high risk behavior and experienced reinfection secondary to high risk behavior.

Required Medical Information

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy, AND 3) documentation of previous HCV therapies to confirm appropriate duration of treatment. Authorization for retreatment requires the following: 1)Evidence of failure to achieve a sustained virologic response (SVR) or lack of efficacy during treatment (polymerase chain reaction (PCR) assay, 12 or more weeks after completing treatment or a 10-fold increase of viral load at week 6 of treatment) OR evidence of adverse event that required therapy discontinuation (Laboratory results and/or clinical presentation), AND 2) Member was adherent to previous therapy as evidenced by pharmacy claims, AND 3) Submission of psychological support/treatment for a minimum of six months for substance abuse related failure (i.e. NA, AA), AND 4)Patient has abstained from the use of illicit drugs and alcohol for a minimum of 3 months as evidenced by negative urine or blood confirmation tests, collected monthly for the past 90 days prior to initiation of therapy.

Age Restriction

Prescriber Restriction

Prescribed by, a gastroenterologist, hepatologist, or infectious disease physician.

Coverage Duration

based on the AASLD treatment guidelines

Other Criteria

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

Drugs MEKTOVI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDAapproved test and that Mektovi will be used in combination with encorafenib.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs MENEST

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For treatment of vaginal atrophy - must have documented failure, intolerance or contraindication to at least 1 formulary vaginal estrogen. (Estrace cream, Premarin cream, Vagifem tab)

Miglustat

Drugs miglustat

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of mild to moderate type 1 Gaucher disease AND patient is symptomatic (i.e. radiologic evidence of skeletal disease, platelet count less than 60,000 microL, liver greater than 2.5 times normal size, spleen greater than 15 times normal size) AND enzyme replacement therapy is not a therapeutic option (e.g. due to allergy, hypersensitivity, or poor venous access).

Age Restriction 18 years of age or older

Prescriber Restriction

Coverage Duration 3 Months

Drugs modafinil

Covered Uses All FDA-Approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and prior therapies used

Age Restriction

Prescriber Restriction

Idiopathic hypersomnia-- approve if the diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)

Coverage Duration Plan year

Drugs MOVANTIK

Covered Uses

All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Known or suspected gastrointestinal obstruction and at increased risk of recurrent obstruction, due to the potential for gastrointestinal perforation. Concomitantly taking strong CYP3A4 inhibitors (e.g., clarithromycin, ketoconazole)

Required Medical Information

Documentation of diagnosis and treatment history.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Approve if member has been taking an opioid for at least 4 weeks and has tried lifestyle changes (e.g. maintaining a diet rich in fiber and/or fiber supplementation along with adequate fluid intake) and has tried a bowel regimen of an osmostic laxative (e.g. PEG 3350) or a stimulant laxative (e.g. bisacodyl) with or without a stool softener (e.g. docusate).

Drugs MULPLETA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used for the treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure and had a platelet count less than 50 × 10^9/L

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Drugs NAGLAZYME

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs NERLYNX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

NERLYNX is indicated for the extended adjuvant treatment of adult patients with early stage HER2overexpressed/amplified breast cancer, to follow adjuvant trastuzumab based therapy.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration Through the end of benefit year

Drugs NEULASTA

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis.

Age Restriction

Prescriber Restriction

Coverage Duration

3 months and is renewable in situations where it continues to provide clinical benefit

Drugs NEUPOGEN

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Varies by indication.

Drugs nevirapine

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs NEXAVAR

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Combination use with other tyrosine kinase inhibitors such as sorafenib, sunitinib. Squamous cell lung cancer being treated with carboplatin and paclitaxel.

Required Medical Information

Diagnosis of one of the following: A) Advanced renal cell carcinoma, or B) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma refractory to radioactive iodine treatment, or C) Unresectable hepatocellular carcinoma.

Age Restriction

Patient must be at least 18 years old or older.

Prescriber Restriction

Coverage Duration Initial: 3 months, Renewal: through end of benefit year w/ stable disease

Drugs NEXLETOL

Covered Uses

Exclusion Criteria

Required Medical Information

Clinical documentation (e.g., chart notes, laboratory values) required for initial authorization includes: 1. Documentation of one of the following diagnoses: A) Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia), OR B) Atherosclerotic cardiovascular disease (ASCVD), AND 2. Documentation of any one of the following: a) Patient has been receiving statin therapy at a maximally tolerated dose and (maximally tolerated dose may include no statin therapy) OR b) Patient has a documented labeled contraindication to all statins OR c) Patient has experienced rhabdomyolysis, AND 3. Patient has tried and failed ezetimibe. Clinical documentation required for reauthorization includes: 1. Physician attestation of medication's efficacy.

Age Restriction

18 years or older

Prescriber Restriction

Coverage Duration Through end of benefit year

Other Criteria

Other Criteria: Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia): The patient has a lowdensity lipoprotein cholesterol (LDL-C level) greater than or equal to 100 mg/dL. C) ASCVD: The patient has a lowdensity lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin. Drugs NEXLIZET

Covered Uses

Exclusion Criteria

Required Medical Information

Clinical documentation (e.g., chart notes, laboratory values) required for initial authorization includes: 1. Documentation of one of the following diagnoses: A) Heterozygous familial hypercholesterolemia, OR B) Atherosclerotic cardiovascular disease (ASCVD) established, AND 2. Documentation of any one of the following: a) Patient has been receiving statin therapy at a maximally tolerated dose (maximally tolerated dose may include no statin therapy), OR b) Patient has a documented labeled contraindication to all statins, OR c) Patient has experienced rhabdomyolysis. Clinical documentation required for reauthorization includes: 1. Physician attestation of medication's efficacy.

Age Restriction

18 years and older.

Prescriber Restriction

Coverage Duration

Through end of benefit year

Other Criteria

Other Criteria: Primary Hyperlipidemia (Including Heterozygous Familial Hypercholesterolemia): The patient has a lowdensity lipoprotein cholesterol (LDL-C level) greater than or equal to 100 mg/dL. C) ASCVD: The patient has a lowdensity lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin.

Drugs NORPACE CR

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis and treatment history including reason why disopyramide IR cannot be used.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs NORVIR ORAL CAPSULE, NORVIR ORAL SOLUTION

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs NOXAFIL

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 13 years and older.

Prescriber Restriction

Coverage Duration

Per treatment OR up to through plan year.

Other Criteria

Fluconazole preferred for candida. Voriconazole preferred for aspergillus.

Drugs NUBEQA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documenation of non-metastatic castration resistant prostate cancer (nmCRPC).

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 months

Drugs NUCALA SUBCUTANEOUS AUTO-INJECTOR

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that either A) patient has asthma with an eosinophilic phenotype defined as blood eosinophils greater than or equal to 300 cells/µL within previous 12 months or greater than or equal to 150 cells/µL within 6 weeks of dosing AND medication will be used in combination with a corticosteroid inhaler and long acting beta2-agonist AND Patient must have experienced two or more exacerbations in the previous year OR require daily oral corticosteroids OR B)patient has eosinophilic granulomatosis with polyangiitis (EGPA)AND documented trial and failure of or contraindication to treatment with at least one immunosuppressants (azathioprine, cyclophosphamide, or methotrexate).

Age Restriction

Prescriber Restriction

Coverage Duration 6 months

Drugs NUCYNTA, NUCYNTA ER

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and treatment history.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented failure to tramadol or tramadol extended-release. Drugs NULOJIX

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Transplant recipients who are Epstein-Barr virus (EBV) seronegative or with unknown EBV serostatus.

Required Medical Information

For prophylaxis of organ rejection in adults receiving kidney transplant, in combination with basiliximab induction, mycophenolate mofetil, and corticosteroids, AND documentation of patient's EBV serostatus.

Age Restriction 18 years of age or older

Prescriber Restriction

Coverage Duration Plan year

Drugs NURTEC ODT

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used for the acute treatment of migraine with or without aura in adults AND the member has tried and failed three alternatives two of which were triptans, unless contraindicated.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Other Criteria

Reauthorization requires documentation of medication efficacy.

Drugs NYMALIZE ORAL SYRINGE

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosis of Subarachnoid hemorrhage: For the improvement of neurological outcome by reducing the incidence and severity of ischemic deficits in adult patients with subarachnoid hemorrhage (SAH) from ruptured intracranial berry aneurysms regardless of their postictus neurological condition (ie, Hunt and Hess grades I to V)

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs

octreotide acetate injection solution

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses, or B) Metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes, or C) Vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Acromegaly: Documentation of inadequate response to surgery and/or radiotherapy, or documentation that patient is not a candidate for surgery and/or radiotherapy. Reauthorization will require statement indicating growth hormone (GH) levels are stabilized at less than 5.0ng/mL and IGF-1 levels are normalized (male less than 1.9U/mL or female less than 2.2 U/mL) as matched by age and gender, or the patient has a documented clinical response defined by a reduction of tumor mass, a reduction in the signs and symptoms of acromegaly, or an improvement in significant comorbidities.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

Drugs ODEFSEY

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs OMNIPOD DASH 5 PACK POD, OMNIPOD DASH PDM KIT, OMNIPOD INSULIN MANAGEMENT, OMNIPOD INSULIN REFILL

Covered Uses

Exclusion Criteria

Required Medical Information

Initial Therapy 1)Diagnosis of Type 1 Diabetes2)Member has previously been on a maintenance program involving at least three injections of insulin per day and frequent self-adjustments of insulin dosage or current use of an insulin pump3)Member performs glucose self-testing at least three times per day4)History of suboptimal blood sugar control despite appropriate management5)Member or caregiver has completed a physician-directed comprehensive diabetes management programContinued Therapy1)There is documented evidence of improvement in control of diabetes (specific to baseline status of disease for individual members)

Age Restriction None

Prescriber Restriction Endocrinologist

Coverage Duration 12 months

Other Criteria

If request is for more than 10 pods per 30 days additional documentation is required to provide clinical rationale for higher quantity

Drugs ORFADIN ORAL CAPSULE 10 MG, 2 MG, 5 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Medical statement of diagnosis of hereditary tyrosinemia type 1 (HT-1) AND current patient weight as dose must be within FDA approved dosing range: maximum dosage for all patients is 2 mg/kg/day. When initiating therapy, Serum tyrosine should be below 500 mmol/L to avoid toxic effects, and urinary succinylacetone levels should be undetectable.

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

ORKAMBI (lumacaftor-ivacaftor)

Drugs ORKAMBI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Use in combination with Kalydeco

Required Medical Information

The patient is positive for the F508del mutation on both alleles of the CFTR gene.

Age Restriction 6 years of age or older

Prescriber Restriction

Coverage Duration Through end of benefit year.

Drugs OTEZLA, OTEZLA STARTER

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented trial and failure of preferred TNF inhibitors (Enbrel and Humira)AND patient is free of any clinically important active infections.

Age Restriction

Prescriber Restriction

Coverage Duration 12 months

Drugs

oxandrolone oral tablet 10 mg, 2.5 mg

Covered Uses

All medically accepted indications not otherwise excluded from Health Plan.

Exclusion Criteria

Pregnancy Category X

Required Medical Information

Statement indicating use to promote weight gain after weight loss following extensive surgery, chronic infections, or severe trauma, and in some patients who without definite pathophysiologic reasons fail to gain or to maintain normal weight, and to offset the protein catabolism associated with prolonged administration of corticosteroids. Statement indicating use for orphan drug indication, short stature associated with Turner syndrome, constitutional delay of growth and puberty, moderate or severe acute alcoholic hepatitis, Duchenne and Becker muscular dystrophy. Initial Therapy for AIDS Wasting: Diagnosis of AIDS wasting/cachexia. For treatment of anorexia associated with weight loss in patients with HIV: 1. Patient is receiving AIDS anti-retroviral therapy AND 2. experienced as least a. 7.5% unintentional weight loss over 6 months b. 10% unintentional weight loss over 12 months c. 5% body cell mass (BCM) loss within 6 months d. BMI less than 20 kg/m2 e. BCM less than 35% male (less than 23% female) and a BMI less than27 kg/m2 AND 3. documentation of trial and failure, contraindication, or intolerance to megestrol at doses up to 800mg daily.

Age Restriction

Prescriber Restriction

Coverage Duration HIV Wasting: 3 months. All other indications: Through the Benefit Year

Drugs OXERVATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Member has a diagnosis (documented in chart notes) of stage 2 (recurrent/persistent epithelial defect) or stage 3 (corneal ulcer) neurotrophic keratitis in the affected eye(s) AND Member is refractory to at least ONE conventional non-surgical treatment for neurotrophic keratitis (e.g. preservative-free artificial tears, topical antibiotic eyedrops, therapeutic contact lenses, etc.)

Age Restriction Member is 2 years of age or older

Prescriber Restriction

The medication is prescribed by an ophthalmologist

Coverage Duration 8 weeks

Drugs PALYNZIQ

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Not to be used in combination with sapropterin dihydrochloride (Kuvan)

Required Medical Information

Documented Diagnosis of Phenylketonuria (PKU), AND blood phenylalanine concentration greater than 600 micromol/L, AND physician agrees to assess patient tolerability, blood phenylalanine concentration, dietary protein and phenylalanine intake throughout treatment, AND prescriber and patient must be enrolled with the Palynziq REMS Program.

Age Restriction 18 years of age and older

Prescriber Restriction

Coverage Duration 3 Months

PAMIDRONATE

Drugs

pamidronate

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment and treatment history. For treatment of hypercalcemia of malignancy, documentation of corrected total serum calcium greater than or equal to 12 mg/dL. For treament of bone metastases, diagnosis of breast cancer or multiple myeloma. For Paget's disease, must have symptomatic form of disease.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For Paget's disease, must have documented failure, intolerance or contraindication to oral agents: alendronate or risedronate.

Drugs PANRETIN

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction 18 years and older

Prescriber Restriction Oncologist or HIV specialist

Coverage Duration Plan year

paricalcitol (ZEMPLAR)

Drugs

paricalcitol oral

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and treatment history.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented failure or intolerance to calcitriol.

Drugs PEGINTRON SUBCUTANEOUS KIT 50 MCG/0.5 ML

Covered Uses

All medically accepted indications not otherwise excluded from Health Plan.

Exclusion Criteria

Uncontrolled depression. Autoimmune hepatitis. Known hypersensitivity reactions (urticaria, angioedema, bronchoconstriction, anaphylaxis, or Stevens-Johnson syndrome) to alpha interferons or any of its components. Hepatic decompensation in cirrhotic patients.

Required Medical Information

Documentation of diagnosis

Age Restriction

Prescriber Restriction

All patients with hepatitis C or hepatitis B, peginterferon must be prescribed by an infectious disease physician, gastroenterologist, hepatologist, or a transplant physician or in consultation with these physicians

Coverage Duration

12 Weeks to 12 Months

Drugs PEMAZYRE

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation that the member has been diagnosed with previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 Months

Penicillamine capsule

Drugs penicillamine oral capsule, penicillamine oral tablet

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Drugs PENTAM

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 4 months and older

Prescriber Restriction

Coverage Duration Per treatment

Drugs PERSERIS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

Age Restriction

Prescriber Restriction

Coverage Duration Through the benefit year

Drugs PICATO

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years or older

Prescriber Restriction

Coverage Duration Per treatment

Other Criteria Must have failed 5-FU cream

Drugs PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the patient is a postmenopausal female or a male AND has advanced or metastatic breast cancer AND has HR-positive disease AND has HER2-negative disease AND has PIK3CA-mutated breast cancer as detected by a FDA approved test AND has progressed on or after at least one prior endocrine-based regimen AND the medication will be used in combination with fulvestrant.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 Months

Drugs PLEGRIDY

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondary-progresive MS)

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs POMALYST

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Patient has a diagnosis of multiple myleoma and the patient has received two prior therapies, including Revlimid and Velcade unless the patient has a contraindication or intolerance to Revlimid or Velcade and the patient has demonstrated disease progression on or within 60 days of completion of last therapy.

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

Other Criteria

Prescriber, pharmacist, and patient must be enrolled in the Pomalyst REMS program.

Drugs POTELIGEO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of relapsed or refractory mycosis fungoides (MF) or Sézary syndrome (SS) after at least one prior systemic therapy.

Age Restriction 18 years or older

Prescriber Restriction

Coverage Duration 3 Months

Drugs

pramipexole oral tablet extended release 24 hr 0.375 mg, 0.75 mg, 1.5 mg, 3 mg, 4.5 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documented diagnosis and treatment history.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Failure to pramipexole IR Drugs PREZCOBIX

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs PREZISTA ORAL SUSPENSION, PREZISTA ORAL TABLET 150 MG, 600 MG, 75 MG, 800 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs

PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Uncontrolled hypertension. Pure red cell aplasia that begins after ESA treatment.

Required Medical Information

Pre-treatment hemoglobin level less than 10 g/dL AND Patient has adequate iron stores prior to initiation of therapy defined as ferritin more than 100 mcg/L or serum transferrin saturation greater than 20% AND other causes of anemia such as iron deficiency, folate deficiency or B12 deficiency, hemolysis, gastrointestinal bleeding, other active or occult bleeding, or underlying hematologic disease (such as sickle cell anemia, thalassemia, and porphyria) have been ruled out AND Diagnosis of one of the following: A) Anemia due to chronic kidney disease (CKD) with or without hemodialysis, OR B) Anemia in patients with non-myeloid malignancies where anemia is due to the effect of concomitant myelosuppressive chemotherapy and two additional months of chemotherapy is anticipated, C) Treatment of anemic in a patient at high risk for perioperative blood loss from elective, noncardiac, nonvascular surgery to reduce the need for allogeneic blood transfusion, D) Anemia in zidovudine-treated HIV infection with serum erythropoietin levels 500 mUnits/mL or less and zidovudine doses 4,200 mg/week or less.

Age Restriction

Prescriber Restriction

CKD - prescribed by a nephrologist or hematologist. Non-myeloid malignancies - prescribed by an oncologist/hematologist. Surgery - Prescribed by a surgeon. HIV - Prescribed by an infectious disease specialist.

Coverage Duration

Initial: 3 months. Renewal: CKD-12 months, Non-myeloid cancers, HIV-4 months. Surgery-3 months

Other Criteria

For renewal of CKD (dialysis patients): Hb less than 11 g/dL or physician will decrease or interrupt dose. For renewal of CKD (non-dialysis patients): Hb less than 10 g/dL or physician will decrease or interrupt dose. For renewal of non-myeloid malignancies: Concurrent myelosuppressive chemotherapy and Hb is 12g/dL or less and there is measurable response after eight weeks (defined as an increase in Hb 1 g/dL or more or a reduction in red blood cell transfusion requirements). For renewal of zidovudine-treated HIV, Hb is 12g/dL or less AND Zidovudine dose remains 4,200 mg/week or less and there is a measurable response after eight weeks (defined as an increase in Hb or a reduction in RBC transfusion requirements or documented dose escalation [up to max of 300 units/kg/dose]).

Drugs PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG, PROMACTA ORAL TABLET

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Relapsed/refractory chronic immune (idiopathic) thrombocytopenic purpura (ITP) for greater than 6 months AND Baseline platelet count is less than 50,000/mcL AND Degree of thrombocytopenia and clinical condition increase the risk of bleeding AND Patient had an insufficient response, intolerance, contraindication to corticosteroids or immune globulin or inadequate response or contraindication to splenectomy, or B) Chronic hepatitis C and patient has thrombocytopenia defined as platelets less than 90,000/mcL for initiation (pre-treatment) of interferon therapy, or C) Severe aplastic anemia and patient has insufficient response to immunosupressive therapy.

Age Restriction

Prescriber Restriction

Coverage Duration 3 Months

PROTOPIC (tacrolimus topical)

Drugs

tacrolimus topical

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 16 years and older (0.1%) or 2 years and older (0.03%)

Prescriber Restriction

Coverage Duration 8 weeks

Drugs PULMOZYME

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of Cystic Fibrosis diagnosis

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Pulmozyme should be used in conjunction with standard therapies for CF. For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations).

Drugs QINLOCK

Covered Uses

Exclusion Criteria Must have previously received treatment with more than 3 kinase inhibitors, including imatinib

Required Medical Information Required diagnosis of Gastrointestinal stromal tumor, advanced.

Age Restriction For adults only

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs REGRANEX

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis

Age Restriction 16 years and older

Prescriber Restriction

Coverage Duration Plan year

Drugs RELENZA DISKHALER

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Per treatment (up to 28 days)

Drugs REMODULIN

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration 3 months

Drugs REPATHA PUSHTRONEX, REPATHA SURECLICK, REPATHA SYRINGE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Concurrent use of Repatha with Praluent (alirocumab injection for SC use), Juxtapid (lomitapide capsules) or Kynamro (mipomersen injection). The efficacy and safety of Praluent, Juxtapid and Kynamro in combination with Repatha have not been established.

Required Medical Information

Clinical documentation (e.g., chart notes, laboratory values) required for "initial" authorization includes: 1. Documentation of one of the following diagnoses: A) Homozygous familial hypercholesterolemia (HoFH), B) Heterozygous familial hypercholesterolemia (HeFH), OR C) Atherosclerotic cardiovascular disease (ASCVD), AND 2. Documentation of any one of the following: a) Patient has been receiving at least 12 consecutive weeks of high intensity statin therapy and will continue to receive high-intensity statin at maximally tolerated dose, b) Patient is unable to tolerate high-intensity statin, c) Patient has been receiving at least 12 consecutive weeks of moderate-intensity statin therapy and will continue to receive a moderate-intensity statin d) Patient has a documented labeled contraindication to all statins, e)Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment, f)Patient has undergone a trial of statin rechallenge with another low-intensity statin with documented reappearance of muscle symptoms (only 2 trials of a statin can be required prior to the approval of a PCSK9). Clinical documentation required for "reauthorization" includes: 1. Patient continues to receive statin at maximally tolerated dose (unless patient has documented inability to take statins), 2. Submission of medical records documenting LDL-C reduction while on Repatha therapy.

Age Restriction

HeFH,ASCVD: 18 years and older. HoFH: 13 years and older

Prescriber Restriction

Prescribed by or in consultation with or recommendation of a Cardiologist, Endocrinologist, or Lipid specialist.

Coverage Duration

Initial Authorization will be for 6 months. Reauthorization will be for 1 year.

Other Criteria

A) HoFH: Patient meets one of the following: a) Patient has genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9 or LDLRAP1 gene locus OR b) Patient has an untreated LDL-C level greater than 500 mg/dL (prior to treatment with antihyperlipidemic agents) OR c) Patient has a treated LDL-C level greater than or equal to 300 mg/dL (after treatment with antihyperlipidemic agents but prior to agents such as Repatha, Kynamro, or Juxtapid OR d) Patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas or xanthelasma B) HeFH: The patient has a low-density lipoprotein cholesterol (LDL-C level) greater than or equal to 160 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C level) greater than or equal to 160 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. C) ASCVD: The patient has a low-density lipoprotein cholesterol (LDL-C) greater than or equal to 70 mg/dL (after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy. ASCVD diagnosis is confirmed by one of the following: acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, peripheral arterial disease presumed to be of atherosclerotic origin. Examples of high-intensity statin therapy include atorvastatin 40-80 mg, Crestor (rosuvastatin) 20-40 mg. Examples of moderate-intensity statin therapy include atorvastatin 10-20 mg, Crestor (rosuvastatin) 20-40 mg. Examples of moderate-intensity statin therapy include atorvastatin 10-20 mg, Crestor (rosuvastatin) 5-10 mg, simvastatin greater than or equal to 20 mg, pravastatin gre

Drugs RETEVMO ORAL CAPSULE 40 MG, 80 MG

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following, and treatment in the following populations:A) Non-small cell lung cancer, metastatic, RET fusion-positive: Treatment of metastatic RET fusion-positive non-small cell lung cancer (NSCLC) in adults. B) Thyroid cancer, medullary, RET-mutant: Treatment of advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in adults and pediatric patients 12 years of age or older who require systemic therapy. OR C)Thyroid cancer, RET fusion-positive: Treatment of advanced or metastatic RET fusion-positive thyroid cancer in adults and pediatric patients 12 years of age or older who are refractory to radioactive iodine (if radioactive iodine is appropriate).

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs RETROVIR INTRAVENOUS

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs REVLIMID

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria Pregnancy (category X)

Required Medical Information

Diagnosis of one of the following: A) Multiple myeloma used in combination with dexamethasone, or B) Diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, or C) Mantle cell lymphoma and patient's disease has relapsed or progressed after trying at least two prior therapies (Velcade and one of the following: bendamustine, cladribine, fludarabine, rituximab) AND patient is enrolled in the Revlimid REMS Program.

Age Restriction

18 years and older

Prescriber Restriction

Hematologist/oncologist. Registered in Revlimid REMS.

Coverage Duration 3 Months

Drugs REYATAZ ORAL CAPSULE 150 MG, 200 MG, 300 MG, REYATAZ ORAL POWDER IN PACKET

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs REYVOW

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used for the acute treatment of migraine with or without aura in adults AND the member has tried and failed three alternatives two of which were triptans, unless contraindicated.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Other Criteria

Reauthorization requires documentation of medication efficacy.

RIBAVIRIN

Drugs

ribavirin oral capsule, ribavirin oral tablet 200 mg

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Per treatment (up to 48 weeks)

Drugs RISPERDAL CONSTA

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment. Documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Must be unable or unwilling to tolerate oral medications. Drugs SABRIL ORAL TABLET, vigabatrin oral powder in packet

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis of refractory complex partial seizures (CPS) or infantile spasms (IS). Previously tried and failed two medications for the diagnosis of refractory complex partial seizures including carbamazepine, ethotoin, felbamate, fosphenytoin, gabapentin, lacosamide, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, pregabalin, primidone, tiagabine, topiramate, valproic acid, divalproex sodium, zonisamide.

Age Restriction

10 years and older for CPS diagnosis. Children aged 1 month to 2 years old for IS.

Prescriber Restriction

Neurologist registered with the Sabril REMS program

Coverage Duration Plan year

Drugs SAMSCA

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Samsca not approved as an intervention to raise serum sodium urgently to prevent or to treat serious neurological symptoms. Samsca cannot be initiated or re-initiated outside of a hospital setting.

Required Medical Information

Serum sodium levels. Initial therapy for hyponatremia (hypervolemic and euvolemic): 1. Diagnosis of significant hyponatremia (euvolemic or hypervolemic), AND 2. Treatment has been initiated or re-initiated in a hospital setting prior to discharge. Reauthorization for hypervolemic and euvolemic hyponatremia: 1. Documentation of clinical benefit, AND 2. Treatment hasbeen initiated or re-initiated or re-initiated in a hospital setting prior to discharge.

Age Restriction

Prescriber Restriction

Coverage Duration 1 month

Other Criteria Documentation of trial and failure of fluid restriction required.

Drugs SANDOSTATIN LAR DEPOT INTRAMUSCULAR SUSPENSION, EXTENDED REL RECON

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses, or B) Metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes, or C) Vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Acromegaly: Documentation of inadequate response to surgery and/or radiotherapy, or documentation that patient is not a candidate for surgery and/or radiotherapy. Reauthorization will require statement indicating growth hormone (GH) levels are stabilized at less than 5.0ng/mL and IGF-1 levels are normalized (male less than 1.9U/mL or female less than 2.2 U/mL) as matched by age and gender, or the patient has a documented clinical response defined by a reduction of tumor mass, a reduction in the signs and symptoms of acromegaly, or an improvement in significant comorbidities.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

Drugs SAPHRIS, SECUADO

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and treatment history

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Member needs to have documented trial of any of the two following drugs: aripiprazole, clozapine, olanzapine, risperidone, quetiapine, ziprasidone

Drugs SELZENTRY ORAL TABLET 150 MG, 25 MG, 300 MG, 75 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented lab assay verifying HIV is CCR5-tropic positive strain.

SEROQUEL XR (Quetiapine)

Drugs

quetiapine oral tablet extended release 24 hr, SEROQUEL XR ORAL TABLET EXTENDED RELEASE 24 HR

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis and treatment history

Age Restriction 10 years and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Documentation of reason why quetiapine IR cannot be used.

Drugs SIGNIFOR LAR

Covered Uses

All FDA-approved indications not otherwise excluded from Part D. All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Documented diagnosis of acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option

Age Restriction 18 years of age and older

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

For renewal, patient's growth hormone level or insulin-like growth factor 1 (IGF-1) level for age and gender has normalized/improved.

SILDENAFIL (REVATIO)

Drugs

sildenafil (pulm.hypertension) oral tablet

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Should not be used in combination with organic nitrates. This product is only indicated for Pulmonary Hypertension and is not to be used for Erectile Dysfunction.

Required Medical Information

Statement of FDA approved diagnosis of pulmonary arterial hypertension

Age Restriction 18 years and older

Prescriber Restriction Cardiologist or Pulmonologist

Coverage Duration Plan year

Drugs SOFOSBUVIR-VELPATASVIR

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Provider must submit medical records documenting the following: 1) medical diagnosis of Chronic Hepatitis C with labs documenting genotype and subtype, AND 2) medical records documenting viral load taken within 6 months of beginning therapy

Age Restriction

18 years of age and older

Prescriber Restriction

Prescribed by, or in consultation with, a gastroenterologist, hepatologist, or infectious disease physician.

Coverage Duration

12 weeks or as defined by current AASLD/IDSA guidance

Other Criteria

Criteria and coverage durations will be applied consistent with current AASLD/IDSA guidance.

Drugs SOMATULINE DEPOT

Covered Uses

All FDA-approved indications not otherwise exluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of A) acromegaly AND Inadequate response to surgery and/or radiation therapy or patient cannot be treated with surgery and/or radiotherapy, or B) unresectable, well or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs).

Age Restriction

18 years of age and older

Prescriber Restriction

Coverage Duration 3 months initial. Continuation 6 months if no progression

Other Criteria

For renewal, patient's IGF-1 levels has normalized or improved.

SOMAVERT (Pegvisomant)

Drugs

Somavert subcutaneous recon soln 10 mg, SOMAVERT SUBCUTANEOUS RECON SOLN 15 MG, 20 MG, 25 MG, 30 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Plan year

Drugs SPRYCEL

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, or B) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML AND failure, resistance, or intolerance to imatinib, or C) Philadelphia chromosome-positive (Ph+) acute lymphoblastic leukemia (ALL) with resistance or intolerance to imatinib.

Age Restriction 18 years or older

Prescriber Restriction Prescriber must be an oncologist.

Coverage Duration Plan year

Stavudine (Zerit)

Drugs

stavudine oral capsule

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs STRIBILD

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs SUNOSI

Covered Uses

Exclusion Criteria

Required Medical Information

Clinical documentation of narcolepsy or obstructive sleep apnea AND failed at least TWO alternatives (e.g. methylphenidate, dextroamphetamine, modafinil and armodafinil).

Age Restriction

Prescriber Restriction

Coverage Duration Through the plan year

Drugs SUSTIVA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs SUTENT

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Combination use with other kinase inhibitors (for example, sorafenib, etc).

Required Medical Information

Diagnosis of one of the following: A) Advanced/metastatic renal cell carcinoma, or B) Gastrointestinal stromal tumors after disease progression on or intolerance to Gleevec, or C) Progressive, well-differentiated pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease.

Age Restriction

Patient must be at least 18 years of age.

Prescriber Restriction Must be prescribed by oncologist

Coverage Duration

3 months initial, then renewable in 6 month increments

Drugs SYNERA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs TABLOID

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration

Through chemotherapy remission inducation and consolidation treatment

Drugs TAFINLAR

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of unresectable or metastatic melanoma along with BRAF V600E or BRAF V600K mutation status as detected by a US Food and Drug Administration-approved test.

Age Restriction 18 years or older

Prescriber Restriction Oncologist

Coverage Duration 3 Months

Other Criteria

For authorization, please submit to EviCore at evicore.com or call at 877-825-7722.

Drugs TAKHZYRO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that medication is being used to prevent attacks of hereditary angioedema (HAE).

Age Restriction 12 years of age or older

Prescriber Restriction

Coverage Duration 3 Months

Drugs TALZENNA ORAL CAPSULE 0.25 MG, 1 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm) human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs ERLOTINIB, TARCEVA

Covered Uses

All FDA approved indications not otherwise excluded by Health Plan. First line for Non-Small Cell Lung Cancer (NSCLC).

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Locally advanced, unresectable, or metastatic pancreatic cancer and Tarceva will be used in combination with gemcitabine, or B) First-line treatment of metastatic non-small cell lung cancer in which tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an Food and Drug Administration (FDA)-approved test, or C) Maintenance treatment of locally advanced or metastatic non-small cell lung cancer when disease has not progressed after 4 cycles of platinum-based first-line chemotherapy, or D) Treatment of locally advanced or metastatic non-small cell lung cancer after failure of at least 1 prior chemotherapy regimen.

Age Restriction

18 years or older

Prescriber Restriction Prescriber must be an oncologist.

Coverage Duration Plan year

TARGRETIN (Bexarotene)

Drugs

bexarotene, TARGRETIN TOPICAL

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of diagnosis and treatment history.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Drugs TASIGNA ORAL CAPSULE 150 MG, 200 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Long QT syndrome. Uncorrected hypokalemia. Uncorrected hypomagnesemia. Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors.

Required Medical Information

Diagnosis of one of the following: A) Newly diagnosed adult patients with Philadephia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase, or B) Ph+ chronic or accelerated phase chronic myeloid leukemia (CML) in adult patients resistant to or intolerant to prior therapy that included imatinib.

Age Restriction 18 years and older

Prescriber Restriction Must be prescribed by Oncologist

Coverage Duration Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Drugs TAVALISSE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented Platelet count less than 30x10^9/L and member had an insufficient response to previous treatment (corticosteroids, immunoglobulins, splenectomy, and/or a thrombopoietin receptor agonists)

Age Restriction 18 years or older

Prescriber Restriction Hematologist

Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs TAZVERIK

Covered Uses

Exclusion Criteria

Required Medical Information

Documentation of metastatic or locally advanced epithelioid sarcoma that is not eligible for complete resection.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs TECFIDERA ORAL CAPSULE, DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Treatment of primary progressive MS is not covered. Combination therapy with a beta interferon product, Gilenya, Aubagio, Tecfidera, Tysabri or Copaxone is not covered.

Required Medical Information

Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondaryprogresive MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs

temozolomide

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration 3 Months

Drugs THALOMID

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Pregnancy (category X)

Required Medical Information

Diagnosis of one of the following: A) Multiple myeloma that is newly diagnosed and is receiving concurrent dexamethasone, or B) Acute treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) AND the medication will not be used as monotherapy if the member has moderate to severe neuritis, or C) Maintenance therapy for prevention and suppression of cutaneous manifestations of ENL recurrence.

Age Restriction

12 years of age and older

Prescriber Restriction

Candidates must follow Thalomid REMS program requirements. Provider and pharmacy must be registered with this program.

Coverage Duration 3 months

Drugs THYMOGLOBULIN

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration Up to 14 days.

Drugs TIBSOVO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of relapsed or refractory acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs TIGAN INTRAMUSCULAR

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction Use not recommended in children

Prescriber Restriction

Coverage Duration Per treatment

Drugs TIVICAY

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Tobramycin Inhalant Solution (TOBI)

Drugs

tobramycin in 0.225 % NaCl

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information Documented diagnosis of cystic fibrosis with Pseudomonas

Age Restriction

Prescriber Restriction

Coverage Duration Plan yer

Drugs

bosentan, TRACLEER

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Receiving concomitant cyclosporine A or glyburide therapy. Aminotransferase elevations are accompanied by signs or symptoms of liver dysfunction or injury or increases in bilirubin at least 2 times the upper limit of normal. For female patients, pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with reliable contraception.

Required Medical Information

Diagnosis of pulmonary arterial hypertension (PAH) WHO Group I AND New York Heart Association (NYHA) Functional Class II-IV

Age Restriction Greater than 12 years of age

Prescriber Restriction

Available only to those enrolled in the Tracleer REMS Program. Prescription is written by or in consultation with a pulmonologist or cardiologist

Coverage Duration

3 Months

Other Criteria

Liver aminotransferases will be measured prior to initiation of treatment and then monthly.

Tranexamic acid (CYKLOKAPRON)

Drugs

tranexamic acid intravenous

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Documentation of hemophilia diagnosis as appropriate

Age Restriction

Prescriber Restriction

Coverage Duration 8 days

Drugs

ALTRENO, tretinoin (antineoplastic)

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Patients diagnosed with acne vulgaris after trying and failing at least 1 preferred alternatives (such as generic acne products - erythromycin/benzoyl peroxide, clindamycin, etc) or other non-cosmetic diagnosis.

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Drugs TRIKAFTA

Covered Uses

Exclusion Criteria Use in combination with other CFTR modulator (Orkambi, Kalydeco, or Symdeko)

Required Medical Information

Documentation of Cystic Fibrosis AND confirmation of presence of at least one F508del mutation in CFTR gene through genetic testing.

Age Restriction 12 years of age or older

Prescriber Restriction Prescribed by pulmonologist or a physician who specializes in the treatment of Cystic fibrosis

Coverage Duration Through end of benefit year

Drugs TRIUMEQ

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

TRIZIVIR (abacavir/Lamivudine/Zidovudine)

Drugs

abacavir-lamivudine-zidovudine

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented negative HLA-B*5701 screening. Drugs TRUVADA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs TUKYSA

Covered Uses

Exclusion Criteria

Required Medical Information

Used for medical treatment of Breast cancer, human epidermal growth factor receptor 2 positive, advanced unresectable or metastatic:advanced unresectable or metastatic human epidermal growth factor receptor 2 (HER2)-positive breast cancer (in combination with trastuzumab and capecitabine) in adults with or without brain metastases who have received 1 or more prior anti-HER2-based regimens in the metastatic setting.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs TURALIO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 months

Drugs TYKERB

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND 1. the medication will be used in combination with Xeloda in a patient with advanced or metastatic disease and the patient has received prior therapy including an anthracycline, a taxane, and trastuzumab, or 2) The medication will be used in combination with Femara for the treatment of a postmenopausal woman with hormone receptor-positive metastatic disease for whom hormonal therapy is indicated.

Age Restriction

18 years or older

Prescriber Restriction

Must be prescribed by Oncologist and Oncologist must monitor treatment

Coverage Duration

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Drugs TYMLOS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Osteoporosis in a postmenopausal female AND one or more of the following: 1) History of osteoporotic fracture, or 2) Documented trial and failure of bisphosphonate or 3) Documented contraindication or intolerance to bisphosphonate therapy. Patient has not received more than 2 years of therapy with Tymlos.

Age Restriction

Prescriber Restriction

Coverage Duration

Initial: 1 year. Renewal: 1 year not to exceed 2 years of total

Drugs UBRELVY

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used for the acute treatment of migraine with or without aura in adults AND the member has tried and failed three alternatives two of which were triptans, unless contraindicated.

Age Restriction

Prescriber Restriction

Coverage Duration

Through end of benefit year

Other Criteria

Reauthorization requires documentation of medication efficacy.

Drugs VALCYTE ORAL RECON SOLN, valganciclovir oral tablet

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 4 months and older.

Prescriber Restriction

Coverage Duration Plan year

Drugs

vancomycin oral capsule

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of A) Clostridium difficile-associated diarrhea, AND Stool culture report within the previous 30 days indicating positive C. difficile toxin, AND documented trial and failure or contraindication to preferred agent, metronidazole, or B) Staphylococcus aureus (including methicillin-resistant strains)enterocolitis

Age Restriction

Prescriber Restriction

Coverage Duration

14 days, Patients with multiple relapses: 6 weeks

Drugs

calcitriol topical

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Must have tried and failed at least 2 topical steroids (at least one mid potency and at least one high potency)

Drugs VENTAVIS

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of Pulmonary Arterial Hypertension (PAH) AND has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class III or IV. Trial and failure of Revatio or Adcirca.

Age Restriction 18 years or older

To years of older

Prescriber Restriction

Must be prescribed by a cardiologist or pulmonologist

Coverage Duration 3 months.

Drugs VERZENIO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented diagnosis of one of the following: A)hormone receptor -positive, human epidermal growth factor receptor 2 - negative advanced or metastatic breast cancer and must ne used in combination with fulvestrant unless there is disease progression following endocrine therapy and prior chemotherapy in the metastatic setting.

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 Months

Drugs VIBERZI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinically diagnosed with irritable bowel syndrome with diarrhea supported by documentation from the patient's medical records AND Other GI medical conditions that could explain the symptoms have been ruled out AND Failed conventional non-pharmacological therapies including (Dietary changes, stress reduction, or behavioral changes)AND Failed conventional pharmacological therapies including: Antidiarrheals, Antidepressants, and Antispasmodics AND Must have tried and failed rifaximin

Age Restriction

Prescriber Restriction

Coverage Duration

Initial Authorization will be for 3 months. Reauthorization will be for 1 year.

Drugs VIMPAT ORAL SOLUTION, VIMPAT ORAL TABLET, VIMPAT ORAL TABLETS, DOSE PACK

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction 17 years of age or older

Prescriber Restriction

Coverage Duration Through the benefit year

Other Criteria Max dose 400mg/day

Drugs VIRACEPT ORAL TABLET

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

VIRAMUNE (Nevirapine)

Drugs nevirapine

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs VIREAD

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

VITEKTA (Elvitegravir)

Drugs tenofovir disoproxil fumarate

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Drugs VITRAKVI ORAL CAPSULE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinical documentation of unresectable or metastatic solid tumors that have a neurotrophic receptor tyrosine kinase (NTRK) gene fusion and have no satisfactory alternative treatments or that have progressed following treatment.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs VIZIMPRO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that medication will be used for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs

voriconazole oral

Covered Uses

All medically-accepted indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Documentation of invasive aspergillosis, bronchopulmonary aspergillosis, candidemia and disseminated candidiasis in skin, abdomen, kidney, bladder wall and wounds, esophageal candidiasis, and serious Candida infections, infections caused by the emerging pathogens Scedosporium sp. and Fusarium sp., or rare and refractory fungal infections should be provided. Preferred alternative for Candida: oral fluconazole

Age Restriction 12 years or older

Prescriber Restriction

Coverage Duration 6 months

Drugs VOTRIENT

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) Advanced/metastatic renal cell carcinoma, or B) Advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., doxorubicin, dacarbazine, ifosfamide, epirubicin, gemicitabine, docetaxel, or vinorelbine).

Age Restriction 18 years of age and older

Prescriber Restriction Oncologist

Coverage Duration Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Drugs VUMERITY

Covered Uses

Exclusion Criteria

Concurrent use with other MS disease modifying agents (including Aubagio, Avonex, Betaseron, Copaxone/Glatiramer/Glatopa, Extavia, Gilenya, Lemtrada, Mavenclad, Mayzent, Ocrevus, Plegridy, Rebif, Tecfidera and Tysabri) OR Member with moderate or severe renal impairment (creatinine clearance less than 60 mL/min.

Required Medical Information

Documentation that the medication will be used for the treatment of relapsing forms of multiple sclerosis (MS)(including clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease).

Age Restriction

Prescriber Restriction Neurologist

Coverage Duration Benefit Year

Drugs VYVANSE ORAL CAPSULE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Diagnosis of one of the following: A) ADHD and tried and failed two alternative medications FDA approved for the treatment of ADHD, or B) Moderate to severe Binge Eating Disorder AND the patient is receiving psychological counseling AND the patient must have tried and failed at least two antidepressant medications.

Age Restriction ADHD: Must be older than 6 years of age, BED: Must be 18 years of age or older

Prescriber Restriction

BED: The medication must be prescribed by a psychiatrist or a psychiatric specialist.

Coverage Duration

12 months

Drugs VYXEOS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Documented diagnosis of newly-diagnosed therapy-related acute myeloid leukemia (t-AML) or AML with myelodysplasiarelated changes (AML-MRC).

Required Medical Information

Age Restriction

Prescriber Restriction oncologist or hematologist

Coverage Duration 3 Months

Drugs WAKIX

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosis of one of Excessive daytime sleepiness in patients with narcolepsy AND 1. Submission of sleep study with narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants AND 3. has experienced inadequate response or intolerable side effects to two preferred products (modafinil, armodofinil, methylphenidate, or dextroamphetamine).

Age Restriction

Prescriber Restriction

Coverage Duration

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Other Criteria

For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy).

Drugs XALKORI

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Stated diagnosis of late-stage (locally advanced or metastatic), non-small cell lung cancers (NSCLC) with expression of the abnormal anaplastic lymphoma kinase (ALK) gene as detected by an FDA approved test.

Age Restriction 18 years and older

Prescriber Restriction Oncologist

Coverage Duration 3 months

Drugs XCOPRI MAINTENANCE PACK, XCOPRI TITRATION PACK

Covered Uses

Exclusion Criteria

Required Medical Information

Diagnosed with partial-onset seizures and treatment failure of at least two other formulary medications used in the treatment of partial onset seizures.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Xeljanz

Drugs XELJANZ ORAL TABLET 10 MG, 5 MG, XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented trial and failure of preferred TNF inhibitors (Enbrel and Humira)AND negative Tuberculin test prior to therapy AND patient is free of any clinically important active infections.

Age Restriction

Prescriber Restriction

Coverage Duration Through end of benefit year

Drugs XIFAXAN ORAL TABLET 200 MG, 550 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Allergy to rifamycin agents

Required Medical Information

Diagnosis of traveler's diarrhea and patient does not have fever or blood in the stool OR Diagnosis of hepatic encephalopathy and tried and failed lactulose therapy OR Diagnosis of irritable bowel syndrome with diarrhea (IBS-D) and failed at least TWO alternatives from TWO different classes such as anitdiarrheals (e.g. loperamide, diphenoxylateatropine), antispasmodics (e.g. dicyclomine), bile acid sequestrants (e.g. cholestryramine, colestipol).

Age Restriction

Traveler's diarrhea: 12 years of age or older, Hepatic encephalopathy and IBS-D: 18 years of age or older

Prescriber Restriction

Coverage Duration Traveler's diarrhea: 3 days, Hepatic encephalopathy: 6 months, IBS-D: 6 weeks

Drugs XOLAIR

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of A) moderate to severe persistent allergic asthma AND Evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. radioallergosorbent test) for a specific IgE or in vitro reactivity to a perennial aeroallergen AND Pretreatment serum IgE levels greater than 30 and less than 700 IU/mL AND Symptoms are not adequately controlled with at least ONE inhaled corticosteroid (ICS) plus long-acting beta2-agonist (LABA) (e.g. Advair, Symbicort) for at least 3 months unless patient is intolerant to such treatment or such treatment is contraindicated, or B) Treatment of chronic idiopathic urticaria in adults and adolescents 12 years and older who remain symptomatic despite H1 antihistamine treatment (i.e. loratidine, cetirizine, levocetirizine, fexofenadine, etc.).

Age Restriction

Patient must be 12 years of age or older

Prescriber Restriction

Initial drug order must be by an allergist/immunologist, dermatologist, or a pulmonologist

Coverage Duration 6 months

Drugs XOSPATA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Clinical documentation of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test.

Age Restriction

Prescriber Restriction Oncology

Coverage Duration 3 months

Drugs

XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK)

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation that the medication will be used in 1) combination with dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma (RRMM) who have received at least four prior therapies and whose disease is refractory to at least two proteasome inhibitors, at least two immunomodulatory agents, and an anti-CD38 monoclonal antibody. OR 2)monotherapy for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma

Age Restriction

Prescriber Restriction Oncologist

Coverage Duration 3 months

Drugs XTANDI

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of metastatic castration-resistant prostate cancer AND the patient has tried and had an inadequate response, contraindication or intolerance to Zytiga.

Age Restriction

Prescriber Restriction Oncologist or urologist

Coverage Duration 3 months

Other Criteria Must try and fail Zytiga first.

Drugs XYREM

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Contraindications: Patient is being treated with sedative hypnotic agents, other CNS depressants, or using alcohol. Patient has succinic semialdehyde dehydrogenase deficiency. Patient has a history of drug abuse.

Required Medical Information

Diagnosis of one of the following: A) Excessive daytime sleepiness in patients with narcolepsy AND 1. Submission of sleep study with narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants AND 3. has experienced inadequate response or intolerable side effects to two preferred products (modafinil, armodofinil, methylphenidate, or dextroamphetamine) AND 4. The requested dose does not exceed the FDA indicated maximum (9gm/night), or B) Cataplexy in patients with narcolepsy AND 1. Submission of sleep study showing narcolepsy diagnosis, AND 2. currently NOT taking any sedative hypnotics or other CNS depressants, AND 3. does not have sleep apnea, AND 4. The dose does not exceed the FDA indicated maximum (9gm/night).

Age Restriction

18 years of age and older

Prescriber Restriction

Neurologist

Coverage Duration

Initial: 3 months, Renewal: 6 months with documentation of continued benefit

Other Criteria

Patient and physician must adhere to all regulations of the Xyrem REMS Program. For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy).

Drugs YONSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documented diagnosis of metastatic castration resistant prostate cancer (CRPC)AND the medication is being used in combination with methylprednisolone AND medication not being used as dual therapy with another androgen receptor inhibitor.

Age Restriction 18 years of age or older

Prescriber Restriction

Coverage Duration 3 Months

Drugs ZAVESCA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Medical statement of approved diagnosis: mild to moderate type 1 Gaucher disease for whom enzyme replacement therapy is not a therapeutic option (e.g., due to constraints such as allergy, hypersensitivity, or poor venous access).

Age Restriction 18 years or older

Prescriber Restriction

Coverage Duration Plan year

Drugs ZEMAIRA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration 3 months.

Drugs ZEPOSIA, ZEPOSIA STARTER KIT, ZEPOSIA STARTER PACK

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Documentation of diagnosis of clinically-isolated syndrome multiple sclerosis OR relapsing multiple sclerosis, including relapsing-remitting disease or active secondary progressive disease AND the patient has had a trial AND a complete blood cell count and liver function test was completed, reviewed, and deemed appropriate for Zeposia treatment by the prescriber AND patient has had ECG to assess for preexisting cardiac conduction abnormalities prior to starting Zeposia AND inadequate response or intolerance to at least one alternative drug indicated for the treatment of multiple sclerosis.

Age Restriction 18 Years of age and older

Prescriber Restriction Neurologist

Coverage Duration Initial 3 months, Continuation of therapy: 12 months

ZIAGEN (Abacavir)

Drugs

abacavir oral tablet, ZIAGEN ORAL SOLUTION

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Documented negative HLA-B*5701 screening.

Zidovudine (RETROVIR)

Drugs zidovudine

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information Clinical documentation of FDA approved indication for treatment.

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Drugs

zoledronic acid-mannitol-water intravenous piggyback 5 mg/100 mL

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment. For treatment of hypercalcemia of malignancy, must have documentation of corrected total serum calcium greater than or equal to 12 mg/dL. For Paget's disease, must have symptomatic form of disease.

Age Restriction 18 years and older

Prescriber Restriction

Coverage Duration

Per treatment

Other Criteria

For Paget's disease, must have documented failure, intolerance or contraindication to oral agent: alendronate OR risedronate. For osteoporosis, must have documented failure, intolerance or contraindication to at least 2 oral bisphosphonates.

Drugs ZOLINZA

Covered Uses All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria

Treatment may be continued as long as there is no evidence of progressive disease or unacceptable toxicity.

Drugs

everolimus (immunosuppressive), ZORTRESS ORAL TABLET 0.25 MG, 0.5 MG, 0.75 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Clinical documentation of FDA approved indication for treatment.

Age Restriction 18 years and older.

Prescriber Restriction

Coverage Duration 6 months.

Other Criteria

Drugs ZYDELIG

Covered Uses All FDA-approved indications not otherwise excluded from Health Plan.

Exclusion Criteria

Required Medical Information

Age Restriction

Prescriber Restriction

Coverage Duration Plan year

Other Criteria For authorization, please submit to EviCore at evicore.com or call at 877-825-7722. Drugs ZYFLO

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Any pertinent clinical situation as defined by the product label that could affect patient safety and/or therapeutic efficacy (i.e. contraindications, warnings, precautions, adverse effects, renal or hepatic function, drug interactions, lab values, required prior or concomitant therapy, inappropriate dosing and/or duration, etc).

Required Medical Information

Documentation of diagnosis and treatment history.

Age Restriction 12 years and older.

Prescriber Restriction

Coverage Duration Plan year

Other Criteria Must have failed montelukast and zafirlukast.

Drugs

abiraterone, ZYTIGA ORAL TABLET 500 MG

Covered Uses

All FDA-approved indications not otherwise excluded by Health Plan.

Exclusion Criteria

Required Medical Information

Diagnosis of metastatic prostate cancer AND Patient has castration-resistant disease (defined by tumor growth/disease progression, risk in PsA levels, new metastases) AND Zytiga will be used in combination with prednisone.

Age Restriction

Prescriber Restriction

Coverage Duration 3 months

Other Criteria

Index

abacavir oral tablet	
abacavir-lamivudine	
abacavir-lamivudine-zidovudine	
abiraterone	
acitretin	
ACTIMMUNE	
adefovir	
ADEMPAS	6
AFINITOR ORAL TABLET 10 MC	
2.5 MG, 5 MG, 7.5 MG	
AIMOVIG AUTOINJECTOR	
AJOVY AUTOINJECTOR	
AJOVY SYRINGE	39
AKYNZEO (FOSNETUPITANT)	
INTRAVENOUS RECON SOLN	86
AKYNZEO (NETUPITANT)	86
ALECENSA	
ALINIA	
ALIQOPA	
alosetron	
ALTRENO	
ALUNBRIG ORAL TABLET 180	
MG, 30 MG, 90 MG	86
ALUNBRIG ORAL	
TABLETS, DOSE PACK	86
ambrisentan	
APTIOM	
APTIVUS	
APTIVUS (WITH VITAMIN E)	
ARCALYST	14
armodafinil	
ATRIPLA	
AUBAGIO	
AUSTEDO ORAL TABLET 12 M	G.
6 MG, 9 MG	18
AVONEX INTRAMUSCULAR PE	Ν
INJECTOR KIT	19
AVONEX INTRAMUSCULAR	
SYRINGE KIT	19
AYVAKIT	20
BALVERSA	21
BANZEL ORAL SUSPENSION	
BANZEL ORAL TABLET 200 MC	
400 MG	
BARACLUDE ORAL SOLUTION	
BAVENCIO	
BELSOMRA	25
BENLYSTA SUBCUTANEOUS	
AUTO-INJECTOR	
benznidazole	
BETASERON SUBCUTANEOUS	
КІТ	
bexarotene	
bosentan	. 245
BOSULIF ORAL TABLET 100	-
MG, 400 MG, 500 MG	86
BRAFTOVI ORAL CAPSULE 50	
MG, 75 MG	
BROVANA	33

BRUKINSA	34
BUPHENYL ORAL TABLET	35
buprenorphine HCl sublingual	36
buprenorphine-naloxone sublingual	.00
film 12-3 mg, 2-0.5 mg, 4-1 mg, 8-2	
mg	36
BUSULFEX	.37
BYNFEZIA	
CABOMETYX ORAL TABLET 20	00
	~ ~
MG, 40 MG, 60 MG	
calcitriol topical2	259
CALQUENCE	40
capecitabine	
CAPLYTA	31
CAPRELSA ORAL TABLET 100	
MG, 300 MG	42
caspofungin	
CAYSTON	
CESAMET	
cinacalcet	
CLOVIQUE	47
clozapine oral tablet, disintegrating	
COMETRIQ ORAL CAPSULE 100	.40
MG/DAY(80 MG X1-20 MG X1),	
140 MG/DAY(80 MG X1-20 MG	
X3), 60 MG/DAY (20 MG X 3/DAY).	86
COPAXONE SUBCUTANEOUS	
	40
SYRINGE 20 MG/ML, 40 MG/ML	
COPIKTRA	50
CORLANOR ORAL SOLUTION	51
COSENTYX	52
COSENTYX (2 SYRINGES)	
COSENTIX (2 STRINGES)	52
COSENTYX PEN	52
COSENTYX PEN (2 PENS)	.52
COTELLIC	86
CRIXIVAN ORAL CAPSULE 200	
MG	53
CYCLOSET	
CYSTAGON	
dalfampridine	11
DAURISMO ORAL TABLET 100	
MG, 25 MG	56
DESCOVY	
DIACOMIT ORAL CAPSULE	
DIAZEPAM INTENSOL	28
diazepam oral concentrate	28
diazoxide	
diclofenac sodium topical gel 3 %	60
didanosine oral capsule,delayed	
release(DR/EC) 250 mg, 400 mg	.61
DIFICID	
doxercalciferol oral	
	03
dronabinol oral capsule 10 mg, 2.5	
mg, 5 mg	
DUPIXENT PEN	65
DUPIXENT SYRINGE	-
SUBCUTANEOUS SYRINGE 200	
	05
MG/1.14 ML, 300 MG/2 ML	
EDECRIN	66

EDURANT	
ELIGARD	
ELIGARD (3 MONTH)	
ELIGARD (4 MONTH)	
ELIGARD (6 MONTH)	
ELZONRIS	
EMCYT	. 70
EMEND ORAL CAPSULE 40 MG,	
80 MG	.71
EMEND ORAL CAPSULE, DOSE	
PACK	
EMSAM	
EMTRIVA	.73
ENBREL SUBCUTANEOUS	75
RECON SOLN ENBREL SUBCUTANEOUS	. 75
	74
	. 74
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50	
	75
MG/ML (1 ML) ENBREL SURECLICK	. 75
ENDREL SURECLICK	
entecavir	
ENTRESTO	
EPCLUSA ORAL TABLET 400-	. //
100 MG	78
EPIDIOLEX	
EPIVIR HBV ORAL SOLUTION	
EPZICOM	
ERIVEDGE	
ERLEADA	
ERLOTINIB	
ERTACZO	
ESBRIET	
EVENITY	
everolimus (antineoplastic)	
everolimus (immunosuppressive)	
EVOTAZ	
EXJADE	
FANAPT ORAL TABLET	
FANAPT ORAL TABLETS, DOSE	
PACK	.89
FARYDAK	
FASENRA PEN	
fentanyl citrate buccal lozenge on a	
handle	. 91
FERRIPROX ORAL TABLET 500	
MG	
FINTEPLA	
FIRDAPSE	
fondaparinux subcutaneous syringe	;
10 mg/0.8 mL, 2.5 mg/0.5 mL, 5	
mg/0.4 mL, 7.5 mg/0.6 mL	
FORTEO	. 97
fosamprenavir	136
FUZEON SUBCUTANEOUS	
RECON SOLN	. 98
FYCOMPA ORAL SUSPENSION	. 99

FYCOMPA ORAL TABLET 10

MG, 12 MG, 2 MG, 4 MG, 6 MG, 8
MG
GAMMAGARD LIQUID 121
GENVOYA 100
GILENYA ORAL CAPSULE 0.5
MG 101
GILOTRIF
GLEOSTINE ORAL CAPSULE 10
MG, 100 MG, 40 MG86
HARVONI ORAL TABLET 90-400
MG 103
HERCEPTIN HYLECTA 104
HORIZANT
HUMIRA PEN
HUMIRA PEN CROHNS-UC-HS
START
HUMIRA PEN PSOR-UVEITS-
ADOL HS
HUMIRA SUBCUTANEOUS
SYRINGE KIT 10 MG/0.2 ML, 20
MG/0.4 ML, 40 MG/0.8 ML
HUMIRA(CF) PEDI CROHNS
STARTER
HUMIRA(CF) PEN CROHNS-UC-
HS
HUMIRA(CF) PEN PSOR-UV-
ADOL HS
HUMIRA(CF) PEN
SUBCUTANEOUS PEN
INJECTOR KIT 40 MG/0.4 ML, 80
NC/0 0 MI 100
MG/0.8 ML
HUMIRA(CF) SUBCUTANEOUS
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML HYCAMTIN
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML HYCAMTIN 86 IBRANCE 107 ICATIBANT
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG,
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 107 ICATIBANT 104 ICLUSIG ORAL TABLET 15 MG, 45 MG IDAMYCIN PFS
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG IDAMYCIN PFS 86 IDHIFA
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 1DAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 1DAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 10AMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 110 IMBRUVICA ORAL TABLET 110
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 107 ICATIBANT 104 ICLUSIG ORAL TABLET 15 MG, 45 MG 10AMYCIN PFS 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG IMBRUVICA ORAL TABLET 110 INBRIJA
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG 110 IMBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG 110 INBRUVICA ORAL TABLET 110 INBRIJA 111 111 INCRELEX 112 112
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 110 IMBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 109 IMBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113 INTELENCE ORAL TABLET 100
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 109 IMBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INREBIC 113 INTELENCE ORAL TABLET 100 109 MG, 200 MG, 25 MG 114
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INTELENCE ORAL TABLET 100 113 MG, 200 MG, 25 MG 114 INTRON A INJECTION 115
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113 INTELENCE ORAL TABLET 100 113 MG, 200 MG, 25 MG 114 INTRON A INJECTION 115 INVEGA SUSTENNA 116
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INTELENCE ORAL TABLET 100 113 MG, 200 MG, 25 MG 114 INTRON A INJECTION 115
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113 INTELENCE ORAL TABLET 100 MG, 200 MG, 25 MG MG, 200 MG, 25 MG 114 INTRON A INJECTION 115 INVEGA SUSTENNA 116 INVIRASE ORAL TABLET 117 IRESSA 86
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 IMBRUVICA ORAL TABLET 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113 INTELENCE ORAL TABLET 100 MG, 200 MG, 25 MG MG, 200 MG, 25 MG 114 INTRON A INJECTION 115 INVEGA SUSTENNA 116 INVIRASE ORAL TABLET 117 IRESSA 86 ISENTRESS HD 118
HUMIRA(CF) SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML 106 HYCAMTIN 86 IBRANCE 107 ICATIBANT 94 ICLUSIG ORAL TABLET 15 MG, 45 MG 45 MG 86 IDAMYCIN PFS 86 IDHIFA 108 imatinib oral tablet 100 mg, 400 mg 109 IMBRUVICA ORAL CAPSULE 140 MG, 70 MG MG, 70 MG 110 INBRIJA 111 INCRELEX 112 INLYTA 86 INREBIC 113 INTELENCE ORAL TABLET 100 MG, 200 MG, 25 MG MG, 200 MG, 25 MG 114 INTRON A INJECTION 115 INVEGA SUSTENNA 116 INVIRASE ORAL TABLET 117 IRESSA 86

ISENTRESS ORAL TABLET 118
ISENTRESS ORAL
TABLET, CHEWABLE 118
ISTURISA ORAL TABLET 1 MG,
10 MG, 5 MG
itraconazole oral capsule 120
JAKAFI ORAL TABLET 10 MG,
15 MG, 20 MG, 25 MG, 5 MG
JYNARQUE ORAL TABLETS,
SEQUENTIAL 45 MG (AM)/ 15 MG
(PM), 60 MG (AM)/ 30 MG (PM),
90 MG (AM)/ 30 MG (PM) 123
KALETRA
KALYDECO ORAL GRANULES
IN PACKET 50 MG, 75 MG
KALYDECO ORAL TABLET 125
KESIMPTA PEN
KISQALI
KISQALI FEMARA CO-PACK 127
KOSELUGO ORAL CAPSULE 10
MG, 25 MG
KUVAN ORAL
TABLET,SOLUBLE
lamivudine oral solution 130
lamivudine oral tablet 100 mg, 150
<i>mg, 300 mg</i> 130
LEDIPASVIR-SOFOSBUVIR 131
LENVIMA
LETAIRIS
LEUKINE INJECTION RECON
SOLN
leuprolide
LEXIVA
lidocaine topical adhesive
patch,medicated 5 % 137
lidocaine topical ointment
linezolid
LOKELMA
LONSURF86 LORBRENA ORAL TABLET 100
MG, 25 MG
LUPRON DEPOT
LUPRON DEPOT (3 MONTH) 141
LUPRON DEPOT (4 MONTH)141
LUPRON DEPOT (6 MONTH) 141
LUPRON DEPOT-PED141
LUPRON DEPOT-PED (3 MONTH)
LYNPARZA ORAL TABLET 142
MAKENA (PF) 143
MAKENA INTRAMUSCULAR OIL
250 MG/ML (1 ML) 143
MAVENCLAD (10 TABLET PACK)
MAVENCLAD (4 TABLET PACK) 144
MAVENCLAD (5 TABLET PACK) 144
MAVENCLAD (6 TABLET PACK) 144
MAVENCLAD (7 TABLET PACK) 144
MAVENCLAD (8 TABLET PACK) 144
MAVENCLAD (9 TABLET PACK) 144

MAVYRET	145
MEKINIST ORAL TABLET 0.5	~ ~
MG, 2 MG	
MEKTOVI	146
MENEST	
miglustat	
MOVANTIK	
NAGLAZYME	
NEULASTA	
NEUPOGEN	
nevirapine	
NEXAVAR	
NEXETOL	
NEXLIZET	
NINLARO	
NORPACE CR	
NORVIR ORAL CAPSULE	
NORVIR ORAL SOLUTION	
NOXAFIL	
NUBEQA	
NUCALA SUBCUTANEOUS	
AUTO-INJECTOR	164
NUCYNTA	
NUCYNTA ER	
NULOJIX	
NURTEC ODT	
NYMALIZE ORAL SYRINGE	168
octreotide acetate injection solution	7
	169
ODEFSEY	170
ODOMZO	
OMNIPOD DASH 5 PACK POD	
OMNIPOD DASH PDM KIT	171
OMNIPOD INSULIN	
MANAGEMENT	
OMNIPOD INSULIN REFILL	
OMNITROPE	102
ORFADIN ORAL CAPSULE 10	
MG, 2 MG, 5 MG	
ORKAMBI	
OTEZLA	
OTEZLA STARTER	.174
oxandrolone oral tablet 10 mg, 2.5	475
mg	
OXERVATE	170
paliperidone oral tablet extended	
release 24hr 1.5 mg, 3 mg, 6 mg, 9	
mg PALYNZIQ	
pamidronate	
PANRETIN	
paricalcitol oral	
PEGINTRON SUBCUTANEOUS	100
KIT 50 MCG/0.5 ML	181
PEMAZYRE	
penicillamine oral capsule	
penicillamine oral tablet	100

PENTAM	.184
PERSERIS	185
PICATO	186
PIQRAY ORAL TABLET 200	
MG/DAY (200 MG X 1), 250	
MG/DAY (200 MG X1-50 MG X1),	
300 MG/DAY (150 MG X 2)	187
PLEGRIDY	
POMALYST	100
POMALTST	
	190
pramipexole oral tablet extended	
release 24 hr 0.375 mg, 0.75 mg,	
1.5 mg, 3 mg, 4.5 mg	
PREZCOBIX	
PREZISTA ORAL SUSPENSION.	193
PREZISTA ORAL TABLET 150	
MG, 600 MG, 75 MG, 800 MG	
PROCRIT INJECTION SOLUTION	
10,000 UNIT/ML, 2,000 UNIT/ML,	
20,000 UNIT/ML, 3,000 UNIT/ML,	
4,000 UNIT/ML, 40,000 UNIT/ML.	194
PROMACTA ORAL POWDER IN	
PACKET 12.5 MG, 25 MG	.195
PROMACTA ORAL TABLET	195
PULMOZYME	197
QINLOCK	
quetiapine oral tablet extended	
release 24 hr	215
raloxifene	
REGRANEX	
RELENZA DISKHALER	
REMODULIN	
REPATHA PUSHTRONEX	
REPATHA SURECLICK	
REPATHA SYRINGE	
RETEVMO ORAL CAPSULE 40	202
MG, 80 MG	203
RETROVIR INTRAVENOUS	203
REVLIMID	
REVEIMID REYATAZ ORAL CAPSULE 150	205
	000
MG, 200 MG, 300 MG REYATAZ ORAL POWDER IN	.206
	~~~
PACKET	
REYVOW	
ribavirin oral capsule	
ribavirin oral tablet 200 mg	
RISPERDAL CONSTA	209
ROZLYTREK ORAL CAPSULE	
100 MG, 200 MG	
RUBRACA	
RUZURGI	
RYDAPT	86
SABRIL ORAL TABLET	
SAMSCA	
SANCUSO	86
SANDOSTATIN LAR DEPOT	
INTRAMUSCULAR	
SUSPENSION, EXTENDED REL	
RECON	212
SAPHRIS	

SECUADO	213
SELZENTRY ORAL TABLET 150	
MG, 25 MG, 300 MG, 75 MG	
SEROQUEL XR ORAL TABLET	
EXTENDED RELEASE 24 HR	045
SIGNIFOR LAR	
sildenafil (pulm.hypertension) oral	
tablet	
sodium phenylbutyrate oral tablet.	
SOFOSBUVIR-VELPATASVIR	
SOMATULINE DEPOT	
Somavert subcutaneous recon sol	In
10 mg	. 220
SOMAVERT SUBCUTANEOUS	
<b>RECON SOLN 15 MG, 20 MG, 25</b>	5
MG, 30 MG	. 220
SPRYCEL	. 221
stavudine oral capsule	. 222
STIVARGA	
STRIBILD	. 223
SUNOSI	
SUSTIVA	
SUTENT	
SYNERA	
TABLOID	
tacrolimus topical	
tadalafil (pulm. hypertension)	
tadalafil oral tablet 20 mg	
•	
	. 229
TAGRISSO ORAL TABLET 40	
	00
MG, 80 MG	
TAKHZYRO	. 230
TAKHZYRO TALZENNA ORAL CAPSULE 0.2	. 230 <b>25</b>
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG	. 230 <b>25</b> . 231
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen	. 230 2 <b>5</b> . 231 32
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA	.230 25 .231 32 232
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL	.230 25 .231 32 232
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150	. 230 25 . 231 32 232 . 233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG	.230 25 .231 32 .232 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE	.230 25 .231 32 .232 .233 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK	.230 25 .231 32 .232 .233 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE	.230 25 .231 32 .232 .233 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK	.230 25 .231 32 .232 .233 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL	.230 25 .231 32 .232 .233 .233
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED	.230 25 .231 232 .233 .233 .234 .235 .236
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120	.230 25 .231 32 .232 .233 .234 .235 .236
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide	.230 25 .231 32 .232 .233 .234 .235 .236
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG	.230 25 .231 32 .232 .233 .234 .235 .236 236
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 237 238 267 .239
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 237 238 267 .239 .240
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 237 237 238 267 .239 .240 241
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 .236 .237 .238 .267 .239 .240 .241 .242
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 .236 .237 .238 .267 .239 .240 .241 .242 .243
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl.	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 .236 .237 .238 .267 .239 .240 .241 .242 .243 .244
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl.	.230 <b>25</b> .231 .232 .233 .233 .234 .235 .236 .236 .237 .238 .267 .239 .240 .241 .242 .243 .244 86
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl	.230 <b>25</b> .231 .232 .233 .234 .235 .236 .236 .237 .238 .236 .237 .238 .240 .241 .242 .243 .244 86 .245
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl TOPOSAR TRACLEER tranexamic acid intravenous	.230 <b>25</b> .231 .232 .233 .234 .235 .236 .236 .237 .238 .267 .239 .240 .241 .242 .243 .244 86 .245 .246
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide temofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl TOPOSAR TRACLEER tranexamic acid intravenous tretinoin (antineoplastic)	.230 <b>25</b> .231 32 .232 .233 .235 .236 237 237 238 237 238 237 239 .240 241 .242 243 .244 86 .245 .246 247
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide tenofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl TOPOSAR TRACLEER tranexamic acid intravenous tretinoin (antineoplastic) TRIKAFTA	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 .236 .237 .238 .237 .238 .240 .241 .242 .241 .242 .243 .244 86 .245 .246 .247 .248
TAKHZYRO TALZENNA ORAL CAPSULE 0.2 MG, 1 MG tamoxifen TARCEVA TARGRETIN TOPICAL TASIGNA ORAL CAPSULE 150 MG, 200 MG TAVALISSE TAZVERIK TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG temozolomide temofovir disoproxil fumarate THALOMID THYMOGLOBULIN TIBSOVO TIGAN INTRAMUSCULAR TIVICAY tobramycin in 0.225 % NaCl TOPOSAR TRACLEER tranexamic acid intravenous tretinoin (antineoplastic)	.230 <b>25</b> .231 32 .232 .233 .234 .235 .236 .236 .237 .238 .236 .237 .238 .240 .241 .242 .243 .244 86 .245 .246 .245 .246 .247 .248 .249

TUKYSA	. 252
TURALIO	. 253
TYKERB	. 254
TYMLOS	. 255
UBRELVY	.256
VALCHLOR	86
VALCYTE ORAL RECON SOLN.	
valganciclovir oral tablet	
vancomycin oral capsule	
VARUBI	
VENCLEXTA	
VENCLEXTA STARTING PACK	
VENTAVIS	
VERZENIO	
VIBERZI	
vigabatrin oral powder in packet	
VIMPAT ORAL SOLUTION	
VIMPAT ORAL TABLET	. 263
VIMPAT ORAL TABLETS, DOSE	
PACK	.263
VIRACEPT ORAL TABLET	264
VIREAD	
VITRAKVI ORAL CAPSULE	.268
VIZIMPRO	.269
voriconazole oral	.270
VOTRIENT	
VUMERITY	
VYVANSE ORAL CAPSULE	
VYXEOS	
ωδκιχ	
	. 275
XALKORI	. 275 . 276
XALKORI XCOPRI MAINTENANCE PACK	. 275 . 276 . 277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK	.275 .276 .277 .277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG	.275 .276 .277 .277 .277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG	.275 .276 .277 .277 .277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET	.275 .276 .277 .277 .277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11	.275 .276 .277 .277 .277
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG	. 275 . 276 . 277 . 277 . 278
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO	. 275 . 276 . 277 . 277 . 278
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200	. 275 . 276 . 277 . 277 . 278 
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG	. 275 . 276 . 277 . 277 . 278 
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR	. 275 .276 .277 .277 .278 . 278 86 86 .279 .280
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA	. 275 .276 .277 .277 .278 . 278 86 86 .279 .280
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100	. 275 .276 .277 .277 .278 . 278 86 86 .279 .280
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40	. 275 .276 .277 .277 .278 . 278 86 86 .279 .280
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG	. 275 .276 .277 .277 . 278 . 278 . 278 . 278 . 278 . 278 . 280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40	. 275 .276 .277 .277 . 278 . 278 . 278 . 278 . 278 . 278 . 280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG	. 275 .276 .277 .277 . 278 . 278 . 278 . 278 . 278 . 278 . 280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60	. 275 .276 .277 .277 . 278 . 278 . 278 . 278 . 278 . 278 . 280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK),	.275 .276 .277 .277 .277 .278 86 .279 .280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG	. 275 .276 .277 .277 . 278 86 86 .279 .280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK),	.275 .276 .277 .277 .278 .278 .278 .278 .278 .280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO. XIFAXAN ORAL TABLET 200 MG, 550 MG. XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK)	.275 .276 .277 .277 .278 .278 .278 .278 .278 .280 .281
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO. XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR. XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM.	.275 .276 .277 .277 .277 .278 .278 .278 .278 .278
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM YONSA	.275 .276 .277 .277 .277 .278 .278 .278 .278 .278
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM YONSA ZAVESCA	.275 .276 .277 .277 .277 .278 .278 .278 .278 .278
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO XIFAXAN ORAL TABLET 200 MG, 550 MG XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK). XTANDI XYREM YONSA ZAVESCA ZEJULA	.275 .276 .277 .277 .277 .278 .278 .278 .278 .279 .280 .281 .281 .281 .282 .283 .284 .285 .286 86
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO. XIFAXAN ORAL TABLET 200 MG, 550 MG. XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM YONSA ZAVESCA ZEJULA ZELBORAF	.275 .276 .277 .277 .277 .278 .278 .278 .278 .280 .280 .281 .281 .281 .282 .283 .284 .285 .286 86 86 86
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO. XIFAXAN ORAL TABLET 200 MG, 550 MG. XOLAIR. XOSPATA. XOSPATA. XOSPATA. XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM YONSA. ZAVESCA ZEJULA. ZELBORAF ZEMAIRA.	.275 .276 .277 .277 .277 .278 .278 .278 .278 .280 .280 .281 .281 .281 .281 .282 .283 .284 .285 .284 .285 .286 86 86 86 86
XALKORI XCOPRI MAINTENANCE PACK XCOPRI TITRATION PACK XELJANZ ORAL TABLET 10 MG 5 MG XELJANZ XR ORAL TABLET EXTENDED RELEASE 24 HR 11 MG XERMELO. XIFAXAN ORAL TABLET 200 MG, 550 MG. XOLAIR XOSPATA XPOVIO ORAL TABLET 100 MG/WEEK (20 MG X 5), 40 MG/WEEK (20 MG X 2), 40MG TWICE WEEK (80 MG/WEEK), 60 MG/WEEK (20 MG X 3), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (20 MG X 4), 80MG TWICE WEEK (160 MG/WEEK) XTANDI XYREM YONSA ZAVESCA ZEJULA ZELBORAF	.275 .276 .277 .277 .277 .278 .278 .278 .278 .280 .280 .281 .281 .281 .282 .283 .284 .283 .284 .285 .286 86 86 86 86 86 86

ZEPOSIA STARTER PACK	.289
zoledronic acid-mannitol-water intravenous piggyback 5 mg/100	
<i>mL</i> <b>ZOLINZA</b>	
ZORTRESS ORAL TABLET 0.25	
MG, 0.5 MG, 0.75 MG	
ZYDELIG	
ZYKADIA ORAL TABLET	
ZYTIGA ORAL TABLET 500 MG	

This formulary was updated on 03/01/2019. For more recent information or other questions, please call Customer Service toll-free at 1.855.443.4735 (TTY/TDD relay: 1.800.955.8771) Monday through Friday from 8 a.m. to 6 p.m.

You must generally use network pharmacies to use your prescription drug benefit. The Formulary or pharmacy network may change at any time. You will receive notice when necessary.

Health First Commercial Plans, Inc. is doing business under the name of Health First Health Plans. Health First Health Plans does not discriminate on the basis of race, color, national origin, disability, age, sex, gender identity, sexual orientation, or health status in the administration of the plan, including enrollment and benefit determinations.



6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

# **Nondiscrimination Notice**

Health First Health Plans complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Health First Health Plans does not exclude people or treat them differently because of race, color, national origin, age, disability, or sex.

Health First Health Plans:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, accessible electronic formats)
- Provides free language services to people whose primary language is not English, such as:
  - Qualified interpreters
  - Information written in other languages

If you need these services, please contact our Civil Rights Coordinator.

If you believe that Health First Health Plans has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with: Civil Rights Coordinator, 6450 US Highway 1, Rockledge, FL 32955, 321-434-4521, 1-800-955- 8771 (TTY), Fax: 321-434-4362, <u>civilrightscoordinator@hf.org</u>. You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance our Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights electronically through the Office for Civil Rights Complaint Portal, available at <u>https://ocrportal.hhs.gov/ocr/portal/lobby.jsf</u>, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Avenue SW., Room 509F, HHH Building, Washington, DC 20201, 1-800-368-1019, 800-537-7697 (TDD).

Complaint forms are available at http://www.hhs.gov/ocr/office/file/index.html.

Health First Commercial Plans, Inc. is doing business under the name of Health First Health Plans. Health First Health Plans does not discriminate on the basis of race, color, national origin, disability, age, sex, gender identity, sexual orientation, or health status in the administration of the plan, including enrollment and benefit determinations.

36194_MPINFO324 (05/2019)



6450 U.S. Highway 1 Rockledge, FL 32955 myHFHP.org

# English:

If you, or someone you're helping, has questions about Health First Health Plans, you have the right to get help and information in your language at no cost. To talk to an interpreter, call 855-443-4735.

# Spanish:

En caso que usted, o alguien a quien usted ayude, tenga cualquier duda o pregunta acerca de Health First Health Plans, usted tiene derecho a obtener ayuda e información en su idioma sin costo alguno. Para hablar con un intérprete, llame al 855-443-4735.

# Haitian Creole:

Si oumenm oswa yon moun w ap ede gen kesyon konsènan Health First Health Plans, se dwa w pou resevwa asistans ak enfòmasyon nan lang ou pale a, san ou pa gen pou peye pou sa. Pou pale avèk yon entèprèt, rele nan 855-443-4735.

# Vietnamese:

Nếu Quý vị, hay người mà Quý vị đang giúp đỡ, có câu hỏi về Health First Health Plans thì Quý vị có quyền được trợ giúp và được biết thêm thông tin bằng ngôn ngữ của mình miễn phí. Để nói chuyện với thông dịch viên, xin gọi số 855-443-4735.

# **Portuguese:**

Você ou alguém que você estiver ajudando tem o direito de tirar dúvidas e obter informações sobre os Health First Health Plans no seu idioma e sem custos. Para falar com um tradutor, ligue para 855-443-4735.

# Chinese:

如果您,或是您正在協助的對象,有與 Health First Health Plans 相關的問題,您有權以您的母語免費取得幫助和資訊。請致電 855-443-4735 與翻譯員洽談。

# French:

Si vous, ou quelqu'un que vous êtes en train d'aider, a des questions à propos de Health First Health Plans, vous avez le droit d'obtenir de l'aide et l'information dans votre langue à aucun coût. Pour parler à un interprète, appelez 855-443-4735.

# Tagalog:

Kung ikaw, o ang iyong tinutulangan, ay may mga katanungan tungkol sa Health First Health Plans, may karapatan ka na humingi ng tulong at impormasyon sa iyong wika nang libre. Upang makausap ang isang tagasalin, tumawag sa 855-443-4735.

# **Russian:**

Если у вас или лица, которому вы помогаете, имеются вопросы по поводу Health First Health Plans, то вы имеете право на бесплатное получение помощи и информации на вашем языке. Для разговора с переводчиком позвоните по телефону 855-443-4735.

# Arabic:

إن كان لديك أو لدى شخص تساعده أسئلة بخصوص Health First Health Plans، فلديك الحق في الحصول على المساعدةوالمعلومات الضرورية بلغتك من دون أية تكلفة. للتحدث مع مترجم اتصل بالرقم 4735-443-855.

# Italian:

Se lei o qualcuno che sta aiutando avete domande su Health First Health Plans, ha il diritto di ottenere aiuto e informazioni nella sua lingua gratuitamente. Per parlare con un interprete, può chiamare il numero 855-443-4735.

# German:

Falls Sie oder jemand, dem Sie helfen, Fragen zum Health First Health Plans haben, haben Sie das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Um mit einem Dolmetscher zu sprechen, rufen Sie bitte die Nummer 855-443-4735 an.

# Korean:

만약 귀하 또는 귀하가 돕고 있는 어떤 사람이 Health First Health Plans 에 관해서 질문이 있다면 귀하는 그러한 도움과 정보를 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 그렇게 통역사와 얘기하기 위해서는 855-443-4735 로 전화하십시오.

# Polish:

Jeśli Ty lub osoba, której pomagasz, macie pytania na temat Health First Health Plans, macie Państwo prawo do bezpłatnego uzyskania informacji i pomocy w języku ojczystym. Aby porozmawiać z tłumaczem, prosimy zadzwonić pod numer 855-443-4735.

# Gujarati:

જો તમે અથવા તમે કોઇને મદદ કરી રહ્યા હો તેમાંથી કોઇને હૅલ્થ ફર્સ્ટ હૅલ્થ પ્લાન્સ વિશે પ્રશ્નો હોય તો તમને તમારી ભાષામાં વિના મૂલ્યે મદદ અને માહિતી મેળવવાનો અધિકાર છે. દુભાષિયા સાથે વાત કરવા માટે 855-443-4735 પર કૉલ કરો.

# Thai:

หากคุณหรือคนที่คุณกำลังช่วยเหลือมีคำถามเกี่ยวกับ Health First Health Plansคุณมีสิทธิที่จะได้รับความช่วยเหลือและข้อมูลในภาษาของคุณได้โดยไม่มีค่าใช้จ่าย หากต้องการพูดคุยกับล่าม โปรดโทร 855-443- 4735.

Health First Commercial Plans, Inc. is doing business under the name of Health First Health Plans. Health First Health Plans does not discriminate on the basis of race, color, national origin, disability, age, sex, gender identity, sexual orientation, or health status in the administration of the plan, including enrollment and benefit determinations.

36194-77150_MPINFO109 (08/2016)