



**Keystone 65 Rx HMO
Personal Choice 65SM Rx PPO
Select Option[®] Rx PDP
2024 Utilization Management
Criteria: Prior Authorization**

**PLEASE READ: THIS DOCUMENT CONTAINS INFORMATION
ABOUT SOME OF THE DRUGS WE COVER IN THIS PLAN**

This document was updated on 11/19/2024. For more recent information or other questions, please contact our Member Help Team: Keystone 65 Rx at 1-844-352-1699, Personal Choice 65 Rx at 1-888-879-4293, Select Option Rx at 1-888-678-7009 or, for TTY/TDD users, 711, seven days a week from 8 a.m. to 8 p.m. Please note that on weekends and holidays from April 1 through September 30, your call may be sent to voicemail. Or, visit www.ibxmedicare.com to use our *Formulary (List of Covered Drugs)* search tool or view a downloadable document.

When this document refers to “we,” “us,” or “our,” it means Independence Blue Cross. When it refers to “plan” or “our plan,” it means Keystone 65 Rx, Personal Choice 65 Rx, and Select Option Rx.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2025, and from time to time during the year. Independence Blue Cross offers Medicare Advantage plans with a Medicare contract. Enrollment in Independence Medicare Advantage plans depends on contract renewal.

Independence Blue Cross offers Medicare Advantage plans with a Medicare contract. Enrollment in Independence Medicare Advantage plans depends on contract renewal.

Keystone 65: Benefits underwritten by Keystone Health Plan East, a subsidiary of Independence Blue Cross — independent licensees of the Blue Cross and Blue Shield Association.

Personal Choice 65: Benefits underwritten by QCC Insurance Company, a subsidiary of Independence Blue Cross — independent licensees of the Blue Cross and Blue Shield Association.

Select Option: Benefits underwritten by QCC Insurance Company, a subsidiary of Independence Blue Cross — independent licensees of the Blue Cross and Blue Shield Association.

There may be restrictions to your drug coverage

Some covered drugs may have additional requirements or limits on coverage. We call this “utilization management.” These requirements and limits may include:

- **Prior Authorization (PA):** Our plan requires you or your physician to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don’t get approval, our plan may not cover the drug. Drugs that require prior authorization are listed in this document.
- **Step Therapy (ST):** In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. Drugs that require step therapy are listed in *2024 Utilization Management Criteria: Step Therapy*.
- **Quantity Limits (QL):** For certain drugs, our plan limits the amount of the drug that our plan will cover. Drugs that have quantity limits are listed in the *Keystone 65 Rx, Personal Choice 65 Rx, and Select Option Rx Formulary (List of Covered Drugs)*.

You can find out if your drug has any additional requirements or limits by looking in your plan *Formulary (List of Covered Drugs)*. You can also get more information about the restrictions applied to specific covered drugs by visiting www.ibxmedicare.com.

You can ask our plan to make an exception to these restrictions or limits, or for a list of other similar drugs that may treat your health condition. Your *Formulary (List of Covered Drugs)* and *Evidence of Coverage* will have more information about the exception request process.

How to use this document

This document, along with *2024 Utilization Management Criteria: Step Therapy*, is intended to be used with your *Formulary (List of Covered Drugs)*. If your prescription drug has the note “PA” in the “Requirements” column of the *Keystone 65 Rx, Personal Choice 65 Rx, and Select Option Rx Formulary (List of Covered Drugs)*, you can find more information on the restriction(s) in this document.

Locate your drug in the index on page 303. The restriction information includes:

- **Prior Authorization**
 - Covered uses
 - Exclusion criteria
 - Required medical information
 - Age restrictions
 - Prescriber restrictions
 - Coverage duration
 - Other criteria

Be sure to read all the information listed for your affected drug. If you have any questions or need assistance with the information contained in this document, please call our Member Help Team: Keystone 65 Rx at 1-844-352-1699, Personal Choice 65 Rx at 1-888-879-4293, or Select Option Rx at 1-888-678-7009 or, for TTY/TDD users, 711.

ABILIFY MYCITE

Products Affected

- ABILIFY MYCITE MAINTENANCE KIT ORAL THERAPY PACK 10 MG
TABLET THERAPY PACK 15 MG, 2 MG, 20 MG,
30 MG, 5 MG
- ABILIFY MYCITE STARTER KIT ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Schizophrenia: (1) Attestation that tracking ingestion of the medication is medically necessary. Bipolar 1 Disorder (BP): (1) Attestation that tracking ingestion of the medication is medically necessary. Adjunctive Treatment for Major Depressive Disorder (MDD): (1) Attestation that medication will be used as adjunct therapy. (2) Attestation that tracking ingestion of the medication is medically necessary.
Age Restrictions	(Schizophrenia, BP, MDD): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ABUSE DETERRENT OPIOID

Products Affected

- *hydrocodone bitartrate er oral capsule extended release 12 hour*
- *hydrocodone bitartrate er oral tablet er 24 hour abuse-deterrent*
- HYSINGLA ER
- NUCYNTA ER
- OXYCONTIN ORAL TABLET ER 12 HOUR ABUSE-DETERRENT
- ROXYBOND ORAL TABLET ABUSE-DETERRENT 15 MG, 5 MG
- XTAMPZA ER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of one of the following: (1) Prior use of TWO generic opioids OR (2) BOTH of the following: (a) there is a history of or a potential for drug abuse among the individual or a member of the individual's household AND (b) documentation of current patient-prescriber opioid treatment agreement
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ACTEMRA SQ

Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS
- TYENNE SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	(PJIA, SJIA, RA, GCA, SSc-ILD): Concurrent therapy with any other biologic disease modifying antirheumatic drug (DMARD), e.g. tumor necrosis factor antagonists.
Required Medical Information	Part D is medically necessary when there is documentation of the following: Polyarticular Juvenile Idiopathic Arthritis (PJIA): (1) Diagnosis of PJIA. (2) Inadequate response or inability to tolerate TWO of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) Xeljanz, (d) Orencia, (e) Cyltezo, (f) Yuflyma, (g) Rinvoq/Rinvoq LQ OR documentation demonstrating that a trial may be inappropriate. Systemic Juvenile Idiopathic Arthritis (SJIA): (1) Diagnosis of SJIA. (2) Inadequate response or inability to tolerate ONE of the following: (a) Non-steroidal anti-inflammatory drug (NSAID), (b) Systemic glucocorticoid, (c) Methotrexate. Moderate to severe rheumatoid arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) Cyltezo, (g) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Giant Cell Arteritis (GCA): (1) Diagnosis of GCA. (2) Documentation of inadequate response/inability to tolerate oral corticosteroids. Systemic Sclerosis (SSc-ILD) (1): Diagnosis of SSc-ILD (2) Documentation of inadequate response/inability to tolerate a 3-month trial of ONE of the following: (a) mycophenolate mofetil, (b) cyclophosphamide, OR (c) azathioprine (3) Diagnosis of SSc-ILD confirmed by a high-resolution CT scan or biopsy
Age Restrictions	(PJIA, SJIA): Member is 2 years of age or older (RA, GCA, SSc-ILD): Member is 18 years of age or older
Prescriber Restrictions	(PJIA, SJIA, RA, GCA): Prescribed by or in consultation with a Rheumatologist (SSc-ILD) -Prescribed by or in consultation with a rheumatologist or pulmonologist
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ACTHAR HP

Products Affected

- ACTHAR
- ACTHAR GEL
- CORTROPHIN

PA Criteria	Criteria Details
Exclusion Criteria	(All Indications): Scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction, sensitivity to proteins or porcine origin, or where congenital infections are suspected in infants. OR Administration of live or live attenuated vaccines in members receiving immunosuppressive doses of H.P. Acthar Gel.
Required Medical Information	Part D is medically necessary when ONE of the following is present: (1) Infantile Spasms (IS): (A) Diagnosis of IS. (2) Acute Exacerbation of Multiple Sclerosis (AEMS): (A) Diagnosis of an AEMS, (B) Currently receiving maintenance treatment for MS (e.g. Avonex, Betaseron, Copaxone, Tecfidera, etc.), (C) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (3) Acute Exacerbation of Psoriatic Arthritis (AEPsA): (A) Diagnosis of an AEPsA, (B) Currently receiving a DMARD, (C) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (4) Acute Exacerbation of Rheumatoid Arthritis (AERA): (A) Diagnosis of an AERA, (B) Currently receiving a DMARD, (C) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (5) Acute Exacerbation of Juvenile Rheumatoid Arthritis (AEJRA): (A) Diagnosis of an AEJRA, (B) Currently receiving a DMARD, (C) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (6) Acute Exacerbation of Ankylosing Spondylitis (AEAS): (A) Diagnosis of an AEAS, (B) Currently receiving a DMARD, (C) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (7) Nephrotic Syndrome (NS): (A) Diagnosis of NS, (B) Proteinuria greater than 3.5g/ 24 hours, (C) serum albumin less than 3 mg/dL, (D) Peripheral edema. (F) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (8) Systemic Lupus Erythematosus (SLE): (A) Diagnosis of SLE, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone).
Age Restrictions	(IS): Member is younger than 2 years of age. (MS): Member is 18 years of age and older. (All Other Indications): Member is 2 years of age and older

PA Criteria	Criteria Details
Prescriber Restrictions	(IS): Prescribed by or in consultation with an neurologist or neonatologist. (All Other Indications): Prescribed by or in consultation with a neurologist, rheumatologist, nephrologist, pulmonologist, ophthalmologist, dermatologist, allergist, immunologist.
Coverage Duration	(IS): 1 year (All Other Indications): 1 month
Other Criteria	Subject to Part B vs Part D review. (9) Systemic Dermatomyositis (SDM): (A) Diagnosis of SDM (polymyositis), (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (10) Severe Erythema Multiforme (SEM): (A) Diagnosis of SEM, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (11) Stevens-Johnson Syndrome (SJS): (A) Diagnosis of SJS, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (12) Serum Sickness (SS): (A) Diagnosis of SS, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (13) Inflammatory Ophthalmic Disease (IOD): (A) Diagnosis of IOD, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (14) Symptomatic Sarcoidosis (SSD): (A) Diagnosis of SSD, (B) Inadequate response or inability to tolerate two systemic steroids (e.g. prednisone, methylprednisolone). (ALL INDICATIONS): Dosing is in accordance with the United States Food and Drug Administration (FDA) approved labeling
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ACUTE HAE AGENTS

Products Affected

- BERINERT
- FIRAZYR SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE
- *icatibant acetate subcutaneous solution prefilled*
- *syringe*
- RUCONEST
- SAJAZIR SUBCUTANEOUS SOLUTION PREFILLED
SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when the following inclusion criteria is met: Hereditary Angioedema (HAE): (1) Used for the treatment of acute abdominal, facial or laryngeal attacks of HAE.
Age Restrictions	
Prescriber Restrictions	(HAE): Prescribed by or in consultation with an allergist or immunologist
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ACUTE SEIZURE ACTIVITY AGENTS

Products Affected

- LIBERVANT
- NAYZILAM
- VALTOCO 10 MG DOSE
- VALTOCO 15 MG DOSE
- VALTOCO 20 MG DOSE
- VALTOCO 5 MG DOSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Prescriber is a neurologist/epilepsy specialist
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ADALIMUMAB NON-PREFERRED PRODUCTS

Products Affected

- ABRILADA (1 PEN)
- ABRILADA (2 SYRINGE)
- *adalimumab-aacf (2 pen)*
- *adalimumab-aaty (1 pen) subcutaneous auto-injector kit 80 mg/0.8ml*
- *adalimumab-aaty (2 pen)*
- *adalimumab-aaty (2 syringe)*
- *adalimumab-adaz subcutaneous solution auto-injector*
- *adalimumab-adaz subcutaneous solution prefilled syringe 40 mg/0.4ml*
- *adalimumab-adbm (2 pen)*
- *adalimumab-adbm (2 syringe)*
- *adalimumab-adbm(cd/uc/hs strt)*
- *adalimumab-adbm(ps/uv starter)*
- *adalimumab-fkjp (2 pen)*
- *adalimumab-fkjp (2 syringe)*
- *adalimumab-ryvk (2 pen)*
- *adalimumab-ryvk (2 syringe)*
- AMJEVITA SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- AMJEVITA SUBCUTANEOUS SOLUTION
- PREFILLED SYRINGE 40 MG/0.4ML, 40 MG/0.8ML
- AMJEVITA-PED 10KG TO <15KG
- AMJEVITA-PED 15KG TO <30KG
- HADLIMA
- HADLIMA PUSHTOUCH
- HULIO (2 PEN)
- HULIO (2 SYRINGE)
- HYRIMOZ
- HYRIMOZ-CROHNS/UC STARTER
- HYRIMOZ-PED<40KG CROHN STARTER
- HYRIMOZ-PED>/=40KG CROHN START
- HYRIMOZ-PLAQ PSOR/UVEIT START
- IDACIO (2 PEN)
- IDACIO (2 SYRINGE)
- IDACIO-CROHNS/UC STARTER
- IDACIO-PSORIASIS STARTER
- SIMLANDI (2 PEN)
- YUSIMRY SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	(RA, AS, JIA, PsA, PsO, CD, UC, HS, UV): Concurrent therapy with any other biologic disease modifying antirheumatic drug (DMARD), e.g. tumor necrosis factor antagonists

PA Criteria	Criteria Details
Required Medical Information	<p>Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) Cyltezo, (g) Yuflyma OR documentation demonstrating that a trial may be inappropriate.</p> <p>Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate TWO of the following: (a) Cosentyx, (b) Humira, (c) Enbrel, (d) Xeljanz/Xeljanz XR, (e) Rinvoq, (f) Cyltezo, (g) Yuflyma or documentation demonstrating that a trial may be inappropriate.</p> <p>Juvenile Idiopathic Arthritis (JIA): (1) Diagnosis of moderate to severe JIA (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Xeljanz, (d) Orencia, (e) Cyltezo, (f) Yuflyma, (g) Rinvoq/Rinvoq LQ OR documentation demonstrating that a trial may be inappropriate.</p> <p>Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Xeljanz/Xeljanz XR (d) Rinvoq/Rinvoq LQ, (e) Skyrizi, (f) Cosentyx, (g) Stelara, (h) Orencia, (i) Otezla, (j) Cyltezo, (k) Yuflyma OR documentation demonstrating that a trial may be inappropriate.</p> <p>Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe chronic PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Skyrizi (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate.</p>
Age Restrictions	
Prescriber Restrictions	<p>(RA, JIA, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist. (PsO, HS): Prescribed by or in consultation with a dermatologist. (CD, UC): Prescribed by or in consultation with a gastroenterologist. (UV): Prescribed by or in consultation with a rheumatologist or ophthalmologist.</p>
Coverage Duration	Indefinite

PA Criteria	Criteria Details
Other Criteria	Crohn's Disease (CD): (1) Diagnosis of CD. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Stelara, (c) Skyrizi, (d) Cyltezo, (e) Yuflyma, (f) Rinvoq or documentation demonstrating that a trial may be inappropriate. Ulcerative Colitis (UC): (1) Diagnosis of UC. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Stelara, (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Cyltezo, (f) Yuflyma, (g) Skyrizi OR documentation demonstrating that a trial may be inappropriate. Hidradenitis Suppurativa (HS): (1) Diagnosis of HS (2) Inadequate response or inability to tolerate one of the following: (a) Humira, (b) Cyltezo, (c) Yuflyma, (d) Cosentyx OR documentation demonstrating that a trial may be inappropriate. Uveitis (UV) (1) Diagnosis of non-infectious intermediate, posterior, or pan- uveitis. (2) Inadequate response or inability to tolerate BOTH of the following: (a) corticosteroid or immunosuppressive drug (e.g., azathioprine, cyclosporine, methotrexate, mycophenolate), AND (b) One of the following: (i) Humira, (ii) Cyltezo, (iii) Yuflyma OR documentation demonstrating that a trial may be inappropriate.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ADALIMUMAB PREFERRED PRODUCTS

Products Affected

- CYLTEZO (2 PEN)
- CYLTEZO (2 SYRINGE)
- CYLTEZO-CD/UC/HS STARTER
- CYLTEZO-PSORIASIS/UV STARTER
- YUFLYMA (1 PEN)
- YUFLYMA (2 SYRINGE)
- YUFLYMA-CD/UC/HS STARTER

PA Criteria	Criteria Details
Exclusion Criteria	(RA, AS, JIA, PsA, PsO, CD, UC, HS, UV): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g., tumor necrosis factor antagonists
Required Medical Information	Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate one DMARD: (e.g., methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate one other treatment such as NSAIDs, COX2 inhibitors or methotrexate. Juvenile Idiopathic Arthritis (JIA): (1) Diagnosis of moderate to severe JIA. (2) Inadequate response or inability to tolerate one of the following: methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least one DMARD: (e.g., methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe chronic PsO.
Age Restrictions	Member is within the age group listed in the FDA labeling for the indication
Prescriber Restrictions	(RA, JIA, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist. (PsO, HS): Prescribed by or in consultation with a dermatologist. (CD, UC): Prescribed by or in consultation with a gastroenterologist. (UV): Prescribed by or in consultation with a rheumatologist or ophthalmologist.
Coverage Duration	Indefinite

PA Criteria	Criteria Details
Other Criteria	Crohn's Disease (CD): (1) Diagnosis of CD. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g., corticosteroid, methotrexate, azathioprine or 6-mercaptopurine). Ulcerative Colitis (UC): (1) Diagnosis of UC. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g., corticosteroid, Aminosalicylate, azathioprine or 6-mercaptopurine). Hidradenitis Suppurativa (HS): (1) Diagnosis of HS. Uveitis (UV)(Initial): (1) Diagnosis of non-infectious intermediate, posterior, or pan-uveitis. (2) Inadequate response or inability to tolerate corticosteroids and at least one immunosuppressive drug (e.g., azathioprine, cyclosporine, methotrexate, mycophenolate).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ADBRY

Products Affected

- ADBRY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Atopic Dermatitis (AD): (1) Diagnosis of moderate to severe AD (2) inadequate response or inability to tolerate BOTH of the following: (a) one topical steroid (medium potency or higher) AND (b) topical tacrolimus.
Age Restrictions	Member is 12 years of age or older
Prescriber Restrictions	(AD): Prescribed by or in consultation with a dermatologist, allergist, immunologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ADEMPAS

Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	(PAH, CTEPH) (Initial, Reauth): Concurrent use of phosphodiesterase inhibitors, nitrates or nitric oxide donors, pregnancy, concurrent use with other soluble guanylate cyclase stimulators (e.g. Verquvo)
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (initial): (1) Diagnosis of PAH WHO Group I with New York Heart Association (NYHA) Functional Class II - IV. (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography. (3) Mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion. Chronic Thromboembolic Pulmonary Hypertension (CTEPH) (initial): (1) Diagnosis of persistent/recurrent CTPH (WHO Group 4) after surgical treatment or inoperable CTEPH.
Age Restrictions	(PAH, CTEPH) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(PAH, CTEPH) (initial, Reauth): Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	(Initial): 6 months (Reauth):12 months
Other Criteria	(PAH, CPTEH) (Reauth): Stabilization or improvement.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AFREZZA

Products Affected

- AFREZZA INHALATION POWDER 12 UNIT, 4 UNIT, 60X4 & 60X8 & 60X12 UNIT, 8 UNIT, 90 X 4 UNIT & 90X8 UNIT, 90 X 8 UNIT & 90X12 UNIT

PA Criteria	Criteria Details
Exclusion Criteria	(DM1, DM2) (Initial, Reauth): Deny for any of the following (1) Member is currently a smoker or recently stopped smoking (past 6 months), (2) Member has chronic lung disease such as asthma or chronic obstructive pulmonary disease (3) active lung cancer
Required Medical Information	Type 1 Diabetes Mellitus (DM1) (Initial): (1) Diagnosis of type 1 diabetes mellitus, (2) Used in combination with a long-acting insulin (e.g. Lantus, Levemir), (3) Documented FEV1 within the last 60 days greater than or equal to 70% of expected normal as determined by the physician, (4) Spirometry (FEV1) has been completed prior to initiation of therapy to identify potential lung disease (must provide the result). Type 2 Diabetes Mellitus (DM2) (Initial): (1) Diagnosis of type 2 diabetes mellitus, (2) Documented FEV1 within the last 60 days greater than or equal to 70% of expected normal as determined by the physician, (3) Spirometry (FEV1) has been completed prior to initiation of therapy to identify potential lung disease (must provide the result).
Age Restrictions	(DM1, DM2) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial): 6 months (Reauth): Indefinite
Other Criteria	(DM1, DM2)(Reauth): (1) Spirometry value (FEV1) that has not declined greater than or equal to 20% from baseline. (2) Documentation of positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AGAMREE 2024

Products Affected

- AGAMREE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(DMD)(Initial): (1) Diagnosis of Duchenne muscular dystrophy (DMD) (2) Member has received genetic testing for a mutation of the dystrophin gene (3) One of the following: (a) Member has a confirmed mutation of the dystrophin gene (b) Muscle biopsy confirmed an absence of the dystrophin gene (4) Inadequate response or inability to tolerate ONE of the following: (a) prednisone (b) prednisolone (5) One of the following: (a) for members less than or equal to 50kg, dose will not exceed 6mg/kg of body weight once daily (b) For members greater than 50kg, dose will not exceed 300mg/day
Age Restrictions	
Prescriber Restrictions	(DMD) (Initial, Reauth): Prescribed by or in consultation with a neurologist who has experience treating children with DMD.
Coverage Duration	(DMD) (Initial, Reauth): 12 months
Other Criteria	(DMD)(Reauth): (1) Member has experienced a benefit from therapy (e.g., improvement or preservation of muscle strength) (2) One of the following: (a) for members less than or equal to 50kg, dose will not exceed 6mg/kg of body weight once daily (b) For members greater than 50kg, dose will not exceed 300mg/day
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AIMOVIG

Products Affected

- AIMOVIG

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of concomitant use with another injectable CGRP inhibitor.
Required Medical Information	Migraines (Initial): Approved when ONE of the following inclusion criteria are met: (1) Diagnosis of episodic migraines defined as 4 to 14 headache days per month AND inadequate response or inability to tolerate a 4-week trial of TWO of the following prophylactic medications (a) topiramate (b) divalproex sodium/ valproic acid (c) beta-blocker: metoprolol, propranolol, timolol, atenolol, nadolol (d) tricyclic antidepressants: amitriptyline, nortriptyline (e) SNRI antidepressants: venlafaxine, duloxetine. OR (2) Diagnosis of chronic migraines defined as greater than or equal to 15 headache days per month AND inadequate response or inability to tolerate a 4 week trial of TWO of the following prophylactic medications (i) topiramate (ii) divalproex sodium/ valproic acid (iii) beta-blocker: metoprolol, propranolol, timolol, atenolol, nadolol (iv) tricyclic antidepressants: amitriptyline, nortriptyline (v) SNRI antidepressants: venlafaxine, duloxetine.
Age Restrictions	(Migraines)(Initial, Reauth): Member 18 years of age or older
Prescriber Restrictions	(Migraines)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist
Coverage Duration	(Migraines)(Initial): 6 months (Migraines)(Reauth): 12 months
Other Criteria	(Migraines)(REAUTH): (1) Response to therapy as defined by a reduction in headache days per month (defined as at least 4 hours duration and moderate intensity).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AJOVY

Products Affected

- AJOVY

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of concomitant use with another injectable CGRP inhibitor.
Required Medical Information	Migraines (Initial): Approved when ONE of the following inclusion criteria are met: (1) Diagnosis of episodic migraines defined as 4 to 14 headache days per month AND inadequate response or inability to tolerate Aimovig or Emgality. OR (2) Diagnosis of chronic migraines defined as greater than or equal to 15 headache days per month AND inadequate response or inability to tolerate Aimovig or Emgality.
Age Restrictions	(Migraines)(Initial, Reauth): Member 18 years of age or older
Prescriber Restrictions	(Migraines)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist
Coverage Duration	(Migraines)(Initial): 6 months (Migraines)(Reauth): 12 months
Other Criteria	(Migraines)(REAUTH): (1) Response to therapy as defined by a reduction headache days per month (defined as at least 4 hours duration and moderate intensity).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ALLERGEN SPECIFIC IMMUNOTHERAPY (SL)

Products Affected

- GRASTEK
- ODACTRA
- ORALAIR

PA Criteria	Criteria Details
Exclusion Criteria	(Initial): Deny with documentation of any of the following: (1) Severe, unstable or uncontrolled asthma (2) History of eosinophilic esophagitis
Required Medical Information	(Initial): (1) Member has a positive skin test or in vitro test for the listed pollen-specific IgE antibody. (2) Inadequate response or inability to tolerate an intranasal corticosteroid and an oral or intranasal antihistamine.
Age Restrictions	
Prescriber Restrictions	(Initial, Reauth): Prescribed by or in consultation with an allergist or immunologist.
Coverage Duration	(Initial, Reauth): Remainder of contract year
Other Criteria	(Reauth): (1) Member has experienced improvement in the symptoms of their allergic rhinitis or a decrease in the number of medications needed to control allergy symptoms
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ALVAIZ 2024

Products Affected

- ALVAIZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic Idiopathic Thrombocytopenic Purpura (ITP)(Initial): (1) Diagnosis of relapsed/refractory, persistent, or chronic ITP. (2) Baseline platelet count is less than 30,000/mcL. (3) Member's degree of thrombocytopenia and clinical condition increase the risk of bleeding. (4) Inadequate response or inability to tolerate ONE of the following: (a) corticosteroids, (b) immune globulin, (c) splenectomy. Refractory Severe Aplastic Anemia (RSAA)(Initial): (1) Diagnosis of refractory severe aplastic anemia. (2) Member has a platelet count less than 30,000/mcL. (3) Inadequate response or inability to tolerate immunosuppressive therapy with antithymocyte globulin and cyclosporine. Chronic Hepatitis C-Associated Thrombocytopenia (HEPC-TP): (1) Diagnosis of chronic hepatitis C-associated thrombocytopenia. (2) One of the following: (a) Planning to initiate and maintain interferon-based treatment, or (b) currently receiving interferon-based treatment.
Age Restrictions	
Prescriber Restrictions	(ITP, RSAA)(Initial, Continuation): Prescribed by or in consultation with hematologist/oncologist. (HEPC-TP)(Initial, Continuation): Prescribed by or in consultation with Hematologist/oncologist, Gastroenterologist, Hepatologist, Infectious disease specialist, HIV specialist.
Coverage Duration	(ITP)(Initial):12mo (RSAA)(Initial):6mo (HEPC-TP)(Initial):3mo (ITP,RSAA,HEPC-TP)(Continuation):12mo

PA Criteria	Criteria Details
Other Criteria	(ITP)(Continuation): (1) Positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. RSAA (Continuation): (1) Positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. (HEPC-TP)(Continuation): (1) ONE of the following: (A) For members that started treatment with Alvaiz prior to initiation of treatment with interferon, both of the following: (i) Member is currently on antiviral interferon therapy for treatment of chronic hepatitis C, AND (ii) documentation that member reached a threshold platelet count that allows initiation of antiviral interferon therapy with Alvaiz treatment by week 9. OR (B) For members that started treatment with Alviaz while on concomitant treatment with interferon, member is currently on antiviral interferon therapy for treatment of chronic hepatitis C.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AMPYRA

Products Affected

- AMPYRA
- *dalfampridine er*

PA Criteria	Criteria Details
Exclusion Criteria	MS (Initial): Deny if member has history of seizure or moderate to severe renal impairment (CrCL less than or equal to 50 mL/min)
Required Medical Information	Multiple Sclerosis (MS) (Initial): (1) Diagnosis of multiple sclerosis. (2) Confirmation that member has difficulty walking. (3) Inadequate response or inability to tolerate dalfampridine (applies to brand Ampyra).
Age Restrictions	
Prescriber Restrictions	(MS) (Initial and Reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	(Initial, Reauth): Remainder of contract year
Other Criteria	(MS) (Reauth): Improvement in walking speed
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

APOKYN

Products Affected

- APOKYN SUBCUTANEOUS SOLUTION CARTRIDGE
- *apomorphine hcl subcutaneous*

PA Criteria	Criteria Details
Exclusion Criteria	(PD): Member not using medication with any 5-HT3 antagonist (e.g. ondansetron, granisetron, dolasetron, palonosetron, alosetron)
Required Medical Information	Parkinson's Disease (PD): (1) Diagnosis of Parkinson's disease, (2) Member is experiencing intermittent OFF Episodes, (3) Concomitant use of medication with other medications for the treatment of Parkinson's disease (e.g. carbidopa/levodopa, pramipexole, ropinirole, etc.)
Age Restrictions	
Prescriber Restrictions	(PD): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ARIKAYCE

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(MAC): (1) Diagnosis of Mycobacterium avium complex (MAC) lung disease (2) Medication is being used as part of a combination antibacterial drug regimen (3) Used in patients who do not achieve at least two negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy
Age Restrictions	Member is 18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or an infectious disease specialist
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ARMODAFINIL

Products Affected

- *armodafinil*
- NUVIGIL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NARCOLEPSY: (1) Diagnosis of Narcolepsy confirmed by a sleep study (unless prescriber provides justification that a sleep study is not feasible). OBSTRUCTIVE SLEEP APNEA/HYPOPNEA SYNDROME (OSAHS): (1) Diagnosis confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). SHIFT WORK SLEEP DISORDER (SWSD): (1) One of the following: (a) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR (b) A sleep study demonstrating loss of a normal sleep-wake pattern (i.e., disturbed chronobiologic rhythmicity). (2) Documentation that the member has no other medical or mental disorder accounting for the symptoms.
Age Restrictions	
Prescriber Restrictions	(Narcolepsy, SWSD, OSAHS): Prescribed by or in consultation with a neurologist or sleep specialist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AUSTEDO

Products Affected

- AUSTEDO & 18 & 24 & 30 MG
- AUSTEDO XR
- AUSTEDO XR PATIENT TITRATION ORAL
TABLET EXTENDED RELEASE THERAPY PACK 12

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Tardive Dyskinesia (TD)(Initial): (1) Diagnosis of TD, (2) Documentation is provided of ONE of the following: (a) Persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering or discontinuation of the offending medication, or (b) Member is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Chorea-Huntington's Disease (CHD): (1) Diagnosis of chorea associated with Huntington's disease
Age Restrictions	
Prescriber Restrictions	(CHD): Prescribed by or in consultation with a neurologist or a psychiatrist. (TD)(Initial, Continuation): Prescribed by or in consultation with a neurologist or a psychiatrist.
Coverage Duration	(TD)(Initial): 3 months. (TD)(Reauth): Indefinite. (CHD): Indefinite.
Other Criteria	(TD)(Reauth): Positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AUTHORIZED GENERICS-AUTHORIZED BRAND ALTERNATIVES

Products Affected

- *dapagliflozin pro-metformin er* 25 mcg/act
- *dapagliflozin propanediol* • *fluticasone-salmeterol inhalation aerosol*
- *fluticasone furoate-vilanterol inhalation aerosol* • *mirabegron er*
powder breath activated 100-25 mcg/act, 200-

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(1) At least 3 months of use of the brand product within the previous 365 days (document drug, duration, dose and date of use) (2) Both of the following: (a) Documentation provided stating that brand product has not been effective (b) Justification provided for why the target drug is expected to provide benefit when the brand product has not been shown to be effective
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until end of the contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

AUVELITY

Products Affected

- AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Major Depressive Disorder (MDD): (1) Diagnosis of major depressive disorder (MDD) (2) Inadequate response or inability to tolerate two formulary antidepressants indicated for MDD (e.g., bupropion, venlafaxine extended-release tablet or capsule, sertraline tablet, etc.)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(MDD): Approve if for continuation of therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

BENLYSTA SC

Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when the following are met: Systemic Lupus Erythematosus (SLE): (1) Diagnosis of active, autoantibody-positive SLE. (2) Member is receiving concurrent treatment with at least ONE of the following: steroids, antimalarials, immunosuppressants, nonsteroidal anti-inflammatory drugs (NSAIDS). Lupus Nephritis (LN): (1) Diagnosis of active lupus nephritis confirmed by kidney biopsy. (2) Member is receiving concurrent standard therapy (e.g. corticosteroids, immunosuppressants, azathioprine).
Age Restrictions	
Prescriber Restrictions	(SLE): Prescribed by or in consultation with a rheumatologist. (LN): Prescribed by or in consultation with a nephrologist or rheumatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

BESREMI

Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when ANY of the following inclusion criteria is met: (1) Drug is FDA approved for indication and regimen requested, (2) The indication and regimen is classified as Category 1 or 2A by National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium, (3) narrative text in The American Hospital Formulary Service-Drug Information (AHFS-DI) or Clinical Pharmacology Compendium is supportive for the specific condition(s) requested, (4) The Micromedex Compendium and the strength of recommendation is listed as Class I, Class IIa, or Class IIb for the specific condition(s) requested, (5) Indication is listed in Lexi-Drugs as "off label" with evidence level A, (6) supported by Peer-Reviewed Medical Literature as defined in Chapter 15 Section 50.4.5 of the Medicare Benefit Policy Manual
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

BIMZELX

Products Affected

- BIMZELX

PA Criteria	Criteria Details
Exclusion Criteria	(PsO): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Skyrizi (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(PsO): Member is 18 years of age or older
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

BYLVAY

Products Affected

- BYLVAY
- BYLVAY (PELLETS)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pruritus associated with progressive familial intrahepatic cholestasis (Pruritus with PFIC) (Initial): (1) Diagnosis of pruritus associated with progressive familial intrahepatic cholestasis (PFIC), (2) Confirmed molecular diagnosis of PFIC type 1, 2, or 3, (3) Prescribed dose is consistent with FDA-approved package labeling and does not exceed a total daily dose of 6mg Cholestatic Pruritus with Alagille Syndrome (CPALGS) (Initial): (1) Diagnosis of Alagille Syndrome. (2) Molecular genetic testing confirms mutations in the JAG1 or NOTCH2 gene (3) Member is experiencing both of the following: (a) moderate to severe cholestatic pruritus (b) member has a serum bile acid concentrations above the upper limit of the normal reference for the reporting laboratory (4) An inadequate response or inability to tolerate at least two of the following treatments used for the relief of pruritus: (a) Ursodeoxycholic acid (e.g., Ursodiol) (b) Antihistamines (e.g., diphenhydramine, hydroxyzine) (c) Rifampin (d) Bile acid sequestrants (e.g., Questran, Colestid, Welchol).
Age Restrictions	(Pruritus with PFIC) (Initial, Reauth): Member is 3 months of age or older (CPALGS)(Initial, Reauth): Member is 1 year of age or older
Prescriber Restrictions	(Pruritus with PFIC)(CPALGS)(Initial, Reauth): Prescribed by or in consultation with a hepatologist or gastroenterologist
Coverage Duration	(All)(Initial):6 months (Pruritus w/PFIC)(Reauth):Indefinite (CPALGS)(Reauth):End of contract year
Other Criteria	(Pruritus with PFIC) (Reauth): (1) Documentation of positive clinical response to therapy (e.g. improvement in pruritus symptoms), (2) Prescribed dose is consistent with FDA-approved package labeling and does not exceed a total daily dose of 6mg. (CPALGS) (Reauth): Positive clinical response to therapy (e.g., reduced bile acids, reduction in pruritis symptoms or ItchRO pruritis score).
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

CAMZYOS

Products Affected

- CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Obstructive hypertrophic cardiomyopathy (HCM) (1) Diagnosis of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM). (2) Member's baseline left ventricular ejection fraction (LVEF) is greater than or equal to 55%. (3) Documentation of Valsalva left ventricular outflow tract (LVOT) gradient assessment at baseline. (4) Inadequate response or inability to tolerate one of the following: (a) one non-vasodilating beta-blocker (e.g., bisoprolol, propranolol), (b) one calcium-channel blocker (e.g., diltiazem, verapamil)
Age Restrictions	(HCM) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(HCM) (Initial, Reauth): Prescribed by or in consultation with a cardiologist
Coverage Duration	(Initial): 6 months. (Reauth): 12 months
Other Criteria	(HCM) (Reauth): (1) Documentation of improvement in functional capacity and symptoms. (2) Member's left ventricular ejection fraction (LVEF) is greater than or equal to 50%. (3) Member does not have worsening heart failure symptoms.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CAPLYTA

Products Affected

- CAPLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Schizophrenia: (1) Diagnosis of schizophrenia (2) Inadequate response or inability to tolerate two generic antipsychotic products (e.g. aripiprazole, olanzapine, quetiapine, paliperidone). Bipolar Depression (BD): (1) Diagnosis of bipolar depression (2) Medication will be used as monotherapy or as adjunctive therapy with lithium or valproate (3) Inadequate response or inability to tolerate two medications indicated for bipolar depression (e.g. fluoxetine, Latuda, quetiapine, olanzapine)
Age Restrictions	(Schizophrenia, BP): member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Schizophrenia, BP): Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CARBAGLU

Products Affected

- CARBAGLU ORAL TABLET SOLUBLE
- *carglumic acid oral tablet soluble*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hyperammonemia Type III (HTIII): (1) Hyperammonemia due to the deficiency of the hepatic enzyme N-acetyl glutamate synthase (NAGS). Acute Hyperammonemia due to Propionic Acidemia or Methylmalonic Acidemia (AH): (1) Hyperammonemia due to propionic acidemia or methylmalonic acidemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CAYSTON

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis (CF): (1) Diagnosis of CF. (2) Evidence of Pseudomonas Aeruginosa in the lungs. (3) Susceptibility results indicating that the Pseudomonas aeruginosa is sensitive to aztreonam. (4) FEV1 between 25% and 75% of predicted
Age Restrictions	(CF): Member is 7 years of age or older
Prescriber Restrictions	(CF): Prescribed by or in consultation with a pulmonologist OR infectious disease specialist or Specialist affiliated with a CF care center.
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CERDELGA

Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	(GD): Member is CYP2D6 Ultra Rapid Metabolizer (URM)
Required Medical Information	Gaucher disease (GD): (1) Diagnosis of Type 1 Gaucher disease and member is CYP2D6 extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM) as detected by an FDA-cleared test for determining CYP2D6 genotype.
Age Restrictions	(GD): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CHOLBAM

Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	(BASD, PD)(Initial, Reauth): Deny if there is documentation of extrahepatic manifestations of bile acid synthesis disorders due to single enzyme defects or peroxisomal disorders including Zellweger spectrum disorders
Required Medical Information	Bile Acid Synthesis Disorder (BASD) (initial): (1) Diagnosis of bile acid synthesis disorder due to a single enzyme defect. Peroxisomal disorder (PD) (initial): (1) Diagnosis of peroxisomal disorder. (2) Used as adjunctive treatment. (3) Member exhibits manifestations of liver disease, steatorrhea, or complications from decreased fat soluble vitamin absorption.
Age Restrictions	
Prescriber Restrictions	(BASD, PD): Hepatologist, Gastroenterologist, Medical geneticist, other specialist that treats inborn errors of metabolism
Coverage Duration	(Initial): 3 months. (Reauth): Indefinite
Other Criteria	(BASD,PD) (Reauth): Documentation of improved liver function tests from the start of treatment.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CIALIS

Products Affected

- CIALIS ORAL TABLET 5 MG
- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	(BPH): Concurrent use of nitrates.
Required Medical Information	Benign prostatic hyperplasia (BPH): (1) Diagnosis of BPH. (2) Inadequate response or inability to tolerate an alpha blocker (e.g. tamsulosin, terazosin) or a 5-alpha reductase inhibitor (e.g. dutasteride, finasteride).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CIBINQO

Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	(AD): Concurrent use with any other biologic immunomodulator, Janus Kinase (JAK) inhibitors, or other immunosuppressants (e.g., azathioprine, cyclosporine)
Required Medical Information	Atopic Dermatitis (AD): (1) Diagnosis of refractory, moderate to severe AD (2) Inadequate response or inability to tolerate one systemic drug product, including biologics (e.g., Dupixent, methylprednisolone, prednisone) or documentation that a trial may be inappropriate.
Age Restrictions	(AD): Member is 12 years of age or older
Prescriber Restrictions	(AD): Prescribed by or in consultation with a dermatologist, allergist, immunologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CIMZIA

Products Affected

- CIMZIA (2 SYRINGE)
- CIMZIA SUBCUTANEOUS KIT 2 X 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	(AS, PsA, PsO, RA, CD, nr-axSpA): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate two of the following: (a) secukinumab (Cosentyx), (b) adalimumab (Humira), (c) etanercept (Enbrel), (d) tofacitinib (Xeljanz/Xeljanz XR), (e) upadacitinib (Rinvoq), (f) adalimumab-adbm (Cyltezo), (g) adalimumab-aaty (Yuflyma) or documentation demonstrating that a trial may be inappropriate. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Xeljanz/Xeljanz XR (d) Rinvoq/Rinvoq LQ, (e) Skyrizi, (f) Cosentyx, (g) Stelara, (h) Orencia, (i) Otezla, (j) Cyltezo, (k) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Skyrizi (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) Cyltezo, (g) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Crohn's Disease (CD): (1) Diagnosis of moderate to severe CD. (2) Inadequate response or inability to tolerate two of the following: (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) risankizumab (Skyrizi), (d) adalimumab-adbm (Cyltezo), (e) adalimumab-aaty (Yuflyma), (f) upadacitinib (Rinvoq) or documentation demonstrating that a trial may be inappropriate. Nonradiographic axial Spondyloarthritis (nr-axSpA): (1) Diagnosis of nr-axSpA. (2) Inadequate response or inability to tolerate one of the following: (a) two NSAIDs (b) Cosentyx (c) Rinvoq or (d) documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(AS, PsA, PsO, RA, CD, nraxSpA): Member is 18 years of age or older
Prescriber Restrictions	(CD): Prescribed by or in consultation with a gastroenterologist. (RA, AS, nr-axSpA): Prescribed by or in consultation with a rheumatologist. (PsA, PsO): prescribed by or in consultation with a dermatologist or rheumatologist.

PA Criteria	Criteria Details
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CINRYZE

Products Affected

- CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when: Hereditary Angioedema (HAE): (1) Diagnosis of hereditary angioedema (HAE) (2) For prophylaxis against HAE attacks.
Age Restrictions	(HAE): Member is 6 years of age or older
Prescriber Restrictions	(HAE): Prescribed by or in consultation with an allergist, immunologist, pulmonologist
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

COMBINATION NSAID PRODUCTS

Products Affected

- *ibuprofen-famotidine*
- *naproxen-esomeprazole mg*
- VIMOVO ORAL TABLET DELAYED RELEASE 500-20 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(All Indications)(Initial): An inadequate response or inability to tolerate a two-week trial of BOTH of the following: (1) Concurrent administration of each of the components of the requested product, and (2) At least ONE generic alternative (when available) of each of the individual components of the requested product.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(All Indication) (Initial, Reauth): 1 year
Other Criteria	(All Indications)(Reauth): Positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CORLANOR

Products Affected

- CORLANOR
- *ivabradine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic Heart Failure (CHF): (1)Diagnosis of chronic heart failure. (2) stable, symptomatic chronic heart failure (3) left ventricular ejection fraction less than or equal to 35% and (4) sinus rhythm with resting heart rate greater than or equal to 70 beats per minute. (5) Member is clinically stable for at least 4 weeks on an optimized regimen which includes: (a) maximally tolerated doses of beta blockers or inability to tolerate beta blockers, (b) ACE inhibitors or ARBs or inability to tolerate ACE inhibitor or ARB. Chronic Heart Failure due to Dilated Cardiomyopathy in pediatric patients ages 6 months and older (CHF-DC): (1) Diagnosis of chronic heart failure due to dilated cardiomyopathy (2) Member is in sinus rhythm with an elevated heart rate.
Age Restrictions	
Prescriber Restrictions	(CHF, CHF-DC): Prescribed by or in consultation with a cardiologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

COSENTYX

Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)
- COSENTYX SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE 75 MG/0.5ML
- COSENTYX UNOREADY

PA Criteria	Criteria Details
Exclusion Criteria	(PsA, PsO, AS, nr-axSpA, ERA, HS): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least ONE DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe plaque psoriasis. Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate one other treatment such as NSAIDs, COX2 inhibitors or methotrexate. Non-Radiographic Axial Spondyloarthritis (nr-axSpA): (1) Diagnosis of nr-axSpA, (2) Inadequate response or inability to tolerate two NSAIDs. Active Enthesitis-related Arthritis (ERA): (1) Diagnosis of active enthesitis-related arthritis (ERA), (2) Inadequate response or inability to tolerate methotrexate, (3) Inadequate response or inability to tolerate at least one NSAID (e.g. ibuprofen, naproxen, meloxicam, celecoxib). Hidradenitis Suppurativa (HS): (1) Diagnosis of HS.
Age Restrictions	(AS, nr-axSpA, HS): Member is 18 years of age or older. (PsO): Member is 6 years of age or older (PsA): Member is 2 years of age or older. (ERA): Member is 4 years of age or older.
Prescriber Restrictions	(PsO, HS): Prescribed by or in consultation with a dermatologist. (PsA): Prescribed by or in consultation with a rheumatologist or dermatologist. (AS, nr-axSpA, ERA): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CRESEMBA [ORAL]

Products Affected

- CRESEMBA ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Invasive Aspergillosis (IA): (1) For use in the treatment of invasive aspergillosis after inadequate response or inability to tolerate Voriconazole (oral Vfend). Mucormycosis (MC): (1) Diagnosis of invasive mucormycosis
Age Restrictions	(IA, MC): Member is 6 years of age or older
Prescriber Restrictions	(All Indications): Prescribed by or in consultation with an infectious disease specialist, oncologist, OR prescribed as part of chemotherapy prophylaxis protocol
Coverage Duration	Remainder of contract year
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

CYSTEAMINE PRODUCTS

Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystinosis: (1) Diagnosis of cystinosis, (2) Member has corneal cystine crystal accumulation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DAYBUE

Products Affected

- DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Rett syndrome (RS)(Initial): (1) Diagnosis of Rett syndrome (2) One of the following: (a) Presence of all of the following clinical signs and symptoms: (i) a pattern of development, regression, then recovery or stabilization (ii) partial or complete loss of purposeful hand skills such as grasping with fingers, reaching for things, or touching things on purpose (iii) partial or complete loss of spoken language (iv) repetitive hand movements, such as wringing the hands, washing, squeezing, clapping, or rubbing (v) gait abnormalities, including walking on toes or with an unsteady, wide-based, stiff-legged gait (b) Molecular genetic testing confirms mutations in the MECP2 gene
Age Restrictions	(RS)(Initial)(Reauth): member is 2 years of age or older
Prescriber Restrictions	(RS)(Initial)(Reauth): Prescribed by or in consultation with one of the following specialists: geneticist, pediatrician, neurologist
Coverage Duration	(Initial)(Reauth): 12 months
Other Criteria	(RS)(Reauth): (1) Documentation of positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms from baseline (e.g., hand behavior, walking/standing, speech, quality of life)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DAYVIGO

Products Affected

- DAYVIGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Insomnia: (1) Diagnosis of insomnia. (2) Inadequate response or inability to tolerate generic ramelteon (Rozerem).
Age Restrictions	(Insomnia): Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DEFERASIROX

Products Affected

- *deferasirox granules*
- *deferasirox oral tablet*
- *deferasirox oral tablet soluble*
- EXJADE
- JADENU
- JADENU SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	(NTDT, CIO-BT)(Initial, Continuation): CrCl less than 40 mL/min or serum creatinine more than 2 times the age-appropriate ULN, platelet counts less than 50,000/mcL
Required Medical Information	Chronic Iron Overload in nontransfusion-dependent thalassemia (NTDT) (Initial): (1) Diagnosis of Chronic iron overload in nontransfusion-dependent thalassemia syndromes,(2) liver iron concentration (LIC) of at least 5 mg of iron per gram of liver dry weight (mg Fe/g dw), (3) serum ferritin greater than 300 mcg/L. (4) Serum ferritin levels consistently greater than 300 mcg/L (as demonstrated with at least two lab values within the previous two months). Chronic Iron Overload Caused by Blood Transfusions (CIO-BT)(Initial): (1) Diagnosis of chronic iron overload caused by blood transfusions (transfusional hemosiderosis). (2) Serum ferritin levels consistently greater than 1,000 mcg/L (as demonstrated with at least two lab values within the previous two months).
Age Restrictions	(NTDT)(Initial, Continuation): Member is 10 years of age or older. (CIO-BT) (Initial, Continuation): Member is 2 years of age or older.
Prescriber Restrictions	
Coverage Duration	(Initial):3 months. (Continuation): 6 months
Other Criteria	(CIO-BT)(Continuation): (1) Decreased serum ferritin level compared with the baseline level for transfusion- dependent anemia. (NTDT)(Continuation): (1) Decreased serum ferritin level compared with the baseline level or reduction in LIC (liver iron concentration).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DIACOMIT

Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dravet syndrome (DS): (1) Diagnosis of seizures associated with Dravet syndrome (DS), (2) Used in combination with clobazam.
Age Restrictions	Member is 6 months of age or older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC 3% PRODUCTS

Products Affected

- *diclofenac sodium external gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Actinic Keratoses: (1) Diagnosis of Actinic Keratoses
Age Restrictions	Actinic Keratoses: Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	90 days
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC EPOLAMINE

Products Affected

- *diclofenac epolamine external*
- FLECTOR EXTERNAL
- LICART EXTERNAL

PA Criteria	Criteria Details
Exclusion Criteria	Previously experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Use for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery.
Required Medical Information	(1): Inadequate response or inability to tolerate at least 2 prescription strength topical NSAIDs (i.e. Diclofenac Gel 1%, Diclofenac Topical Solution 1.5%)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DOJOLVI

Products Affected

- DOJOLVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Long-Chain Fatty Acid Oxidation Disorders (LC-FAOD): (1) Diagnosis of molecularly confirmed LC-FAOD, (2) Will be used as a source of calories and fatty acids.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

DOPTELET

Products Affected

- DOPTELET ORAL TABLET 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic Liver Disease (CLD): (1) Baseline platelet count less than 50,000/mcL. Member is scheduled to undergo a procedure. Chronic Immune Thrombocytopenia (ITP)(Initial): (1) Baseline platelet count less than 30,000/mcL. (2) Inadequate response or inability to tolerate one of the following: (a) corticosteroids (b) immunoglobulins (c) splenectomy (d) Rituxan (rituximab)
Age Restrictions	
Prescriber Restrictions	(ITP)(Initial, Continuation): Prescribed by or in consultation with hematologist/oncologist
Coverage Duration	(CLD): 1 month. (ITP): 12 months
Other Criteria	(ITP)(Continuation): (1) Positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	Yes

DUPIXENT

Products Affected

- DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR MG/2ML
- DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 200 MG/1.14ML, 300

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Atopic Dermatitis (AD): (1) Diagnosis of moderate-severe atopic dermatitis. (2) inadequate response or inability to tolerate ONE of the following: (a) one topical steroid (medium potency or higher), (b) topical tacrolimus, (c) topical pimecrolimus, (d) topical Eucrisa (crisaborole). Asthma (Initial): (1) ONE of the following: (a) Member has oral corticosteroid-dependent asthma or (b) Member has blood eosinophil levels of at least 150 cells/mcL at baseline or at least 300 cells/mcL within the past 12 months, (2) the requested medication will be used in addition to BOTH of the following: (a) medium to high dose inhaled corticosteroid (e.g. greater than or equal to 500mcg fluticasone propionate equivalent/day) AND (b) one additional controller medication. Chronic Rhinosinusitis with Nasal Polyposis(CRSwNP)(Initial): (1) Diagnosis of chronic rhinosinusitis with nasal polyposis, (2) concurrent use of intranasal corticosteroid. Eosinophilic esophagitis (EoE)(Initial): (1) Diagnosis of eosinophilic esophagitis (EoE) (2) Member has symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, gastroesophageal reflux disease [GERD]/heartburn symptoms, chest pain, abdominal pain (3) Member has at least 15 intraepithelial eosinophils per high power (HPF) (4) Other causes of esophageal eosinophilia have been excluded (5) Member weighs at least 15 kg (6) Inadequate response or inability to tolerate at least an 8 week trial of one of the following: (a) proton pump inhibitors (e.g., pantoprazole, omeprazole (b) Topical (esophageal) corticosteroids (e.g., budesonide, fluticasone). Prurigo Nodularis (PN)(Initial): (1) Diagnosis of Prurigo Nodularis (PN) (2) inadequate response or inability to tolerate one medium or higher potency topical corticosteroid.</p>
Age Restrictions	<p>(Asthma)(Initial, Reauth): Member is 6 years old or older. (AD): Member is 6 months of age or older. (CRSwNP)(Initial, Reauth): Member is 12 years of age or older (PN)(COPD)(Initial, Reauth): Member is 18 years of age or older (EoE)(Initial, Reauth): Member is 1 year of age or older</p>

PA Criteria	Criteria Details
Prescriber Restrictions	(Initial/Reauth) Prescribed by/in consultation with: (AD): Dermatologist, allergist, or immunologist (Asthma)(COPD): Allergist, immunologist or pulmonologist (CRSwNP): Allergist, immunologist or ENT specialist (EoE): Gastroenterologist, allergist, or immunologist (PN): Allergist/immunologist or dermatologist
Coverage Duration	(AD): Indefinite (Asthma, CRSwNP, EoE, PN, COPD): 12 months
Other Criteria	<p>Chronic obstructive pulmonary disease (COPD)(Initial): (1) Diagnosis of COPD (2) Presence of Type 2 inflammation evidenced by blood eosinophils greater than or equal to 300 cells/mcL at baseline (3) Member is receiving ONE of the following therapies for at least 3 months: (a) Triple therapy (i.e., an inhaled corticosteroid (ICS), a long-acting muscarinic antagonist (LAMA) and a long-acting beta agonist (LABA) (b) If ICS are contraindicated, a LAMA and a LABA (4) Member has had ONE of the following within the past 12 months: (a) At least two exacerbations where systemic corticosteroids [intramuscular, intravenous, or oral (e.g., prednisone)] were required at least once (b) COPD-related hospitalization (5) Member experiences dyspnea during everyday activities (e.g., needs to stop for breath when walking on level ground).</p> <p>(Asthma)(Reauth): (1) Documentation of positive clinical response to therapy (e.g. reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications) (2) the requested medication will be used in addition to BOTH of the following: (a) medium to high dose inhaled corticosteroid (e.g. greater than or equal to 500mcg fluticasone propionate equivalent/day) AND (b) one additional controller medication. (CRSwNP) (Reauth): (1) Documentation of positive clinical response to therapy (e.g. reduction of nasal polyps score [NPS: 0-8 scale], improvement in nasal congestion/obstruction score [NC: 0-3 scale]), (2) Used in combination with another agent for CRSwNP. (EoE)(Reauth): (1) Documentation of a positive clinical response to therapy as evidenced by improvement of at least one of the following from baseline: (a)Symptoms (e.g., dysphagia, food impaction, heartburn, chest pain), (b) Histologic measures (e.g., esophageal intraepithelial eosinophil count), (c) Endoscopic measures (e.g., edema, furrows, exudates, rings, strictures). (PN)(Reauth): (1) Documentation of a positive clinical response to therapy (e.g., reduction in the number of nodular lesions from baseline, or improvement in symptoms from baseline) (COPD)(Reauth): (1) Member demonstrates a positive clinical response to therapy (e.g., improved lung function, a reduction in COPD exacerbations) (2) Member continues to receive ONE of the following therapies: (a) Triple therapy (i.e., an inhaled corticosteroid (ICS), a long-acting muscarinic antagonist (LAMA) and a long-acting beta agonist (LABA) (b) If ICS are contraindicated, a LAMA and a LABA</p>
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

DUVYZAT 2024

Products Affected

- DUVYZAT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Duchenne muscular dystrophy (DMD) (Initial): (1) Diagnosis of Duchenne muscular dystrophy (DMD) (2) Diagnosis confirmed by ONE of the following: (a) Mutation of the dystrophin gene (b) Absence of the dystrophin protein confirmed by muscle biopsy (3) Member is ambulatory without needing an assistive device (e.g., without side-by-side assist, cane, walker, wheelchair, etc.) prior to initiating Duvyzat (4) Duvyzat will be used concomitantly with a corticosteroid regimen
Age Restrictions	
Prescriber Restrictions	(DMD) (Initial) (Reauth): Prescribed by or in consultation with a neurologist who has experience treating children with Duchenne Muscular Dystrophy (DMD)
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(DMD) (Reauth): (1) Member has experienced a benefit from therapy (e.g., improvement in preservation of muscle strength) (2) Member is maintaining ambulatory status without needing an assistive device (e.g., without side-by-side assist, cane, walker, wheelchair, etc.) (3) Member continues to receive concomitant corticosteroid regimen.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EGRIFTA

Products Affected

- EGRIFTA SV

PA Criteria	Criteria Details
Exclusion Criteria	(HIV-L): (1) hypothalamic-pituitary axis due to hypophysectomy, hypopituitarism, pituitary tumor/surgery, or head irradiation or trauma, (2) hypersensitivity to tesamorelin and/or mannitol, (3) malignancy, active (either newly diagnosed or recurrent) malignancies should be inactive and completely treated prior to initiating therapy, (4) pregnancy.
Required Medical Information	HIV-Associated Lipodystrophy (HIV-L): (1) Diagnosis of HIV-associated lipodystrophy, (2) one of the following: (a) waist-circumference of greater than or equal to 95 cm (37.4 inches) in men, OR (b) waist-circumference of greater than or equal to 94 cm (37 inches) for women, (3) one of the following: (a) Waist-to-hip ratio of greater than or equal to 0.94 for men, OR (b) waist-to-hip ratio of greater than or equal to 0.88 for women, (4) body mass index (BMI) greater than 20 kg/m ² , (5) fasting blood glucose (FBG) levels less than or equal to 150 mg/dL (8.33 mmol/L), (6) Member has been on a stable regimen of antiretrovirals (e.g. NRTIs, NNRTI, Protease Inhibitors, Integrase Inhibitors) for at least 8 weeks.
Age Restrictions	
Prescriber Restrictions	(HIV-L): Prescribed by or in consultation with HIV-infection specialist OR endocrinologist.
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EMFLAZA

Products Affected

- *deflazacort*
- EMFLAZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Duchenne Muscular Dystrophy (MDM): (1) Diagnosis confirmed by ONE of the following: (a) Mutation of the dystrophin gene (b) Absence of the dystrophin protein confirmed by muscle biopsy (2) Inadequate response or inability to tolerate prednisone or prednisolone
Age Restrictions	(DMD): Member is 2 years of age or older
Prescriber Restrictions	(DMD): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EMGALITY

Products Affected

- EMGALITY
- EMGALITY (300 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of concomitant use with another injectable CGRP inhibitor.
Required Medical Information	Migraines (Initial): Approved when ONE of the following inclusion criteria are met: (1) Diagnosis of episodic migraines defined as 4 to 14 headache days per month AND inadequate response or inability to tolerate a 4 week trial of TWO of the following prophylactic medications (a) topiramate (b) divalproex sodium/ valproic acid (c) beta-blocker: metoprolol, propranolol, timolol, atenolol, nadolol (d) tricyclic antidepressants: amitriptyline, nortriptyline (e)SNRI antidepressants: venlafaxine, duloxetine. OR (2) Diagnosis of chronic migraines defined as greater than or equal to 15 headache days per month AND inadequate response or inability to tolerate a 4 week trial of TWO of the following prophylactic medications (i) topiramate (ii) divalproex sodium/ valproic acid (iii) beta-blocker: metoprolol, propranolol, timolol, atenolol, nadolol (iv) tricyclic antidepressants: amitriptyline, nortriptyline (v)SNRI antidepressants: venlafaxine, duloxetine. Episodic Cluster Headaches (ECH) (Initial): (1) Diagnosis of episodic cluster headache, (2) Member has experienced at least 2 cluster periods lasting 6 days to 365 days, separated by pain-free periods lasting at least three months
Age Restrictions	(Migraine, ECH)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(Migraine, ECH)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, pain specialist
Coverage Duration	(Migraine, ECH)(Initial): 6 months, (Migraine, ECH)(Reauth):12 months
Other Criteria	(Migraine)(REAUTH): (1) Response to therapy as defined by a reduction in headache days per month (defined as at least 4 hours duration and moderate intensity). (ECH)(REAUTH): (1) Response to therapy as defined by a reduction in weekly cluster headache attacks.
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

EMSAM

Products Affected

- EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Major depressive disorder (MDD): (1) Diagnosis of major depressive disorder. (2) Inadequate response or inability to tolerate ONE SSRI or SNRI. (3) 4-5 half-lives (approximately 1 week) has elapsed after discontinuation of antidepressants without long half-lives OR at least 5 weeks has elapsed after discontinuation with antidepressants with long half-lives (e.g. fluoxetine).
Age Restrictions	(MDD): Member is 18 years of age or older.
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED
- SYRINGE
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	(RA, PsA, PJIA, PsO, AS): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate ONE DMARD: (e.g. Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine). Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least ONE DMARD: (e.g. Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine). Polyarticular Juvenile Idiopathic Arthritis (PJIA): (1) Diagnosis of moderate to severe PJIA, (2) Inadequate response or inability to tolerate ONE of the following: Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine. Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe chronic PsO. Ankylosing Spondylitis (AS): (1) Diagnosis of AS, (2) Inadequate response or inability to tolerate one other treatment such as NSAIDs, COX2 inhibitors or methotrexate.
Age Restrictions	(PJIA, PsA): Member is 2 years of age or older. (RA, AS): Member is 18 years of age or older. (PsO): Member is 4 years of age or older
Prescriber Restrictions	(RA, PJIA, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a rheumatologist or dermatologist. (PsO): Prescribed by or in consultation with a dermatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ENDARI

Products Affected

- ENDARI
- *l-glutamine oral packet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Sickle Cell Disease (SC)(Initial): (1) One of the following: (A) Member is using L-Glutamine with concurrent hydroxyurea therapy, OR (B) Member has an inadequate response or inability to tolerate hydroxyurea. (2) Member has had 2 or more painful sickle cell crises within the past 12 months.
Age Restrictions	
Prescriber Restrictions	(SC)(Initial, Reauth): Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(Reauth): Documentation of positive clinical response to therapy (e.g. reduction in the number of sickle cell crises)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ENSPRYNG

Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Neuromyelitis Optica Spectrum Disorder (NMOSD)(Initial): (1) Diagnosis of NMSOD, (2) Member is anti-aquaporin-4 (AQP4) antibody positive.
Age Restrictions	
Prescriber Restrictions	(NMOSD)(Initial, Reauth): Prescribed by or in consultation with a neurologist or ophthalmologist
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(NMOSD)(Reauth): Positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ENTADFI

Products Affected

- ENTADFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Benign Prostatic Hyperplasia (BPH): (1) Diagnosis of Benign Prostatic Hyperplasia (BPH) (2) inadequate response or inability to tolerate one of the following generics (alfuzosin, doxazosin, tamsulosin, terazosin, silodosin) (3) Inadequate response or inability to tolerate one of the following: (a) 5-alpha-reductase inhibitor (i.e., finasteride 5mg, dutasteride) (b) phosphodiesterase type 5 inhibitor (i.e., tadalafil 2.5mg or 5mg)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	26 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ENTYVIO SQ 2024

Products Affected

- ENTYVIO PEN

PA Criteria	Criteria Details
Exclusion Criteria	(UC, CD): Concurrent therapy with any other biologic disease modifying antirheumatic drug (DMARD), e.g., tumor necrosis factor antagonists
Required Medical Information	<p>Ulcerative Colitis (UC): (1) Diagnosis of moderately to severely active Ulcerative Colitis (2) One of the following: (a) Inadequate response or inability to tolerate two of the following: (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) upadacitinib (Rinvoq), (d) tofacitinib (Xeljanz/Xeljanz XR), (e) adalimumab-adbm (Cyltezo), (f) adalimumab-aaty (Yuflyma), (g) risankizumab (Skyrizi) OR documentation demonstrating that a trial may be inappropriate.</p> <p>(b) One of the following: (i) Medication will be used as a maintenance dose following two doses of Entyvio IV for induction (ii) Member is currently established on Entyvio IV.</p> <p>Crohn's Disease (CD): (1) Diagnosis of moderately to severely active CD (2) One of the following: (a) Inadequate response or inability to tolerate TWO of the following (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) Risankizumab (Skyrizi), (d) adalimumab-adbm (Cyltezo), (e) adalimumab-aaty (Yuflyma), (f) upadacitinib (Rinvoq) OR documentation demonstrating that a trial may be inappropriate (b) One of the following: (i) Medication will be used as a maintenance dose following two doses of Entyvio IV for induction (ii) Member is currently established on Entyvio IV.</p>
Age Restrictions	(UC, CD): Member is 18 years of age or older
Prescriber Restrictions	UC, CD): Prescribed by or in consultation with a gastroenterologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EPIDIOLEX

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dravet Syndrome (DS): (1) Inadequate response or inability to tolerate ONE of the following: (a) clobazam (b) valproic acid or (c) topiramate. (2) Baseline CBC, serum transaminases and total bilirubin prior to initiating therapy. Lennox-Gastaut Syndrome (LGS): (1) Inadequate response or inability to tolerate ONE of the following: (a) clobazam (b) valproic acid, or (c) topiramate. (2) Baseline CBC, serum transaminases and total bilirubin prior to initiating therapy. Tuberous Sclerosis Complex (TSC): (1) Concurrent use with additional anti-epileptic(s). (2) Baseline CBC, serum transaminases and total bilirubin prior to initiating therapy
Age Restrictions	(DS, LGS, TCS): Member is 1 year of age or older
Prescriber Restrictions	(DS, LGS, TCS): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EPSOLAY

Products Affected

- EPSOLAY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Rosacea): (1) Diagnosis of rosacea (2) Member has inflammatory lesions (3) inadequate response or inability to tolerate one of the formulary topical products for rosacea (e.g., azelaic acid gel, metronidazole cream or gel) for sufficient duration (minimum 30-day supply)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ESBRIET

Products Affected

- ESBRIET
- *pirfenidone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Idiopathic Pulmonary Fibrosis (IPF): (1) Diagnosis of IPF confirmed by high resolution CT scan or biopsy.
Age Restrictions	
Prescriber Restrictions	(IPF): Prescribed by or in consultation a pulmonologist or lung transplant specialist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(REAUTH) (IPF): BOTH of the following (1) stabilization from baseline AND (2) no elevations in AST or ALT greater than 5 times upper limit of normal or greater than 3 times upper limit of normal with signs or symptoms of severe liver damage.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EUCRISA

Products Affected

- EUCRISA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Atopic Dermatitis (AD): (1) Inadequate response or inability to tolerate at least ONE of the following: (a) topical tacrolimus OR topical pimecrolimus , OR (b) generic, prescription medium potency or higher topical steroid.
Age Restrictions	(AD): Member is 3 months of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EVEKEO

Products Affected

- *amphetamine sulfate*
- EVEKEO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Attention Deficit Hyperactivity Disorder (ADHD): (1) Diagnosis of ADHD. (2) Inadequate response or inability to tolerate TWO generic stimulant products (e.g. amphetamine/dextroamphetamine, methylphenidate) Narcolepsy: (1) Diagnosis of narcolepsy.
Age Restrictions	(ADHD): Member is 3 years of age or older. (Narcolepsy): Member is 6 years of age or older.
Prescriber Restrictions	
Coverage Duration	Remainder of Contract Year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EVENITY

Products Affected

- EVENITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary for: Post Menopausal Osteoporosis (PMO): (1) Diagnosis of PMO defined as ONE of the following: (a) Bone mineral density T-score less than or equal to -2.50 OR (b) Documented history of an osteoporotic non- collision fracture [e.g. vertebral, hip, nonvertebral]) AND (3) Member is at high risk of fracture as defined by one of the following: (a) Member has risk factors for a fracture (e.g. endocrine disorders, gastrointestinal disorders, use of medications associated with low bone mass or bone loss such as corticosteroids) OR (b) Inadequate response or inability to tolerate ONE of the following: (i) bisphosphonates, (ii) hormone replacement therapy, (iii) selective-estrogen receptor modulators (SERMs), OR (iv) Denosumab (Prolia). (4) Cumulative lifetime therapy does not exceed 12 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EVRYSDI

Products Affected

- EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Spinal Muscular Atrophy (SMA)(Initial): (1) Diagnosis of SMA, (2) Member has confirmed mutations in chromosome 5q that leads to SMN protein deficiency.
Age Restrictions	
Prescriber Restrictions	(SMA)(Initial, Reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	(Initial)(Reauth): 12 months
Other Criteria	(SMA)(Reauth): (1) Positive clinical response to therapy (e.g. improvement in ability to sit without support)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EXTENDED RELEASE METFORMIN

Products Affected

- GLUMETZA
- *metformin hcl er (mod)*

PA Criteria	Criteria Details
Exclusion Criteria	(DM2)(Initial, Reauth): (1) Serum creatinine levels greater than or equal to 1.5 mg/dL in males, or serum creatinine levels greater than or equal to 1.4 mg/dL in females. (2) Hepatic impairment. (3) Metabolic acidosis, including diabetic ketoacidosis. (4) Used for preventing weight gain.
Required Medical Information	Diabetes Mellitus Type 2 (DM2)(Initial): (1) Diagnosis of DM2. (2) Member has a HgbA1C greater than 7.0%. All Indications: (1) Inadequate response or inability to tolerate both of the following: (a) Immediate release metformin, and (b) Extended-release metformin (generic Glucophage XR).
Age Restrictions	(DM2 for tablets) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): End of contract year.
Other Criteria	(DM2)(Reauth): Member has had a positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

EYSUVIS

Products Affected

- EYSUVIS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dry Eye Disease (DED)(Initial): (1) Diagnosis of DED
Age Restrictions	
Prescriber Restrictions	(DED)(Initial, Reauth): Prescribed by or in consultation with an ophthalmologist or optometrist.
Coverage Duration	(Initial, Reauth): 14 days
Other Criteria	(DED)(Reauth): (1) Positive clinical response to therapy (e.g. improvement in dry eye symptoms).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FABHALTA

Products Affected

- FABHALTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Paroxysmal Nocturnal Hemoglobinuria (PNH) (Initial): (1) Diagnosis of PNH, (2) Member's Hemoglobin (Hb) level is less than 10 g/dL despite prior eculizumab (Soliris) or ravulizumab-cwvz (Ultomiris) therapy
Age Restrictions	
Prescriber Restrictions	(PNH) (Initial, Continuation): Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	(Initial): 6 months, (Continuation): 12 months
Other Criteria	(PNH) (Continuation): (1) Positive clinical response to therapy (e.g., improvement in hemoglobin level, hemoglobin stabilization, decrease in the number of red blood cell transfusions).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FASENRA

Products Affected

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when there is a documentation of the following: Severe Asthma (SA)(Initial): (1) Diagnosis of severe asthma with eosinophilic phenotype as defined by ONE of the following at initiation of therapy (a) blood eosinophil levels are at least 150 cells/microliter if dependent on daily oral corticosteroids or (b) blood eosinophil levels are at least 300 cells/microliter AND, (2) ONE of the following: (a) Member has had at least one asthma exacerbation requiring systemic corticosteroids within the past 12 months, or (b) Any prior intubation for asthma exacerbation, or (c) prior asthma-related hospitalization within the past 12 months, AND (3) Member is currently treated with ONE of the following unless there is a contraindication or intolerance to these medications: (a) high-dose inhaled corticosteroids (ICS) (e.g. greater than or equal to 500 mcg fluticasone propionate equivalent/day) AND an additional asthma controller medication (e.g. leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline) OR (b) one maximally-dosed combination ICS/LABA product).
Age Restrictions	(SA)(Initial): Member is 6 years of age or older
Prescriber Restrictions	(SA)(Initial): Prescribed by or in consultation with pulmonologist or allergy/immunology specialist
Coverage Duration	(Initial): 12 months. (Reauth): 12 months
Other Criteria	Subject to Part B vs Part D review (SA) (Reauth): (1) Member is currently treated with ONE of the following unless there is a contraindication or intolerance to these medications: (a) high-dose inhaled corticosteroids (ICS) (e.g. greater than or equal to 500 mcg fluticasone propionate equivalent/day) AND an additional asthma controller medication (e.g. leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline) OR (b) one maximally-dosed combination ICS/LABA product). (2) Positive clinical response to therapy (e.g. reduction in exacerbations, decreased use of rescue medications)

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FERRIPROX

Products Affected

- *deferiprone*
- FERRIPROX
- FERRIPROX TWICE-A-DAY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Transfusional Iron Overload (TIO): (1) Diagnosis of transfusional iron overload due to one of the following: (a) Thalassemia syndromes, (b) sickle cell disease, (c) other transfusion-dependent anemias. (2) Inadequate response or inability to tolerate current chelation therapy.
Age Restrictions	(TIO): Member is 3 years of age or older
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FILSPARI

Products Affected

- FILSPARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Immunoglobulin A nephropathy (IgAN)(Initial): (1) Diagnosis of primary immunoglobulin A nephropathy (IgAN) as confirmed by a kidney biopsy (2) Member is at risk of rapid disease progression [e.g., generally a urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g, or by other criteria such as clinical risk scoring using the International IgAN Prediction Tool] (3) Used to reduce proteinuria (4) Member has an estimated glomerular filtration rate (eGFR) of greater than or equal to 30 mL/min/1.75 m ² (5) Member has had an inadequate response or inability to tolerate at minimum 90-day trial of a maximally tolerated dose of one of the following (a) Angiotensin-receptor blockers (ARB) (e.g., losartan, valsartan), (b) Angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril) (6) Medication will not be used in combination with any of the following (a) Angiotensin receptor blockers, (b) Endothelin receptor antagonists (ERAs) (e.g., ambrisentan, bosentan, Opsumit), (c) Aliskiren)
Age Restrictions	(IgAN)(Initial)(Continuation): Member is 18 years of age or older
Prescriber Restrictions	(IgAN)(Initial)(Continuation): Prescribed by or in consultation with a nephrologist
Coverage Duration	(Initial)(Continuation): 12 months
Other Criteria	(IgAN)(Continuation): (1) Documentation of positive clinical response to therapy from baseline as demonstrated by a decrease in urine protein-to-creatinine ration (UPCR).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FILSUEZ 2024

Products Affected

- FILSUEZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(EB)(Initial): (1) Diagnosis of one of the following: (a) Dystrophic epidermolysis bullosa (DEB) (b) Junctional epidermolysis bullosa (JEB) (2) Disease is confirmed by one of the following: (a) genetic testing confirms mutation in one of the following genes: (i) for DEB, collagen type VII (COL7A1) (ii) For JEB, one of the following mutations (ITGA6, ITGB4, collagen type XVII (COL17A1), LAMA3, LAMB3, LAMC2, ITGA3, LAMA3A) (b) skin biopsy (3) Medication is being used for the treatment of wounds (4) Wounds associated with DEB or JEB are present for at least 21 days and less than 9 months old (5) Member does not have signs of infection for wound being treated (6) Member has no evidence or history of basal or squamous cell carcinoma for wound being treated (7) Member does not have history of stem cell transplant or gene therapy (i.e. Vyjuvek) for the treatment of epidermolysis bullosa
Age Restrictions	
Prescriber Restrictions	(EB) (Initial, Reauth): Prescribed by or in consultation with a dermatologist with expertise in the treatment of epidermolysis bullosa.
Coverage Duration	(EB)(Initial): 3 months, (Reauth): 6 months
Other Criteria	(EB)(Reauth): (1) member demonstrates positive clinical response to therapy (2) Member does not have signs of infection for wound being treated (3) Member has no evidence or history of basal squamous cell carcinoma for would being treated
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FINTEPLA

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dravet Syndrome (DS): Inadequate response or inability to tolerate ONE of the following: (a) clobazam, (b) valproic acid, (c) divalproex sodium, (d) topiramate. Lennox-Gastaut Syndrome (LGS): (1) Diagnosis of seizures associated with Lennox-Gastaut Syndrome (LGS) (2) Inadequate response or inability to tolerate ONE of the following: valproic acid, lamotrigine, topiramate, felbamate, clobazam.
Age Restrictions	(DS, LGS): Member is 2 years of age or older.
Prescriber Restrictions	(DS, LGS): Prescribed by or in consultation with a neurologist.
Coverage Duration	Indefinite.
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

FIRDAPSE

Products Affected

- FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	(LEMS)(Initial, Continuation): history of seizures
Required Medical Information	Lambert-Eaton Myasthenic Syndrome (LEMS)(Initial): (1) Diagnosis of LEMS. (2) Neurological symptoms persist after treatment of malignancy, when malignancy is present.
Age Restrictions	(LEMS)(Initial, Continuation): Member is 6 years of age or older
Prescriber Restrictions	(LEMS)(Initial, Continuation): Prescribed by or in consultation with a neurologist.
Coverage Duration	(Initial):90 Days, (Continuation): Indefinite
Other Criteria	(LEMS)(CONTINUATION): positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GALAFOLD

Products Affected

- GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	(FD)(Initial, Reauth): Will not be used in combination with Fabrazyme (agalsidase beta)
Required Medical Information	Fabry Disease (FD)(Initial): (1) Member has amenable galactosidase alpha gene (GLA) variant based on in vitro assay data
Age Restrictions	(FD)(Initial, Reauth): Member is 16 years of age or older
Prescriber Restrictions	(FD)(Initial, Reauth): Prescribed by or in consultation with a clinical genetics specialist OR a nephrologist
Coverage Duration	(Initial): 6 months, (Reauth): Indefinite
Other Criteria	(FD)(Reauth): Positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GATTEX

Products Affected

- GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Short Bowel Syndrome (SBS)(Initial): (1) Diagnosis of Short Bowel Syndrome, (2) individual receives parenteral support at least three times per week for at least 12 months.
Age Restrictions	
Prescriber Restrictions	(SBS): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	(Initial, Reauth): 6 months
Other Criteria	(SBS)(REAUTH): (1)Reduction in parenteral support from baseline (prior to initiation of Gattex therapy)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GOCOVRI

Products Affected

- GOCOVRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dyskinesia in Parkinson's disease (DPD): (1) Diagnosis of PD (2) Member is experiencing dyskinesia. (3) Member is receiving levodopa based therapy (4) Inadequate response or inability to tolerate amantadine immediate-release. Parkinson's Disease with OFF episodes (PD with OFF episodes): (1) Diagnosis of Parkinson's disease. (2) Concurrent use of carbidopa/levodopa containing product. (3) Member is experiencing intermittent OFF episodes. (4) Member had inadequate response or inability to tolerate ONE of the following: (a) MAO-B inhibitor (e.g. rasagiline, selegiline), (b) Dopamine agonist (e.g. pramipexole, ropinirole), (c) COMT inhibitor (e.g. entacapone).
Age Restrictions	
Prescriber Restrictions	(DPD, PD with OFF episodes): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GRALISE

Products Affected

- *gabapentin (once-daily)*
- GRALISE ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Post Herpetic Neuralgia (PHN): (1) Diagnosis of post herpetic neuralgia, (2) Inadequate response to gabapentin or pregabalin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

GROWTH HORMONES

Products Affected

- GENOTROPIN MINIQUICK SUBCUTANEOUS PREFILLED SYRINGE
- GENOTROPIN SUBCUTANEOUS CARTRIDGE
- HUMATROPE INJECTION CARTRIDGE
- NORDITROPIN FLEXPOR SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 10 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 20 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 5 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED
- ZOMACTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(1) Growth Failure in Children (GFC)(Initial): (A) Diagnosis of growth hormone deficiency confirmed by one of the following: (I) Height is documented by one of the following (utilizing age and gender growth charts related to height): (a) Height is greater than 2.0 standard deviations [SD] below midparental height (b) Height is greater than 2.25 SD below population mean (below the 1.2 percentile for age and gender) (II) Growth velocity is greater than 2 SD below mean for age and gender (III) Delayed skeletal maturation of greater than 2 SD below mean for age and gender (eg, delayed is greater than 2 years compared with chronological age), (B) documentation of bone age, (C) abnormal response from provocative testing, such as insulin-induced hypoglycemia test, levodopa, or clonidine. (2) Small for Gestational Age (SGA)(Initial): (A) Diagnosis of SGA, (B) Clinical documentation of no catch-up growth by 2 to 4 years of age. (3) Growth Failure Associated with Chronic Kidney Disease (GF-CKD) (Initial), (4) Growth failure associated with Noonan Syndrome, Prader-Willi Syndrome, Turner Syndrome or short stature homeobox-containing gene (SHOX) deficiency (Initial), (5) Diagnostically confirmed Growth Hormone Deficiency in adults (GHDA)(Initial), OR (6) Idiopathic Short Stature (ISS)(Initial): (A) Diagnosis of ISS defined by height standard deviation score (SDS) less than or equal to -2.25, (B) Documentation of growth velocity less than 25th percentile for bone age.
Age Restrictions	
Prescriber Restrictions	(All Other Indications)(Initial): Prescribed by or in consultation with an endocrinologist. (GF-CKD)(Initial) : Prescribed by or in consultation with an endocrinologist or nephrologist
Coverage Duration	(Initial, Continuation): 12 months

PA Criteria	Criteria Details
Other Criteria	(GHDA)(Continuation): Annual clinical re-evaluation by the treating endocrinologist. (GFC)(Continuation): (1) Annual clinical re-evaluation by the treating endocrinologist, (2) Growth velocity greater than or equal to 2.5cm/year. (SGA, ISS)(Continuation): (1) Increase in growth velocity from baseline, (2) Annual clinical re-evaluation by the treating endocrinologist. (GF-CKD)(Continuation): (1) No history of renal transplant, (2) Annual clinical re-evaluation by the treating endocrinologist. (Growth Failure Associated with Noonan Syndrome, Prader-Willi Syndrome, Turner Syndrome or Short Stature Homeobox-Containing Gene (SHOX) Deficiency)(Continuation): (1) Annual clinical re-evaluation by the treating endocrinologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HAEGARDA

Products Affected

- HAEGARDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hereditary Angioedema (HAE): (1) Diagnosis of hereditary angioedema (HAE), (2) For prophylaxis against HAE attacks.
Age Restrictions	(HAE): Member is 6 years of age or older
Prescriber Restrictions	(HAE): Prescribed by or in consultation with an allergist, immunologist, pulmonologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HETLIOZ

Products Affected

- HETLIOZ
- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Non-24 Hour Sleep-Wake Cycle (Non-24)(Initial): (1) Diagnosis of a circadian period greater than 24 hours (also known as non-24-hour sleep-wake disorder), (2) Member is totally blind (has no light perception). Nighttime Sleep Disturbances in Smith-Magenis Syndrome (SMS)(Initial): (1) Diagnosis of SMS, (2) Member is experiencing nighttime sleep disturbances.
Age Restrictions	(SMS)(Initial): Member is 16 years of age or older
Prescriber Restrictions	(Non-24, SMS)(Initial, Reauth): Prescribed by or in consultation with a sleep specialist or neurologist.
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(Non-24)(Reauth): Documentation of positive clinical response (SMS)(Reauth): Documentation of positive clinical response to therapy (e.g. improvement in nighttime total sleep time, improvement in nighttime sleep quality)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HETLIOZ LQ

Products Affected

- HETLIOZ LQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Nighttime sleep disturbances in Smith-Magenis Syndrome (SMS)(Initial): (1) Diagnosis of SMS, (2) Member is experiencing nighttime sleep disturbances
Age Restrictions	(SMS)(Initial): Member is 3 to 15 years of age
Prescriber Restrictions	(SMS)(Initial, Reauth): Prescribed by or in consultation with a sleep specialist or neurologist.
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(SMS)(Reauth): Documentation of positive clinical response to therapy (e.g. improvement in nighttime total sleep time, improvement in nighttime sleep quality)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH DOSE OPIOIDS

Products Affected

- BELBUCA BUCCAL FILM 300 MCG, 450 MCG, 600 MCG, 750 MCG, 900 MCG
- DILAUDID ORAL TABLET 4 MG, 8 MG
- *fentanyl transdermal patch 72 hour 100 mcg/hr, 25 mcg/hr, 37.5 mcg/hr, 50 mcg/hr, 62.5 mcg/hr, 75 mcg/hr, 87.5 mcg/hr*
- *hydrocodone bitartrate er oral tablet er 24 hour abuse-deterrent*
- *hydromorphone hcl er oral tablet extended release 24 hour*
- *hydromorphone hcl oral tablet 4 mg, 8 mg*
- HYSINGLA ER
- *levorphanol tartrate oral tablet 3 mg*
- *methadone hcl oral solution*
- *methadone hcl oral tablet*
- *morphine sulfate er beads oral capsule extended release 24 hour 120 mg*
- *morphine sulfate er oral capsule extended release 24 hour 100 mg, 60 mg, 80 mg*
- *morphine sulfate er oral tablet extended release 100 mg, 200 mg, 60 mg*
- MS CONTIN ORAL TABLET EXTENDED RELEASE 100 MG, 200 MG, 60 MG
- NUCYNTA ER
- NUCYNTA ORAL TABLET 100 MG, 75 MG
- *oxycodone hcl oral tablet 30 mg*
- OXYCONTIN ORAL TABLET ER 12 HOUR ABUSE-DETERRENT
- *oxymorphone hcl er oral tablet extended release 12 hour 20 mg, 30 mg*
- *oxymorphone hcl er oral tablet extended release 12 hour 40 mg*
- *oxymorphone hcl oral tablet 10 mg*
- ROXICODONE ORAL TABLET 30 MG
- ROXYBOND ORAL TABLET ABUSE-DETERRENT 30 MG
- XTAMPZA ER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NEW TO HIGH DOSE OPIOID THERAPY : ONE of the following (A) pain associated with cancer, (B) chronic non-cancer pain and BOTH of the following (1) member is opioid tolerant (e.g. at least one week where total daily dose is at least one of the following: 30mg of oxycodone, 60mg of oral morphine, 8mg of oral hydromorphone, 25mg of oral oxymorphone or an equianalgesic dose of another opioid) AND (2) the member has been evaluated for non-opioid prescription pharmacologic treatment prior to initiation of high dose opioid therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year

PA Criteria	Criteria Details
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable. CONTINUING HIGH DOSE OPIOID THERAPY (morphine equivalent dose 90mg per day or greater): ONE of the following (1) pain associated with cancer OR (2) chronic non-cancer pain and ALL (a) member's pain has been assessed within the last 6 months AND (b) member has clinically meaningful improvement in pain and functioning that outweighs risks to patient safety AND (c) member is not being treated for substance abuse
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HORIZANT

Products Affected

- HORIZANT ORAL TABLET EXTENDED RELEASE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Postherpetic neuralgia (PHN): (1) Diagnosis of PHN and (2) Inadequate response or inability to tolerate gabapentin or pregabalin. Restless legs syndrome (RLS): (1) Diagnosis of RLS and (2) Inadequate response or inability to tolerate pramipexole or ropinirole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HRM

Products Affected

- ALLZITAL
- ASCOMP-CODEINE
- *butalbital-acetaminophen oral capsule*
- *butalbital-acetaminophen oral tablet 50-300 mg, 50-325 mg*
- *butalbital-apap-caff-cod*
- *butalbital-apap-caffeine oral capsule*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*
- *carbinoxamine maleate oral solution*
- *carbinoxamine maleate oral tablet*
- *chlordiazepoxide-clidinium*
- *chlorzoxazone oral*
- *clemastine fumarate oral syrup*
- *clemastine fumarate oral tablet 2.68 mg*
- DEMEROL INJECTION SOLUTION 25 MG/ML, 50 MG/ML
- *dipyridamole oral*
- ESGIC ORAL TABLET
- FIORICET ORAL CAPSULE
- FIORICET/CODEINE ORAL CAPSULE 50-300-40-30 MG
- INDOCIN ORAL
- INDOCIN RECTAL
- *indomethacin oral suspension*
- *indomethacin rectal suppository 50 mg*
- LIBRAX
- *meperidine hcl injection solution 100 mg/ml, 25 mg/ml, 50 mg/ml*
- *meperidine hcl oral solution*
- *meperidine hcl oral tablet 50 mg*
- *meprobamate*
- *metaxalone*
- *orphenadrine citrate er*
- *pentazocine-naloxone hcl*
- *promethazine hcl oral solution*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal suppository 12.5 mg, 25 mg*
- *promethazine vc*
- PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG
- RYVENT
- TENCON ORAL TABLET 50-325 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly
Age Restrictions	Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	2 years
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

HRM ESTROGENS

Products Affected

- ACTIVELLA ORAL TABLET 1-0.5 MG
- ANGELIQ
- BIJUVA
- CLIMARA
- CLIMARA PRO
- COMBIPATCH
- DIVIGEL
- DOTTI
- ELESTRIN
- ESTRACE ORAL
- *estradiol transdermal*
- *estradiol-norethindrone acet*
- ESTROGEL
- EVAMIST
- FYAVOLV
- JINTELI
- LYLLANA
- MENOSTAR
- MIMVEY
- MINIVELLE
- *norethindrone-eth estradiol*
- PREMARIN ORAL
- VIVELLE-DOT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ONE of the following: (1) Inadequate response or inability to tolerate vaginal estrogen preparations (e.g. vaginal tablets, rings, or cream, etc.) OR (2) Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly
Age Restrictions	Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	2 years
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HRM KETOROLAC

Products Affected

- *ketorolac tromethamine oral*
- SPRIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ONE of the following: (1) Inadequate response or inability to tolerate TWO alternative NSAIDs such as meloxicam, naproxen, celecoxib, ibuprofen, etc. OR (2) Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly
Age Restrictions	Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	One Month
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HRM NON BENZODIAZEPINE HYPNOTICS

Products Affected

- AMBIEN CR ORAL TABLET EXTENDED RELEASE 12.5 MG
- AMBIEN ORAL TABLET 10 MG
- EDLUAR SUBLINGUAL TABLET SUBLINGUAL 10 MG
- *eszopiclone oral tablet 3 mg*
- LUNESTA ORAL TABLET 3 MG
- *zolpidem tartrate er oral tablet extended release 12.5 mg*
- *zolpidem tartrate oral capsule*
- *zolpidem tartrate oral tablet 10 mg*
- *zolpidem tartrate sublingual tablet sublingual 3.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Initial): Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly
Age Restrictions	(Initial, Reauth): Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	(Initial): 3 months. (Reauth): 2 years
Other Criteria	(REAUTH): Prescriber is aware of the risk versus benefit of using the medication chronically (greater than 90 days)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HRM NORGESIC

Products Affected

- NORGESIC
- *norgesic forte*
- *orphenadrine-aspirin-caffeine oral tablet 25-385-30 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(1) For age 65 and older: Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly. (2) Inadequate response or inability to tolerate two generic skeletal muscle relaxants (e.g., tizanidine, chlorzoxazone 500mg, etc.)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	2 years
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HRM SHORT TERM SKELETAL MUSCLE RELAXANTS

Products Affected

- AMRIX
- *carisoprodol oral*
- *cyclobenzaprine hcl er*
- *cyclobenzaprine hcl oral*
- FEXMID
- SOMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Acute Muscle Spasms (AMS): Prescriber attestation that drug will be used only for short periods (up to 2 or 3 weeks). All Indications: Risk versus benefit has been assessed for this request of a high risk medication (HRM) in elderly.
Age Restrictions	Apply if member is greater than or equal to 65 years
Prescriber Restrictions	
Coverage Duration	(AMS): 1 year. (All other indications): 2 years
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HUMIRA

Products Affected

- HUMIRA (2 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA-CD/UC/HS STARTER SUBCUTANEOUS AUTO-INJECTOR KIT 80 MG/0.8ML
- HUMIRA-PSORIASIS/UVEIT STARTER SUBCUTANEOUS AUTO-INJECTOR KIT

PA Criteria	Criteria Details
Exclusion Criteria	(RA, AS, JIA, PsA, PsO, CD, UC, HS, UV): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate one other treatment such as NSAIDs, COX2 inhibitors or methotrexate. Juvenile Idiopathic Arthritis (JIA): (1) Diagnosis of moderate to severe JIA. (2) Inadequate response or inability to tolerate one of the following: methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe chronic PsO.
Age Restrictions	(JIA, UV): Member is 2 years of age or older. (CD): Member is 6 years of age or older. (HS): Member is 12 years of age or older. (RA, AS, PsA, PsO): Member is 18 years of age or older. (UC): Member is 5 years of age or older.
Prescriber Restrictions	(RA, JIA, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist. (PsO, HS): Prescribed by or in consultation with a dermatologist. (CD, UC): Prescribed by or in consultation with a gastroenterologist. (UV): Prescribed by or in consultation with a rheumatologist or ophthalmologist.
Coverage Duration	Indefinite

PA Criteria	Criteria Details
Other Criteria	Crohn's Disease (CD): (1) Diagnosis of CD. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. corticosteroid, methotrexate, azathioprine or 6-mercaptopurine). Ulcerative Colitis (UC): (1) Diagnosis of UC. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. corticosteroid, Aminosalicylate, azathioprine or 6-mercaptopurine). Hidradenitis Suppurativa (HS): (1) Diagnosis of HS. Uveitis (UV)(Initial): (1) Diagnosis of non-infectious intermediate, posterior, or pan-uveitis. (2) Inadequate response or inability to tolerate corticosteroids and at least one immunosuppressive drug (e.g. azathioprine, cyclosporine, methotrexate, mycophenolate).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HUMULIN U500

Products Affected

- HUMULIN R U-500 (CONCENTRATED)
- HUMULIN R U-500 KWIKPEN SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Type 1 Diabetes (T1DM): (1) Diagnosis of type 1 diabetes mellitus (2) Insulin requirement exceeding 200 units per day. Type 2 Diabetes (T1DM): (1) Diagnosis of Type 2 diabetes mellitus (2) Insulin requirement exceeding 200 units per day.
Age Restrictions	
Prescriber Restrictions	(T1DM, T1DM2): Prescribed by or in consultation with an endocrinologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

HYFTOR

Products Affected

- HYFTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Facial Angiofibroma (FA)(Initial): (1) Diagnosis of facial angiofibroma associated with tuberous sclerosis complex
Age Restrictions	(FA) (Initial, Reauth): Member is 6 years of age or older
Prescriber Restrictions	(FA) (Initial, Reauth): Prescribed by or in consultation with a dermatologist, neurologist, or geneticist.
Coverage Duration	(Initial): 6 months, (Reauth): Indefinite
Other Criteria	(FA) (Reauth): (1) Documentation of positive clinical response to therapy (e.g., improvement in size or redness of facial angiofibroma)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ILUMYA

Products Affected

- ILUMYA

PA Criteria	Criteria Details
Exclusion Criteria	(PsO): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Skyrizi, (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(PsO): Member is 18 years of age or older.
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INBRIJA

Products Affected

- INBRIJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Parkinson's Disease (PD): (1) Diagnosis of Parkinson's disease. (2) concurrent use of carbidopa/levodopa containing product, (3) Member is experiencing intermittent OFF episodes, (4) member had inadequate response or inability to tolerate ONE of the following: (a) MAO-B inhibitor (e.g. rasagiline, selegiline), (b) Dopamine agonist (e.g. pramipexole, ropinirole), (c) COMT inhibitor (e.g. entacapone)
Age Restrictions	
Prescriber Restrictions	(PD): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INCRELEX

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	(GHGD, PIGF-1D) (Initial and Continuation): Known or suspected malignancy, closed epiphyses, concurrent GH therapy
Required Medical Information	Growth Hormone Gene Deletion (GHGD): (Initial) (1) Diagnosis of growth hormone gene deletion who have developed neutralizing antibodies to growth hormone. Severe Primary IGF-1 Deficiency (PIGF-1D): (Initial) (1) Diagnosis of Severe primary IGF-1 deficiency (2) height standard deviation score less than or equal to -3.0 (3) basal IGF-1 standard deviation score less than or equal to -3.0 AND normal or elevated growth hormone.
Age Restrictions	(GHGD, PIGF-1D) (Initial and Continuation): Member is 2 years of age or older
Prescriber Restrictions	(GHGD, PIGF-1D) (Initial and Continuation) Prescribed by or in consultation with an endocrinologist
Coverage Duration	(Initial and continuation): 12 months
Other Criteria	(GHGD, PIGF-1D)(CONTINUATION): (1) Documentation of increase in growth velocity from baseline AND (2) Annual clinical re-evaluation by an endocrinologist AND (3) BOTH of the following: (a) Expected adult height is not obtained (b) documentation of expected adult height goal
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INGREZZA

Products Affected

- INGREZZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Tardive Dyskinesia (TD) (Initial): (1) Diagnosis of moderate to severe tardive dyskinesia (2) documentation of baseline AIMS and (3) one of the following: (a) persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication, or (b) member is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Chorea- Huntington's Disease (CHD): (1) Diagnosis of chorea associated with Huntington's disease
Age Restrictions	
Prescriber Restrictions	(TD) (Initial, Reauth): Prescribed by or in consultation with a neurologist or a psychiatrist. (CHD): Prescribed by or in consultation with a neurologist or a psychiatrist.
Coverage Duration	(TD) (Initial): 3 months (TD)(Reauth): indefinite (CHD): Indefinite
Other Criteria	(TD) (Reauthorization): Valbenazine (Ingrezza) is reapproved with documentation of positive clinical response, as demonstrated by improvement in AIMS, to Ingrezza therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INHALED TOBRAMYCIN

Products Affected

- TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis (CF): (1) Diagnosis of cystic fibrosis AND (2) evidence of P. aeruginosa in the lungs (3) FEV1 between 25% and 80%, (4) Member not colonized with Burkholderia cepacia.
Age Restrictions	(CF): Member is 6 years of age or older
Prescriber Restrictions	(CF): Prescribed by or in consultation with a pulmonologist, infectious disease specialist or Specialist affiliated with a CF care center.
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INJECTABLE METHOTREXATE

Products Affected

- OTREXUP SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.4ML, 12.5 MG/0.4ML, 15 MG/0.4ML, 17.5 MG/0.4ML, 20 MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML
- RASUVO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Rheumatoid arthritis (RA): (1) Diagnosis of severe, active rheumatoid arthritis (RA), (2) Inadequate response or inability to tolerate oral methotrexate. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA, (2) Inadequate response or inability to tolerate oral methotrexate. Polyarticular juvenile idiopathic arthritis (pJIA): (1) Diagnosis of pJIA, (2) Inadequate response or inability to tolerate oral methotrexate. Psoriasis: (1) Diagnosis of severe psoriasis, (2) Inadequate response to BOTH of the following: (a) oral methotrexate AND (b) topical steroids (e.g. clobetasol propionate cream/ointment, betamethasone cream/ointment, etc.).
Age Restrictions	(Psoriasis): Member is 18 years of age or older
Prescriber Restrictions	(RA, pJIA): Recommended by rheumatologist. (PSA): Recommended by a rheumatologist or dermatologist. (Psoriasis): Recommended by dermatologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INSULIN GLARGINE

Products Affected

- *insulin glargine max solostar*
- *insulin glargine solostar subcutaneous solution pen-injector 300 unit/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(DM): (1) Diagnosis of Diabetes Mellitus (2) inadequate response or inability to tolerate two of the following: (a) brand Lantus (b) Levemir (c) Toujeo (d) Tresiba (3) Documentation provided stating that the Brand products has not been effective (4) Justification provided for why the target drug is expected to provide benefit when the Brand products have not been shown to be effective
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

INTRAVENOUS IMMUNE GLOBULINS (IVIG)

Products Affected

- BIVIGAM INTRAVENOUS SOLUTION 5 GM/50ML
- GAMMAGARD INJECTION SOLUTION 2.5 GM/25ML
- GAMMAGARD S/D LESS IGA
- GAMMAKED INJECTION SOLUTION 1 GM/10ML
- GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML
- GAMUNEX-C INJECTION SOLUTION 1 GM/10ML
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 2 GM/20ML
- PANZYGA
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML

PA Criteria	Criteria Details
Exclusion Criteria	(Initial): History of hypersensitivity (including anaphylaxis or severe systemic reaction) to immune globulin or any component of the preparation.
Required Medical Information	Part D is medically necessary when ONE of the following is present: (1) Autoimmune mucocutaneous blistering disease pemphigus vulgaris, pemphigus foliaceus, bullous pemphigoid, mucous membrane (cicatricial) pemphigoid, benign mucous membrane pemphigoid, epidermolysis bullosa acquisita and ONE of the following: (a) inadequate response or inability to tolerate conventional therapy (i.e. steroids, immunosuppressants) OR (b) rapidly progressive disease in conjunction with conventional therapy (i.e. steroids, immunosuppressants). (2) Acute Idiopathic Thrombocytopenia Purpura (ITP) and ONE of the following (a) management of acute bleeding (b) used to increase platelet count prior to surgical procedures (c) severe thrombocytopenia (platelets less than 20,000 per uL) or (d) high risk for intracerebral hemorrhage. (3) Chronic ITP and ALL of the following (a) inadequate response or inability to tolerate corticosteroids (b) duration of illness greater than 6 months (c) platelets persistently less than 20,000/ uL. (4) Chronic B-cell lymphocytic leukemia with IgG less than 600mg/dL and recurrent, serious bacterial infections requiring antibiotic therapy. (5) Hematopoietic stem cell transplant and IgG less than 400mg/dL. (6) HIV and all of the following (a) less than 14 years of age (b) evidence of qualitative or quantitative humoral immunologic defects and (c) current bacterial infection despite antimicrobial prophylaxis. (7) Solid organ transplant. (8) Chronic Inflammatory Demyelinating Polyneuritis confirmed by electrodiagnostic testing or nerve biopsy and an inadequate response or inability to tolerate corticosteroids. (9) Dermatomyositis or Polymyositis diagnosed by laboratory testing (antinuclear or myositis specific antibodies), biopsy, EMG, or MRI) and inadequate response or inability to tolerate steroids or immunosuppressants. (10) Guillain Barre syndrome with impaired function (i.e. unable to stand or walk without aid).
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g. immunologist, hematologist, neurologist).
Coverage Duration	(All Indications): 6 months
Other Criteria	Subject to Part B vs Part D review. (11) Lambert Eaton myasthenic syndrome and an inadequate response or inability to tolerate steroids, immunosuppressants, or cholinesterase inhibitors (12) Multifocal motor neuropathy diagnosed by electrodiagnostic studies. (13) Acute exacerbations of multiple sclerosis (MS) unresponsive to steroids. (14) Myasthenia gravis and an inadequate response or inability to tolerate at least 8 weeks of standard therapy (e.g., corticosteroids, azathioprine, cyclosporine, cyclophosphamide, cholinesterase inhibitors) (15) Myasthenic crisis (16) Stiff person syndrome and an inadequate response or inability to tolerate standard therapy (e.g., muscle relaxants, benzodiazepines, and gabapentin-related medications) (17) Severe, active SLE and an inadequate response or inability to tolerate steroids (18) Kawasaki disease. (All Indications)(CONTINUATION): (1) Documentation of clinical improvement as appropriate to the diagnosis (e.g. clinical improvement in symptoms, Rankin score and Activities of Daily Living (ADL) scores, etc.)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

IQIRVO 2024

Products Affected

- IQIRVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Primary biliary cholangitis (PBC)(Initial): (1) Diagnosis of primary biliary cholangitis (PBC) (also known as primary biliary cirrhosis) (2) One of the following: (a) Both of the following: (i) Member has failed to achieve an alkaline phosphatase (ALP) level of less than 1.67 times the upper limit of normal (ULN) after treatment with ursodeoxycholic acid (UDCA) (e.g., Urso, Urso Forte, ursodiol) (ii) Used in combination with ursodeoxycholic acid (UDCA) (b) History of contraindication or intolerance to ursodeoxycholic acid (UDCA)
Age Restrictions	
Prescriber Restrictions	(PBC)(Initial, Reauth): Prescribed by or in consultation with hepatologist or gastroenterologist
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(PBC)(Reauth): (1) Member demonstrates positive clinical response to therapy (e.g., ALP level less than 1.67 times ULN, total bilirubin less than or equal to ULN, ALP decrease greater than or equal to 15 percent from baseline).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ISTURISA

Products Affected

- ISTURISA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cushing's disease (CD): (1) Diagnosis of (pituitary) Cushing's disease. (2) Pituitary surgery is not an option or has not been curative (3) Member has inadequate response or inability to tolerate Signifor [LAR].
Age Restrictions	(CD): Member is 18 years of age or older
Prescriber Restrictions	(CD): Prescribed by or in consultation with an endocrinologist
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

JOENJA

Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(APDS)(Initial): (1) Diagnosis of activated phosphoinositide 3-kinase delta syndrome (APDS) (2) Molecular genetic testing confirms mutations in the PIK3CD or PIK3R1 gene (3) Member weighs greater than or equal to 45kg (4) Both of the following (a) Presence of nodal and/or extranodal proliferation (e.g., lymphadenopathy, splenomegaly, hepatomegaly) (b) Presence of other clinical findings and manifestations consistent with APDS (e.g., recurrent sino-pulmonary infections, bronchiectasis, enteropathy) (5) Inadequate response or inability to tolerate at least one standard of care treatment for APDS (e.g., Immunoglobulin replacement therapy)
Age Restrictions	(APDS)(Initial)(Reauth): Member is 12 years of age or older
Prescriber Restrictions	(APDS)(Initial)(Reauth): Prescribed by or in consultation with hematologist, geneticist or immunologist
Coverage Duration	(Initial): 6 months. (Reauth): 12 months
Other Criteria	(APDS)(Reauth): (1) Documentation of positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

JUXTAPID

Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	(HoFH)(Initial, Reauth): Moderate-severe liver impairment, active liver disease, unexplained LFT abnormalities. Juxtapid: Pregnancy, concomitant use with strong or moderate CYP3A4 inhibitors.
Required Medical Information	Homozygous Familial Hypercholesterolemia (HoFH): (1) Diagnosis of HoFH with ONE of the following: (a) Genetic confirmation of 2 mutant alleles at the LDL receptor. Apo B, PCSK9, or ARH adaptor protein gene locus (b) Untreated LDL-C greater than 500 mg/dL or treated LDL cholesterol greater than or equal to 300 mg/dL or treated non-HDL cholesterol greater than or equal to 330 mg/dL together with either of the following: (i) Tendinous xanthoma prior to 10 years of age (ii) Elevated LDL cholesterol prior to lipid-lowering therapy consistent with HeFH in both parents AND (2) Inadequate response or inability to tolerate the combination of BOTH of the following: (a) Either ezetimibe or a Bile Acid Sequestrant (e.g. cholestyramine) AND (b) ONE of the following: (i) ONE high potency statin at the maximally tolerated dose (e.g. atorvastatin, rosuvastatin) OR (ii) Inability to tolerate statin therapy as determined by one of the following: (A) member had rhabdomyolysis or symptoms with creatine kinase (CK) exceeding 10 times the upper limit of normal (ULN) or (B) either of the following with TWO statins: myalgia (no CK elevation) or myositis (CK less than 10 times ULN or (C) hepatotoxicity from statin use (increased AST/ALT exceeding 3 times ULN) or (D) liver disease documented by Child Pugh A or worse OR AST/ALT exceeding 3 times ULN for at least 6 weeks. (3) Inadequate response or inability to tolerate BOTH of the following: (a) Repatha and (b) Praluent.
Age Restrictions	
Prescriber Restrictions	(HoFH)(Initial, Reauth): Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.
Coverage Duration	(Initial, Reauth): 6 months
Other Criteria	(HoFH)(REAUTH): (1) Documentation of reduction in LDL level since initiation of therapy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

JYNARQUE

Products Affected

- JYNARQUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Initial): Baseline serum transaminases and bilirubin prior to initiation of therapy
Age Restrictions	(Initial and Reauth): Member is 18 years of age or older
Prescriber Restrictions	(Initial and Reauth): Prescribed by or in consultation with a nephrologist or kidney transplant specialist
Coverage Duration	(Initial): 3 months. (Reauth): 12 months.
Other Criteria	(REAUTH): (1) ONE of the following (a) decline in kidney function has slowed or (b) kidney pain has improved and (2) serum transaminase less than 3 times the upper limit of normal and (3) bilirubin less than 2 times upper limit of normal
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KALYDECO

Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis (CF): (1) Diagnosis of Cystic Fibrosis with documentation of one mutation in the CFTR gene that is responsive to ivacaftor based on clinical and/or in vitro assay data, (2) Mutation was documented by an FDA-cleared CF mutation test to detect the presence of the CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions.
Age Restrictions	(CF): Member is 1 month of age or older for granules. Member is 6 years of age or older for tablets
Prescriber Restrictions	(CF): Prescribed by or in consultation with is a pulmonologist or Specialist affiliated with a CF care center
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KERENDIA

Products Affected

- KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic kidney disease associated with type 2 diabetes (CKD with T2D): (1) Diagnosis of chronic kidney disease (CKD) associated with type 2 diabetes (T2D), (2) One of the following (a) Minimum 30-day trial of a maximally tolerated dose and member will continue therapy with ONE of the following (i) generic angiotensin-converting enzyme (ACE) inhibitor (e.g. benazepril, lisinopril), (ii) generic angiotensin II receptor blocker (ARB) (e.g. losartan, valsartan), (b) Member has contraindication or intolerance to ACE inhibitors or ARBs
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KEVEYIS

Products Affected

- KEVEYIS
- ORMALVI

PA Criteria	Criteria Details
Exclusion Criteria	(Initial, Reauth): Concomitant use of high dose Aspirin, severe pulmonary disease and hepatic insufficiency
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	(Initial, Reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	(Initial): 3 months (Reauth): Indefinite
Other Criteria	(Reauth): Member has had a positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KEVZARA

Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	(RA, PMR, PJIA): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Rheumatoid arthritis (RA): (1) Diagnosis of RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) Rinvoq, (d)Xeljanz/Xeljanz XR, (e) Orencia, (f) adalimumab-adbm (Cyltezo), (g) adalimumab-aaty (Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Polymyalgia Rheumatica (PMR): (1) Diagnosis of polymyalgia rheumatica (2) Inadequate response or inability to tolerate corticosteroids (e.g., prednisone). Polyarticular Juvenile Idiopathic Arthritis (PJIA): (1) Diagnosis of active polyarticular juvenile idiopathic arthritis (PJIA) (2) Member weighs at least 63 kg (3) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Xeljanz, (d) Orencia, (e) Cyltezo, (f) Yuflyma, (g) Rinvoq/Rinvoq LQ OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(RA, PMR): Member is 18 years of age or older
Prescriber Restrictions	(RA, PMR, PJIA): Prescribed by or in consultation with a rheumatologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KINERET

Products Affected

- KINERET SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	(RA, NOMID, DIRA): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Rheumatoid Arthritis (RA): (1) Diagnosis of RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) adalimumab-adbm (Cyltezo), (g)adalimumab-aaty (Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Neonatal-Onset Multisystem Inflammatory Disease (NOMID): (1) Diagnosis of NOMID. (2) Diagnosis has been confirmed by one of the following: (a) NLRP-3 (nucleotide-binding domain, leucine rich family (NLR), pyrin domain containing 3] gene (also known as Cold-Induced Auto-inflammatory Syndrome-1 [CIAS1]) mutation (b) Evidence of active inflammation including both of the following: (i) clinical symptoms (e.g. rash, fever, arthralgia) (ii) elevated acute phase reactants (e.g. ESR, CRP). Deficiency of Interleukin-1 Receptor Antagonist (DIRA): (1) Diagnosis of DIRA.
Age Restrictions	
Prescriber Restrictions	(RA, NOMID, DIRA): Prescribed by or in consultation with a rheumatologist or pediatric specialist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

KORLYM

Products Affected

- KORLYM
- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	(HCS): Pregnancy
Required Medical Information	Hyperglycemia in members with Cushing Syndrome (HCS): (1) Hyperglycemia secondary to hypercortisolism in adult member with endogenous Cushing syndrome, (2) member has type 2 diabetes mellitus or glucose intolerance, (3) member has failed surgery or is not a candidate for surgery.
Age Restrictions	
Prescriber Restrictions	(HCS): Prescribed by or in consultation with an endocrinologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LETAIRIS

Products Affected

- *ambrisentan*
- LETAIRIS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (Initial): (1) Documentation of a diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV. (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography. (3) Documentation of mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion. (4) Inadequate response or inability to tolerate ambrisentan (applies to brand Letairis only)
Age Restrictions	
Prescriber Restrictions	(PAH) (Initial): Prescribed by or in consultation with a Cardiologist or Pulmonologist.
Coverage Duration	(Initial): 6 months. (Continuation):12 months.
Other Criteria	(PAH)(Continuation): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE TRANSDERMAL PATCH

Products Affected

- *lidocaine external patch 5 %*
- LIDOCAN
- LIDODERM
- TRIDACAINE II

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Post-Herpetic Neuralgia (PHN): (1) Diagnosis of post-herpetic neuralgia. Diabetic Peripheral Neuropathy (DPN): (1) Diagnosis of diabetic peripheral neuropathy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LITFULO

Products Affected

- LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	(Alopecia Areata): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g., tumor necrosis factor antagonists JAK inhibitors, or potent immunosuppressants such as azathioprine, cyclosporine
Required Medical Information	Alopecia Areata (AA): (1) Diagnosis of alopecia areata (2) Member has at least 50% scalp hair loss (3) Other causes of hair loss have been ruled out (e.g., androgenetic alopecia, trichotillomania, tinea capitis, psoriasis) (4) Inadequate response or inability to tolerate one previous treatment for alopecia areata (e.g., topical, intralesional, or systemic corticosteroids, topical immunotherapy)
Age Restrictions	(Alopecia Areata): Member is 12 years of age or older
Prescriber Restrictions	(AA): Prescribed by or in consultation with a dermatologist
Coverage Duration	(AA): Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIVMARLI

Products Affected

- LIVMARLI ORAL SOLUTION 9.5 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cholestatic Pruritus with Alagille Syndrome (CPALGS) (Initial): (1) Diagnosis of Cholestatic Pruritus with Alagille Syndrome. (2) Diagnosis of ALGS confirmed by one of the following: (A) liver histology showing bile duct scarcity, OR (B) THREE of the following clinical features (i) hepatic: cholestasis, jaundice, hepatomegaly (ii) Facial: high prominent forehead, pointed chin, deep-set eyes (iii) Ocular: posterior embryotoxon, optic disc drusen (iv) Cardiac: pulmonary stenosis, tetralogy of Fallot (v) Skeletal: butterfly vertebrae, pathological fractures (vi) Renal: renal dysplasia, renal tubular acidosis (vii) Vascular: intracranial bleeding, CNS/pulmonary vascular malformations, OR (C) genetic mutation in the JAG1 or NOTCH2 genes. (3) An inadequate response or inability to tolerate at least two of the following treatments used for the relief of pruritus: (a) Ursodeoxycholic acid (e.g. Ursodiol) (b) Antihistamines (e.g. diphenhydramine, hydroxyzine) (c) Rifampin (d) Bile acid sequestrants (e.g. Questran).
Age Restrictions	(CPALGS): Member is 3 months of age or older(PFIC): Member is 5 years of age or older
Prescriber Restrictions	(CPALGS, PFIC) (Initial, Reauth): Prescribed by or in consultation with a gastroenterologist or hepatologist.
Coverage Duration	(CPALGS, PFIC)(Initial): 6 months (CPALGS, PFIC)(Reauth): End of contract year
Other Criteria	Progressive Familial Intrahepatic Cholestasis (PFIC) (Initial): (1) BOTH of the following: (A) Diagnosis of Progressive familial intrahepatic cholestasis (PFIC) (B) Molecular genetic testing confirms mutations in the ATP8B1, ABCB11, ABCB4, TJP2, NR1H4, or MYO5B gene (2) Member is experiencing moderate to severe cholestatic pruritus (3) Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus: (i) Ursodeoxycholic acid (e.g., Ursodiol) (ii) Antihistamines (e.g., diphenhydramine, hydroxyzine) (iii) Rifampin (iv) Bile acid sequestrants (e.g., Questran) (CPALGS, PFIC) (Reauth): Positive clinical response to therapy (e.g. reduced bile acids, reduction in pruritis severity score).

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIVTENCITY

Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cytomegalovirus (CMV): (1) Diagnosis of cytomegalovirus (CMV) infection/disease, (2) Member is a recipient of one of the following (a) Hematopoietic stem cell transplant, (b) solid organ transplant, (3) Inadequate response to a minimum 2 weeks duration or inability to tolerate one of the following therapies at an appropriately indicated dose (a) intravenous (IV) ganciclovir, (b) oral valganciclovir, (c) intravenous (IV) foscarnet, (d) intravenous (IV) cidofovir, (4) Member weighs greater than or equal to 35kg
Age Restrictions	(CMV): Member is 12 years of age or older
Prescriber Restrictions	(CMV): Prescribed by or in consultation with a provider who specializes in one of the following areas (1) transplant, (2) infectious disease
Coverage Duration	8 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LODOCO

Products Affected

- LODOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(CV)(Initial): (1) Diagnosis of cardiovascular disease (CV) (2) Used for the secondary prevention of CV disease (e.g., very high-risk patients) (3) Member is on guideline therapy management for multiple risk factors (e.g., dyslipidemia, hypertension, hyperglycemia) associated with CV disease
Age Restrictions	(CV)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): 6 months
Other Criteria	(CV)(Reauth): (1) Documentation of positive clinical response is provided
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LUMRYZ

Products Affected

- LUMRYZ

PA Criteria	Criteria Details
Exclusion Criteria	CN, EDSN (Initial, Reauth): Concurrent use of sedative hypnotics and alcohol
Required Medical Information	Cataplexy in Narcolepsy (CN)(Initial): (1) Diagnosis of cataplexy in narcolepsy, (2) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT) or prescriber provides justification that a sleep study is not feasible. Excessive Daytime Sleepiness in Narcolepsy (EDSN)(Initial): (1) Diagnosis of EDSN (Narcolepsy Type 2), (2) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT) or prescriber provides justification that a sleep study is not feasible and (3) Inadequate response or inability to modafinil or armodafinil.
Age Restrictions	(CN, EDSN)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(CN, EDSN)(Initial, Reauth): Prescribed by or in consultation with a neurologist, psychiatrist, or sleep specialist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(CN, EDSN)(Reauth): (1) Documentation to support the efficacy associated with the current regimen (including but not limited to reduction in the frequency of cataplexy attacks or an improvement in the Epworth sleepiness scale), (2) Member is re-evaluated every 12 months.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LYRICA CR

Products Affected

- LYRICA CR
- *pregabalin er*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Neuropathic Pain Associated with Diabetic Peripheral Neuropathy (DPN): (1) Diagnosis of neuropathic pain associated with diabetic peripheral neuropathy, (2) inadequate response or inability to tolerate gabapentin or pregabalin. Post-Herpetic Neuralgia (PHN): (1) Diagnosis of post-herpetic neuralgia (2) inadequate response or inability to tolerate gabapentin or pregabalin.
Age Restrictions	(DPN, PHN): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

METFORMIN IR

Products Affected

- *metformin hcl oral tablet 625 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(DM) (Initial): (1) Diagnosis of Type 2 Diabetes Mellitus (DM2) (2) Inadequate response to at least a 12-week trial of generic metformin 500mg, metformin 850mg, or metformin 1000mg as evidenced by Hemoglobin A1C level above the member's goal, or inability to tolerate generic metformin 500mg, metformin 850mg, or metformin 1000mg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(DM) (Reauth): (1) Member has experienced an objective response to therapy demonstrated by an improvement in HbA1c from baseline
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

METHOCARBAMOL 1000MG TAB

Products Affected

- *methocarbamol oral tablet 1000 mg*
- TANLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Inadequate response or inability to tolerate two generic skeletal muscle relaxants (e.g., tizanidine, chlorzoxazone 500mg, etc.)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MODAFINIL

Products Affected

- *modafinil oral*
- PROVIGIL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>NARCOLEPSY: (1) Diagnosis of Narcolepsy confirmed by a sleep study (unless prescriber provides justification that a sleep study is not feasible).</p> <p>OBSTRUCTIVE SLEEP APNEA/HYPOPNEA SYNDROME (OSAHS): (1) Diagnosis confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). SHIFT WORK SLEEP DISORDER (SWSD): (1) One of the following: (a) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR (b) A sleep study demonstrating loss of a normal sleep-wake pattern (i.e., disturbed chronobiologic rhythmicity). (2) Documentation that the member has no other medical or mental disorder accounting for the symptoms. Multiple Sclerosis (MS) Related Fatigue: (1) Diagnosis of Multiple Sclerosis (MS) related fatigue, (2) Used in combination with standard educational therapies (e.g. psychoeducation, behavioral programs, scheduled naps, additional non-pharmacological therapies, etc.) (All Indications): Inadequate response or inability to tolerate generic modafinil (applies to Brand Provigil)</p>
Age Restrictions	
Prescriber Restrictions	(Narcolepsy, SWSD, OSAHS): Prescribed by or in consultation with a neurologist or sleep specialist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MULPLETA

Products Affected

- MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic Liver Disease (CLD): (1) Member is scheduled to undergo a procedure (2) Documentation of baseline platelet count less than 50,000/mcL
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MYALEPT

Products Affected

- MYALEPT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Generalized Lipodystrophy (GL): (1) Diagnosis of congenital or acquired generalized lipodystrophy.
Age Restrictions	
Prescriber Restrictions	(GL): Prescribed by or in consultation with an endocrinologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MYCAPSSA

Products Affected

- MYCAPSSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Acromegaly)(Initial): (1) One of the following: (a) Inadequate response to surgical resection and/or pituitary irradiation, (b) member is not a candidate for surgical resection or pituitary irradiation (2) Inadequate response or inability to tolerate a dopamine agonist (e.g. bromocriptine or cabergoline) at maximally tolerated doses (3) Member has responded to and tolerated treatment with octreotide or lanreotide.
Age Restrictions	(Acromegaly): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(Acromegaly) (Reauth): Positive clinical response to therapy (e.g. reduction or normalization of IGF-1/GH level for same age and sex)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

MYFEMBREE

Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Uterine Leiomyomas (UL)(Initial): (1) Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas (fibroids), (2) Member is premenopausal, (3) ONE of the following: (a) Inadequate response or inability to tolerate one of the following for no more than 30 days: combination (estrogen/progestin) contraceptive, progestins, tranexamic acid, OR (b) Member has had a previous interventional therapy to reduce bleeding (e.g. uterine-artery embolization and magnetic resonance-guided focused ultrasonography), (4) Treatment duration of therapy has not exceeded a total of 24 months.</p> <p>(Endometriosis)(Initial) (1) Diagnosis of moderate to severe pain associated with endometriosis (2) member is premenopausal (3) One of the following (a) Inadequate response or inability to tolerate one of the following for at least 3 months (danazol, combination (estrogen/progestin) contraceptive, progestins (b) member has had surgical ablation to prevent recurrence (4) Treatment duration of therapy has not exceeded a total of 24 months</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	<p>(UL) (Reauth): (1) Member has improvement in bleeding associated with uterine leiomyomas (fibroids) (e.g. significant/sustained reduction in menstrual blood loss per cycle, improved quality of life, etc.), (2) Treatment duration of therapy has not exceeded a total of 24 months.</p> <p>(Endometriosis)(Reauth) (1) Member has improvement in pain associated with endometriosis (e.g., improvement in dysmenorrhea and nonmenstrual pelvic pain) (2) Treatment duration of therapy has not exceeded a total of 24 months</p>
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

MYTESI

Products Affected

- MYTESI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Noninfectious Diarrhea associated with HIV/AIDS (NID): (1) Diagnosis of HIV/AIDS and member is on antiretroviral therapy. (2) Member requires symptomatic relief of non-infectious diarrhea. (3) Inadequate response or inability to tolerate at least one anti-diarrheal medication (e.g. loperamide, atropine/diphenoxylate, etc.). (4) Infectious diarrhea (e.g. cryptosporidiosis, C. Difficile, etc.) has been ruled out.
Age Restrictions	(NID): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NEXLETOL/NEXLIZET

Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Heterozygous Familial Hypercholesterolemia (HeFH) OR Primary hyperlipidemia (PH). (Initial): (1) One of the following: (A) Diagnosis of HeFH, OR (B) Diagnosis of Primary hyperlipidemia (2) ONE of the following: (A) LDL-C 70 mg/dL or greater after at least 8 weeks of one low, moderate or high-intensity statin therapy and member will continue to receive statin therapy at maximally tolerated dose OR (B) Inability to tolerate statin therapy as documented by one of the following: (i) member had rhabdomyolysis or symptoms with creatine kinase (CK) exceeding 10 times the upper limit of normal (ULN) or (ii) either of the following with TWO statins: myalgia (no CK elevation) or myositis (CK less than 10 times ULN or (iii) hepatotoxicity from statin use (increased AST/ALT exceeding 3 times ULN) or (iv) liver disease documented by Child Pugh A or worse OR AST/ALT exceeding 3 times ULN for at least 6 weeks. (3) One of the following: (A) Member has been receiving at least 8 weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy OR (B) Member has contraindication or intolerance to ezetimibe (Zetia).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial): 6 months. (Continuation): 12 months

PA Criteria	Criteria Details
Other Criteria	Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD (Initial): (1) One of the following: (A) Diagnosis of established cardiovascular disease (CVD) (e.g., coronary artery disease, symptomatic peripheral arterial disease, cerebrovascular atherosclerotic disease) OR (B) Diagnosis of a high risk for a CVD event but without established CVD [e.g., diabetes mellitus (type 1 or type 2) in females over 65 years of age or males over 60 years of age] (2) One of the following: (A) Member is statin intolerant as evidenced by an inability to tolerate at least two statins, with at least one started at the lowest starting daily dose, due to intolerable symptoms or clinically significant biomarker changes of liver function or muscle function (e.g., creatine kinase) (B) Member has a contraindication to all statins (3) ONE of the following LDL-C values within the last 120 days (A) LDL-C greater than or equal to 55 mg/dL with ASCVD OR (B) LDL-C greater than or equal to 100 mg/dL without ASCVD (4) One of the following (A) For Nexletol, ONE of the following: (i) Member has been receiving at least 12 weeks of generic ezetimibe therapy (ii) Patient has a history of contraindication, or intolerance to ezetimibe OR (B) For Nexlizet, member has been receiving at least 12 weeks of generic ezetimibe therapy (HeFH, PH) (Continuation): (1) Positive Clinical response to therapy (e.g. reduction in LDL-C levels). (2) One of the following: (A) Member continues to receive other lipid-lowering therapy (e.g. statins, ezetimibe) at the maximally tolerated dose OR (B) Member has inability to tolerate other lipid-lowering therapy (e.g. statins, ezetimibe) (CVD) (Continuation): (1) Positive Clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NGENLA

Products Affected

- NGENLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Growth Failure in Children (GFC)(Initial): (1) Diagnosis of growth hormone deficiency confirmed by one of the following: (a) Height is documented by one of the following (utilizing age and gender growth charts related to height): (i) Height is greater than 2.0 standard deviations [SD] below midparental height (ii) Height is greater than 2.25 SD below population mean (below the 1.2 percentile for age and gender) (b) Growth velocity is greater than 2 SD below mean for age and gender (c) Delayed skeletal maturation of greater than 2 SD below mean for age and gender (e.g., delayed is greater than 2 years compared with chronological age) (2) documentation of bone age, (3) abnormal response from provocative testing, such as insulin-induced hypoglycemia test, levodopa, or clonidine.
Age Restrictions	(GFC) (Initial, Continuation): Member is 3 years of age or greater.
Prescriber Restrictions	(GFC) (Initial, Continuation): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	(Initial, Continuation): 12 months
Other Criteria	(GFC) (Continuation): (1) Annual clinical re-evaluation by the treating endocrinologist (2) Growth velocity greater than or equal to 2.5 cm/year.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NON-ORAL ANTIBIOTICS

Products Affected

- DALVANCE
- NUZYRA
- SIVEXTRO
- VABOMERE
- ZEMDRI
- ZERBAXA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Initial): Part D is medically necessary when documentation of a current bacterial infection and an inadequate response or inability to tolerate at least TWO drugs to which the organism is susceptible OR the requested agent is the only antibiotic to which the organism is susceptible.
Age Restrictions	
Prescriber Restrictions	(Initial, Reauth): Prescribed by or in consultation with an infectious disease specialist OR oncologist
Coverage Duration	(Initial, Reauth): 1 month
Other Criteria	Subject to Part B vs Part D review. Subject to additional clinical review for ESRD-related use - if applicable. (REAUTH): Prescriber attests that an infectious disease consult determines that a longer duration of therapy is required.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NON-ORAL CHEMO AGENTS

Products Affected

- TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when ANY of the following inclusion criteria is met: (1) Drug is FDA approved for indication and regimen requested, (2) The indication and regimen is classified as Category 1 or 2A by National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium, (3) narrative text in The American Hospital Formulary Service-Drug Information (AHFS-DI) or Clinical Pharmacology Compendium is supportive for the specific condition(s) requested, (4) The Micromedex Compendium and the strength of recommendation is listed as Class I, Class IIa, or Class IIb for the specific condition(s) requested, (5) Indication is listed in Lexi-Drugs as "off label" with evidence level A, (6) supported by Peer-Reviewed Medical Literature as defined in Chapter 15 Section 50.4.5 of the Medicare Benefit Policy Manual
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy. Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NON-PREFERRED HEPATITIS C AGENTS

Products Affected

- SOVALDI
- VOSEVI
- ZEPATIER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(1) Documentation of member's Hepatitis C genotype. (2) Prescribed regimen is consistent with the current AASLD/ IDSA guidance. (3) Inability to tolerate ONE of the following when appropriate per the current AASLD/IDSA guidance: Harvoni, Epclusa, or Mavyret.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Duration will be applied consistent with AASLD/ IDSA guidance
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NOURIANZ

Products Affected

- NOURIANZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Parkinson's Disease (PD): (1) Diagnosis of Parkinson's disease. (2) concurrent use of carbidopa/levodopa containing product, (3) Member is experiencing intermittent OFF episodes, (4) member had inadequate response or inability to tolerate ONE of the following: (a) MAO-B inhibitor (e.g. rasagiline, selegiline), (b) Dopamine agonist (e.g. pramipexole, ropinirole), (c) COMT inhibitor (e.g. entacapone)
Age Restrictions	
Prescriber Restrictions	(PD): Prescribed by or in consultation with a neurologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NOXAFIL

Products Affected

- NOXAFIL ORAL SUSPENSION
- *posaconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Treatment of invasive aspergillosis (TAI): Diagnosis of TAI. Prophylaxis of Invasive Aspergillus Infections (AI): (1) For use in prophylaxis of invasive Aspergillus infection due to being severely immunocompromised after inadequate response or inability to tolerate voriconazole (Vfend). Prophylaxis of Invasive Candida Infections(CI): (1) For use in prophylaxis of invasive Candida infection due to being severely immunocompromised after inadequate response or inability to tolerate voriconazole (Vfend). Oropharyngeal Candidiasis (OC): (1) Diagnosis of oropharyngeal candidiasis or (2) Diagnosis of Oropharyngeal candidiasis refractory to itraconazole and /or fluconazole.
Age Restrictions	(TAI): Member is 13 years of age or older (OC): Member is 13 years of age or older. (AI, CI): Member is 2 years of age or older
Prescriber Restrictions	(All Indications): Prescribed by or in consultation with an infectious disease specialist, transplant specialist, hematologist, oncologist, OR prescribed as part of chemotherapy prophylaxis protocol
Coverage Duration	Remainder of contract year
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NOXAFIL 300MG PAK

Products Affected

- NOXAFIL ORAL PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Prophylaxis of Invasive Aspergillus Infections (AI): (1) For use in prophylaxis of invasive Aspergillus infection due to being severely immunocompromised after inadequate response or inability to tolerate voriconazole (Vfend). Prophylaxis of Invasive Candida Infections(CI): (1) For use in prophylaxis of invasive Candida infection due to being severely immunocompromised after inadequate response or inability to tolerate voriconazole (Vfend).
Age Restrictions	(AI, CI): Member is 2 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist, transplant specialist, hematologist, oncologist, OR prescribed as part of chemotherapy prophylaxis protocol
Coverage Duration	Remainder of contract year
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NUCALA

Products Affected

- NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	(SA, EGPA, CRSwNP) Will not be used in combination with other biologics for asthma/allergic conditions (e.g., benralizumab [Fasenra], dupilumab [Dupixent], omalizumab [Xolair], reslizumab [Cinqair])
Required Medical Information	Part D is medically necessary when there is documentation of ONE of the following: Severe Asthma with Eosinophilic Phenotype (SA) (Initial): (1) Diagnosis of severe asthma with eosinophilic phenotype as defined by one of the following: a) blood eosinophil levels are at least 150 cells/microliter at initiation of therapy (measured within 6 weeks of dosing), or b) blood eosinophil levels at least 300 cells/microliter within the past 12 months. (2) Member has had at least two or more asthma exacerbations requiring systemic corticosteroids within the past 12 months OR member has had any prior intubation for an asthma exacerbation OR Member has had a prior asthma-related hospitalization within the past 12 months, AND (3) Member is currently treated with high-dose inhaled corticosteroids (ICS) (e.g. greater than or equal to 500 mcg fluticasone propionate equivalent/day) AND an additional controller medication (e.g. long-acting beta agonist such as salmeterol or formoterol, leukotriene inhibitor such as montelukast, theophylline), unless the individual is intolerant of or has a contraindication to two of the additional controller agents. Eosinophilic Granulomatosis with Polyangiitis (EGPA) (Initial): (1) Diagnosis of Eosinophilic Granulomatosis with Polyangiitis (EGPA), (2) Member's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy), and (3) Member is currently receiving corticosteroid therapy (e.g. prednisolone, prednisone). Chronic Rhinosinusitis with Nasal Polyps (CRSwNP) (Initial): (1) Diagnosis of chronic rhinosinusitis with nasal polyps (CRSwNP). (2) Unless contraindicated, the member has had an inadequate response to at least a 2 month treatment with an intranasal corticosteroid (e.g. fluticasone, mometasone). (3) Used in combination with another agent for CRSwNP.
Age Restrictions	(SA) (Initial, Reauth): Member is 6 years of age or older. (HES) (Initial, Reauth): Member is 12 years of age or older. (CRSwNP, EGPA) (Initial, Reauth): Member is 18 years of age or older.

PA Criteria	Criteria Details
Prescriber Restrictions	(SA): Prescribed by or in consultation with pulmonologist or allergy/immunology specialist. (EGPA): Prescribed by or in consultation with a rheumatologist. (HES): Prescribed by or in consultation with either allergist/immunologist or hematologist. (CRSwNP): Prescribed by or in consultation with allergist, immunologist, otolaryngologist or pulmonologist.
Coverage Duration	(Initial): 12 months. (Reauth): 12 months.
Other Criteria	Subject to Part B vs Part D review. Hypereosinophilic Syndrome (HES) (Initial): (1) Diagnosis of HES. (2) All of the following: (a) Member has been diagnosed for at least 6 months, (b) Verification that other non-hematologic secondary causes have been ruled out (e.g. drug hypersensitivity, parasitic helminth infection, HIV infection, non-hematologic malignancy), (c) member is FIP1-like1-platelet derived growth factor receptor alpha kinase (FIP1L1-PDGFR kinase)-negative. (3) Member has uncontrolled HES defined by both of the following: (a) Pre-treatment blood eosinophil count greater than or equal to 1000 cells/microliter, (b) Member has experienced 2 or more flares within the past 12 months. (4) Inadequate response or inability to tolerate one of the following: (a) corticosteroid therapy (e.g. prednisone), (b) cytotoxic/immunosuppressive therapy (e.g. hydroxyurea, cyclosporine, imatinib). (SA)(Reauth): (1) Member is currently treated with high-dose inhaled corticosteroids (ICS) (e.g. greater than or equal to 500mcg fluticasone propionate equivalent/day) AND an additional controller medication (e.g. long-acting beta agonist such as salmeterol or formoterol, leukotriene inhibitor such as montelukast, theophylline), unless the individual is intolerant of or has a contraindication to two of the additional controller agents. (EGPA) (Reauth): (1) Positive clinical response to therapy (e.g. increase in remission time). (HES) (Reauth): (1) Positive clinical response (e.g. reduction in corticosteroid use, reduction of blood eosinophil count, reduction in flares). (CRSwNP) (Reauth): (1) Documentation of positive clinical response to therapy (e.g. reduction of nasal polyps score [NPS: 0-8 scale], improvement in nasal obstruction symptoms via visual analog scale [VAS: 0-10 scale]), (2) Used in combination with another agent for CRSwNP.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NUEDEXTA

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	(PBA): Presence of cardiac rhythm disorder documented by a cardiac test (e.g. electrocardiogram)
Required Medical Information	Pseudobulbar Affect (PBA): (1) Diagnosis of Pseudobulbar affect (2) Member has one of the following underlying conditions: (a) Amyotrophic lateral sclerosis (b) Multiple sclerosis (c) Alzheimer's disease (d) Parkinson's disease (e) Stroke (f) Traumatic brain injury
Age Restrictions	(PBA): Member is 18 years of age or older
Prescriber Restrictions	(PBA): Prescribed by or in consultation with a neurologist or psychiatrist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NUPLAZID

Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hallucinations and Delusions Parkinson Disease Psychosis (HDPDP): Inadequate response or inability to tolerate ONE of the following (a) quetiapine or (b) clozapine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NURTEC

Products Affected

- NURTEC

PA Criteria	Criteria Details
Exclusion Criteria	(AM)(Initial, Reauth): Will be used for preventive treatment of migraine. Medication will be used in combination with another oral CGRP inhibitor. (MP) (Initial, Reauth): Will be used for acute treatment of migraine. Medication used in combination with another CGRP inhibitor for the preventive treatment of migraines.
Required Medical Information	Acute Treatment of Migraines (AM)(Initial): (1) Diagnosis of migraine with or without aura in adults. (2) Member has fewer than 15 headache days per month. (3) inadequate response or inability to tolerate ONE generic formulary triptan. Preventative Treatment of Migraines (MP) (Initial): (1) Diagnosis of episodic migraines defined as 4 to 18 migraine days per month AND inadequate response or inability to tolerate a 4 week trial of TWO of the following prophylactic medications (a) topiramate (b) divalproex sodium/ valproic acid (c) beta-blocker: metoprolol, propranolol, timolol, atenolol, nadolol (d) tricyclic antidepressants: amitriptyline, nortriptyline (e) SNRI antidepressants: venlafaxine, duloxetine
Age Restrictions	(AM)(MP)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(AM)(MP)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist.
Coverage Duration	(AM)(MP)(Initial): 6 months. (AM)(MP)(Reauth): 12 months
Other Criteria	(AM)(REAUTH): (1) Positive response to therapy (e.g. reduction in pain, photophobia, phonophobia, nausea). (MP)(Reauth): (1) Positive clinical response to therapy (e.g. reduction in headache frequency and/or intensity, use of acute migraine medications [e.g. nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g. ibuprofen, naproxen), triptans (e.g. eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

OALIVA

Products Affected

- OALIVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Primary biliary cholangitis (PBC): (1) Used in combination with ursodeoxycholic acid (e.g. Urso, Urso Forte, ursodiol), OR (2) inability to tolerate ursodeoxycholic acid. (2) Member has one of the following: (a) no cirrhosis or (b) compensated cirrhosis with no evidence of portal hypertension.
Age Restrictions	
Prescriber Restrictions	(PBC) (Initial, Reauth): Prescribed by or in consultation with a hepatologist or gastroenterologist
Coverage Duration	(Initial): 6 months. (Reauth): Indefinite
Other Criteria	(PCB)(Reauth): Positive clinical response to Ocaliva therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OFEV

Products Affected

- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Idiopathic Pulmonary Fibrosis (IPF): (1) Diagnosis of IPF confirmed by high resolution CT scan or biopsy. Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD): (1) Diagnosis of SSc-ILD confirmed by a high resolution CT scan or biopsy. Chronic Fibrosing Interstitial Lung Disease (ILDs) with progressive phenotype (Initial): (1) Confirmed by high resolution CT scan (HRCT) showing at least 10% of lung volume with fibrotic features, (2) Disease has a progressive phenotype as observed by one of the following: (a) decline of forced vital capacity (FVC), (b) worsening of respiratory symptoms, (c) increased extent of fibrosis seen on imaging.
Age Restrictions	
Prescriber Restrictions	(IPF, SSc-ILD, ILDs): Prescribed by or in consultation a pulmonologist or lung transplant specialist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(REAUTH) (IPF, ILDs, SSc-ILD): BOTH of the following (1) stabilization from baseline AND (2) no elevations in AST or ALT greater than 5 times upper limit of normal or greater than 3 times upper limit of normal with signs or symptoms of severe liver damage
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OLUMIANT

Products Affected

- OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	(RA, COVID-19, Alopecia Areata): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g., tumor necrosis factor antagonists JAK inhibitors, or potent immunosuppressants such as azathioprine, cyclosporine
Required Medical Information	Rheumatoid arthritis (RA): (1) Diagnosis of RA. (2) Inadequate response or inability to tolerate two of the following: (a) Humira, (b) Enbrel (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) Cyltezo, (g) Yuflyma OR documentation demonstrating that a trial may be inappropriate. (COVID-19): (1) Diagnosis of COVID-19 (2) Member is hospitalized (3) Member requires one of the following: (a) supplemental oxygen, (b) non-invasive mechanical ventilation, (c) invasive mechanical ventilation, (d) extracorporeal membrane oxygenation (ECMO). Alopecia Areata (AA): (1) Diagnosis of alopecia areata (2) Member has at least 50% scalp hair loss (3) Other causes of hair loss have been ruled out (e.g., androgenetic alopecia, trichotillomania, tinea capitis, psoriasis) (4) Inadequate response or inability to tolerate one previous treatment for alopecia areata (e.g., topical, intralesional, or systemic corticosteroids, topical immunotherapy)
Age Restrictions	(RA, COVID-19, Alopecia Areata): Member is 18 years of age or older
Prescriber Restrictions	(RA): Prescribed by or in consultation with a rheumatologist (AA): Prescribed by or in consultation with a dermatologist
Coverage Duration	(RA, Alopecia Areata): Indefinite. (COVID-19): 14 days
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OMVOH SQ

Products Affected

- OMVOH SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	(UC): Concurrent therapy with biologic disease modifying anti-rheumatic drug (DMARD), e.g., tumor necrosis factor antagonists
Required Medical Information	Ulcerative Colitis (UC): (1) Diagnosis of moderately to severely active Ulcerative Colitis (2) One of the following: (A) Inadequate response or inability to tolerate two of the following: (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) upadacitinib (Rinvoq), (d) tofacitinib (Xeljanz/Xeljanz XR), (e) adalimumab-adbm (Cyltezo), (f) adalimumab-aaty (Yuflyma), (g) risankizumab (Skyrizi) OR documentation demonstrating that a trial may be inappropriate (B) Will be used as a maintenance dose following the intravenous induction doses.
Age Restrictions	(UC): Member is 18 years of age or older
Prescriber Restrictions	UC): Prescribed by or in consultation with a gastroenterologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ONGENTYS

Products Affected

- ONGENTYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Parkinson's Disease (PD): (1) Diagnosis of Parkinson's disease. (2) concurrent use of carbidopa/levodopa containing product, (3) Member is experiencing intermittent OFF episodes, (4) member had inadequate response or inability to tolerate ONE of the following: (a) MAO-B inhibitor (e.g. rasagiline, selegiline), (b) Dopamine agonist (e.g. pramipexole, ropinirole), (c) COMT inhibitor (e.g. entacapone)
Age Restrictions	
Prescriber Restrictions	(PD): Prescribed by or in consultation with a neurologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ONYCHOMYCOSIS AGENTS

Products Affected

- JUBLIA
- *tavaborole*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Onychomycosis: (1) Diagnosis of onychomycosis of the toenail(s) due to Trichophyton rubrum and Trichophyton mentagrophytes. (2) Diagnosis confirmed by one of the following: (a) positive potassium hydroxide (KOH) preparation, (b) culture, or (c) histology. (3) Condition is causing debility or a disruption in member's activities of daily living. (4) Inadequate response or inability to tolerate BOTH of the following: (a) oral generic terbinafine, AND (b) oral generic itraconazole OR topical generic ciclopirox.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OPSYNVI 2024

Products Affected

- OPSYNVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PAH)(Initial): (1) Documentation of a diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV (2) Diagnosis is confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography (3) Member is currently on both of the following for the treatment of pulmonary arterial hypertension (a) macitentan (Opsumit) (b) Tadalafil
Age Restrictions	
Prescriber Restrictions	(PAH) (Initial, Continuation): Prescribed by or in consultation with pulmonologist or cardiologist
Coverage Duration	(Initial): 6 months. (Continuation): 12 months
Other Criteria	(PAH)(Continuation): Stabilization or improvement as evaluated by a cardiologist or pulmonologist
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OPZELURA

Products Affected

- OPZELURA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Atopic Dermatitis (AD) (Initial): (1) For short-term, non-continuous treatment of chronic atopic dermatitis. (2) Inadequate response or inability to tolerate at least TWO of the following: (a) topical tacrolimus OR topical pimecrolimus, OR (b) generic, prescription medium potency or higher topical steroid, OR (c) Eucrisa. (3) Will not be used in combination with therapeutic biologics, other Janus kinase inhibitors, or potent immunosuppressants (e.g. azathioprine or cyclosporine). Nonsegmental Vitiligo (NV)(Initial): (1) Diagnosis of nonsegmental vitiligo (2) Inadequate response or inability to tolerate one of the following (a) medium or higher potency topical corticosteroid (b) pimecrolimus cream (c) tacrolimus ointment (3) Will not be used in combination with therapeutic biologics, other Janus kinase inhibitors, or potent immunosuppressants (e.g., azathioprine or cyclosporine).
Age Restrictions	(AD, NV): Member is 12 years of age or older
Prescriber Restrictions	(AD) (Initial, Reauth): Prescribed by or in consultation with a dermatologist or allergist/immunologist (NV) (Initial, Reauth): Prescribed by or in consultation with a dermatologist.
Coverage Duration	(AD)(Initial): 8 Weeks, (Reauth): End of contract year (NV)(Initial): 6 months, (Reauth): 12 months
Other Criteria	(AD) (Reauth): Positive clinical response to therapy. (NV)(Reauth): (1) Documentation of positive clinical response to therapy (2) Will not be used in combination with therapeutic biologics, other Janus kinase inhibitors, or potent immunosuppressants (e.g., azathioprine or cyclosporine).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORAL ANTIBIOTICS

Products Affected

- NUZYRA
- SIVEXTRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Initial): Documentation of a current bacterial infection and an inadequate response or inability to tolerate TWO antibiotics to which the organism is susceptible OR the requested medication is the only antibiotic to which the organism is susceptible.
Age Restrictions	
Prescriber Restrictions	(Initial, Reauth): Prescribed by or in consultation with an infectious disease specialist OR oncologist
Coverage Duration	(Initial, Reauth): 1 month
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable. (REAUTH): Prescriber attests that an infectious disease consult determines that a longer duration of therapy is required.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORAL CHEMO AGENTS

Products Affected

- *abiraterone acetate*
- AFINITOR
- AFINITOR DISPERZ
- AKEEGA
- ALECENSA
- ALUNBRIG
- AUGTYRO ORAL CAPSULE 40 MG
- AYVAKIT
- BALVERSA
- *bexarotene*
- BOSULIF
- BRAFTOVI ORAL CAPSULE 75 MG
- BRUKINSA
- CABOMETYX
- CALQUENCE ORAL TABLET
- CAPRELSA
- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)
- COPIKTRA
- COTELLIC
- *dasatinib*
- DAURISMO
- ERIVEDGE
- ERLEADA
- *erlotinib hcl*
- *everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus oral tablet soluble*
- FOTIVDA
- FRUZAQLA
- GAVRETO
- *gefitinib*
- GILOTRIF
- GLEEVEC
- GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG
- IBRANCE
- ICLUSIG
- IDHIFA
- *imatinib mesylate*
- IMBRUVICA ORAL CAPSULE
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG
- INLYTA
- INQOVI
- INREBIC
- IRESSA
- IWILFIN
- JAKAFI
- JAYPIRCA
- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)
- KOSELUGO
- KRAZATI
- *lapatinib ditosylate*
- LAZCLUZE
- *lenalidomide*
- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)
- LONSURF
- LORBRENA
- LUMAKRAS ORAL TABLET 120 MG, 320 MG
- LYNPARZA ORAL TABLET
- LYTGobi (12 MG DAILY DOSE)
- LYTGobi (16 MG DAILY DOSE)
- LYTGobi (20 MG DAILY DOSE)
- MEKINIST
- MEKTOVI
- NERLYNX
- NEXAVAR
- NINLARO
- NUBEQA
- ODOMZO
- OGSIVEO
- OJEMDA ORAL SUSPENSION RECONSTITUTED

- OJEMDA ORAL TABLET 100 MG
- OJJAARA
- ONUREG
- ORGOVYX
- ORSERDU
- *pazopanib hcl*
- PEMAZYRE
- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)
- POMALYST
- QINLOCK
- RETEVMO ORAL TABLET
- REVLIMID
- REZLIDHIA
- ROZLYTREK
- RUBRACA
- RYDAPT
- SCEMBLIX
- *sorafenib tosylate*
- SPRYCEL
- STIVARGA
- *sunitinib malate*
- SUTENT
- TABRECTA
- TAFINLAR
- TAGRISSO
- TALZENNA
- TARGRETIN
- TASIGNA
- TAZVERIK
- TEPMETKO
- THALOMID ORAL CAPSULE 100 MG, 50 MG
- TIBSOVO
- TORPENZ
- TRUQAP ORAL TABLET
- TUKYSA
- TURALIO ORAL CAPSULE 125 MG
- TYKERB
- VANFLYTA
- VENCLEXTA
- VENCLEXTA STARTING PACK
- VERZENIO
- VITRAKVI
- VIZIMPRO
- VONJO
- VORANIGO
- VOTRIENT
- WELIREG
- XALKORI
- XOSPATA
- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (80 MG TWICE WEEKLY)
- XTANDI
- YONSA
- ZEJULA ORAL TABLET
- ZELBORAF
- ZOLINZA
- ZYDELIG
- ZYKADIA ORAL TABLET
- ZYTIGA

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	Approved when ANY of the following inclusion criteria is met: (1) Drug is FDA approved for indication and regimen requested, (2) The indication and regimen is classified as Category 1 or 2A by National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium, (3) narrative text in The American Hospital Formulary Service-Drug Information (AHFS-DI) or Clinical Pharmacology Compendium is supportive for the specific condition(s) requested, (4) The Micromedex Compendium and the strength of recommendation is listed as Class I, Class IIa, or Class IIb for the specific condition(s) requested, (5) Indication is listed in Lexi-Drugs as "off label" with evidence level A, (6) supported by Peer-Reviewed Medical Literature as defined in Chapter 15 Section 50.4.5 of the Medicare Benefit Policy Manual
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORAL PAH AGENTS

Products Affected

- OPSUMIT
- ORENITRAM
- ORENITRAM MONTH 1
- ORENITRAM MONTH 2
- ORENITRAM MONTH 3

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (Initial): (1) Documentation of a diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV. (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography. (3) Documentation of mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion.
Age Restrictions	
Prescriber Restrictions	(PAH) (Initial): Prescribed by or in consultation with a Cardiologist or Pulmonologist
Coverage Duration	(Initial): 6 months. (Continuation):12 months.
Other Criteria	(PAH)(Continuation): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORENCIA SQ

Products Affected

- ORENCIA CLICKJECT
- ORENCIA SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	(RA, PsA, pJIA): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Rheumatoid arthritis (RA): (1) Diagnosis of RA. (2) Inadequate response or inability to tolerate one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Psoriatic arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Polyarticular Juvenile idiopathic arthritis (pJIA): (1) Diagnosis of JIA. (2) Inadequate response or inability to tolerate one of the following: methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine.
Age Restrictions	(Pjia, PsA): Member is 2 years of age or older. (RA): member is 18 years of age or older
Prescriber Restrictions	(RA, pJIA): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORIAHNN

Products Affected

- ORIAHNN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Uterine Leiomyomas (UL)(Initial): (1) Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas (fibroids), (2) Member is premenopausal, (3) ONE of the following: (a) Inadequate response or inability to tolerate one of the following for no more than 30 days: combination (estrogen/progestin) contraceptive, progestins, tranexamic acid, OR (b) Member has had a previous interventional therapy to reduce bleeding (e.g. uterine-artery embolization and magnetic resonance-guided focused ultrasonography), (4) Treatment duration of therapy has not exceeded a total of 24 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(UL) (Reauth): (1) Member has improvement in bleeding associated with uterine leiomyomas (fibroids) (e.g. significant/sustained reduction in menstrual blood loss per cycle, improved quality of life, etc.), (2) Treatment duration of therapy has not exceeded a total of 24 months.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORILISSA

Products Affected

- ORILISSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pain Associated with Endometriosis (PAE): (1) Documentation of ONE of the following, (a) Inadequate response or inability to tolerate BOTH of the following (i) one nonsteroidal anti-inflammatory drug AND (ii) one contraceptive OR (b) Member has had surgical ablation to prevent recurrence. (2) Treatment duration does not exceed 24 months (150mg tablet) or 6 months (200mg tablet).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months for 150mg tablet, 6 months for 200mg tablet
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORKAMBI

Products Affected

- ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	(CF): Diagnosis of CF other than those homozygous for the F508del mutation
Required Medical Information	Cystic Fibrosis (CF): (1) Diagnosis of CF, (2) Member is homozygous for the F508del mutation in the CFTR gene (3) If the member's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene
Age Restrictions	(CF): Member is 1 year of age or older
Prescriber Restrictions	(CF): Prescribed by or in consultation with pulmonologist or Specialist affiliated with a CF care center
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ORLADEYO

Products Affected

- ORLADEYO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hereditary Angioedema (HAE): (1) Diagnosis of HAE, (2) For prophylaxis against HAE attacks.
Age Restrictions	(HAE): Member is 12 years of age or older
Prescriber Restrictions	(HAE): Prescribed by or in consultation with an immunologist, allergist or pulmonologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OSMOLEX

Products Affected

- OSMOLEX ER ORAL TABLET EXTENDED
RELEASE 24 HOUR 129 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Parkinson's Disease (PD): (1) Diagnosis of PD, (2) inadequate response or inability to tolerate amantadine immediate release, (3) Inadequate response or inability to tolerate one of the following: (a) Carbidopa-levodopa (b) MAO-B Inhibitor (e.g. rasagiline, selegiline) (c) Dopamine Agonist (e.g. pramipexole, ropinirole) . Drug-induced extrapyramidal symptoms (DIEPS): (1) BOTH of the following: (A) ONE of the following: (i) Member has persistent extrapyramidal symptoms despite a trial of dose reduction, tapering, or discontinuation of the offending medication or (ii) Member is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. (B) Inadequate response or inability to tolerate amantadine immediate release
Age Restrictions	
Prescriber Restrictions	(DIEPS): Prescribed by or in consultation with a neurologist or psychiatrist (PD): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OTEZLA

Products Affected

- OTEZLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Oral Ulcers Associated with Behcet's Disease (OU-BD): (1) Diagnosis of OU-BD. (2) Inadequate response or inability to tolerate ONE topical corticosteroid (e.g. triamcinolone acetonide dental paste) AND ONE systemic corticosteroid. Psoriatic arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least one DMARD. Plaque psoriasis (PsO): (1) Diagnosis of PsO.
Age Restrictions	(PsA, OUBD): Member is 18 years of age or older. (PsO): Member is 6 years of age or older.
Prescriber Restrictions	(PsA, OU-BD): Prescribed by or in consultation with a Rheumatologist or Dermatologist. (PsO): Prescribed by or in consultation with dermatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

OXERVATE

Products Affected

- OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Neurotrophic keratitis (NK)(Initial): (1) Diagnosis of NK. (2) Submission of chart documentation indicating treatment of left eye, right eye, or both
Age Restrictions	
Prescriber Restrictions	(NK) (Initial, Reauth) Prescribed by or in consultation with an ophthalmologist or Optometrist.
Coverage Duration	8 weeks
Other Criteria	(NK)(Reauth): (1)Submission of chart documentation indicating treatment of left eye, right eye, or both, (2) Member has received less than or equal to 8 weeks of therapy (one course of therapy) per affected eye(s), (3) Documentation of clinical rational for treatment greater than 8 weeks (e.g. member has a recurrence of neurotrophic keratitis, or treatment of a different eye), (4) Documentation of clinical response to prior Oxervate therapy, (5) Member will not exceed a total of 16 weeks of Oxervate therapy per affected eye(s).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PALYNZIQ

Products Affected

- PALYNZIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Phenylketonuria (PK)(Initial): (1) Member has uncontrolled blood phenylalanine concentrations greater than 600 micromol/L on existing management, (2) Member will continue to have phenylalanine blood levels measured periodically during therapy.
Age Restrictions	(PK)(Initial, Continuation): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial): 6 months, (Continuation): Indefinite
Other Criteria	(PK)(CONTINUATION): (1) A positive clinical response to Palynziq therapy as determined by prescriber.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PART D VS EXCLUDED

Products Affected

- AURYXIA
- CRINONE
- IMVEXXY MAINTENANCE PACK
- IMVEXXY STARTER PACK
- INTRAROSA
- OSPHENA
- VFEND IV
- *voriconazole intravenous*
- ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	Subject to additional clinical review for ESRD-related use - if applicable.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PDE INHIBITOR AGENTS FOR PAH

Products Affected

- ADCIRCA
- ALYQ
- *tadalafil (pah)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH)(Initial): (1) Diagnosis of PAH WHO Group I with New York Heart Association (NYHA) Functional Class II to IV, (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography, (3) Mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion, (4) inadequate response or inability to tolerate generic tadalafil (Applies to Brand Adcirca only)
Age Restrictions	
Prescriber Restrictions	(PAH)(Initial): Prescribed by or in consultation with Cardiologist or Pulmonologist
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(PAH)(Reauth): (1) Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PRALUENT

Products Affected

- PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Hyperlipidemia (HLA) (Initial): (1) Diagnosis of hyperlipidemia, (2) ONE of the following: (A) LDL-C 70 mg/dL or greater after a minimum 8-week trial of at least moderate-intensity statin therapy OR (B) Inability to tolerate statin therapy as documented by one of the following: (i) member had rhabdomyolysis or symptoms with creatine kinase (CK) exceeding 10 times the upper limit of normal (ULN) or (ii) either of the following with TWO statins: myalgia (no CK elevation) or myositis (CK less than 10 times ULN or (iii) hepatotoxicity from statin use (increased AST/ALT exceeding 3 times ULN) or (iv) liver disease documented by Child Pugh A or worse OR AST/ALT exceeding 3 times ULN for at least 6 weeks</p> <p>Atherosclerotic cardiovascular disease (ASCVD) (Initial): (1) Diagnosis of atherosclerotic cardiovascular disease (ASCVD) as diagnosed by either stress test, angiography, atherosclerotic event (e.g. MI, angina, stroke, claudication, carotid stenosis) or arterial intervention for atherosclerotic disease (e.g. coronary, peripheral, carotid), (2) ONE of the following: (A) LDL-C 70 mg/dL or greater after a minimum 8-week trial of at least moderate-intensity statin therapy OR (B) Inability to tolerate statin therapy as documented by one of the following: (i) member had rhabdomyolysis or symptoms with creatine kinase (CK) exceeding 10 times the upper limit of normal (ULN) or (ii) either of the following with TWO statins: myalgia (no CK elevation) or myositis (CK less than 10 times ULN or (iii) hepatotoxicity from statin use (increased AST/ALT exceeding 3 times ULN) or (iv) liver disease documented by Child Pugh A or worse OR AST/ALT exceeding 3 times ULN for at least 6 weeks</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial): 6 months (Continuation): 12 months

PA Criteria	Criteria Details
Other Criteria	<p>Homozygous Familial Hypercholesterolemia (HoFH) (Initial): (1) Diagnosis of HoFH. (2) One of the following: (a) Untreated LDL-C greater than 500 mg/dL or (b) Treated LDL-C greater than 300 mg/dL. (3) Member is receiving other lipid-lowering therapy (e.g. statin, ezetimibe). (4) Not used in combination with Juxtapid (lomitapide). (HLA, ASVCD)(CONTINUATION): (1) Positive Clinical response to therapy (e.g. reduction in LDL-C levels). (HoFH)(Continuation): (1) Positive Clinical response to therapy (e.g. reduction in LDL-C levels). (2) Member continues to receive other lipid-lowering therapy (e.g. statin, ezetimibe). (3) Not used in combination with Juxtapid (lomitapide).</p>
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PREFERRED HEPATITIS C AGENTS

Products Affected

- EPCLUSA
- HARVONI ORAL PACKET
- HARVONI ORAL TABLET 90-400 MG
- *ledipasvir-sofosbuvir*
- MAVYRET
- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(1) Documentation of member's Hepatitis C Genotype. (2) Prescribed regimen is consistent with the current AASLD/IDSA guidance.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Duration will be applied consistent with AASLD/ IDSA guidance
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PRETOMANID

Products Affected

- *pretomanid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Multidrug Resistant Tuberculosis (MDRTB): (1) Diagnosis of pulmonary extensively drug resistant (XDR), treatment-intolerant or nonresponsive MDRTB. (2) Medication will be used as part of a combination regimen with bedaquiline (Sirturo) and linezolid.
Age Restrictions	(MDRTB): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	26 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PROCYSBI

Products Affected

- PROCYSBI ORAL PACKET

PA Criteria	Criteria Details
Exclusion Criteria	(NC) (Initial, Reauth): Hypersensitivity to penicillamine
Required Medical Information	Nephrotic Cystinosis (NC) (Initial): (1) Diagnosis of nephrotic cystinosis, (2) inadequate response or titration from cysteamine bitartrate immediate-release capsules (Cystagon), (3) documentation of baseline WBC and alkaline phosphatase levels.
Age Restrictions	(NC): Member is 1 year of age or older
Prescriber Restrictions	(NC)(Initial, Reauth): Prescribed by or in consultation with a nephrologist.
Coverage Duration	(Initial): 3 months. (Reauth): 6 months.
Other Criteria	(NCB)(REAUTH): The prescriber has evaluated all of the following since the initiation of treatment: (1) ONE of the following: (a) WBC cysteine level or (b) plasma cysteamine level, (2) WBC count, AND (3) alkaline phosphatase level.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PROLIA

Products Affected

- PROLIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Part D is medically necessary when all of the following criteria are met:</p> <p>Osteoporosis (OS) (Initial): ALL of the following: (1) Diagnosis of osteoporosis confirmed by one of the following: (a) T-score less than or equal to -2.5 (b) history of osteoporotic fracture (e.g. vertebral, hip, non-vertebral) or (c) multiple risk factors for fracture. (2) ONE of the following: (a) inadequate response or inability to tolerate at least one other osteoporosis medicine (e.g. oral bisphosphonates, calcitonin, estrogens, selective estrogen receptor modulator (SERMs)) or (b) severely deteriorated condition indicating that the osteoporosis is so significant that a trial of oral bisphosphonates is not medically warranted or (c) documented moderate to severe renal insufficiency (glomerular filtration rate [GFR] 30 to 35 ml/min or less). Osteopenia (OPN) (Initial): BOTH of the following: (1) Diagnosis of osteopenia confirmed by a T-score less than -1.0, but greater than -2.5. (2) One of the following: (a) Member is receiving adjuvant aromatase inhibitor therapy for breast cancer (b) member is receiving androgen deprivation therapy for non-metastatic prostate cancer. Prophylaxis of Postmenopausal Osteoporosis (PO) (Initial): BOTH of the following: (1) BMD T score less than -1.0 and greater than -2.5. (2) ONE of the following: (a) Inadequate response or inability to tolerate an oral bisphosphonates or a selective estrogen receptor modulator (SERMs) or (b) documented moderate to severe renal insufficiency (glomerular filtration rate [GFR] 30 to 35 ml/min or less) not receiving dialysis or diagnosed with stage 5 kidney disease.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite

PA Criteria	Criteria Details
Other Criteria	Subject to Part B vs Part D review. Glucocorticoid Induced Osteoporosis (GCO)(Initial): ALL of the following: (1) Diagnosis of CGO in men or women. (2) Daily dose of prednisone is greater than or equal to 5mg for at least 3 months. (3) ONE of the following: (a)Inadequate response or inability to tolerate ONE of the following: (i) bisphosphonates, (ii) hormone replacement therapy, (iii) selective-estrogen receptor modulators (SERMs)., (b) severely deteriorated condition indicating that the osteoporosis is so significant that a trial of oral bisphosphonates is not medically warranted, (c) documented moderate to severe renal insufficiency (glomerular filtration rate [GFR] 30 to 35 ml/min or less)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PROMACTA

Products Affected

- PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Chronic Idiopathic Thrombocytopenic Purpura (ITP)(Initial): (1) Diagnosis of relapsed/refractory, persistent, or chronic ITP. (2) Baseline platelet count is less than 30,000/mcL. (3) Member's degree of thrombocytopenia and clinical condition increase the risk of bleeding. (4) Inadequate response or inability to tolerate ONE of the following: (a) corticosteroids, (b) immune globulin, (c) splenectomy. First-Line for Severe Aplastic Anemia(FLSAA): (1) Diagnosis of severe aplastic anemia. (2) Used for first-line treatment (i.e., member has not received prior immunosuppressive therapy). (3) Member meets at least TWO of the following: (a) absolute neutrophil count less than 500/mcL, (b) platelet count less than 20,000/mcL, (c) absolute reticulocyte count less than 60,000/mcL. (4) Used in combination with standard immunosuppressive therapy (e.g. Atgam [antithymocyte globulin equine] and cyclosporine). Refractory Severe Aplastic Anemia (RSAA)(Initial): (1) Diagnosis of refractory severe aplastic anemia. (2) Member has a platelet count less than 30,000/mcL. (3) Inadequate response or inability to tolerate immunosuppressive therapy with antithymocyte globulin and cyclosporine. Chronic Hepatitis C-Associated Thrombocytopenia (HEPC-TP): (1) Diagnosis of chronic hepatitis C-associated thrombocytopenia. (2) One of the following: (a) Planning to initiate and maintain interferon-based treatment, or (b) currently receiving interferon-based treatment.</p>
Age Restrictions	
Prescriber Restrictions	(ITP, FLSAA, RSAA)(Initial, Continuation): Prescribed by or in consultation with hematologist/oncologist. (HEPC-TP)(Initial, Continuation): Prescribed by or in consultation with Hematologist/oncologist, Gastroenterologist, Hepatologist, Infectious disease specialist, HIV specialist.
Coverage Duration	(ITP)Initial,Cont.=12mo.(FLSAA)=6mo.(RSAA)Initial=6mo.(HEPC-TP)Initial=3mo.(RSAA,HEPC-TP)Cont.=12mo.

PA Criteria	Criteria Details
Other Criteria	(ITP, RSAA)(Continuation): (1) Positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. (HEPC-TP)(Continuation): (1) ONE of the following: (A) For members that started treatment with Promacta prior to initiation of treatment with interferon, both of the following: (i) Member is currently on antiviral interferon therapy for treatment of chronic hepatitis C, AND (ii) member reached a threshold platelet count that allows initiation of antiviral interferon therapy with Promacta treatment by week 9. (B) For members that started treatment with Promacta while on concomitant treatment with interferon, member is currently on antiviral interferon therapy for treatment of chronic hepatitis C.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PYRUKYND

Products Affected

- PYRUKYND
- PYRUKYND TAPER PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hemolytic anemia with pyruvate kinase deficiency (HAWPKD) (Initial): (1) Diagnosis of hemolytic anemia confirmed by the presence of chronic hemolysis (e.g., increased indirect bilirubin, elevated lactated dehydrogenase [LDH], decreased haptoglobin, increased reticulocyte count) (2) Diagnosis of pyruvate kinase deficiency confirmed by molecular testing of ALL of the following mutations on the PKLR gene: (a) Presence of at least 2 variant alleles in the pyruvate kinase liver and red blood cell (PKLR) gene, of which at least 1 was a missense variant, (b) Member is not homozygous for the c. 1436G A (p.R479H) variant, (c) Member does not have 2 non-missense variants (without the presence of another missense variant) in the PKLR gene (3) Hemoglobin is less than or equal to 10g/dL (4) Member has symptomatic anemia or is transfusion dependent (5) Other causes of hemolytic anemia (e.g., infections, toxins, drugs) have been ruled out.
Age Restrictions	(HAWPKD) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(HAWPKD) (Initial, Reauth): Prescribed by or in consultation with a hematologist
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(HAWPKD)(Reauth): (1) Documentation of positive clinical response to therapy, improvement in markers of hemolysis from baseline (e.g., bilirubin, lactated dehydrogenase [LDH], haptoglobin, reticulocyte count).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

QBREXZA

Products Affected

- QBREXZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hyperhidrosis: (1) Hyperhidrosis Disease Severity Scale grade 3 or 4.
Age Restrictions	(Hyperhidrosis): Member is 9 years of age or older
Prescriber Restrictions	(Hyperhidrosis): Prescribed by or in consultation with a dermatologist, primary care physician, internist, or pediatrician.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

QUALAQUIN

Products Affected

- QUALAQUIN
- *quinine sulfate oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Uncomplicated Malaria: (1) Diagnosis of uncomplicated malaria due to (a) plasmodium falciparum malaria or (b) plasmodium vivax . (2) One of the following: (a) Both of the following: (i) treatment in areas of chloroquine-sensitive malaria (ii) Inadequate response or inability to tolerate chloroquine or hydroxychloroquine, or (b) treatment in areas of chloroquine-resistant malaria. Babesiosis: (1) Diagnosis of babesiosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Babesiosis: 10 days Uncomplicated Malaria: 14 Days
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

QULIPTA

Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	(CMP, EMP) (Initial, Reauth): Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.
Required Medical Information	Episodic Migraine Prevention (EMP)(Initial): (1) Diagnosis of episodic migraines defined as 4 to 14 headache days per month AND (2) Inadequate response or inability to tolerate BOTH of the following: (a) Emgality AND (b) Aimovig. Chronic Migraine Prevention (CMP)(Initial): (1) Diagnosis of chronic migraines defined as greater than or equal to 15 headache days per month AND (2) Inadequate response or inability to tolerate BOTH of the following: (a) Emgality AND (b) Aimovig.
Age Restrictions	(EMP, CMP) (Initial, Reauth): Member 18 years of age
Prescriber Restrictions	(EMP, CMP) (Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist
Coverage Duration	(EMP, CMP) (Initial): 6 months (Migraine Prevention EMP, CMP) (Reauth): 12 months
Other Criteria	(EMP, CMP) (Reauth): (1) Response to therapy as defined by a reduction in headache days per month (defined as at least 4 hours duration and moderate intensity).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

QUVIVIQ

Products Affected

- QUVIVIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Insomnia): (1) Diagnosis of insomnia characterized by difficulties with sleep onset and/or sleep maintenance (2) inadequate response or inability to tolerate ramelteon and Belsomra
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RADICAVA

Products Affected

- RADICAVA ORS STARTER KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Amyotrophic Lateral Sclerosis (ALS)(Initial): (1) Diagnosis of definite or probable ALS per the revised El Escorial World Federation of Neurology criteria. (2) Time from symptom onset is 2 years or less. (3) Normal respiratory function defined as forced vital capacity (FVC) of greater than or equal to 80% at the start of treatment. (4) Scores of 2 points or greater on each individual item of the ALS Functional Rating Scale-revised (ALSFRS-R).
Age Restrictions	
Prescriber Restrictions	(ALS)(Initial and Reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	(Initial, Reauth): 6 months.
Other Criteria	(ALS)(Reauth): (1) Member shows benefit from therapy (e.g. slowing of decline of functional abilities).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RAVICTI

Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	(UCD): Acute hyperammonemia. N-acetylglutamate synthase (NAGS) deficiency
Required Medical Information	Urea Cycle Disorder (UCD): (1) Diagnosis of urea cycle disorder involving deficiencies of carbamoyl phosphate synthetase (CPS), ornithine transcarbamoylase (OTC), or argininosuccinic acid synthetase (AAS) confirmed via enzymatic, biochemical, or genetic testing. (2) Inadequate response or inability to tolerate sodium phenylbutyrate.
Age Restrictions	(UCD): Member is 2 months of age or older
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RECORLEV

Products Affected

- RECORLEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cushing's Syndrome (CS) (Initial): (1) Diagnosis of Cushing's syndrome (2) Member is being treated for endogenous hypercortisolemia (e.g., pituitary adenoma, ectopic tumor, adrenal adenoma) (3) One of the following: (a) Member is not a candidate for surgery, (b) surgery has not been curative (4) Inadequate response or inability to tolerate oral ketoconazole
Age Restrictions	(CS) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(CS) (Initial) (Reauth): Prescribed by or in consultation with an endocrinologist
Coverage Duration	(Initial) (Reauth): 12 months
Other Criteria	(CS) (Reauth): (1) Documentation of positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

REPATHA

Products Affected

- REPATHA
- REPATHA PUSHTRONEX SYSTEM
- REPATHA SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hyperlipidemia (HLA) (Initial): (1) Diagnosis of hyperlipidemia. Atherosclerotic cardiovascular disease (ASCVD)(Initial): (1) Diagnosis of ASCVD confirmed by either stress test, angiography, atherosclerotic event (e.g. MI, angina, stroke, claudication, carotid stenosis) or arterial intervention for atherosclerotic disease (e.g. coronary, peripheral, carotid). (HLA, ASCVD) (Initial): ONE of the following: (A) LDL-C 70 mg/dL or greater after a minimum 8-week trial of at least moderate-intensity statin therapy OR (B) Inability to tolerate statin therapy as documented by one of the following: (i) member had rhabdomyolysis or symptoms with creatine kinase (CK) exceeding 10 times the upper limit of normal (ULN) or (ii) either of the following with TWO statins: myalgia (no CK elevation) or myositis (CK less than 10 times ULN or (iii) hepatotoxicity from statin use (increased AST/ALT exceeding 3 times ULN) or (iv) liver disease documented by Child Pugh A or worse OR AST/ALT exceeding 3 times ULN for at least 6 weeks
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(Initial): 6 months. (Continuation): 12 months.
Other Criteria	Homozygous Familial Hypercholesterolemia (HoFH) (Initial): (1) Diagnosis of HoFH. (2) One of the following: (a) Untreated LDL-C greater than 500 mg/dL or (b) Treated LDL-C greater than 300 mg/dL. (3) Member is receiving other lipid-lowering therapy (e.g. statin, ezetimibe). (4) Not used in combination with Juxtapid (lomitapide). (HLA, ASCVD)(Continuation): (1) Positive Clinical response to therapy (e.g. reduction in LDL-C levels). (HoFH)(Continuation): (1) Positive Clinical response to therapy (e.g. reduction in LDL-C levels). (2) Member continues to receive other lipid-lowering therapy (e.g. statin, ezetimibe). (3) Not used in combination with Juxtapid (lomitapide).
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

RESPIRATORY ENZYMES

Products Affected

- ARALAST NP INTRAVENOUS SOLUTION
RECONSTITUTED 1000 MG
- GLASSIA
- PROLASTIN-C INTRAVENOUS SOLUTION
- ZEMAIRA INTRAVENOUS SOLUTION
RECONSTITUTED 1000 MG

PA Criteria	Criteria Details
Exclusion Criteria	(ATT): (1) IgA deficiency with known anti-IgA antibody.
Required Medical Information	Part D is medically necessary when ALL of the following are met: Congenital Alpha-1 Antitrypsin Deficiency (ATT): (1) Diagnosis of congenital alpha1-antitrypsin deficiency confirmed by one of the following: (a) PiZZ, PiZ(null) or Pi(null)(null) protein phenotypes (homozygous) or (b) Other rare AAT disease-causing alleles associated with serum alpha1-antitrypsin (AAT) level less than 11?mol/L. (2) Low serum concentration of alpha-1 antitrypsin defined as less than 35 percent of normal (less than 80 mg/dL or less than 11 uM/L or less than 0.8 g/L). (3) the member has progressive panacinar emphysema with a documented rate of decline in forced expiratory volume in 1 second (FEV 1)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

REZDIFFRA 2024

Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(MASH) (Initial): (1) Diagnosis of metabolic dysfunction associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH) (2) Documentation of ONE of the following: (a) FibroScan-aspartate aminotransferase (FAST) (b) MRI-aspartate aminotransferase (MAST) (c) Liver biopsy (3) Documentation that disease is fibrosis stage F2 or F3 as confirmed by ONE of the following: (a) FibroScan (b) Fibrosis-4 index (FIB-4) (c) Magnetic Resonance Elastography (MRE). (4) Presence of greater than or equal to 3 metabolic risk factors (e.g., Type 2 diabetes, hypertension, obesity) (5) Drug is used as adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program)
Age Restrictions	
Prescriber Restrictions	(Initial, Reauth): Prescribed by or in consultation with a gastroenterologist or hepatologist.
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(MASH) (Reauth): (1) member demonstrates positive response to therapy (e.g., MASH resolution, fibrosis stage improvement, etc.) (2) Member has not progressed to cirrhosis.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

REZUROCK

Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic graft-versus-host disease (cGVHD) (Initial): (1) Diagnosis of chronic graft-versus-host disease, (2) Inadequate response or inability to tolerate two or more lines of systemic therapy (e.g. corticosteroids, mycophenolate, etc.)
Age Restrictions	
Prescriber Restrictions	(cGVHD) (Initial/Reauth): Prescribed by or in consultation with one of the following (1) Hematologist, (2) Oncologist, (3) physician experienced in the management of transplant patients
Coverage Duration	(Initial/Reauth): 12 months
Other Criteria	(cGVHD) (Reauth): (1) Member does not show evidence of progressive disease while on therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RINVOQ

Products Affected

- RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	(RA, PsA, AD, UC, AS, nr-AxSPA, CD, PJIA): Concurrent use with any other biologic disease modifying antirheumatic drug (DMARD), other JAK inhibitors, or potent immunosuppressants (e.g. azathioprine, cyclosporine).
Required Medical Information	Rheumatoid Arthritis (RA): (1) Diagnosis of moderate to severely active RA. (2) Member has inadequate response or inability to tolerate one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine) (3) Member had inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA, (2) Inadequate response or inability to tolerate at least one DMARD (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine) (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate. Atopic Dermatitis (AD): (1) Diagnosis of moderate to severe atopic dermatitis, (2) Inadequate response or inability to tolerate at least one of the following: (a) medium or higher potency topical corticosteroid, (b) pimecrolimus cream, (c) tacrolimus cream, (d) Eucrisa (crisaborole) ointment (3) Inadequate response or inability to tolerate at least one systemic drug product for the treatment of atopic dermatitis (examples include, but are not limited to Dupixent, Adbry or documentation that a trial may be inappropriate. Ulcerative Colitis (UC): (1) Diagnosis of Ulcerative Colitis (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. corticosteroid, Aminosalicylate, azathioprine or 6-mercaptopurine). (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(RA, UC, AS, nr-AxSPA, CD): Member is 18 years of age or older (AD): Member is 12 years of age or older. (PsA, PJIA): Member is 2 years of age or older.
Prescriber Restrictions	(RA, AS, nr-AxSPA, PJIA): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a rheumatologist or dermatologist. (AD): Prescribed by or in consultation with a dermatologist, allergist, immunologist. (UC, CD): Prescribed by or in consultation with a gastroenterologist.

PA Criteria	Criteria Details
Coverage Duration	Indefinite
Other Criteria	<p>Ankylosing Spondylitis (AS): (1) Diagnosis of active ankylosing spondylitis. (2) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g., Enbrel, Humira, Cyltezo, Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Non-radiographic axial spondyloarthritis (nr-AxSPA): (1) Diagnosis of nr-AxSPA (2) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g., Enbrel, Humira, Cimzia, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate (3) Inadequate response or inability to tolerate one NSAID. Crohn's Disease (CD): (1) Diagnosis of CD. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g., corticosteroid, methotrexate, azathioprine or 6-mercaptopurine) (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g., Enbrel, Humira, Cyltezo, Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Polyarticular Juvenile Idiopathic Arthritis (PJIA): (1) Diagnosis of active PJIA, (2) Inadequate response or inability to tolerate ONE of the following: Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate.</p>
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RINVOQ LQ 2024

Products Affected

- RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	(PsA, pJIA): Concurrent use with any other biologic disease modifying antirheumatic drug (DMARD), other JAK inhibitors, or potent immunosuppressants (e.g. azathioprine, cyclosporine).
Required Medical Information	Psoriatic Arthritis (PsA): (1) Diagnosis of PsA, (2) Inadequate response or inability to tolerate at least one DMARD (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate. Polyarticular Juvenile Idiopathic Arthritis (PJIA): (1) Diagnosis of active PJIA, (2) Inadequate response or inability to tolerate ONE of the following: Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine (3) Member had an inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(PsA, pJIA): Member is 2 years of age or older
Prescriber Restrictions	(PsA): Prescribed by or in consultation with a rheumatologist or dermatologist. (pJIA): Prescribed by or in consultation with a rheumatologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

RIVFLOZA 2024

Products Affected

- RIVFLOZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PH1)(Initial): (1) Diagnosis of primary hyperoxaluria type 1 (PH1) (2) Disease has been confirmed by both of the following: (a) One of the following: (i) elevated urinary oxalate excretion (ii) elevated plasma oxalate concentration (iii) spot urinary oxalate to creatinine molar ratio greater than normal for age (b) One of the following: (i) genetic testing demonstrating a mutation in the alanine: glyoxylate aminotransferase (AGXT) gene (ii) Liver biopsy demonstrating absence or reduced alanine: glyoxylate aminotransferase (AGT) activity (3) Member has preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2)
Age Restrictions	
Prescriber Restrictions	(PH1) (Initial, Reauth): Prescribed by or in consultation with one of the following: hepatologist, nephrologist, urologist, geneticist, specialist with expertise in the treatment of PH1
Coverage Duration	(PH1) (Initial, Reauth): 12 months
Other Criteria	(PH1) (Reauth): Member demonstrates positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SAMSCA

Products Affected

- SAMSCA
- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	Hyponatremia: (1) Members who are unable to sense or appropriately respond to thirst, (2) hypovolemic hyponatremia, (3) concomitant use of strong CYP3A inhibitors
Required Medical Information	Hyponatremia: (1) Diagnosis of clinically significant hypervolemic or euvolemic hyponatremia.
Age Restrictions	Hyponatremia: Member is 18 years of age or older.
Prescriber Restrictions	Hyponatremia: Prescribed by or in consultation with a cardiologist, endocrinologist or nephrologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SEROSTIM

Products Affected

- SEROSTIM SUBCUTANEOUS SOLUTION
RECONSTITUTED 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Wasting or Cachexia Associated with HIV (WC-HIV): (1) Diagnosis of wasting or cachexia associated with HIV. (2) Member is receiving concomitant antiretroviral therapy. (3) Nutritional evaluation since onset of wasting first occurred
Age Restrictions	
Prescriber Restrictions	(WC-HIV): Prescribed by or in consultation with a HIV specialist or infectious disease specialist
Coverage Duration	48 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SIGNIFOR

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cushing's disease (CD)(Initial): (1) Diagnosis of (pituitary) Cushing's disease. (2) Pituitary surgery is not an option or has not been curative
Age Restrictions	(CD)(Initial): Member is 18 years of age or older
Prescriber Restrictions	(CD)(Initial, Reauth): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(CD)(Reauth): Documentation of positive clinical response (e.g. clinically meaningful reduction in 24- hour urinary free cortisol levels, improvement in signs or symptoms of the disease)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SILDENAFIL

Products Affected

- REVATIO ORAL TABLET
- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*

PA Criteria	Criteria Details
Exclusion Criteria	(PAH, RP): Documentation of concomitant nitrate use
Required Medical Information	Pulmonary Arterial Hypertension (PAH): (1) Diagnosis of PAH WHO Group I with New York Heart Association (NYHA) Functional Class II to IV (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography (3) Mean pulmonary artery pressure greater than or equal 25 mm Hg at rest or greater than 30 mm Hg with exertion. Raynaud's Phenomenon (RP): (1) Diagnosis of secondary Raynaud's phenomenon. (2) Inadequate response or inability to tolerate a calcium channel blocker.
Age Restrictions	
Prescriber Restrictions	(PAH, RP): Prescribed by or in consultation with a Cardiologist or Pulmonologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SILIQ

Products Affected

- SILIQ

PA Criteria	Criteria Details
Exclusion Criteria	(PsO)(Initial, Reauth): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Plaque psoriasis (PsO)(Initial): (1) Diagnosis of moderate to severe PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) Skyrizi, (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate. (3) Member has been evaluated for depression and suicidal ideations using the PHQ-9.
Age Restrictions	(PsO)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(PsO)(Initial, Reauth): Prescribed by or in consultation with a dermatologist
Coverage Duration	(Initial): 16 weeks (Reauth) 1 year
Other Criteria	(PsO) (Reauth): (1) Member has positive response to therapy as evidenced by one of the following: (i) Reduction in the body surface area (BSA) involvement from baseline, (ii) Improvement in symptoms (e.g. pruritus, inflammation) from baseline , (2) Member has been evaluated for depression and suicidal ideations using the PHQ-9
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SIMPONI

Products Affected

- SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	(AS, PsA, RA, UC): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Ankylosing Spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate two of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) secukinumab (Cosentyx), (d) tofacitinib (Xeljanz/Xeljanz XR), (e) upadacitinib (Rinvoq), (f) adalimumab-adbm (Cyltezo), (g) adalimumab-aaty (Yuflyma) or documentation demonstrating that a trial may be inappropriate. Psoriatic Arthritis (PsA): (1) Diagnosis of active PsA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Xeljanz/Xeljanz XR, (d) Rinvoq/Rinvoq LQ, (e) Skyrizi, (f) Cosentyx, (g) Stelara, (h) Orencia, (i) Otezla, (j) Cyltezo, (k) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Rheumatoid arthritis (RA): (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Rinvoq, (d) Xeljanz/Xeljanz XR, (e) Orencia, (f) Cyltezo, (g) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Ulcerative Colitis (UC): (1) Diagnosis of moderate to severe UC. (2) Inadequate response or inability to tolerate two of the following: (a) Humira, and (b) Xeljanz/Xeljanz XR, (c) Stelara, (d) Rinvoq, (e) Cyltezo, (f) Yuflyma, (g) Skyrizi OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(AS, PsA, RA, UC): Member is 18 years of age or older
Prescriber Restrictions	(RA, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a rheumatologist or dermatologist. (UC): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

SIRTURO

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Multi-Drug Resistant Tuberculosis (MDR-TB): (1) Diagnosis of MDR-TB. (2) Member weighs at least 15kg (applies to members 5 to less than 18 years of age). (3) One of the following: (a) Medication will be used in combination with at least 3 other drugs to which the member's MDR-TB isolate has been shown to be susceptible in vitro OR (b) If in vitro testing results are unavailable, treatment will be initiated in combination with at least 4 other drugs to which the member's MDR-TB isolate is likely to be susceptible. (4) Sirturo (bedaquiline) will be administered by directly observed therapy (DOT).
Age Restrictions	(MDR-TB): Member is 5 years of age or older.
Prescriber Restrictions	(MDR-TB): Prescribed by or in consultation with infectious disease specialist or pulmonologist
Coverage Duration	24 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SKYCLARYS

Products Affected

- SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Friedreich Ataxia (FA)(Initial): (1) Diagnosis of Friedreich Ataxia (FA) confirmed by genetic testing demonstrating mutation in the FXN gene (2) Member has a Modified Friedreich's Ataxia Rating Scale (mFARS) score of greater than or equal to 20 and less than or equal to 80 (3) Member has B-type natriuretic peptide value less than or equal to 200 pg/ml
Age Restrictions	(FA)(Initial)(Reauth): member is 16 years of age or older
Prescriber Restrictions	(FA)(Initial)(Reauth): Prescribed by or in consultation with one of the following specialists: neurologist, neurogeneticist, Physiatrist (Physical Medicine and Rehabilitation Specialist)
Coverage Duration	(Initial): 12 months (Reauth): 12 months
Other Criteria	(FA)(Reauth): (1) Documentation of positive clinical response to therapy as evidenced by one of the following: (a) an increase in peak work (in Watts/kg) during exercise testing from baseline (b) a decrease in the rate of progression of Modified Friedreich's Ataxia Rating Scale (mFARS)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SKYRIZI SC

Products Affected

- SKYRIZI PEN
- SKYRIZI SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	(PsO, PsA, CD, UC): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Plaque psoriasis (PsO):Diagnosis of moderate to severe PsO. Psoriatic Arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least ONE DMARD. Crohn's Disease (CD): (1) Diagnosis of CD. (2) One of the following: (a) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g., corticosteroid, methotrexate, azathioprine or 6-mercaptopurine). (b) Will be used as a maintenance dose following the intravenous induction doses. Ulcerative Colitis (UC): (1) Diagnosis of moderately to severely active UC. (2) One of the following: (a) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. prednisone, mesalamine) (b) will be used as a maintenance dose following the intravenous induction doses.
Age Restrictions	(PsO, PsA, CD, UC): Member is 18 years of age or older
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist (PsA): Prescribed by or consultation with a rheumatologist or dermatologist (CD, UC): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	Indefinite
Other Criteria	Part B drug applies only to beneficiaries enrolled in an MA-PD plan.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SKYTROFA

Products Affected

- SKYTROFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Growth Failure in Children (GFC)(Initial): (1) Diagnosis of growth hormone deficiency confirmed by one of the following: (a) Height is documented by one of the following (utilizing age and gender growth charts related to height): (i) Height is greater than 2.0 standard deviations [SD] below midparental height (ii) Height is greater than 2.25 SD below population mean (below the 1.2 percentile for age and gender). (2) Growth velocity is greater than 2 SD below mean for age and gender. (3) Delayed skeletal maturation of greater than 2 SD below mean for age and gender (e.g. delayed is greater than 2 years compared with chronological age). (4) documentation of bone age, abnormal response from provocative testing, such as insulin-induced hypoglycemia test, levodopa, or clonidine.
Age Restrictions	(GFC) (Initial, Reauth): Member is 1 years of age or greater
Prescriber Restrictions	(GFC)(Initial): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	(Initial, Continuation): 12 months
Other Criteria	(GFC) (Continuation): (1) Annual clinical re-evaluation by the treating endocrinologist
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SOGROYA

Products Affected

- SOGROYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Growth Failure in Children (GFC)(Initial): (1) Diagnosis of growth hormone deficiency confirmed by one of the following: (a) Height is documented by one of the following (utilizing age and gender growth charts related to height): (i) Height is greater than 2.0 standard deviations [SD] below midparental height (ii) Height is greater than 2.25 SD below population mean (below the 1.2 percentile for age and gender). (b) Growth velocity is greater than 2 SD below mean for age and gender.(c) Delayed skeletal maturation of greater than 2 SD below mean for age and gender (e.g., delayed is greater than 2 years compared with chronological age).(2) documentation of bone age, (3) abnormal response from provocative testing, such as insulin-induced hypoglycemia test, levodopa, or clonidine. Diagnostically confirmed Growth Hormone Deficiency in adults (GHDA)(Initial)
Age Restrictions	(GFC) (Initial, Continuation): Member is 2.5 years of age or greater. (GHDA)(Initial, Continuation): Member is 18 years of age or older.
Prescriber Restrictions	(GFC, GHDA)(Initial, Continuation): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	(Initial, Continuation): 12 months
Other Criteria	(GFC) (Continuation): (1) Annual clinical re-evaluation by the treating endocrinologist (2) Growth velocity greater than or equal to 2.5 cm/year. (GHDA)(Continuation): Annual clinical re-evaluation by the treating endocrinologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SOHONOS

Products Affected

- SOHONOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(FOP)(Initial): (1) Diagnosis of Fibrodysplasia Ossificans Progressiva (FOP) (2) Molecular generic testing confirms mutations in the ACVR1 gene
Age Restrictions	(FOP) (Initial, Reauth): For female members: 8 years of age or older. For male members: 10 years of age or older
Prescriber Restrictions	(FOP) (Initial, Reauth): Prescribed by or in consultation with geneticist or orthopedic physician
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(FOP)(Reauth): (1) Documentation is provided that member demonstrates positive clinical response to therapy (e.g., reduction in volume in new abnormal bone growth)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SOTATERCEPT

Products Affected

- WINREVAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (Initial): (1) Diagnosis of PAH WHO Group I with New York Heart Association (NYHA) Functional Class II to III (2) inadequate response or inability to tolerate TWO of the following: (a) Endothelin Receptor Antagonist (bosentan, ambrisentan, macitentan) (b) Phosphodiesterase 5 inhibitor (tadalafil, sildenafil) (c) IV prostacyclin therapy (treprostinil, epoprostenol) (3) Member continues to receive other PAH therapies (e.g., ambrisentan, tadalafil)
Age Restrictions	
Prescriber Restrictions	(PAH) (Initial) (Reauth): Prescribed by or in consultation with cardiologist or pulmonologist.
Coverage Duration	(Initial) 6 months (Reauth) 12 months
Other Criteria	(PAH)(Reauth): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SOTYKTU

Products Affected

- SOTYKTU

PA Criteria	Criteria Details
Exclusion Criteria	Plaque Psoriasis (PsO): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	(PsO): (1) Diagnosis of moderate to severe PsO. (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira, (b) Enbrel, (c) Skyrizi (d) Cosentyx, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(PsO): Member is 18 years of age or older
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SPEVIGO SQ 2024

Products Affected

- SPEVIGO SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(GPP)(Initial): (1) Diagnosis of generalized pustular psoriasis (GPP) as defined by both of the following: (a) Primary, sterile, macroscopically visible pustules on non-acral skin (excluding cases where pustulation is restricted to psoriatic plaques) (b) Disease is relapsing (more than 1 episode) or persistent (more than 3 months) (2) Subcutaneous formulation will not be used to treat GPP flare (3) Member weighs at least 40 kg
Age Restrictions	(GPP) (Initial, Reauth): Member is 12 years of age or older
Prescriber Restrictions	(GPP): Prescribed by or in consultation with a dermatologist
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(GPP)(Reauth): Documentation of positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

STELARA SQ

Products Affected

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	(CD, UC, PsA, PsO): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), e.g. tumor necrosis factor antagonists
Required Medical Information	Part D is medically necessary when: Plaque psoriasis (PsO): Diagnosis of moderate to severe PsO. Psoriatic arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine). Crohn's Disease (CD): (1) Diagnosis of moderate to severe CD. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. corticosteroid, Aminosalicylate, azathioprine or 6-mercaptopurine). Ulcerative Colitis (UC): (1) Diagnosis of moderate to severe UC. (2) Inadequate response or inability to tolerate at least ONE conventional treatment: (e.g. corticosteroid, Aminosalicylate, azathioprine or 6-mercaptopurine).
Age Restrictions	(CD, UC): Member is 18 years of age or older. (PsO, PsA): Member is 6 years of age or older.
Prescriber Restrictions	(CD, UC): prescribed by or in consultation with a Gastroenterologist. (PsO): prescribed by or in consultation with a Dermatologist. (PsA): prescribed by or in consultation with a Dermatologist or Rheumatologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SUNOSI

Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NARCOLEPSY: (1) One of the following: (2) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT) (b) prescriber provides justification that a sleep study is not feasible, (2) Inadequate response or inability to tolerate modafinil or armodafinil. OBSTRUCTIVE SLEEP APNEA/HYPOPNEA SYNDROME (OSAHS): (1) Diagnosis confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), (2) documentation that the medication is being used as an adjunct treatment for the underlying obstruction, (3) inadequate response or inability to tolerate modafinil or armodafinil.
Age Restrictions	
Prescriber Restrictions	(Narcolepsy, OSAHS): Prescribed by or in consultation with a neurologist or sleep specialist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SYMDEKO

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis (CF): (1) Diagnosis of Cystic Fibrosis. (2) One of the following: a) Member is homozygous for the F508del mutation, or b) Member has at least one tezacaftor/ivacaftor responsive mutation in the CFTR gene. (3) If the member's genotype is unknown, an FDA-cleared test must be used to detect the presence of CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test
Age Restrictions	(CF): Member is 6 years of age or older
Prescriber Restrictions	(CF): Prescribed by or in consultation with a pulmonologist or Specialist affiliated with a CF care center
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

SYMLIN

Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION
PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION
PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	(DM): Gastroparesis.
Required Medical Information	Diabetes Mellitus (DM): (1) Diagnosis of diabetes (Type 1 or Type 2). (2) inadequate response to optimal insulin monotherapy. (3) concurrent use of mealtime insulin.
Age Restrictions	(DM): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TADLIQ

Products Affected

- TADLIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH)(Initial): (1) Diagnosis of PAH WHO Group I with New York Heart Association (NYHA) Functional Class II to IV, (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography, (3) Mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion, (4) inadequate response or inability to tolerate generic tadalafil
Age Restrictions	
Prescriber Restrictions	(PAH)(Initial): Prescribed by or in consultation with Cardiologist or Pulmonologist
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(PAH)(Reauth): (1) Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TAFAMIDIS

Products Affected

- VYNDAMAX
- VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM)(Initial): (1) Diagnosis of ATTR-CM confirmed by one of the following: (a) Member has a transthyretin (TTR) mutation (e.g. V122I), (b) cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits, or (c) all of the following: echocardiogram or cardiac magnetic resonance image suggestive of amyloidosis, scintigraphy scan suggestive of cardiac TTR amyloidosis, and absence of light-chain amyloidosis, (2) One of the following: (a) History of heart failure (HF), with at least one prior hospitalization for HF, or (b) presence of clinical signs and symptoms of HF (e.g. dyspnea, edema), (3) Member has New York Heart Association (NYHA) Functional Class I, II, or III heart failure
Age Restrictions	(ATTR-CM) (Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(ATTR-CM) (Initial, Reauth): Prescribed by or in consultation with a cardiologist
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(ATTR-CM)(Reauth): (1) Positive clinical response to therapy, (2) Member continues to have NYHA Functional Class I, II, or III heart failure.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TAKHZYRO

Products Affected

- TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when: Hereditary Angioedema (HAE): (1) Diagnosis of HAE. (2) For prophylaxis against HAE attacks.
Age Restrictions	
Prescriber Restrictions	(HAE): Prescribed by or in consultation with an allergist, immunologist, pulmonologist
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TALTZ

Products Affected

- TALTZ

PA Criteria	Criteria Details
Exclusion Criteria	(PsO, PsA, AS, nr-axSpA): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	<p>Plaque Psoriasis (PsO): (1) Diagnosis of moderate to severe PsO. (2) ONE of the following: (A) For members 6 to 17 years of age: an inadequate response or inability to tolerate two of the following: (a) Enbrel, (b) Cosentyx, (c) Stelara or documentation demonstrating that a trial may be inappropriate, OR (B) For members 18 years of age or older: an inadequate response or inability to tolerate two of the following: (a) Cosentyx (b) Enbrel (c) Humira (d) Skyrizi, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma or documentation demonstrating that a trial may be inappropriate. Psoriatic arthritis (PsA): (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate TWO of the following: (a) Enbrel (b) Humira (c) Cosentyx (d) Rinvoq/Rinvoq LQ (e) Skyrizi (f) Stelara (g) Xeljanz/Xeljanz XR, (h) Orencia, (i) Otezla, (j) Cyltezo, (k) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Ankylosing spondylitis (AS): (1) Diagnosis of AS. (2) Inadequate response or inability to tolerate TWO of the following: (a) adalimumab (Humira), (b) etanercept (Enbrel), (c) secukinumab (Cosentyx), (d) upadacitinib (Rinvoq), (e) tofacitinib (Xeljanz/Xeljanz XR), (f) adalimumab-adbm (Cyltezo), (g) adalimumab-aaty (Yuflyma) OR documentation demonstrating that a trial may be inappropriate. Non-Radiographic Axial Spondyloarthritis (nr-axSpA): (1) Diagnosis of nr-axSpA. (2) Inadequate response or inability to tolerate Rinvoq OR Cosentyx OR documentation demonstrating that a trial may be inappropriate (3) Inadequate response or inability to tolerate one NSAID (e.g. ibuprofen, meloxicam, naproxen).</p>
Age Restrictions	(PsA, AS, nraxSpA): Member is 18 years of age or older. (PsO): Member is 6 years of age or older.
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist. (AS, nr-axSpA): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	Indefinite

PA Criteria	Criteria Details
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TARPEYO

Products Affected

- TARPEYO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Immunoglobulin A nephropathy (IgAN): (1) Diagnosis of primary immunoglobulin A nephropathy (IgAN) (2) Member is at risk of rapid disease progression [e.g., generally a urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g, or by other criteria such as clinical risk scoring using the international IgAN Prediction Tool] (3) Used to reduce proteinuria (4) Estimated glomerular filtration rate (eGFR) greater than or equal to 35 ml/min/1.73 m ² (5) One of the following: (a) Member has been on a minimum 90-day trial of maximally tolerated dose and will continue to receive therapy with one of the following: (i) an angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril), (ii) An angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan), (b) Member is unable to tolerate both ACE inhibitors and ARBs (6) Inadequate response or inability to tolerate another glucocorticoid (e.g., prednisone, methylprednisolone)
Age Restrictions	(IgAN): Member is 18 years of age or older
Prescriber Restrictions	(IgAN): Prescribed by or in consultation with a nephrologist
Coverage Duration	9 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TAVALISSE

Products Affected

- TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Chronic Immune Thrombocytopenia (ITP)(Initial): (1) Documentation of baseline platelet count less than 30,000/mcL, (2) Inadequate response or inability to tolerate ONE of the following: (a) Corticosteroids, (b) Immunoglobulins, (c) Splenectomy, (d) Thrombopoietin receptor agonists (e.g. Nplate, Promacta), or (e) rituximab (Rituxan).
Age Restrictions	(ITP)(Initial, Continuation): Member is 18 years of age or older
Prescriber Restrictions	(ITP)(Initial, Continuation): Prescribed by or in consultation with hematologist/oncologist
Coverage Duration	(Initial, Continuation): 12 months
Other Criteria	(ITP)(Continuation): (1) Positive clinical response to Tavalisse therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	Yes

TAVNEOS

Products Affected

- TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Severe Active Anti-neutrophil Cytoplasmic Autoantibody (ANCA)-associated Vasculitis (granulomatosis with polyangiitis [GPA] and microscopic polyangiitis [MPA]) (ANCA-V(GPA)(MPA))(Initial): (1) Diagnosis of Severe Active Anti-neutrophil Cytoplasmic Autoantibody (ANCA)-associated Vasculitis (granulomatosis with polyangiitis [GPA] or microscopic polyangiitis [MPA]) AND 2) Used as adjunct to standard therapy, and glucocorticoids
Age Restrictions	(ANCA-V(GPA)(MPA))(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(ANCA-V(GPA)(MPA))(Initial, Reauth): Prescribed by or in consultation with a nephrologist, pulmonologist, or rheumatologist.
Coverage Duration	(Initial): 6 Months (Reauth): 1 year
Other Criteria	(ANCA-V(GPA)(MPA))(Reauth): (1) Positive clinical response to therapy defined as sustained remission as assessed by the Birmingham Vasculitis Activity Score (BVAS). (2) Used as adjunct to standard therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TERIPARATIDE

Products Affected

- FORTEO SUBCUTANEOUS SOLUTION PEN-INJECTOR 600 MCG/2.4ML
- *teriparatide subcutaneous solution pen-injector*
620 mcg/2.48ml

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Primary or Hypogonadal Osteoporosis (HGO)(Initial): (1) Diagnosis of HGO in men. (2) Member has severe osteoporosis or is at a high to very high risk for fracture as defined by ONE of the following: (a) Member has a history of multiple vertebral fractures, (b) T-score of the individual's bone mineral density (BMD) is at least -2.5 standard deviations below the young adult mean OR (c) history of osteoporotic fracture (i.e. hip, spine, etc.). (3) Inadequate response or inability to tolerate (a) bisphosphonates or (b) hormone replacement therapy. Postmenopausal Osteoporosis (PMO) (Initial): (1) Diagnosis of PMO. (2) Member has severe osteoporosis or is at a high to very high risk for fracture as defined by ONE of the following: (a) Member has a history of multiple vertebral fractures, (b) T score of the individual's bone mineral density (BMD) is at least -2.5 standard deviations below the young adult mean OR (c) history of osteoporotic fracture (i.e. hip, spine, etc.). (3) Inadequate response or inability to tolerate ONE of the following: (a) bisphosphonates, (b) hormone replacement therapy, (c) selective-estrogen receptor modulators (SERMs), OR (d) Denosumab (Prolia). Glucocorticoid Induced Osteoporosis (GCO)(Initial): (1) Diagnosis of CGO in men or women. (2) Daily dose of prednisone is greater than or equal to 5mg for at least 3 months. (3) Inadequate response or inability to tolerate ONE of the following: (a) bisphosphonates, (b) hormone replacement therapy, (c) selective-estrogen receptor modulators (SERMs), or (d) Denosumab (Prolia).</p>
Age Restrictions	(HGO, PMO, GCO) (Initial and Reauth): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial and Reauth): Remainder of contract year
Other Criteria	(HGO, PMO, GCO) (Reauth): One of the following: (1) Cumulative lifetime therapy does not exceed 2 years [applies to Teriparatide and Forteo], OR (2) member remains at or has returned to having a high risk for fracture despite a total of 24 months of use of parathyroid hormones [applies to Forteo only].

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TESTOSTERONE PRODUCTS

Products Affected

- AVEED
- DEPO-TESTOSTERONE INTRAMUSCULAR SOLUTION
- JATENZO
- *testosterone cypionate intramuscular solution 100 mg/ml, 200 mg/ml*
- *testosterone enanthate intramuscular solution*
- *testosterone transdermal gel 10 mg/act (2%), 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)*
- *testosterone transdermal solution*
- TLANDO
- XYOSTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Delayed Puberty (DP): (1) Diagnosis of delayed puberty in male members (Applies to Testosterone Enanthate only). Breast Cancer (BC): (1) Diagnosis of inoperable breast cancer in female members (Applies to Testosterone Enanthate only). Hypogonadism (HG)(New starts only): (1) Attestation that diagnosis was initially confirmed by ALL of the following: (A) Two early morning total testosterone levels below 300 ng/dL measured on separate occasions, (B) Normal Prolactin Level, and (C) Physical or cognitive symptoms of testosterone deficiency (e.g. physical: fatigue, sleep disturbances, decreased activity, cognitive: depressive symptoms, cognitive dysfunction, loss of concentration, poor memory, irritability), (2) ONE of the following (A) negative history of prostate and breast cancer OR (B) Both of the following (i) history of prostate cancer with stable PSA less than or equal to 4ng/dL for 2 years and (ii) documentation that the risk versus benefit has been assessed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(DP, BC): Indefinite (HG)(New Starts, Continuation): Indefinite
Other Criteria	(HG)(Continuation): ONE of the following (1) negative history of prostate and breast cancer OR (2) Both of the following (a) history of prostate cancer with stable PSA less than or equal to 4ng/dL for 2 years and (b) documentation that the risk versus benefit has been assessed.
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

TOPICAL CHEMO AGENTS

Products Affected

- *bexarotene*
- TARGRETIN
- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Approved when ANY of the following inclusion criteria is met: (1) Drug is FDA approved for indication and regimen requested, (2) The indication and regimen is classified as Category 1 or 2A by National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium, (3) narrative text in The American Hospital Formulary Service-Drug Information (AHFS-DI) or Clinical Pharmacology Compendium is supportive for the specific condition(s) requested, (4) The Micromedex Compendium and the strength of recommendation is listed as Class I, Class IIa, or Class IIb for the specific condition(s) requested, (5) Indication is listed in Lexi-Drugs as "off label" with evidence level A, (6) supported by Peer-Reviewed Medical Literature as defined in Chapter 15 Section 50.4.5 of the Medicare Benefit Policy Manual
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	Indefinite
Other Criteria	(All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TOPICAL RETINOID PRODUCTS

Products Affected

- *adapalene external cream*
- *adapalene external gel 0.3 %*
- *adapalene external pad*
- *adapalene-benzoyl peroxide external gel*
- AKLIEF
- ALTRENO
- ATRALIN
- CABTREO
- *clindamycin-tretinoin*
- DIFFERIN EXTERNAL CREAM
- DIFFERIN EXTERNAL GEL 0.3 %
- DIFFERIN EXTERNAL LOTION
- EPIDUO
- EPIDUO FORTE
- RETIN-A
- RETIN-A MICRO
- RETIN-A MICRO PUMP EXTERNAL GEL 0.06 %, 0.08 %
- *tretinoin external*
- *tretinoin microsphere external gel 0.04 %, 0.1 %*
- *tretinoin microsphere pump external gel 0.08 %*
- TWYNEO
- ZIANA

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TRACLEER

Products Affected

- *bosentan*
- TRACLEER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (Initial): (1) Documentation of a diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV. (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography. (3) Documentation of mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion. (4) Inadequate response or inability to tolerate bosentan (applies to brand Tracleer only)
Age Restrictions	
Prescriber Restrictions	(PAH) (Initial): Prescribed by or in consultation with a Cardiologist or Pulmonologist.
Coverage Duration	(Initial): 6 months. (Continuation):12 months.
Other Criteria	(PAH)(Continuation): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TREMFYA

Products Affected

- TREMFYA SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- TREMFYA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	(PsO, PsA): Concurrent therapy with biological DMARDs or other tumor necrosis factor antagonists
Required Medical Information	Plaque psoriasis (PsO): (1) Diagnosis or moderate to severe PsO. (2) Inadequate response or inability to tolerate two of the following: (a) Cosentyx, (b) Enbrel, (c) Humira, (d) Skyrizi, (e) Stelara, (f) Otezla, (g) Cyltezo, (h) Yuflyma OR documentation demonstrating that a trial may be inappropriate. Psoriatic arthritis (PsA): (1) Diagnosis of PsA, (2) Inadequate response or inability to tolerate TWO of the following: (a) Humira (b) Enbrel (c) Cosentyx (d) Rinvoq/Rinvoq LQ (e) Skyrizi (f) Stelara (g) Xeljanz/Xeljanz XR, (h) Orencia, (i) Otezla, (j) Cyltezo, (k) Yuflyma or documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(PsO, PsA): Member is 18 years of age or older
Prescriber Restrictions	(PsO): Prescribed by or in consultation with a dermatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TRIKAFTA

Products Affected

- TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis (CF): (1) Member has at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by an FDA-cleared cystic fibrosis mutation test OR a mutation in the CFTR gene that is responsive based on in vitro data.
Age Restrictions	(CF): Member is 2 years of age or older
Prescriber Restrictions	(CF): Prescribed by or in consultation with a pulmonologist or Specialist affiliated with a CF care center
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TYMLOS

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Postmenopausal Osteoporosis (PMO) (Initial): (1) Diagnosis of PMO. (2) Member has severe osteoporosis or is at a high to very high risk for fracture as defined by ONE of the following: (a) Member has a history of multiple vertebral fractures, (b) T score of the individual's bone mineral density (BMD) is at least -2.5 standard deviations below the young adult mean OR (c) history of osteoporotic fracture (i.e. hip, spine, etc.). (3) Inadequate response or inability to tolerate ONE of the following: (a) bisphosphonates, (b) hormone replacement therapy, (c) selective-estrogen receptor modulators (SERMs), OR (d) Denosumab (Prolia). Osteoporosis at high risk for fracture in men (OSTm) (Initial): (1) Diagnosis of primary or hypogonadal osteoporosis (2) Both of the following: (a) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) (b) One of the following (i) History of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm (ii) inadequate response or inability to tolerate at least one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab])
Age Restrictions	(PMO, OSTm) (Initial and Reauth): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial and Reauth): Remainder of contract year
Other Criteria	(PMO, OSTm)(Reauth): Cumulative lifetime therapy does not exceed 2 years
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TYRVAYA

Products Affected

- TYRVAYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Dry eye disease (DED) (Initial) (Reauth): (1) Diagnosis of dry eye disease confirmed by ONE of the following diagnostic tests (a) Schirmer test, (b) ocular surface dye staining (e.g. rose Bengal, fluorescein, lissamine green), (c) tear function index/fluorescein clearance test, (d) tear break up time, (e) tear film osmolarity, (f) slit lamp lid evaluation, (g) lacrimal gland function, (2) Inadequate response or inability to tolerate Cyclosporin 0.05% (Restasis)
Age Restrictions	
Prescriber Restrictions	(DED) (Initial) (Reauth): Prescribed by or in consultation with Ophthalmologist or Optometrist
Coverage Duration	(Initial) (Reauth): 12 months
Other Criteria	(DED) (Reauth): (1) Documentation of positive clinical response to therapy (e.g. increased tear production or improvement in dry eye symptoms).
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

TYVASO DPI

Products Affected

- TYVASO DPI MAINTENANCE KIT INHALATION
POWDER 16 MCG, 32 MCG, 48 MCG, 64 MCG
- TYVASO DPI TITRATION KIT INHALATION
POWDER 16 & 32 & 48 MCG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH) (Initial): (1) Diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV. (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography. (3) Mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion. Pulmonary Hypertension Associated with Interstitial Lung Disease (PH-ILD) (Initial): (1) Diagnosis of pulmonary hypertension associated with interstitial lung disease (PHILD) WHO Group 3. (2) Diagnosis confirmed by diagnostic test(s) (e.g., right heart catheterization, doppler echocardiogram, computerized tomography imaging)
Age Restrictions	
Prescriber Restrictions	(PAH)(PH-ILD)(Initial): Prescribed by or in consultation with a Cardiologist or Pulmonologist
Coverage Duration	(Initial): 6 months. (Continuation): 12 months.
Other Criteria	(PAH)(PH-ILD)(CONTINUATION): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

UBRELVY

Products Affected

- UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	(AM)(Initial, Reauth): Will be used for preventative treatment of migraine. Medication will be used in combination with another oral CGRP inhibitor.
Required Medical Information	Acute Treatment of Migraines (AM)(Initial): (1) Diagnosis of migraine with or without aura in adults. (2) Member has fewer than 15 headache days per month. (3) inadequate response or inability to tolerate ONE generic formulary triptan.
Age Restrictions	(AM)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(AM)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist.
Coverage Duration	(AM)(Initial): 6 months. (AM)(Reauth): 12 months
Other Criteria	(AM)(REAUTH): (1) Positive response to therapy (e.g. reduction in pain, photophobia, phonophobia, nausea)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

UPTRAVI

Products Affected

- UPTRAVI ORAL
- UPTRAVI TITRATION

PA Criteria	Criteria Details
Exclusion Criteria	(PAH)(Initial, Reauth): Taken in combination with a prostanoid/prostacyclin analogue (e.g. epoprostenol, iloprost, treprostinil)
Required Medical Information	Pulmonary Arterial Hypertension (PAH)(Initial): (1) Diagnosis of pulmonary arterial hypertension (PAH) WHO Group I with New York Heart Association (NYHA) Functional Class II to IV AND (2) Diagnosis confirmed by catheterization (right-heart or Swan-Ganz) or echocardiography AND (3) Documentation of mean pulmonary artery pressure greater than or equal to 25 mm Hg at rest or greater than 30 mm Hg with exertion AND (4) inadequate response or inability to tolerate TWO of the following (a) an endothelin receptor antagonist (Tracleer if naive to the class) (b) a phosphodiesterase inhibitor (sildenafil if naive to the class) (c) riociguat (Adempas).
Age Restrictions	
Prescriber Restrictions	(PAH)(Initial): Prescribed by or in consultation with a Cardiologist or Pulmonologist
Coverage Duration	(Initial): 6 months. (Reauth): 12 months.
Other Criteria	(PAH)(REAUTH): Stabilization or improvement as evaluated by a cardiologist or pulmonologist.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VECAMEYL

Products Affected

- VECAMEYL

PA Criteria	Criteria Details
Exclusion Criteria	(EHTN, MHTN): Uremia, glaucoma, organic pyloric stenosis, coronary insufficiency, recent myocardial infarction
Required Medical Information	Essential Hypertension (EHTN): (1) Diagnosis of moderately severe to severe essential hypertension (2) an inadequate response or inability to tolerate at least two antihypertensive medications in different classes. Malignant Hypertension (MHTN): (1) Diagnosis of malignant hypertension, (2) An inadequate response or inability to tolerate at least two antihypertensive medications in different classes.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VELSIPITY

Products Affected

- VELSIPITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Ulcerative Colitis (UC): (1) Diagnosis of moderately to severely active Ulcerative Colitis (2) Inadequate response or inability to tolerate TWO of the following (a) Humira (b) Xeljanz/Xeljanz XR (c) Stelara (d) Rinvoq (e) Cyltezo, (f) Yuflyma, (g) Skyrizi OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(UC): Member is 18 years of age or older
Prescriber Restrictions	(UC): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VEOZAH

Products Affected

- VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(VMS)(Initial): (1) Diagnosis of moderate to severe vasomotor symptoms due to menopause (2) Inadequate response or inability to tolerate one of the following (a) menopausal hormone therapy (e.g., estradiol tablets) (b) non-hormonal therapy (e.g., paroxetine, venlafaxine, clonidine, etc.)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(VMS)(Initial): 6 months, (Reauth): 12 months
Other Criteria	(VMS)(Reauth): (1) Documentation of positive clinical response to therapy (e.g., decrease in frequency and severity of vasomotor symptoms from baseline, etc.)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VERKAZIA

Products Affected

- VERKAZIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Initial): (1) Diagnosis of moderate to severe vernal keratoconjunctivitis confirmed by the presence of clinical signs and symptoms (e.g., itching, photophobia, giant papillae at the upper tarsal conjunctiva or at the limbus, thick mucus discharge, conjunctival hyperemia). (2) Inadequate response or inability to tolerate one of the following: (a) Topical ophthalmic "dual-acting" mast cell stabilizer and antihistamine (e.g., olopatadine, azelastine), (b) Topical ophthalmic mast cell stabilizers (e.g., cromolyn). (3) Inadequate response or inability to tolerate short term use (up to 2 to 3 weeks), of topical ophthalmic corticosteroids (e.g., dexamethasone, prednisolone, fluorometholone).
Age Restrictions	(Initial, reauth): Member is 4 years of age or older
Prescriber Restrictions	(Initial, reauth): Prescribed by or in consultation with an ophthalmologist or optometrist
Coverage Duration	(Initial): 6 months, (reauth): 12 months
Other Criteria	(Reauth): Documentation of positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms (e.g., itching, photophobia, papillary hypertrophy, mucus discharge, conjunctival hyperemia)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VIJOICE

Products Affected

- VIJOICE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PROS) (Initial): (1) Diagnosis of PIK3CA-Related Overgrowth Spectrum (PROS) (2) Documentation of mutation in the PIK3CA gene (3) Documentation of severe clinical manifestations (e.g., Congenital Lipomatous Overgrowth, Vascular malformations, Epidermal nevi, Scoliosis/skeletal and spinal [CLOVES], Facial Infiltrating Lipomatosis [FIL], Klippel-Trenaunay Syndrome [KTS], Megalencephaly-Capillary Malformation Polymicrogyria [MCAP])
Age Restrictions	(PROS) (Initial, Reauth): member is 2 years of age or older
Prescriber Restrictions	(PROS) (Initial, Reauth): Prescribed by or in consultation with a physician who specializes in the treatment of PROS
Coverage Duration	Initial: 6 months. Reauthorization: 12 months
Other Criteria	(PROS) (Reauth): (1) Documentation of positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VIVJOA

Products Affected

- VIVJOA

PA Criteria	Criteria Details
Exclusion Criteria	Recurrent vulvovaginal candidiasis (RVVC): Member is of reproductive potential
Required Medical Information	(RVVC): (1) Diagnosis of recurrent vulvovaginal candidiasis (RVVC) (2) Diagnosis of RVVC confirmed by one of the following: (a) positive potassium hydroxide (KOH) preparation (b) vaginal fungal culture (3) Member has experienced 3 or more symptomatic episodes of vulvovaginal candidiasis (VVC) within the past 12 months (4) Inadequate response or inability to tolerate BOTH of the following: (a) one intravaginal product (e.g., clotrimazole, miconazole, terconazole) (b) oral fluconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	4 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VOQUEZNA

Products Affected

- VOQUEZNA DUAL PAK
- VOQUEZNA TRIPLE PAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(H. pylori): (1) Diagnosis of Helicobacter pylori infection (2) Inadequate response or inability to tolerate ONE of the following first line treatment regimens: (a) Clarithromycin based therapy (e.g., clarithromycin based triple therapy, clarithromycin based concomitant therapy), (b) Bismuth quadruple therapy (e.g., bismuth and metronidazole and tetracycline and proton pump inhibitor [PPI])
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VOQUEZNA TABLETS

Products Affected

- VOQUEZNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(H. pylori): (1) Diagnosis of Helicobacter pylori infection (2) One of the following: (a) Used in combination with amoxicillin and clarithromycin for the treatment of H. pylori infection OR (b) Used in combination with amoxicillin for the treatment of H. pylori infection (3) An inadequate response or inability to tolerate ONE of the following: (a) Clarithromycin based therapy (e.g., clarithromycin based triple therapy, clarithromycin based concomitant therapy) OR (b) Bismuth quadruple therapy (e.g., bismuth and metronidazole and tetracycline and proton pump inhibitor [PPI]). Erosive Esophagitis: (1) ONE of the following: (a) Used for healing of all grades of erosive esophagitis and relief of heartburn associated with erosive esophagitis OR (b) Used to maintain healing and relief of heartburn associated with erosive esophagitis (2) An inadequate response or inability to tolerate TWO of the following generic proton pump inhibitors (PPI's): omeprazole, esomeprazole, pantoprazole, lansoprazole, rabeprazole, and dexlansoprazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(H. pylori): 1 month. Healing erosive esophagitis (EE): 8 weeks. Maintenance of EE: 6 months.
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VOWST

Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Prevention of the recurrence of Clostridioides difficile infection (PCDI): (1) Diagnosis of recurrent Clostridioides difficile infection as defined by both of the following: (a) Presence of diarrhea defined as passage of 3 or more loose bowel movements within a 24-hour period for at least 2 consecutive days (b) a positive stool test for C. difficile toxin or toxigenic C. difficile (2) Member has a history of two or more recurrent episodes of CDI within 12 months (3) Member has completed at least 10 consecutive days of one of the following antibiotic therapies 2-4 days prior to initiating Vowst: (a) oral vancomycin (b) Difacid (fidaxomicin) (4) Member has completed the recommended bowel prep (e.g. 296mL of magnesium citrate) the day before and at least 8 hours prior to initiating Vowst (5) Previous episode of CDI is under control (e.g., less than 3 unformed or loose [i.e., Bristol Stool Scale type 6-7] stools per day for at least 2 consecutive days)
Age Restrictions	(PCDI): Member is 18 years of age or older
Prescriber Restrictions	(PCDI): Prescribed by or in consultation with gastroenterologist or infectious disease specialist
Coverage Duration	(PCDI): 14 days
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VOXZOGO

Products Affected

- VOXZOGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Achondroplasia) (Initial): (1) Member has open epiphyses, (2) Diagnosis of achondroplasia as confirmed by both of the following, (i) Member has clinical manifestations characteristic of achondroplasia (e.g. macrocephaly, frontal bossing, midface retrusion, disproportionate short stature with rhizomelic shortening of the arms and the legs, brachydactyly, trident configuration of the hands, thoracolumbar kyphosis, and accentuated lumbar lordosis), (ii) Member has radiographic findings characteristic of achondroplasia (e.g. large calvaria and narrowing of the foramen magnum region, undertubulated, shortened long bones with metaphyseal abnormalities, narrowing of the interpedicular distance of the caudal spine, square ilia and horizontal acetabula, small sacrosciatic notches, proximal scooping of the femoral metaphysis, and short and narrow chest)
Age Restrictions	
Prescriber Restrictions	(Achondroplasia) (Initial) (Reauth): Prescribed by or in consultation with one of the following: (1) clinical geneticist, (2) endocrinologist, (3) a physician who has specialized expertise in the management of achondroplasia
Coverage Duration	(Initial) (Reauth): 12 months
Other Criteria	(Achondroplasia) (Reauth): (1) member has open epiphyses, (2) Documentation of positive clinical response to therapy [e.g. improvement in annualized growth velocity (AGV) compared to baseline].
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VOYDEYA 2024

Products Affected

- VOYDEYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Paroxysmal nocturnal hemoglobinuria (PNH)(Initial): (1) Diagnosis of Paroxysmal nocturnal hemoglobinuria (PNH) (2) Medication will be used as add-on therapy to ravulizumab (Ultomiris) or eculizumab (Soliris) (3) Hemoglobin levels are less than or equal to 9.5 g/dL (4) Absolute reticulocyte count greater than or equal to 120 x 10 ⁹ /L.
Age Restrictions	
Prescriber Restrictions	(PNH)(Initial, Continuation): Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	(Initial, Continuation): 12 months
Other Criteria	(PNH)(Continuation): (1) Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) (2) Positive clinical response to therapy (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) (3) Will be used as add-on therapy to ravulizumab (Ultomiris) or eculizumab (Soliris)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VTAMA

Products Affected

- VTAMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PsO) (Initial): (1) Diagnosis of plaque psoriasis (2) Inadequate response or inability to tolerate TWO of the following topical therapies for sufficient duration (e.g., minimum of 4 weeks): (a) corticosteroids (e.g., betamethasone, clobetasol) (b) Vitamin D analogs (e.g., calcitriol, calcipotriene) (c) Concurrent combination of Vitamin D analog and corticosteroid (e.g., Enstilar, Taclonex) (d) tazarotene (e) Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
Age Restrictions	
Prescriber Restrictions	(PsO) (Initial, Reauth): Prescribed by or in consultation with a dermatologist
Coverage Duration	(Initial): 6 months. (Reauthorization): 12 months
Other Criteria	(PsO) (Reauthorization): (1) Documentation of positive clinical response to therapy (e.g., reduction in the body surface area (BSA) involvement from baseline, improvement in symptoms such as pruritus or inflammation from baseline)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VUITY

Products Affected

- VUITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(Presbyopia) (Initial): (1) Diagnosis of presbyopia (2) Provider confirms valid clinical rationale, which excludes lifestyle choice, as to why patient is unable to use corrective lenses (e.g., eyeglasses or contact lenses)
Age Restrictions	
Prescriber Restrictions	(Presbyopia) (Initial) (Reauth): prescribed by or in consultation with one of the following (1) Ophthalmologist, (2) Optometrist
Coverage Duration	(Initial): 3 months, (Reauth): 6 months
Other Criteria	(Presbyopia) (Reauth): (1) Documentation of positive clinical response to therapy (e.g. improvement in near vision in low light conditions without loss of distance vision)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

VYVANSE

Products Affected

- *lisdexamfetamine dimesylate*
- VYVANSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Attention Deficit Hyperactivity Disorder (ADHD): (1) Diagnosis of ADHD (2) inadequate response or inability to tolerate one of the following: immediate release formulations of amphetamine, dextroamphetamine, or methylphenidate Binge Eating Disorder (BED): (1)Diagnosis of BED. (2) Member has BED for 3 months or longer.
Age Restrictions	(ADHD): Member is 6 years of age or older (BED): Member is 18 years of age or older.
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

WAINUA 2024

Products Affected

- WAINUA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(hATTR amyloidosis)(Initial): (1) Diagnosis of hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) with polyneuropathy (2) Member has a transthyretin (TTR) mutation (e.g., V30M) (3) One of the following: (a) Member has a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2 (b) member has a baseline neuropathy impairment score (NIS) greater than or equal to 10 and less than or equal to 130 (c) Member has a baseline Karnofsky Performance Status score greater than 50 percent (4) Presence of clinical signs and symptoms of the disease (e.g., neuropathy)
Age Restrictions	
Prescriber Restrictions	(hATTR amyloidosis)(Initial, Cont): Prescribed by or in consultation with a neurologist
Coverage Duration	(hATTR amyloidosis)(Initial, Cont): 12 months
Other Criteria	(hATTR amyloidosis)(Cont): (1) Member demonstrates positive clinical response to therapy (2) One of the following: (a) Member continues to have a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2 (b) member continues to have a baseline neuropathy impairment score (NIS) greater than or equal to 10 and less than or equal to 130 (c) Member continues to have a baseline Karnofsky Performance Status score greater than 50 percent
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

WAKIX

Products Affected

- WAKIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Narcolepsy type 1: (1) Diagnosis of cataplexy with narcolepsy (Type 1). (2) ONE of the following: (a) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT), (b) prescriber provides justification confirming that a sleep study would not be feasible and (3) Symptoms of cataplexy are present. Narcolepsy (Type 2): (1) Diagnosis of excessive daytime sleepiness in Narcolepsy (Type 2). (2) ONE of the following: (a) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT), (b) prescriber provides justification confirming that a sleep study would not be feasible and (3) Inadequate response or inability to tolerate modafinil or armodafinil.
Age Restrictions	
Prescriber Restrictions	(Narcolepsy): Prescribed by or in consultation with a neurologist or sleep specialist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

WEGOVY

Products Affected

- WEGOVY

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of other GLP-1 receptor agonist (e.g., Adlyxin, Byetta, Bydureon, Mounjaro, Ozempic, Rybelsus, Trulicity, Victoza)
Required Medical Information	(Initial): (1) Documentation of both of the following: (a) BMI greater than or equal to 27 kg/m ² (b) member has a history of cardiovascular disease, as evidenced by at least one of the following: (i) prior myocardial infarction (ii) prior stroke (ischemic and hemorrhagic stroke) (iii) symptomatic peripheral arterial disease, as evidenced by intermittent claudication with ankle-brachial index less than 0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease and (2) Documentation showing member is on guideline therapy management for multiple risk factors (e.g., dyslipidemia, hypertension) associated with cardiovascular disease as evidenced by presence of all of the following: (a) one of the following: (i) platelet aggregation inhibitor (e.g., acetylsalicylic acid, P2Y ₁₂ receptor inhibitors) or anti-thrombotic medication (e.g., vitamin K antagonists, direct oral anticoagulants) (ii) member has inability to tolerate platelet aggregation inhibitor or anti-thrombotic medication (b) one of the following: (i) lipid-lowering medications (e.g., statins, ezetimibe, fibrate, PCSK-9 inhibitors) or (ii) member has inability to tolerate lipid-lowering medications (c) one of the following: (i) other cardiac medications including beta blocker, angiotensin-converting-enzyme inhibitor, angiotensin-receptor blocker, calcium channel blocker (ii) member has inability to tolerate other cardiac medications including beta blocker, angiotensin-converting-enzyme inhibitor, angiotensin-receptor blocker, calcium channel blocker (3) Member does not have a history of Type 1 or Type 2 Diabetes Mellitus
Age Restrictions	Member is 45 years of age or older
Prescriber Restrictions	
Coverage Duration	(Initial, Reauth): End of contract year

PA Criteria	Criteria Details
Other Criteria	<p>(1) Documentation showing member is on guideline therapy management for multiple risk factors (e.g., dyslipidemia, hypertension) associated with cardiovascular disease as evidenced by presence of all of the following: (a) one of the following: (i) platelet aggregation inhibitor (e.g., acetylsalicylic acid, P2Y12 receptor inhibitors) or anti-thrombotic medication (e.g., vitamin K antagonists, direct oral anticoagulants) (ii) member has inability to tolerate platelet aggregation inhibitor or anti-thrombotic medication (b) one of the following: (i) lipid-lowering medications (e.g., statins, ezetimibe, fibrates, PCSK-9 inhibitors) or (ii) member has inability to tolerate lipid-lowering medications (c) one of the following: (i) other cardiac medications including beta blocker, angiotensin-converting-enzyme inhibitor, angiotensin-receptor blocker, calcium channel blocker (ii) member has inability to tolerate other cardiac medications including beta blocker, angiotensin-converting-enzyme inhibitor, angiotensin-receptor blocker, calcium channel blocker (2) Documentation is provided confirming one of the following: (a) Member has not experienced a cardiovascular event since starting treatment (i.e., nonfatal myocardial infarction, nonfatal stroke) (b) Member has experienced a non-fatal myocardial infarction or nonfatal stroke since starting treatment, but the prescriber attests the benefit outweighs risk for member to continue treatment with semaglutide (Wegovy)</p>
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

WILSONS DISEASE

Products Affected

- CUVRIOR
- *trientine hcl oral capsule 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Wilson's disease (WD): (1) Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration) (2) Inadequate response or inability to tolerate a penicillamine product (e.g., Depen, Cuprimine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XDEMVY

Products Affected

- XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of treatment of demodex blepharitis
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with an ophthalmologist or optometrist
Coverage Duration	6 weeks
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XELJANZ

Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	(RA, PsA, UC, PJI, AS): Concurrent therapy with any other biologic disease modifying anti-rheumatic drug (DMARD), Janus kinase (JAK) inhibitors or potent immunosuppressants (e.g. azathioprine, cyclosporine)
Required Medical Information	Rheumatoid arthritis (RA) [applies to Xeljanz/Xeljanz XR tablets]: (1) Diagnosis of moderate to severe RA. (2) Inadequate response or inability to tolerate one DMARD: (e.g. methotrexate, hydroxychloroquine, leflunomide, azathioprine, sulfasalazine), (3) Member had inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate. Psoriatic Arthritis (PsA) [applies to Xeljanz/Xeljanz XR tablets]: (1) Diagnosis of PsA. (2) Inadequate response or inability to tolerate at least ONE DMARD: (e.g. Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine) (3) Inadequate response or inability to tolerate one or more TNF inhibitors (e.g. Enbrel, Humira, Cyltezo, Yuflyma) or documentation demonstrating that a trial may be inappropriate. Ulcerative Colitis (UC) [applies to Xeljanz/Xeljanz XR tablets]: (1) Diagnosis of moderate to severe UC. (2) Inadequate response or inability to tolerate at least ONE conventional therapy (e.g. corticosteroid, aminosalicylate, azathioprine or 6-mercaptopurine) (3) Inadequate response or inability to tolerate one or more TNF inhibitors (e.g. adalimumab (Humira), adalimumab-adbm (Cyltezo, adalimumab-aaty (Yuflyma)) or documentation demonstrating that a trial may be inappropriate. Polyarticular Juvenile Idiopathic Arthritis (PJI) [applies to Xeljanz tablets/oral solution]: (1) Diagnosis of PJI. (2) Inadequate response or inability to tolerate one or more TNF inhibitors (e.g. adalimumab (Humira), etanercept (Enbrel), adalimumab-adbm (Cyltezo), adalimumab-aaty (Yuflyma)) OR documentation demonstrating that a trial may be inappropriate. (3) Inadequate response or inability to tolerate ONE of the following: Methotrexate, Hydroxychloroquine, Leflunomide, Azathioprine, Sulfasalazine.
Age Restrictions	(RA, PsA, UC, AS): Member is 18 years of age or older. (PJI): Member is 2 years of age or older.
Prescriber Restrictions	(RA, PJI, AS): Prescribed by or in consultation with a rheumatologist. (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist. (UC): Prescribed by or in consultation with a gastroenterologist.

PA Criteria	Criteria Details
Coverage Duration	Indefinite
Other Criteria	Ankylosing Spondylitis (AS): (1) Diagnosis of ankylosing spondylitis. (2) Inadequate response or inability to tolerate one or more TNF inhibitors (e.g. adalimumab (Humira), adalimumab-adbm (Cyltezo), adalimumab-aaty (Yuflyma), etanercept (Enbrel)) OR documentation that a trial may be inappropriate.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XENAZINE

Products Affected

- *tetrabenazine*
- XENAZINE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Tardive Dyskinesia (TD)(Initial): (1) Diagnosis of TD, (2) Documentation is provided of ONE of the following: (a) Persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering or discontinuation of the offending medication, or (b) Member is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Tourette's Syndrome (TS): (1) Diagnosis of TS, (2) Member has tics associated with Tourette's syndrome, (3) Inadequate response or inability to tolerate haloperidol or risperidone. Chorea- Huntington's Disease (CHD): (1) Diagnosis of chorea associated with Huntington's disease
Age Restrictions	
Prescriber Restrictions	(TS, CHD): Prescribed by or in consultation with a neurologist or a psychiatrist. (TD)(Initial, Reauth): Prescribed by or in consultation with a neurologist or a psychiatrist.
Coverage Duration	(TD)(Initial): 3 months. (TD)(Reauth): Indefinite. (TS, CHD): Indefinite.
Other Criteria	(TD)(Reauth): Positive clinical response to therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XERMELO

Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Carcinoid Syndrome Diarrhea (CSD)(Initial): (1) Diagnosis of carcinoid syndrome diarrhea, (2) Diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g. octreotide [Sandostatin, Sandostatin LAR], lanreotide [Somatuline Depot]) for at least 3 months, (3) Used in combination with SSA therapy
Age Restrictions	(CSD)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(CSD)(Initial, Reauth): Prescribed by or in consultation with an Oncologist, Endocrinologist, or Gastroenterologist
Coverage Duration	(Initial): 12 months (Reauth): Indefinite
Other Criteria	(CSD)(Reauth): (1) Positive clinical response to Xermelo therapy, (2) Xermelo will continue to be used in combination with SSA therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XGEVA

Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Part D is medically necessary when: Prevention of Skeletal Related Events in Multiple Myeloma or Bone Metastases from Solid Tumors (MM-BMST) : (1) For prevention of skeletal related events in multiple myeloma or bone metastases from solid tumors. Giant Cell Tumor of the Bone (GCTB): (1) Diagnosis of GCTB. (2) Member is (a) adult or (b) adolescent that is skeletally mature. (3) Tumor is unresectable or surgical resection is likely to result in severe morbidity. Hypercalcemia of Malignancy Refractory to Bisphosphonates (HCMRB): (1) Diagnosis of HCMRB.
Age Restrictions	
Prescriber Restrictions	(GCTB, HCMRB): Prescribed by or in consultation with an urologist, oncologist or hematologist.
Coverage Duration	Indefinite
Other Criteria	Subject to Part B vs Part D review. (All Indications): Approve if for continuation of therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XIFAXAN

Products Affected

- XIFAXAN ORAL TABLET 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Prophylaxis for Hepatic Encephalopathy (HE): (1) Diagnosis of hepatic disease with risk for hepatic encephalopathy (i.e., previous episode of hepatic encephalopathy, advanced liver disease, hepatocellular carcinoma), (2) Inadequate response or inability to tolerate lactulose. Irritable Bowel Syndrome (IBS): (1) Diagnosis of irritable bowels syndrome- diarrhea, (2) Inadequate response or inability to tolerate BOTH of the following: (A) ONE of the following: (i) ONE Tricyclic antidepressant or (ii)selective serotonin reuptake inhibitor and (B) dicyclomine (3): BOTH of the following: (A) Member does not exceed 3 total courses (42 days in total) of therapy, (B) Member experiences irritable bowel syndrome with diarrhea (IBS-D) symptoms. Small Bowel Bacterial Overgrowth (SBBO)/Small Intestinal Bacterial Overgrowth (SIBO) (Initial): (1) Diagnosis of Small Bowel Bacterial Overgrowth (SBBO)/Small Intestinal Bacterial Overgrowth (SIBO)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	(HE): Indefinite. (IBS): 2 weeks. (SBBO/SIBO): Initial and Reauth: 2 weeks.
Other Criteria	(SBBO/SIBO) (Reauth): (1) Documentation of Small Bowel Bacterial Overgrowth (SBBO)/Small Intestinal Bacterial Overgrowth (SIBO) recurrence following successful initial treatment
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XOLAIR

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Part D is medically necessary for: Moderate to Severe Persistent Allergic Asthma (PAA)(Initial): (1) DIAGNOSIS OF MODERATE TO SEVERE PERSISTENT ALLERGIC ASTHMA, (2) The individual has a positive skin test or in vitro reactivity to a perennial aeroallergen, (3) The individual has a baseline serum IgE level of between 30 IU/mL and 1300 IU/mL, (4) Inadequate response or inability to tolerate a combination of high-dose inhaled corticosteroids (ICS)with a long-acting beta-agonist (LABA). Chronic Urticaria (CU)(Initial): (1) DIAGNOSIS OF CHRONIC URTICARIA, (2) an inadequate response, contraindication or inability to tolerate ONE second- generation H1 antihistamine at the maximally tolerated dose in addition to ANY of the following: (a) leukotriene receptor antagonist (e.g. montelukast) (b) histamine H2-receptor antagonist (e.g. ranitidine, cimetidine, famotidine), (c) substituting to a different second- generation antihistamine, (d) systemic glucocorticosteroids or (e) cyclosporine, (3) will be used concurrently with an h1 antihistamine, unless there is contraindication or intolerance to H1 antihistamines. Nasal Polyps (NP)(Initial): (1) Diagnosis of nasal polyps. (2) Member will use concurrently with nasal corticosteroid. (3) Member had inadequate response or inability to tolerate an intranasal corticosteroid. (4) Member has a baseline serum IgE level of between 30 IU/mL and 1500 IU/mL</p>
Age Restrictions	<p>(PAA)(Initial, Reauth): Member is 6 years of age or older (CU)(Initial, Reauth): Member is 12 years of age and older (NP)(Initial, Reauth): Member is 18 years of age and older (IMFA)(Initial, Reauth): Member is 1 year of age and older</p>
Prescriber Restrictions	<p>(PAA)(Initial, Reauth): Prescribed by or in consultation with an Allergist, Immunologist, or Pulmonologist. (CU)(Initial, Reauth): Prescribed by or in consultation with allergist/immunologist, dermatologist. (NP)(Initial, Reauth): Prescribed by or in consultation with an allergist, immunologist, pulmonologist, or ENT specialist. (IMFA)(Initial, Reauth): Prescribed by or in consultation with an Allergist or Immunologist</p>
Coverage Duration	<p>(Initial): 12 months. (Reauth): 12 months.</p>

PA Criteria	Criteria Details
Other Criteria	<p>Subject to Part B vs Part D review. IgE-Mediated Food Allergy (IMFA) (Initial): (1) One of the following: (A) Both of the following (i) Diagnosis of IgE-Mediated Food Allergy (ii) Clinical history of IgE Mediated Food Allergy (B) Documentation that the member has a history of severe allergic response, including anaphylaxis, following exposure to one or more foods (2) Used in conjunction with food allergen avoidance (3) Both of the following (A) Baseline (pre- Xolair treatment) serum total IgE level is greater than or equal to 30 IU/mL and less than or equal to 1850 IU/mL (B) Dosing is according to serum total IgE levels and body weight (PAA)(Reauth): (1) Documentation of positive clinical response to therapy (e.g. reduction in number of asthma exacerbations, improvement in forced expiratory volume in 1 second (FEV1), or decreased use of rescue medications). (CU)(Reauth): (1) Member's disease status has been re- evaluated since the last authorization to confirm the member's condition warrants continued treatment (2) member has experienced one or both of the following: (a) Reduction in itching severity from baseline (b) Reduction in number of hives from baseline. (NP)(Reauth): (1) Documentation of positive clinical response to therapy (e.g. reduction in nasal polyps score [NPS: 0-8 scale], improvement in nasal congestion/obstruction score [NCS: 0-3 scale]) (2) Used in combination with another agent for nasal polyps. (IMFA)(Reauth) (1) Documentation of positive clinical response to therapy (2) Used in conjunction with food allergen avoidance. (3) Dosing will continue to be based on body weight and pretreatment total IgE serum levels</p>
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XOLREMDI 2024

Products Affected

- XOLREMDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(WHIM)(Initial): (1) Diagnosis of WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome (2) Member has genotype confirmed variant of CXCR4 as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA) (3) Member has an absolute neutrophil count (ANC) less than or equal to 500 cells /uL
Age Restrictions	
Prescriber Restrictions	(WHIM) (Initial) (Reauth): Prescribed by or in consultation with immunologist, hematologist, geneticist or allergist
Coverage Duration	(Initial) 6 months (Reauth) 12 months
Other Criteria	(WHIM)(Reauth): Documentation of positive clinical response to therapy (e.g., improvement in ANC, reduction in infections)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XYREM

Products Affected

- *sodium oxybate*
- XYREM

PA Criteria	Criteria Details
Exclusion Criteria	CN, EDSN (Initial, Reauth): Concurrent use of sedative hypnotics and alcohol
Required Medical Information	Cataplexy in Narcolepsy (CN)(Initial): (1) Diagnosis of cataplexy in narcolepsy, (2) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT) or prescriber provides justification that a sleep study is not feasible. Excessive Daytime Sleepiness in Narcolepsy (EDSN)(Initial): (1) Diagnosis of EDSN (Narcolepsy Type 2), (2) Diagnosis Confirmed by Polysomnography (PSG) or Multiple sleep latency test (MSLT) or prescriber provides justification that a sleep study is not feasible and (3) Inadequate response or inability to modafinil or armodafinil. (adult use only)
Age Restrictions	
Prescriber Restrictions	(CN, EDSN)(Initial, Reauth): Prescribed by or in consultation with a neurologist, psychiatrist, or sleep specialist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(CN, EDSN)(Reauth): (1) Documentation to support the efficacy associated with the current regimen (including but not limited to reduction in the frequency of cataplexy attacks or an improvement in the Epworth sleepiness scale), (2) Member is re-evaluated every 12 months.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

XYWAV

Products Affected

- XYWAV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Cataplexy in Narcolepsy (CN)(Initial): (1) Diagnosis of cataplexy in narcolepsy. Excessive Daytime Sleepiness in Narcolepsy (EDSN)(Initial): (1) Diagnosis of EDSN (Narcolepsy Type 2), (2) Inadequate response or inability to modafinil. Idiopathic Hypersomnia (IH) (Initial): (1) Diagnosis of Idiopathic Hypersomnia as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), (2) Symptoms of excessive daytime sleepiness (e.g. nap duration of longer than 60 minutes) are present
Age Restrictions	
Prescriber Restrictions	(CN, EDSN, IH)(Initial, Reauth): Prescribed by or in consultation with a neurologist, psychiatrist, or sleep specialist.
Coverage Duration	(Initial, Reauth): 12 months
Other Criteria	(CN, EDSN)(Reauth): (1) Documentation to support the efficacy associated with the current regimen (including but not limited to reduction in the frequency of cataplexy attacks or an improvement in the Epworth sleepiness scale), (2) Member is re-evaluated every 12 months. (IH) (Reauth): (1) Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZAVESCA

Products Affected

- *miglustat*
- YARGESA
- ZAVESCA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Type 1 Gaucher's Disease (T1GD): (1) Diagnosis of mild to moderate T1GD, (2) enzyme replacement therapy is not a therapeutic option (e.g. because of allergy, hypersensitivity, or poor venous access).
Age Restrictions	(T1GD): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Remainder of contract year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZAVZPRET

Products Affected

- ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	(AM)(Initial, Reauth): Will be used for preventative treatment of migraine. Medication will be used in combination with another oral CGRP inhibitor
Required Medical Information	Acute Treatment of Migraines (AM)(Initial): (1) Diagnosis of migraine with or without aura in adults. (2) Member has fewer than 15 headache days per month. (3) inadequate response or inability to tolerate ONE generic formulary triptan (4) Inadequate response or inability to tolerate ONE of the following: (a) Nurtec ODT (b) Ubrelvy
Age Restrictions	(AM)(Initial, Reauth): Member is 18 years of age or older
Prescriber Restrictions	(AM)(Initial, Reauth): Prescribed by or in consultation with a neurologist or headache specialist certified by the United Council for Neurologic Subspecialties, or pain specialist.
Coverage Duration	(AM)(Initial): 6 months. (AM)(Reauth): 12 months
Other Criteria	(AM)(REAUTH): (1) Positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZEPOSIA

Products Affected

- ZEPOSIA
- ZEPOSIA 7-DAY STARTER PACK
- ZEPOSIA STARTER KIT ORAL CAPSULE
THERAPY PACK 0.23MG & 0.46MG 0.92MG(21)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Multiple Sclerosis (MS): (1) Diagnosis of relapsing form of multiple sclerosis (MS) (e.g. clinically isolated syndrome, relapsing remitting disease, secondary progressive disease, including active disease with new brain lesions) and inadequate response or inability to tolerate two of the following medications: (a) Avonex (interferon beta-1a), (b) Plegridy (peginterferon beta-1a), (c) Betaseron (interferon beta-1b), (d) Glatopa (glatiramer acetate), (e) Tecfidera (Dimethyl Fumarate), (f) Gilenya (fingolimod), (g) Aubagio (teriflunomide), or (h) Rebif (interferon beta 1a), OR documentation demonstrating that a trial may be inappropriate. Ulcerative Colitis (UC): (1) Diagnosis of ulcerative colitis, (2) Inadequate response or inability to tolerate TWO of the following (a) Humira (b) Xeljanz/Xeljanz XR (c) Stelara (d) Rinvoq (e) Cyltezo, (f) Yuflyma, (g) Skyrizi OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(MS) (UC): Member is 18 years of age or older
Prescriber Restrictions	UC): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZILBRYSQ 2024

Products Affected

- ZILBRYSQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(gMG)(Initial): (1) Diagnosis of generalized myasthenia gravis (gMG) (2) Member is anti-acetylcholine receptor (AChR) antibody positive (3) One of the following: (a) Inadequate response or inability to tolerate two immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus) (b) Both of the following: (i) Inadequate response or inability to tolerate one immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus) (ii) Inadequate response or inability to tolerate one of the following: (1) Chronic plasmapheresis or plasma exchange (PE) (2) Intravenous immunoglobulin (IVIG)
Age Restrictions	
Prescriber Restrictions	(gMG) (Initial, Reauth): Prescribed by or in consultation with a neurologist
Coverage Duration	(gMG) (Initial, Reauth): 12 months
Other Criteria	(gMG) (Reauth): Member demonstrates positive clinical response to therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZOKINVY

Products Affected

- ZOKINVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Hutchinson-Gilford Progeria Syndrome (HGPS): (1) Diagnosis of Hutchinson-Gilford Progeria Syndrome. (2) Member has a body surface area of 0.39 m ² or above. Processing-deficient Progeroid Laminopathies (PDPL): (1) For treatment of processing-deficient Progeroid Laminopathies. (2) One of the following: (a) Heterozygous LMNA mutation with progerin-like protein accumulation (b) Homozygous or compound heterozygous ZMPSTE24 mutations. (3) Member has a body surface area of 0.39 m ² or above
Age Restrictions	(HGPS, PDPL): Member is 12 months of age or older
Prescriber Restrictions	
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZORYVE

Products Affected

- ZORYVE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PsO) (Initial): (1) Diagnosis of plaque psoriasis (2) Inadequate response or inability to tolerate TWO of the following topical therapies for sufficient duration (e.g., minimum of 4 weeks): (a) corticosteroids (e.g., betamethasone, clobetasol) (b) Vitamin D analogs (e.g., calcitriol, calcipotriene) (c) Concurrent combination of Vitamin D analog and corticosteroid (e.g., Enstilar, Taclonex) (d) Tazarotene (e) Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
Age Restrictions	
Prescriber Restrictions	(PsO) (Initial, Reauth): Prescribed by or in consultation with a dermatologist
Coverage Duration	(Initial): 6 months. (Reauthorization): 12 months
Other Criteria	(PsO) (Reauthorization): (1) Documentation of positive clinical response to therapy (e.g., reduction in the body surface area (BSA) involvement from baseline, improvement in symptoms such as pruritus or inflammation from baseline)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZORYVE FOAM

Products Affected

- ZORYVE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(SD)(Initial): (1) Diagnosis of seborrheic dermatitis (2) Inadequate response or inability to tolerate a minimum of a 4-week trial of TWO of the following generic topical therapies: (a) corticosteroids (e.g., betamethasone, clobetasol) (b) Antifungals (e.g., ciclopirox, ketoconazole) (c) calcineurin inhibitors (e.g., tacrolimus)
Age Restrictions	
Prescriber Restrictions	SD) (Initial, Reauth): Prescribed by or in consultation with a dermatologist
Coverage Duration	(Initial): 6 months (Reauth): 12 months
Other Criteria	(SD) (Reauth): Documentation of positive clinical response to therapy (e.g., reduction in the body surface area (BSA) involvement from baseline, improvement in symptoms such as pruritus or inflammation from baseline)
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZTALMY

Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(CDKL5): (1) Diagnosis of cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (2) Documentation of mutation in the CDKL5 gene (3) member is experiencing motor seizures (e.g., bilateral tonic, generalized tonic-clonic, bilateral clonic, atonic, focal, or bilateral tonic-clonic) (4) Inadequate response or inability to tolerate two formulary anticonvulsants (e.g., valproic acid, levetiracetam, lamotrigine)
Age Restrictions	(CDKL5): Member is 2 years of age or older
Prescriber Restrictions	(CDKL5): Prescribed by or in consultation with a neurologist
Coverage Duration	Indefinite
Other Criteria	Approve if for continuation of therapy
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZURZUVAE

Products Affected

- ZURZUVAE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	(PPD): (1) Diagnosis of Postpartum depression (PPD) (2) Inadequate response or inability to tolerate (a) one generic Selective serotonin reuptake inhibitor (SSRI) (b) One generic Serotonin-Norepinephrine reuptake inhibitor (SNRI) (3) Medication will not be used for greater than 14 days
Age Restrictions	(PPD): Member is 18 years of age or older
Prescriber Restrictions	
Coverage Duration	14 days
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

ZYMFENTRA SQ 2024

Products Affected

- ZYMFENTRA (2 PEN)
- ZYMFENTRA (2 SYRINGE)

PA Criteria	Criteria Details
Exclusion Criteria	(UC, CD): Concurrent therapy with any other biologic disease modifying antirheumatic drug (DMARD), e.g., tumor necrosis factor antagonists
Required Medical Information	Ulcerative Colitis (UC): (1) Diagnosis of moderately to severely active Ulcerative Colitis (2) Inadequate response or inability to tolerate two of the following: (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) upadacitinib (Rinvoq), (d) tofacitinib (Xeljanz/Xeljanz XR), (e) adalimumab-adbm (Cyltezo), (f) adalimumab-aaty (Yuflyma), (g) risankizumab (Skyrizi) OR documentation demonstrating that a trial may be inappropriate. Crohn's Disease (CD): (1) Diagnosis of moderately to severely active CD (2) Inadequate response or inability to tolerate TWO of the following (a) adalimumab (Humira), (b) ustekinumab (Stelara), (c) Risankizumab (Skyrizi), (d) adalimumab-adbm (Cyltezo), (e) adalimumab-aaty (Yuflyma), (f) upadacitinib (Rinvoq) OR documentation demonstrating that a trial may be inappropriate.
Age Restrictions	(UC, CD): Member is 18 years of age or older
Prescriber Restrictions	(UC, CD): Prescribed by or in consultation with a gastroenterologist
Coverage Duration	Indefinite
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Index

ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET THERAPY PACK 15 MG, 2 MG, 20 MG, 30 MG, 5 MG	3	AKEEGA	178
ABILIFY MYCITE STARTER KIT ORAL TABLET THERAPY PACK 10 MG	3	AKLIEF	252
<i>abiraterone acetate</i>	178	ALECENSA	178
ABRILADA (1 PEN)	11	ALLZITAL	104
ABRILADA (2 SYRINGE)	11	ALTRENO	252
ACTEMRA ACTPEN	5	ALUNBRIG	178
ACTEMRA SUBCUTANEOUS	5	ALVAIZ	23
ACTHAR	7	ALYQ	192
ACTHAR GEL	7	AMBIEN CR ORAL TABLET EXTENDED RELEASE 12.5 MG	108
ACTIVELLA ORAL TABLET 1-0.5 MG	106	AMBIEN ORAL TABLET 10 MG	108
<i>adalimumab-aacf (2 pen)</i>	11	<i>ambrisentan</i>	136
<i>adalimumab-aaty (1 pen) subcutaneous auto- injector kit 80 mg/0.8ml</i>	11	AMJEVITA SUBCUTANEOUS SOLUTION AUTO-INJECTOR	11
<i>adalimumab-aaty (2 pen)</i>	11	AMJEVITA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 40 MG/0.4ML, 40 MG/0.8ML	11
<i>adalimumab-aaty (2 syringe)</i>	11	AMJEVITA-PED 10KG TO <15KG	11
<i>adalimumab-adaz subcutaneous solution auto- injector</i>	11	AMJEVITA-PED 15KG TO <30KG	11
<i>adalimumab-adaz subcutaneous solution prefilled syringe 40 mg/0.4ml</i>	11	<i>amphetamine sulfate</i>	79
<i>adalimumab-adbm (2 pen)</i>	11	AMPYRA	25
<i>adalimumab-adbm (2 syringe)</i>	11	AMRIX	110
<i>adalimumab-adbm(cd/uc/hs strt)</i>	11	ANGELIQ	106
<i>adalimumab-adbm(ps/uv starter)</i>	11	APOKYN SUBCUTANEOUS SOLUTION CARTRIDGE	26
<i>adalimumab-fkjp (2 pen)</i>	11	<i>apomorphine hcl subcutaneous</i>	26
<i>adalimumab-fkjp (2 syringe)</i>	11	ARALAST NP INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG	212
<i>adalimumab-ryvk (2 pen)</i>	11	ARIKAYCE	27
<i>adalimumab-ryvk (2 syringe)</i>	11	<i>armodafinil</i>	28
<i>adapalene external cream</i>	252	ASCOMP-CODEINE	104
<i>adapalene external gel 0.3 %</i>	252	ATRALIN	252
<i>adapalene external pad</i>	252	AUGTYRO ORAL CAPSULE 40 MG	178
<i>adapalene-benzoyl peroxide external gel</i>	252	AURYXIA	191
ADBRY	16	AUSTEDO	29
ADCIRCA	192	AUSTEDO XR	29
ADEMPAS	17	AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG	29
AFINITOR	178	AUVELITY	31
AFINITOR DISPERZ	178	AVEED	249
AFREZZA INHALATION POWDER 12 UNIT, 4 UNIT, 60X4 & 60X8 & 60X12 UNIT, 8 UNIT, 90 X 4 UNIT & 90X8 UNIT, 90 X 8 UNIT & 90X12 UNIT	18	AYVAKIT	178
AGAMREE	19	BALVERSA	178
AIMOVIG	20	BELBUCA BUCCAL FILM 300 MCG, 450 MCG, 600 MCG, 750 MCG, 900 MCG	101
AJOVY	21	BENLYSTA SUBCUTANEOUS	32

BERINERT	9	<i>clindamycin-tretinoin</i>	252
BESREMI	33	COMBIPATCH	106
<i>bexarotene</i>	178, 251	COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG	178
BIJUVA	106	COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG	178
BIMZELX	34	COMETRIQ (60 MG DAILY DOSE)	178
BIVIGAM INTRAVENOUS SOLUTION 5 GM/50ML	122	COPIKTRA	178
<i>bosentan</i>	253	CORLANOR	49
BOSULIF	178	CORTROPHIN	7
BRAFTOVI ORAL CAPSULE 75 MG	178	COSENTYX (300 MG DOSE)	50
BRUKINSA	178	COSENTYX SENSOREADY (300 MG)	50
<i>butalbital-acetaminophen oral capsule</i>	104	COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 75 MG/0.5ML	50
<i>butalbital-acetaminophen oral tablet 50-300 mg, 50-325 mg</i>	104	COSENTYX UNOREADY	50
<i>butalbital-apap-caff-cod</i>	104	COTELLIC	178
<i>butalbital-apap-caffeine oral capsule</i>	104	CRESEMBA ORAL	51
<i>butalbital-apap-caffeine oral tablet 50-325- 40 mg</i>	104	CRINONE	191
<i>butalbital-asa-caff-codeine</i>	104	CUVRIOR	279
<i>butalbital-aspirin-caffeine oral capsule</i>	104	<i>cyclobenzaprine hcl er</i>	110
BYLVAY	35	<i>cyclobenzaprine hcl oral</i>	110
BYLVAY (PELLETS)	35	CYLTEZO (2 PEN)	14
CABOMETYX	178	CYLTEZO (2 SYRINGE)	14
CABTREO	252	CYLTEZO-CD/UC/HS STARTER	14
CALQUENCE ORAL TABLET	178	CYLTEZO-PSORIASIS/UV STARTER	14
CAMZYOS	37	CYSTADROPS	52
CAPLYTA	38	CYSTARAN	52
CAPRELSA	178	<i>dalfampridine er</i>	25
CARBAGLU ORAL TABLET SOLUBLE	39	DALVANCE	157
<i>carbinoxamine maleate oral solution</i>	104	<i>dapagliflozin pro-metformin er</i>	30
<i>carbinoxamine maleate oral tablet</i>	104	<i>dapagliflozin propanediol</i>	30
<i>carglumic acid oral tablet soluble</i>	39	<i>dasatinib</i>	178
<i>carisoprodol oral</i>	110	DAURISMO	178
CAYSTON	40	DAYBUE	53
CERDELGA	41	DAYVIGO	54
<i>chlordiazepoxide-clidinium</i>	104	<i>deferasirox granules</i>	55
<i>chlorzoxazone oral</i>	104	<i>deferasirox oral tablet</i>	55
CHOLBAM	42	<i>deferasirox oral tablet soluble</i>	55
CIALIS ORAL TABLET 5 MG	43	<i>deferiprone</i>	87
CIBINQO	44	<i>deflazacort</i>	66
CIMZIA (2 SYRINGE)	45	DEMEROL INJECTION SOLUTION 25 MG/ML, 50 MG/ML	104
CIMZIA SUBCUTANEOUS KIT 2 X 200 MG	45	DEPO-TESTOSTERONE INTRAMUSCULAR SOLUTION	249
CINRYZE	47	DIACOMIT	56
<i>clemastine fumarate oral syrup</i>	104	<i>diclofenac epolamine external</i>	58
<i>clemastine fumarate oral tablet 2.68 mg</i>	104	<i>diclofenac sodium external gel 3 %</i>	57
CLIMARA	106		
CLIMARA PRO	106		

DIFFERIN EXTERNAL CREAM.....	252	ESTROGEL.....	106
DIFFERIN EXTERNAL GEL 0.3 %.....	252	<i>eszopiclone oral tablet 3 mg</i>	108
DIFFERIN EXTERNAL LOTION.....	252	EUCRISA.....	78
DILAUDID ORAL TABLET 4 MG, 8 MG.....	101	EVAMIST.....	106
<i>dipyridamole oral</i>	104	EVEKEO.....	79
DIVIGEL.....	106	EVENITY.....	80
DOJOLVI.....	59	<i>everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5</i>	
DOPTELET ORAL TABLET 20 MG.....	60	<i>mg</i>	178
DOTTI.....	106	<i>everolimus oral tablet soluble</i>	178
DUPIXENT SUBCUTANEOUS SOLUTION		EVRYSDI.....	81
AUTO-INJECTOR.....	61	EXJADE.....	55
DUPIXENT SUBCUTANEOUS SOLUTION		EYSUVIS.....	83
PREFILLED SYRINGE 200 MG/1.14ML, 300		FABHALTA.....	84
MG/2ML.....	61	FASENRA.....	85
DUVYZAT.....	64	FASENRA PEN.....	85
EDLUAR SUBLINGUAL TABLET SUBLINGUAL		<i>fentanyl transdermal patch 72 hour 100</i>	
10 MG.....	108	<i>mcg/hr, 25 mcg/hr, 37.5 mcg/hr, 50 mcg/hr,</i>	
EGRIFTA SV.....	65	<i>62.5 mcg/hr, 75 mcg/hr, 87.5 mcg/hr</i>	101
ELESTRIN.....	106	FERRIPROX.....	87
EMFLAZA.....	66	FERRIPROX TWICE-A-DAY.....	87
EMGALITY.....	67	FEXMID.....	110
EMGALITY (300 MG DOSE).....	67	FILSPARI.....	88
EMSAM.....	69	FILSUVEZ.....	89
ENBREL MINI.....	70	FINTEPLA.....	90
ENBREL SUBCUTANEOUS SOLUTION 25		FIORICET ORAL CAPSULE.....	104
MG/0.5ML.....	70	FIORICET/CODEINE ORAL CAPSULE 50-300-	
ENBREL SUBCUTANEOUS SOLUTION		40-30 MG.....	104
PREFILLED SYRINGE.....	70	FIRAZYR SUBCUTANEOUS SOLUTION	
ENBREL SURECLICK SUBCUTANEOUS		PREFILLED SYRINGE.....	9
SOLUTION AUTO-INJECTOR.....	70	FIRDAPSE.....	91
ENDARI.....	71	FLECTOR EXTERNAL.....	58
ENSPRYNG.....	72	<i>fluticasone furoate-vilanterol inhalation</i>	
ENTADFI.....	73	<i>aerosol powder breath activated 100-25</i>	
ENTYVIO PEN.....	74	<i>mcg/act, 200-25 mcg/act</i>	30
EPCLUSA.....	195	<i>fluticasone-salmeterol inhalation aerosol</i>	30
EPIDIOLEX.....	75	FORTEO SUBCUTANEOUS SOLUTION PEN-	
EPIDUO.....	252	INJECTOR 600 MCG/2.4ML.....	247
EPIDUO FORTE.....	252	FOTIVDA.....	178
EPSOLAY.....	76	FRUZAQLA.....	178
ERIVEDGE.....	178	FYAVOLV.....	106
ERLEADA.....	178	<i>gabapentin (once-daily)</i>	95
<i>erlotinib hcl</i>	178	GALAFOLD.....	92
ESBRIET.....	77	GAMMAGARD INJECTION SOLUTION 2.5	
ESGIC ORAL TABLET.....	104	GM/25ML.....	122
ESTRACE ORAL.....	106	GAMMAGARD S/D LESS IGA.....	122
<i>estradiol transdermal</i>	106	GAMMAKED INJECTION SOLUTION 1	
<i>estradiol-norethindrone acet</i>	106	GM/10ML.....	122

GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML	122	<i>hydrocodone bitartrate er oral tablet er 24 hour abuse-deterrent</i>	4, 101
GAMUNEX-C INJECTION SOLUTION 1 GM/10ML	122	<i>hydromorphone hcl er oral tablet extended release 24 hour</i>	101
GATTEX	93	<i>hydromorphone hcl oral tablet 4 mg, 8 mg</i>	101
GAVRETO	178	HYFTOR	114
<i>gefitinib</i>	178	HYRIMOZ	11
GENOTROPIN MINIQUICK SUBCUTANEOUS PREFILLED SYRINGE	96	HYRIMOZ-CROHNS/UC STARTER	11
GENOTROPIN SUBCUTANEOUS CARTRIDGE	96	HYRIMOZ-PED<40KG CROHN STARTER	11
GILOTRIF	178	HYRIMOZ-PED>=40KG CROHN START	11
GLASSIA	212	HYRIMOZ-PLAQ PSOR/UEIT START	11
GLEEVEC	178	HYSINGLA ER	4, 101
GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG	178	IBRANCE	178
GLUMETZA	82	<i>ibuprofen-famotidine</i>	48
GOCOVRI	94	<i>icatibant acetate subcutaneous solution prefilled syringe</i>	9
GRALISE ORAL TABLET	95	ICLUSIG	178
GRASTEK	22	IDACIO (2 PEN)	11
HADLIMA	11	IDACIO (2 SYRINGE)	11
HADLIMA PUSHTOUCH	11	IDACIO-CROHNS/UC STARTER	11
HAEGARDA	98	IDACIO-PSORIASIS STARTER	11
HARVONI ORAL PACKET	195	IDHIFA	178
HARVONI ORAL TABLET 90-400 MG	195	ILUMYA	115
HETLIOZ	99	<i>imatinib mesylate</i>	178
HETLIOZ LQ	100	IMBRUVICA ORAL CAPSULE	178
HORIZANT ORAL TABLET EXTENDED RELEASE	103	IMBRUVICA ORAL SUSPENSION	178
HULIO (2 PEN)	11	IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG	178
HULIO (2 SYRINGE)	11	IMVEXXY MAINTENANCE PACK	191
HUMATROPE INJECTION CARTRIDGE	96	IMVEXXY STARTER PACK	191
HUMIRA (2 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT	111	INBRIJA	116
HUMIRA (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML	111	INCRELEX	117
HUMIRA-CD/UC/HS STARTER SUBCUTANEOUS AUTO-INJECTOR KIT 80 MG/0.8ML	111	INDOCIN ORAL	104
HUMIRA-PSORIASIS/UEIT STARTER SUBCUTANEOUS AUTO-INJECTOR KIT	111	INDOCIN RECTAL	104
HUMULIN R U-500 (CONCENTRATED)	113	<i>indomethacin oral suspension</i>	104
HUMULIN R U-500 KWIKPEN SUBCUTANEOUS SOLUTION PEN-INJECTOR ..	113	<i>indomethacin rectal suppository 50 mg</i>	104
<i>hydrocodone bitartrate er oral capsule extended release 12 hour</i>	4	INGREZZA	118
		INLYTA	178
		INQOVI	178
		INREBIC	178
		<i>insulin glargine max solostar</i>	121
		<i>insulin glargine solostar subcutaneous solution pen-injector 300 unit/ml</i>	121
		INTRAROSA	191
		IQIRVO	124
		IRESSA	178
		ISTURISA ORAL TABLET 1 MG, 5 MG	125

<i>ivabradine hcl</i>	49	<i>lisdexamfetamine dimesylate</i>	274
IWILFIN.....	178	LITFULO.....	138
JADENU.....	55	LIVMARLI ORAL SOLUTION 9.5 MG/ML.....	139
JADENU SPRINKLE.....	55	LIVTENCITY.....	141
JAKAFI.....	178	LODOCO.....	142
JATENZO.....	249	LONSURF.....	178
JAYPIRCA.....	178	LORBRENA.....	178
JINTELI.....	106	LUMAKRAS ORAL TABLET 120 MG, 320 MG... ..	178
JOENJA.....	126	LUMRYZ.....	143
JUBLIA.....	174	LUNESTA ORAL TABLET 3 MG.....	108
JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG.....	127	LYLLANA.....	106
JYNARQUE.....	129	LYNPARZA ORAL TABLET.....	178
KALYDECO.....	130	LYRICA CR.....	144
KERENDIA.....	131	LYTGOBI (12 MG DAILY DOSE).....	178
<i>ketorolac tromethamine oral</i>	107	LYTGOBI (16 MG DAILY DOSE).....	178
KEVEYIS.....	132	LYTGOBI (20 MG DAILY DOSE).....	178
KEVZARA.....	133	MAVYRET.....	195
KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE.....	134	MEKINIST.....	178
KISQALI (200 MG DOSE).....	178	MEKTOVI.....	178
KISQALI (400 MG DOSE).....	178	MENOSTAR.....	106
KISQALI (600 MG DOSE).....	178	<i>meperidine hcl injection solution 100 mg/ml, 25 mg/ml, 50 mg/ml</i>	104
KORLYM.....	135	<i>meperidine hcl oral solution</i>	104
KOSELUGO.....	178	<i>meperidine hcl oral tablet 50 mg</i>	104
KRAZATI.....	178	<i>meprobamate</i>	104
<i>lapatinib ditosylate</i>	178	<i>metaxalone</i>	104
LAZCLUZE.....	178	<i>metformin hcl er (mod)</i>	82
<i>ledipasvir-sofosbuvir</i>	195	<i>metformin hcl oral tablet 625 mg</i>	145
<i>lenalidomide</i>	178	<i>methadone hcl oral solution</i>	101
LENVIMA (10 MG DAILY DOSE).....	178	<i>methadone hcl oral tablet</i>	101
LENVIMA (12 MG DAILY DOSE).....	178	<i>methocarbamol oral tablet 1000 mg</i>	146
LENVIMA (14 MG DAILY DOSE).....	178	<i>mifepristone oral tablet 300 mg</i>	135
LENVIMA (18 MG DAILY DOSE).....	178	<i>miglustat</i>	292
LENVIMA (20 MG DAILY DOSE).....	178	MIMVEY.....	106
LENVIMA (24 MG DAILY DOSE).....	178	MINIVELLE.....	106
LENVIMA (4 MG DAILY DOSE).....	178	<i>mirabegron er</i>	30
LENVIMA (8 MG DAILY DOSE).....	178	<i>modafinil oral</i>	147
LETAIRIS.....	136	<i>morphine sulfate er beads oral capsule extended release 24 hour 120 mg</i>	101
<i>levorphanol tartrate oral tablet 3 mg</i>	101	<i>morphine sulfate er oral capsule extended release 24 hour 100 mg, 60 mg, 80 mg</i>	101
<i>l-glutamine oral packet</i>	71	<i>morphine sulfate er oral tablet extended release 100 mg, 200 mg, 60 mg</i>	101
LIBERVANT.....	10	MS CONTIN ORAL TABLET EXTENDED RELEASE 100 MG, 200 MG, 60 MG.....	101
LIBRAX.....	104	MULPLETA.....	148
LICART EXTERNAL.....	58	MYALEPT.....	149
<i>lidocaine external patch 5 %</i>	137		
LIDOCAN.....	137		
LIDODERM.....	137		

MYCAPSSA.....	150	OMNITROPE SUBCUTANEOUS SOLUTION	
MYFEMBREE.....	151	CARTRIDGE.....	96
MYTESI.....	153	OMNITROPE SUBCUTANEOUS SOLUTION	
<i>naproxen-esomeprazole mg</i>	48	RECONSTITUTED.....	96
NAYZILAM.....	10	OMVOH SUBCUTANEOUS.....	172
NERLYNX.....	178	ONGENTYS.....	173
NEXAVAR.....	178	ONUREG.....	178
NEXLETOL.....	154	OPSUMIT.....	181
NEXLIZET.....	154	OPSYNVI.....	175
NGENLA.....	156	OPZELURA.....	176
NINLARO.....	178	ORALAIR.....	22
NORDITROPIN FLEXPPO SUBCUTANEOUS		ORENCIA CLICKJECT.....	182
SOLUTION PEN-INJECTOR.....	96	ORENCIA SUBCUTANEOUS SOLUTION	
<i>norethindrone-eth estradiol</i>	106	PREFILLED SYRINGE.....	182
NORGESIC.....	109	ORENITRAM.....	181
<i>norgesic forte</i>	109	ORENITRAM MONTH 1.....	181
NOURIANZ.....	160	ORENITRAM MONTH 2.....	181
NOXAFIL ORAL PACKET.....	162	ORENITRAM MONTH 3.....	181
NOXAFIL ORAL SUSPENSION.....	161	ORGOVYX.....	178
NUBEQA.....	178	ORIAHNN.....	183
NUCALA.....	163	ORILISSA.....	184
NUCYNTA ER.....	4, 101	ORKAMBI.....	185
NUCYNTA ORAL TABLET 100 MG, 75 MG.....	101	ORLADEYO.....	186
NUDEXTA.....	165	ORMALVI.....	132
NUPLAZID ORAL CAPSULE.....	166	<i>orphenadrine citrate er</i>	104
NUPLAZID ORAL TABLET 10 MG.....	166	<i>orphenadrine-aspirin-caffeine oral tablet 25-</i>	
NURTEC.....	167	<i>385-30 mg</i>	109
NUTROPIN AQ NUSPIN 10 SUBCUTANEOUS		ORSERDU.....	178
SOLUTION PEN-INJECTOR.....	96	OSMOLEX ER ORAL TABLET EXTENDED	
NUTROPIN AQ NUSPIN 20 SUBCUTANEOUS		RELEASE 24 HOUR 129 MG.....	187
SOLUTION PEN-INJECTOR.....	96	OSPHENA.....	191
NUTROPIN AQ NUSPIN 5 SUBCUTANEOUS		OTEZLA.....	188
SOLUTION PEN-INJECTOR.....	96	OTREXUP SUBCUTANEOUS SOLUTION	
NUVIGIL.....	28	AUTO-INJECTOR 10 MG/0.4ML, 12.5	
NUZYRA.....	157, 177	MG/0.4ML, 15 MG/0.4ML, 17.5 MG/0.4ML, 20	
OALIVA.....	169	MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML.....	120
OCTAGAM INTRAVENOUS SOLUTION 1		OXERVATE.....	189
GM/20ML, 2 GM/20ML.....	122	<i>oxycodone hcl oral tablet 30 mg</i>	101
ODACTRA.....	22	OXYCONTIN ORAL TABLET ER 12 HOUR	
ODOMZO.....	178	ABUSE-DETERRENT.....	4, 101
OFEV.....	170	<i>oxymorphone hcl er oral tablet extended release</i>	
OGSIVEO.....	178	<i>12 hour 20 mg, 30 mg</i>	101
OJEMDA ORAL SUSPENSION RECONSTITUTED		<i>oxymorphone hcl er oral tablet extended release</i>	
.....	178	<i>12 hour 40 mg</i>	101
OJEMDA ORAL TABLET 100 MG.....	178	<i>oxymorphone hcl oral tablet 10 mg</i>	101
OJJAARA.....	178	PALYNZIQ.....	190
OLUMIANT.....	171	PANZYGA.....	122

<i>pazopanib hcl</i>	178	REPATHA SURECLICK.....	210
PEMAZYRE.....	178	RETEVMO ORAL TABLET.....	178
<i>pentazocine-naloxone hcl</i>	104	RETIN-A.....	252
PIQRAY (200 MG DAILY DOSE).....	178	RETIN-A MICRO.....	252
PIQRAY (250 MG DAILY DOSE).....	178	RETIN-A MICRO PUMP EXTERNAL GEL 0.06 %, 0.08 %.....	252
PIQRAY (300 MG DAILY DOSE).....	178	REVATIO ORAL TABLET.....	222
<i>pirfenidone</i>	77	REVLIMID.....	178
POMALYST.....	178	REZDIFFRA.....	213
<i>posaconazole oral</i>	161	REZLIDHIA.....	178
PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR.....	193	REZUROCK.....	214
<i>pregabalin er</i>	144	RINVOQ.....	215
PREMARIN ORAL.....	106	RINVOQ LQ.....	217
<i>pretomanid</i>	196	RIVFLOZA.....	218
PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML.....	122	ROXICODONE ORAL TABLET 30 MG.....	101
PROCYSBI ORAL PACKET.....	197	ROXYBOND ORAL TABLET ABUSE- DETERRENT 15 MG, 5 MG.....	4
PROLASTIN-C INTRAVENOUS SOLUTION.....	212	ROXYBOND ORAL TABLET ABUSE- DETERRENT 30 MG.....	101
PROLIA SUBCUTANEOUS SOLUTION REFILLED SYRINGE.....	198	ROZLYTREK.....	178
PROMACTA.....	200	RUBRACA.....	178
<i>promethazine hcl oral solution</i>	104	RUCONEST.....	9
<i>promethazine hcl oral tablet</i>	104	RYDAPT.....	178
<i>promethazine hcl rectal suppository 12.5 mg, 25 mg</i>	104	RYVENT.....	104
<i>promethazine vc</i>	104	SAJAZIR SUBCUTANEOUS SOLUTION REFILLED SYRINGE.....	9
PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG.....	104	SAMSCA.....	219
PROVIGIL.....	147	SCEMBLIX.....	178
PYRUKYND.....	202	SEROSTIM SUBCUTANEOUS SOLUTION RECONSTITUTED 4 MG, 5 MG, 6 MG.....	220
PYRUKYND TAPER PACK.....	202	SIGNIFOR.....	221
QBREXZA.....	203	<i>sildenafil citrate oral suspension reconstituted</i>	222
QINLOCK.....	178	<i>sildenafil citrate oral tablet 20 mg</i>	222
QUALAQUIN.....	204	SILIQ.....	223
<i>quinine sulfate oral</i>	204	SIMLANDI (2 PEN).....	11
QULIPTA.....	205	SIMPONI SUBCUTANEOUS SOLUTION AUTO- INJECTOR.....	224
QUVIVIQ.....	206	SIMPONI SUBCUTANEOUS SOLUTION REFILLED SYRINGE.....	224
RADICAVA ORS STARTER KIT.....	207	SIRTURO.....	226
RASUVO SUBCUTANEOUS SOLUTION AUTO- INJECTOR 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML.....	120	SIVEXTRO.....	157, 177
RAVICTI.....	208	SKYCLARYS.....	227
RECORLEV.....	209	SKYRIZI PEN.....	228
REPATHA.....	210	SKYRIZI SUBCUTANEOUS.....	228
REPATHA PUSHTRONEX SYSTEM.....	210	SKYTROFA.....	229
		<i>sodium oxybate</i>	290
		<i>sofosbuvir-velpatasvir</i>	195

SOGROYA.....	230	<i>testosterone transdermal gel 10 mg/act (2%),</i>	
SOHONOS.....	231	<i>12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%),</i>	
SOMA.....	110	<i>20.25 mg/act (1.62%), 25 mg/2.5gm (1%),</i>	
<i>sorafenib tosylate</i>	178	<i>40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%).....</i>	249
SOTYKTU.....	233	<i>testosterone transdermal solution</i>	249
SOVALDI.....	159	<i>tetrabenazine</i>	283
SPEVIGO SUBCUTANEOUS.....	234	THALOMID ORAL CAPSULE 100 MG, 50 MG...	178
SPRIX.....	107	TIBSOVO.....	178
SPRYCEL.....	178	TLANDO.....	249
STELARA SUBCUTANEOUS SOLUTION 45		TOBI PODHALER.....	119
MG/0.5ML.....	235	<i>tolvaptan</i>	219
STELARA SUBCUTANEOUS SOLUTION		TORPENZ.....	178
PREFILLED SYRINGE.....	235	TRACLEER.....	253
STIVARGA.....	178	TRELSTAR MIXJECT.....	158
<i>sunitinib malate</i>	178	TREMFYA SUBCUTANEOUS SOLUTION	
SUNOSI.....	236	AUTO-INJECTOR.....	254
SUTENT.....	178	TREMFYA SUBCUTANEOUS SOLUTION	
SYMDEKO.....	237	PREFILLED SYRINGE.....	254
SYMLINPEN 120 SUBCUTANEOUS SOLUTION		<i>tretinoin external</i>	252
PEN-INJECTOR.....	238	<i>tretinoin microsphere external gel 0.04 %, 0.1</i>	
SYMLINPEN 60 SUBCUTANEOUS SOLUTION		<i>%.....</i>	252
PEN-INJECTOR.....	238	<i>tretinoin microsphere pump external gel 0.08 %</i>	
TABRECTA.....	178	<i>.....</i>	252
<i>tadalafil (pah)</i>	192	TRIDACAINE II.....	137
<i>tadalafil oral tablet 2.5 mg, 5 mg</i>	43	<i>trientine hcl oral capsule 500 mg</i>	279
TADLIQ.....	239	TRIKAFTA.....	255
TAFINLAR.....	178	TRUQAP ORAL TABLET.....	178
TAGRISSO.....	178	TUKYSA.....	178
TAKHZYRO.....	241	TURALIO ORAL CAPSULE 125 MG.....	178
TALTZ.....	242	TWYNEO.....	252
TALZENNA.....	178	TYENNE SUBCUTANEOUS.....	5
TANLOR.....	146	TYKERB.....	178
TARGRETIN.....	178, 251	TYMLOS.....	256
TARPEYO.....	244	TYRVAYA.....	257
TASIGNA.....	178	TYVASO DPI MAINTENANCE KIT INHALATION	
<i>tasimelteon</i>	99	POWDER 16 MCG, 32 MCG, 48 MCG, 64 MCG	258
<i>tavaborole</i>	174	TYVASO DPI TITRATION KIT INHALATION	
TAVALISSE.....	245	POWDER 16 & 32 & 48 MCG.....	258
TAVNEOS.....	246	UBRELVY.....	259
TAZVERIK.....	178	UPTRAVI ORAL.....	260
TENCON ORAL TABLET 50-325 MG.....	104	UPTRAVI TITRATION.....	260
TEPMETKO.....	178	VABOMERE.....	157
<i>teriparatide subcutaneous solution pen-injector</i>		VALCHLOR.....	251
<i>620 mcg/2.48ml</i>	247	VALTOCO 10 MG DOSE.....	10
<i>testosterone cypionate intramuscular solution</i>		VALTOCO 15 MG DOSE.....	10
<i>100 mg/ml, 200 mg/ml</i>	249	VALTOCO 20 MG DOSE.....	10
<i>testosterone enanthate intramuscular solution</i>	249	VALTOCO 5 MG DOSE.....	10

VANFLYTA.....	178	XOSPATA.....	178
VECAMYL.....	261	XPOVIO (100 MG ONCE WEEKLY) ORAL	
VELSIPITY.....	262	TABLET THERAPY PACK 50 MG.....	178
VENCLEXTA.....	178	XPOVIO (40 MG ONCE WEEKLY) ORAL	
VENCLEXTA STARTING PACK.....	178	TABLET THERAPY PACK 40 MG.....	178
VEOZAH.....	263	XPOVIO (40 MG TWICE WEEKLY) ORAL	
VERKAZIA.....	264	TABLET THERAPY PACK 40 MG.....	178
VERZENIO.....	178	XPOVIO (60 MG ONCE WEEKLY) ORAL	
VFEND IV.....	191	TABLET THERAPY PACK 60 MG.....	178
VIJOICE.....	265	XPOVIO (60 MG TWICE WEEKLY).....	178
VIMOVO ORAL TABLET DELAYED RELEASE		XPOVIO (80 MG ONCE WEEKLY) ORAL	
500-20 MG.....	48	TABLET THERAPY PACK 40 MG.....	178
VITRAKVI.....	178	XPOVIO (80 MG TWICE WEEKLY).....	178
VIVELLE-DOT.....	106	XTAMPZA ER.....	4, 101
VIVJOA.....	266	XTANDI.....	178
VIZIMPRO.....	178	XYOSTED.....	249
VONJO.....	178	XYREM.....	290
VOQUEZNA.....	268	XYWAV.....	291
VOQUEZNA DUAL PAK.....	267	YARGESA.....	292
VOQUEZNA TRIPLE PAK.....	267	YONSA.....	178
VORANIGO.....	178	YUFLYMA (1 PEN).....	14
<i>voriconazole intravenous</i>	191	YUFLYMA (2 SYRINGE).....	14
VOSEVI.....	159	YUFLYMA-CD/UC/HS STARTER.....	14
VOTRIENT.....	178	YUSIMRY SUBCUTANEOUS SOLUTION AUTO-	
VOWST.....	269	INJECTOR.....	11
VOXZOGO.....	270	ZAVESCA.....	292
VOYDEYA.....	271	ZAVZPRET.....	293
VTAMA.....	272	ZEJULA ORAL TABLET.....	178
VUITY.....	273	ZELBORAF.....	178
VYNDAMAX.....	240	ZEMAIRA INTRAVENOUS SOLUTION	
VYNDAQEL.....	240	RECONSTITUTED 1000 MG.....	212
VYVANSE.....	274	ZEMDRI.....	157
WAINUA.....	275	ZEPATIER.....	159
WAKIX.....	276	ZEPOSIA.....	294
WEGOVY.....	277	ZEPOSIA 7-DAY STARTER PACK.....	294
WELIREG.....	178	ZEPOSIA STARTER KIT ORAL CAPSULE	
WINREVAIR.....	232	THERAPY PACK 0.23MG &0.46MG	
XALKORI.....	178	0.92MG(21).....	294
XDEMVI.....	280	ZERBAXA.....	157
XELJANZ.....	281	ZIANA.....	252
XELJANZ XR.....	281	ZILBRYSQ.....	295
XENAZINE.....	283	ZOKINVY.....	296
XERMELO.....	284	ZOLINZA.....	178
XGEVA.....	285	<i>zolpidem tartrate er oral tablet extended</i>	
XIFAXAN ORAL TABLET 550 MG.....	286	<i>release 12.5 mg</i>	108
XOLAIR.....	287	<i>zolpidem tartrate oral capsule</i>	108
XOLREMDI.....	289	<i>zolpidem tartrate oral tablet 10 mg</i>	108

<i>zolpidem tartrate sublingual tablet sublingual</i>	
3.5 mg.....	108
ZOMACTON.....	96
ZORYVE.....	297, 298
ZTALMY.....	299
ZTLIDO.....	191
ZURZUVAE.....	300
ZYDELIG.....	178
ZYKADIA ORAL TABLET.....	178
ZYMFENTRA (2 PEN).....	301
ZYMFENTRA (2 SYRINGE).....	301
ZYTIGA.....	178