

2022 Prior Authorization Criteria

Updated 12/01/2022

ABIRATERONE

Products Affected

- Abiraterone Acetate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Node-positive (N1), non-metastatic (M0) prostate cancer |

ACITRETIN

Products Affected

- Acitretin

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Psoriasis: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to methotrexate or cyclosporine. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Prevention of non-melanoma skin cancers in high risk individuals, Lichen planus, Keratosis follicularis (Darier Disease) |

ACTIMMUNE

Products Affected

- Actimmune

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Mycosis fungoides, Sezary syndrome. |

ADEMPAS

Products Affected

- Adempas

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. For chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4): 1) Patient has persistent or recurrent CTEPH after pulmonary endarterectomy (PEA), OR 2) Patient has inoperable CTEPH with the diagnosis confirmed by right heart catheterization AND by computed tomography (CT), magnetic resonance imaging (MRI), or pulmonary angiography. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

AIMOVIG

Products Affected

- Aimovig

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline, OR 2) The patient experienced an inadequate treatment response with a 4-week trial of any of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants, OR 3) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Initial 3 months, Reauthorization Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ALDURAZYME

Products Affected

- Aldurazyme

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For mucopolysaccharidosis I: Diagnosis of mucopolysaccharidosis I was confirmed by an enzyme assay demonstrating a deficiency of alpha-L-iduronidase enzyme activity or by genetic testing. Patients with Scheie syndrome must have moderate to severe symptoms. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ALECENSA

Products Affected

- Alecensa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent ALK-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC. |

ALOSETRON

Products Affected

- Alosetron HCl

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for a biological female or a person that self-identifies as a female with a diagnosis of severe diarrhea-predominant irritable bowel syndrome (IBS) AND 2) Chronic IBS symptoms lasting at least 6 months AND 3) Gastrointestinal tract abnormalities have been ruled out AND 4) Inadequate response to one conventional therapy (e.g., antispasmodics, antidepressants, antidiarrheals). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ALPHA1-PROTEINASE INHIBITOR

Products Affected

- Aralast NP Intravenous Solution Reconstituted
1000 MG, 500 MG
- Prolastin-C
- Zemaira

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For alpha1-proteinase inhibitor deficiency: Patient must have 1) clinically evident emphysema and 2) pretreatment serum alpha1-proteinase inhibitor level less than 11 micromol/L (80 mg/dL by radial immunodiffusion or 50 mg/dL by nephelometry). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ALUNBRIG

Products Affected

- Alunbrig

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent ALK-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC. |

AMBRISENTAN

Products Affected

- Ambrisentan

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

AMPHETAMINES

Products Affected

- Amphetamine-Dextroamphet ER
- Amphetamine-Dextroamphetamine

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of Attention-Deficit Hyperactivity Disorder (ADHD) or Attention Deficit Disorder (ADD) OR 2) The patient has a diagnosis of narcolepsy confirmed by a sleep study. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

APOKYN

Products Affected

- Apokyn Subcutaneous Solution Cartridge
- Apomorphine HCl Subcutaneous

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For continuation treatment of off episodes in Parkinson's disease: The patient is experiencing improvement on the requested drug. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ARCALYST

Products Affected

- Arcalyst

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (new starts): 1) two or more gout flares within the previous 12 months, AND 2) inadequate response, intolerance or contraindication to maximum tolerated doses of a non-steroidal anti-inflammatory drug (NSAID) and colchicine, AND 3) concurrent use with urate-lowering therapy. For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (continuation): 1) patient must have achieved or maintained a clinical benefit (i.e., a fewer number of gout attacks or fewer flare days) compared to baseline, AND 2) continued use of urate-lowering therapy concurrently with the requested drug. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | For prevention of gout flares: 4 months. Other: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Prevention of gout flares in patients initiating or continuing urate-lowering therapy. |

ARMODAFINIL

Products Affected

- Armodafinil

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of narcolepsy and the diagnosis is confirmed by sleep lab evaluation OR 2) The patient has a diagnosis of Shift Work Disorder (SWD) OR 3) The patient has a diagnosis of obstructive sleep apnea (OSA) and the diagnosis is confirmed by polysomnography. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

AUSTEDO

Products Affected

- Austedo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Tourette's syndrome |

AVASTIN

Products Affected

- Avastin

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. For all indications except ophthalmic-related disorders: The patient had an intolerable adverse event to both Mvasi AND Zirabev and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | <p>Breast cancer, central nervous system (CNS) tumor types: adult low-grade (WHO Grade II) infiltrative supratentorial astrocytoma/oligodendroglioma, adult intracranial and spinal ependymoma, anaplastic gliomas, adult medulloblastoma, primary central nervous system lymphoma, meningiomas, limited and extensive brain metastases, metastatic spine tumors, malignant pleural mesothelioma, ovarian cancer/fallopian tube cancer/primary peritoneal cancer types: carcinosarcoma (malignant mixed Mullerian tumors), clear cell carcinoma, mucinous carcinoma, grade 1 endometrioid carcinoma, low-grade serous carcinoma, ovarian borderline epithelial tumors (low malignant potential) with invasive implants, and malignant sex cord-stromal tumors, soft tissue sarcoma types: angiosarcoma and solitary fibrous tumor/hemangiopericytoma, uterine neoplasms, endometrial carcinoma, vulvar squamous cell carcinoma, and ophthalmic-related disorders: diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma and retinopathy of prematurity, small bowel adenocarcinoma.</p> |

AYVAKIT

Products Affected

- Ayvakit

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For myeloid and lymphoid neoplasms with eosinophilia, the patient meets all of the following criteria: 1) the disease is FIP1L1- PDGFRA rearrangement-positive, AND 2) The disease harbors a PDGFRA D842A mutation, AND 3) The disease is resistant to imatinib. For GIST, the patient meets either of the following criteria: 1) The disease harbors PDGFRA exon 18 mutation, including PDGFRA D842V mutations, OR 2) The requested drug will be used after failure on at least two Food and Drug Administration (FDA)-approved therapies in unresectable, recurrent, or metastatic disease without PDGFRA exon 18 mutation. For advanced systemic mastocytosis (AdvSM): 1) the patient has a diagnosis of advanced systemic mastocytosis including aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), and mast cell leukemia (MCL) AND 2) the patient has a platelet count of greater than or equal to 50,000/mcL. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Myeloid and lymphoid neoplasms with eosinophilia, gastrointestinal stromal tumor (GIST) for unresectable, recurrent, or metastatic disease without platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation |

B VS. D

Products Affected

- Abelcet
- Abraxane
- Acetylcysteine Inhalation
- Acyclovir Sodium Intravenous Solution
- Adriamycin Intravenous Solution
- Albuterol Sulfate Inhalation Nebulization Solution (2.5 MG/3ML) 0.083%, 0.63 MG/3ML, 1.25 MG/3ML, 2.5 MG/0.5ML
- Alimta
- AmBisome
- Amphotericin B Intravenous
- Amphotericin B Liposome
- Aprepitant Oral Capsule
- Arformoterol Tartrate
- azaCITIDine
- azaTHIOprine Oral Tablet 50 MG
- Bendeka
- Brovana
- Budesonide Inhalation Suspension 0.25 MG/2ML, 0.5 MG/2ML
- Calcitonin (Salmon) Nasal
- Calcitriol Intravenous Solution 1 MCG/ML
- Calcitriol Oral
- CARBOplatin Intravenous Solution
- Cinacalcet HCl
- CISplatin Intravenous Solution 100 MG/100ML, 200 MG/200ML, 50 MG/50ML
- Clinimix/Dextrose (4.25/10)
- Clinimix/Dextrose (4.25/5)
- Clinimix/Dextrose (5/15)
- Clinimix/Dextrose (5/20)
- Clinimix/Dextrose (6/5)
- Clinimix/Dextrose (8/10)
- Clinimix/Dextrose (8/14)
- Clinisol SF
- Clinolipid
- Cromolyn Sodium Inhalation
- Cyclophosphamide Injection
- Cyclophosphamide Intravenous
- Cyclophosphamide Oral Capsule
- Cyclophosphamide Oral Tablet
- CycloSPORINE Intravenous
- CycloSPORINE Modified
- CycloSPORINE Oral Capsule
- Cytarabine Injection Solution
- Dextrose Intravenous Solution 50 %, 70 %
- Diphtheria-Tetanus Toxoids DT
- DOCEtaxel CONCENTRATE 160 MG/8ML Intravenous
- DOCEtaxel CONCENTRATE 80 MG/4ML Intravenous
- DOCEtaxel Intravenous Concentrate 160 MG/8ML, 20 MG/ML, 80 MG/4ML
- DOCEtaxel Intravenous Solution 160 MG/16ML, 20 MG/2ML, 80 MG/8ML
- DOCEtaxel SOLUTION 160 MG/16ML Intravenous
- DOCEtaxel SOLUTION 20 MG/2ML Intravenous
- DOCEtaxel SOLUTION 80 MG/8ML Intravenous
- Doxercalciferol Oral
- DOXOrubicin HCl Intravenous Solution
- DOXOrubicin HCl Liposomal
- Dronabinol
- Engerix-B Injection Suspension 20 MCG/ML
- Engerix-B Injection Suspension Prefilled Syringe
- epiRUBicin HCl Intravenous Solution 200 MG/100ML, 50 MG/25ML
- Etoposide Intravenous Solution 100 MG/5ML, 500 MG/25ML
- Everolimus Oral Tablet 0.25 MG, 0.5 MG, 0.75 MG, 1 MG
- Fluorouracil Intravenous
- Formoterol Fumarate Inhalation
- FreAmine III Intravenous Solution 10 %
- Fulvestrant Intramuscular Solution Prefilled Syringe
- GamaSTAN
- Ganciclovir Sodium Intravenous Solution Reconstituted
- Gemcitabine HCl Intravenous Solution 1 GM/26.3ML, 2 GM/52.6ML, 200 MG/5.26ML
- Gemcitabine HCl Intravenous Solution Reconstituted

- Gengraf Oral Capsule 100 MG, 25 MG
- Gengraf Oral Solution
- Granisetron HCl Oral
- Heparin Sodium (Porcine) Injection Solution 1000 UNIT/ML, 10000 UNIT/ML, 20000 UNIT/ML, 5000 UNIT/ML
- Hepatamine
- HumuLIN R U-500 (CONCENTRATED)
- Ibandronate Sodium
- Imovax Rabies Intramuscular Suspension Reconstituted
- Intralipid
- Intron A
- Ipratropium Bromide Inhalation
- Ipratropium-Albuterol
- Irinotecan HCl
- Kadcyca
- Leucovorin Calcium Injection Solution 500 MG/50ML
- Leucovorin Calcium Injection Solution Reconstituted
- Levalbuterol HCl Inhalation
- levOCARNitine Oral Solution
- levOCARNitine Oral Tablet
- Lidocaine HCl (PF) Injection Solution 0.5 %, 1 %, 1.5 %
- Lidocaine HCl Injection Solution 0.5 %, 1 %, 2 %
- Methotrexate Sodium (PF) Injection Solution 1 GM/40ML, 250 MG/10ML, 50 MG/2ML
- Methotrexate Sodium Injection Solution 250 MG/10ML, 50 MG/2ML
- Methotrexate Sodium Injection Solution Reconstituted
- Morphine Sulfate (PF) Injection Solution 10 MG/ML, 2 MG/ML, 4 MG/ML, 5 MG/ML, 8 MG/ML
- Morphine Sulfate (PF) Intravenous Solution 10 MG/ML, 2 MG/ML
- Morphine Sulfate (PF) Intravenous Solution 4 MG/ML, 8 MG/ML
- Morphine Sulfate (PF) SOLUTION 4 MG/ML Intravenous
- Morphine Sulfate (PF) SOLUTION 8 MG/ML Intravenous
- Morphine Sulfate Intravenous Solution 1 MG/ML, 10 MG/ML, 4 MG/ML, 8 MG/ML
- Mycophenolate Mofetil Oral
- Mycophenolate Sodium
- Nulojix
- Nutrilipid
- Oxaliplatin
- PACLitaxel Intravenous Concentrate 100 MG/16.7ML, 150 MG/25ML, 30 MG/5ML, 300 MG/50ML
- PACLitaxel Protein-Bound Part
- Pamidronate Disodium Intravenous Solution 30 MG/10ML, 90 MG/10ML
- Pamidronate Disodium Intravenous Solution 6 MG/ML
- Pamidronate Disodium Intravenous Solution Reconstituted
- Paraplatin Intravenous Solution 1000 MG/100ML
- Paricalcitol Oral
- PEMEtrexed Disodium Intravenous Solution Reconstituted
- Pentamidine Isethionate Inhalation
- Plenamine
- PreHevbrio
- Premasol Intravenous Solution 10 %
- Procalamine
- Prograf Oral Packet
- Prosol
- RabAvert
- Recombivax HB
- SandIMMUNE Oral Solution
- Sirolimus Oral
- Tacrolimus Oral
- TDVAX
- Tenivac
- Toposar Intravenous Solution 1 GM/50ML, 100 MG/5ML
- TPN Electrolytes Intravenous Concentrate
- Travasol
- TrophAmine Intravenous Solution 10 %
- vinCRISTine Sulfate Intravenous
- Vinorelbine Tartrate
- Zoledronic Acid Intravenous Concentrate
- Zoledronic Acid Intravenous Solution
- Zortress Oral Tablet 1 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | N/A |
| Other Criteria | This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

BALVERSA

Products Affected

- Balversa

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BANZEL

Products Affected

- Rufinamide

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 1 year of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BENLYSTA

Products Affected

- Benlysta

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | For patients new to therapy: severe active central nervous system lupus. |
| Required Medical Information | For systemic lupus erythematosus (SLE): 1) Patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid or antimalarial) for SLE OR 2) patient is not currently receiving stable standard therapy regimen for SLE because patient tried and had an inadequate response or intolerance to stable standard therapy regimen. For lupus nephritis: 1) Patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid) for lupus nephritis OR 2) patient is not currently receiving a stable standard therapy regimen for lupus nephritis because patient tried and had an inadequate response or intolerance to a stable standard therapy regimen. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BERINERT

Products Affected

- Berinert

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For hereditary angioedema (HAE): The requested drug is being used for the treatment of acute angioedema attacks. Patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has HAE with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiotensin-1, plasminogen, or kininogen-1 (KNG1) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month. |
| Age Restrictions | |
| Prescriber Restrictions | Immunologist, allergist, rheumatologist |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BESREMI

Products Affected

- Besremi

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BETASERON

Products Affected

- Betaseron Subcutaneous Kit

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BEXAROTENE

Products Affected

- Bexarotene

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Mycosis fungoides, Sezary syndrome, CD30-positive primary cutaneous anaplastic large cell lymphoma, CD30-positive lymphomatoid papulosis. |

BOSENTAN

Products Affected

- Bosentan

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BOSULIF

Products Affected

- Bosulif

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL): Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant: patient has experienced resistance or intolerance to imatinib or dasatinib. If patient experienced resistance to an alternative tyrosine kinase inhibitor for CML, patient is negative for all of the following mutations: T315I, G250E, V299L, and F317L. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) |

BRAFTOVI

Products Affected

- Braftovi Oral Capsule 75 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For colorectal cancer: The patient must meet both of the following criteria: 1) Tumor is positive for BRAF V600E mutation, 2) The requested drug will be used for either of the following: a) as subsequent therapy for advanced or metastatic disease, or b) as primary treatment for unresectable metachronous metastases. For cutaneous melanoma: The patient must meet all of the following criteria: 1) Tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), 2) The requested drug will be used in combination with binimetinib, and 3) The requested drug will be used for either of the following: a) unresectable or metastatic disease, or b) adjuvant systemic therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Adjuvant systemic therapy for cutaneous melanoma |

BRIVIACT

Products Affected

- Briviact

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) If the patient is 4 years of age or older, the patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam. |
| Age Restrictions | 1 month of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BRIVIACT INJ

Products Affected

- Briviact

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) If the patient is 4 years of age or older, the patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam. |
| Age Restrictions | 1 month of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BRUKINSA

Products Affected

- Brukinsa

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

BUDESONIDE CAP

Products Affected

- Budesonide Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Patient has had a clinical relapse after cessation of treatment (induction) therapy for use in maintenance of microscopic colitis. |
| Age Restrictions | Crohn's, treatment: 8 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Microscopic colitis, maintenance: 12 months, all other indications: 3 months |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Treatment and maintenance of microscopic colitis in adults |

BUPRENORPHINE

Products Affected

- Buprenorphine HCl Sublingual

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug is being prescribed for the treatment of opioid use disorder AND patient meets one of the following: 1) The patient is pregnant or breastfeeding, and the requested drug is being prescribed for induction therapy and/or subsequent maintenance therapy for treatment of opioid use disorder OR 2) The requested drug is being prescribed for induction therapy for transition from opioid use to treatment of opioid use disorder OR 3) The requested drug is being prescribed for maintenance therapy for treatment of opioid use disorder in a patient who is intolerant to naloxone. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CABOMETYX

Products Affected

- Cabometyx

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For renal cell carcinoma: The disease is advanced, relapsed, or stage IV. For non-small cell lung cancer: 1) The disease is rearranged during transfection (RET) positive AND 2) the disease is recurrent, advanced, or metastatic. For hepatocellular carcinoma: the requested drug will be used as subsequent treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Non-small cell lung cancer |

CALCIPOTRIENE

Products Affected

- Calcipotriene External Ointment
- Calcipotriene External Solution
- Calcitrene
- Enstilar

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for the treatment of psoriasis AND 2) The patient experienced an inadequate treatment response, intolerance, or the patient has a contraindication to a topical steroid. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CALQUENCE

Products Affected

- Calquence

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For chronic lymphocytic leukemia or small lymphocytic lymphoma: the patient has experienced an intolerable adverse event with ibrutinib. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma |

CAPLYTA

Products Affected

- Caplyta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For treatment of schizophrenia: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, asenapine, olanzapine, quetiapine, risperidone, ziprasidone, AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: Latuda, Rexulti, Secuado, Vraylar. For treatment of depressive episodes associated with bipolar I: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: olanzapine, quetiapine, AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: Latuda, Vraylar. For treatment of depressive episodes associated with bipolar II: The patient experienced an inadequate treatment response, intolerance, or contraindication to generic quetiapine. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CAPRELSA

Products Affected

- Caprelsa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell. |

CARBAGLU

Products Affected

- Carbaglu Oral Tablet Soluble
- Carglumic Acid Oral Tablet Soluble

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For N-acetylglutamate synthase (NAGS) deficiency: Diagnosis of NAGS deficiency was confirmed by enzymatic or genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CAYSTON

Products Affected

- Cayston

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For treatment of respiratory symptoms in cystic fibrosis patients: 1) Pseudomonas aeruginosa is present in the patient's airway cultures OR 2) The patient has a history of pseudomonas aeruginosa infection or colonization in the airways. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CERDELGA

Products Affected

- Cerdelga

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For Gaucher disease, the diagnosis was confirmed by an enzyme assay demonstrating a deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing. The patient's CYP2D6 metabolizer status has been established using an FDA-cleared test. The patient is a CYP2D6 extensive metabolizer, an intermediate metabolizer, or a poor metabolizer. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CEREZYME

Products Affected

- Cerezyme Intravenous Solution Reconstituted
400 UNIT

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For Gaucher disease, the diagnosis was confirmed by an enzyme assay demonstrating a deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Type 2 Gaucher disease, Type 3 Gaucher disease |

CLOBAZAM

Products Affected

- CloBAZam

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 2 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CLOMIPRAMINE

Products Affected

- clomiPRAMINE HCl Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for one of the following: Obsessive-Compulsive Disorder (OCD) or Panic Disorder AND 2) The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to any of the following: a serotonin and norepinephrine reuptake inhibitor (SNRI) or a selective serotonin reuptake inhibitor (SSRI) OR 3) The requested drug is being prescribed for Depression AND 4) The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to two of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Depression, Panic Disorder |

CLORAZEPATE

Products Affected

- Clorazepate Dipotassium

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For all indications: the prescriber must acknowledge the benefit of therapy with the requested drug outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) the requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety OR 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs) OR b) serotonin-norepinephrine reuptake inhibitors (SNRIs). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Short-term relief anxiety-1 month, Anxiety Disorders-4 months, All other Diagnoses-Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 65 years of age or older. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CLOZAPINE ODT

Products Affected

- cloZAPine Oral Tablet Dispersible

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

COMETRIQ

Products Affected

- Cometriq (100 MG Daily Dose) Oral Kit 80 & 20 MG
- Cometriq (60 MG Daily Dose) MG
- Cometriq (140 MG Daily Dose) Oral Kit 3 x 20 MG & 80 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC: The requested medication is used for NSCLC when the patient's disease expresses rearranged during transfection (RET) gene rearrangements. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Non-small cell lung cancer (NSCLC), differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell. |

COPIKTRA

Products Affected

- Copiktra

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

COTELLIC

Products Affected

- Cotellic

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): The patient must meet both of the following criteria: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with vemurafenib. For unresectable or metastatic melanoma: The patient must meet both of the following criteria: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with vemurafenib (with or without atezolizumab). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma) |

CYSTADROPS

Products Affected

- Cystadrops

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cystinosis: 1) Diagnosis of cystinosis was confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing, and 2) Patient has corneal cystine crystal accumulation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CYSTAGON

Products Affected

- Cystagon

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For nephropathic cystinosis: Diagnosis of nephropathic cystinosis was confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

CYSTARAN

Products Affected

- Cystaran

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cystinosis: 1) Diagnosis of cystinosis was confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing, and 2) Patient has corneal cystine crystal accumulation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DALFAMPRIDINE

Products Affected

- Dalfampridine ER

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For multiple sclerosis, patient must meet the following: For new starts, prior to initiating therapy, patient meets the following: patient demonstrates sustained walking impairment. For continuation of therapy, patient meets the following: patient must have experienced an improvement in walking speed OR other objective measure of walking ability since starting the requested drug. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DAURISMO

Products Affected

- Daurismo

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For acute myeloid leukemia: 1) the requested medication must be used in combination with cytarabine, 2) the patient is 75 years of age or older OR has comorbidities that preclude intensive chemotherapy, and 3) the requested medication will be used as treatment for induction therapy, post-induction therapy, or relapsed or refractory disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Post induction therapy following response to previous therapy with the same regimen for acute myeloid leukemia (AML). Relapsed/refractory AML as a component of repeating the initial successful induction regimen. |

DEFERASIROX

Products Affected

- Deferasirox Granules
- Deferasirox Oral Tablet
- Deferasirox Oral Tablet Soluble

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For chronic iron overload due to blood transfusions: pretreatment serum ferritin level is greater than 1000 mcg/L. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DEMSER

Products Affected

- metyroSINE

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DESVENLAFAXINE

Products Affected

- Desvenlafaxine Succinate ER

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DEXMETHYLPHENIDATE

Products Affected

- Dexmethylphenidate HCl

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of Attention-Deficit Hyperactivity Disorder (ADHD) or Attention Deficit Disorder (ADD) OR 2) The requested drug is being prescribed for the treatment of cancer-related fatigue after other causes of fatigue have been ruled out. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Cancer-related fatigue |

DHE NASAL

Products Affected

- Dihydroergotamine Mesylate Nasal

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has experienced an inadequate treatment response to one triptan 5-HT1 receptor agonist OR 2) The patient has experienced an intolerance to one triptan 5-HT1 receptor agonist OR 3) The patient has a contraindication that would prohibit a trial of triptan 5-HT1 receptor agonists. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DIACOMIT

Products Affected

- Diacomit

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DIAZEPAM

Products Affected

- diazePAM Intensol
- diazePAM Oral Concentrate
- diazePAM Oral Solution 5 MG/5ML
- diazePAM Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For all indications: the prescriber must acknowledge the benefit of therapy with the requested drug outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) the requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Short-term relief anx-1 mo, skeletal muscle spasm-3 mo, Anx Disorders-4 mo, Other Diagnoses-PlanYR |
| Other Criteria | This Prior Authorization requirement only applies to patients 65 years of age or older. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DOPTELET

Products Affected

- Doptelet

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For thrombocytopenia associated with chronic liver disease: Baseline platelet count prior to a scheduled procedure is less than 50,000/mcL. For chronic immune thrombocytopenia (ITP): 1) For new starts: a) Patient has had an inadequate response or is intolerant to a prior therapy such as corticosteroids or immunoglobulins, AND b) Untransfused platelet count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000 to 50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes patient to trauma). 2) For continuation of therapy, platelet count response to the requested drug: a) Current platelet count is less than or equal to 200,000/mcL OR b) Current platelet count is greater than 200,000/mcL and less than or equal to 400,000/mcL and dosing will be adjusted to a platelet count sufficient to avoid clinically important bleeding. |
| Age Restrictions | 18 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Chronic liver disease: 1 month, ITP initial: 6 months, ITP reauthorization: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

DRIZALMA

Products Affected

- Drizalma Sprinkle

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has tried duloxetine capsules OR 2) The patient is unable to take duloxetine capsules for any reason (e.g., difficulty swallowing capsules, requires nasogastric administration). |
| Age Restrictions | Generalized Anxiety Disorder - 7 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Cancer pain, chemotherapy-induced neuropathic pain |

EMGALITY

Products Affected

- Emgality
- Emgality (300 MG Dose)

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for the preventive treatment of migraine in an adult patient AND 2) The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline, OR 3) The patient experienced an inadequate treatment response with a 4-week trial of any of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants, OR 4) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants OR 5) The requested drug is being prescribed for the treatment of episodic cluster headaches in an adult patient AND 6) The patient received the requested drug for at least 3 weeks of treatment and had a reduction in weekly cluster headache attack frequency from baseline OR 7) The patient experienced an inadequate treatment response, intolerance, or contraindication to a triptan medication (i.e., 5-HT ₁ receptor agonist). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Initial 3 months, Reauthorization Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

EMSAM

Products Affected

- Emsam

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) Patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion OR 2) Patient is unable to swallow oral formulations. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ENBREL

Products Affected

- Enbrel Mini
- Enbrel Subcutaneous Solution 25 MG/0.5ML
- Enbrel Subcutaneous Solution Prefilled Syringe
- Enbrel Subcutaneous Solution Reconstituted
- Enbrel SureClick Subcutaneous Solution Auto-Injector

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial, OR 2) Intolerance or contraindication to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) Patient meets any of the following: a) Patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR b) Pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, OR c) Patient has severe psoriasis that warrants a biologic DMARD as first-line therapy (i.e. at least 10% of the BSA or crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected). For hidradenitis suppurativa (new starts only): patient has severe, refractory disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Hidradenitis suppurativa |

ENDARI

Products Affected

- Endari

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 5 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

EPCLUSA

Products Affected

- Epclusa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current AASLD treatment guidelines. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Criteria will be applied consistent with current AASLD-IDSA guidance. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

EPIDIOLEX

Products Affected

- Epidiolex

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ERGOTAMINE

Products Affected

- Ergotamine-Caffeine

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ERIVEDGE

Products Affected

- Erivedge

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For adult medulloblastoma: patient has received chemotherapy previously AND has tumor(s) with mutations in the sonic hedgehog pathway |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Adult medulloblastoma |

ERLEADA

Products Affected

- Erleada

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ERLOTINIB

Products Affected

- Erlotinib HCl

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC (including brain metastases from NSCLC): 1) the disease is recurrent, advanced, or metastatic and 2) the patient has sensitizing EGFR mutation-positive disease. For pancreatic cancer: the disease is locally advanced, unresectable, or metastatic. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced non-small cell lung cancer (NSCLC), recurrent chordoma, relapsed or stage IV renal cell carcinoma (RCC), brain metastases from NSCLC. |

ESBRIET

Products Affected

- Esbriet
- Pirfenidone

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For idiopathic pulmonary fibrosis (Initial Review Only): 1) a high-resolution computed tomography (HRCT) study of the chest or a lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, OR 2) HRCT study of the chest reveals a result other than the UIP pattern (e.g., probable UIP, indeterminate for UIP) and the diagnosis is supported either by a lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if a lung biopsy has not been conducted. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

EVEROLIMUS

Products Affected

- Afinitor Disperz
- Afinitor Oral Tablet 10 MG
- Everolimus Oral Tablet 10 MG, 2.5 MG, 5 MG, 7.5 MG
- Everolimus Oral Tablet Soluble

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For breast cancer: 1) The disease is recurrent or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, AND 2) The requested medication is prescribed in combination with exemestane, fulvestrant, or tamoxifen, AND 3) The requested medication is used for subsequent treatment. For renal cell carcinoma: The disease is relapsed, advanced, or stage IV. For subependymal giant cell astrocytoma (SEGA): The requested drug is given as adjuvant treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Classic Hodgkin lymphoma, thymomas and thymic carcinomas, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, soft tissue sarcoma (perivascular epithelioid cell tumors (PEComa) and lymphangiomyomatosis subtypes), gastrointestinal stromal tumors, neuroendocrine tumors of the thymus, thyroid carcinoma (papillary, Hurthle cell, and follicular), endometrial carcinoma. |

EXKIVITY

Products Affected

- Exkivity

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FABRAZYME

Products Affected

- Fabrazyme

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Diagnosis of Fabry disease was confirmed by an enzyme assay demonstrating a deficiency of alpha-galactosidase enzyme activity or by genetic testing, or the patient is a symptomatic obligate female carrier. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FANAPT

Products Affected

- Fanapt
- Fanapt Titration Pack

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For treatment of schizophrenia: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, asenapine, olanzapine, quetiapine, risperidone, ziprasidone AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: Latuda, Rexulti, Secuado, Vraylar. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FASENRA

Products Affected

- Fasenra
- Fasenra Pen

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For severe asthma: For initial therapy: 1) Either a) Patient has baseline blood eosinophil count of at least 150 cells per microliter OR b) Patient is dependent on systemic corticosteroids, and 2) Patient has a history of severe asthma despite current treatment with both of the following medications at optimized doses: a) inhaled corticosteroid and b) additional controller (long-acting beta2-agonist, leukotriene modifier, or sustained-release theophylline). For continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose. |
| Age Restrictions | 12 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FEBUXOSTAT

Products Affected

- Febuxostat

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response to a maximally titrated dose of allopurinol OR the patient has experienced an intolerance to allopurinol OR the patient has a contraindication that would prohibit a trial of allopurinol. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FENTANYL PATCH

Products Affected

- FentaNYL Transdermal Patch 72 Hour 100 MCG/HR, 25 MCG/HR, 75 MCG/HR
- fentaNYL Transdermal Patch 72 Hour 12 MCG/HR, 50 MCG/HR

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR 2) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 3) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 4) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 5) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FETZIMA

Products Affected

- Fetzima
- Fetzima Titration

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FINTEPLA

Products Affected

- Fintepla

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FLUCYTOSINE

Products Affected

- Flucytosine Oral

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 weeks |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FORM ALT PA SUCRALFATE

Products Affected

- Carafate Oral Suspension
- Sucralfate Oral Suspension

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an intolerance to one other formulary product such as sucralfate tablets. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FORTEO

Products Affected

- Forteo Subcutaneous Solution Pen-Injector 600 MCG/2.4ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For postmenopausal osteoporosis: patient has ONE of the following (1 or 2): 1) a history of fragility fracture, OR 2) A pre-treatment T-score of less than or equal to -2.5 or pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment Fracture Risk Assessment Tool (FRAX) fracture probability AND patient has ANY of the following: a) indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) patient has failed prior treatment with or is intolerant to a previous injectable osteoporosis therapy OR c) patient has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate. For primary or hypogonadal osteoporosis in men: patient has one of the following: 1) a history of osteoporotic vertebral or hip fracture, OR 2) pre-treatment T-score of less than or equal to -2.5, OR 3) pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability. For glucocorticoid-induced osteoporosis: Patient has had an oral bisphosphonate trial of at least 1-year duration unless patient has a contraindication or intolerance to an oral bisphosphonate, AND patient meets ANY of the following: 1) patient has a history of fragility fracture, OR 2) a pre-treatment T-score of less than or equal to -2.5, OR 3) pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 months total unless the patient remains at high risk for fracture and benefit outweighs risk |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Other Criteria | Patient has high FRAX fracture probability if the 10 year probability is either greater than or equal to 20 percent for any major osteoporotic fracture or greater than or equal to 3 percent for hip fracture. If glucocorticoid treatment is greater than 7.5 mg (prednisone equivalent) per day, the estimated risk score generated with FRAX should be multiplied by 1.15 for major osteoporotic fracture and 1.2 for hip fracture. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FOTIVDA

Products Affected

- Fotivda

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For advanced renal cell carcinoma: The following criteria must be met: 1) The disease is relapsed or refractory, 2) The requested medication must be used after at least two prior systemic therapies, and 3) The patient has experienced disease progression or an intolerable adverse event with a trial of cabozantinib. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

FYCOMPA

Products Affected

- Fycompa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For treatment of partial-onset seizures: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam. For adjunctive treatment of primary generalized tonic-clonic seizures: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following: Vimpat, Spritam. |
| Age Restrictions | Partial-onset seizures: 4 years of age or older. Primary generalized tonic-clonic seizures: 12 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GATTEX

Products Affected

- Gattex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For short bowel syndrome (SBS) initial therapy: Adult patients were dependent on parenteral support for at least 12 months. For SBS continuation: Requirement for parenteral support has decreased from baseline while on therapy with the requested medication. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GAVRETO

Products Affected

- Gavreto

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced, or metastatic, and 2) The tumor is rearranged during transfection (RET) fusion-positive or RET rearrangement-positive. |
| Age Restrictions | Non-small cell lung cancer: 18 years of age or older. Medullary thyroid cancer and thyroid cancer: 12 years of age or older. |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced rearranged during transfection (RET) rearrangement-positive non-small cell lung cancer |

GILENYA

Products Affected

- Gilenya Oral Capsule 0.5 MG

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GILOTRIF

Products Affected

- Gilotrif

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For non-small cell lung cancer (NSCLC): Patient meets either of the following: 1) Patient has metastatic squamous NSCLC that progressed after platinum-based chemotherapy, OR 2) Patient has sensitizing EGFR mutation-positive disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GLATIRAMER

Products Affected

- Glatiramer Acetate
- Glatopa

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GRALISE

Products Affected

- Gralise Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response to gabapentin immediate-release or the patient has experienced an intolerance to gabapentin immediate-release. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

GROWTH HORMONE

Products Affected

- Genotropin MiniQuick Subcutaneous Prefilled Syringe
- Genotropin Subcutaneous Cartridge

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Pediatric patients with closed epiphyses (except in patients with PWS). |
| Required Medical Information | Pediatric growth hormone deficiency (GHD): Patient (pt) meets any of the following: 1) younger than 2.5 years old (yo) with pre-treatment (pre-tx) height (ht) more than 2 standard deviations (SD) below mean and slow growth velocity OR 2) 2.5 yo or older AND one of the following: a) pre-tx 1-year ht velocity more than 2 SD below mean OR b) pre-tx ht more than 2 SD below mean and 1-year ht velocity more than 1 SD below mean, AND patient meets any of the following: 1) failed 2 pre-tx growth hormone (GH) stimulation tests (peak below 10 ng/mL), OR 2) pituitary/central nervous system (CNS) disorder (e.g., genetic defects, CNS tumors, congenital structural abnormalities) and pre-tx insulin-like growth factor-1 (IGF-1) more than 2 SD below mean, OR 3) pt is a neonate or was diagnosed with GHD as a neonate. Turner syndrome: 1) Confirmed by karyotyping AND 2) pre-tx ht is less than the 5th percentile for age. Small for gestational age (GA): 1) Birth weight (wt) less than 2500g at GA greater than 37 weeks, OR birth wt or length below 3rd percentile for GA or at least 2 SD below mean for GA, AND 2) did not manifest catch-up growth by age 2. |
| Age Restrictions | SGA: 2 years of age or older |
| Prescriber Restrictions | Endocrinologist, pediatric endocrinologist, pediatric nephrologist, infectious disease specialist, gastroenterologist/nutritional support specialist, geneticist. |
| Coverage Duration | Plan Year |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Other Criteria | <p>Adult GHD: Pt meets any of the following: 1) failed 2 pre-tx GH stimulation tests, OR 2) pre-tx IGF-1 more than 2 SD below mean AND failed 1 pre-tx GH stimulation test. (Note: Stimulation tests include: a) insulin tolerance test [ITT] [peak GH less than or equal to 5 ng/ml], or b) Macrelin-stimulation test [peak GH level less than 2.8ng/ml], or c) glucagon-stimulation test [GST] [peak GH level less than or equal to 3 ng/ml] for pt with a body mass index [BMI] 25-30 kg/m2 and high pretest probability of GHD [e.g., acquired structural abnormalities] or BMI less than 25 kg/m2, or d) GST [peak GH level less than or equal to 1 ng/ml] in pt with BMI 25-30 kg/m2 and low pretest probability of GHD or BMI greater than 30 kg/m2), OR 3) organic hypothalamic-pituitary disease (e.g., suprasellar mass with previous surgery and cranial irradiation) with 3 or more pituitary hormone deficiencies AND pre-tx IGF-1 more than 2 SD below mean, OR 4) genetic or structural hypothalamic-pituitary defects, OR 5) childhood-onset GHD with congenital (genetic or structural) abnormality of the hypothalamus/pituitary/CNS. Renewal for pediatric GHD, TS, SGA, and adult GHD: Patient is experiencing improvement.</p> |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

HAEGARDA

Products Affected

- Haegarda

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug is being used for the prevention of acute angioedema attacks. Patient has hereditary angioedema (HAE) with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiopoietin-1, plasminogen, or kininogen-1 (KNG1) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month. |
| Age Restrictions | |
| Prescriber Restrictions | Immunologist, allergist, rheumatologist |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HARVONI

Products Affected

- Harvoni

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current AASLD treatment guidelines. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Criteria applied consistent w/ current AASLD-IDSa guidance. Reminder for 8wk option if appropriate. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HERCEPTIN

Products Affected

- Herceptin Intravenous Solution Reconstituted
150 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. The patient had an intolerable adverse event to Trazimera and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

HERCEPTIN HYLECTA

Products Affected

- Herceptin Hylecta

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months, Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer. |

HERZUMA

Products Affected

- Herzuma

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. The patient had an intolerable adverse event to Trazimera and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

HETLIOZ

Products Affected

- Hetlioz

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For Non-24-Hour Sleep-Wake Disorder: 1) for initial therapy and continuation of therapy: a) diagnosis of total blindness in both eyes (e.g., nonfunctioning retinas) and b) unable to perceive light in either eye, AND 2) if currently on therapy with the requested drug, patient must meet at least one of the following: a) increased total nighttime sleep or b) decreased daytime nap duration. For nighttime sleep disturbances in Smith-Magenis Syndrome (SMS): 1) for initial therapy and continuation therapy, the patient has a confirmed diagnosis of SMS AND 2) if currently on therapy with the requested drug, the patient experiences improvement in the quality of sleep since starting therapy. |
| Age Restrictions | Non-24: 18 years of age or older. SMS: 16 years of age or older |
| Prescriber Restrictions | Sleep disorder specialist or neurologist |
| Coverage Duration | Initiation: 6 Months, Renewal: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-ANTICONVULSANTS

Products Affected

- PHENobarbital Oral Elixir
- PHENobarbital Oral Tablet
- PHENobarbital Sodium Injection

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Epilepsy |

HRM-ANTIPARKINSON

Products Affected

- Benztropine Mesylate Oral
- Trihexyphenidyl HCl

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | EPS (extrapyramidal symptoms): 1) The patient has not tried the non-HRM alternative drug amantadine AND 2) The patient has a contraindication to the non-HRM alternative drug amantadine OR 3) The patient has tried the non-HRM alternative drug amantadine AND 4) The patient experienced an inadequate treatment response OR intolerance to the non-HRM alternative drug amantadine. Parkinson's: 1) The patient has tried two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole. AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-CYPROHEPTADINE

Products Affected

- Cyproheptadine HCl Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) The prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Pruritus, spasticity due to spinal cord injury |

HRM-DIPYRIDAMOLE

Products Affected

- Dipyridamole Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-GUANFACINE ER

Products Affected

- guanFACINE HCl ER

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-GUANFACINE IR

Products Affected

- guanFACINE HCl Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-HYDROXYZINE

Products Affected

- hydroXYzine HCl Oral Syrup
- hydroXYzine HCl Oral Tablet
- hydroXYzine Pamoate Oral Capsule 25 MG, 50 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety. If the patient is taking one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, cyclobenzaprine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.]. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-HYDROXYZINE INJ

Products Affected

- hydrOXYzine HCl Intramuscular

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Alcohol Withdrawal Syndrome: 1) The patient has not tried one of the following alternative drugs: clorazepate or lorazepam AND 2) The patient has a contraindication to one of the following alternative drugs: clorazepate or lorazepam OR 3) The patient has tried one of the following alternative drugs: clorazepate or lorazepam AND 4) The patient experienced an inadequate treatment response OR intolerance to one of the following alternative drugs: clorazepate or lorazepam. Anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-HYPNOTICS

Products Affected

- Zolpidem Tartrate Oral

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a contraindication to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) OR 2) The non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) has been tried AND 3) The patient experienced an inadequate treatment response OR intolerance to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) AND 4) If the patient is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, quetiapine, sertraline, clonazepam, escitalopram, alprazolam) with the requested drug, the prescriber has determined that taking multiple central nervous system (CNS) active medications is medically necessary for the patient [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls.]. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-PROMETHAZINE

Products Affected

- Promethazine HCl Injection
- Promethazine HCl Oral Syrup
- Promethazine HCl Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HRM-SCOPOLAMINE

Products Affected

- Scopolamine

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Excessive salivation |

HRM-SKELETAL MUSCLE RELAXANTS

Products Affected

- Cyclobenzaprine HCl Oral Tablet 10 MG, 5 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | If the patient is using one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, hydroxyzine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.]. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 3 months |
| Other Criteria | This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

HUMIRA

Products Affected

- Humira Pediatric Crohns Start Subcutaneous Prefilled Syringe Kit 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- Humira Pen Subcutaneous Pen-Injector Kit
- Humira Pen-CD/UC/HS Starter
- Humira Pen-Pediatric UC Start
- Humira Pen-Ps/UV/Adol HS Start Subcutaneous Pen-Injector Kit 40 MG/0.8ML
- Humira Pen-Psor/Uveit Starter
- Humira Subcutaneous Prefilled Syringe Kit 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis and axial spondyloarthritis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR 2) Intolerance or contraindication to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) Patient meets any of the following: a) Patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR b) Pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, OR c) Patient has severe psoriasis that warrants a biologic DMARD as first-line therapy (i.e. at least 10% of the BSA or crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy. For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |

| | |
|-----------------------|--|
| PA Criteria | Criteria Details |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Axial spondyloarthritis, Behcet's syndrome |

HYPNOTIC BENZODIAZEPINES

Products Affected

- Temazepam Oral Capsule 15 MG, 7.5 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Prescriber must acknowledge the benefit of therapy with the requested drug outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 65 years of age or older. APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

IBRANCE

Products Affected

- Ibrance

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Unresectable well-differentiated/dedifferentiated liposarcoma of the retroperitoneum. |

ICATIBANT

Products Affected

- Icatibant Acetate
- Sajazir

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For hereditary angioedema (HAE): The requested drug is being used for the treatment of acute angioedema attacks. Patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has HAE with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiotensin-1, plasminogen, or kininogen-1 (KNG1) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month. |
| Age Restrictions | 18 years of age or older |
| Prescriber Restrictions | Immunologist, allergist, rheumatologist |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ICLUSIG

Products Affected

- Iclusig

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL): diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, including patients who have received a hematopoietic stem cell transplant: 1) patient has accelerated or blast phase CML and no other kinase inhibitor is indicated, OR 2) patient has chronic phase CML and has experienced resistance or intolerance to at least 2 prior kinase inhibitors AND at least one of those was imatinib or dasatinib, OR 3) patient is positive for the T315I mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Therapy after hematopoietic stem cell transplant (HSCT) for chronic myeloid leukemia (CML) and acute lymphoblastic leukemia (ALL) patients |

IDHIFA

Products Affected

- IDHIFA

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation: 1) patient has a physiologic age of 60 years or older with newly-diagnosed AML and meets one of the following: a) patient is not a candidate for intensive induction therapy, or b) patient declines intensive induction chemotherapy, OR 2) patient has a physiologic age of 60 years or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug OR 3) patient has relapsed or refractory AML. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Newly-diagnosed acute myeloid leukemia |

IMATINIB

Products Affected

- Imatinib Mesylate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML) or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL): diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML: patient did not fail (excluding failure due to intolerance) prior therapy with a tyrosine kinase inhibitor. For melanoma: c-Kit mutation is positive. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Desmoid tumors, pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT), recurrent chordoma, melanoma, AIDS-related Kaposi sarcoma, chronic myelomonocytic leukemia, chronic graft versus host disease (cGVHD), T-cell acute lymphoblastic leukemia, aggressive systemic mastocytosis when eosinophilia is present with FIP1L1-PDGFR α fusion gene |

IMBRUVICA

Products Affected

- Imbruvica

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For mantle cell lymphoma: 1) the requested drug will be used in a patient who has received at least one prior therapy, OR 2) the requested drug will be used in combination with rituximab as pretreatment to induction therapy with RHyperCVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone) regimen. For marginal zone lymphoma (including gastric mucosa-associated lymphoid tissue [MALT] lymphoma, non-gastric MALT lymphoma, nodal marginal zone lymphoma, and splenic marginal zone lymphoma): the patient has received at least one prior therapy. For hairy cell leukemia: the requested drug will be used as a single agent for disease progression. For primary central nervous system lymphoma: 1) the disease is relapsed or refractory OR 2) the requested drug is used for induction therapy as a single agent. For histologic transformation of nodal marginal zone lymphoma to diffuse large B-cell lymphoma: the requested drug will be used in patients who have received prior chemoimmunotherapy. For diffuse large B-cell lymphoma: the requested drug will be used as second-line or subsequent therapy. For AIDS-related B-cell lymphoma: the requested drug will be used as a single agent and as second-line or subsequent therapy for relapsed disease. For post-transplant lymphoproliferative disorders: the requested drug will be used in patients who have received prior chemoimmunotherapy. For high-grade B-cell lymphoma: the requested drug will be used as second-line or subsequent therapy. For follicular lymphoma: the requested drug will be used as a single agent.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | Hairy cell leukemia, lymphoplasmacytic lymphoma, follicular lymphoma, primary central nervous system lymphoma, AIDS-related B-cell lymphoma, histologic transformation of nodal marginal zone lymphoma to diffuse large B-cell lymphoma, diffuse large B-cell lymphoma, post-transplant lymphoproliferative disorders, high-grade B-cell lymphoma. |

INCRELEX

Products Affected

- Increlex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Pediatric patients with closed epiphyses |
| Required Medical Information | For growth failure due to severe primary insulin-like growth factor-1 (IGF-1) deficiency or growth hormone gene deletion in patients who have developed neutralizing antibodies to growth hormone, must meet all of the following prior to beginning therapy with the requested drug (new starts only): 1) height 3 or more standard deviations (SD) below the mean for children of the same age and gender AND 2) basal IGF-1 level 3 or more SD below the mean for children of the same age and gender AND 3) provocative growth hormone test showing a normal or elevated growth hormone level. For renewal, patient is experiencing improvement. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

INGREZZA

Products Affected

- Ingrezza

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

INLYTA

Products Affected

- Inlyta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For renal cell carcinoma: the disease is advanced, relapsed, or stage IV. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Thyroid carcinoma (papillary, Hurthle cell, or follicular). |

INQOVI

Products Affected

- Inqovi

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

INREBIC

Products Affected

- Inrebic

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and janus kinase 2 (JAK2) rearrangement |

IR BEFORE ER

Products Affected

- Hysingla ER
- Methadone HCl IntensoI
- Methadone HCl Oral Solution
- Methadone HCl Oral Tablet
- Morphine Sulfate ER Oral Tablet Extended Release

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR 2) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 3) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 4) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 5) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

IRESSA

Products Affected

- Iressa

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC: 1) disease must be metastatic, advanced, or recurrent and 2) patient must have a sensitizing EGFR mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent or advanced non-small cell lung cancer (NSCLC). |

ISOTRETINOIN

Products Affected

- Accutane 40 MG
- Amnesteem
- Claravis
- Myorisan
- Zenatane
- ISOTretinoin Oral Capsule 10 MG, 20 MG, 30 MG,

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Refractory acne vulgaris, severe refractory rosacea, neuroblastoma, cutaneous T-cell lymphoma (CTCL) (e.g., mycosis fungoides, Sezary syndrome), high risk for developing skin cancer (squamous cell cancers), transient acantholytic dermatosis (Grover's Disease), keratosis follicularis (Darier Disease), lamellar ichthyosis, pityriasis rubra pilaris. |

ITRACONAZOLE

Products Affected

- Itraconazole Oral Capsule

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | If for the treatment of onychomycosis due to dermatophytes (Tinea unguium), the diagnosis has been confirmed by a fungal diagnostic test (e.g., potassium hydroxide [KOH] preparation, fungal culture, or nail biopsy). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Disseminated/CNS histoplasmosis, Histoplasmosis/Coccidioidomycosis ppx: 12 mths. Others: 6 mths |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Coccidioidomycosis, Coccidioidomycosis prophylaxis in HIV infection, Cryptococcosis, Histoplasmosis prophylaxis in HIV infection, invasive fungal infection prophylaxis in liver transplant patients, Microsporidiosis, Talaromycosis (formerly Penicilliosis), Pityriasis versicolor/Tinea versicolor, Sporotrichosis, Tinea corporis, Tinea cruris, Tinea capitis, Tinea manuum, Tinea pedis |

IVERMECTIN TAB

Products Affected

- Ivermectin Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug is not being prescribed for the prevention or treatment of coronavirus disease 2019 (COVID-19). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 1 month |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Ascariasis, Cutaneous larva migrans, Mansonelliasis, Scabies, Gnathostomiasis, Pediculosis |

IVIG

Products Affected

- Bivigam GM/100ML, 20 GM/200ML, 5 GM/50ML
- Flebogamma DIF Intravenous Solution 10 GM/100ML, 10 GM/200ML, 2.5 GM/50ML, 20 GM/200ML, 20 GM/400ML, 5 GM/100ML, 5 GM/50ML
- Gammagard
- Gammagard S/D Less IgA
- Gammaked Injection Solution 1 GM/10ML, 10 GM/100ML, 20 GM/200ML, 5 GM/50ML
- Gammaplex Intravenous Solution 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 20 GM/400ML, 5 GM/100ML, 5 GM/50ML
- Gamunex-C
- Octagam
- Panzyga
- Privigen

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For B-cell chronic lymphocytic leukemia (CLL): 1) serum IgG less than 500 mg/dL OR 2) a history of recurrent bacterial infections. For bone marrow transplant/hematopoietic stem cell transplant (BMT/HSCT): 1) IVIG is requested within the first 100 days post-transplant OR 2) serum IgG less than 400 mg/dL. For pediatric human immunodeficiency virus (HIV) infection: 1) serum IgG less than 400 mg/dL, OR 2) history of recurrent bacterial infections. For dermatomyositis and polymyositis: 1) at least one standard first-line treatment (corticosteroid or immunosuppressant) has been tried but was unsuccessful or not tolerated OR 2) patient is unable to receive standard therapy because of a contraindication or other clinical reason. For pure red cell aplasia (PRCA): PRCA is secondary to parvovirus B19 infection. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

JAKAFI

Products Affected

- Jakafi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For polycythemia vera: patient had an inadequate response or intolerance to interferon therapy or hydroxyurea. For acute lymphoblastic leukemia: patient has a cytokine receptor-like factor 2 (CRLF2) mutation or a mutation associated with activation of the Janus kinase/signal transducers and activators of transcription (JAK/STAT) pathway. For CMML-2: the requested drug is used in combination with a hypomethylating agent. For BCR-ABL negative aCML: the requested drug is used as a single agent or in combination with a hypomethylating agent. For essential thrombocythemia: patient had an inadequate response or loss of response to hydroxyurea, interferon therapy, or anagrelide. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Lower-risk myelofibrosis, accelerated phase myelofibrosis, blast phase myelofibrosis/acute myeloid leukemia, acute lymphoblastic leukemia (ALL), chronic myelomonocytic leukemia (CMML)-2, BCR-ABL negative atypical chronic myeloid leukemia (aCML), essential thrombocythemia, and myeloid, lymphoid or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement |

KALYDECO

Products Affected

- Kalydeco

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor. |
| Age Restrictions | 4 months of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

KANJINTI

Products Affected

- Kanjinti

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. The patient had an intolerable adverse event to Trazimera and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

KETOCONAZOLE

Products Affected

- Ketoconazole Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Acute or chronic liver disease. Concurrent use with drugs that are contraindicated with ketoconazole tablets: dofetilide, quinidine, pimozone, cisapride, methadone, disopyramide, dronedarone, ranolazine, ergot alkaloids, irinotecan, lurasidone, oral midazolam, alprazolam, triazolam, felodipine, nisoldipine, tolvaptan, eplerenone, lovastatin, simvastatin, or colchicine. |
| Required Medical Information | The potential benefits outweigh the risks of treatment with oral ketoconazole. For systemic fungal infections, the patient has any of the following diagnoses: blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis. For Cushing's syndrome: the requested drug is being prescribed for a patient who cannot tolerate surgery or where surgery has not been curative. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Cushing's syndrome |

KEYTRUDA

Products Affected

- Keytruda Intravenous Solution

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

KISQALI

Products Affected

- KISQALI (200 MG Dose)
- KISQALI (400 MG Dose)
- KISQALI (600 MG Dose)
- KISQALI Femara (400 MG Dose)
- KISQALI Femara (600 MG Dose)
- KISQALI Femara(200 MG Dose)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | For treatment of breast cancer using KISQALI (ribociclib) in combination with an aromatase inhibitor or KISQALI Femara Co-Pack (ribociclib and letrozole) as initial endocrine-based therapy, one of the following criteria must be met: 1) the patient is pre- or peri-menopausal OR 2) the patient is postmenopausal OR male AND the patient has experienced an intolerable adverse event to Ibrance (palbociclib) AND Verzenio (abemaciclib) or has a contraindication to Ibrance (palbociclib) AND Verzenio (abemaciclib). For treatment of breast cancer with KISQALI (ribociclib) in combination with fulvestrant, one of the following criteria must be met: 1) the requested drug is being used with fulvestrant as initial endocrine-based therapy in a postmenopausal patient or in a male, OR 2) the requested drug is being used following disease progression on endocrine therapy in a postmenopausal patient or in a male and the patient has experienced an intolerable adverse event to Ibrance (palbociclib) AND Verzenio (abemaciclib) OR has a contraindication to Ibrance (palbociclib) AND Verzenio (abemaciclib). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

KORLYM

Products Affected

- Korlym

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | Endocrinologist |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

KYNMOBI

Products Affected

- Kynmobi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For continuation treatment of off episodes in Parkinson's disease: The patient is experiencing improvement on the requested drug. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

LAPATINIB

Products Affected

- Lapatinib Ditosylate

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For HER2-positive breast cancer, the requested drug will be used in combination with any of the following: 1) aromatase inhibitor, 2) capecitabine, OR 3) trastuzumab. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Brain metastases from human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent epidermal growth factor receptor (EGFR)-positive chordoma, HER2-amplified and RAS and BRAF wild-type colorectal cancer in combination with trastuzumab. |

LENVIMA

Products Affected

- Lenvima (10 MG Daily Dose)
- Lenvima (12 MG Daily Dose)
- Lenvima (14 MG Daily Dose)
- Lenvima (18 MG Daily Dose)
- Lenvima (20 MG Daily Dose)
- Lenvima (24 MG Daily Dose)
- Lenvima (4 MG Daily Dose)
- Lenvima (8 MG Daily Dose)

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For differentiated thyroid cancer (follicular, papillary, or Hurthle cell): disease is not amenable to radioactive iodine therapy and unresectable, locally recurrent, persistent, or metastatic. For hepatocellular carcinoma: disease is unresectable or inoperable, local, metastatic or with extensive liver tumor burden. For renal cell carcinoma, the disease is advanced, relapsed, or stage IV. For endometrial carcinoma, the patient meets ALL of the following: 1) The disease is advanced, recurrent, or metastatic, 2) The patient experienced disease progression following prior systemic therapy, AND 3) The patient is not a candidate for curative surgery or radiation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Medullary thyroid carcinoma, recurrent endometrial carcinoma, thymic carcinoma |

LEUPROLIDE

Products Affected

- Leuprolide Acetate Injection

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Use in combination with growth hormone for children with growth failure and advancing puberty, recurrent androgen receptor positive salivary gland tumors. |

LIDOCAINE PATCHES

Products Affected

- Lidocaine External Patch 5 %

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Pain associated with diabetic neuropathy, pain associated with cancer-related neuropathy (including treatment-related neuropathy [e.g., neuropathy associated with radiation treatment or chemotherapy]). |

LONSURF

Products Affected

- Lonsurf

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For colorectal cancer: The disease is advanced or metastatic. For gastric or gastroesophageal junction adenocarcinoma, all of the following criteria must be met: 1) The disease is unresectable locally advanced, recurrent, or metastatic, and 2) The patient has been previously treated with at least two prior lines of chemotherapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

LORBRENA

Products Affected

- Lorbrena

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Anaplastic lymphoma kinase (ALK)-positive recurrent or advanced non-small cell lung cancer (NSCLC). Repressor of silencing (ROS)-1 rearrangement-positive recurrent, advanced, or metastatic NSCLC following progression on crizotinib, entrectinib, or ceritinib. |

LUMAKRAS

Products Affected

- Lumakras

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

LUMIZYME

Products Affected

- Lumizyme

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Diagnosis of Pompe disease was confirmed by an enzyme assay demonstrating a deficiency of acid alpha-glucosidase (GAA) enzyme activity or by genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

LUPRON PED

Products Affected

- Lupron Depot-Ped (1-Month)
- Lupron Depot-Ped (3-Month)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For central precocious puberty (CPP), patients not currently receiving therapy must meet all of the following criteria: 1) Diagnosis of CPP was confirmed by a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test OR a pubertal level of a third generation luteinizing hormone (LH) assay, 2) Assessment of bone age versus chronological age supports the diagnosis of CPP, and 3) The onset of secondary sexual characteristics occurred prior to 8 years of age for female patients OR prior to 9 years of age for male patients. |
| Age Restrictions | CPP: Patient must be less than 12 years old if female and less than 13 years old if male. |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

LUPRON-ENDOMETRIOSIS

Products Affected

- Lupron Depot (1-Month) Intramuscular Kit 3.75 MG
- Lupron Depot (3-Month) Intramuscular Kit 11.25 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For uterine fibroids, patient must meet one of the following: 1) Diagnosis of anemia (e.g., hematocrit less than or equal to 30 percent and/or hemoglobin less than or equal to 10g/dL), OR 2) the requested medication will be used prior to surgery for uterine fibroids. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Fibroids: 3 months (mo), max 6 mo total. Endometriosis: 6 mo, max 12 mo total. Others: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Breast cancer, malignant sex cord-stromal tumors, epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer |

LYNPARZA

Products Affected

- Lynparza Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For recurrent or metastatic breast cancer: the disease is BRCA 1/2-germline mutated. For prostate cancer: The patient has progressed on prior treatment with an androgen receptor-directed therapy. For epithelial ovarian, fallopian tube, or primary peritoneal cancer: 1) The requested drug is used for maintenance therapy for stage II-IV or recurrent disease who are in complete or partial response to chemotherapy OR 2) The patient has deleterious or suspected deleterious germline BRCA-mutated advanced, recurrent, or persistent disease after two or more prior chemotherapy regimens. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent HER2-negative, BRCA 1/2-germline mutated breast cancer, recurrent or metastatic HER2-positive, BRCA 1/2-germline mutated breast cancer |

LYRICA CR

Products Affected

- Lyrica CR

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response to gabapentin, or the patient has experienced an intolerance to gabapentin, or the patient has a contraindication to gabapentin. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

MAVYRET

Products Affected

- Mavyret

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh [CTP] class B or C). |
| Required Medical Information | For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [CTP class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases (AASLD) treatment guidelines. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Criteria will be applied consistent with current AASLD-IDSA guidance |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

MEGESTROL

Products Affected

- Megestrol Acetate Oral Suspension 625 MG/5ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Patient has experienced an inadequate treatment response or intolerance to megestrol 40 mg/mL oral suspension. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Cancer-related cachexia in adults |

MEKINIST

Products Affected

- Mekinist

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For brain metastasis from melanoma, adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with dabrafenib. For unresectable or metastatic melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used as a single agent or in combination with dabrafenib. For non-small cell lung cancer, anaplastic thyroid cancer, and solid tumors: 1) The tumor is positive for a BRAF V600E mutation, and 2) The requested drug will be used in combination with dabrafenib. For uveal melanoma, the requested drug will be used as a single agent. For low grade serous ovarian cancer: The requested drug will be used to treat persistent or recurrent disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Brain metastases from melanoma, uveal melanoma, central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma), low grade serous ovarian cancer. |

MEKTOVI

Products Affected

- Mektovi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cutaneous melanoma: The patient must meet all of the following criteria: 1) Tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), 2) The requested drug will be used in combination with encorafenib, and 3) The requested drug will be used for either of the following: a) unresectable or metastatic disease, or b) adjuvant systemic therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Adjuvant systemic therapy for cutaneous melanoma |

MEMANTINE

Products Affected

- Memantine HCl ER
- Memantine HCl Oral Solution 2 MG/ML
- Memantine HCl Oral Tablet 10 MG, 5 MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This edit only applies to patients less than 30 years of age. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

METHYLPHENIDATE

Products Affected

- Metadate ER Oral Tablet Extended Release 20 MG
- Methylphenidate HCl ER Oral Tablet Extended Release
- Methylphenidate HCl Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of Attention-Deficit Hyperactivity Disorder (ADHD) or Attention Deficit Disorder (ADD) OR 2) The patient has a diagnosis of narcolepsy confirmed by a sleep study OR 3) The requested drug is being prescribed for the treatment of cancer-related fatigue after other causes of fatigue have been ruled out. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |

MIGLUSTAT

Products Affected

- Miglustat

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For Gaucher disease: the diagnosis was confirmed by an enzyme assay demonstrating a deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

MODAFINIL

Products Affected

- Modafinil

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of narcolepsy and the diagnosis is confirmed by sleep lab evaluation OR 2) The patient has a diagnosis of Shift Work Disorder (SWD) OR 3) The patient has a diagnosis of obstructive sleep apnea (OSA) and the diagnosis is confirmed by polysomnography. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

MONJUVI

Products Affected

- Monjuvi

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

MVASI

Products Affected

- Mvasi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Off Label Uses | <p>Breast cancer, central nervous system (CNS) tumor types: adult low-grade (WHO Grade II) infiltrative supratentorial astrocytoma/oligodendroglioma, adult intracranial and spinal ependymoma, anaplastic gliomas, adult medulloblastoma, primary central nervous system lymphoma, meningiomas, limited and extensive brain metastases, metastatic spine tumors, malignant pleural mesothelioma, epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer, including the following cancer types: carcinosarcoma (malignant mixed Mullerian tumors), clear cell carcinoma, mucinous carcinoma, grade 1 endometrioid carcinoma, low-grade serous carcinoma, ovarian borderline epithelial tumors (low malignant potential) with invasive implants, and malignant sex cord-stromal tumors, soft tissue sarcoma types: angiosarcoma and solitary fibrous tumor/hemangiopericytoma, uterine neoplasms, endometrial carcinoma, vulvar squamous cell carcinoma, and ophthalmic-related disorders: diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma and retinopathy of prematurity, hepatocellular carcinoma, small bowel adenocarcinoma.</p> |

NAGLAZYME

Products Affected

- Naglazyme

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For mucopolysaccharidosis VI disease: Diagnosis was confirmed by an enzyme assay demonstrating a deficiency of N-acetylgalactosamine 4-sulfatase (arylsulfatase B) enzyme activity or by genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NATPARA

Products Affected

- Natpara

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Acute postsurgical hypoparathyroidism (within 6 months of surgery) and expected recovery from hypoparathyroidism. |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NERLYNX

Products Affected

- Nerlynx

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer, Brain metastases from HER2-positive breast cancer. |

NEXAVAR

Products Affected

- NexAVAR
- SORafenib Tosylate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For thyroid carcinoma: Histology is follicular, papillary, Hurthle cell or medullary. For acute myeloid leukemia, any of the following criteria must be met: 1) The requested drug is used in combination with azacitidine or decitabine for low-intensity treatment induction or post-induction therapy AND the patient is 60 years of age or older with FLT3-ITD mutation, OR 2) The disease is relapsed/refractory AND the requested drug is a component of repeating the initial successful induction if late relapse (greater than or equal to 12 months), OR 3) The disease is relapsed/refractory AND the requested drug is used in combination with azacitidine or decitabine if the patient is FLT3-ITD mutation positive. For renal cell carcinoma, the patient meets ALL of the following: 1) The disease is advanced, AND 2) The patient has experienced disease progression or an intolerable adverse event with a trial of cabozantinib or axitinib. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Acute myeloid leukemia, soft tissue sarcoma (angiosarcoma, desmoid tumors/aggressive fibromatosis, and solitary fibrous tumor subtypes), gastrointestinal stromal tumor, medullary thyroid carcinoma, osteosarcoma, recurrent chordoma, epithelial ovarian cancer, fallopian tube cancer, primary peritoneal cancer. |

NINLARO

Products Affected

- Ninlaro

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For multiple myeloma: The requested drug will be used in combination with lenalidomide and dexamethasone OR pomalidomide and dexamethasone OR dexamethasone OR cyclophosphamide and dexamethasone OR as a single agent. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Systemic light chain amyloidosis, Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma |

NITISINONE

Products Affected

- Nitisinone

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For hereditary tyrosinemia type 1: Diagnosis of hereditary tyrosinemia type 1 is confirmed by one of the following: 1) biochemical testing (e.g., detection of succinylacetone in urine) or 2) DNA testing (mutation analysis). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NORTHERA

Products Affected

- Droxidopa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For neurogenic orthostatic hypotension (nOH): Prior to initial therapy, patient has a persistent, consistent decrease in systolic blood pressure of at least 20 mmHg OR decrease in diastolic blood pressure of at least 10 mmHg within 3 minutes of standing or head-up tilt test. For continuation of therapy for nOH, patient experienced benefit from therapy (e.g., a sustained decrease in dizziness, lightheadedness, or feeling faint). For both initial and continuation of therapy for nOH, the requested drug will be used for patients with neurogenic orthostatic hypotension associated with one of the following diagnoses: 1) Primary autonomic failure due to Parkinson's disease, multiple system atrophy, or pure autonomic failure, OR 2) Dopamine beta-hydroxylase deficiency, OR 3) Non-diabetic autonomic neuropathy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 3 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NOXAFIL SUSP

Products Affected

- Noxafil Oral Suspension

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used orally. For treatment of oropharyngeal candidiasis: patient has experienced an inadequate treatment response, intolerance, or has a contraindication to fluconazole. |
| Age Restrictions | 13 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Oropharyngeal candidiasis: 1 month. All other indications: 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NUBEQA

Products Affected

- Nubeqa

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NUCALA

Products Affected

- Nucala

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | <p>For initial therapy for severe asthma: 1) Either a) Patient has baseline blood eosinophil count of at least 150 cells per microliter OR b) Patient is dependent on systemic corticosteroids, and 2) Patient has a history of severe asthma despite current treatment with both of the following medications at optimized doses: a) inhaled corticosteroid and b) additional controller (long-acting beta2-agonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For continuation therapy for severe asthma: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose. For initial therapy for eosinophilic granulomatosis with polyangiitis (EGPA): Patient has a history or the presence of an eosinophil count of more than 1000 cells per microliter or a blood eosinophil level of greater than 10 percent. For continuation of therapy for EGPA: Patient has a beneficial response to treatment with the requested drug, as demonstrated by any of the following: 1) a reduction in the frequency of relapses, 2) a reduction in the daily oral corticosteroid dose, or 3) no active vasculitis. For initial therapy for hypereosinophilic syndrome (HES): 1) Patient has had HES for greater than or equal to 6 months, 2) Patient has HES without an identifiable non-hematologic secondary cause, 3) Patient does not have FIP1L1-PDGFRα kinase-positive HES, 4) Patient has a history or presence of a blood eosinophil count of at least 1000 cells per microliter, AND 5) Patient has been on a stable dose of at least one HES therapy (e.g., oral corticosteroid, immunosuppressive, and/or cytotoxic therapy). For continuation of therapy for HES: Patient has a beneficial response to treatment as demonstrated by a reduction in HES flares.</p> |
| Age Restrictions | Asthma: 6 years of age or older, EGPA and chronic rhinosinusitis with nasal polyps: 18 years of age or older, HES: 12 years of age or older |
| Prescriber Restrictions | |

| PA Criteria | Criteria Details |
|--------------------------|-------------------------------|
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NUEDEXTA

Products Affected

- Nuedexta

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NUPLAZID

Products Affected

- Nuplazid Oral Capsule
- Nuplazid Oral Tablet 10 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For hallucinations and delusions associated with Parkinson's disease psychosis, the diagnosis of Parkinson's disease must be made prior to the onset of psychotic symptoms. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

NURTEC

Products Affected

- Nurtec

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | Acute migraine treatment: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to at least one triptan 5-HT1 receptor agonist. Preventive treatment of episodic migraine: 1) The patient received at least 3 months of preventive treatment with the requested drug and the patient had a reduction in migraine days per month from baseline OR 2) The patient meets either of the following: a) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants OR b) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Preventive treatment of migraine - initial: 3 months, All other indications: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OCREVUS

Products Affected

- Ocrevus

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OCTREOTIDE

Products Affected

- Octreotide Acetate Injection Solution 100 MCG/ML, 1000 MCG/ML, 200 MCG/ML, 50 MCG/ML, 500 MCG/ML
- Octreotide Acetate Subcutaneous

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly (continuation of therapy): patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control of thymomas and thymic carcinomas, the requested drug will be used as second-line systemic therapy in patients with unresectable or extrathoracic metastatic disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Tumor control of thymomas and thymic carcinomas. |

ODOMZO

Products Affected

- Odomzo

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OFEV

Products Affected

- Ofev

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OGIVRI

Products Affected

- Ogivri

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. The patient had an intolerable adverse event to Trazimera and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

OMNIPOD

Products Affected

- Omnipod 5 G6 Intro (Gen 5)
- Omnipod 5 G6 Pod (Gen 5)
- Omnipod Classic PDM (Gen 3)
- Omnipod Classic Pods (Gen 3)
- Omnipod DASH Intro (Gen 4)
- Omnipod DASH Pods (Gen 4)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has diabetes requiring insulin management with multiple daily injections AND 2) The patient is self-testing glucose levels 4 or more times per day OR the patient is using a continuous glucose monitor AND 3) The patient has experienced any of the following with the current diabetes regimen: inadequate glycemic control, recurrent hypoglycemia, wide fluctuations in blood glucose, dawn phenomenon with persistent severe early morning hyperglycemia, severe glycemic excursions. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | For continuation of therapy with an insulin pump, the patient has stable or improved glycemic control. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ONTRUZANT

Products Affected

- Ontruzant

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. The patient had an intolerable adverse event to Trazimera and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

ONUREG

Products Affected

- Onureg

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OPSUMIT

Products Affected

- Opsumit

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ORAL-INTRANASAL FENTANYL

Products Affected

- FentaNYL Citrate Buccal Lozenge On A Handle

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is indicated for the treatment of breakthrough CANCER-related pain only. The requested drug is being prescribed for the management of breakthrough pain in a CANCER patient with underlying CANCER pain AND 2) The International Classification of Diseases (ICD) diagnosis code provided supports the CANCER-RELATED diagnosis. [Note: For drug coverage approval, ICD diagnosis code provided MUST support the CANCER-RELATED diagnosis.] AND 3) The patient is currently receiving, and will continue to receive, around-the-clock opioid therapy for underlying CANCER pain AND 4) The requested drug is intended only for use in opioid tolerant patients. The patient can safely take the requested dose based on their current opioid use history. [Note: Patients considered opioid tolerant are those who are taking around-the-clock medicine consisting of at least 60 mg of oral morphine per day, at least 25 mcg per hour of transdermal fentanyl, at least 30 mg of oral oxycodone per day, at least 60 mg of oral hydrocodone per day, at least 8 mg of oral hydromorphone per day, at least 25 mg of oral oxymorphone per day, or an equianalgesic dose of another opioid medication daily for one week or longer.]. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ORGOVYX

Products Affected

- Orgovyx

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ORKAMBI

Products Affected

- Orkambi Oral Packet 100-125 MG, 150-188 MG
- Orkambi Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor. |
| Age Restrictions | 1 year of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OTEZLA

Products Affected

- Otezla

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For plaque psoriasis (new starts only): Patient meets either of the following: 1) Inadequate response or intolerance to ANY of the following: a) a topical therapy (e.g., a topical corticosteroid, calcineurin inhibitor, vitamin D analog), b) phototherapy (e.g., UVB, PUVA), or c) pharmacologic treatment with methotrexate, cyclosporine, or acitretin OR 2) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

OXANDROLONE

Products Affected

- Oxandrolone Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Turners Syndrome: Plan Year, All other diagnoses: 6 months |
| Other Criteria | Coverage will be denied if request is for an indication excluded from Medicare Part D. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Cachexia associated with AIDS (HIV wasting), To enhance growth in patients with Turners Syndrome |

PANRETIN

Products Affected

- Panretin

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Topical treatment of cutaneous lesions in patients with non-AIDS-related Kaposi sarcoma |

PEGASYS

Products Affected

- Pegasys Subcutaneous Solution 180 MCG/ML
- Pegasys Subcutaneous Solution Prefilled Syringe

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For chronic hepatitis C: Hepatitis C virus (HCV) confirmed by presence of hepatitis C virus HCV RNA in serum prior to starting treatment and the planned treatment regimen. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | HCV: 12-48 weeks depending on regimen. HBV: 48 weeks. All Other: Plan Year. |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Myeloproliferative neoplasm (essential thrombocythemia, polycythemia vera, symptomatic low risk myelofibrosis), systemic mastocytosis, adult T-cell leukemia/lymphoma, mycosis fungoides/Sezary syndrome, primary cutaneous CD30+ T-cell lymphoproliferative disorders. |

PEMAZYRE

Products Affected

- Pemazyre

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PHENYL BUTYRATE

Products Affected

- Sodium Phenylbutyrate Oral Powder 3 GM/TSP
- Sodium Phenylbutyrate Oral Tablet

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For urea cycle disorder: Diagnosis of urea cycle disorder (UCD) was confirmed by enzymatic, biochemical or genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PHESGO

Products Affected

- Phesgo

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer |

PHYTONADIONE TABLETS

Products Affected

- Phytonadione Tablet 5 MG Oral

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PIQRAY

Products Affected

- Piqray (200 MG Daily Dose)
- Piqray (250 MG Daily Dose)
- Piqray (300 MG Daily Dose)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated breast cancer in combination with fulvestrant. |

POMALYST

Products Affected

- Pomalyst

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For multiple myeloma: The patient has previously received at least two prior therapies for multiple myeloma, including an immunomodulatory agent AND a proteasome inhibitor. For Kaposi sarcoma, patient meets one of the following: 1) patient has acquired immunodeficiency syndrome (AIDS), or 2) patient is negative for human immunodeficiency virus (HIV). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Systemic light chain amyloidosis, primary central nervous system (CNS) lymphoma, POEMS syndrome. |

POSACONAZOLE

Products Affected

- Posaconazole

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used orally. |
| Age Restrictions | Treatment of Invasive Aspergillosis: 13 years of age or older, Prophylaxis of Invasive Aspergillus and Candida Infections: 2 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PRALUENT

Products Affected

- Praluent Subcutaneous Solution Auto-Injector

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PREVYMIS

Products Affected

- Prevyomis Oral

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For prophylaxis of cytomegalovirus (CMV) infection and disease: 1) The patient is CMV-seropositive, 2) the patient is a recipient of an allogeneic hematopoietic stem cell transplant (HSCT), AND 3) The requested medication will not be used beyond Day 100 post-transplantation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PROCRIT

Products Affected

- Procrit

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Patients receiving chemotherapy with curative intent. Patients with myeloid cancer. |
| Required Medical Information | Requirements regarding hemoglobin (Hgb) values exclude values due to a recent transfusion. For initial approval: 1) for all uses except anemia due to chemotherapy or myelodysplastic syndrome (MDS): patient has adequate iron stores AND 2) for all uses except surgery: pretreatment (no erythropoietin treatment in previous month) Hgb is less than 10 g/dL (less than 9 g/dL for anemia in congestive heart failure), AND 3) for MDS: pretreatment serum erythropoietin level is 500 international units/L or less. For reauthorizations (patient received erythropoietin treatment in previous month) in all uses except surgery: 1) patient has received at least 12 weeks of erythropoietin therapy, AND 2) patient responded to erythropoietin therapy, AND 3) current Hgb is less than 12 g/dL, AND 4) for all uses except anemia due to chemotherapy or MDS: patient has adequate iron stores. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 16 weeks |
| Other Criteria | Coverage includes use in anemia in patients whose religious beliefs forbid blood transfusions. Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual (e.g., used for treatment of anemia for a patient with chronic renal failure who is undergoing dialysis, or furnished from physician's supply incident to a physician service). |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | Anemia due to myelodysplastic syndromes (MDS), anemia in congestive heart failure (CHF), anemia in rheumatoid arthritis (RA), anemia due to hepatitis C treatment (ribavirin in combination with either interferon alfa or peginterferon alfa) |

PROMACTA

Products Affected

- Promacta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For chronic or persistent immune thrombocytopenia (ITP): 1) For new starts: a) Patient has had an inadequate response or is intolerant to a prior therapy such as corticosteroids or immunoglobulins, b) Untransfused platelet (plt) count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000-50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes patient to trauma) AND c) For chronic ITP only: patient has had an inadequate response or intolerance to avatrombopag. 2) For continuation of therapy, plt count response to the requested drug: a) Current plt count is less than or equal to 200,000/mcL OR b) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to a plt count sufficient to avoid clinically important bleeding. For thrombocytopenia associated with chronic hepatitis C: 1) For new starts: the requested drug is used for initiation and maintenance of interferon-based therapy. 2) For continuation of therapy: patient is receiving interferon-based therapy. For severe aplastic anemia (AA): For continuation of therapy following the initial 6 month approval for severe aplastic anemia: The patient must meet one of the following: 1) Current plt count is 50,000-200,000/mcL OR 2) Current plt count is less than 50,000/mcL and patient has not received appropriately titrated therapy for at least 16 weeks, OR 3) Current plt count is less than 50,000/mcL and patient is transfusion-independent, OR 4) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to achieve and maintain an appropriate target plt count.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |

| PA Criteria | Criteria Details |
|--------------------------|--|
| Coverage Duration | HCV: 6mo, ITP/AA initial: 6mo, ITP reauth: Plan Year, AA reauth: APR-Plan Year, IPR-16 wks |
| Other Criteria | APR: adequate platelet response (greater than 50,000/mcL), IPR: inadequate platelet response (less than 50,000/mcL). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

PULMOZYME

Products Affected

- Pulmozyme Inhalation Solution 2.5 MG/2.5ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For cystic fibrosis: Diagnosis of cystic fibrosis was confirmed by appropriate diagnostic or genetic testing. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

QINLOCK

Products Affected

- Qinlock

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

QUETIAPINE XR

Products Affected

- QUETiapine Fumarate ER

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For schizophrenia, acute treatment of manic or mixed episodes associated with bipolar I disorder, both as monotherapy and as an adjunct to lithium or divalproex, the acute treatment of depressive episodes associated with bipolar disorder, maintenance treatment of bipolar I disorder, as an adjunct to lithium or divalproex, adjunctive treatment of major depressive disorder, or maintenance monotherapy treatment in bipolar I disorder: The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: A) aripiprazole, B) asenapine, C) olanzapine, D) quetiapine immediate-release, E) risperidone, F) ziprasidone. For all indications: If the patient is 65 years of age or older AND is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, sertraline, clonazepam, escitalopram, alprazolam, zolpidem) with the requested drug, the prescriber determined that taking multiple central nervous system (CNS) active medications is medically necessary. [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls.]. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Maintenance monotherapy treatment in bipolar I disorder, monotherapy treatment of generalized anxiety disorder, monotherapy treatment of major depressive disorder |

QUININE SULFATE

Products Affected

- QuiNINE Sulfate Oral

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 1 month |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Babesiosis, uncomplicated Plasmodium vivax malaria |

REGRANEX

Products Affected

- Regranex

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For the treatment of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 20 weeks |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

RELISTOR INJ

Products Affected

- Relistor Subcutaneous Solution

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for opioid-induced constipation in an adult patient with advanced illness or pain caused by active cancer who requires opioid dosage escalation for palliative care OR 2) The requested drug is being prescribed for opioid-induced constipation in an adult patient with chronic non-cancer pain, including chronic pain related to prior cancer or its treatment who does not require frequent (e.g., weekly) opioid dosage escalation AND 3) The patient is unable to tolerate oral medications OR 4) An oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain (e.g., Movantik) has been tried AND 5) The patient experienced an inadequate treatment response or intolerance to an oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain (e.g., Movantik) OR 6) The patient has a contraindication that would prohibit a trial of an oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain (e.g., Movantik). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 4 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

REMICADE

Products Affected

- inFLIXimab
- Remicade

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | <p>For moderately to severely active Crohn's disease (new starts only): 1) Pt has fistulizing disease, OR 2) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 3) Intolerance or contraindication (CI) to conventional therapy. For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids) OR 2) Intolerance or CI to conventional therapy. For moderately to severely active rheumatoid arthritis (new starts only): 1) Pt meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) or leflunomide OR b) intolerance or CI to MTX AND leflunomide AND 2) pt meets ANY of the following: a) inadequate response, intolerance or CI to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis (new starts only): Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR intolerance or CI to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at time of diagnosis, AND 2) Pt meets ANY of the following: a) pt has experienced inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with MTX, cyclosporine, or acitretin, OR b) pharmacologic treatment with MTX, cyclosporine, or acitretin is contraindicated, OR c) pt has severe psoriasis that warrants a biologic DMARD as first-line therapy (i.e. at least 10% of BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Other Criteria | For hidradenitis suppurativa (new starts only): pt has severe, refractory disease. For uveitis (new starts only): Inadequate response or intolerance or has a CI to a trial of immunosuppressive therapy for uveitis. For FDA-approved indications and off-label uses that overlap: the patient had an intolerable adverse event to Renflexis and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Behcet's syndrome, granulomatosis with polyangiitis (Wegener's granulomatosis), hidradenitis suppurativa, juvenile idiopathic arthritis, pyoderma gangrenosum, sarcoidosis, Takayasu's arteritis, uveitis |

RENFLEXIS

Products Affected

- Renflexis

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | <p>For moderately to severely active Crohn's disease (new starts only): 1) Pt has fistulizing disease, OR 2) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 3) Intolerance or contraindication (CI) to conventional therapy. For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids) OR 2) Intolerance or CI to conventional therapy. For moderately to severely active rheumatoid arthritis (new starts only): 1) Pt meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) or leflunomide OR b) intolerance or CI to MTX AND leflunomide AND 2) pt meets ANY of the following: a) inadequate response, intolerance or CI to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis (new starts only): Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR intolerance or CI to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at time of diagnosis, AND 2) Pt meets ANY of the following: a) pt has experienced inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with MTX, cyclosporine, or acitretin, OR b) pharmacologic treatment with MTX, cyclosporine, or acitretin is contraindicated, OR c) pt has severe psoriasis that warrants a biologic DMARD as first-line therapy (i.e. at least 10% of BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Other Criteria | For hidradenitis suppurativa (new starts only): pt has severe, refractory disease. For uveitis (new starts only): Inadequate response or intolerance or has a CI to a trial of immunosuppressive therapy for uveitis. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Behcet's syndrome, granulomatosis with polyangiitis (Wegener's granulomatosis), hidradenitis suppurativa, juvenile idiopathic arthritis, pyoderma gangrenosum, sarcoidosis, Takayasu's arteritis, uveitis |

RETEVMO

Products Affected

- Retevmo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced or metastatic, and 2) Tumor is RET fusion-positive or RET rearrangement-positive. |
| Age Restrictions | Non-small cell lung cancer: 18 years of age or older. Medullary thyroid cancer and thyroid cancer: 12 years of age or older. |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced rearranged during transfection (RET)-rearrangement positive non-small cell lung cancer |

REVLIMID

Products Affected

- Lenalidomide
- Revlimid

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For myelodysplastic syndrome (MDS): Lower risk MDS with symptomatic anemia per the Revised International Prognostic Scoring System (IPSS-R), International Prognostic Scoring System (IPSS), or World Health organization (WHO) classification-based Prognostic Scoring System (WPSS). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | <p>Systemic light chain amyloidosis, classical Hodgkin lymphoma, myelodysplastic syndrome without the 5q deletion cytogenetic abnormality, myelofibrosis-associated anemia, POEMS syndrome, myeloproliferative neoplasms, non-Hodgkin's lymphoma with the following subtypes: acquired immunodeficiency syndrome (AIDS)-related non-germinal center diffuse large B-cell lymphoma, primary central nervous system (CNS) lymphoma, monomorphic post-transplant lymphoproliferative disorder, chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), diffuse large B-cell lymphoma, multicentric Castleman's disease, adult T-cell leukemia/lymphoma, mycosis fungoides (MF)/Sezary syndrome (SS), angioimmunoblastic T-cell lymphoma (AITL), peripheral T-cell lymphoma not otherwise specified (PTCL NOS), enteropathy-associated T-cell lymphoma, monomorphic epitheliotropic intestinal T-cell lymphoma, nodal peripheral T-cell lymphoma, primary cutaneous anaplastic large cell lymphoma (ALCL), hepatosplenic T-cell lymphoma, high-grade B-cell lymphomas, histologic transformation of nodal marginal zone lymphoma to diffuse large B-cell lymphoma, histologic transformation of follicular lymphoma to diffuse large B-cell lymphoma, AIDS-related Kaposi sarcoma, smoldering myeloma</p> |

REZUROCK

Products Affected

- Rezurock

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 12 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

RIABNI

Products Affected

- Riabni

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis and 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year |
| Other Criteria | The patient had an intolerable adverse event to both Truxima AND Ruxience and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Off Label Uses | <p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, gastric mucosa-associated lymphoid tissue [MALT], nongastric MALT), Burkitt lymphoma, primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma with translocations of MYC and BCL2 and/or BCL6 (double/triple hit lymphoma), high-grade B-cell lymphoma not otherwise specified, histological transformation from follicular lymphoma to diffuse large B-cell lymphoma, histological transformation from nodal marginal zone lymphoma to diffuse large B-cell lymphoma, Castleman's disease, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis, immune checkpoint inhibitor-related toxicities, moderately to severely active rheumatoid arthritis, pemphigus vulgaris, pediatric Burkitt-like lymphoma (BLL) and pediatric mature B-cell acute leukemia (B-AL).</p> |

RINVOQ

Products Affected

- Rinvoq

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): patient has experienced an inadequate treatment response or intolerance to at least one tumor necrosis factor (TNF) inhibitor. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

RITUXAN

Products Affected

- Rituxan Intravenous Solution

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis and 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year |
| Other Criteria | The patient had an intolerable adverse event to both Truxima AND Ruxience and that adverse event was NOT attributed to the active ingredient as described in the prescribing information. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | <p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, gastric mucosa-associated lymphoid tissue [MALT], nongastric MALT), primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma with translocations of MYC and BCL2 and/or BCL6 (double/triple hit lymphoma), high-grade B-cell lymphoma not otherwise specified, histological transformation from follicular lymphoma to diffuse large B-cell lymphoma, histological transformation from nodal marginal zone lymphoma to diffuse large B-cell lymphoma, Castleman's disease, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis, and immune checkpoint inhibitor-related toxicities.</p> |

RITUXAN HYCELA

Products Affected

- Rituxan Hycela

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Malignancies must be CD20 positive. Patient must receive at least one full dose of a rituximab product by intravenous infusion without experiencing severe adverse reactions. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Castleman's disease (CD), high-grade B-cell lymphoma, histologic transformation of nodal marginal zone lymphoma to diffuse large B-cell lymphoma, marginal zone lymphomas (nodal marginal zone lymphoma, gastric mucosa-associated lymphoid tissue (MALT) lymphoma, nongastric MALT lymphoma, and splenic marginal zone lymphoma), mantle cell lymphoma, post-transplant lymphoproliferative disorder (PTLD), primary cutaneous B-cell lymphoma (e.g., cutaneous marginal zone lymphoma or cutaneous follicle center lymphomas), hairy cell leukemia, small lymphocytic lymphoma (SLL). |

ROZLYTREK

Products Affected

- Rozlytrek

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced ROS1-positive non-small cell lung cancer (NSCLC), advanced, recurrent, or persistent neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors. |

RUBRACA

Products Affected

- Rubraca

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For metastatic castration-resistant prostate cancer with a deleterious breast cancer susceptibility gene (BRCA) mutation (germline and/or somatic): 1) patient has been treated with androgen receptor-directed therapy, 2) patient has been treated with a taxane-based chemotherapy or the patient is not fit for chemotherapy, 3) the requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy, and 4) patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib). For maintenance treatment of patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy, patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib). For treatment of patients with a deleterious breast cancer susceptibility gene (BRCA) mutation (germline and/or somatic)-associated epithelial ovarian, fallopian tube, or primary peritoneal cancer who have been treated with two or more chemotherapies: if prescribed for deleterious germline BRCA-mutated advanced ovarian cancer treated with two or more prior chemotherapies, the patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

RUXIENCE

Products Affected

- Ruxience

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis and 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | <p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, gastric mucosa-associated lymphoid tissue [MALT], nongastric MALT), Burkitt lymphoma, primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma with translocations of MYC and BCL2 and/or BCL6 (double/triple hit lymphoma), high-grade B-cell lymphoma not otherwise specified, histological transformation from follicular lymphoma to diffuse large B-cell lymphoma, histological transformation from nodal marginal zone lymphoma to diffuse large B-cell lymphoma, Castleman's disease, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis, immune checkpoint inhibitor-related toxicities, pemphigus vulgaris, pediatric Burkitt-like lymphoma (BLL), and pediatric mature B-cell acute leukemia (B-AL).</p> |

RYDAPT

Products Affected

- Rydapt

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For acute myeloid leukemia (AML): AML must be FLT3 mutation-positive. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 or FLT3 rearrangements: the disease is in chronic or blast phase. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Relapsed or refractory acute myeloid leukemia (AML), myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 or FLT3 rearrangements, post-remission maintenance therapy for acute myeloid leukemia (AML), re-induction in residual disease for acute myeloid leukemia (AML) |

SAPROPTERIN

Products Affected

- Javygtor Oral Packet 100 MG
- Javygtor Oral Tablet
- Sapropterin Dihydrochloride Oral Packet
- Sapropterin Dihydrochloride Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For phenylketonuria: For patients who have not yet received a therapeutic trial of the requested drug, the patient's pretreatment, including before dietary management, phenylalanine level is greater than 6 mg/dL (360 micromol/L). For patients who completed a therapeutic trial of the requested drug, the patient must have experienced improvement (for example, reduction in blood phenylalanine levels, improvement in neuropsychiatric symptoms). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Initial: 2 months. All others: Plan Year. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SAVELLA

Products Affected

- Savella
- Savella Titration Pack

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to duloxetine or pregabalin. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SCSEMBLIX

Products Affected

- Scemblix

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML) in the chronic phase: 1) the diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene AND the patient meets either of the following: A) the patient has previously been treated with 2 or more tyrosine kinase inhibitors (TKIs) AND at least one of those was imatinib or dasatinib, OR B) the patient is positive for the T315I mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SIGNIFOR

Products Affected

- Signifor

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | Endocrinologist |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SILDENAFIL

Products Affected

- Sildenafil Citrate Oral Tablet 20 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SIRTURO

Products Affected

- Sirturo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | Prescribed by or in consultation with an infectious disease specialist. |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SKYRIZI

Products Affected

- Skyrizi
- Skyrizi (150 MG Dose)
- Skyrizi Pen

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) patient meets any of the following: a) patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, or b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, or c) patient has severe psoriasis that warrants a biologic disease-modifying antirheumatic drug (DMARD) as first-line therapy (i.e. at least 10% of the body surface area or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SKYRIZI-CD

Products Affected

- Skyrizi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) patient meets any of the following: a) patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, or b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, or c) patient has severe psoriasis that warrants a biologic disease-modifying antirheumatic drug (DMARD) as first-line therapy (i.e. at least 10% of the body surface area or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SOMATULINE DEPOT

Products Affected

- Somatuline Depot Subcutaneous Solution 120 MG/0.5ML, 60 MG/0.2ML, 90 MG/0.3ML

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly continuation of therapy: patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control, the requested drug will be used for any of the following: 1) neuroendocrine tumor of the thymus or lung in patients with locoregional unresectable disease and/or distant metastatic disease, OR 2) unresected primary gastrinoma, OR 3) pheochromocytomas and paragangliomas, used for either of the following: a) symptomatic locally unresectable disease with somatostatin receptor positive imaging OR b) secreting tumor in metastatic disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Tumor control of neuroendocrine tumors (NETs) of the lung, thymus (carcinoid tumors) or unresected primary gastrinoma, and pheochromocytoma/paraganglioma. |

SOMAVERT

Products Affected

- Somavert

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly continuation of therapy: patient's IGF-1 level has decreased or normalized since initiation of therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SPRYCEL

Products Affected

- Sprycel

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML), including patients who have received a hematopoietic stem cell transplant: diagnosis was confirmed by detection of the Philadelphia (Ph) chromosome or BCR-ABL gene. If patient experienced resistance to an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I/A, F317L/V/I/C, and V299L mutations. For acute lymphoblastic leukemia (ALL), the patient has a diagnosis of one of the following: 1) Philadelphia chromosome positive ALL that has been confirmed by detection of the Ph chromosome or BCR-ABL gene, OR 2) Ph-like B-ALL with ABL-class kinase fusion, OR 3) relapsed or refractory T-cell ALL with ABL-class translocation. For GIST, patient must have progressed on imatinib, sunitinib, and regorafenib. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Gastrointestinal stromal tumor (GIST), metastatic chondrosarcoma, recurrent chordoma, T-cell acute lymphoblastic leukemia (ALL), Philadelphia (Ph)-like B-ALL |

STELARA

Products Affected

- Stelara Subcutaneous Solution 45 MG/0.5ML
- Stelara Subcutaneous Solution Prefilled Syringe

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) Patient had an inadequate response, intolerance, or contraindication to two of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Skyrizi (risankizumab-rzaa). For active psoriatic arthritis (PsA) (new starts only): patient had an inadequate response, intolerance, or contraindication to two of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Rinvoq (upadacitinib), Skyrizi (risankizumab-rzaa), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For moderately to severely active Crohn's disease (new starts only): patient had an inadequate response, intolerance, or contraindication to one of the following products: Humira (adalimumab) or Skyrizi (risankizumab-rzaa). For moderately to severely active ulcerative colitis (new starts): patient had an inadequate response, intolerance, or contraindication to Humira (adalimumab). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

STIVARGA

Products Affected

- Stivarga

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For gastrointestinal stromal tumors: The disease is progressive, locally advanced, unresectable, or metastatic. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Progressive gastrointestinal stromal tumors (GIST), osteosarcoma, glioblastoma, angiosarcoma, retroperitoneal/intra-abdominal soft tissue sarcoma, rhabdomyosarcoma, solitary fibrous tumor, and soft tissue sarcomas of the extremities, body wall, head and neck, advanced colorectal cancer. |

SUTENT

Products Affected

- SUNItinib Malate

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For renal cell carcinoma, the disease is relapsed, advanced, or stage IV. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Thyroid carcinoma (follicular, medullary, papillary, and Hurthle cell), soft tissue sarcoma (angiosarcoma, solitary fibrous tumor, and alveolar soft part sarcoma subtypes), recurrent chordoma, thymic carcinoma. |

SYMDEKO

Products Affected

- Symdeko

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The requested medication will not be used in combination with other medications containing ivacaftor. |
| Age Restrictions | 6 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SYMPAZAN

Products Affected

- Sympazan

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 2 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

SYNRIBO

Products Affected

- Synribo

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Follow-up therapy for chronic myeloid leukemia (CML) patients after hematopoietic stem cell transplant (HSCT) |

TABRECTA

Products Affected

- Tabrecta

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For recurrent, advanced, or metastatic NSCLC: Tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Treatment of recurrent or advanced non-small cell lung cancer (NSCLC). |

TADALAFIL (PAH)

Products Affected

- Adcirca
- Alyq
- Tadalafil (PAH)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TAFINLAR

Products Affected

- Tafinlar

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For brain metastases from melanoma, adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with trametinib. For unresectable or metastatic melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used as a single agent or in combination with trametinib. For non-small cell lung cancer and solid tumors: 1) The tumor is positive for a BRAF V600E mutation, and 2) The requested drug will be used in combination with trametinib. For thyroid carcinoma with papillary, follicular, or Hurthle histology: The tumor is positive for BRAF activating mutation (e.g., V600E or V600K). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Brain metastases from melanoma, thyroid carcinoma (papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma) |

TAGRISO

Products Affected

- Tagrisso

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC, the requested drug is used in any of the following settings: 1) The patient meets both of the following: a) patient has metastatic, advanced, or recurrent NSCLC (including brain and/or leptomeningeal metastases from NSCLC) and b) patient has a sensitizing EGFR mutation OR 2) Patient meets both of the following: a) request is for adjuvant treatment of NSCLC following tumor resection and b) patient has EGFR mutation-positive disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent or advanced non-small cell lung cancer (NSCLC), brain metastases from sensitizing EGFR mutation-positive NSCLC, leptomeningeal metastases from EGFR mutation-positive NSCLC |

TALTZ

Products Affected

- Taltz

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) The patient had an inadequate response, intolerance, or contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Skyrizi (risankizumab-rzaa). For active ankylosing spondylitis (new starts only): the patient had an inadequate response, intolerance, or contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active psoriatic arthritis (PsA) (new starts only): the patient had an inadequate response, intolerance, or contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Rinvoq (upadacitinib), Skyrizi (risankizumab-rzaa), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active axial spondyloarthritis (new starts only): Patient meets any of the following: 1) has had an inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial or 2) has an intolerance or contraindication to NSAIDs. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TALZENNA

Products Affected

- Talzenna

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For germline BRCA-mutated (gBRCAm) metastatic or recurrent breast cancer, the patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent germline breast cancer susceptibility gene (BRCA)-mutated breast cancer |

TARGRETIN TOPICAL

Products Affected

- Bexarotene
- Targretin External

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Mycosis fungoides, chronic or smoldering adult T-cell leukemia/lymphoma, primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma. |

TASIGNA

Products Affected

- Tasigna

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL), diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant: patient has experienced resistance or intolerance to imatinib or dasatinib. If patient experienced resistance to an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I, Y253H, E255K/V, and F359V/C/I mutations. For GIST, patient must have progressed on imatinib, sunitinib, and regorafenib. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), gastrointestinal stromal tumor (GIST) |

TAZAROTENE

Products Affected

- Tazarotene External Cream
- Tazorac External Cream 0.05 %

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For plaque psoriasis: 1) The requested drug is being prescribed to treat less than 20 percent of the patient's body surface area AND 2) The patient experienced an inadequate treatment response or intolerance to at least one topical corticosteroid OR has a contraindication that would prohibit a trial of topical corticosteroids. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TAZVERIK

Products Affected

- Tazverik

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | Epithelioid sarcoma: 16 years of age or older, Follicular lymphoma: 18 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TECENTRIQ

Products Affected

- Tecentriq

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For urothelial carcinoma, patient meets one of the following criteria: 1) Patient is ineligible for cisplatin therapy and tumors express PD-L1 (defined as PD-L1 stained tumor-infiltrating immune cells [IC] covering greater than or equal to 5 percent of the tumor area) OR 2) Patient is ineligible for any platinum containing chemotherapy. For non-small cell lung cancer (NSCLC): 1) the patient has recurrent, advanced or metastatic disease AND the requested drug will be used as any of the following: a) first-line treatment of tumors with high PD-L1 expression (defined as PD-L1 stained greater than or equal to 50 percent of tumor cells or PD-L1 stained tumor-infiltrating immune cells [IC] covering greater than or equal to 10 percent of the tumor area) and no EGFR or ALK genomic tumor aberrations, b) used in combination with carboplatin, paclitaxel, and bevacizumab, or in combination with carboplatin and albumin-bound paclitaxel for nonsquamous NSCLC, or c) the requested drug will be used as subsequent therapy or continuation maintenance therapy, OR 2) the patient has stage II to IIIA disease AND the requested drug will be used as adjuvant treatment following resection and platinum-based chemotherapy for tumors with PD-L1 expression on greater than or equal to 1 percent of tumor cells. For hepatocellular carcinoma, the requested drug will be used as initial treatment in combination with bevacizumab.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced non-small cell lung cancer, PD-L1 positive triple negative recurrent breast cancer in combination with paclitaxel protein-bound |

TECFIDERA

Products Affected

- Tecfidera

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TEMAZEPAM 30MG

Products Affected

- Temazepam Oral Capsule 30 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Prescriber must acknowledge that the benefit of therapy with the requested drug outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg). |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | This Prior Authorization requirement only applies to patients 65 years of age or older. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TEPMETKO

Products Affected

- Tepmetko

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TETRABENAZINE

Products Affected

- Tetrabenazine

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For treatment of chorea associated with Huntington's disease: The patient must have a prior inadequate response or intolerable adverse event with deutetrabenazine therapy. For treatment of tardive dyskinesia: The patient must have a prior inadequate response or intolerable adverse event with deutetrabenazine or valbenazine therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Tic disorders, tardive dyskinesia, hemiballismus, chorea not associated with Huntington's disease. |

TETRACYCLINE

Products Affected

- Tetracycline HCl Oral

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient will use the requested drug orally. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

THALOMID

Products Affected

- Thalomid

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For cachexia: Cachexia must be due to cancer or HIV infection. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Myelofibrosis-related anemia, recurrent aphthous stomatitis, recurrent human immunodeficiency virus (HIV)-associated aphthous ulcers, cachexia, HIV-associated diarrhea, acquired immunodeficiency syndrome (AIDS)-related Kaposi's sarcoma, Behcet's syndrome, chronic graft-versus-host disease, Crohn's disease, multicentric Castleman's disease. |

TIBSOVO

Products Affected

- Tibsovo

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Patient has disease with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation. For acute myeloid leukemia (AML): 1) patient has newly-diagnosed AML and meets one of the following: a) 75 years of age or older, b) patient has comorbidities that preclude use of intensive induction chemotherapy, or c) patient is 60 physiologic years of age or older and declines intensive induction chemotherapy, OR 2) patient is 60 physiologic years of age or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug, OR 3) patient has relapsed or refractory AML. For unresectable or metastatic cholangiocarcinoma: the requested drug will be used as subsequent treatment for progression on or after systemic treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Conventional (grades 1-3) or dedifferentiated chondrosarcoma |

TOBRAMYCIN

Products Affected

- Tobramycin Inhalation Nebulization Solution 300 MG/5ML

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For cystic fibrosis and non-cystic fibrosis bronchiectasis, the patient must meet one of the following: 1) Pseudomonas aeruginosa is present in the patient's airway cultures, OR 2) the patient has a history of Pseudomonas aeruginosa infection or colonization in the airways. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Non-cystic fibrosis bronchiectasis |

TOPICAL LIDOCAINE

Products Affected

- Glydo External Prefilled Syringe
- Lidocaine External Ointment 5 %
- Lidocaine HCl External Solution
- Lidocaine HCl Urethral/Mucosal External Gel
- Lidocaine-Prilocaine External Cream

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being used for topical anesthesia, AND 2) If the requested drug will be used as part of a compounded product, then all the active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 3 months |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TOPICAL TESTOSTERONES

Products Affected

- Androderm Transdermal Patch 24 Hour
- Testosterone Transdermal Gel 12.5 MG/ACT (1%), 25 MG/2.5GM (1%), 50 MG/5GM (1%)

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Primary or hypogonadotropic hypogonadism: 1) Request is for continuation of testosterone therapy and the patient had a confirmed low morning testosterone level according to current practice guidelines or your standard lab reference values before starting testosterone therapy [Note: Safety and efficacy of testosterone products in patients with age-related hypogonadism (also referred to as late-onset hypogonadism) have not been established.] OR 2) Request is not for continuation of testosterone therapy and the patient has at least two confirmed low morning testosterone levels according to current practice guidelines or your standard lab reference values [Note: Safety and efficacy of testosterone products in patients with age-related hypogonadism (also referred to as late-onset hypogonadism) have not been established.]. Gender dysphoria: The patient is able to make an informed decision to engage in hormone therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Gender Dysphoria |

TOPICAL TRETINOIN

Products Affected

- Avita
- Tretinoin External Cream
- Tretinoin External Gel 0.01 %, 0.025 %

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TRAZIMERA

Products Affected

- Trazimera

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year. |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neoadjuvant treatment for human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent or advanced unresectable HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive esophageal and esophagogastric junction cancer, HER2-positive advanced or recurrent uterine serous carcinoma, HER2-amplified colorectal cancer in combination with pertuzumab or lapatinib. |

TRELSTAR

Products Affected

- Trelstar Mixject Intramuscular Suspension
Reconstituted 11.25 MG, 3.75 MG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For gender dysphoria, patient meets either of the following (1 or 2): 1) the requested drug is used to suppress puberty and the patient is at Tanner stage 2 or greater, OR 2) patient is undergoing gender transition, and the patient will receive the requested drug concomitantly with gender-affirming hormones. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Gender dysphoria |

TREPROSTINIL INJ

Products Affected

- Treprostinil

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (World Health Organization [WHO] Group 1), the diagnosis was confirmed by right heart catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TRIENTINE

Products Affected

- Trientine HCl

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TRIKAFTA

Products Affected

- Trikafta

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The requested medication will not be used in combination with other medications containing ivacaftor. |
| Age Restrictions | 6 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TRUSELTIQ

Products Affected

- Truseltiq (100MG Daily Dose)
- Truseltiq (125MG Daily Dose)
- Truseltiq (50MG Daily Dose)
- Truseltiq (75MG Daily Dose)

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TRUXIMA

Products Affected

- Truxima

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis and 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | <p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, gastric mucosa-associated lymphoid tissue [MALT], nongastric MALT), Burkitt lymphoma, primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma with translocations of MYC and BCL2 and/or BCL6 (double/triple hit lymphoma), high-grade B-cell lymphoma not otherwise specified, histological transformation from follicular lymphoma to diffuse large B-cell lymphoma, histological transformation from nodal marginal zone lymphoma to diffuse large B-cell lymphoma, Castleman's disease, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis, immune checkpoint inhibitor-related toxicities, pemphigus vulgaris, pediatric Burkitt-like lymphoma (BLL), and pediatric mature B-cell acute leukemia (B-AL).</p> |

TUKYSA

Products Affected

- Tukysa

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer, including patients with brain metastases, who have received one or more lines of prior HER2-targeted therapy in the metastatic setting. |

TURALIO

Products Affected

- Turalio

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

TYMLOS

Products Affected

- Tymlos

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For postmenopausal osteoporosis: patient has ONE of the following: 1) a history of fragility fracture, OR 2) a pre-treatment T-score of less than or equal to -2.5 or pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment Fracture Risk Assessment Tool (FRAX) fracture probability AND patient has ANY of the following: a) indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) patient has failed prior treatment with or is intolerant to a previous injectable osteoporosis therapy, OR c) patient has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 months lifetime total for parathyroid hormone analogs (e.g., abaloparatide or teriparatide) |
| Other Criteria | Patient has high Fracture Risk Assessment Tool (FRAX) fracture probability if the 10 year probability is either greater than or equal to 20 percent for any major osteoporotic fracture or greater than or equal to 3 percent for hip fracture. If glucocorticoid treatment is greater than 7.5 mg (prednisone equivalent) per day, the estimated risk score generated with FRAX should be multiplied by 1.15 for major osteoporotic fracture and 1.2 for hip fracture. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

UBRELVY

Products Affected

- Ubrelyvy

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to one triptan 5-HT1 receptor agonist. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

UCERIS

Products Affected

- Budesonide ER Oral Tablet Extended Release 24 Hour

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one 5-aminosalicylic acid (5-ASA) therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 2 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VALCHLOR

Products Affected

- Valchlor

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Chronic or smoldering adult T-cell leukemia/lymphoma, Stage 2 or higher mycosis fungoides/Sezary syndrome, primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma, lymphomatoid papulosis. |

VELCADE

Products Affected

- Bortezomib Injection Solution Reconstituted 1 MG, 2.5 MG
- Bortezomib Injection Solution Reconstituted 3.5 MG
- Bortezomib Intravenous
- Velcade Injection

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Systemic light chain amyloidosis, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, multicentric Castleman's disease, adult T-cell leukemia/lymphoma, acute lymphoblastic leukemia, AIDS-related Kaposi's sarcoma, Hodgkin lymphoma, POEMS syndrome |

VEMLIDY

Products Affected

- Vemlidy

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | For chronic hepatitis B virus infection, the requested drug will be used in a patient who meets either of the following (new starts only): 1) inadequate virologic response or intolerable adverse event to tenofovir disoproxil fumarate OR 2) bone loss and mineralization defects or is at risk for bone loss and mineralization defects (for example, history of fragility fractures, advanced age, frailty, chronic glucocorticoid use, low T-scores, or increased fall risk). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VENCLEXTA

Products Affected

- Venclexta
- Venclexta Starting Pack

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For acute myeloid leukemia (AML), any of the following criteria must be met: 1) the patient's physiologic age is 60 years of age or older OR 2) the requested drug will be used as a component of repeating the initial successful induction regimen if late relapse OR 3) the patient has comorbidities that preclude use of intensive induction chemotherapy OR 4) the requested drug will be used for relapsed or refractory disease. For blastic plasmacytoid dendritic cell neoplasm (BPDCN), any of the following criteria must be met: 1) patient has systemic disease treated with palliative intent OR 2) patient has relapsed or refractory disease. For multiple myeloma, all of the following must be met: 1) the disease is relapsed or progressive AND 2) the requested drug will be used in combination with dexamethasone AND 3) patient has t(11:14) translocation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Mantle cell lymphoma, blastic plasmacytoid dendritic cell neoplasm (BPDCN), multiple myeloma, relapsed or refractory acute myeloid leukemia (AML), AML in patients 60 physiologic years of age or older. |

VENTAVIS

Products Affected

- Ventavis

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For pulmonary arterial hypertension (World Health Organization [WHO] Group 1), the diagnosis was confirmed by right heart catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VERSACLOZ

Products Affected

- Versacloz

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For the treatment of a severely ill patient with schizophrenia who failed to respond adequately to standard antipsychotic treatment (i.e., treatment-resistant schizophrenia), 1) the patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: A) aripiprazole, B) asenapine, C) olanzapine, D) quetiapine, E) risperidone, F) ziprasidone AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: A) Latuda, B) Rexulti, C) Secuado, D) Vraylar. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VERZENIO

Products Affected

- Verzenio

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer in combination with fulvestrant or an aromatase inhibitor, or as a single agent if progression on prior endocrine therapy and prior chemotherapy in the metastatic setting. |

V-GO

Products Affected

- V-Go 20
- V-Go 30
- V-Go 40

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has diabetes requiring insulin management with multiple daily injections AND 2) The patient is self-testing glucose levels 4 or more times per day OR the patient is using a continuous glucose monitor AND 3) The patient has experienced any of the following with the current diabetes regimen: inadequate glycemic control, recurrent hypoglycemia, wide fluctuations in blood glucose, dawn phenomenon with persistent severe early morning hyperglycemia, severe glycemic excursions. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | For continuation of therapy with an insulin pump, the patient has stable or improved glycemic control. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VIGABATRIN

Products Affected

- Vigabatrin
- Vigadrone

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For complex partial seizures (CPS): patient had an inadequate response to at least 2 antiepileptic drugs for CPS. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VITRAKVI

Products Affected

- Vitrakvi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Advanced, recurrent, or persistent neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors. |

VIZIMPRO

Products Affected

- Vizimpro

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For non-small cell lung cancer (NSCLC): 1) the disease is recurrent, advanced or metastatic, and 2) the patient has sensitizing EGFR mutation-positive disease. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced non-small cell lung cancer (NSCLC). |

VONJO

Products Affected

- Vonjo

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VORICONAZOLE

Products Affected

- Voriconazole Intravenous
- Voriconazole Oral

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The patient will use the requested drug orally or intravenously. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VOSEVI

Products Affected

- Vosevi

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh class B or C) |
| Required Medical Information | For hepatitis C: Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current AASLD treatment guidelines. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Criteria will be applied consistent with current AASLD-IDSA guidance. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VOTRIENT

Products Affected

- Votrient

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For renal cell carcinoma: The disease is advanced, relapsed, or stage IV. For soft tissue sarcoma (STS): The patient does not have an adipocytic soft tissue sarcoma. For uterine sarcoma: The disease is recurrent or metastatic. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Thyroid carcinoma (follicular, papillary, Hurthle cell, or medullary), uterine sarcoma. |

VUMERITY

Products Affected

- Vumerity

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

VYVANSE

Products Affected

- Vyvanse

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) The patient has a diagnosis of Attention-Deficit Hyperactivity Disorder (ADHD) or Attention Deficit Disorder (ADD) OR 2) The requested drug is being prescribed for the treatment of moderate to severe binge eating disorder (BED) in an adult. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

WELIREG

Products Affected

- Welireg

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XALKORI

Products Affected

- Xalkori

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC, the requested drug is used in any of the following settings: 1) the patient has recurrent, advanced or metastatic ALK-positive NSCLC, 2) the patient has recurrent, advanced or metastatic ROS-1 positive NSCLC, or 3) the patient has NSCLC with high-level MET amplification or MET exon 14 skipping mutation. For IMT, the disease is ALK-positive. For anaplastic large cell lymphoma, the disease is relapsed or refractory and ALK-positive. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent non-small cell lung cancer (NSCLC), NSCLC with high-level MET amplification or MET exon 14 skipping mutation, inflammatory myofibroblastic tumors (IMT). |

XELJANZ

Products Affected

- Xeljanz
- Xeljanz XR

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For moderately to severely active rheumatoid arthritis (new starts only): patient has experienced an inadequate treatment response or intolerance to at least one tumor necrosis factor (TNF) inhibitor. For active psoriatic arthritis (new starts only): 1) Patient has experienced an inadequate treatment response or intolerance to at least one TNF inhibitor AND 2) The requested drug is used in combination with a nonbiologic DMARD. For moderately to severely active ulcerative colitis (new starts only): Inadequate response, intolerance or contraindication to a tumor necrosis factor (TNF) blocker. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XERMELO

Products Affected

- Xermelo

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XGEVA

Products Affected

- Xgeva

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For hypercalcemia of malignancy: condition is refractory to intravenous (IV) bisphosphonate therapy or there is a clinical reason to avoid IV bisphosphonate therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XIFAXAN

Products Affected

- Xifaxan Oral Tablet 550 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed to reduce the risk of overt hepatic encephalopathy (HE) recurrence OR 2) The patient has the diagnosis of irritable bowel syndrome with diarrhea (IBS-D) AND 3) If the patient has previously received treatment with the requested drug, the patient has experienced a recurrence of symptoms AND 4) The patient has not already received an initial 14-day course of treatment and two additional 14-day courses of treatment with the requested drug OR 5) The patient has not previously received treatment with the requested drug. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Reduction in risk of overt HE recurrence: 6 months, IBS-D: 14 days |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XOLAIR

Products Affected

- Xolair

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For allergic asthma initial therapy: 1) Patient has positive skin test (or blood test) to at least 1 perennial aeroallergen, 2) Patient has baseline IgE level greater than or equal to 30 IU/mL, and 3) Patient has inadequate asthma control despite current treatment with both of the following medications at optimized doses: a) Inhaled corticosteroid, and b) Additional controller (long acting beta2-agonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For allergic asthma continuation therapy only: Patient's asthma control has improved on treatment with the requested drug since initiation of therapy. For chronic idiopathic urticaria (CIU) initial therapy: 1) Patient has been evaluated for other causes of urticaria, including bradykinin-related angioedema and IL-1-associated urticarial syndromes (auto-inflammatory disorders, urticarial vasculitis), and 2) Patient has experienced a spontaneous onset of wheals, angioedema, or both, for at least 6 weeks. For CIU continuation therapy: Patient has experienced a response (e.g., improved symptoms) since initiation of therapy. |
| Age Restrictions | For CIU: 12 years of age or older. For allergic asthma: 6 years of age or older. For nasal polyps: 18 years of age or older. |
| Prescriber Restrictions | |
| Coverage Duration | Allergic asthma and nasal polyps: Plan Year. CIU initial: 6 months. CIU continuation: Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XOSPATA

Products Affected

- Xospata

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FLT3 rearrangement: the disease is in chronic or blast phase. |
| Age Restrictions | 18 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FLT3 rearrangement |

XPOVIO

Products Affected

- Xpovio (100 MG Once Weekly)
- Xpovio (40 MG Once Weekly)
- Xpovio (40 MG Twice Weekly)
- Xpovio (60 MG Once Weekly)
- Xpovio (60 MG Twice Weekly)
- Xpovio (80 MG Once Weekly)
- Xpovio (80 MG Twice Weekly)

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XTANDI

Products Affected

- Xtandi

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

XYREM

Products Affected

- Xyrem

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | 1) The requested drug is being prescribed for the treatment of excessive daytime sleepiness in a patient 7 years of age or older with narcolepsy AND 2) The diagnosis has been confirmed by sleep lab evaluation AND 3)The patient experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, or methylphenidate) OR has a contraindication that would prohibit a trial of central nervous system (CNS) stimulant drugs (e.g., amphetamine, dextroamphetamine, or methylphenidate) [Note: Coverage of amphetamines may require prior authorization.] AND 4) If the patient is 18 years of age or older, the patient experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) wakefulness promoting drug (e.g., armodafinil) OR has a contraindication that would prohibit a trial of central nervous system (CNS) wakefulness promoting drugs (e.g., armodafinil) [Note: coverage of armodafinil may require prior authorization.] OR 5) The requested drug is being prescribed for the treatment of cataplexy in a patient 7 years of age or older with narcolepsy AND 6) The diagnosis has been confirmed by sleep lab evaluation. |
| Age Restrictions | 7 years of age or older |
| Prescriber Restrictions | Prescribed by or in consultation with a sleep disorder specialist or neurologist. |
| Coverage Duration | Plan Year |
| Other Criteria | If the request is for a continuation of therapy, then the patient experienced a decrease in daytime sleepiness with narcolepsy or a decrease in cataplexy episodes with narcolepsy. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ZARXIO

Products Affected

- Zarxio

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Use of the requested product within 24 hours prior to or following chemotherapy. |
| Required Medical Information | For prophylaxis or treatment of myelosuppressive chemotherapy-induced febrile neutropenia (FN) patient must meet both of the following: 1) Patient has a solid tumor or non-myeloid cancer, and 2) Patient has received, is currently receiving, or will be receiving treatment with myelosuppressive anti-cancer therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 months |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Neutropenia in myelodysplastic syndromes (MDS), agranulocytosis, neutropenia in aplastic anemia, human immunodeficiency virus (HIV)-related neutropenia, neutropenia related to renal transplant. |

ZEJULA

Products Affected

- Zejula

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | <p>For ovarian, fallopian tube, or primary peritoneal cancer, the requested drug is used in any of the following settings: 1) as maintenance treatment of stage II-IV epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in a complete or partial response to first-line platinum-based chemotherapy AND if it is known that the patient has breast cancer susceptibility gene (BRCA)-mutated disease, the patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib), 2) as maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in a complete or partial response to chemotherapy AND the patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib), 3) as treatment of advanced, persistent, or recurrent ovarian, fallopian tube, or primary peritoneal cancer in patients treated with three or more prior chemotherapy regimens and whose cancer is associated with homologous recombination deficiency (HRD) positive status defined by either a) a deleterious or suspected deleterious BRCA mutation AND if prescribed for advanced, persistent, or recurrent ovarian cancer with deleterious or suspected deleterious germline BRCA mutation, the patient experienced an unacceptable toxicity with a trial of Lynparza (olaparib), or b) genomic instability and progression more than six months after response to the last platinum-based chemotherapy, or 4) in combination with bevacizumab for platinum-sensitive persistent or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer.</p> |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|--|
| Off Label Uses | In combination with bevacizumab for persistent or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer for platinum-sensitive disease. |

ZELBORAF

Products Affected

- Zelboraf

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K) and 2) The requested drug will be used in combination with cobimetinib. For unresectable or metastatic melanoma: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K) and 2) the requested drug will be used as a single agent, or in combination with cobimetinib (with or without atezolizumab). For Erdheim-Chester Disease: Tumor is positive for BRAF V600 mutation. For non-small cell lung cancer: 1) Tumor is positive for the BRAF V600E mutation, and 2) The patient has recurrent, advanced, or metastatic disease. For thyroid carcinoma: 1) Tumor is positive for BRAF mutation, and 2) Patient has radioiodine refractory follicular, Hurthle cell, or papillary thyroid carcinoma. For hairy cell leukemia: The requested drug will be used for subsequent therapy. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Non-small cell lung cancer, hairy cell leukemia, thyroid carcinoma (i.e., papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system cancer (i.e., glioma, meningioma, astrocytoma), adjuvant systemic therapy for cutaneous melanoma. |

ZIRABEV

Products Affected

- Zirabev

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual. |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |

| PA Criteria | Criteria Details |
|-----------------------|---|
| Off Label Uses | <p>Breast cancer, central nervous system (CNS) tumor types: adult low-grade (WHO Grade II) infiltrative supratentorial astrocytoma/oligodendroglioma, adult intracranial and spinal ependymoma, anaplastic gliomas, adult medulloblastoma, primary central nervous system lymphoma, meningiomas, limited and extensive brain metastases, metastatic spine tumors, malignant pleural mesothelioma, epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer, including the following cancer types: carcinosarcoma (malignant mixed Mullerian tumors), clear cell carcinoma, mucinous carcinoma, grade 1 endometrioid carcinoma, low-grade serous carcinoma, ovarian borderline epithelial tumors (low malignant potential) with invasive implants, and malignant sex cord-stromal tumors, soft tissue sarcoma types: angiosarcoma and solitary fibrous tumor/hemangiopericytoma, uterine neoplasms, endometrial carcinoma, vulvar squamous cell carcinoma, and ophthalmic-related disorders: diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma and retinopathy of prematurity, hepatocellular carcinoma, small bowel adenocarcinoma.</p> |

ZOLINZA

Products Affected

- Zolinda

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Mycosis fungoides, Sezary syndrome. |

ZONISADE

Products Affected

- Zonisade

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | For adjunctive treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom, Xcopri, Spritam OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules). |
| Age Restrictions | 16 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ZTALMY

Products Affected

- Ztalmy

| PA Criteria | Criteria Details |
|-------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 2 years of age or older |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

ZYDELIG

Products Affected

- Zydelig

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Relapsed or refractory chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), relapsed or refractory follicular lymphoma, and marginal zone lymphomas [nodal marginal zone lymphoma, gastric mucosa associated lymphoid tissue (MALT) lymphoma, non-gastric MALT lymphoma, and splenic marginal zone lymphoma]. |

ZYKADIA

Products Affected

- Zykadia Oral Tablet

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | For NSCLC: the patient has recurrent, advanced, or metastatic ALK-positive or ROS1-positive disease. For inflammatory myofibroblastic tumor: the disease is ALK-positive. For brain metastases from NSCLC: the patient has ALK-positive NSCLC. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications, Some Medically-accepted Indications. |
| Off Label Uses | Recurrent or advanced ALK-positive non-small cell lung cancer (NSCLC), recurrent, advanced, or metastatic ROS1-positive NSCLC, inflammatory myofibroblastic tumor (IMT), brain metastases from NSCLC. |

ZYPREXA RELPREVV

Products Affected

- ZyPREXA Relprevv

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | Tolerability with oral olanzapine has been established. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | Plan Year |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

