

Prior Authorization Criteria
Corticotropin (H.P. Acthar, Purified Cortrophin Gel)

All requests for Corticotropin (H.P. Acthar, Purified Cortrophin Gel) require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

For all requests for Corticotropin (H.P. Acthar, Purified Cortrophin Gel) all of the following criteria must be met:

- Documentation of current height (cm) and weight (kg) are required
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines. If a requested dose is above these recommendations, medical rationale must be submitted.
- For infused products, must provide member's weight, dose, frequency and titration schedule.
- For H.P. Acthar Gel requests excluding diagnosis of infantile spasms, member must use Purified Cortrophin Gel unless contraindicated or clinically significant adverse effects are experienced.

Coverage may be provided with a diagnosis of infantile spasms (West Syndrome) and the following criteria is met (H.P. Acthar **ONLY**):

- Member is less than 24 months of age
- Prescribed by a neurologist
- Medication is used as monotherapy
- Diagnosis is supported by documentation of epileptic spasms, arrest of psychomotor development, and EEG pattern of hypsarrhythmia (Hypsarrhythmia, which does not typically occur with other forms of epilepsy, can help to confirm a diagnosis of infantile spasms)
- **Initial Duration of Approval:** 1 month
- **Reauthorization criteria**
 - Dosing does not exceed FDA labeled dosing
 - Requires documentation showing the member's EEG with continued hypsarrhythmia after 2 weeks of treatment **OR** the member is continuing to experience spasms.
- **Reauthorization Duration of Approval:** 1 month

Coverage may be provided with a diagnosis of acute exacerbation of multiple sclerosis and the following criteria is met:

- Member is 18 years of age or older
- Must be prescribed by a neurologist or physician that specializes in the treatment of multiple sclerosis

- Member has tried and failed oral methylprednisone 0.5g daily for 5 days and intravenous methylprednisolone 1g/day for 3 to 5 days within the last 45 days or has a contraindication to corticosteroid therapy
- There is documentation or claims verifying the member is on a medication for the treatment of multiple sclerosis. If not on a disease modifying therapy, refer to care management
- **Initial Duration of Approval:** 3 weeks
- **Reauthorization criteria**
 - Dosing does not exceed FDA labeled dosing
 - Documentation of disease response with treatment as indicated by resolution of symptoms
 - Absence of unacceptable toxicity from the drug (e.g. GI bleeding, gastric ulcer, hypertension, hypokalemia, severe depression, frank psychotic manifestations, posterior subcapsular cataracts, glaucoma)
- **Reauthorization Duration of Approval:** 3 weeks

Coverage may be provided with a diagnosis of moderately to severely active psoriatic arthritis and the following criteria is met:

- Must provide documentation of at least 6 tender and 6 swollen joint counts and ≥ 30 minutes of morning joint stiffness
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Member has tried and failed three disease modifying anti-rheumatic drugs (DMARD) with a different mechanism of action for at least 12 weeks or has a contraindication to DMARD therapy
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Must provide documentation of 70% improvement in tender or swollen joint counts were achieved as well as a 70% improvement in at least three of the following criteria:
 - Patient assessment
 - Physician assessment
 - Pain scale
 - Disability/functional questionnaire
 - Acute phase reactant (ESR or CRP)
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided with a diagnosis rheumatoid arthritis or juvenile rheumatoid arthritis and the following criteria is met:

- Must be prescribed by a rheumatologist
- Member must have rheumatoid arthritis of at least 2 years duration
- Member has tried and failed three biologics with a different mechanism of action for at least 12 weeks or has contraindications to biologic therapy
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy

- Must have active disease as defined by at least 6 tender and 6 swollen joints
- Must provide laboratory documentation with an ESR rate of at least 28mm/hr or CRP at least 1.2 times the upper limit of normal
- Must be stable on a DMARD and prednisone for at least 4 weeks
- **Initial Duration of Approval:** 3 months
- **Reauthorization criteria**
 - Must provide documentation of reductions in tender joint count and swollen joint count
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided with a diagnosis of ankylosing spondylitis and the following criteria is met:

- Must be prescribed by a rheumatologist
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Member has tried and failed two nonsteroidal anti-inflammatory drugs (NSAIDs) or has a contraindication to NSAID therapy
- Member has tried and failed two tumor necrosis factor alpha antagonists (anti-TNF-alpha) or has a contraindication to anti-TNF-alpha therapy
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Must provide chart documentation demonstrating clinical benefit and tolerance to therapy
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided with a diagnosis of systemic lupus erythematosus and the following criteria is met:

- Must be prescribed by a rheumatologist
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Member has tried and failed prednisone or has a contraindication to prednisone
- Member has tried and failed two steroid-sparing immunosuppressive agents (e.g. azathioprine, cyclophosphamide, or methotrexate) or has a contraindication to all immunosuppressive agents
- Must provide chart documentation confirming at least 4 of the American College of Rheumatology (ACR) SLE Criteria (See Table 1)
- Must provide baseline Physician Global Assessment, Patient Global Assessment, or SLEDAI-2K score
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Must provide chart documentation demonstrating a decrease in Physician Global Assessment, Patient Global Assessment, or SLEDAI-2K score
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided with a diagnosis of systemic dermatomyositis (polymyositis) and the following criteria is met:

- Must be prescribed by a rheumatologist
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Member has tried and failed two steroid-sparing immunosuppressive agents (e.g. azathioprine, tacrolimus, or methotrexate) for at least 12 weeks or has a contraindication to all immunosuppressive agents
- Member has tried and failed rituximab for at least 12 weeks or has a contraindication to rituximab
- Member has tried and failed intravenous immune globulin (IVIG) therapy for at least 12 weeks or has a contraindication to IVIG
- Must have documentation of active myositis defined as one of the following:
 - Baseline Manual Muscle Testing (MMT-8) no greater than 125/150 and at least 2 of the following:
 - Patient global with a minimum value of 2.0 cm on a 10 cm visual analog scale (VAS)
 - Physician global with a minimum value of 2.0 cm on a 10 cm VAS scale
 - Health Assessment Questionnaire (HAQ) disability index with a minimum value of 0.25
 - Elevation of at least one of the muscle enzymes [which includes creatine kinase (CK), aldolase, lactate dehydrogenase (LDH), alanine aminotransferase (ALT) and aspartate aminotransferase (AST)] at a minimum level of 1.3 times the upper limit of normal.
 - Global extramuscular disease activity score with a minimum value of 1.0 cm on a 10 cm VAS scale [this measure is the physician's composite evaluation and is based on assessments of activity scores on the constitutional, cutaneous, skeletal, gastrointestinal, pulmonary and cardiac scales of the Myositis Disease Activity Assessment Tool (MDAAT)].
 - Cutaneous VAS score on MDAAT > 3 cm on a 10 cm VAS scale and at least 3 of the following:
 - Patient global with a minimum value of 2.0 cm on a 10 cm visual analog scale (VAS)
 - Physician global with a minimum value of 2.0 cm on a 10 cm VAS scale
 - Health Assessment Questionnaire (HAQ) disability index with a minimum value of 0.25
 - Elevation of at least one of the muscle enzymes [which includes creatine kinase (CK), aldolase, lactate dehydrogenase (LDH), alanine aminotransferase (ALT) and aspartate aminotransferase (AST)] at a minimum level of 1.3 times the upper limit of normal.
 - Global extramuscular disease activity score with a minimum value of 1.0 cm on a 10 cm VAS scale [this measure is the physician's composite evaluation and is based on assessments of activity scores on the constitutional, cutaneous, skeletal, gastrointestinal, pulmonary and cardiac scales of the Myositis Disease Activity Assessment Tool (MDAAT)].
- **Initial Duration of Approval:** 6 months

- **Reauthorization criteria**
 - Must provide chart documentation demonstrating a decrease in Physician Global Assessment, Patient Global Assessment, or SLEDAI-2K score
- **Reauthorization Duration of Approval: 1 year**

Coverage may be provided with a diagnosis of severe erythema multiforme or Stevens-Johnson syndrome and the following criteria is met:

- Must be prescribed by a rheumatologist, allergist, or dermatologist
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- **Initial Duration of Approval: 3 months**
- **Reauthorization criteria**
 - Must provide chart documentation demonstrating clinical benefit and tolerance to therapy
- **Reauthorization Duration of Approval: 3 months**

Coverage may be provided with a diagnosis of serum sickness and the following criteria is met:

- Must be prescribed by an allergist or immunologist
- Must provide laboratory documentation demonstrating neutropenia, development of reactive plasmacytoid lymphocytes, and elevated erythrocyte sedimentation rate or C-reactive protein.
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- **Initial Duration of Approval: 3 months**
- **Reauthorization criteria**
 - Must provide chart documentation demonstrating clinical benefit and tolerance to therapy
- **Reauthorization Duration of Approval: 3 months**

Coverage may be provided with a diagnosis of severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: keratitis, iritis, iridocyclitis, diffuse posterior uveitis and chorioiditis, optic neuritis, chorioretinitis, and anterior segment inflammation and the following criteria is met:

- Must be prescribed by an ophthalmologist
- Member has tried and failed two topical glucocorticoids or has a contraindication to topical glucocorticoid therapy
- Member has tried and failed two steroid-sparing immunosuppressive agents (e.g. azathioprine, mycophenolate, or methotrexate) or has a contraindication to all immunosuppressive agents
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- **Initial Duration of Approval: 3 months**
- **Reauthorization criteria**
 - Must provide chart documentation demonstrating clinical benefit and tolerance to therapy

- **Reauthorization Duration of Approval:** 3 months

Coverage may be provided with a diagnosis of sarcoidosis and the following criteria is met:

- Must be prescribed by a rheumatologist or pulmonologist
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Member must have been on a stable dose of ≥ 5 mg prednisone for ≥ 3 months
- Must have documentation of deterioration of pulmonary disease as defined by a decrease of 5% forced vital capacity (FVC) in the previous year
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Documentation of reduction in concurrent prednisone dose
 - Must provide chart documentation demonstrating clinical benefit and tolerance to therapy
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided with a diagnosis of nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus and the following criteria is met:

- Must be used to induce a diuresis or remission of proteinuria
- Must be prescribed by a nephrologist
- Must be experiencing an acute exacerbation of nephrotic syndrome
- Must have a documented trial and failure of, intolerance to, or contraindication to treatment with a cytotoxic/immunosuppressive regimen (e.g. cyclophosphamide, cyclosporine, mycophenolate)
- Must currently be using conventional symptomatic therapy regimen (diuretics, ACE inhibitors, Angiotensin Receptor Blockers (ARBs), albumin)
- Member has tried and failed two intravenous corticosteroids or has a contraindication to intravenous corticosteroid therapy
- Must provide baseline 24-hour proteinuria level or urine protein:creatinine ratio
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Must provide chart documentation of reduction in 24-hour proteinuria level or urine protein:creatinine ratio
- **Reauthorization Duration of Approval:** 1 year

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.

Table 1. 1997 Update of the 1982 American College of Rheumatology Revised Criteria for Classification of Systemic Lupus Erythematosus

Criterion	Definition
Malar Rash	Fixed erythema, flat or raised, over the malar eminences, tending to spare the nasolabial folds
Discoid Rash	Erythematous raised patches with adherent keratotic scaling and follicular plugging; atrophic scarring may occur in older lesions
Photosensitivity	Skin rash as a result of unusual reaction to sunlight, by patient's history or physician's observation
Oral Ulcers	Oral or nasopharyngeal ulceration, usually painless, observed by a physician
Arthritis	Nonerosive arthritis involving two or more peripheral joints, characterized by tenderness, swelling or effusion
Serositis	a) Pleuritis-convincing history of pleuritic pain or rub heard by a physician or evidence of pleural effusion <i>or</i> b) Pericarditis-documented by ECG or rub or evidence of pericardial effusion
Renal Disorder	a) Persistent proteinuria greater than 0.5 g per day or greater than 3+ if quantitation not performed <i>or</i> b) Cellular casts-may be red cell, hemoglobin, granular, tubular or mixed
Neurologic Disorder	a) Seizures-in the absence of offending drugs or known metabolic derangements; eg uremia, ketoacidosis, or electrolyte imbalance <i>or</i> b) Psychosis-in the absence of offending drugs or known metabolic derangements, eg uremia, ketoacidosis, or electrolyte imbalance
Hematologic Disorder	a) Hemolytic anemia-with reticulocytosis <i>or</i> b) Leukopenia-less than 4000=mm ³ total on two or more occasions <i>or</i> c) Lymphopenia-less than 1500=mm ³ on two or more occasions <i>or</i> d) Thrombocytopenia-less than 100 1000=mm ³ in the absence of offending drugs
Immunologic Disorder	a) Anti-DNA-antibody to native DNA in abnormal titer <i>or</i> b) Anti-SM-presence of antibody to Sm nuclear antigen <i>or</i> c) Positive finding of antiphospholipid antibodies based on: 1. An abnormal serum level of IgG or IgM anticardiolipin antibodies <i>or</i> 2. A positive test result for lupus anticoagulant using a standard method <i>or</i> 3. 3 A false positive serologic test for syphilis known to be positive for at least six months and confirmed by <i>Treponema pallidum</i> immobilization or fluorescent treponemal antibody absorption test
Antinuclear Antibody	An abnormal titer of antinuclear antibody by immunofluorescence or an equivalent assay at any point in time and in the absence of drugs known to be associated with drug induced lupus syndrome

**CORTICOTROPIN (H.P. ACTHAR, PURIFIED CORTROPHIN GEL)
PRIOR AUTHORIZATION FORM- PAGE 1 of 2**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

PROVIDER INFORMATION

Requesting Provider:	Provider NPI:
Provider Specialty:	Office Contact:
State license #:	Office NPI:
Office Address:	Office Phone:
	Office Fax:

MEMBER INFORMATION

Member Name:	DOB:
Member ID:	Member weight: Height:

REQUESTED DRUG INFORMATION

Medication:	Strength:
Directions:	Quantity: Refills:
Is the member currently receiving requested medication? <input type="checkbox"/> Yes <input type="checkbox"/> No	Date Medication Initiated:

Billing Information

This medication will be billed: at a pharmacy **OR** medically, JCODE: _____

Place of Service: Hospital Provider's office Member's home Other

Place of Service Information

Name:	NPI:
Address:	Phone:

MEDICAL HISTORY (Complete for ALL requests)

Diagnosis:	ICD Code:
------------	-----------

Please check the applicable diagnosis:

Infantile Spasms (West Syndrome)

- Does the member have epileptic spasms? Yes No
- Is treatment being used as monotherapy? Yes No
- Has an EEG pattern shown hypsarrhythmia? Yes No
- Is there an arrest of psychomotor development? Yes No

Multiple Sclerosis (MS), acute exacerbation

- Has the member tried and failed or have a contraindication to corticosteroids? Yes No
- Is the member on a medication for the treatment of multiple sclerosis? Yes No

Other: _____

For H.P. Acthar Gel requests excluding diagnosis of infantile spasms, is the member using Purified Cortrophin Gel first unless contraindicated or clinically significant adverse effects are experienced? Yes No

Will this be administered via intravenous route? Yes No

If the patient is less than 2 years old, are they suspected to have any congenital infections? Yes, provide suspected congenital infection _____ No

Has the patient had surgery recently? Yes, Provide date: _____ No

Does the patient have a sensitivity of porcine origin? Yes No

CURRENT or PREVIOUS THERAPY

Medication Name	Strength/ Frequency	Dates of Therapy	Status (Discontinued & Why/Current)



**CORTICOTROPIN (H.P. ACTHAR, PURIFIED CORTROPHIN GEL)
PRIOR AUTHORIZATION FORM (CONTINUED)– PAGE 2 of 2**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

MEMBER INFORMATION

Member Name:	DOB:
Member ID:	Member weight: Height:

REAUTHORIZATION

Has the member experienced a significant improvement with treatment? Yes No

Please describe or provide documentation: _____

If used for Infantile Spasms (West Syndrome):

Is the member still experiencing spasms? Yes No

Does the member have continued hysarrhythmia after 2 weeks of treatment? Yes No

If yes, please provide the member's EEG report.

SUPPORTING INFORMATION or CLINICAL RATIONALE

Prescribing Provider Signature

Date



Updated: 7/2022
PARP Approved: 8/2022