

Remicade (infliximab), Infliximab (unbranded)

Override(s)	Approval Duration
Prior Authorization	1 Year

Medications	Dosing Limit
Remicade (infliximab) 100 mg vial	5 mg/kg as frequently as every 8 weeks
Infliximab (unbranded) 100 mg vial	

Dosing Override Criteria:

- I. For initiation of therapy, may approve up to 5 mg/kg at weeks 0, 2, and 6; **OR**
- II. For Ankylosing Spondylitis (AS) may approve 5 mg/kg as frequent as every 6 weeks; **OR**
- III. For Rheumatoid Arthritis (RA), may approve dose escalation up to 10 mg/kg every 8 weeks **OR** 5 mg/kg every 4 weeks for individuals who have an incomplete response; **OR**
- IV. For Crohn’s Disease (CD), may approve dose escalation up to 10 mg/kg every 8 weeks if the individual has previously achieved response to infliximab at standard dosing and subsequently lost response; **OR**
- V. For pediatric individuals less than 18 years of age with severe Crohn’s Disease (CD) or severe Ulcerative Colitis (UC), may approve up to 10 mg/kg every 4 weeks for initial or continuation of therapy. Adults with CD or UC who initiated treatment at less than 18 years of age may continue current dosage (up to 10 mg/kg every 4 weeks) if stable.

For Ulcerative Colitis (UC), may approve increased dosing, up to 10 mg/kg every 8 weeks if the following criteria are met:

- A. Individual has been treated with standard maintenance dosing (i.e. 5 mg/kg every 8 weeks) for *at least* 2 doses or 16 weeks; **AND**
- B. The increased dosing is being prescribed by or in consultation with a gastroenterologist;
- AND**
- C. Individual initially achieved an adequate response to standard maintenance dosing but has subsequently lost response, as determined by the prescriber; **OR**
- D. Individual partially responded but had an inadequate response to standard maintenance dosing as determined by the prescriber;
- AND**
- E. Symptoms, if present, are not due to active infections or any other gastrointestinal disorder other than the primary disease; **AND**
- F. Requested dosing does not exceed up to up to 10 mg/kg every 8 weeks.

Initial approval duration for increased dosing for UC: 16 weeks

Requests for continued escalated dosing for UC may be approved if the following criteria are met:

- A. Requested dosing does not exceed up to 10 mg/kg every 8 weeks; **AND**
- B. Individual has subsequently regained response or achieved adequate response following increased dosing, as shown by improvement in signs and symptoms of the disease

(including but not limited to reduction in stool frequency/bloody stools, improvement abdominal pain, or endoscopic response); **AND**

- C. Individual is not experiencing unacceptable adverse effects from increased dosing; **AND**
- D. Individual will be assessed regularly for dose de-escalation.

Continued approval duration for increased dosing for UC: 6 months

For UC, Increased dosing may not be approved for the following:

- A. Individual has had no response to infliximab at standard maintenance dosing (i.e. 5 mg/kg every 8 weeks); **OR**

Individual is requesting dose escalation in absence of signs and symptoms of the disease (for example, requesting based on results of therapeutic drug level or anti-drug antibody testing alone).

APPROVAL CRITERIA

Initial requests for Remicade (infliximab) or Infliximab (unbranded) may be approved for the following:

- I. Crohn's disease (CD) when each of the following criteria are met:
 - A. Individual is 6 year of age or older with moderate to severe CD; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy (such as systemic corticosteroids or immunosuppressants [such as thiopurines or methotrexate]); **OR**
 - C. Individual has a contraindication to systemic corticosteroids or thiopurines or methotrexate;

OR

- D. Individual is 6 years of age of older with fistulizing CD;

OR

- II. Ulcerative colitis (UC) when each of the following criteria are met:
 - A. Individual is 6 years of age or older with moderate to severe UC; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy (such as 5-Aminosalicylic acid products, systemic corticosteroids, or immunosuppressants [such as thiopurines]); **OR**
 - C. Individual has a contraindication to 5-ASA products or systemic corticosteroids or thiopurines;

OR

- III. Rheumatoid arthritis (RA) when each of the following criteria are met:
 - A. Individual is 18 years of age or older with moderate to severe RA; **AND**
 - B. Documentation is provided that individual has had an inadequate response to methotrexate titrated to maximally tolerated dose (ACR 2021); **OR**
 - C. Documentation is provided that if methotrexate is not tolerated, individual has had an inadequate response to or is intolerant of other conventional therapy (sulfasalazine, leflunomide, or hydroxychloroquine); **OR**
 - D. Documentation is provided that individual has a contraindication to methotrexate, sulfasalazine, leflunomide, and hydroxychloroquine;

OR

- IV. Ankylosing spondylitis (AS) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe AS; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy [such as NSAIDs or nonbiologic disease modifying anti-rheumatoid drugs (DMARDs) (such as sulfasalazine)] (ACR 2019); **OR**
 - C. Individual has a contraindication to NSAIDs or sulfasalazine;

OR

- V. Psoriatic arthritis (PsA) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe PsA; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy [nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate, sulfasalazine, cyclosporine, or leflunomide)]; **OR**
 - C. Individual has a contraindication to methotrexate, sulfasalazine, cyclosporine, and leflunomide;

OR

- VI. Plaque psoriasis (Ps) (Psoriasis vulgaris) when each of the following criteria are met:
- A. Individual is 18 years of age or older with chronic moderate to severe (that is, extensive or disabling) plaque Ps with either of the following (AAD 2019):
 - 1. Plaque Ps (psoriasis vulgaris) involving greater than three percent (3%) body surface area (BSA); **OR**
 - 2. Plaque Ps (psoriasis vulgaris) involving less than or equal to three percent (3%) BSA involving sensitive areas or areas that significantly impact daily function (such as palms, soles of feet, head/neck, or genitalia);

AND

- B. Individual has had an inadequate response to or is intolerant of phototherapy or other systemic therapies (such as acitretin, cyclosporine, or methotrexate); **OR**
- C. Individual has a contraindication to phototherapy, acitretin, cyclosporine, and methotrexate;

OR

- VII. Polyarticular juvenile idiopathic arthritis (PJIA) when each of the following criteria are met (DP B IIb, Lahdenne 2003, Gerloni 2005):
- A. Individual is 2 years of age or older with moderate to severe PJIA; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy [nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate)]; **OR**
 - C. Individual has a contraindication to methotrexate;

OR

- VIII. Non-infectious uveitis (UV) when each of the following criteria are met (Levy-Clarke 2014):
- A. Individual has chronic, recurrent, treatment-refractory or vision-threatening disease; **AND**
 - B. Individual has had an inadequate response to or is intolerant of conventional therapy [such as corticosteroids or immunosuppressants (azathioprine, cyclosporine, or methotrexate)]; **OR**
 - C. Individual has a contraindication to corticosteroids, azathioprine,

cyclosporine, and methotrexate;

OR

- IX. Immune checkpoint inhibitor therapy-related toxicities in an individual with any of the following conditions (NCCN 2A):
- A. Moderate to severe diarrhea or colitis unresponsive to high-dose systemic corticosteroids; **OR**
 - B. Moderate to severe pneumonitis if no improvement after 48 hours of high-dose systemic corticosteroids; **OR**
 - C. Acute kidney injury/elevated serum creatinine if toxicity remains greater than stage 2 after 4-6 weeks of corticosteroids or if creatinine increases during steroid taper (or once off steroids); **OR**
 - D. Myocarditis if unresponsive to high-dose systemic corticosteroids; **OR**
 - E. Moderate to severe inflammatory arthritis unresponsive to corticosteroids or nonbiologic DMARDs; **OR**
 - F. Grade 1-4 uveitis that is refractory to high-dose systemic corticosteroids;

OR

- X. Acute Graft-versus-host disease (GVHD) when each of the following criteria are met (NCCN 2A)
- A. Individual has a diagnosis of steroid-refractory acute GVHD; **AND**
 - B. Individual is initiating infliximab in combination with systemic corticosteroids;

OR

- XI. Sarcoidosis when each of the following criteria are met (Baughman 2006):
- A. Individual is 18 years of age or older; **AND**
 - B. Individual has chronic, progressive, treatment-refractory disease; **AND**
 - C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to systemic corticosteroids; **AND**
 - D. Individual has had an inadequate response to or is intolerant of nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate or azathioprine); **OR**
 - E. Individual has a contraindication to methotrexate and azathioprine.

Continuation requests for Remicade (infliximab) or Infliximab (unbranded) may be approved if the following criterion is met:

- I. Individual has been receiving and is maintained on a stable dose of Remicade (infliximab) or Infliximab (unbranded). Medication samples/coupons/discount cards are excluded from consideration as a trial.; **AND**
- II. There is clinically significant improvement or stabilization in clinical signs and symptoms of the disease.

Requests for Remicade (infliximab) or Infliximab (unbranded) may not be approved for the following:

- I. In combination with oral or topical JAK inhibitors, ozanimod, etrasimod, apremilast, deucravacitinib, or any of the following biologic immunomodulators: Other TNF antagonists, IL-23 inhibitors, IL-17 inhibitors, IL-6 inhibitors, IL-1 inhibitors, vedolizumab, ustekinumab, abatacept, rituximab, or natalizumab; **OR**

- II. Tuberculosis, other active serious infections, or a history of recurrent infections [repeat testing not required for ongoing authorization]; **OR**
- III. If initiating therapy, individual has not had a tuberculin skin test (TST), or a Center for Disease Control (CDC-) and Prevention -recommended equivalent, to evaluate for latent tuberculosis (unless switching therapy from another targeted immune modulator and no new risk factors); **OR**
- IV. When the above criteria are not met and for all other indications.

Note:

TNFi have black box warnings for serious infections and malignancy. Individuals treated with TNFi are at increased risk for developing serious infections that may lead to hospitalization or death. Most individuals who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. TNFi should be discontinued if an individual develops a serious infection or sepsis. Individuals should be tested for latent tuberculosis (TB) before TNFi use and during therapy. Treatment for latent TB should be initiated prior to TNFi use. Risks and benefits of TNFi should be carefully considered prior to initiation of therapy in individuals with chronic or recurrent infection. Lymphoma and other malignancies have been reported in children and adolescents treated with TNFi. Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL) have been reported in individuals treated with TNFi. Almost all cases had received treatment with azathioprine or 6-mercaptopurine concomitantly with a TNFi at or prior to diagnosis. It is uncertain whether HSTCL is related to the use of a TNFi or a TNFi in combination with these other immunosuppressants.

Key References:

1. Alikhan A, Sayed C, Alavi A et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations: Part II: Topical, intralesional, and systemic medical management. *J Am Acad Dermatol.* 2019; 81:91-101.
2. Brunner HI, Ruperto N, Tzaribachev N, et al. Subcutaneous golimumab for children with active polyarticular-course juvenile idiopathic arthritis: results of a multicentre, double-blind, randomised-withdrawal trial. *Ann Rheum Dis.* 2018; 77(1):21-29.
3. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: October 20, 2022.
4. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
5. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2022; Updated periodically.
6. NCCN Drugs & Biologics Compendium (NCCN Compendium®) 2022 National Comprehensive Cancer Network, Inc. Available at: NCCN.org. Updated periodically. Accessed on: October 14, 2022.
7. Centers for Disease Control and Prevention (CDC). Tuberculosis (TB). Available at: <https://www.cdc.gov/tb/topic/basics/risk.htm>. Last updated: March 18, 2016. Accessed October 28, 2022.
8. Fraenkel L, Bathon JM, England BR et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care & Research.* 2021;73(7):924-939.
9. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019; 80: 1029-72.
10. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum.* 2019; 71(1): 5-32.
11. Feuerstein JD, Ho EY, Shmidt E et al. American Gastroenterological Association Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology* 2021; 160:2496-2508.
12. Feuerstein JD, Issacs KL, Schneider Y, et al. American Gastroenterological Association Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology* 2020; 158:1450-1461.
13. Rubin DT, Ananthakrishnan AN, Siegel CA et al. American College of Gastroenterology Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol* 2019; 114:384-413.
14. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/ Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019; 71(10):1599-1613.

15. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Oligoarthritis, Temporomandibular Joint Arthritis, and Systemic Juvenile Idiopathic Arthritis. *Arthritis Rheum.* 2022; 74(4):553-569.
16. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Entesitis. *Arthritis Rheum.* 2019; 71(6):846-863.
17. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care & Research.* 2011; 63(4):465-482.
18. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders; American Uveitis Society subcommittee. *Ophthalmology.* 2014; 121(3):785-796.
19. Baughman RP, Drent M, et al. Infliximab therapy in patients with chronic sarcoidosis and pulmonary involvement. *Am J Respir Crit Care Med.* 2006; 174:795-802.
20. Lahdenne P, Vahasalo P, & Honkanen V: Infliximab or etanercept in the treatment of children with refractory juvenile idiopathic arthritis: an open label study. *Ann Rheum Dis* 2003; 62(3):245-247.
21. Gerloni V, Pontikaki I, Gattinara M, et al: Efficacy of repeated intravenous infusions of an anti-tumor necrosis factor alpha monoclonal antibody, infliximab, in persistently active, refractory juvenile idiopathic arthritis: results of an open-label prospective study. *Arthritis Rheum* 2005; 52(2):548-553.
22. Genovese MC, Sanchez-Burson J, Oh M, Balazs E, Neal J, Everding A, Hala T, Wojciechowski R, Fanjiang G, Cohen S. Comparative clinical efficacy and safety of the proposed biosimilar ABP 710 with infliximab reference product in patients with rheumatoid arthritis. *Arthritis Res Ther.* 2020 Mar 26;22(1):60. doi: 10.1186/s13075-020-2142-1. PMID: 32216829; PMCID: PMC7098142.
23. Smolen, J.S. et al. Safety, immunogenicity and efficacy after switching from reference infliximab to biosimilar SB2 compared with continuing reference infliximab and SB2 in patients with rheumatoid arthritis: results of a randomised, double-blind, phase III transition study. *Ann. Rheum. Dis.* 77, 234–240 (2018).
24. Jorgensen KK, Olsen IC, Goll GL, et al. Switching from originator infliximab to biosimilar CT-P13 compared with maintained treatment with originator infliximab (NOR-SWITCH): a 52-week, randomised, double-blind, non-inferiority trial. *Lancet.* 2017;389:2304-2316.
25. Alten R, Batko B, Hala T, et al. Randomised, double-blind, phase III study comparing the infliximab biosimilar, PF-06438179/GP1111, with reference infliximab: efficacy, safety and immunogenicity from week 30 to week 54. *RMD Open.* 2019 Mar 28;5(1):e000876. doi: 10.1136/rmdopen-2018-000876. PMID: 30997153; PMCID: PMC6446180.
26. Cohen S, Pablos JL, Pavelka K, et al. An open-label extension study to demonstrate long-term safety and efficacy of ABP 501 in patients with rheumatoid arthritis. *Arthritis Res Ther.* 2019;21:84. doi: 10.1186/s13075-019-1857-3
27. Wiland P, Jeka S, Dokoupilová E, et al. Switching to Biosimilar SDZ-ADL in Patients with Moderate-to-Severe Active Rheumatoid Arthritis: 48-Week Efficacy, Safety and Immunogenicity Results From the Phase III, Randomized, Double-Blind ADMYRA Study. *BioDrugs.* 2020 Dec;34(6):809-823. doi: 10.1007/s40259-020-00447-6. PMID: 33119861; PMCID: PMC7669771.

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.