

Medicare Part D: 5 Tier Closed Performance Formulary 2026

Please click here.

For Medicare Part D: Prior Authorization Criteria

Please click here.

For Medicare Part D: Step Therapy Criteria

Please click here.

Plans covered in Pennsylvania:

Community Blue Medicare HMO Distinct (HMO)

Community Blue Medicare HMO Merit (HMO)

Community Blue Medicare HMO Signature (HMO)

Community Blue Medicare PPO Signature (PPO)

Community Blue Medicare Plus PPO Signature (PPO)

Complete Blue HMO Distinct (HMO)

Complete Blue PPO Distinct (PPO)

Complete Blue PPO Merit (PPO)

Complete Blue PPO Signature (PPO)

Complete Blue Plus PPO Distinct (PPO)

Complete Blue Plus PPO Merit (PPO)

Freedom Blue PPO ValueRx (PPO)

Together Blue Medicare HMO Signature (HMO)

Security Blue HMO-POS ValueRx (HMO-POS)

For more recent information or other questions, please contact:

Community Blue Medicare HMO Pharmacy Service at 1-888-234-5397.

Complete Blue HMO Pharmacy Service at 1-888-234-5397.

Together Blue Medicare HMO Pharmacy Service at 1-888-328-5704.

Security Blue HMO-POS Pharmacy Service at 1-800-935-2583.

Community Blue Medicare PPO / Community Blue Medicare Plus PPO Pharmacy Service at 1-888-757-2946.

Complete Blue PPO / Complete Blue Plus PPO Pharmacy Service at 1-833-227-9375

Freedom Blue PPO Pharmacy Service at 1-800-550-8722.

For TTY users, *711 National Relay Service*, Monday through Sunday, 8:00 a.m. to 8:00 p.m. Visit medicare.highmark.com.

Formulary ID: 26018 Version: 16

Updated: 6/2026

Note to existing members: This formulary has changed since last year. Please review this document to make sure that it still contains the drugs you take.

When this Drug List (formulary) refers to “we,” “us,” or “our,” it means Highmark Senior Health Company, Highmark Choice Company, or Highmark Health Insurance Company.

When it refers to “plan” or “our plan,” it means Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO.

This document includes a Drug List (formulary) for our plan, which is current as of January 1, 2026. For an updated Drug List (formulary), please contact us. Our contact information, along with the date we last updated the Drug List (formulary), appears on the front and back cover pages.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2027, and from time to time during the year.

What is the Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO formulary?

In this document, we use the terms Drug List and formulary to mean the same thing. A formulary is a list of covered drugs selected by our plan in consultation with a team of health care providers, which represents the prescription therapies believed to be a necessary part of a quality treatment program. Our plan will generally cover the drugs listed in our formulary as long as the drug is medically necessary, the prescription is filled at a Community Blue Medicare HMO, Complete Blue HMO, Security Blue HMO-POS, Together Blue HMO, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, or Freedom Blue PPO network pharmacy, and other plan rules are followed. For more information on how to fill your prescriptions, please review your Evidence of Coverage.

Can the formulary change?

Most changes in drug coverage happen on January 1, but we may add or remove drugs on the formulary during the year, move them to different cost-sharing tiers, or add new restrictions. We must follow the Medicare rules in making these changes. Updates to the formulary are posted monthly to our website: [medicare.highmark.com/formulary](https://www.medicare.highmark.com/formulary).

Changes that can affect you this year: In the below cases, you will be affected by coverage changes during the year:

- **Immediate substitutions of certain new versions of brand name drugs and original biological products.** We may immediately remove a drug from our formulary if we are replacing it with a certain new version of that drug that will appear on the same or lower cost-sharing tier and with the same or fewer restrictions. When we add a new version of a drug to our formulary, we may decide to keep the brand name drug or original biological product on our formulary, but immediately move it to a different cost-sharing tier or add new restrictions.

We can make these immediate changes only if we are adding a new generic version of a brand name drug, or adding certain new biosimilar versions of an original biological product, that was already on the formulary (for example, adding an interchangeable biosimilar that can be substituted for an original biological product by a pharmacy without a new prescription.)

If you are currently taking the brand name drug or original biological product, we may not tell you in advance before we make an immediate change, but we will later provide you with information about the specific change(s) we have made.

If we make such a change, you or your prescriber can ask us to make an exception and continue to cover for you the drug that is being changed. For more information, see the section below titled “How do I request an exception to the Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO’s formulary?”

Some of these drug types may be new to you. For more information, see the section below titled “What are original biological products and how are they related to biosimilars?”

- **Drugs removed from the market.** If a drug is withdrawn from sale by the manufacturer or the Food and Drug Administration (FDA) determines to be withdrawn for safety or effectiveness reasons, we may immediately remove the drug from our formulary and later provide notice to members who take the drug.
- **Other changes.** We may make other changes that affect members currently taking a drug. For instance, we may remove a brand name drug from the formulary when adding a generic equivalent or remove an original biological product when adding a biosimilar. We may also apply new restrictions to the brand name drug or original biological product, or move it to a different cost-sharing tier, or both. We may make changes based on new clinical guidelines. If we remove drugs from our formulary, add prior authorization, quantity limits and/or step therapy restrictions on a drug, or move a drug to a higher cost-sharing tier, we must notify affected members of the change at least 30 days before the change becomes effective. Alternatively, when a member requests a refill of the drug, they may receive a 31-day supply of the drug and notice of the change.

If we make these other changes, you or your prescriber can ask us to make an exception for you and continue to cover the drug you have been taking. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled “How do I request an exception to the Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO’s formulary?”

Changes that will not affect you if you are currently taking the drug. Generally, if you are taking a drug on our 2026 formulary that was covered at the beginning of the year, we will not discontinue or reduce coverage of the drug during the 2026 coverage year except as described above. This means these drugs will remain available at the same cost-sharing and with no new restrictions for those members taking them for the remainder of the coverage year. You will not get direct notice this year about changes that do not affect you. However, on January 1 of the next year, such changes would affect you, and it is important to check the formulary for the new benefit year for any changes to drugs.

The enclosed formulary is current as of June 1, 2026. To get updated information about the drugs covered by our plan, please contact us. Our contact information appears on the front and back cover pages. In the event of mid-year non-maintenance formulary changes, members will be notified by mail and prospective members will receive an update with this formulary. The most up-to-date formulary is available on our website [medicare.highmark.com/formulary](https://www.medicare.highmark.com/formulary).

How do I use the formulary?

There are two ways to find your drug within the formulary:

Medical Condition

The formulary begins on page 9. The drugs in this formulary are grouped into categories depending on the type of medical conditions that they are used to treat. For example, drugs used to treat a heart condition are listed under the category, “Cardiovascular – Hypertension & Lipids.” If you know what your drug is used for, look for the category name in the list that begins on page 9. Then look under the category name for your drug.

Alphabetical Listing

If you are not sure what category to look under, you should look for your drug in the Index that begins on page 9. The Index provides an alphabetical list of all of the drugs included in this document. Both brand name drugs and generic drugs are listed in the Index. Look in the Index and find your drug. Next to your drug, you will see the page number where you can find coverage information. Turn to the page listed in the Index and find the name of your drug in the first column of the list.

What are generic drugs?

Our plan covers both brand name drugs and generic drugs. A generic drug is approved by the FDA as having the same active ingredient as the brand name drug. Generally, generic drugs work just as well as and usually cost less than brand name drugs. There are generic drug substitutes available for many brand name drugs. Generic drugs usually can be substituted for the brand name drug at the pharmacy without needing a new prescription, depending on state laws.

What are original biological products and how are they related to biosimilars?

On the formulary, when we refer to drugs, this could mean a drug or a biological product. Biological products are drugs that are more complex than typical drugs. Since biological products are more complex than typical drugs, instead of having a generic form, they have alternatives that are called biosimilars. Generally, biosimilars work just as well as the original biological product and may cost less. There are biosimilar alternatives for some original biological products. Some biosimilars are interchangeable biosimilars and, depending on state laws, may be substituted for the original biological product at the pharmacy without needing a new prescription, just like generic drugs can be substituted for brand name drugs.

For discussion of drug types, please see the Evidence of Coverage, Chapter 3, Section 3.1, “The ‘Drug List’ tells which Part D drugs are covered.”

Are there any restrictions on my coverage?

Some covered drugs may have additional requirements or limits on coverage. These requirements and limits may include:

- **Prior Authorization:** Our plan requires you or your prescriber to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don't get approval, our plan may not cover the drug.
- **Quantity Limits:** For certain drugs, our plan limits the amount of the drug that we will cover. For example, our plan provides 31 tablets, per prescription for 100mg Drug A. This may be in addition to a standard one-month or three-month supply.
- **Step Therapy:** In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. For example, if Drug A and Drug B both treat your medical condition, our plan may not cover Drug B unless you try Drug A first. If Drug A does not work for you, our plan will then cover Drug B.

You can find out if your drug has any additional requirements or limits by looking in the formulary that begins on page 9. You can also get more information about the restrictions applied to specific covered drugs by visiting our website. We have posted online document(s) that explain(s) our prior authorization and step therapy restrictions. You may also ask us to send you a copy. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You can ask our plan to make an exception to these restrictions or limits or for a list of other, similar drugs that may treat your health condition. See the section, “How do I request an exception to the Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO’s formulary?” on page 6 for information about how to request an exception.

What if my drug is not on the formulary?

If your drug is not included in this formulary (list of covered drugs), you should first contact Pharmacy Service and ask if your drug is covered.

If you learn that our plan does not cover your drug, you have two options:

- You can ask Pharmacy Service for a list of similar drugs that are covered by our plan. When you receive the list, show it to your doctor and ask them to prescribe a similar drug that is covered by our plan.
- You can ask our plan to make an exception and cover your drug. See below for information about how to request an exception.

How do I request an exception to the Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO’s formulary?

You can ask our plan to make an exception to our coverage rules. There are several types of exceptions that you can ask us to make.

- You can ask us to cover a drug even if it is not on our formulary. If approved, this drug will be covered at a pre-determined cost-sharing level, and you would not be able to ask us to provide the drug at a lower cost-sharing level.
- You can ask us to waive a coverage restriction including prior authorization, step therapy, or a quantity limit on your drug. For example, for certain drugs, our plan limits the amount of the drug that we will cover. If your drug has a quantity limit, you can ask us to waive the limit and cover a greater amount.
- You can ask us to cover a formulary drug at a lower cost-sharing level unless the drug is on the specialty tier. If approved, this would lower the amount you must pay for your drug.

Generally, our plan will only approve your request for an exception if the alternative drugs included on the plan’s formulary, the lower cost-sharing drug, or applying the restriction would not be as effective for you and/or would cause you to have adverse effects.

You or your prescriber should contact us to ask for a tiering or, formulary exception, including an exception to a coverage restriction. ***When you request an exception, your prescriber will need to explain the medical reasons why you need the exception.*** Generally, we must make our decision within 72 hours of getting your prescriber's supporting statement. You can ask for an expedited (fast) decision if you believe, and we agree, that your health could be seriously harmed by waiting up to 72 hours for a decision. If we agree, or if your prescriber asks for a fast decision, we must give you a decision no later than 24 hours after we get your prescriber's supporting statement.

What can I do if my drug is not on the formulary or has a restriction?

As a new or continuing member in our plan you may be taking drugs that are not on our formulary. Or, you may be taking a drug that is on our formulary but has a coverage restriction, such as prior authorization. You should talk to your prescriber about requesting a coverage decision to show that you meet the criteria for approval, switching to an alternative drug that we cover, or requesting a formulary exception so that we will cover the drug you take. While you and your doctor determine the right course of action for you, we may cover your drug in certain cases during the first 90 days you are a member of our plan.

For each of your drugs that is not on our formulary or has a coverage restriction, we will cover a temporary 31-day supply. If your prescription is written for fewer days, we'll allow refills to provide up to a maximum 31-day supply of medication. If coverage is not approved, after your first 31-day supply, we will not pay for these drugs, even if you have been a member of the plan less than 90 days.

If you are a resident of a long-term care facility and you need a drug that is not on our formulary or if your ability to get your drugs is limited, but you are past the first 90 days of membership in our plan, we will cover a 31-day emergency supply of that drug while you pursue a formulary exception.

The above transition process will be implemented to accommodate you if you have an immediate need for a non-formulary drug or a drug that requires prior authorization due to a change in your level of care while you are waiting for an exception request to be processed.

For more information

For more detailed information about your plan's prescription drug coverage, please review your Evidence of Coverage and other plan materials.

If you have questions about your plan, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

If you have general questions about Medicare prescription drug coverage, please call Medicare at 1-800-MEDICARE (1-800-633-4227) 24 hours a day/ seven days a week. TTY users should call 1-877-486-2048. Or visit <http://www.medicare.gov>.

Community Blue Medicare HMO, Complete Blue HMO, Together Blue HMO, Security Blue HMO-POS, Community Blue Medicare PPO, Community Blue Medicare Plus PPO, Complete Blue PPO, Complete Blue Plus PPO, and Freedom Blue PPO formulary

The formulary that begins on the next page provides coverage information about the drugs covered by your plan. If you have trouble finding your drug in the list, turn to the Index that begins on page 9.

The first column of the chart lists the drug name. Brand name drugs are capitalized (e.g., ABELCET) and generic drugs are listed in lower-case italics (e.g., *abacavir*).

The information in the Requirements/Limits column tells you if your plan has any special requirements for coverage of your drug.

The following is a formulary Format Example Only:

Drug Name	Performance Drug Tier	Requirements/Limits
Anti-Infectives		
<i>XYZ DRUG</i>	NF	QL-28

Table of Contents

Anti - Infectives.....	3
Antineoplastic / Immunosuppressant Drugs.....	11
Autonomic / Cns Drugs, Neurology / Psych.....	18
Cardiovascular, Hypertension / Lipids.....	31
Dermatologicals/Topical Therapy.....	36
Diagnostics / Miscellaneous Agents.....	40
Ear, Nose / Throat Medications.....	41
Endocrine/Diabetes.....	42
Gastroenterology.....	46
Immunology, Vaccines / Biotechnology.....	48
Miscellaneous Supplies.....	50
Musculoskeletal / Rheumatology.....	50
Obstetrics / Gynecology.....	52
Ophthalmology.....	55
Respiratory And Allergy.....	56
Urologicals.....	59
Vitamins, Hematinics / Electrolytes.....	60

Requirements/Limits

LA = Limited access

PA = Prior authorization required

PA-BvD = This drug may be covered under Medicare part B or D depending on the circumstance.

Information may need to be submitted describing the use and setting of the drug to make the determination.

PA-NS = Prior authorization required for new starts only

QL = Quantity limit applies. The quantity limit is noted for each drug. For example, if the quantity limit is QL (90 EA per 180 days), the quantity limit would be 90 units per 180-day supply.

ST = Step therapy applies

ST-NS = Step therapy applies to new starts only

Drug Tier**T1** = Cost-Sharing Tier 1 includes preferred generic drugs. This is the lowest cost-sharing tier.**T2** = Cost-Sharing Tier 2 includes generic drugs.**T3** = Cost-Sharing Tier 3 includes preferred brand name drugs and may include some single-sourced drugs (those generic drugs made by a single manufacturer).**T4** = Cost-Sharing Tier 4 includes non-preferred brand name drugs and may include some single-sourced generic drugs (those generic drugs made by a single manufacturer).**T5** = Cost-Sharing Tier 5 includes specialty drugs. This is the highest cost-sharing tier.**lowercase italics** = Generic drugs**UPPERCASE BOLD** = Brand name drugs

Drug Name	Drug Tier	Requirements/Limits
Anti - Infectives		
<i>abacavir</i>	T4	
<i>abacavir-lamivudine</i>	T4	
<i>acyclovir oral capsule</i>	T2	
<i>acyclovir oral suspension 200 mg/5 ml</i>	T4	
<i>acyclovir oral tablet</i>	T2	
<i>acyclovir sodium intravenous solution</i>	T4	PA-BvD
<i>adefovir</i>	T4	
<i>albendazole</i>	T4	
<i>amantadine hcl oral capsule</i>	T3	QL (124 EA per 31 days)
<i>amantadine hcl oral solution</i>	T3	
<i>amantadine hcl oral tablet</i>	T3	
<i>amikacin injection solution 500 mg/2 ml</i>	T4	
<i>amoxicillin oral capsule</i>	T1	
<i>amoxicillin oral suspension for reconstitution</i>	T1	
<i>amoxicillin oral tablet</i>	T2	
<i>amoxicillin oral tablet, chewable 125 mg, 250 mg</i>	T1	
<i>amoxicillin-pot clavulanate oral suspension for reconstitution</i>	T2	
<i>amoxicillin-pot clavulanate oral tablet</i>	T2	
<i>amphotericin b</i>	T4	PA-BvD
<i>amphotericin b liposome</i>	T5	PA-BvD

Drug Name	Drug Tier	Requirements/Limits
<i>ampicillin oral capsule 500 mg</i>	T2	
<i>ampicillin sodium injection recon soln 1 gram, 10 gram</i>	T4	
<i>ampicillin sodium injection recon soln 2 gram</i>	T2	
<i>ampicillin-sulbactam injection</i>	T4	
APTIVUS	T5	
ARIKAYCE	T5	PA
<i>atazanavir</i>	T4	
<i>atovaquone</i>	T4	
<i>atovaquone-proguanil</i>	T3	
<i>azithromycin intravenous</i>	T2	
<i>azithromycin oral tablet</i>	T2	
<i>aztreonam</i>	T4	
BICILLIN C-R	T3	
BICILLIN L-A INTRAMUSCULAR SYRINGE 600,000 UNIT/ML	T3	
BIKTARVY	T5	QL (31 EA per 31 days)
BLUJEP	T4	QL (20 EA per 5 days)
<i>caspofungin</i>	T4	
CAYSTON	T5	PA
<i>cefaclor oral capsule 500 mg</i>	T2	
<i>cefadroxil oral capsule</i>	T2	
<i>cefadroxil oral suspension for reconstitution 250 mg/5 ml</i>	T2	
<i>cefadroxil oral suspension for reconstitution 500 mg/5 ml</i>	T4	
<i>cefadroxil oral tablet</i>	T3	
<i>cefazolin injection recon soln 1 gram, 500 mg</i>	T2	
<i>cefazolin intravenous recon soln 10 gram</i>	T4	
<i>cefdinir oral capsule</i>	T2	
<i>cefepime injection</i>	T4	
<i>cefixime oral capsule</i>	T2	
<i>cefoxitin</i>	T4	
<i>cefpodoxime oral suspension for reconstitution 100 mg/5 ml</i>	T4	
<i>cefpodoxime oral suspension for reconstitution 50 mg/5 ml</i>	T3	
<i>cefpodoxime oral tablet</i>	T3	
<i>cefprozil</i>	T2	
<i>ceftaroline fosamil</i>	T5	

Drug Name	Drug Tier	Requirements/Limits
<i>ceftazidime</i>	T4	
<i>ceftriaxone injection recon soln 1 gram, 10 gram, 2 gram, 250 mg, 500 mg</i>	T2	
<i>cefuroxime axetil oral tablet</i>	T2	
<i>cefuroxime sodium injection recon soln 750 mg</i>	T3	
<i>cefuroxime sodium intravenous recon soln 1.5 gram</i>	T3	
<i>cephalexin oral capsule 250 mg, 500 mg</i>	T2	
<i>cephalexin oral suspension for reconstitution</i>	T2	
<i>chloroquine phosphate oral tablet 250 mg</i>	T3	QL (50 EA per 30 days)
<i>chloroquine phosphate oral tablet 500 mg</i>	T3	QL (25 EA per 30 days)
CIMDUO	T5	QL (31 EA per 31 days)
<i>ciprofloxacin hcl oral tablet 250 mg, 500 mg, 750 mg</i>	T1	
<i>ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml</i>	T2	
<i>clarithromycin oral suspension for reconstitution</i>	T4	
<i>clarithromycin oral tablet</i>	T2	
<i>clarithromycin oral tablet extended release 24 hr</i>	T2	
<i>clindamycin hcl</i>	T2	
<i>clindamycin in 5 % dextrose</i>	T2	
CLINDAMYCIN PEDIATRIC	T4	
<i>clindamycin phosphate injection</i>	T2	
<i>clotrimazole mucous membrane</i>	T2	
COARTEM	T4	
<i>colistin (colistimethate na)</i>	T4	
CRESEMBA ORAL	T5	
<i>dapsone oral</i>	T3	
<i>daptomycin intravenous recon soln 350 mg</i>	T4	
<i>daptomycin intravenous recon soln 500 mg</i>	T5	
<i>darunavir oral tablet 600 mg</i>	T4	
<i>darunavir oral tablet 800 mg</i>	T5	
DELSTRIGO	T5	QL (31 EA per 31 days)
DESCOVY	T5	QL (31 EA per 31 days)
<i>dicloxacillin</i>	T2	
DOVATO	T5	QL (31 EA per 31 days)
DOXY-100	T4	
<i>doxycycline hyclate intravenous</i>	T4	
<i>doxycycline hyclate oral capsule</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>doxycycline hyclate oral tablet 100 mg, 20 mg</i>	T2	
<i>doxycycline hyclate oral tablet, delayed release (dr/ec) 100 mg</i>	T4	
<i>doxycycline monohydrate oral capsule 100 mg, 50 mg</i>	T2	
<i>doxycycline monohydrate oral tablet 100 mg, 50 mg</i>	T2	
EDURANT	T5	
EDURANT PED	T5	
<i>efavirenz oral tablet</i>	T4	
<i>efavirenz-emtricitabin-tenofof</i>	T5	
<i>efavirenz-lamivuv-tenofof disop</i>	T5	QL (31 EA per 31 days)
<i>emtricitabine</i>	T4	
<i>emtricitabine-tenofovir (tdf) oral tablet 100-150 mg, 167-250 mg, 200-300 mg</i>	T4	
<i>emtricitabine-tenofovir (tdf) oral tablet 133-200 mg</i>	T5	
<i>emtricitabine-rilpivirine-tenofof df</i>	T5	
EMTRIVA ORAL SOLUTION	T3	
<i>entecavir</i>	T4	
<i>ertapenem</i>	T4	
<i>erythromycin oral tablet</i>	T4	
<i>ethambutol</i>	T2	
<i>etravirine</i>	T5	
EVOTAZ	T5	
<i>famciclovir</i>	T3	
<i>fidaxomicin</i>	T5	QL (20 EA per 10 days)
<i>fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml</i>	T2	
<i>fluconazole in nacl (iso-osm) intravenous piggyback 400 mg/200 ml</i>	T3	
<i>fluconazole oral suspension for reconstitution</i>	T3	
<i>fluconazole oral tablet</i>	T2	
<i>flucytosine</i>	T5	
<i>fosamprenavir</i>	T5	
<i>fosfomycin tromethamine</i>	T4	
<i>gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml</i>	T2	
<i>gentamicin injection</i>	T4	

Drug Name	Drug Tier	Requirements/Limits
GENVOYA	T5	
<i>griseofulvin microsize</i>	T4	
<i>griseofulvin ultramicrosize oral tablet 125 mg, 250 mg</i>	T4	
<i>hydroxychloroquine oral tablet 200 mg</i>	T2	QL (93 EA per 31 days)
<i>imipenem-cilastatin</i>	T4	
IMPAVIDO	T5	
INTELENCE ORAL TABLET 25 MG	T4	
ISENTRESS HD	T5	
ISENTRESS ORAL POWDER IN PACKET	T5	
ISENTRESS ORAL TABLET	T5	
ISENTRESS ORAL TABLET,CHEWABLE 100 MG	T5	
ISENTRESS ORAL TABLET,CHEWABLE 25 MG	T3	
<i>isoniazid oral solution</i>	T2	
<i>isoniazid oral tablet</i>	T1	
<i>itraconazole oral capsule</i>	T4	PA
<i>ivermectin oral</i>	T3	PA
JULUCA	T5	
KALETRA ORAL SOLUTION	T5	
<i>ketoconazole oral</i>	T2	
LAGEVRIO (EUA)	T3	
<i>lamivudine oral solution</i>	T2	
<i>lamivudine oral tablet 100 mg</i>	T3	
<i>lamivudine oral tablet 150 mg, 300 mg</i>	T2	
<i>lamivudine-zidovudine</i>	T3	
<i>levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml</i>	T2	
<i>levofloxacin oral solution</i>	T4	
<i>levofloxacin oral tablet</i>	T2	
<i>linezolid in dextrose 5%</i>	T4	
<i>linezolid oral tablet</i>	T4	
LIVTENCITY	T5	PA; QL (372 EA per 31 days)
<i>lopinavir-ritonavir oral tablet</i>	T4	
<i>maraviroc oral tablet 150 mg</i>	T5	
<i>maraviroc oral tablet 300 mg</i>	T3	
MAVYRET ORAL PELLETS IN PACKET	T5	PA; QL (140 EA per 28 days)
MAVYRET ORAL TABLET	T5	PA; QL (84 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
<i>mefloquine</i>	T2	
<i>meropenem intravenous recon soln 1 gram, 500 mg</i>	T3	
<i>methenamine hippurate</i>	T4	
<i>metronidazole in nacl (iso-os)</i>	T2	
<i>metronidazole oral tablet 250 mg, 500 mg</i>	T2	
<i>micafungin</i>	T4	
<i>minocycline oral capsule</i>	T2	
<i>minocycline oral tablet</i>	T2	
<i>moxifloxacin oral</i>	T3	
<i>moxifloxacin-sod.chloride(iso)</i>	T4	
<i>nafcillin injection recon soln 1 gram, 2 gram</i>	T4	
<i>nafcillin injection recon soln 10 gram</i>	T5	
<i>neomycin</i>	T2	
<i>nevirapine oral suspension</i>	T2	
<i>nevirapine oral tablet</i>	T2	
<i>nevirapine oral tablet extended release 24 hr 400 mg</i>	T4	
<i>nitazoxanide</i>	T5	
<i>nitrofurantoin macrocrystal oral capsule 100 mg</i>	T2	QL (90 EA per 365 days)
<i>nitrofurantoin macrocrystal oral capsule 50 mg</i>	T2	QL (180 EA per 365 days)
<i>nitrofurantoin monohyd/m-cryst</i>	T2	QL (90 EA per 365 days)
NORVIR ORAL POWDER IN PACKET	T4	
<i>nystatin oral suspension</i>	T2	
<i>nystatin oral tablet</i>	T2	
ODEFSEY	T5	QL (31 EA per 31 days)
<i>ofloxacin oral tablet 300 mg</i>	T4	
<i>ofloxacin oral tablet 400 mg</i>	T2	
ORLYNVAH	T4	QL (10 EA per 5 days)
<i>oseltamivir oral capsule 30 mg</i>	T2	QL (170 EA per 365 days)
<i>oseltamivir oral capsule 45 mg, 75 mg</i>	T2	QL (90 EA per 365 days)
<i>oseltamivir oral suspension for reconstitution</i>	T3	QL (1080 ML per 365 days)
<i>oxacillin injection recon soln 1 gram, 2 gram</i>	T2	
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (10)- 100 MG (10)	T3	QL (180 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (6)- 100 MG (5)	T3	QL (99 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 300 MG (150 MG X 2)-100 MG	T3	QL (270 EA per 365 days)

Drug Name	Drug Tier	Requirements/Limits
<i>penicillin g pot in dextrose intravenous piggyback 2 million unit/50 ml, 3 million unit/50 ml</i>	T4	
<i>penicillin g potassium injection recon soln 20 million unit</i>	T4	
<i>penicillin v potassium oral tablet</i>	T1	
<i>pentamidine inhalation</i>	T4	PA-BvD
<i>pentamidine injection</i>	T4	
PIFELTRO	T5	QL (62 EA per 31 days)
<i>piperacillin-tazobactam intravenous recon soln 2.25 gram, 3.375 gram, 4.5 gram, 40.5 gram</i>	T4	
<i>posaconazole oral tablet, delayed release (dr/ec)</i>	T5	PA
<i>praziquantel</i>	T4	
PREVYMIS ORAL PELLETS IN PACKET	T5	PA; QL (124 EA per 31 days)
PREVYMIS ORAL TABLET	T5	QL (31 EA per 31 days)
PREZCOBIX	T5	
PREZISTA ORAL SUSPENSION	T5	
PREZISTA ORAL TABLET 150 MG	T5	
PREZISTA ORAL TABLET 75 MG	T4	
PRIFTIN	T3	
<i>primaquine</i>	T3	
<i>pyrazinamide</i>	T4	
<i>pyrimethamine</i>	T5	PA
<i>quinine sulfate</i>	T4	PA; QL (42 EA per 28 days)
RELENZA DISKHALER	T3	
REYATAZ ORAL POWDER IN PACKET	T5	
<i>ribavirin oral capsule</i>	T3	
<i>ribavirin oral tablet 200 mg</i>	T3	
<i>rifabutin</i>	T4	
<i>rifampin</i>	T4	
<i>rimantadine</i>	T4	
<i>ritonavir</i>	T3	
RUKOBIA	T5	QL (62 EA per 31 days)
SELZENTRY ORAL SOLUTION	T5	
SIRTURO	T5	PA
<i>sofosbuvir-velpatasvir</i>	T5	PA; QL (28 EA per 28 days)
<i>streptomycin</i>	T5	
STRIBILD	T5	
<i>sulfadiazine</i>	T4	
<i>sulfamethoxazole-trimethoprim oral suspension</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>sulfamethoxazole-trimethoprim oral tablet</i>	T1	
SUNLENCA ORAL	T5	
SYMTUZA	T5	QL (31 EA per 31 days)
<i>tenofovir disoproxil fumarate</i>	T3	
<i>terbinafine hcl oral</i>	T1	QL (90 EA per 180 days)
<i>tetracycline oral capsule</i>	T4	
<i>tigecycline</i>	T4	
<i>tinidazole</i>	T2	
TIVICAY ORAL TABLET 50 MG	T5	
TIVICAY PD	T5	
TOBI PODHALER	T5	PA; QL (224 EA per 56 days)
<i>tobramycin in 0.225 % nacl</i>	T5	PA
<i>tobramycin inhalation</i>	T5	PA
<i>tobramycin sulfate injection solution</i>	T4	
<i>trimethoprim</i>	T2	
TRIUMEQ	T5	
TRIUMEQ PD	T4	QL (186 EA per 31 days)
<i>valacyclovir</i>	T2	
<i>valganciclovir oral recon soln</i>	T4	
<i>valganciclovir oral tablet</i>	T3	
<i>vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg</i>	T4	
<i>vancomycin oral capsule 125 mg</i>	T4	QL (124 EA per 31 days)
<i>vancomycin oral capsule 250 mg</i>	T4	QL (248 EA per 31 days)
VEMLIDY	T5	QL (31 EA per 31 days)
VIRACEPT ORAL TABLET	T5	
VIREAD ORAL POWDER	T5	
VIREAD ORAL TABLET 150 MG, 200 MG, 250 MG	T5	
VIVJOA	T4	PA; QL (18 EA per 84 days)
<i>voriconazole intravenous recon soln</i>	T5	PA
<i>voriconazole oral</i>	T4	
VOSEVI	T5	PA; QL (28 EA per 28 days)
XIFAXAN ORAL TABLET 200 MG	T4	QL (27 EA per 365 days)
XIFAXAN ORAL TABLET 550 MG	T5	PA; QL (62 EA per 31 days)
XOFLUZA ORAL TABLET 40 MG, 80 MG	T3	QL (9 EA per 365 days)
<i>zidovudine</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
Antineoplastic / Immunosuppressant Drugs		
<i>abiraterone oral tablet 250 mg</i>	T5	PA-NS; QL (124 EA per 31 days)
<i>abiraterone oral tablet 500 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
ABIRTEGA	T3	PA-NS; QL (124 EA per 31 days)
AKEEGA	T5	PA-NS; QL (62 EA per 31 days)
ALECENSA	T5	PA-NS; QL (248 EA per 31 days)
ALUNBRIG ORAL TABLET 180 MG, 90 MG	T5	PA-NS; QL (31 EA per 31 days)
ALUNBRIG ORAL TABLET 30 MG	T5	PA-NS; QL (186 EA per 31 days)
ALUNBRIG ORAL TABLETS,DOSE PACK	T5	PA-NS; QL (60 EA per 365 days)
<i>anastrozole</i>	T2	
AUGTYRO ORAL CAPSULE 160 MG	T5	PA-NS; QL (62 EA per 31 days)
AUGTYRO ORAL CAPSULE 40 MG	T5	PA-NS; QL (186 EA per 31 days)
AVMAPKI-FAKZYNJA	T5	PA-NS; QL (66 EA per 28 days)
AYVAKIT	T5	PA-NS; QL (31 EA per 31 days)
<i>azathioprine oral tablet 50 mg</i>	T2	PA-BvD
BALVERSA	T5	PA-NS
<i>bexarotene oral</i>	T5	PA-NS
<i>bexarotene topical</i>	T5	PA-NS; QL (60 GM per 28 days)
<i>bicalutamide</i>	T2	
BOSULIF ORAL CAPSULE 100 MG	T5	PA-NS; QL (186 EA per 31 days)
BOSULIF ORAL CAPSULE 50 MG	T5	PA-NS; QL (341 EA per 31 days)
BOSULIF ORAL TABLET 100 MG	T5	PA-NS; QL (93 EA per 31 days)
BOSULIF ORAL TABLET 400 MG, 500 MG	T5	PA-NS; QL (31 EA per 31 days)
BRAFTOVI	T5	PA-NS; QL (186 EA per 31 days)
BRUKINSA ORAL TABLET	T5	PA-NS; QL (62 EA per 31 days)
CABOMETYX	T5	PA-NS; QL (31 EA per 31 days)
CALQUENCE (ACALABRUTINIB MAL)	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 300 MG	T5	PA-NS; QL (31 EA per 31 days)
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
COMETRIQ ORAL CAPSULE 140 MG/DAY(80 MG X1-20 MG X3)	T5	PA-NS; QL (112 EA per 28 days)
COMETRIQ ORAL CAPSULE 60 MG/DAY (20 MG X 3/DAY)	T5	PA-NS; QL (84 EA per 28 days)
COPIKTRA	T5	PA-NS; QL (62 EA per 31 days)
COTELLIC	T5	PA-NS; LA; QL (63 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
<i>cyclophosphamide oral capsule</i>	T3	PA-BvD
<i>cyclophosphamide oral tablet 50 mg</i>	T3	PA-BvD
<i>cyclosporine modified oral capsule</i>	T2	PA-BvD
<i>cyclosporine modified oral solution</i>	T5	PA-BvD
<i>cyclosporine oral capsule</i>	T2	PA-BvD
DANZITEN	T5	PA-NS; QL (124 EA per 31 days)
<i>dasatinib oral tablet 100 mg, 140 mg, 50 mg, 70 mg, 80 mg</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>dasatinib oral tablet 20 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
DAURISMO ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
DAURISMO ORAL TABLET 25 MG	T5	PA-NS; QL (62 EA per 31 days)
ELIGARD	T4	ST-NS; QL (1 EA per 30 days)
ELIGARD (3 MONTH)	T4	ST-NS; QL (1 EA per 90 days)
ELIGARD (4 MONTH)	T4	ST-NS; QL (1 EA per 120 days)
ELIGARD (6 MONTH)	T4	ST-NS; QL (1 EA per 180 days)
ENSACOVE	T5	PA-NS; QL (62 EA per 31 days)
ENVARUSUS XR ORAL TABLET EXTENDED RELEASE 24 HR 0.75 MG, 1 MG	T4	PA-BvD
ENVARUSUS XR ORAL TABLET EXTENDED RELEASE 24 HR 4 MG	T5	PA-BvD
ERIVEDGE	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 240 MG	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 60 MG	T5	PA-NS; QL (93 EA per 31 days)
<i>erlotinib</i>	T5	PA-NS; QL (31 EA per 31 days)
EULEXIN	T4	
<i>everolimus (antineoplastic) oral tablet</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 2 mg, 5 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 3 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>everolimus (immunosuppressive) oral tablet 0.25 mg</i>	T4	PA-BvD
<i>everolimus (immunosuppressive) oral tablet 0.5 mg, 0.75 mg, 1 mg</i>	T5	PA-BvD
<i>exemestane</i>	T4	
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 120 MG	T5	PA-NS
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 80 MG	T4	PA-NS
FOTIVDA	T5	PA-NS; QL (21 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
FRUZAQLA ORAL CAPSULE 1 MG	T5	PA-NS; QL (84 EA per 28 days)
FRUZAQLA ORAL CAPSULE 5 MG	T5	PA-NS; QL (21 EA per 28 days)
GAVRETO	T5	PA-NS; QL (124 EA per 31 days)
<i>gefitinib</i>	T5	PA-NS; QL (31 EA per 31 days)
GENGRAF ORAL CAPSULE	T2	PA-BvD
GILOTRIF	T5	PA-NS; QL (31 EA per 31 days)
GOMEKLI ORAL CAPSULE 1 MG	T5	PA-NS; QL (126 EA per 28 days)
GOMEKLI ORAL CAPSULE 2 MG	T5	PA-NS; QL (84 EA per 28 days)
GOMEKLI ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (168 EA per 28 days)
HERNEXEOS	T5	PA-NS; QL (93 EA per 31 days)
<i>hydroxyurea</i>	T2	
HYRNUO	T5	PA-NS; QL (124 EA per 31 days)
IBRANCE ORAL CAPSULE 125 MG	T5	PA-NS; QL (21 EA per 28 days)
IBRANCE ORAL TABLET	T5	PA-NS; QL (21 EA per 28 days)
IBTROZI	T5	PA-NS; QL (93 EA per 31 days)
ICLUSIG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA	T5	PA-NS; QL (31 EA per 31 days)
<i>imatinib oral tablet 100 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>imatinib oral tablet 400 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
IMBRUVICA ORAL CAPSULE 140 MG	T5	PA-NS; QL (124 EA per 31 days)
IMBRUVICA ORAL CAPSULE 70 MG	T5	PA-NS; QL (31 EA per 31 days)
IMBRUVICA ORAL SUSPENSION	T5	PA-NS; QL (216 ML per 25 days)
IMBRUVICA ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
IMKELDI	T5	PA-NS; QL (280 ML per 28 days)
INLURIYO	T5	PA-NS; QL (62 EA per 31 days)
INLYTA	T5	PA-NS; QL (124 EA per 31 days)
INQOVI	T5	PA-NS; QL (5 EA per 28 days)
INREBIC	T5	PA-NS; QL (124 EA per 31 days)
ITOVEBI ORAL TABLET 3 MG	T5	PA-NS; QL (62 EA per 31 days)
ITOVEBI ORAL TABLET 9 MG	T5	PA-NS; QL (31 EA per 31 days)
IWILFIN	T5	PA-NS; QL (248 EA per 31 days)
JAKAFI	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG	T5	PA-NS; QL (70 EA per 28 days)
KISQALI FEMARA CO-PACK ORAL TABLET 600 MG/DAY(200 MG X 3)-2.5 MG	T5	PA-NS; QL (91 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (21 EA per 28 days)
KISQALI ORAL TABLET 400 MG/DAY (200 MG X 2)	T5	PA-NS; QL (42 EA per 28 days)
KISQALI ORAL TABLET 600 MG/DAY (200 MG X 3)	T5	PA-NS; QL (63 EA per 28 days)
KOSELUGO ORAL CAPSULE 10 MG	T5	PA-NS; QL (279 EA per 31 days)
KOSELUGO ORAL CAPSULE 25 MG	T5	PA-NS; QL (124 EA per 31 days)
KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG	T5	PA-NS; QL (600 EA per 30 days)
KOSELUGO ORAL CAPSULE, SPRINKLE 7.5 MG	T5	PA-NS; QL (360 EA per 30 days)
KRAZATI	T5	PA-NS; QL (186 EA per 31 days)
<i>lapatinib</i>	T5	PA-NS; QL (186 EA per 31 days)
LAZCLUZE ORAL TABLET 240 MG	T5	PA-NS; QL (30 EA per 30 days)
LAZCLUZE ORAL TABLET 80 MG	T5	PA-NS; QL (60 EA per 30 days)
<i>lenalidomide</i>	T5	PA-NS; QL (21 EA per 28 days)
LENVIMA	T5	PA-NS
<i>letrozole</i>	T2	
<i>leucovorin calcium oral tablet 10 mg, 15 mg, 25 mg</i>	T3	
<i>leucovorin calcium oral tablet 5 mg</i>	T2	
LEUKERAN	T5	
<i>leuprolide acetate (3 month)</i>	T4	QL (1 EA per 30 days)
<i>leuprolide subcutaneous kit</i>	T4	QL (2 EA per 28 days)
<i>lomustine oral capsule 10 mg, 40 mg</i>	T4	PA-NS
<i>lomustine oral capsule 100 mg</i>	T5	PA-NS
LONSURF	T5	PA-NS
LORBRENA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
LORBRENA ORAL TABLET 25 MG	T5	PA-NS; QL (93 EA per 31 days)
LUMAKRAS ORAL TABLET 120 MG, 240 MG	T5	PA-NS; QL (62 EA per 31 days)
LUMAKRAS ORAL TABLET 320 MG	T5	PA-NS; QL (93 EA per 31 days)
LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	T5	QL (1 EA per 90 days)
LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 3.75 MG	T5	QL (1 EA per 30 days)
LYNPARZA	T5	PA-NS; QL (124 EA per 31 days)
LYSODREN	T5	

Drug Name	Drug Tier	Requirements/Limits
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3)	T5	PA-NS; QL (93 EA per 31 days)
LYTGOBI ORAL TABLET 16 MG/DAY (4 MG X 4)	T5	PA-NS; QL (124 EA per 31 days)
LYTGOBI ORAL TABLET 20 MG/DAY (4 MG X 5)	T5	PA-NS; QL (155 EA per 31 days)
MATULANE	T5	
<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml)</i>	T2	PA
<i>megestrol oral suspension 625 mg/5 ml (125 mg/ml)</i>	T4	PA
<i>megestrol oral tablet</i>	T2	PA-NS
MEKINIST ORAL RECON SOLN	T5	PA-NS; QL (1260 ML per 31 days)
MEKINIST ORAL TABLET 0.5 MG	T5	PA-NS; QL (93 EA per 31 days)
MEKINIST ORAL TABLET 2 MG	T5	PA-NS; QL (31 EA per 31 days)
MEKTOVI	T5	PA-NS; QL (186 EA per 31 days)
<i>mercaptopurine oral suspension</i>	T5	
<i>mercaptopurine oral tablet</i>	T2	
<i>mesna oral</i>	T4	
<i>methotrexate sodium</i>	T2	PA-BvD
<i>methotrexate sodium (pf) injection solution</i>	T2	PA-BvD
MODEYSO	T5	PA-NS; QL (20 EA per 28 days)
<i>mycophenolate mofetil oral capsule</i>	T2	PA-BvD
<i>mycophenolate mofetil oral suspension for reconstitution</i>	T5	PA-BvD
<i>mycophenolate mofetil oral tablet</i>	T2	PA-BvD
<i>mycophenolate sodium</i>	T4	PA-BvD
NEMLUVIO	T5	PA; QL (2 EA per 28 days)
NERLYNX	T5	PA-NS; QL (186 EA per 31 days)
<i>nilotinib hcl</i>	T5	PA-NS; QL (124 EA per 31 days)
<i>nilutamide</i>	T5	
NINLARO	T5	PA-NS; QL (3 EA per 28 days)
NUBEQA	T5	PA-NS; QL (124 EA per 31 days)
<i>octreotide acetate injection solution</i>	T4	PA
ODOMZO	T5	PA-NS; LA; QL (31 EA per 31 days)
OGSIVEO ORAL TABLET 100 MG, 150 MG	T5	PA-NS; QL (62 EA per 31 days)
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	T5	PA-NS; QL (96 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4)	T5	PA-NS; QL (16 EA per 28 days)
OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)	T5	PA-NS; QL (20 EA per 28 days)
OJEMDA ORAL TABLET 600 MG/WEEK (100 MG X 6)	T5	PA-NS; QL (24 EA per 28 days)
OJJAARA	T5	PA-NS; QL (31 EA per 31 days)
ONUREG	T5	PA-NS; QL (14 EA per 28 days)
ORGOVYX	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 345 MG	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 86 MG	T5	PA-NS; QL (93 EA per 31 days)
<i>pazopanib oral tablet 200 mg</i>	T5	PA-NS; QL (124 EA per 31 days)
PEMAZYRE	T5	PA-NS; QL (14 EA per 21 days)
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (28 EA per 28 days)
PIQRAY ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	T5	PA-NS; QL (56 EA per 28 days)
<i>pomalidomide</i>	T5	PA-NS; QL (21 EA per 28 days)
PROGRAF ORAL GRANULES IN PACKET	T4	PA-BvD
QINLOCK	T5	PA-NS; QL (93 EA per 31 days)
RETEVMO ORAL TABLET 120 MG, 160 MG, 80 MG	T5	PA-NS; QL (62 EA per 31 days)
RETEVMO ORAL TABLET 40 MG	T5	PA-NS; QL (93 EA per 31 days)
REVUFORJ ORAL TABLET 110 MG	T5	PA-NS; QL (124 EA per 31 days)
REVUFORJ ORAL TABLET 160 MG	T5	PA-NS; QL (62 EA per 31 days)
REVUFORJ ORAL TABLET 25 MG	T5	PA-NS; QL (248 EA per 31 days)
REZLIDHIA	T5	PA-NS; QL (62 EA per 31 days)
ROMVIMZA	T5	PA-NS; QL (8 EA per 28 days)
ROZLYTREK ORAL CAPSULE 100 MG	T5	PA-NS; QL (155 EA per 31 days)
ROZLYTREK ORAL CAPSULE 200 MG	T5	PA-NS; QL (93 EA per 31 days)
ROZLYTREK ORAL PELLETS IN PACKET	T5	PA-NS; QL (372 EA per 31 days)
RUBRACA	T5	PA-NS; QL (124 EA per 31 days)
RYDAPT	T5	PA-NS; QL (248 EA per 31 days)
SCEMBLIX ORAL TABLET 100 MG	T5	PA-NS; QL (124 EA per 31 days)
SCEMBLIX ORAL TABLET 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SCEMBLIX ORAL TABLET 40 MG	T5	PA-NS; QL (248 EA per 31 days)
SIGNIFOR	T5	PA
<i>sirolimus</i>	T4	PA-BvD
SOLTAMOX	T4	

Drug Name	Drug Tier	Requirements/Limits
<i>sorafenib</i>	T5	PA-NS; QL (124 EA per 31 days)
STIVARGA	T5	PA-NS; QL (84 EA per 28 days)
<i>sunitinib malate</i>	T5	PA-NS; QL (31 EA per 31 days)
TABLOID	T4	
TABRECTA	T5	PA-NS; QL (124 EA per 31 days)
<i>tacrolimus oral capsule 0.5 mg, 1 mg</i>	T2	PA-BvD
<i>tacrolimus oral capsule 5 mg</i>	T4	PA-BvD
TAFINLAR ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
TAFINLAR ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (930 EA per 31 days)
TAGRISSE	T5	PA-NS; LA; QL (31 EA per 31 days)
TALZENNA	T5	PA-NS; QL (31 EA per 31 days)
<i>tamoxifen</i>	T1	
TEPMETKO	T5	PA-NS; QL (62 EA per 31 days)
THALOMID ORAL CAPSULE 100 MG	T5	PA-NS; QL (112 EA per 28 days)
THALOMID ORAL CAPSULE 50 MG	T5	PA-NS; QL (28 EA per 28 days)
TIBSOVO	T5	PA-NS; QL (62 EA per 31 days)
<i>toremifene</i>	T4	
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 11.25 MG	T4	ST-NS; QL (1 EA per 84 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 22.5 MG	T4	ST-NS; QL (1 EA per 168 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 3.75 MG	T4	ST-NS; QL (1 EA per 28 days)
<i>tretinoin (antineoplastic)</i>	T5	
TRUQAP	T5	PA-NS; QL (64 EA per 28 days)
TUKYSA ORAL TABLET 150 MG	T5	PA-NS; QL (124 EA per 31 days)
TUKYSA ORAL TABLET 50 MG	T5	PA-NS; QL (248 EA per 31 days)
TURALIO	T5	PA-NS; QL (124 EA per 31 days)
VANFLYTA	T5	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 10 MG	T3	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 100 MG	T5	PA-NS; QL (186 EA per 31 days)
VENCLEXTA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
VENCLEXTA STARTING PACK	T5	PA-NS; QL (84 EA per 365 days)
VERZENIO	T5	PA-NS; QL (62 EA per 31 days)
VIJOICE ORAL GRANULES IN PACKET	T5	PA-NS; QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
VIJOICE ORAL TABLET 125 MG, 50 MG	T5	PA-NS; QL (28 EA per 28 days)
VIJOICE ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
VITRAKVI ORAL CAPSULE 100 MG	T5	PA-NS; QL (62 EA per 31 days)
VITRAKVI ORAL CAPSULE 25 MG	T5	PA-NS; QL (186 EA per 31 days)
VITRAKVI ORAL SOLUTION	T5	PA-NS; QL (310 ML per 31 days)
VIZIMPRO	T5	PA-NS; QL (31 EA per 31 days)
VONJO	T5	PA-NS; QL (124 EA per 31 days)
VORANIGO ORAL TABLET 10 MG	T5	PA-NS; QL (62 EA per 31 days)
VORANIGO ORAL TABLET 40 MG	T5	PA-NS; QL (31 EA per 31 days)
WELIREG	T5	PA-NS; QL (93 EA per 31 days)
XALKORI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XALKORI ORAL PELLETT 150 MG	T5	PA-NS; QL (186 EA per 31 days)
XALKORI ORAL PELLETT 20 MG, 50 MG	T5	PA-NS; QL (124 EA per 31 days)
XATMEP	T4	PA-BvD
XERMELO	T5	PA; QL (93 EA per 31 days)
XGEVA	T5	PA-NS
XOSPATA	T5	PA-NS; QL (124 EA per 31 days)
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40MG TWICE WEEK (40 MG X 2), 80 MG/WEEK (40 MG X 2)	T5	PA-NS; QL (8 EA per 28 days)
XPOVIO ORAL TABLET 40 MG/WEEK (10 MG X 4)	T5	PA-NS; QL (16 EA per 28 days)
XPOVIO ORAL TABLET 60 MG/WEEK (60 MG X 1), 80 MG/WEEK (80 MG X 1)	T5	PA-NS; QL (4 EA per 28 days)
XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK)	T5	PA-NS; QL (24 EA per 28 days)
XPOVIO ORAL TABLET 80MG TWICE WEEK (160 MG/WEEK)	T5	PA-NS; QL (32 EA per 28 days)
XTANDI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 40 MG	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 80 MG	T5	PA-NS; QL (62 EA per 31 days)
YONSA	T5	PA-NS; QL (124 EA per 31 days)
ZEJULA ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
ZELBORAF	T5	PA-NS; QL (248 EA per 31 days)
ZOLINZA	T5	PA-NS
ZYKADIA	T5	PA-NS; QL (93 EA per 31 days)
Autonomic / Cns Drugs, Neurology / Psych		
ABILIFY MAINTENA	T5	QL (1 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i>	T2	PA; QL (5167 ML per 31 days)
<i>acetaminophen-codeine oral tablet</i>	T2	PA; QL (403 EA per 31 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML	T3	PA; QL (1 ML per 28 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 70 MG/ML	T3	PA; QL (2 ML per 28 days)
<i>alprazolam oral tablet 0.25 mg, 0.5 mg</i>	T2	PA; QL (93 EA per 31 days)
<i>alprazolam oral tablet 1 mg, 2 mg</i>	T2	PA; QL (155 EA per 31 days)
<i>amitriptyline</i>	T2	PA-NS
<i>amoxapine</i>	T2	
<i>aripiprazole oral solution</i>	T4	PA-NS
<i>aripiprazole oral tablet</i>	T3	
<i>aripiprazole oral tablet, disintegrating 10 mg</i>	T4	PA-NS
<i>aripiprazole oral tablet, disintegrating 15 mg</i>	T5	PA-NS
<i>armodafinil</i>	T4	PA; QL (31 EA per 31 days)
<i>asenapine maleate</i>	T4	PA-NS; QL (62 EA per 31 days)
<i>atomoxetine oral capsule 10 mg, 25 mg, 40 mg</i>	T3	QL (62 EA per 31 days)
<i>atomoxetine oral capsule 100 mg, 60 mg, 80 mg</i>	T3	QL (31 EA per 31 days)
<i>atomoxetine oral capsule 18 mg</i>	T3	QL (124 EA per 31 days)
AUSTEDO ORAL TABLET 12 MG, 6 MG	T5	PA; QL (124 EA per 31 days)
AUSTEDO ORAL TABLET 9 MG	T5	PA; QL (155 EA per 31 days)
AUSTEDO XR	T5	PA; QL (31 EA per 31 days)
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG	T5	PA; QL (56 EA per 365 days)
AUVELITY	T5	PA-NS; QL (62 EA per 31 days)
<i>baclofen oral tablet 10 mg, 20 mg</i>	T2	
BAFIERTAM	T5	PA; QL (124 EA per 31 days)
<i>benztropine oral</i>	T2	PA
<i>brivaracetam oral solution</i>	T5	QL (620 ML per 31 days)
<i>brivaracetam oral tablet</i>	T5	QL (62 EA per 31 days)
<i>bromocriptine</i>	T4	
<i>buprenorphine</i>	T4	PA; QL (4 EA per 28 days)
<i>buprenorphine hcl sublingual</i>	T3	
<i>buprenorphine-naloxone sublingual film</i>	T2	
<i>buprenorphine-naloxone sublingual tablet</i>	T4	
<i>bupropion hcl oral tablet</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>bupropion hcl oral tablet extended release 24 hr 150 mg</i>	T2	QL (93 EA per 31 days)
<i>bupropion hcl oral tablet extended release 24 hr 300 mg</i>	T2	QL (31 EA per 31 days)
<i>bupropion hcl oral tablet sustained-release 12 hr</i>	T2	QL (62 EA per 31 days)
<i>bupirone</i>	T2	
<i>butorphanol nasal</i>	T4	QL (5 ML per 28 days)
CAPLYTA	T5	PA-NS; QL (31 EA per 31 days)
<i>carbamazepine oral capsule, er multiphase 12 hr</i>	T3	
<i>carbamazepine oral suspension 100 mg/5 ml</i>	T2	
<i>carbamazepine oral tablet</i>	T2	
<i>carbamazepine oral tablet extended release 12 hr</i>	T3	
<i>carbamazepine oral tablet, chewable 100 mg</i>	T2	
<i>carbidopa-levodopa oral tablet</i>	T2	
<i>carbidopa-levodopa oral tablet extended release</i>	T2	
<i>carbidopa-levodopa oral tablet, disintegrating</i>	T2	
<i>carbidopa-levodopa-entacapone</i>	T4	
<i>celecoxib oral capsule 100 mg, 200 mg, 50 mg</i>	T2	QL (62 EA per 31 days)
<i>celecoxib oral capsule 400 mg</i>	T3	QL (62 EA per 31 days)
<i>chlorpromazine oral</i>	T4	
<i>citalopram oral solution</i>	T3	
<i>citalopram oral tablet</i>	T1	
<i>clobazam oral suspension</i>	T4	PA-NS; QL (496 ML per 31 days)
<i>clobazam oral tablet</i>	T3	PA-NS; QL (62 EA per 31 days)
<i>clomipramine</i>	T4	PA-NS
<i>clonazepam oral tablet 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>clonazepam oral tablet 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet, disintegrating 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>clonazepam oral tablet, disintegrating 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clorazepate dipotassium oral tablet 15 mg</i>	T3	PA-NS; QL (186 EA per 31 days)
<i>clorazepate dipotassium oral tablet 3.75 mg, 7.5 mg</i>	T3	PA-NS; QL (93 EA per 31 days)
<i>clozapine oral tablet 100 mg</i>	T3	QL (279 EA per 31 days)
<i>clozapine oral tablet 200 mg</i>	T3	QL (124 EA per 31 days)
<i>clozapine oral tablet 25 mg</i>	T2	QL (279 EA per 31 days)
<i>clozapine oral tablet 50 mg</i>	T2	QL (93 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>clozapine oral tablet, disintegrating 100 mg, 25 mg</i>	T4	QL (279 EA per 31 days)
<i>clozapine oral tablet, disintegrating 12.5 mg</i>	T4	QL (93 EA per 31 days)
<i>clozapine oral tablet, disintegrating 150 mg</i>	T4	QL (186 EA per 31 days)
<i>clozapine oral tablet, disintegrating 200 mg</i>	T4	QL (124 EA per 31 days)
COBENFY	T5	PA-NS; QL (62 EA per 31 days)
COBENFY STARTER PACK	T5	PA-NS; QL (112 EA per 365 days)
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
COPAXONE SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
<i>cyclobenzaprine oral tablet 10 mg</i>	T2	QL (93 EA per 31 days)
<i>cyclobenzaprine oral tablet 5 mg</i>	T2	QL (155 EA per 31 days)
<i>dalfampridine</i>	T5	PA; QL (62 EA per 31 days)
<i>dantrolene oral</i>	T4	
<i>desipramine oral tablet 10 mg, 25 mg</i>	T3	
<i>desipramine oral tablet 100 mg, 150 mg, 50 mg, 75 mg</i>	T4	
<i>desvenlafaxine succinate</i>	T3	QL (31 EA per 31 days)
<i>dexmethylphenidate oral capsule, er biphasic 50-50</i>	T4	QL (31 EA per 31 days)
<i>dexmethylphenidate oral tablet 10 mg</i>	T2	QL (62 EA per 31 days)
<i>dexmethylphenidate oral tablet 2.5 mg, 5 mg</i>	T2	QL (93 EA per 31 days)
<i>dextroamphetamine-amphetamine oral capsule, extended release 24hr</i>	T3	QL (31 EA per 31 days)
<i>dextroamphetamine-amphetamine oral tablet 10 mg, 15 mg, 30 mg, 5 mg, 7.5 mg</i>	T1	QL (62 EA per 31 days)
<i>dextroamphetamine-amphetamine oral tablet 20 mg</i>	T1	QL (93 EA per 31 days)
DIACOMIT ORAL CAPSULE 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL CAPSULE 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIAZEPAM INTENSOL	T2	PA-NS; QL (248 ML per 31 days)
<i>diazepam oral solution 5 mg/5 ml (1 mg/ml)</i>	T2	PA-NS; QL (1500 ML per 31 days)
<i>diazepam oral tablet</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>diazepam rectal</i>	T4	
<i>diclofenac potassium oral tablet 50 mg</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>diclofenac sodium oral</i>	T2	
<i>diclofenac sodium topical drops</i>	T2	QL (450 ML per 28 days)
<i>diflunisal</i>	T2	
<i>dihydroergotamine nasal</i>	T5	PA; QL (8 ML per 28 days)
DILANTIN	T3	
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg (14)- 240 mg (46)</i>	T5	PA; QL (120 EA per 365 days)
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 240 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>divalproex oral capsule, delayed rel sprinkle</i>	T2	
<i>divalproex oral tablet extended release 24 hr 250 mg</i>	T2	
<i>divalproex oral tablet extended release 24 hr 500 mg</i>	T3	
<i>divalproex oral tablet, delayed release (dr/ec)</i>	T2	
<i>donepezil oral tablet 10 mg, 5 mg</i>	T1	
<i>donepezil oral tablet 23 mg</i>	T3	QL (31 EA per 31 days)
<i>donepezil oral tablet, disintegrating</i>	T2	
<i>doxepin oral capsule</i>	T2	PA-NS
<i>doxepin oral concentrate</i>	T2	PA-NS
<i>doxepin oral tablet</i>	T3	PA
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG	T3	PA-NS; QL (93 EA per 31 days)
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 30 MG, 60 MG	T3	PA-NS; QL (62 EA per 31 days)
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 40 MG	T3	PA-NS; QL (31 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 20 mg, 60 mg</i>	T2	QL (62 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 30 mg</i>	T2	QL (31 EA per 31 days)
EMGALITY PEN	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 300 MG/3 ML (100 MG/ML X 3)	T5	PA; QL (3 ML per 28 days)
EMSAM	T5	QL (30 EA per 30 days)
<i>entacapone</i>	T3	
EPIDIOLEX	T5	PA-NS

Drug Name	Drug Tier	Requirements/Limits
<i>ergotamine-caffeine</i>	T3	PA
<i>escitalopram oxalate oral capsule</i>	T4	PA-NS; QL (31 EA per 31 days)
<i>escitalopram oxalate oral solution</i>	T4	QL (620 ML per 31 days)
<i>escitalopram oxalate oral tablet 10 mg</i>	T2	QL (45 EA per 30 days)
<i>escitalopram oxalate oral tablet 20 mg, 5 mg</i>	T2	QL (30 EA per 30 days)
<i>eslicarbazepine oral tablet 200 mg</i>	T4	QL (186 EA per 31 days)
<i>eslicarbazepine oral tablet 400 mg</i>	T5	QL (93 EA per 31 days)
<i>eslicarbazepine oral tablet 600 mg, 800 mg</i>	T5	QL (62 EA per 31 days)
<i>ethosuximide oral capsule</i>	T3	
<i>ethosuximide oral solution</i>	T4	
<i>etodolac oral capsule</i>	T3	
<i>etodolac oral tablet</i>	T2	
<i>etodolac oral tablet extended release 24 hr 400 mg, 500 mg</i>	T3	
<i>etodolac oral tablet extended release 24 hr 600 mg</i>	T4	
EXXUA ORAL TABLET EXTENDED RELEASE 24 HR	T4	PA-NS; QL (31 EA per 31 days)
EXXUA ORAL TABLET, EXT REL 24HR DOSE PACK	T4	PA-NS; QL (64 EA per 365 days)
FANAPT ORAL TABLET 1 MG	T4	PA-NS; QL (62 EA per 31 days)
FANAPT ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG	T5	PA-NS; QL (62 EA per 31 days)
FANAPT TITRATION PACK A	T4	PA-NS; QL (16 EA per 365 days)
<i>felbamate</i>	T4	
<i>fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 75 mcg/hr</i>	T4	PA; QL (10 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 25 mcg/hr, 50 mcg/hr</i>	T2	PA; QL (10 EA per 30 days)
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)	T4	PA-NS; QL (56 EA per 365 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 40 MG, 80 MG	T4	PA-NS; QL (31 EA per 31 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 20 MG	T4	PA-NS; QL (93 EA per 31 days)
<i>fingolimod</i>	T5	PA; QL (31 EA per 31 days)
FINTEPLA	T5	PA-NS; QL (360 ML per 30 days)
<i>fluoxetine (pmd)</i>	T2	
<i>fluoxetine oral capsule</i>	T1	
<i>fluoxetine oral solution</i>	T3	
<i>fluoxetine oral tablet 10 mg, 20 mg</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>fluphenazine decanoate</i>	T3	
<i>fluphenazine hcl injection</i>	T4	
<i>fluphenazine hcl oral concentrate</i>	T4	
<i>fluphenazine hcl oral elixir</i>	T4	
<i>fluphenazine hcl oral tablet 1 mg, 2.5 mg, 5 mg</i>	T2	
<i>fluphenazine hcl oral tablet 10 mg</i>	T4	
<i>flurbiprofen oral tablet 100 mg</i>	T2	
<i>fluvoxamine oral tablet</i>	T2	
<i>gabapentin oral capsule 100 mg, 400 mg</i>	T2	QL (270 EA per 30 days)
<i>gabapentin oral capsule 300 mg</i>	T2	QL (360 EA per 30 days)
<i>gabapentin oral solution 250 mg/5 ml</i>	T4	QL (2160 ML per 30 days)
<i>gabapentin oral tablet 600 mg</i>	T2	QL (180 EA per 30 days)
<i>gabapentin oral tablet 800 mg</i>	T2	QL (120 EA per 30 days)
<i>galantamine oral capsule, ext rel. pellets 24 hr</i>	T3	
<i>galantamine oral solution</i>	T4	
<i>galantamine oral tablet</i>	T3	
<i>glatiramer subcutaneous syringe 20 mg/ml</i>	T5	PA; QL (31 ML per 31 days)
<i>glatiramer subcutaneous syringe 40 mg/ml</i>	T5	PA; QL (12 ML per 28 days)
GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
GLATOPA SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
<i>guanfacine oral tablet extended release 24 hr</i>	T2	PA
<i>haloperidol</i>	T2	
<i>haloperidol decanoate</i>	T2	
<i>haloperidol lactate injection</i>	T1	
<i>haloperidol lactate oral</i>	T2	
<i>hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i>	T2	PA; QL (372 EA per 31 days)
<i>hydromorphone oral liquid</i>	T4	PA; QL (1240 ML per 31 days)
<i>hydromorphone oral tablet 2 mg, 4 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>hydromorphone oral tablet 8 mg</i>	T3	PA; QL (155 EA per 31 days)
IBU ORAL TABLET 600 MG, 800 MG	T1	
<i>ibuprofen oral tablet 400 mg, 600 mg, 800 mg</i>	T1	
<i>imipramine hcl</i>	T2	PA-NS
<i>indomethacin oral capsule</i>	T2	
<i>indomethacin oral capsule, extended release</i>	T2	
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,092 MG/3.5 ML	T5	QL (3.5 ML per 180 days)

Drug Name	Drug Tier	Requirements/Limits
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,560 MG/5 ML	T5	QL (5 ML per 180 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 117 MG/0.75 ML	T5	QL (0.75 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 156 MG/ML	T5	QL (1 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 234 MG/1.5 ML	T5	QL (1.5 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 39 MG/0.25 ML	T4	QL (0.25 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 78 MG/0.5 ML	T5	QL (0.5 ML per 28 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 273 MG/0.88 ML	T5	QL (0.88 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 410 MG/1.32 ML	T5	QL (1.32 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 546 MG/1.75 ML	T5	QL (1.75 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 819 MG/2.63 ML	T5	QL (2.63 ML per 84 days)
KESIMPTA PEN	T5	PA; QL (0.4 ML per 28 days)
KLOXXADO	T3	
<i>lacosamide oral</i>	T4	
<i>lamotrigine oral tablet</i>	T2	
<i>lamotrigine oral tablet extended release 24hr</i>	T4	
<i>lamotrigine oral tablet, chewable dispersible</i>	T2	
<i>levetiracetam oral solution 100 mg/ml</i>	T2	
<i>levetiracetam oral tablet</i>	T2	
<i>levetiracetam oral tablet extended release 24 hr</i>	T2	
<i>lithium carbonate oral capsule</i>	T1	
<i>lithium carbonate oral tablet</i>	T1	
<i>lithium carbonate oral tablet extended release</i>	T2	
<i>lithium citrate</i>	T2	
<i>lofexidine</i>	T5	
LORAZEPAM INTENSOL	T2	PA; QL (155 ML per 31 days)
<i>lorazepam oral tablet 0.5 mg</i>	T2	PA; QL (124 EA per 31 days)
<i>lorazepam oral tablet 1 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>lorazepam oral tablet 2 mg</i>	T2	PA; QL (155 EA per 31 days)
<i>loxapine succinate</i>	T2	
<i>lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg</i>	T4	PA-NS; QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>lurasidone oral tablet 80 mg</i>	T4	PA-NS; QL (62 EA per 31 days)
MARPLAN	T4	
<i>meloxicam oral tablet</i>	T1	
<i>memantine oral capsule, sprinkle, er 24hr</i>	T4	
<i>memantine oral solution</i>	T4	
<i>memantine oral tablet</i>	T2	
<i>memantine-donepezil</i>	T4	PA
<i>methadone oral solution 10 mg/5 ml</i>	T2	PA; QL (620 ML per 31 days)
<i>methadone oral solution 5 mg/5 ml</i>	T2	PA; QL (1240 ML per 31 days)
<i>methadone oral tablet 10 mg</i>	T2	PA; QL (124 EA per 31 days)
<i>methadone oral tablet 5 mg</i>	T2	PA; QL (248 EA per 31 days)
<i>methsuximide</i>	T4	
<i>methylphenidate hcl oral tablet</i>	T2	QL (93 EA per 31 days)
<i>mirtazapine oral tablet 15 mg, 30 mg, 45 mg</i>	T2	
<i>mirtazapine oral tablet 7.5 mg</i>	T3	
<i>mirtazapine oral tablet, disintegrating</i>	T3	
<i>modafinil</i>	T3	PA; QL (31 EA per 31 days)
<i>molindone</i>	T4	
<i>morphine concentrate oral solution</i>	T2	PA; QL (310 ML per 31 days)
<i>morphine oral solution 10 mg/5 ml</i>	T2	PA; QL (2800 ML per 31 days)
<i>morphine oral solution 20 mg/5 ml (4 mg/ml)</i>	T2	PA; QL (1400 ML per 31 days)
<i>morphine oral tablet</i>	T2	PA; QL (186 EA per 31 days)
<i>morphine oral tablet extended release 100 mg</i>	T3	PA; QL (62 EA per 31 days)
<i>morphine oral tablet extended release 15 mg, 30 mg, 60 mg</i>	T3	PA; QL (100 EA per 31 days)
<i>morphine oral tablet extended release 200 mg</i>	T3	PA; QL (31 EA per 31 days)
<i>nabumetone</i>	T2	
<i>naloxone injection solution</i>	T2	
<i>naloxone injection syringe</i>	T2	
<i>naltrexone</i>	T2	
<i>naproxen oral tablet</i>	T1	
<i>naproxen oral tablet, delayed release (dr/ec) 375 mg</i>	T2	
<i>naproxen oral tablet, delayed release (dr/ec) 500 mg</i>	T4	
<i>naproxen sodium oral tablet 550 mg</i>	T2	
<i>naratriptan oral tablet 1 mg</i>	T3	QL (20 EA per 28 days)
<i>naratriptan oral tablet 2.5 mg</i>	T3	QL (9 EA per 28 days)
NAYZILAM	T4	PA-NS; QL (10 EA per 30 days)

Drug Name	Drug Tier	Requirements/Limits
<i>nefazodone</i>	T4	
<i>nortriptyline</i>	T2	
NUEDEXTA	T5	PA; QL (62 EA per 31 days)
NUPLAZID	T5	PA-NS; QL (31 EA per 31 days)
NURTEC ODT	T5	PA; QL (18 EA per 28 days)
<i>olanzapine intramuscular</i>	T4	
<i>olanzapine oral tablet</i>	T2	QL (31 EA per 31 days)
<i>olanzapine oral tablet,disintegrating</i>	T4	QL (31 EA per 31 days)
OPIPZA	T5	PA-NS
<i>oxcarbazepine oral suspension</i>	T4	
<i>oxcarbazepine oral tablet</i>	T2	
<i>oxycodone oral capsule</i>	T3	PA; QL (186 EA per 31 days)
<i>oxycodone oral concentrate</i>	T4	PA; QL (180 ML per 31 days)
<i>oxycodone oral tablet 10 mg, 15 mg, 20 mg, 5 mg</i>	T3	PA; QL (186 EA per 31 days)
<i>oxycodone oral tablet 30 mg</i>	T3	PA; QL (138 EA per 31 days)
<i>oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg</i>	T3	PA; QL (372 EA per 31 days)
<i>oxycodone-acetaminophen oral tablet 5-325 mg, 7.5-325 mg</i>	T2	PA; QL (372 EA per 31 days)
<i>paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 9 mg</i>	T4	QL (31 EA per 31 days)
<i>paliperidone oral tablet extended release 24hr 6 mg</i>	T4	QL (62 EA per 31 days)
<i>paroxetine hcl oral suspension</i>	T4	PA-NS
<i>paroxetine hcl oral tablet</i>	T1	PA-NS
<i>perampanel oral suspension</i>	T5	QL (744 ML per 31 days)
<i>perampanel oral tablet 10 mg, 12 mg, 4 mg, 6 mg, 8 mg</i>	T5	QL (31 EA per 31 days)
<i>perampanel oral tablet 2 mg</i>	T4	QL (31 EA per 31 days)
<i>perphenazine</i>	T4	
PERSERIS	T5	QL (1 EA per 28 days)
<i>phenelzine</i>	T2	
<i>phenobarbital</i>	T2	PA-NS
PHENYTEK	T3	
<i>phenytoin oral suspension 125 mg/5 ml</i>	T2	
<i>phenytoin oral tablet,chewable</i>	T2	
<i>phenytoin sodium extended oral capsule 100 mg</i>	T2	
<i>pimozide</i>	T4	
<i>piroxicam</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>pramipexole oral tablet</i>	T2	
<i>pregabalin oral capsule 100 mg, 150 mg, 200 mg, 25 mg, 50 mg, 75 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>pregabalin oral capsule 225 mg, 300 mg</i>	T2	PA-NS; QL (62 EA per 31 days)
<i>pregabalin oral solution</i>	T2	PA-NS; QL (930 ML per 31 days)
<i>primidone oral tablet 125 mg</i>	T3	
<i>primidone oral tablet 250 mg, 50 mg</i>	T2	
<i>protriptyline</i>	T4	
<i>pyridostigmine bromide oral tablet 60 mg</i>	T3	
<i>pyridostigmine bromide oral tablet extended release 180 mg</i>	T4	
<i>quetiapine oral tablet 100 mg, 200 mg, 300 mg, 400 mg, 50 mg</i>	T2	QL (62 EA per 31 days)
<i>quetiapine oral tablet 150 mg</i>	T3	QL (62 EA per 31 days)
<i>quetiapine oral tablet 25 mg</i>	T1	QL (62 EA per 31 days)
<i>quetiapine oral tablet extended release 24 hr</i>	T3	QL (62 EA per 31 days)
QULIPTA	T3	PA; QL (31 EA per 31 days)
RALDESY	T5	
<i>ramelteon</i>	T4	QL (31 EA per 31 days)
<i>rasagiline</i>	T4	
REXULTI ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 12.5 mg/2 ml</i>	T4	QL (2 EA per 28 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 25 mg/2 ml, 37.5 mg/2 ml, 50 mg/2 ml</i>	T5	QL (2 EA per 28 days)
<i>risperidone oral solution</i>	T2	QL (496 ML per 31 days)
<i>risperidone oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T2	QL (31 EA per 31 days)
<i>risperidone oral tablet 3 mg</i>	T2	QL (93 EA per 31 days)
<i>risperidone oral tablet 4 mg</i>	T2	QL (124 EA per 31 days)
<i>risperidone oral tablet,disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T4	QL (31 EA per 31 days)
<i>risperidone oral tablet,disintegrating 3 mg</i>	T4	QL (93 EA per 31 days)
<i>risperidone oral tablet,disintegrating 4 mg</i>	T4	QL (124 EA per 31 days)
<i>rivastigmine</i>	T4	QL (30 EA per 30 days)
<i>rivastigmine tartrate</i>	T3	
<i>rizatriptan oral tablet 10 mg</i>	T2	QL (12 EA per 28 days)
<i>rizatriptan oral tablet 5 mg</i>	T2	QL (24 EA per 28 days)
<i>rizatriptan oral tablet,disintegrating 10 mg</i>	T2	QL (12 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
<i>rizatriptan oral tablet,disintegrating 5 mg</i>	T2	QL (24 EA per 28 days)
<i>ropinirole oral tablet</i>	T2	
ROWEEPRA	T2	
<i>rufinamide oral suspension</i>	T5	PA-NS
<i>rufinamide oral tablet 200 mg</i>	T4	PA-NS
<i>rufinamide oral tablet 400 mg</i>	T5	PA-NS
RYTARY	T3	ST
SECUADO	T5	PA-NS; QL (31 EA per 31 days)
<i>selegiline hcl</i>	T2	
<i>sertraline oral concentrate</i>	T2	
<i>sertraline oral tablet</i>	T1	
SKYCLARYS	T5	PA; QL (93 EA per 31 days)
<i>sodium oxybate</i>	T5	PA; QL (540 ML per 30 days)
SPRITAM ORAL TABLET FOR SUSPENSION 250 MG, 500 MG	T4	
SUBVENITE ORAL TABLET	T2	
<i>sulindac</i>	T2	
<i>sumatriptan nasal spray,non-aerosol 20 mg/actuation</i>	T4	QL (8 EA per 28 days)
<i>sumatriptan nasal spray,non-aerosol 5 mg/actuation</i>	T4	QL (32 EA per 28 days)
<i>sumatriptan succinate oral tablet 100 mg</i>	T2	QL (9 EA per 28 days)
<i>sumatriptan succinate oral tablet 25 mg</i>	T2	QL (36 EA per 28 days)
<i>sumatriptan succinate oral tablet 50 mg</i>	T2	QL (18 EA per 28 days)
<i>sumatriptan succinate subcutaneous pen injector 6 mg/0.5 ml</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan succinate subcutaneous solution</i>	T4	QL (4 ML per 28 days)
SYMPAZAN ORAL FILM 10 MG, 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SYMPAZAN ORAL FILM 5 MG	T4	PA-NS; QL (62 EA per 31 days)
TASCENSO ODT	T5	PA; QL (31 EA per 31 days)
<i>tasimelteon</i>	T5	PA; QL (31 EA per 31 days)
<i>teriflunomide</i>	T5	PA; QL (31 EA per 31 days)
<i>tetrabenazine oral tablet 12.5 mg</i>	T5	PA; QL (93 EA per 31 days)
<i>tetrabenazine oral tablet 25 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>thioridazine</i>	T2	
<i>thiothixene oral capsule 1 mg, 2 mg</i>	T2	
<i>thiothixene oral capsule 10 mg, 5 mg</i>	T3	
<i>tiagabine</i>	T4	
<i>tizanidine oral tablet</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>topiramate oral capsule, sprinkle 15 mg, 25 mg</i>	T2	
<i>topiramate oral solution</i>	T4	PA-NS; QL (496 ML per 31 days)
<i>topiramate oral tablet</i>	T2	
<i>tramadol oral tablet 50 mg</i>	T2	PA; QL (240 EA per 30 days)
<i>tramadol-acetaminophen</i>	T2	PA; QL (372 EA per 31 days)
<i>tranylcypromine</i>	T4	
<i>trazodone oral tablet 100 mg, 150 mg, 50 mg</i>	T1	
<i>trifluoperazine</i>	T2	
<i>trimipramine oral capsule 100 mg</i>	T4	PA-NS
<i>trimipramine oral capsule 25 mg, 50 mg</i>	T3	PA-NS
TRINTELLIX	T4	PA-NS
UBRELVY ORAL TABLET 100 MG	T5	PA; QL (17 EA per 28 days)
UBRELVY ORAL TABLET 50 MG	T5	PA; QL (34 EA per 28 days)
<i>valproic acid</i>	T2	
<i>valproic acid (as sodium salt) oral solution 250 mg/5 ml</i>	T2	
VALTOCO	T5	PA-NS; QL (10 EA per 30 days)
<i>venlafaxine oral capsule,extended release 24hr 150 mg, 37.5 mg</i>	T2	QL (31 EA per 31 days)
<i>venlafaxine oral capsule,extended release 24hr 75 mg</i>	T2	QL (93 EA per 31 days)
<i>venlafaxine oral tablet</i>	T2	
VERSACLOZ	T5	QL (558 ML per 31 days)
<i>vigabatrin</i>	T5	PA-NS
VIGADRONE	T5	PA-NS
<i>vilazodone</i>	T3	PA-NS; QL (31 EA per 31 days)
VIVITROL	T5	
VRAYLAR ORAL CAPSULE	T5	PA-NS; QL (31 EA per 31 days)
VUMERITY	T5	PA; QL (124 EA per 31 days)
XCOPRI	T5	PA-NS
XCOPRI MAINTENANCE PACK	T5	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 12.5 MG (14)- 25 MG (14)	T4	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 150 MG (14)- 200 MG (14), 50 MG (14)- 100 MG (14)	T5	PA-NS
<i>zaleplon oral capsule 10 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>zaleplon oral capsule 5 mg</i>	T4	PA; QL (93 EA per 31 days)
ZAVZPRET	T5	PA; QL (8 EA per 30 days)

Drug Name	Drug Tier	Requirements/Limits
ZEPOSIA	T5	PA; QL (31 EA per 31 days)
ZEPOSIA STARTER KIT (28-DAY)	T5	PA; QL (56 EA per 365 days)
ZEPOSIA STARTER PACK (7-DAY)	T5	PA; QL (14 EA per 365 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML	T5	PA; QL (11.648 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 23 MG/0.574 ML	T5	PA; QL (16.072 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 32.4 MG/0.81 ML	T5	PA; QL (22.68 ML per 28 days)
<i>ziprasidone hcl</i>	T3	QL (62 EA per 31 days)
<i>ziprasidone mesylate</i>	T4	
<i>zolpidem oral tablet</i>	T2	PA; QL (31 EA per 31 days)
ZONISADE	T5	PA-NS; QL (930 ML per 31 days)
<i>zonisamide</i>	T2	
ZTALMY	T5	PA-NS; QL (1100 ML per 30 days)
ZUBSOLV	T3	
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG	T5	PA-NS; QL (28 EA per 180 days)
ZURZUVAE ORAL CAPSULE 30 MG	T5	PA-NS; QL (14 EA per 180 days)
Cardiovascular, Hypertension / Lipids		
<i>acebutolol</i>	T2	
<i>aliskiren</i>	T4	
<i>amiloride</i>	T2	
<i>amiloride-hydrochlorothiazide</i>	T2	
<i>amiodarone oral</i>	T2	
<i>amlodipine</i>	T1	
<i>amlodipine-benazepril</i>	T1	
<i>amlodipine-olmesartan</i>	T2	QL (31 EA per 31 days)
<i>amlodipine-valsartan</i>	T2	
<i>aspirin-dipyridamole</i>	T4	
<i>atenolol</i>	T1	
<i>atenolol-chlorthalidone</i>	T2	
<i>atorvastatin</i>	T1	
ATTRUBY	T5	PA; QL (124 EA per 31 days)
<i>benazepril</i>	T1	
<i>benazepril-hydrochlorothiazide</i>	T1	
<i>bisoprolol fumarate oral tablet 10 mg, 5 mg</i>	T2	
<i>bisoprolol-hydrochlorothiazide</i>	T1	

Drug Name	Drug Tier	Requirements/Limits
<i>bumetanide oral</i>	T2	
<i>candesartan</i>	T2	
<i>candesartan-hydrochlorothiazid</i>	T2	
<i>captopril</i>	T1	
CARTIA XT	T2	
<i>carvedilol</i>	T1	
<i>chlorthalidone oral tablet 25 mg, 50 mg</i>	T1	
<i>cholestyramine (with sugar) oral powder in packet</i>	T2	
CHOLESTYRAMINE LIGHT ORAL POWDER IN PACKET	T2	
<i>cilostazol</i>	T2	
<i>clonidine</i>	T4	
<i>clonidine hcl oral tablet 0.1 mg, 0.2 mg, 0.3 mg</i>	T1	
<i>clopidogrel oral tablet 75 mg</i>	T1	
<i>colesevelam</i>	T4	
<i>colestipol oral packet</i>	T4	
<i>colestipol oral tablet</i>	T3	
<i>dabigatran etexilate oral capsule 110 mg</i>	T4	QL (124 EA per 31 days)
<i>dabigatran etexilate oral capsule 150 mg, 75 mg</i>	T4	QL (62 EA per 31 days)
<i>digoxin oral solution</i>	T3	QL (155 ML per 31 days)
<i>digoxin oral tablet 125 mcg (0.125 mg)</i>	T1	QL (62 EA per 31 days)
<i>digoxin oral tablet 250 mcg (0.25 mg)</i>	T2	QL (31 EA per 31 days)
<i>diltiazem hcl oral capsule,extended release 12 hr</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24 hr 360 mg, 420 mg</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24hr 120 mg, 180 mg, 240 mg, 300 mg</i>	T2	
<i>diltiazem hcl oral tablet</i>	T2	
<i>diltiazem hcl oral tablet extended release 24 hr</i>	T2	
DILT-XR	T2	
<i>dofetilide</i>	T4	
<i>doxazosin</i>	T2	
ELIQUIS DVT-PE TREAT 30D START	T3	QL (74 EA per 30 days)
ELIQUIS ORAL TABLET 2.5 MG	T3	QL (60 EA per 30 days)
ELIQUIS ORAL TABLET 5 MG	T3	QL (74 EA per 30 days)
<i>eltrombopag olamine oral powder in packet 12.5 mg</i>	T5	PA; QL (372 EA per 31 days)
<i>eltrombopag olamine oral powder in packet 25 mg</i>	T5	PA; QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>eltrombopag olamine oral tablet 12.5 mg, 25 mg</i>	T5	PA; QL (31 EA per 31 days)
<i>eltrombopag olamine oral tablet 50 mg, 75 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>enalapril maleate oral tablet</i>	T1	
<i>enalapril-hydrochlorothiazide</i>	T1	
<i>enoxaparin subcutaneous syringe</i>	T4	
ENTRESTO ORAL TABLET 24-26 MG	T4	ST; QL (186 EA per 31 days)
ENTRESTO ORAL TABLET 49-51 MG	T4	ST; QL (93 EA per 31 days)
ENTRESTO ORAL TABLET 97-103 MG	T4	ST; QL (62 EA per 31 days)
ENTRESTO SPRINKLE	T3	PA; QL (248 EA per 31 days)
<i>eplerenone</i>	T4	
<i>ethacrynic acid</i>	T4	
<i>ezetimibe</i>	T2	
<i>ezetimibe-simvastatin</i>	T3	QL (31 EA per 31 days)
<i>felodipine</i>	T2	
<i>fenofibrate micronized oral capsule 134 mg, 200 mg, 67 mg</i>	T2	
<i>fenofibrate nanocrystallized</i>	T2	
<i>fenofibrate oral tablet 160 mg, 54 mg</i>	T2	
<i>flecainide</i>	T2	
<i>fondaparinux subcutaneous syringe 10 mg/0.8 ml, 5 mg/0.4 ml, 7.5 mg/0.6 ml</i>	T5	
<i>fondaparinux subcutaneous syringe 2.5 mg/0.5 ml</i>	T4	
<i>fosinopril</i>	T1	
<i>fosinopril-hydrochlorothiazide</i>	T1	
<i>furosemide injection</i>	T2	
<i>furosemide oral solution</i>	T2	
<i>furosemide oral tablet</i>	T1	
<i>gemfibrozil</i>	T1	
<i>heparin (porcine) injection solution</i>	T3	PA-BvD
<i>hydralazine oral</i>	T1	
<i>hydrochlorothiazide</i>	T1	
<i>icosapent ethyl oral capsule 0.5 gram</i>	T4	QL (248 EA per 31 days)
<i>icosapent ethyl oral capsule 1 gram</i>	T4	QL (124 EA per 31 days)
<i>indapamide</i>	T1	
<i>irbesartan</i>	T1	QL (31 EA per 31 days)
<i>irbesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>isosorbide dinitrate oral tablet 10 mg, 20 mg, 30 mg, 5 mg</i>	T2	
<i>isosorbide mononitrate</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>isradipine</i>	T4	
<i>ivabradine oral tablet 5 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>ivabradine oral tablet 7.5 mg</i>	T4	PA; QL (62 EA per 31 days)
JANTOVEN	T1	
KERENDIA	T4	PA; QL (31 EA per 31 days)
<i>labetalol oral tablet 100 mg, 200 mg, 300 mg</i>	T2	
<i>lisinopril</i>	T1	
<i>lisinopril-hydrochlorothiazide</i>	T1	
<i>losartan oral tablet 100 mg</i>	T1	QL (31 EA per 31 days)
<i>losartan oral tablet 25 mg</i>	T1	QL (93 EA per 31 days)
<i>losartan oral tablet 50 mg</i>	T1	QL (62 EA per 31 days)
<i>losartan-hydrochlorothiazide</i>	T1	
<i>lovastatin</i>	T1	
<i>metolazone</i>	T2	
<i>metoprolol succinate</i>	T2	
<i>metoprolol ta-hydrochlorothiaz</i>	T2	
<i>metoprolol tartrate oral tablet 100 mg, 25 mg, 50 mg</i>	T1	
<i>metoprolol tartrate oral tablet 37.5 mg, 75 mg</i>	T2	
<i>metyrosine</i>	T5	PA
<i>mexiletine</i>	T3	
<i>minoxidil oral</i>	T2	
<i>moexipril</i>	T1	
MULPLETA	T5	PA
MULTAQ	T4	
<i>nadolol</i>	T4	
<i>nebivolol oral tablet 10 mg, 2.5 mg</i>	T2	QL (93 EA per 31 days)
<i>nebivolol oral tablet 20 mg</i>	T2	QL (62 EA per 31 days)
<i>nebivolol oral tablet 5 mg</i>	T2	QL (217 EA per 31 days)
NEXLETOL	T4	PA; QL (31 EA per 31 days)
NEXLIZET	T4	PA; QL (31 EA per 31 days)
<i>niacin oral tablet extended release 24 hr 1,000 mg, 750 mg</i>	T4	
<i>niacin oral tablet extended release 24 hr 500 mg</i>	T4	QL (31 EA per 31 days)
<i>nicardipine oral</i>	T4	
<i>nifedipine oral tablet extended release</i>	T2	
<i>nifedipine oral tablet extended release 24hr</i>	T2	
<i>nimodipine oral capsule</i>	T4	
NITRO-BID	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>nitroglycerin sublingual</i>	T2	
<i>nitroglycerin transdermal patch 24 hour</i>	T2	
<i>nitroglycerin translingual</i>	T4	
<i>olmesartan oral tablet 20 mg, 40 mg</i>	T1	QL (31 EA per 31 days)
<i>olmesartan oral tablet 5 mg</i>	T1	QL (93 EA per 31 days)
<i>olmesartan-amlodipin-hcthiazyd</i>	T3	
<i>olmesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>omega-3 acid ethyl esters</i>	T3	QL (124 EA per 31 days)
PACERONE ORAL TABLET 100 MG, 200 MG, 400 MG	T2	
<i>pentoxifylline</i>	T2	
<i>perindopril erbumine</i>	T2	
<i>pindolol</i>	T4	
<i>pitavastatin calcium</i>	T3	
<i>prasugrel hcl</i>	T3	
<i>pravastatin</i>	T1	
<i>prazosin</i>	T2	
PREVALITE ORAL POWDER IN PACKET	T4	
<i>propafenone oral capsule,extended release 12 hr</i>	T4	
<i>propafenone oral tablet</i>	T2	
<i>propranolol oral capsule,extended release 24 hr</i>	T2	
<i>propranolol oral tablet</i>	T2	
<i>quinapril</i>	T1	
<i>quinapril-hydrochlorothiazide</i>	T1	
<i>quinidine sulfate oral tablet</i>	T2	
<i>ramipril</i>	T1	
<i>ranolazine</i>	T4	QL (62 EA per 31 days)
REPATHA SURECLICK	T3	PA; QL (3 ML per 28 days)
REPATHA SYRINGE	T3	PA; QL (3 ML per 28 days)
<i>rosuvastatin</i>	T1	
<i>sacubitril-valsartan oral tablet 24-26 mg</i>	T2	QL (186 EA per 31 days)
<i>sacubitril-valsartan oral tablet 49-51 mg</i>	T2	QL (93 EA per 31 days)
<i>sacubitril-valsartan oral tablet 97-103 mg</i>	T2	QL (62 EA per 31 days)
<i>simvastatin</i>	T1	
SOTALOL AF	T2	
<i>sotalol oral</i>	T2	
<i>spironolactone oral tablet</i>	T1	
<i>spironolacton-hydrochlorothiaz</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>telmisartan</i>	T1	
<i>telmisartan-amlodipine</i>	T2	
<i>telmisartan-hydrochlorothiazid</i>	T1	
<i>terazosin</i>	T1	
TIADYL ER	T2	
<i>ticagrelor</i>	T2	
<i>timolol maleate oral</i>	T3	
<i>toremide oral</i>	T1	
<i>trandolapril</i>	T1	
<i>triamterene-hydrochlorothiazid</i>	T1	
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 400 MCG, 600 MCG, 800 MCG	T5	PA; QL (62 EA per 31 days)
UPTRAVI ORAL TABLET 200 MCG	T5	PA; QL (224 EA per 28 days)
UPTRAVI ORAL TABLETS,DOSE PACK	T5	PA; QL (400 EA per 365 days)
<i>valsartan oral tablet 160 mg, 40 mg, 80 mg</i>	T1	QL (62 EA per 31 days)
<i>valsartan oral tablet 320 mg</i>	T1	QL (31 EA per 31 days)
<i>valsartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>verapamil oral capsule, 24 hr er pellet ct</i>	T4	
<i>verapamil oral capsule,ext rel. pellets 24 hr 120 mg, 180 mg, 240 mg</i>	T2	
<i>verapamil oral capsule,ext rel. pellets 24 hr 360 mg</i>	T4	
<i>verapamil oral tablet</i>	T2	
<i>verapamil oral tablet extended release</i>	T2	
VERQUVO	T4	PA; QL (31 EA per 31 days)
<i>warfarin</i>	T1	
XARELTO DVT-PE TREAT 30D START	T3	QL (51 EA per 30 days)
XARELTO ORAL SUSPENSION FOR RECONSTITUTION	T3	QL (930 ML per 31 days)
XARELTO ORAL TABLET 10 MG, 20 MG	T3	QL (31 EA per 31 days)
XARELTO ORAL TABLET 15 MG	T3	QL (52 EA per 31 days)
XARELTO ORAL TABLET 2.5 MG	T3	QL (62 EA per 31 days)
Dermatologicals/Topical Therapy		
ACCUTANE ORAL CAPSULE 10 MG, 20 MG, 40 MG	T4	
<i>acitretin</i>	T4	PA
<i>acyclovir topical ointment</i>	T3	QL (30 GM per 30 days)
ALA-CORT TOPICAL CREAM	T1	
<i>alclometasone</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>ammonium lactate</i>	T2	
AMNESTEEM	T4	
ANZUPGO	T5	PA; QL (60 GM per 30 days)
<i>betamethasone dipropionate topical cream</i>	T2	
<i>betamethasone dipropionate topical lotion</i>	T2	
<i>betamethasone valerate topical cream</i>	T2	
<i>betamethasone valerate topical lotion</i>	T3	
<i>betamethasone valerate topical ointment</i>	T2	
<i>betamethasone, augmented topical cream</i>	T2	
<i>betamethasone, augmented topical gel</i>	T4	
<i>betamethasone, augmented topical lotion</i>	T4	
<i>betamethasone, augmented topical ointment</i>	T3	
<i>calcipotriene scalp</i>	T3	QL (60 ML per 28 days)
<i>calcipotriene topical cream</i>	T4	QL (60 GM per 28 days)
<i>calcipotriene topical ointment</i>	T3	QL (60 GM per 28 days)
<i>ciclopirox topical cream</i>	T2	QL (90 GM per 28 days)
<i>ciclopirox topical gel</i>	T3	QL (45 GM per 28 days)
<i>ciclopirox topical shampoo</i>	T3	QL (120 ML per 28 days)
<i>ciclopirox topical solution</i>	T2	
<i>ciclopirox topical suspension</i>	T3	QL (60 ML per 28 days)
CLARAVIS	T4	
<i>clindamycin phosphate topical gel</i>	T2	QL (60 GM per 28 days)
<i>clindamycin phosphate topical lotion</i>	T2	QL (60 ML per 28 days)
<i>clindamycin phosphate topical solution</i>	T2	QL (60 ML per 28 days)
<i>clindamycin phosphate topical swab</i>	T2	
<i>clindamycin-benzoyl peroxide topical gel 1.2 % (1 % base) -5 %</i>	T2	
<i>clotrimazole topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole topical solution</i>	T3	QL (30 ML per 28 days)
<i>clotrimazole-betamethasone topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole-betamethasone topical lotion</i>	T3	QL (60 ML per 28 days)
COSENTYX (2 SYRINGES)	T5	PA; QL (2 ML per 28 days)
COSENTYX PEN (2 PENS)	T5	PA; QL (2 ML per 28 days)
COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
COSENTYX UNOREADY PEN	T5	PA; QL (2 ML per 28 days)
<i>desoximetasone topical cream</i>	T4	QL (100 GM per 28 days)
<i>desoximetasone topical gel</i>	T4	QL (60 GM per 28 days)
<i>diclofenac sodium topical gel 3 %</i>	T4	PA; QL (100 GM per 28 days)

Drug Name	Drug Tier	Requirements/Limits
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
ERY PADS	T3	
<i>erythromycin with ethanol topical gel</i>	T2	QL (60 GM per 28 days)
<i>erythromycin with ethanol topical solution</i>	T2	QL (60 ML per 28 days)
FILSUEVZ	T5	PA
<i>fluocinolone and shower cap</i>	T4	QL (118.28 ML per 28 days)
<i>fluocinolone topical cream 0.01 %</i>	T3	QL (60 GM per 28 days)
<i>fluocinolone topical cream 0.025 %</i>	T3	QL (120 GM per 28 days)
<i>fluocinolone topical ointment</i>	T3	QL (120 GM per 28 days)
<i>fluocinolone topical solution</i>	T3	QL (90 ML per 28 days)
<i>fluocinonide topical cream 0.05 %</i>	T2	QL (60 GM per 28 days)
<i>fluocinonide topical gel</i>	T3	QL (60 GM per 28 days)
<i>fluocinonide topical ointment</i>	T2	QL (60 GM per 28 days)
<i>fluocinonide topical solution</i>	T2	QL (60 ML per 28 days)
<i>fluocinonide-emollient</i>	T4	QL (60 GM per 28 days)
<i>fluorouracil topical cream 5 %</i>	T3	
<i>fluorouracil topical solution 2 %</i>	T2	
<i>fluorouracil topical solution 5 %</i>	T3	
<i>fluticasone propionate topical cream</i>	T2	
<i>gentamicin topical</i>	T3	QL (60 GM per 28 days)
<i>halobetasol propionate topical cream</i>	T4	QL (50 GM per 28 days)
<i>halobetasol propionate topical ointment</i>	T4	QL (50 GM per 28 days)
<i>hydrocortisone topical cream 1 %</i>	T1	
<i>hydrocortisone topical lotion 2.5 %</i>	T2	QL (118 ML per 28 days)
<i>hydrocortisone topical ointment 1 %, 2.5 %</i>	T1	
<i>imiquimod topical cream in packet 5 %</i>	T2	
<i>ketoconazole topical cream</i>	T2	QL (60 GM per 28 days)
<i>ketoconazole topical shampoo</i>	T2	QL (120 ML per 28 days)
<i>lidocaine hcl mucous membrane solution 2 %</i>	T2	
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i>	T2	PA; QL (50 ML per 28 days)
<i>lidocaine topical adhesive patch,medicated 5 %</i>	T4	PA; QL (93 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>lidocaine topical ointment</i>	T4	PA; QL (50 GM per 28 days)
<i>lidocaine-prilocaine topical cream</i>	T2	PA; QL (30 GM per 28 days)
<i>malathion</i>	T4	
<i>metronidazole topical cream</i>	T3	
<i>metronidazole topical gel 0.75 %</i>	T3	
<i>metronidazole topical gel 1 %</i>	T4	
<i>metronidazole topical lotion</i>	T4	
<i>mometasone topical</i>	T2	
<i>mupirocin</i>	T2	
NYAMYC	T2	QL (60 GM per 28 days)
<i>nystatin topical cream</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical ointment</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical powder</i>	T2	QL (60 GM per 28 days)
NYSTOP	T2	QL (60 GM per 28 days)
PANRETIN	T5	PA-NS
<i>penciclovir</i>	T4	QL (5 GM per 28 days)
<i>permethrin</i>	T2	
<i>podofilox topical solution</i>	T4	
<i>selenium sulfide topical lotion</i>	T1	
SILIQ	T5	PA; QL (6 ML per 28 days)
<i>silver sulfadiazine</i>	T1	
SKYRIZI SUBCUTANEOUS PEN INJECTOR	T5	PA; QL (1 ML per 84 days)
SKYRIZI SUBCUTANEOUS SYRINGE	T5	PA; QL (1 ML per 84 days)
SSD	T4	
STELARA SUBCUTANEOUS SOLUTION	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T3	PA; QL (0.5 ML per 84 days)
STEQEYMA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
<i>sulfacetamide sodium (acne)</i>	T2	
SULFAMYLON TOPICAL CREAM	T3	
<i>tacrolimus topical</i>	T4	QL (100 GM per 28 days)
<i>tazarotene topical cream</i>	T4	PA; QL (60 GM per 28 days)
<i>tretinoin topical cream</i>	T2	PA; QL (45 GM per 28 days)

Drug Name	Drug Tier	Requirements/Limits
<i>tretinoin topical gel 0.01 %, 0.025 %</i>	T4	PA; QL (45 GM per 28 days)
<i>triamcinolone acetonide topical cream</i>	T2	
<i>triamcinolone acetonide topical lotion</i>	T2	
<i>triamcinolone acetonide topical ointment 0.025 %, 0.1 %, 0.5 %</i>	T2	
TRIDERM TOPICAL CREAM 0.5 %	T4	
<i>ustekinumab subcutaneous solution</i>	T5	PA; QL (0.5 ML per 84 days)
<i>ustekinumab subcutaneous syringe 45 mg/0.5 ml</i>	T5	PA; QL (0.5 ML per 84 days)
<i>ustekinumab subcutaneous syringe 90 mg/ml</i>	T5	PA; QL (1 ML per 56 days)
VALCHLOR	T5	PA-NS
YESINTEK SUBCUTANEOUS SOLUTION	T3	PA; QL (0.5 ML per 84 days)
YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T3	PA; QL (0.5 ML per 84 days)
YESINTEK SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
ZELSUVMI	T5	PA; QL (31 GM per 28 days)
ZORYVE TOPICAL CREAM 0.15 %	T4	PA; QL (60 GM per 28 days)
Diagnosics / Miscellaneous Agents		
<i>acamprosate</i>	T4	
<i>anagrelide</i>	T4	
<i>bupropion hcl (smoking deter)</i>	T2	QL (62 EA per 31 days)
<i>carglumic acid</i>	T5	PA
CHEMET	T4	
CLINIMIX 4.25%/D5W SULFIT FREE	T4	PA-BvD
<i>d10 %-0.45 % sodium chloride</i>	T2	
<i>d2.5 %-0.45 % sodium chloride</i>	T2	
<i>d5 % and 0.9 % sodium chloride</i>	T2	
<i>d5 %-0.45 % sodium chloride</i>	T2	
<i>deferasirox oral tablet, dispersible 125 mg</i>	T4	PA
<i>deferasirox oral tablet, dispersible 250 mg, 500 mg</i>	T5	PA
<i>dextrose 10 % in water (d10w)</i>	T2	
<i>dextrose 5 % in water (d5w) intravenous parenteral solution</i>	T2	
<i>disulfiram</i>	T3	
<i>droxidopa oral capsule 100 mg</i>	T5	PA; QL (465 EA per 31 days)
<i>droxidopa oral capsule 200 mg, 300 mg</i>	T5	PA; QL (186 EA per 31 days)
DUVYZAT	T5	PA; QL (420 ML per 35 days)
FABHALTA	T5	PA; QL (62 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>glycerol phenylbutyrate</i>	T5	PA
INCRELEX	T5	PA
JOENJA	T5	PA; QL (60 EA per 30 days)
KIONEX ORAL SUSPENSION	T2	
KYGEVVI	T5	PA; QL (496 EA per 30 days)
<i>levocarnitine (with sugar)</i>	T4	PA-BvD
<i>levocarnitine oral tablet</i>	T4	PA-BvD
LITFULO	T5	PA; QL (28 EA per 28 days)
LOARGYS	T5	PA
LOKELMA	T3	PA; QL (93 EA per 31 days)
<i>midodrine</i>	T2	
NICOTROL NS	T4	
<i>nitisinone</i>	T5	PA
PHEBURANE	T5	PA; QL (620 GM per 31 days)
<i>pilocarpine hcl oral</i>	T3	
PROLASTIN-C INTRAVENOUS SOLUTION	T5	PA
REVCOVI	T5	
REZDIFFRA	T5	PA; QL (31 EA per 31 days)
<i>riluzole</i>	T4	
<i>risedronate oral tablet 30 mg</i>	T5	
<i>sevelamer carbonate oral powder in packet</i>	T4	PA-BvD
<i>sevelamer carbonate oral tablet</i>	T3	PA-BvD
<i>sodium chloride 0.9 % intravenous parenteral solution</i>	T2	
<i>sodium chloride irrigation</i>	T2	
<i>sodium phenylbutyrate</i>	T5	PA
<i>sodium polystyrene sulfonate</i>	T2	
SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 2.5 MG, 5 MG	T5	PA; QL (31 EA per 31 days)
SOHONOS ORAL CAPSULE 10 MG	T5	PA; QL (62 EA per 31 days)
SPS (WITH SORBITOL) ORAL	T2	
<i>trientine oral capsule 250 mg</i>	T5	QL (248 EA per 31 days)
<i>varenicline tartrate oral tablet</i>	T4	QL (60 EA per 30 days)
<i>varenicline tartrate oral tablets,dose pack</i>	T4	QL (106 EA per 365 days)
ZOKINVY	T5	PA
Ear, Nose / Throat Medications		
<i>acetic acid otic (ear)</i>	T2	
<i>azelastine nasal spray,non-aerosol 137 mcg (0.1 %)</i>	T2	QL (30 ML per 25 days)

Drug Name	Drug Tier	Requirements/Limits
<i>chlorhexidine gluconate mucous membrane</i>	T1	
<i>ciprofloxacin-dexamethasone</i>	T4	
<i>fluocinolone acetonide oil</i>	T4	
<i>ipratropium bromide nasal spray,non-aerosol 21 mcg (0.03 %)</i>	T2	QL (30 ML per 28 days)
<i>ipratropium bromide nasal spray,non-aerosol 42 mcg (0.06 %)</i>	T2	QL (15 ML per 28 days)
KOURZEQ	T2	
<i>neomycin-polymyxin-hc otic (ear)</i>	T3	
<i>ofloxacin otic (ear)</i>	T2	
<i>olopatadine nasal</i>	T3	QL (30.5 GM per 30 days)
PERIOGARD	T1	
<i>triamcinolone acetonide dental</i>	T2	
Endocrine/Diabetes		
<i>acarbose</i>	T2	QL (93 EA per 31 days)
ALCOHOL PADS	T2	PA
<i>cabergoline</i>	T3	
<i>calcitonin (salmon) nasal</i>	T3	PA-BvD
<i>calcitriol oral capsule</i>	T2	PA-BvD
<i>calcitriol oral solution</i>	T4	PA-BvD
CERDELGA	T5	PA; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 30 mg, 60 mg</i>	T4	PA-BvD; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 90 mg</i>	T4	PA-BvD; QL (124 EA per 31 days)
<i>danazol</i>	T4	
<i>dapagliflozin</i>	T3	QL (31 EA per 31 days)
<i>desmopressin nasal spray,non-aerosol 10 mcg/spray (0.1 ml)</i>	T4	
<i>desmopressin oral</i>	T2	
<i>dexamethasone oral solution</i>	T2	
<i>dexamethasone oral tablet</i>	T1	
<i>diazoxide</i>	T4	
<i>doxercalciferol oral</i>	T4	PA-BvD
FARXIGA	T3	QL (31 EA per 31 days)
FIASP FLEXTOUCH U-100 INSULIN	T3	
FIASP PENFILL U-100 INSULIN	T3	
FIASP U-100 INSULIN	T3	
<i>fludrocortisone</i>	T2	
<i>glimepiride oral tablet 1 mg, 2 mg, 4 mg</i>	T1	

Drug Name	Drug Tier	Requirements/Limits
<i>glipizide oral tablet 10 mg, 5 mg</i>	T1	
<i>glipizide oral tablet extended release 24hr</i>	T1	
<i>glipizide-metformin</i>	T1	
GLUCAGON EMERGENCY KIT (HUMAN)	T3	
<i>glyburide</i>	T2	
<i>glyburide-metformin</i>	T2	
GLYXAMBI	T3	QL (31 EA per 31 days)
GVOKE	T3	
GVOKE HYPOPEN 2-PACK	T3	
GVOKE PFS 1-PACK SYRINGE SUBCUTANEOUS SYRINGE 1 MG/0.2 ML	T3	
HUMALOG JUNIOR KWIKPEN U-100	T3	
HUMALOG KWIKPEN INSULIN	T3	
HUMALOG MIX 50-50 KWIKPEN	T3	
HUMALOG MIX 75-25 KWIKPEN	T3	
HUMALOG MIX 75-25(U-100)INSULN	T3	
HUMALOG U-100 INSULIN	T3	
HUMULIN 70/30 U-100 INSULIN	T3	
HUMULIN 70/30 U-100 KWIKPEN	T3	
HUMULIN N NPH INSULIN KWIKPEN	T3	
HUMULIN N NPH U-100 INSULIN	T3	
HUMULIN R REGULAR U-100 INSULN	T3	
HUMULIN R U-500 (CONC) INSULIN	T3	
HUMULIN R U-500 (CONC) KWIKPEN	T3	
<i>hydrocortisone oral</i>	T1	
<i>insulin lispro</i>	T3	
<i>insulin lispro protamin-lispro</i>	T3	
JANUMET	T3	QL (62 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 100-1,000 MG, 50-500 MG	T3	QL (31 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 50-1,000 MG	T3	QL (62 EA per 31 days)
JANUVIA ORAL TABLET 100 MG, 50 MG	T3	QL (31 EA per 31 days)
JANUVIA ORAL TABLET 25 MG	T3	QL (93 EA per 31 days)
JARDIANCE ORAL TABLET 10 MG	T3	QL (62 EA per 31 days)
JARDIANCE ORAL TABLET 25 MG	T3	QL (31 EA per 31 days)
JAVYGTOR	T5	PA

Drug Name	Drug Tier	Requirements/Limits
JENTADUETO ORAL TABLET 2.5-1,000 MG, 2.5-500 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 5-1,000 MG	T3	QL (31 EA per 31 days)
LANTUS SOLOSTAR U-100 INSULIN	T3	
LANTUS U-100 INSULIN	T3	
<i>levothyroxine oral tablet</i>	T1	
LEVOXYL ORAL TABLET 100 MCG, 112 MCG, 125 MCG, 137 MCG, 150 MCG, 175 MCG, 200 MCG, 25 MCG, 50 MCG, 75 MCG, 88 MCG	T3	
LIOMNY	T2	
<i>liothyronine oral</i>	T2	
<i>liraglutide</i>	T4	PA; QL (9 ML per 30 days)
<i>metformin oral tablet 1,000 mg, 500 mg, 850 mg</i>	T1	
<i>metformin oral tablet extended release 24 hr</i>	T1	
<i>metformin oral tablet extended release 24hr</i>	NF	
<i>metformin oral tablet,er gast.retention 24 hr</i>	NF	
<i>methimazole oral tablet 10 mg, 5 mg</i>	T2	
<i>methylprednisolone</i>	T2	
<i>mifepristone oral tablet 300 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>miglustat</i>	T5	PA; QL (93 EA per 31 days)
MOUNJARO	T3	PA; QL (2 ML per 28 days)
<i>nateglinide</i>	T1	QL (93 EA per 31 days)
NOVOLIN 70/30 U-100 INSULIN	T3	
NOVOLIN 70-30 FLEXPEN U-100	T3	
NOVOLIN N FLEXPEN	T3	
NOVOLIN N NPH U-100 INSULIN	T3	
NOVOLIN R FLEXPEN	T3	
NOVOLIN R REGULAR U100 INSULIN	T3	
NOVOLOG FLEXPEN U-100 INSULIN	T3	
NOVOLOG MIX 70-30 U-100 INSULN	T3	
NOVOLOG MIX 70-30FLEXPEN U-100	T3	
NOVOLOG PENFILL U-100 INSULIN	T3	
NOVOLOG U-100 INSULIN ASPART	T3	
OZEMPIC SUBCUTANEOUS	T3	PA; QL (3 ML per 28 days)
<i>paricalcitol oral</i>	T4	PA-BvD
<i>pioglitazone</i>	T1	QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>pioglitazone-metformin</i>	T2	QL (93 EA per 31 days)
<i>prednisolone oral solution</i>	T2	
<i>prednisolone sodium phosphate oral solution 25 mg/5 ml (5 mg/ml), 5 mg base/5 ml (6.7 mg/5 ml)</i>	T2	
<i>prednisone oral solution</i>	T3	
<i>prednisone oral tablet</i>	T1	
<i>prednisone oral tablets,dose pack</i>	T2	
<i>propylthiouracil</i>	T2	
RECORLEV	T5	
<i>repaglinide oral tablet 0.5 mg, 1 mg</i>	T2	QL (124 EA per 31 days)
<i>repaglinide oral tablet 2 mg</i>	T2	QL (248 EA per 31 days)
RYBELSUS	T3	PA; QL (31 EA per 31 days)
<i>sapropterin</i>	T5	PA
SOLQUA 100/33	T3	QL (18 ML per 30 days)
SOMAVERT	T5	PA
SYNAREL	T5	PA
SYNJARDY	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 12.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 25-1,000 MG	T3	QL (31 EA per 31 days)
SYNTHROID	T3	
<i>testosterone cypionate</i>	T2	PA
<i>testosterone enanthate</i>	T3	PA
<i>testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)</i>	T4	PA
<i>tolvaptan</i>	T5	PA
<i>tolvaptan (polycys kidney dis) oral tablet</i>	T5	PA; QL (112 EA per 28 days)
<i>tolvaptan (polycys kidney dis) oral tablets, sequential</i>	T5	PA; QL (56 EA per 28 days)
TOUJEO MAX U-300 SOLOSTAR	T3	
TOUJEO SOLOSTAR U-300 INSULIN	T3	
TRADJENTA	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-5-1,000 MG, 25-5-1,000 MG	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 12.5-2.5-1,000 MG, 5-2.5-1,000 MG	T3	QL (62 EA per 31 days)
TRULICITY	T3	PA; QL (2 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
UNITHROID	T3	
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 10-500 MG, 5-500 MG	T3	QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
XULTOPHY 100/3.6	T3	QL (15 ML per 30 days)
YARGESA	T5	PA; QL (93 EA per 31 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML	T5	PA; QL (1.12 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 294 MCG/0.98 ML	T5	PA; QL (1.96 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 420 MCG/1.4 ML	T5	PA; QL (2.8 ML per 28 days)
ZELVYSIA ORAL POWDER IN PACKET 500 MG	T5	PA
Gastroenterology		
<i>alosetron oral tablet 0.5 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>alosetron oral tablet 1 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>aprepitant oral capsule 125 mg</i>	T5	PA-BvD
<i>aprepitant oral capsule 40 mg, 80 mg</i>	T4	PA-BvD
<i>aprepitant oral capsule, dose pack</i>	T4	PA-BvD
<i>balsalazide</i>	T4	
<i>betaine</i>	T5	
<i>budesonide oral capsule, delayed, extend. release</i>	T4	
<i>budesonide oral tablet, delayed and ext. release</i>	T5	
COMPRO	T4	
CONSTULOSE	T2	
CREON	T3	
<i>cromolyn oral</i>	T4	
<i>dicyclomine oral capsule</i>	T2	
<i>dicyclomine oral solution</i>	T2	
<i>dicyclomine oral tablet 20 mg</i>	T2	
<i>diphenoxylate-atropine oral liquid</i>	T4	
<i>diphenoxylate-atropine oral tablet</i>	T2	
<i>dronabinol</i>	T4	PA-BvD
ENULOSE	T2	
EOHILIA	T5	PA; QL (1800 ML per 365 days)
<i>esomeprazole magnesium oral capsule, delayed release(dr/ec)</i>	T2	QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>famotidine oral suspension for reconstitution</i>	T2	
<i>famotidine oral tablet 20 mg, 40 mg</i>	T1	
GATTEX 30-VIAL	T5	PA
GAVILYTE-C	T1	
GAVILYTE-G	T1	
GAVILYTE-N	T1	
GENERLAC	T2	
<i>glycopyrrolate oral tablet 1 mg, 2 mg</i>	T2	
<i>granisetron hcl oral</i>	T4	PA-BvD
<i>hydrocortisone rectal</i>	T4	
<i>hydrocortisone topical cream with perineal applicator 2.5 %</i>	T1	
<i>lactulose oral solution</i>	T2	
LINZESS	T3	QL (31 EA per 31 days)
<i>loperamide oral capsule</i>	T2	
<i>lubiprostone</i>	T4	QL (62 EA per 31 days)
<i>meclizine oral tablet 12.5 mg, 25 mg</i>	T2	
<i>mesalamine oral capsule (with del rel tablets)</i>	T4	QL (186 EA per 31 days)
<i>mesalamine oral capsule,extended release 24hr</i>	T4	QL (124 EA per 31 days)
<i>mesalamine oral tablet,delayed release (dr/ec) 1.2 gram</i>	T4	QL (124 EA per 31 days)
<i>mesalamine rectal enema</i>	T4	QL (1860 ML per 31 days)
<i>metoclopramide hcl oral solution</i>	T2	
<i>metoclopramide hcl oral tablet</i>	T2	
<i>misoprostol</i>	T2	
MOVANTI	T3	QL (31 EA per 31 days)
<i>nitroglycerin rectal</i>	T4	
<i>omeprazole oral capsule,delayed release(dr/ec)</i>	T1	
<i>ondansetron hcl oral solution</i>	T3	PA-BvD
<i>ondansetron hcl oral tablet 4 mg, 8 mg</i>	T2	PA-BvD
<i>ondansetron oral tablet,disintegrating 4 mg, 8 mg</i>	T2	PA-BvD
<i>pantoprazole oral tablet,delayed release (dr/ec)</i>	T1	
<i>peg 3350-electrolytes</i>	T1	
<i>peg3350-sod sul-nacl-kcl-asb-c</i>	T4	
<i>peg-electrolyte soln</i>	T1	
<i>prochlorperazine</i>	T4	
<i>prochlorperazine maleate</i>	T2	
PROCTO-MED HC	T2	
PROCTOSOL HC TOPICAL	T2	

Drug Name	Drug Tier	Requirements/Limits
PROCTOZONE-HC	T2	
<i>rabeprazole oral tablet, delayed release (dr/ec)</i>	T2	QL (62 EA per 31 days)
<i>scopolamine base</i>	T3	QL (10 EA per 30 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML)	T5	PA; QL (1.2 ML per 56 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 360 MG/2.4 ML (150 MG/ML)	T5	PA; QL (2.4 ML per 56 days)
<i>sodium, potassium, mag sulfates</i>	T4	
<i>sucralfate oral suspension</i>	T4	
<i>sucralfate oral tablet</i>	T2	
<i>sulfasalazine</i>	T2	
<i>ursodiol oral capsule 300 mg</i>	T4	
<i>ursodiol oral tablet</i>	T3	
VOWST	T5	PA; QL (12 EA per 14 days)
ZENPEP ORAL CAPSULE, DELAYED RELEASE (DR/EC) 10,000-32,000 -42,000 UNIT, 15,000-47,000 -63,000 UNIT, 20,000-63,000- 84,000 UNIT, 25,000-79,000- 105,000 UNIT, 3,000-10,000 -14,000-UNIT, 40,000-126,000- 168,000 UNIT, 5,000-17,000- 24,000 UNIT, 60,000-189,600- 252,600 UNIT	T3	
Immunology, Vaccines / Biotechnology		
ABRYSVO (PF)	T3	QL (1 EA per 365 days)
ACTHIB (PF)	T3	
ACTIMMUNE	T5	PA
ADACEL (TDAP ADOLESN/ADULT) (PF)	T3	
AREXVY (PF)	T3	QL (1 EA per 365 days)
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	T5	PA; QL (1 EA per 28 days)
AVONEX INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (1 EA per 28 days)
<i>bcg vaccine, live (pf)</i>	T3	
BESREMI	T5	PA-NS; QL (2 ML per 28 days)
BETASERON SUBCUTANEOUS KIT	T5	PA; QL (14 EA per 28 days)
BEXSERO	T3	
BOOSTRIX TDAP	T3	
DAPTACEL (DTAP PEDIATRIC) (PF)	T3	
ENGERIX-B (PF)	T3	PA-BvD
ENGERIX-B PEDIATRIC (PF)	T3	PA-BvD
GAMMAGARD LIQUID	T5	PA

Drug Name	Drug Tier	Requirements/Limits
GAMMAGARD LIQUID ERC	T5	PA
GAMMAGARD S-D (IGA < 1 MCG/ML)	T5	PA
GARDASIL 9 (PF)	T3	
HAVRIX (PF)	T3	
HEPLISAV-B (PF)	T3	PA-BvD
HIBERIX (PF)	T3	
IMOVAX RABIES VACCINE (PF)	T3	PA-BvD
INFANRIX (DTAP) (PF)	T3	
IPOL	T3	
IXIARO (PF)	T3	
JYNNEOS (PF)	T3	PA-BvD
KINRIX (PF)	T3	
LEUKINE INJECTION RECON SOLN	T5	PA
MENQUADFI (PF)	T4	
MENVEO A-C-Y-W-135-DIP (PF) INTRAMUSCULAR KIT	T3	
M-M-R II (PF)	T3	
MRESVIA (PF)	T3	QL (1 ML per 365 days)
NIVESTYM	T5	
NORDITROPIN FLEXP SUBCUTANEOUS PEN INJECTOR 10 MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML (10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML)	T5	PA
PEDIARIX (PF)	T3	
PEDVAX HIB (PF)	T3	
PEGASYS	T5	PA
PENBRAYA (PF)	T3	
PENMENVY MEN A-B-C-W-Y (PF)	T3	
PENTACEL (PF)	T3	
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PRIORIX (PF)	T3	
PRIVIGEN	T5	PA
PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML	T3	PA-BvD
PROCRIT INJECTION SOLUTION 20,000 UNIT/ML, 40,000 UNIT/ML	T5	PA-BvD

Drug Name	Drug Tier	Requirements/Limits
PROQUAD (PF)	T3	
QUADRACEL (PF)	T3	
RABAVERT (PF)	T3	PA-BvD
RECOMBIVAX HB (PF)	T3	PA-BvD
RETACRIT	T3	PA-BvD
ROTARIX ORAL SUSPENSION	T3	
ROTATEQ VACCINE	T3	
SHINGRIX (PF)	T3	QL (2 EA per 999 days)
TENIVAC (PF)	T3	
TICOVAC	T3	
TRUMENBA	T3	
TWINRIX (PF)	T3	
TYPHIM VI	T3	
VAQTA (PF)	T3	
VARIVAX (PF)	T3	
VAXCHORA VACCINE	T3	QL (200 ML per 365 days)
VIMKUNYA	T3	
VIVOTIF	T3	
XOLREMDI	T5	PA; QL (124 EA per 31 days)
YF-VAX (PF)	T3	
ZARXIO	T5	
Miscellaneous Supplies		
ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"	T4	
GAUZE PAD TOPICAL BANDAGE 2 X 2 "	T2	PA
<i>insulin syringe-needle u-100 syringe 0.3 ml 29 gauge, 1 ml 29 gauge x 1/2", 1/2 ml 28 gauge</i>	T3	
<i>pen needle, diabetic needle 29 gauge x 1/2"</i>	T4	
Musculoskeletal / Rheumatology		
ACTEMRA ACTPEN	T5	PA; QL (3.6 ML per 28 days)
ACTEMRA SUBCUTANEOUS	T5	PA; QL (3.6 ML per 28 days)
<i>alendronate oral tablet 10 mg, 35 mg, 70 mg</i>	T1	
<i>allopurinol oral tablet 100 mg, 300 mg</i>	T1	
BENLYSTA SUBCUTANEOUS	T5	PA; QL (4 ML per 28 days)
<i>colchicine oral tablet</i>	T2	QL (62 EA per 31 days)
ENBREL MINI	T5	PA; QL (8 ML per 28 days)
ENBREL SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5)	T5	PA; QL (4 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
ENBREL SUBCUTANEOUS SYRINGE 50 MG/ML (1 ML)	T5	PA; QL (8 ML per 28 days)
ENBREL SURECLICK	T5	PA; QL (8 ML per 28 days)
HADLIMA	T5	PA; QL (1.6 ML per 28 days)
HADLIMA PUSHTOUCH	T5	PA; QL (1.6 ML per 28 days)
HADLIMA(CF)	T5	PA; QL (0.8 ML per 28 days)
HADLIMA(CF) PUSHTOUCH	T5	PA; QL (0.8 ML per 28 days)
<i>ibandronate oral</i>	T2	
KINERET	T5	PA; QL (18.76 ML per 28 days)
<i>leflunomide</i>	T3	
ORENCIA CLICKJECT	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 50 MG/0.4 ML	T5	PA; QL (1.6 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 87.5 MG/0.7 ML	T5	PA; QL (2.8 ML per 28 days)
OTEZLA	T5	PA; QL (62 EA per 31 days)
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47)	T5	PA; QL (110 EA per 365 days)
OTEZLA XR	T5	PA; QL (30 EA per 30 days)
OTEZLA XR INITIATION	T5	PA; QL (82 EA per 365 days)
<i>penicillamine oral tablet</i>	T5	
<i>probenecid</i>	T2	
<i>probenecid-colchicine</i>	T2	
PROLIA	T4	PA; QL (1 ML per 180 days)
<i>raloxifene</i>	T3	
RINVOQ LQ	T5	PA; QL (372 ML per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG	T5	PA; QL (31 EA per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 45 MG	T5	PA; QL (168 EA per 365 days)
<i>risedronate oral tablet 150 mg, 35 mg, 35 mg (12 pack), 35 mg (4 pack), 5 mg</i>	T4	
<i>risedronate oral tablet,delayed release (dr/ec)</i>	T4	
SIMLANDI(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
SIMLANDI(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML	T5	PA; QL (2 EA per 28 days)
<i>teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)</i>	T5	PA; QL (2.48 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
TYMLOS	T5	PA; QL (1.56 ML per 30 days)
XELJANZ ORAL SOLUTION	T5	PA; QL (310 ML per 31 days)
XELJANZ ORAL TABLET	T5	PA; QL (62 EA per 31 days)
XELJANZ XR	T5	PA; QL (31 EA per 31 days)
YUFLYMA(CF)	T5	PA; QL (2 EA per 28 days)
YUFLYMA(CF) AI CROHN'S-UC-HS	T5	PA; QL (6 EA per 365 days)
YUFLYMA(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
Obstetrics / Gynecology		
ABIGALE	T2	
ALTAVERA (28)	T2	
ALYACEN 1/35 (28)	T2	
APRI	T2	
ARANELLE (28)	T2	
AVIANE	T2	
AZURETTE (28)	T2	
BALZIVA (28)	T2	
BLISOVI FE 1.5/30 (28)	T2	
BRIELLYN	T2	
CAMILA	T2	
<i>clindamycin phosphate vaginal</i>	T2	
CRYSELLE (28)	T2	
CYRED EQ	T2	
DEPO-SUBQ PROVERA 104	T3	
<i>drospirenone-ethinyl estradiol</i>	T2	
ENSKYCE	T2	
ERRIN	T2	
ESTARYLLA	T2	
<i>estradiol oral</i>	T2	
<i>estradiol transdermal patch weekly 0.025 mg/24 hr, 0.06 mg/24 hr, 0.1 mg/24 hr</i>	T3	
<i>estradiol transdermal patch weekly 0.0375 mg/24 hr, 0.05 mg/24 hr, 0.075 mg/24 hr</i>	T2	
<i>estradiol vaginal</i>	T4	
<i>estradiol-norethindrone acet</i>	T2	
<i>etonogestrel-ethinyl estradiol</i>	T3	
GALLIFREY	T2	
HAILEY 24 FE	T2	
HEATHER	T2	
ICLEVIA	T2	

Drug Name	Drug Tier	Requirements/Limits
IMVEXXY MAINTENANCE PACK	T3	
IMVEXXY STARTER PACK	T3	
INCASSIA	T2	
INTROVALE	T2	
ISIBLOOM	T2	
JASMIEL (28)	T2	
JINTELI	T4	
JULEBER	T2	
JUNEL 1.5/30 (21)	T2	
JUNEL 1/20 (21)	T2	
JUNEL FE 1.5/30 (28)	T2	
JUNEL FE 1/20 (28)	T2	
JUNEL FE 24	T2	
KARIVA (28)	T2	
KELNOR 1/35 (28)	T2	
KURVELO (28)	T2	
LESSINA	T2	
LEVONEST (28)	T2	
<i>levonorgestrel-ethinyl estrad oral tablet 0.1-20 mg-mcg</i>	T2	
<i>levonorgestrel-ethinyl estrad oral tablets,dose pack,3 month</i>	T2	
<i>levonorg-eth estrad triphasic</i>	T2	
LILETTA	T3	
LORYNA (28)	T2	
LOW-OGESTREL (28)	T2	
LUIZZA	T2	
LUTERA (28)	T2	
LYLEQ	T2	
LYZA	T2	
MARLISSA (28)	T2	
<i>medroxyprogesterone</i>	T2	
<i>metronidazole vaginal gel 0.75 % (37.5mg/5 gram)</i>	T3	
MICONAZOLE-3 VAGINAL SUPPOSITORY	T2	
MICROGESTIN 1.5/30 (21)	T2	
MICROGESTIN 1/20 (21)	T2	
MICROGESTIN FE 1.5/30 (28)	T2	

Drug Name	Drug Tier	Requirements/Limits
MICROGESTIN FE 1/20 (28)	T2	
MILI	T2	
NECON 0.5/35 (28)	T2	
NEXPLANON	T3	
<i>norethindrone (contraceptive)</i>	T2	
<i>norethindrone acetate</i>	T2	
<i>norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg</i>	T3	
<i>norethindrone ac-eth estradiol oral tablet 1-20 mg-mcg</i>	T2	
<i>norgestimate-ethinyl estradiol oral tablet 0.18/0.215/0.25 mg-0.035mg (28), 0.25-0.035 mg</i>	T2	
NORTREL 0.5/35 (28)	T2	
NORTREL 1/35 (21)	T2	
NORTREL 1/35 (28)	T2	
NORTREL 7/7/7 (28)	T2	
NYLIA 1/35 (28)	T2	
NYLIA 7/7/7 (28)	T2	
PIMTREA (28)	T2	
PORTIA 28	T2	
PREMARIN VAGINAL	T3	
RECLIPSEN (28)	T2	
SETLAKIN	T2	
SPRINTEC (28)	T2	
SYEDA	T2	
<i>terconazole</i>	T2	
<i>tranexamic acid oral</i>	T3	
TRI-ESTARYLLA	T2	
TRI-MILI	T2	
TRI-SPRINTEC (28)	T2	
TRI-VYLIBRA	T2	
TURQOZ (28)	T2	
VALTYA	T2	
VELIVET TRIPHASIC REGIMEN (28)	T2	
VESTURA (28)	T2	
VIENVA	T2	
VIORELE (28)	T2	
VYFEMLA (28)	T2	
VYLIBRA	T2	

Drug Name	Drug Tier	Requirements/Limits
XELRIA FE	T2	
YUVAFEM	T4	
ZAFEMY	T3	
ZOVIA 1-35 (28)	T2	
Ophthalmology		
<i>acetazolamide</i>	T3	
ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.1 %	T3	
<i>apraclonidine</i>	T3	
<i>azelastine ophthalmic (eye)</i>	T2	
<i>bacitracin-polymyxin b</i>	T2	
BESIVANCE	T4	
<i>betaxolol ophthalmic (eye)</i>	T3	
<i>bimatoprost ophthalmic (eye) drops 0.03 %</i>	T2	
<i>brimonidine ophthalmic (eye) drops 0.1 %</i>	T3	
<i>brimonidine ophthalmic (eye) drops 0.15 %</i>	T4	
<i>brimonidine ophthalmic (eye) drops 0.2 %</i>	T2	
<i>brimonidine-timolol</i>	T3	
<i>brinzolamide</i>	T4	
<i>carteolol</i>	T2	
<i>ciprofloxacin hcl ophthalmic (eye)</i>	T2	
COMBIGAN	T3	
<i>cromolyn ophthalmic (eye)</i>	T2	
<i>cyclosporine ophthalmic (eye)</i>	T3	QL (60 EA per 30 days)
CYSTARAN	T5	PA; QL (60 ML per 28 days)
<i>dexamethasone sodium phosphate ophthalmic (eye)</i>	T2	
<i>diclofenac sodium ophthalmic (eye)</i>	T2	
<i>difluprednate</i>	T4	
<i>dorzolamide</i>	T2	
<i>dorzolamide-timolol</i>	T2	
<i>erythromycin ophthalmic (eye)</i>	T2	
<i>fluorometholone</i>	T3	
<i>flurbiprofen sodium</i>	T2	
<i>gatifloxacin</i>	T3	
<i>gentamicin ophthalmic (eye) drops</i>	T2	
<i>ketorolac ophthalmic (eye) drops 0.4 %</i>	T3	
<i>ketorolac ophthalmic (eye) drops 0.5 %</i>	T2	
<i>latanoprost</i>	T1	

Drug Name	Drug Tier	Requirements/Limits
<i>levobunolol ophthalmic (eye) drops 0.5 %</i>	T2	
LUMIGAN OPHTHALMIC (EYE) DROPS 0.01 %	T3	QL (5 ML per 31 days)
<i>methazolamide</i>	T4	
<i>moxifloxacin ophthalmic (eye) drops</i>	T3	
<i>neomycin-bacitracin-poly-hc</i>	T3	
<i>neomycin-bacitracin-polymyxin</i>	T2	
<i>neomycin-polymyxin b-dexameth</i>	T2	
<i>neomycin-polymyxin-gramicidin</i>	T3	
<i>neomycin-polymyxin-hc ophthalmic (eye)</i>	T3	
<i>ofloxacin ophthalmic (eye)</i>	T2	
<i>pilocarpine hcl ophthalmic (eye) drops 1 %, 2 %, 4 %</i>	T3	
<i>polymyxin b sulf-trimethoprim</i>	T1	
<i>prednisolone acetate</i>	T2	
<i>prednisolone sodium phosphate ophthalmic (eye)</i>	T2	
RESTASIS	T3	QL (60 EA per 30 days)
RESTASIS MULTIDOSE	T3	QL (5.5 ML per 27 days)
RHOPRESSA	T4	ST
ROCKLATAN	T4	ST
SIMBRINZA	T4	
<i>sulfacetamide sodium ophthalmic (eye) drops</i>	T2	
<i>sulfacetamide-prednisolone</i>	T2	
<i>timolol maleate ophthalmic (eye) drops</i>	T1	
<i>timolol maleate ophthalmic (eye) gel forming solution</i>	T3	
<i>tobramycin ophthalmic (eye)</i>	T1	
<i>tobramycin-dexamethasone</i>	T3	
<i>travoprost</i>	T3	
<i>trifluridine</i>	T3	
XDEMVY	T5	PA; QL (10 ML per 42 days)
XIIDRA	T3	QL (60 EA per 30 days)
ZIRGAN	T4	ST
Respiratory And Allergy		
<i>acetylcysteine</i>	T2	PA-BvD
ADEMPAS	T5	PA; QL (93 EA per 31 days)
ADVAIR HFA	T3	QL (12 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation</i>	T3	QL (17 GM per 30 days)

Drug Name	Drug Tier	Requirements/Limits
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020503)</i>	T3	QL (13.4 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020983)</i>	NF	
<i>albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml</i>	T2	PA-BvD
<i>albuterol sulfate oral syrup</i>	T1	
<i>ambrisentan</i>	T5	PA; QL (31 EA per 31 days)
ANORO ELLIPTA	T3	QL (60 EA per 30 days)
ASMANEX HFA	T3	QL (13 GM per 30 days)
ASMANEX TWISTHALER INHALATION AEROSOL POWDR BREATH ACTIVATED 110 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (120), 220 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (60)	T3	QL (1 EA per 30 days)
ATROVENT HFA	T4	QL (25.8 GM per 30 days)
<i>azelastine-fluticasone</i>	T4	QL (23 GM per 30 days)
BREO ELLIPTA	T3	QL (60 EA per 30 days)
BREYNA	T3	QL (10.3 GM per 30 days)
BREZTRI AEROSPHERE	T3	QL (10.7 GM per 30 days)
BRINSUPRI	T5	PA; QL (31 EA per 31 days)
<i>budesonide inhalation</i>	T4	PA-BvD
<i>budesonide-formoterol</i>	T3	QL (10.2 GM per 30 days)
CINRYZE	T5	PA; QL (20 EA per 28 days)
COMBIVENT RESPIMAT	T3	QL (4 GM per 30 days)
<i>cromolyn inhalation</i>	T4	PA-BvD
<i>cyproheptadine oral tablet</i>	T2	PA
<i>desloratadine oral tablet</i>	T2	QL (31 EA per 31 days)
DULERA	T3	QL (13 GM per 30 days)
<i>epinephrine injection auto-injector 0.15 mg/0.3 ml, 0.3 mg/0.3 ml</i>	T3	
FASENRA PEN	T5	PA; QL (1 ML per 28 days)
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (0.5 ML per 56 days)
FASENRA SUBCUTANEOUS SYRINGE 30 MG/ML	T5	PA; QL (1 ML per 28 days)
<i>flunisolide</i>	T3	QL (50 ML per 25 days)
<i>fluticasone propionate inhalation blister with device 100 mcg/actuation, 50 mcg/actuation</i>	T4	ST; QL (60 EA per 30 days)

Drug Name	Drug Tier	Requirements/Limits
<i>fluticasone propionate inhalation blister with device 250 mcg/actuation</i>	T4	ST; QL (240 EA per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 110 mcg/actuation</i>	T4	ST; QL (12 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 220 mcg/actuation</i>	T4	ST; QL (24 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 44 mcg/actuation</i>	T4	ST; QL (10.6 GM per 30 days)
<i>fluticasone propionate nasal</i>	T1	QL (16 GM per 30 days)
<i>fluticasone propion-salmeterol inhalation blister with device</i>	T3	QL (60 EA per 30 days)
<i>hydroxyzine hcl oral tablet</i>	T2	PA
<i>icatibant</i>	T5	PA; QL (18 ML per 30 days)
<i>ipratropium bromide inhalation solution</i>	T2	PA-BvD
<i>ipratropium-albuterol</i>	T2	PA-BvD
JASCAYD	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 5.8 MG, 50 MG, 75 MG	T5	PA; QL (56 EA per 28 days)
KALYDECO ORAL GRANULES IN PACKET 25 MG	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL TABLET	T5	PA; QL (62 EA per 31 days)
<i>levalbuterol hcl inhalation solution for nebulization 1.25 mg/3 ml</i>	T3	PA-BvD
<i>levalbuterol tartrate</i>	T3	QL (30 GM per 30 days)
<i>levocetirizine oral solution</i>	T4	QL (310 ML per 31 days)
<i>levocetirizine oral tablet</i>	T1	QL (31 EA per 31 days)
<i>mometasone nasal</i>	T3	QL (34 GM per 30 days)
<i>montelukast oral tablet</i>	T2	QL (31 EA per 31 days)
<i>montelukast oral tablet,chewable</i>	T2	QL (31 EA per 31 days)
OFEV	T5	PA; QL (62 EA per 31 days)
OPSUMIT	T5	PA; QL (31 EA per 31 days)
OPSYNVI	T5	PA; QL (31 EA per 31 days)
ORKAMBI ORAL GRANULES IN PACKET	T5	PA; QL (62 EA per 31 days)
ORKAMBI ORAL TABLET	T5	PA; QL (124 EA per 31 days)
<i>pirfenidone oral capsule</i>	T5	PA; QL (279 EA per 31 days)
<i>pirfenidone oral tablet</i>	T5	PA; QL (93 EA per 31 days)
<i>promethazine oral tablet</i>	T4	PA
PULMOZYME	T5	PA
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION	T3	QL (10.6 GM per 30 days)

Drug Name	Drug Tier	Requirements/Limits
QVAR REDHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION	T3	QL (21.2 GM per 30 days)
<i>roflumilast</i>	T4	QL (31 EA per 31 days)
SAJAZIR	T5	PA; QL (18 ML per 30 days)
SEREVENT DISKUS	T3	QL (60 EA per 30 days)
<i>sildenafil (pulm.hypertension) oral tablet</i>	T3	PA; QL (372 EA per 31 days)
SPIRIVA RESPIMAT	T3	QL (4 GM per 30 days)
STIOLTO RESPIMAT	T3	QL (4 GM per 30 days)
STRIVERDI RESPIMAT	T4	QL (4 GM per 30 days)
<i>tadalafil (pulm. hypertension)</i>	T5	PA; QL (62 EA per 31 days)
<i>terbutaline oral</i>	T4	
THEO-24	T3	
<i>theophylline oral solution</i>	T2	
<i>theophylline oral tablet extended release 12 hr</i>	T2	
<i>theophylline oral tablet extended release 24 hr</i>	T2	
TRELEGY ELLIPTA	T3	QL (60 EA per 30 days)
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	T5	PA; QL (56 EA per 28 days)
TRIKAFTA ORAL TABLETS, SEQUENTIAL	T5	PA; QL (84 EA per 28 days)
VENTOLIN HFA	T3	QL (36 GM per 30 days)
WINREVAIR	T5	PA
WIXELA INHUB	T3	QL (60 EA per 30 days)
XOLAIR	T5	PA
Urologicals		
<i>alfuzosin</i>	T2	QL (31 EA per 31 days)
<i>bethanechol chloride oral tablet 10 mg, 25 mg, 5 mg</i>	T2	
<i>bethanechol chloride oral tablet 50 mg</i>	T3	
CYSTAGON	T4	
<i>dutasteride</i>	T2	QL (31 EA per 31 days)
<i>dutasteride-tamsulosin</i>	T4	QL (31 EA per 31 days)
ELMIRON	T4	
<i>finasteride oral tablet 5 mg</i>	T2	
GEMTESA	T4	QL (31 EA per 31 days)
MYRBETRIQ ORAL SUSPENSION,EXTENDED REL RECON	T3	QL (300 ML per 30 days)
MYRBETRIQ ORAL TABLET EXTENDED RELEASE 24 HR	T3	QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
<i>oxybutynin chloride oral syrup</i>	T2	
<i>oxybutynin chloride oral tablet 5 mg</i>	T2	
<i>oxybutynin chloride oral tablet extended release 24hr 10 mg, 5 mg</i>	T2	QL (31 EA per 31 days)
<i>oxybutynin chloride oral tablet extended release 24hr 15 mg</i>	T3	QL (62 EA per 31 days)
<i>potassium citrate oral tablet extended release</i>	T3	
RIVFLOZA SUBCUTANEOUS SOLUTION	T5	PA; QL (1 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML	T5	PA; QL (0.8 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 160 MG/ML	T5	PA; QL (1 ML per 28 days)
<i>silodosin</i>	T4	
<i>solifenacin</i>	T4	QL (31 EA per 31 days)
<i>tadalafil oral tablet 2.5 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>tadalafil oral tablet 5 mg</i>	T4	PA; QL (31 EA per 31 days)
<i>tamsulosin</i>	T1	
<i>tolterodine oral capsule,extended release 24hr</i>	T3	QL (31 EA per 31 days)
<i>tolterodine oral tablet</i>	T3	QL (62 EA per 31 days)
<i>trospium oral tablet</i>	T2	QL (93 EA per 31 days)
Vitamins, Hematinics / Electrolytes		
<i>calcium acetate(phosphat bind)</i>	T2	PA-BvD
CLINIMIX 5%/D15W SULFITE FREE	T4	PA-BvD
CLINIMIX 4.25%/D10W SULF FREE	T4	PA-BvD
CLINIMIX 5%-D20W(SULFITE-FREE)	T4	PA-BvD
<i>fluoride (sodium) oral tablet</i>	T2	
INTRALIPID INTRAVENOUS EMULSION 20 %	T4	PA-BvD
ISOLYTE S PH 7.4	T3	PA-BvD
ISOLYTE-P IN 5 % DEXTROSE	T4	PA-BvD
KLOR-CON	T4	
KLOR-CON M10	T2	
KLOR-CON M15	T2	
KLOR-CON M20	T2	
<i>magnesium sulfate injection</i>	T2	
PLENAMINE	T3	PA-BvD
<i>potassium chlorid-d5-0.45%nacl</i>	T2	
<i>potassium chloride in 0.9%nacl intravenous parenteral solution 20 meq/l, 40 meq/l</i>	T2	

Drug Name	Drug Tier	Requirements/Limits
<i>potassium chloride in 5 % dex intravenous parenteral solution 20 meq/l</i>	T2	
<i>potassium chloride in lr-d5</i>	T2	
<i>potassium chloride in water intravenous piggyback 10 meq/100 ml, 20 meq/100 ml, 40 meq/100 ml</i>	T2	
<i>potassium chloride intravenous</i>	T2	
<i>potassium chloride oral capsule, extended release</i>	T2	
<i>potassium chloride oral liquid</i>	T2	
<i>potassium chloride oral tablet extended release 10 meq, 20 meq, 8 meq</i>	T2	
<i>potassium chloride oral tablet,er particles/crystals</i>	T2	
<i>potassium chloride-0.45 % nacl</i>	T2	
<i>potassium chloride-d5-0.2%nacl</i>	T2	
<i>potassium chloride-d5-0.9%nacl</i>	T2	
PRENATAL VITAMIN PLUS LOW IRON	T2	PA
PROSOL 20 %	T4	PA-BvD
<i>sodium chloride 0.45 % intravenous</i>	T2	
<i>sodium chloride 3 % hypertonic</i>	T2	
<i>sodium chloride 5 % hypertonic</i>	T2	
TRAVASOL 10 %	T4	PA-BvD
TROPHAMINE 10 %	T4	PA-BvD

Index of Drugs

<i>abacavir</i>	3	<i>amitriptyline</i>	19	AVONEX	48
<i>abacavir-lamivudine</i>	3	<i>amlodipine</i>	31	AYVAKIT	11
ABIGALE	52	<i>amlodipine-benazepril</i>	31	<i>azathioprine</i>	11
ABILIFY MAINTENA	18	<i>amlodipine-olmesartan</i>	31	<i>azelastine</i>	41, 55
<i>abiraterone</i>	11	<i>amlodipine-valsartan</i>	31	<i>azelastine-fluticasone</i>	57
ABIRTEGA	11	<i>ammonium lactate</i>	37	<i>azithromycin</i>	4
ABRYSVO (PF)	48	AMNESTEEM	37	<i>aztreonam</i>	4
<i>acamprosate</i>	40	<i>amoxapine</i>	19	AZURETTE (28)	52
<i>acarbose</i>	42	<i>amoxicillin</i>	3	<i>bacitracin-polymyxin b</i>	55
ACCUTANE	36	<i>amoxicillin-pot clavulanate</i>	3	<i>baclofen</i>	19
<i>acebutolol</i>	31	<i>amphotericin b</i>	3	BAFIERTAM	19
<i>acetaminophen-codeine</i>	19	<i>amphotericin b liposome</i>	3	<i>balsalazide</i>	46
<i>acetazolamide</i>	55	<i>ampicillin</i>	4	BALVERSA	11
<i>acetic acid</i>	41	<i>ampicillin sodium</i>	4	BALZIVA (28)	52
<i>acetylcysteine</i>	56	<i>ampicillin-sulbactam</i>	4	<i>bcg vaccine, live (pf)</i>	48
<i>acitretin</i>	36	<i>anagrelide</i>	40	<i>benazepril</i>	31
ACTEMRA	50	<i>anastrozole</i>	11	<i>benazepril-hydrochlorothiazide</i> ..	31
ACTEMRA ACTPEN	50	ANORO ELLIPTA	57	BENLYSTA	50
ACTHIB (PF)	48	ANZUPGO	37	<i>benztropine</i>	19
ACTIMMUNE	48	<i>apraclonidine</i>	55	BESIVANCE	55
<i>acyclovir</i>	3, 36	<i>aprepitant</i>	46	BESREMI	48
<i>acyclovir sodium</i>	3	APRI	52	<i>betaine</i>	46
ADACEL(TDAP		APTIVUS	4	<i>betamethasone dipropionate</i>	37
ADOLESN/ADULT)(PF)	48	ARANELLE (28)	52	<i>betamethasone valerate</i>	37
<i>adefovir</i>	3	AREXVY (PF)	48	<i>betamethasone, augmented</i>	37
ADEMPAS	56	ARIKAYCE	4	BETASERON	48
ADVAIR HFA	56	<i>aripiprazole</i>	19	<i>betaxolol</i>	55
AIMOVIG AUTOINJECTOR	19	<i>armodafinil</i>	19	<i>bethanechol chloride</i>	59
AKEEGA	11	<i>asenapine maleate</i>	19	<i>bexarotene</i>	11
ALA-CORT	36	ASMANEX HFA	57	BEXSERO	48
<i>albendazole</i>	3	ASMANEX TWISTHALER ...	57	<i>bicalutamide</i>	11
<i>albuterol sulfate</i>	56, 57	<i>aspirin-dipyridamole</i>	31	BICILLIN C-R	4
<i>alclometasone</i>	36	ASSURE ID INSULIN		BICILLIN L-A	4
ALCOHOL PADS	42	SAFETY	50	BIKTARVY	4
ALECENSA	11	<i>atazanavir</i>	4	<i>bimatoprost</i>	55
<i>alendronate</i>	50	<i>atenolol</i>	31	<i>bisoprolol fumarate</i>	31
<i>alfuzosin</i>	59	<i>atenolol-chlorthalidone</i>	31	<i>bisoprolol-hydrochlorothiazide</i> ..	31
<i>aliskiren</i>	31	<i>atomoxetine</i>	19	BLISOVI FE 1.5/30 (28)	52
<i>allopurinol</i>	50	<i>atorvastatin</i>	31	BLUJEPA	4
<i>alosetron</i>	46	<i>atovaquone</i>	4	BOOSTRIX TDAP	48
ALPHAGAN P	55	<i>atovaquone-proguanil</i>	4	BOSULIF	11
<i>alprazolam</i>	19	ATROVENT HFA	57	BRAFTOVI	11
ALTAVERA (28)	52	ATTRUBY	31	BREO ELLIPTA	57
ALUNBRIG	11	AUGTYRO	11	BREYNA	57
ALYACEN 1/35 (28)	52	AUSTEDO	19	BREZTRI AEROSPHERE	57
<i>amantadine hcl</i>	3	AUSTEDO XR	19	BRIELLYN	52
<i>ambrisentan</i>	57	AUSTEDO XR TITRATION		<i>brimonidine</i>	55
<i>amikacin</i>	3	KT(WK1-4)	19	<i>brimonidine-timolol</i>	55
<i>amiloride</i>	31	AUVELITY	19	BRINSUPRI	57
<i>amiloride-hydrochlorothiazide</i> ...	31	AVIANE	52	<i>brinzolamide</i>	55
<i>amiodarone</i>	31	AVMAPKI-FAKZYNJA	11	<i>brivaracetam</i>	19

<i>bromocriptine</i>	19	CHEMET	40	COMETRIQ	11
BRUKINSA	11	<i>chlorhexidine gluconate</i>	42	COMPRO	46
<i>budesonide</i>	46, 57	<i>chloroquine phosphate</i>	5	CONSTULOSE	46
<i>budesonide-formoterol</i>	57	<i>chlorpromazine</i>	20	COPAXONE	21
<i>bumetanide</i>	32	<i>chlorthalidone</i>	32	COPIKTRA	11
<i>buprenorphine</i>	19	<i>cholestyramine (with sugar)</i>	32	COSENTYX	37
<i>buprenorphine hcl</i>	19	CHOLESTYRAMINE		COSENTYX (2 SYRINGES) ...	37
<i>buprenorphine-naloxone</i>	19	LIGHT	32	COSENTYX PEN (2 PENS)	37
<i>bupropion hcl</i>	19, 20	<i>ciclopirox</i>	37	COSENTYX UNOREADY	
<i>bupropion hcl (smoking deter)</i>	40	<i>cilostazol</i>	32	PEN	37
<i>buspironone</i>	20	CIMDUO	5	COTELLIC	11
<i>butorphanol</i>	20	<i>cinacalcet</i>	42	CREON	46
<i>cabergoline</i>	42	CINRYZE	57	CRESEMBA	5
CABOMETYX	11	<i>ciprofloxacin hcl</i>	5, 55	<i>cromolyn</i>	46, 55, 57
<i>calcipotriene</i>	37	<i>ciprofloxacin in 5 % dextrose</i>	5	CRYSSELLE (28)	52
<i>calcitonin (salmon)</i>	42	<i>ciprofloxacin-dexamethasone</i>	42	<i>cyclobenzaprine</i>	21
<i>calcitriol</i>	42	<i>citalopram</i>	20	<i>cyclophosphamide</i>	12
<i>calcium acetate(phosphat bind)</i> ..	60	CLARAVIS	37	<i>cyclosporine</i>	12, 55
CALQUENCE		<i>clarithromycin</i>	5	<i>cyclosporine modified</i>	12
(ACALABRUTINIB MAL)	11	<i>clindamycin hcl</i>	5	<i>cyproheptadine</i>	57
CAMILA	52	<i>clindamycin in 5 % dextrose</i>	5	CYRED EQ	52
<i>candesartan</i>	32	CLINDAMYCIN		CYSTAGON	59
<i>candesartan-hydrochlorothiazid</i> ..	32	PEDIATRIC	5	CYSTARAN	55
CAPLYTA	20	<i>clindamycin phosphate</i>	5, 37, 52	<i>d10 %-0.45 % sodium chloride</i> ..	40
CAPRELSA	11	<i>clindamycin-benzoyl peroxide</i>	37	<i>d2.5 %-0.45 % sodium chloride</i> ..	40
<i>captopril</i>	32	CLINIMIX 5%/D15W		<i>d5 % and 0.9 % sodium</i>	
<i>carbamazepine</i>	20	SULFITE FREE	60	<i>chloride</i>	40
<i>carbidopa-levodopa</i>	20	CLINIMIX 4.25%/D10W		<i>d5 %-0.45 % sodium chloride</i>	40
<i>carbidopa-levodopa-entacapone</i> ..	20	SULF FREE	60	<i>dabigatran etexilate</i>	32
<i>carglumic acid</i>	40	CLINIMIX 4.25%/D5W		<i>dalfampridine</i>	21
<i>carteolol</i>	55	SULFIT FREE	40	<i>danazol</i>	42
CARTIA XT	32	CLINIMIX 5%-		<i>dantrolene</i>	21
<i>carvedilol</i>	32	D20W(SULFITE-FREE)	60	DANZITEN	12
<i>caspofungin</i>	4	<i>clobazam</i>	20	<i>dapagliflozin</i>	42
CAYSTON	4	<i>clomipramine</i>	20	<i>dapsone</i>	5
<i>cefaclor</i>	4	<i>clonazepam</i>	20	DAPTACEL (DTAP	
<i>cefadroxil</i>	4	<i>clonidine</i>	32	PEDIATRIC) (PF)	48
<i>cefazolin</i>	4	<i>clonidine hcl</i>	32	<i>daptomycin</i>	5
<i>cefdinir</i>	4	<i>clopidogrel</i>	32	<i>darunavir</i>	5
<i>cefepime</i>	4	<i>clorazepate dipotassium</i>	20	<i>dasatinib</i>	12
<i>cefixime</i>	4	<i>clotrimazole</i>	5, 37	DAURISMO	12
<i>cefoxitin</i>	4	<i>clotrimazole-betamethasone</i>	37	<i>deferasirox</i>	40
<i>cefpodoxime</i>	4	<i>clozapine</i>	20, 21	DELSTRIGO	5
<i>cefprozil</i>	4	COARTEM	5	DEPO-SUBQ PROVERA 104 ..	52
<i>ceftaroline fosamil</i>	4	COBENFY	21	DESCOVY	5
<i>ceftazidime</i>	5	COBENFY STARTER PACK ..	21	<i>desipramine</i>	21
<i>ceftriaxone</i>	5	<i>colchicine</i>	50	<i>desloratadine</i>	57
<i>cefuroxime axetil</i>	5	<i>colesevelam</i>	32	<i>desmopressin</i>	42
<i>cefuroxime sodium</i>	5	<i>colestipol</i>	32	<i>desoximetasone</i>	37
<i>celecoxib</i>	20	<i>colistin (colistimethate na)</i>	5	<i>desvenlafaxine succinate</i>	21
<i>cephalexin</i>	5	COMBIGAN	55	<i>dexamethasone</i>	42
CERDELGA	42	COMBIVENT RESPIMAT	57		

<i>dexamethasone sodium phosphate</i>	55	<i>efavirenz-lamivu-tenofovir disoproxil fumarate</i>	6	<i>estradiol-norethindrone acetate</i>	52
<i>dexamethylphenidate</i>	21	ELIGARD	12	<i>ethacrynic acid</i>	33
<i>dextroamphetamine-amphetamine</i>	21	ELIGARD (3 MONTH)	12	<i>ethambutol</i>	6
<i>dextrose 10 % in water (d10w)</i>	40	ELIGARD (4 MONTH)	12	<i>ethosuximide</i>	23
<i>dextrose 5 % in water (d5w)</i>	40	ELIGARD (6 MONTH)	12	<i>etodolac</i>	23
DIACOMIT	21	ELIQUIS	32	<i>etonogestrel-ethinyl estradiol</i>	52
<i>diazepam</i>	21	ELIQUIS DVT-PE TREATMENT	32	<i>etravirine</i>	6
DIAZEPAM INTENSOL	21	ELMIRON	59	EULEXIN	12
<i>diazoxide</i>	42	<i>eltrombopag olamine</i>	32, 33	<i>everolimus (antineoplastic)</i>	12
<i>diclofenac potassium</i>	21	EMGALITY PEN	22	<i>everolimus (immunosuppressive)</i>	12
<i>diclofenac sodium</i>	22, 37, 55	EMGALITY SYRINGE	22	EVOTAZ	6
<i>dicloxacillin</i>	5	EMSAM	22	<i>exemestane</i>	12
<i>dicyclomine</i>	46	<i>emtricitabine</i>	6	EXXUA	23
<i>diflunisal</i>	22	<i>emtricitabine-tenofovir (tdf)</i>	6	<i>ezetimibe</i>	33
<i>difluprednate</i>	55	<i>emtricitabine-tenofovir disoproxil fumarate</i>	6	<i>ezetimibe-simvastatin</i>	33
<i>digoxin</i>	32	EMTRIVA	6	FABHALTA	40
<i>dihydroergotamine</i>	22	<i>enalapril maleate</i>	33	<i>famciclovir</i>	6
DILANTIN	22	<i>enalapril-hydrochlorothiazide</i>	33	<i>famotidine</i>	47
<i>diltiazem hcl</i>	32	ENBREL	50, 51	FANAPT	23
DILT-XR	32	ENBREL MINI	50	FANAPT TITRATION PACK A	23
<i>dimethyl fumarate</i>	22	ENBREL SURECLICK	51	FARXIGA	42
<i>diphenoxylate-atropine</i>	46	ENGERIX-B (PF)	48	FASENRA	57
<i>disulfiram</i>	40	ENGERIX-B PEDIATRIC (PF)	48	FASENRA PEN	57
<i>divalproex</i>	22	<i>enoxaparin</i>	33	<i>felbamate</i>	23
<i>dofetilide</i>	32	ENSACOVE	12	<i>felodipine</i>	33
<i>donepezil</i>	22	ENSKYCE	52	<i>fenofibrate</i>	33
<i>dorzolamide</i>	55	<i>entacapone</i>	22	<i>fenofibrate micronized</i>	33
<i>dorzolamide-timolol</i>	55	<i>entecavir</i>	6	<i>fenofibrate nanocrystallized</i>	33
DOVATO	5	ENTRESTO	33	<i>fentanyl</i>	23
<i>doxazosin</i>	32	ENTRESTO SPRINKLE	33	FETZIMA	23
<i>doxepin</i>	22	ENULOSE	46	FIASP FLEXTOUCH U-100 INSULIN	42
<i>doxercalciferol</i>	42	ENVARUS XR	12	FIASP PENFILL U-100 INSULIN	42
DOXY-100	5	EOHILIA	46	FIASP U-100 INSULIN	42
<i>doxycycline hyclate</i>	5, 6	EPIDIOLEX	22	<i>fidaxomicin</i>	6
<i>doxycycline monohydrate</i>	6	<i>epinephrine</i>	57	FILSUEVZ	38
DRIZALMA SPRINKLE	22	<i>eplerenone</i>	33	<i>finasteride</i>	59
<i>dronabinol</i>	46	<i>ergotamine-caffeine</i>	23	<i>finingolimod</i>	23
<i>drospirenone-ethinyl estradiol</i>	52	ERIVEDGE	12	FINTEPLA	23
<i>droxidopa</i>	40	ERLEADA	12	FIRMAGON KIT W DILUENT SYRINGE	12
DULERA	57	<i>erlotinib</i>	12	<i>flecainide</i>	33
<i>duloxetine</i>	22	ERRIN	52	<i>fluconazole</i>	6
DUPIXENT PEN	38	<i>ertapenem</i>	6	<i>fluconazole in nacl (iso-osm)</i>	6
DUPIXENT SYRINGE	38	ERY PADS	38	<i>flucytosine</i>	6
<i>dutasteride</i>	59	<i>erythromycin</i>	6, 55	<i>fludrocortisone</i>	42
<i>dutasteride-tamsulosin</i>	59	<i>erythromycin with ethanol</i>	38	<i>flunisolide</i>	57
DUVYZAT	40	<i>escitalopram oxalate</i>	23	<i>fluocinolone</i>	38
EDURANT	6	<i>eslicarbazepine</i>	23	<i>fluocinolone acetonide oil</i>	42
EDURANT PED	6	<i>esomeprazole magnesium</i>	46		
<i>efavirenz</i>	6	ESTARYLLA	52		
<i>efavirenz-emtricitabine-tenofovir disoproxil fumarate</i>	6	<i>estradiol</i>	52		

<i>fluocinolone and shower cap</i>	38	<i>glipizide-metformin</i>	43	HUMULIN R REGULAR U-100 INSULN	43
<i>fluocinonide</i>	38	GLUCAGON EMERGENCY KIT (HUMAN)	43	HUMULIN R U-500 (CONC) INSULIN	43
<i>fluocinonide-emollient</i>	38	<i>glyburide</i>	43	HUMULIN R U-500 (CONC) KWIKPEN	43
<i>fluoride (sodium)</i>	60	<i>glyburide-metformin</i>	43	<i>hydralazine</i>	33
<i>fluorometholone</i>	55	<i>glycerol phenylbutyrate</i>	41	<i>hydrochlorothiazide</i>	33
<i>fluorouracil</i>	38	<i>glycopyrrolate</i>	47	<i>hydrocodone-acetaminophen</i>	24
<i>fluoxetine</i>	23	GLYXAMBI	43	<i>hydrocortisone</i>	38, 43, 47
<i>fluoxetine (pmdd)</i>	23	GOMEKLI	13	<i>hydromorphone</i>	24
<i>fluphenazine decanoate</i>	24	<i>granisetron hcl</i>	47	<i>hydroxychloroquine</i>	7
<i>fluphenazine hcl</i>	24	<i>griseofulvin microsize</i>	7	<i>hydroxyurea</i>	13
<i>flurbiprofen</i>	24	<i>griseofulvin ultramicrosize</i>	7	<i>hydroxyzine hcl</i>	58
<i>flurbiprofen sodium</i>	55	<i>guanfacine</i>	24	HYRNUO	13
<i>fluticasone propionate</i>	38, 57, 58	GVOKE	43	<i>ibandronate</i>	51
<i>fluticasone propion-salmeterol</i> ...	58	GVOKE HYPOPEN 2-PACK	43	IBRANCE	13
<i>fluvoxamine</i>	24	GVOKE PFS 1-PACK SYRINGE	43	IBTROZI	13
<i>fondaparinux</i>	33	HADLIMA	51	IBU	24
<i>fosamprenavir</i>	6	HADLIMA PUSH TOUCH	51	<i>ibuprofen</i>	24
<i>fosfomycin tromethamine</i>	6	HADLIMA (CF)	51	<i>icatibant</i>	58
<i>fosinopril</i>	33	HADLIMA (CF) PUSH TOUCH	51	ICLEVIA	52
<i>fosinopril-hydrochlorothiazide</i> ...	33	HAILEY 24 FE	52	ICLUSIG	13
FOTIVDA	12	<i>halobetasol propionate</i>	38	<i>icosapent ethyl</i>	33
FRUZAQLA	13	<i>haloperidol</i>	24	IDHIFA	13
<i>furosemide</i>	33	<i>haloperidol decanoate</i>	24	<i>imatinib</i>	13
<i>gabapentin</i>	24	<i>haloperidol lactate</i>	24	IMBRUVICA	13
<i>galantamine</i>	24	HAVRIX (PF)	49	<i>imipenem-cilastatin</i>	7
GALLIFREY	52	HEATHER	52	<i>imipramine hcl</i>	24
GAMMAGARD LIQUID	48	<i>heparin (porcine)</i>	33	<i>imiquimod</i>	38
GAMMAGARD LIQUID ERC	49	HEPLISAV-B (PF)	49	IMKELDI	13
GAMMAGARD S-D (IGA < 1 MCG/ML)	49	HERNEXEOS	13	IMOVAX RABIES VACCINE (PF)	49
GARDASIL 9 (PF)	49	HIBERIX (PF)	49	IMPAVIDO	7
<i>gatifloxacin</i>	55	HUMALOG JUNIOR KWIKPEN U-100	43	IMVEXXY MAINTENANCE PACK	53
GATTEX 30-VIAL	47	HUMALOG KWIKPEN INSULIN	43	IMVEXXY STARTER PACK	53
GAUZE PAD	50	HUMALOG MIX 50-50 KWIKPEN	43	INCASSIA	53
GAVILYTE-C	47	HUMALOG MIX 75-25 KWIKPEN	43	INCRELEX	41
GAVILYTE-G	47	HUMALOG MIX 75-25(U-100)INSULN	43	<i>indapamide</i>	33
GAVILYTE-N	47	HUMALOG U-100 INSULIN ..	43	<i>indomethacin</i>	24
GAVRETO	13	HUMULIN 70/30 U-100 INSULIN	43	INFANRIX (DTAP) (PF)	49
<i>gefitinib</i>	13	INSULIN	43	INLURIYO	13
<i>gemfibrozil</i>	33	HUMULIN 70/30 U-100 KWIKPEN	43	INLYTA	13
GEMTESA	59	HUMALOG MIX 75-25(U-100)INSULN	43	INQOVI	13
GENERLAC	47	HUMALOG MIX 75-25(U-100)INSULN	43	INREBIC	13
GENGRAF	13	HUMALOG U-100 INSULIN ..	43	<i>insulin lispro</i>	43
<i>gentamicin</i>	6, 38, 55	HUMULIN 70/30 U-100 INSULIN	43	<i>insulin lispro protamin-lispro</i>	43
<i>gentamicin in nacl (iso-osm)</i>	6	INSULIN	43	<i>insulin syringe-needle u-100</i>	50
GENVOYA	7	HUMULIN N NPH INSULIN KWIKPEN	43	INTELENCE	7
GILOTRIF	13	HUMULIN N NPH U-100 INSULIN	43	INTRALIPID	60
<i>glatiramer</i>	24	INSULIN	43	INTROVALE	53
GLATOPA	24				
<i>glimepiride</i>	42				
<i>glipizide</i>	43				

INVEGA HAFYERA	24, 25	<i>ketoconazole</i>	7, 38	<i>levonorg-eth estrad triphasic</i>	53
INVEGA SUSTENNA	25	<i>ketorolac</i>	55	<i>levothyroxine</i>	44
INVEGA TRINZA	25	KINERET	51	LEVOXYL	44
IPOL	49	KINRIX (PF)	49	<i>lidocaine</i>	38, 39
<i>ipratropium bromide</i>	42, 58	KIONEX	41	<i>lidocaine hcl</i>	38
<i>ipratropium-albuterol</i>	58	KISQALI	14	<i>lidocaine-prilocaine</i>	39
<i>irbesartan</i>	33	KISQALI FEMARA CO-		LILETTA	53
<i>irbesartan-hydrochlorothiazide</i> ..	33	PACK	13	<i>linezolid</i>	7
ISENTRESS	7	KLOR-CON	60	<i>linezolid in dextrose 5%</i>	7
ISENTRESS HD	7	KLOR-CON M10	60	LINZESS	47
ISIBLOOM	53	KLOR-CON M15	60	LIOMNY	44
ISOLYTE S PH 7.4	60	KLOR-CON M20	60	<i>liothyronine</i>	44
ISOLYTE-P IN 5 %		KLOXXADO	25	<i>liraglutide</i>	44
DEXTROSE	60	KOSELUGO	14	<i>lisinopril</i>	34
<i>isoniazid</i>	7	KOURZEQ	42	<i>lisinopril-hydrochlorothiazide</i> ...	34
<i>isosorbide dinitrate</i>	33	KRAZATI	14	LITFULO	41
<i>isosorbide mononitrate</i>	33	KURVELO (28)	53	<i>lithium carbonate</i>	25
<i>isradipine</i>	34	KYGEVVI	41	<i>lithium citrate</i>	25
ITOVEBI	13	<i>labetalol</i>	34	LIVTENCITY	7
<i>itraconazole</i>	7	<i>lacosamide</i>	25	LOARGYS	41
<i>ivabradine</i>	34	<i>lactulose</i>	47	<i>lofexidine</i>	25
<i>ivermectin</i>	7	LAGEVRIO (EUA)	7	LOKELMA	41
IWILFIN	13	<i>lamivudine</i>	7	<i>lomustine</i>	14
IXIARO (PF)	49	<i>lamivudine-zidovudine</i>	7	LONSURF	14
JAKAFI	13	<i>lamotrigine</i>	25	<i>loperamide</i>	47
JANTOVEN	34	LANTUS SOLOSTAR U-100		<i>lopinavir-ritonavir</i>	7
JANUMET	43	INSULIN	44	<i>lorazepam</i>	25
JANUMET XR	43	LANTUS U-100 INSULIN	44	LORAZEPAM INTENSOL	25
JANUVIA	43	<i>lapatinib</i>	14	LORBRENA	14
JARDIANCE	43	<i>latanoprost</i>	55	LORYNA (28)	53
JASCAYD	58	LAZCLUZE	14	<i>losartan</i>	34
JASMIEL (28)	53	<i>leflunomide</i>	51	<i>losartan-hydrochlorothiazide</i>	34
JAVYGTOR	43	<i>lenalidomide</i>	14	<i>lovastatin</i>	34
JAYPIRCA	13	LENVIMA	14	LOW-OGESTREL (28)	53
JENTADUETO	44	LESSINA	53	<i>loxapine succinate</i>	25
JENTADUETO XR	44	<i>letrozole</i>	14	<i>lubiprostone</i>	47
JINTELI	53	<i>leucovorin calcium</i>	14	LUIZZA	53
JOENJA	41	LEUKERAN	14	LUMAKRAS	14
JULEBER	53	LEUKINE	49	LUMIGAN	56
JULUCA	7	<i>leuprolide</i>	14	LUPRON DEPOT	14
JUNEL 1.5/30 (21)	53	<i>leuprolide acetate (3 month)</i>	14	LUPRON DEPOT (3	
JUNEL 1/20 (21)	53	<i>levalbuterol hcl</i>	58	MONTH)	14
JUNEL FE 1.5/30 (28)	53	<i>levalbuterol tartrate</i>	58	<i>lurasidone</i>	25, 26
JUNEL FE 1/20 (28)	53	<i>levetiracetam</i>	25	LUTERA (28)	53
JUNEL FE 24	53	<i>levobunolol</i>	56	LYLEQ	53
JYNNEOS (PF)	49	<i>levocarnitine</i>	41	LYNPARZA	14
KALETRA	7	<i>levocarnitine (with sugar)</i>	41	LYSODREN	14
KALYDECO	58	<i>levocetirizine</i>	58	LYTGOBI	15
KARIVA (28)	53	<i>levofloxacin</i>	7	LYZA	53
KELNOR 1/35 (28)	53	<i>levofloxacin in d5w</i>	7	<i>magnesium sulfate</i>	60
KERENDIA	34	LEVONEST (28)	53	<i>malathion</i>	39
KESIMPTA PEN	25	<i>levonorgestrel-ethinyl estrad</i>	53	<i>maraviroc</i>	7

MARLISSA (28)	53	<i>mirtazapine</i>	26	<i>nilotinib hcl</i>	15
MARPLAN	26	<i>misoprostol</i>	47	<i>nilutamide</i>	15
MATULANE	15	M-M-R II (PF)	49	<i>nimodipine</i>	34
MAVYRET	7	<i>modafinil</i>	26	NINLARO	15
<i>meclizine</i>	47	MODEYSO	15	<i>nitazoxanide</i>	8
<i>medroxyprogesterone</i>	53	<i>moexipril</i>	34	<i>nitisinone</i>	41
<i>mefloquine</i>	8	<i>molindone</i>	26	NITRO-BID	34
<i>megestrol</i>	15	<i>mometasone</i>	39, 58	<i>nitrofurantoin macrocrystal</i>	8
MEKINIST	15	<i>montelukast</i>	58	<i>nitrofurantoin monohyd/m-cryst</i> ... 8	
MEKTOVI	15	<i>morphine</i>	26	<i>nitroglycerin</i>	35, 47
<i>meloxicam</i>	26	<i>morphine concentrate</i>	26	NIVESTYM	49
<i>memantine</i>	26	MOUNJARO	44	NORDITROPIN FLEXPRO ... 49	
<i>memantine-donepezil</i>	26	MOVANTIK	47	<i>norethindrone (contraceptive)</i> ... 54	
MENQUADFI (PF)	49	<i>moxifloxacin</i>	8, 56	<i>norethindrone acetate</i>	54
MENVEO A-C-Y-W-135-DIP (PF)	49	<i>moxifloxacin-sod.chloride(iso)</i> 8		<i>norethindrone ac-eth estradiol</i> ... 54	
<i>mercaptapurine</i>	15	MRESVIA (PF)	49	<i>norgestimate-ethinyl estradiol</i> ... 54	
<i>meropenem</i>	8	MULPLETA	34	NORTREL 0.5/35 (28)	54
<i>mesalamine</i>	47	MULTAQ	34	NORTREL 1/35 (21)	54
<i>mesna</i>	15	<i>mupirocin</i>	39	NORTREL 1/35 (28)	54
<i>metformin</i>	44	<i>mycophenolate mofetil</i>	15	NORTREL 7/7/7 (28)	54
<i>methadone</i>	26	<i>mycophenolate sodium</i>	15	<i>nortriptyline</i>	27
<i>methazolamide</i>	56	MYRBETRIQ	59	NORVIR	8
<i>methenamine hippurate</i>	8	<i>nabumetone</i>	26	NOVOLIN 70/30 U-100 INSULIN	44
<i>methimazole</i>	44	<i>nadolol</i>	34	NOVOLIN 70-30 FLEXPEN U-100	44
<i>methotrexate sodium</i>	15	<i>nafacillin</i>	8	NOVOLIN N FLEXPEN	44
<i>methotrexate sodium (pf)</i>	15	<i>naloxone</i>	26	NOVOLIN N NPH U-100 INSULIN	44
<i>methsuximide</i>	26	<i>naltrexone</i>	26	NOVOLIN R FLEXPEN	44
<i>methylphenidate hcl</i>	26	<i>naproxen</i>	26	NOVOLIN R REGULAR U100 INSULIN	44
<i>methylprednisolone</i>	44	<i>naproxen sodium</i>	26	NOVOLOG FLEXPEN U-100 INSULIN	44
<i>metoclopramide hcl</i>	47	<i>naratriptan</i>	26	NOVOLOG MIX 70-30 U-100 INSULIN	44
<i>metolazone</i>	34	<i>nateglinide</i>	44	NOVOLOG MIX 70-30FLEXPEN U-100	44
<i>metoprolol succinate</i>	34	NAYZILAM	26	NOVOLOG PENFILL U-100 INSULIN	44
<i>metoprolol ta-hydrochlorothiaz</i> .. 34		<i>neбиволol</i>	34	NOVOLOG U-100 INSULIN ASPART	44
<i>metoprolol tartrate</i>	34	NECON 0.5/35 (28)	54	NUBEQA	15
<i>metronidazole</i>	8, 39, 53	<i>nefazodone</i>	27	NUEDEXTA	27
<i>metronidazole in nacl (iso-os)</i> 8		NEMLUVIO	15	NUPLAZID	27
<i>metyrosine</i>	34	<i>neomycin</i>	8	NURTEC ODT	27
<i>mexiletine</i>	34	<i>neomycin-bacitracin-poly-hc</i> 56		NYAMYC	39
<i>micafungin</i>	8	<i>neomycin-bacitracin-polymyxin</i> ..56		NYLIA 1/35 (28)	54
MICONAZOLE-3	53	<i>neomycin-polymyxin b-dexameth</i>	56	NYLIA 7/7/7 (28)	54
MICROGESTIN 1.5/30 (21) 53		<i>neomycin-polymyxin-gramicidin</i> 56		<i>nystatin</i>	8, 39
MICROGESTIN 1/20 (21) 53		<i>neomycin-polymyxin-hc</i> 42, 56		NYSTOP	39
MICROGESTIN FE 1.5/30 (28)	53	NERLYNX	15	<i>octreotide acetate</i>	15
MICROGESTIN FE 1/20 (28) . 54		<i>nevirapine</i>	8		
<i>midodrine</i>	41	NEXLETOL	34		
<i>mifepristone</i>	44	NEXLIZET	34		
<i>miglustat</i>	44	NEXPLANON	54		
MILI	54	<i>niacin</i>	34		
<i>minocycline</i>	8	<i>nicardipine</i>	34		
<i>minoxidil</i>	34	NICOTROL NS	41		
		<i>nifedipine</i>	34		

ODEFSEY	8	<i>pen needle, diabetic</i>	50	<i>potassium chloride-d5-</i>	
ODOMZO	15	PENBRAYA (PF)	49	<i>0.2%nacl</i>	61
OFEV	58	<i>penciclovir</i>	39	<i>potassium chloride-d5-</i>	
<i>ofloxacin</i>	8, 42, 56	<i>penicillamine</i>	51	<i>0.9%nacl</i>	61
OGSIVEO	15	<i>penicillin g pot in dextrose</i>	9	<i>potassium citrate</i>	60
OJEMDA	15, 16	<i>penicillin g potassium</i>	9	<i>pramipexole</i>	28
OJJAARA	16	<i>penicillin v potassium</i>	9	<i>prasugrel hcl</i>	35
<i>olanzapine</i>	27	PENMENVY MEN A-B-C-W-		<i>pravastatin</i>	35
<i>olmesartan</i>	35	Y (PF)	49	<i>praziquantel</i>	9
<i>olmesartan-amlodipin-hcthiazyd</i>	35	PENTACEL (PF)	49	<i>prazosin</i>	35
<i>olmesartan-hydrochlorothiazide</i>	35	<i>pentamidine</i>	9	<i>prednisolone</i>	45
<i>olopatadine</i>	42	<i>pentoxifylline</i>	35	<i>prednisolone acetate</i>	56
<i>omega-3 acid ethyl esters</i>	35	<i>perampanel</i>	27	<i>prednisolone sodium phosphate</i>	
<i>omeprazole</i>	47	<i>perindopril erbumine</i>	35	45, 56
<i>ondansetron</i>	47	PERIOGARD	42	<i>prednisone</i>	45
<i>ondansetron hcl</i>	47	<i>permethrin</i>	39	<i>pregabalin</i>	28
ONUREG	16	<i>perphenazine</i>	27	PREMARIN	54
OPIPZA	27	PERSERIS	27	PRENATAL VITAMIN	
OPSUMIT	58	PHEBURANE	41	PLUS LOW IRON	61
OPSYNVI	58	<i>phenelzine</i>	27	PREVALITE	35
ORENCIA	51	<i>phenobarbital</i>	27	PREVYMIS	9
ORENCIA CLICKJECT	51	PHENYTEK	27	PREZCOBIX	9
ORGOVYX	16	<i>phenytoin</i>	27	PREZISTA	9
ORKAMBI	58	<i>phenytoin sodium extended</i>	27	PRIFTIN	9
ORLYNVAH	8	PIFELTRO	9	<i>primaquine</i>	9
ORSERDU	16	<i>pilocarpine hcl</i>	41, 56	<i>primidone</i>	28
<i>oseltamivir</i>	8	<i>pimozide</i>	27	PRIORIX (PF)	49
OTEZLA	51	PIMTREA (28)	54	PRIVIGEN	49
OTEZLA STARTER	51	<i>pindolol</i>	35	<i>probenecid</i>	51
OTEZLA XR	51	<i>pioglitazone</i>	44	<i>probenecid-colchicine</i>	51
OTEZLA XR INITIATION	51	<i>pioglitazone-metformin</i>	45	<i>prochlorperazine</i>	47
<i>oxacillin</i>	8	<i>piperacillin-tazobactam</i>	9	<i>prochlorperazine maleate</i>	47
<i>oxcarbazepine</i>	27	PIQRAY	16	PROCRIT	49
<i>oxybutynin chloride</i>	60	<i>pirfenidone</i>	58	PROCTO-MED HC	47
<i>oxycodone</i>	27	<i>piroxicam</i>	27	PROCTOSOL HC	47
<i>oxycodone-acetaminophen</i>	27	<i>pitavastatin calcium</i>	35	PROCTOZONE-HC	48
OZEMPIC	44	PLEGRIDY	49	PROGRAF	16
PACERONE	35	PLENAMINE	60	PROLASTIN-C	41
<i>paliperidone</i>	27	<i>podofilox</i>	39	PROLIA	51
PANRETIN	39	<i>polymyxin b sulf-trimethoprim</i>	56	<i>promethazine</i>	58
<i>pantoprazole</i>	47	<i>pomalidomide</i>	16	<i>propafenone</i>	35
<i>paricalcitol</i>	44	PORTIA 28	54	<i>propranolol</i>	35
<i>paroxetine hcl</i>	27	<i>posaconazole</i>	9	<i>propylthiouracil</i>	45
PAXLOVID	8	<i>potassium chlorid-d5-</i>		PROQUAD (PF)	50
<i>pazopanib</i>	16	<i>0.45%nacl</i>	60	PROSOL 20 %	61
PEDIARIX (PF)	49	<i>potassium chloride</i>	61	<i>protriptyline</i>	28
PEDVAX HIB (PF)	49	<i>potassium chloride in 0.9%nacl</i>	60	PULMOZYME	58
<i>peg 3350-electrolytes</i>	47	<i>potassium chloride in 5 % dex</i>	61	<i>pyrazinamide</i>	9
<i>peg3350-sod sul-nacl-kcl-asb-c</i>	47	<i>potassium chloride in lr-d5</i>	61	<i>pyridostigmine bromide</i>	28
PEGASYS	49	<i>potassium chloride in water</i>	61	<i>pyrimethamine</i>	9
<i>peg-electrolyte soln</i>	47	<i>potassium chloride-0.45 % nacl</i>	61	QINLOCK	16
PEMAZYRE	16			QUADRACEL (PF)	50

<i>quetiapine</i>	28	<i>rosuvastatin</i>	35	SOLTAMOX	16
<i>quinapril</i>	35	ROTARIX	50	SOMAVERT	45
<i>quinapril-hydrochlorothiazide</i>	35	ROTATEQ VACCINE	50	<i>sorafenib</i>	17
<i>quinidine sulfate</i>	35	ROWEEPR	29	<i>sotalol</i>	35
<i>quinine sulfate</i>	9	ROZLYTREK	16	SOTALOL AF	35
QULIPTA	28	RUBRACA	16	SPIRIVA RESPIMAT	59
QVAR REDHALER	58, 59	<i>rufinamide</i>	29	<i>spironolactone</i>	35
RABAVERT (PF)	50	RUKOBIA	9	<i>spironolacton-hydrochlorothiaz.</i>	35
<i>rabeprazole</i>	48	RYBELSUS	45	SPRINTEC (28)	54
RALDESY	28	RYDAPT	16	SPRITAM	29
<i>raloxifene</i>	51	RYTARY	29	SPS (WITH SORBITOL)	41
<i>ramelteon</i>	28	<i>sacubitril-valsartan</i>	35	SSD	39
<i>ramipril</i>	35	SAJAZIR	59	STELARA	39
<i>ranolazine</i>	35	<i>sapropterin</i>	45	STEQEYMA	39
<i>rasagiline</i>	28	SCSEMBLIX	16	STIOLTO RESPIMAT	59
RECLIPSEN (28)	54	<i>scopolamine base</i>	48	STIVARGA	17
RECOMBIVAX HB (PF)	50	SECUADO	29	<i>streptomycin</i>	9
RECORLEV	45	<i>selegiline hcl</i>	29	STRIBILD	9
RELENZA DISKHALER	9	<i>selenium sulfide</i>	39	STRIVERDI RESPIMAT	59
<i>repaglinide</i>	45	SELZENTRY	9	SUBVENITE	29
REPATHA SURECLICK	35	SEREVENT DISKUS	59	<i>sucralfate</i>	48
REPATHA SYRINGE	35	<i>sertraline</i>	29	<i>sulfacetamide sodium</i>	56
RESTASIS	56	SETLAKIN	54	<i>sulfacetamide sodium (acne)</i>	39
RESTASIS MULTIDOSE	56	<i>sevelamer carbonate</i>	41	<i>sulfacetamide-prednisolone</i>	56
RETACRIT	50	SHINGRIX (PF)	50	<i>sulfadiazine</i>	9
RETEVMO	16	SIGNIFOR	16	<i>sulfamethoxazole-trimethoprim</i>	9, 10
REVCovi	41	<i>sildenafil (pulm.hypertension)</i>	59	SULFAMYLON	39
REVUFORJ	16	SILIQ	39	<i>sulfasalazine</i>	48
REXULTI	28	<i>silodosin</i>	60	<i>sulindac</i>	29
REYATAZ	9	<i>silver sulfadiazine</i>	39	<i>sumatriptan</i>	29
REZDIFFRA	41	SIMBRINZA	56	<i>sumatriptan succinate</i>	29
REZLIDHIA	16	SIMLANDI(CF)	51	<i>sunitinib malate</i>	17
RHOPRESSA	56	SIMLANDI(CF)		SUNLENCA	10
<i>ribavirin</i>	9	AUTOINJECTOR	51	SYEDA	54
<i>rifabutin</i>	9	<i>simvastatin</i>	35	SYMPAZAN	29
<i>rifampin</i>	9	<i>sirolimus</i>	16	SYMTUZA	10
<i>riluzole</i>	41	SIRTURO	9	SYNAREL	45
<i>rimantadine</i>	9	SKYCLARYS	29	SYNJARDY	45
RINVOQ	51	SKYRIZI	39, 48	SYNJARDY XR	45
RINVOQ LQ	51	<i>sodium chloride</i>	41	SYNTHROID	45
<i>risedronate</i>	41, 51	<i>sodium chloride 0.45 %</i>	61	TABLOID	17
<i>risperidone</i>	28	<i>sodium chloride 0.9 %</i>	41	TABRECTA	17
<i>risperidone microspheres</i>	28	<i>sodium chloride 3 % hypertonic</i> .	61	<i>tacrolimus</i>	17, 39
<i>ritonavir</i>	9	<i>sodium chloride 5 % hypertonic</i> .	61	<i>tadalafil</i>	60
<i>rivastigmine</i>	28	<i>sodium oxybate</i>	29	<i>tadalafil (pulm. hypertension)</i>	59
<i>rivastigmine tartrate</i>	28	<i>sodium phenylbutyrate</i>	41	TAFINLAR	17
RIVFLOZA	60	<i>sodium polystyrene sulfonate</i>	41	TAGRISO	17
<i>rizatriptan</i>	28, 29	<i>sodium,potassium,mag sulfates</i> ..	48	TALZENNA	17
ROCKLATAN	56	<i>sofosbuvir-velpatasvir</i>	9	<i>tamoxifen</i>	17
<i>roflumilast</i>	59	SOHONOS	41	<i>tamsulosin</i>	60
ROMVIMZA	16	<i>solifenacin</i>	60	TASCENSO ODT	29
<i>ropinirole</i>	29	SOLQUA 100/33	45		

<i>tasimelteon</i>	29	<i>tramadol-acetaminophen</i>	30	VALTYA	54
<i>tazarotene</i>	39	<i>trandolapril</i>	36	<i>vancomycin</i>	10
<i>telmisartan</i>	36	<i>tranexamic acid</i>	54	VANFLYTA	17
<i>telmisartan-amlodipine</i>	36	<i>tranylcypromine</i>	30	VAQTA (PF)	50
<i>telmisartan-hydrochlorothiazid</i> ..	36	TRAVASOL 10 %	61	<i>varenicline tartrate</i>	41
TENIVAC (PF)	50	<i>travoprost</i>	56	VARIVAX (PF)	50
<i>tenofovir disoproxil fumarate</i>	10	<i>trazodone</i>	30	VAXCHORA VACCINE	50
TEPMETKO	17	TRELEGY ELLIPTA	59	VELIVET TRIPHASIC	
<i>terazosin</i>	36	TRELSTAR	17	REGIMEN (28)	54
<i>terbinafine hcl</i>	10	<i>tretinoin</i>	39, 40	VEMLIDY	10
<i>terbutaline</i>	59	<i>tretinoin (antineoplastic)</i>	17	VENCLEXTA	17
<i>terconazole</i>	54	<i>triamcinolone acetonide</i>	40, 42	VENCLEXTA STARTING	
<i>teriflunomide</i>	29	<i>triamterene-hydrochlorothiazid</i> ..	36	PACK	17
<i>teriparatide</i>	51	TRIDERM	40	<i>venlafaxine</i>	30
<i>testosterone</i>	45	<i>trientine</i>	41	VENTOLIN HFA	59
<i>testosterone cypionate</i>	45	TRI-ESTARYLLA	54	<i>verapamil</i>	36
<i>testosterone enanthate</i>	45	<i>trifluoperazine</i>	30	VERQUVO	36
<i>tetrabenazine</i>	29	<i>trifluridine</i>	56	VERSACLOZ	30
<i>tetracycline</i>	10	TRIJARDY XR	45	VERZENIO	17
THALOMID	17	TRIKAFTA	59	VESTURA (28)	54
THEO-24	59	<i>trimethoprim</i>	10	VIENVA	54
<i>theophylline</i>	59	TRI-MILI	54	<i>vigabatrin</i>	30
<i>thioridazine</i>	29	<i>trimipramine</i>	30	VIGADRONE	30
<i>thiothixene</i>	29	TRINTELLIX	30	VIJOICE	17, 18
TIADYL T ER	36	TRI-SPRINTEC (28)	54	<i>vilazodone</i>	30
<i>tiagabine</i>	29	TRIUMEQ	10	VIMKUNYA	50
TIBSOVO	17	TRIUMEQ PD	10	VIORELE (28)	54
<i>ticagrelor</i>	36	TRI-VYLIBRA	54	VIRACEPT	10
TICOVAC	50	TROPHAMINE 10 %	61	VIREAD	10
<i>tigecycline</i>	10	<i>trospium</i>	60	VITRAKVI	18
<i>timolol maleate</i>	36, 56	TRULICITY	45	VIVITROL	30
<i>tinidazole</i>	10	TRUMENBA	50	VIVJOA	10
TIVICAY	10	TRUQAP	17	VIVOTIF	50
TIVICAY PD	10	TUKYSA	17	VIZIMPRO	18
<i>tizanidine</i>	29	TURALIO	17	VONJO	18
TOBI PODHALER	10	TURQOZ (28)	54	VORANIGO	18
<i>tobramycin</i>	10, 56	TWINRIX (PF)	50	<i>voriconazole</i>	10
<i>tobramycin in 0.225 % nacl</i>	10	TYMLOS	52	VOSEVI	10
<i>tobramycin sulfate</i>	10	TYPHIM VI	50	VOWST	48
<i>tobramycin-dexamethasone</i>	56	UBRELVY	30	VRAYLAR	30
<i>tolterodine</i>	60	UNITHROID	46	VUMERITY	30
<i>tolvaptan</i>	45	UPTRAVI	36	VYFEMLA (28)	54
<i>tolvaptan (polycys kidney dis)</i>	45	<i>ursodiol</i>	48	VYLIBRA	54
<i>topiramate</i>	30	<i>ustekinumab</i>	40	<i>warfarin</i>	36
<i>toremifene</i>	17	<i>valacyclovir</i>	10	WELIREG	18
<i>toremide</i>	36	VALCHLOR	40	WINREVAIR	59
TOUJEO MAX U-300		<i>valganciclovir</i>	10	WIXELA INHUB	59
SOLOSTAR	45	<i>valproic acid</i>	30	XALKORI	18
TOUJEO SOLOSTAR U-300		<i>valproic acid (as sodium salt)</i>	30	XARELTO	36
INSULIN	45	<i>valsartan</i>	36	XARELTO DVT-PE TREAT	
TRADJENTA	45	<i>valsartan-hydrochlorothiazide</i> ...	36	30D START	36
<i>tramadol</i>	30	VALTOCO	30	XATMEP	18

XCOPRI	30	<i>zolpidem</i>	31
XCOPRI MAINTENANCE PACK	30	ZONISADE	31
XCOPRI TITRATION PACK	30	<i>zonisamide</i>	31
XDEMVY	56	ZORYVE	40
XELJANZ	52	ZOVIA 1-35 (28)	55
XELJANZ XR	52	ZTALMY	31
XELRIA FE	55	ZUBSOLV	31
XERMELO	18	ZURZUVAE	31
XGEVA	18	ZYKADIA	18
XIFAXAN	10		
XIGDUO XR	46		
XIIDRA	56		
XOFLUZA	10		
XOLAIR	59		
XOLREMDI	50		
XOSPATA	18		
XPOVIO	18		
XTANDI	18		
XULTOPHY 100/3.6	46		
YARGESA	46		
YESINTEK	40		
YF-VAX (PF)	50		
YONSA	18		
YORVIPATH	46		
YUFLYMA(CF)	52		
YUFLYMA(CF) AI CROHN'S-UC-HS	52		
YUFLYMA(CF) AUTOINJECTOR	52		
YUVAFEM	55		
ZAFEMY	55		
<i>zaleplon</i>	30		
ZARXIO	50		
ZAVZPRET	30		
ZEJULA	18		
ZELBORAF	18		
ZELSUVMI	40		
ZELVYSIA	46		
ZENPEP	48		
ZEPOSIA	31		
ZEPOSIA STARTER KIT (28-DAY)	31		
ZEPOSIA STARTER PACK (7-DAY)	31		
<i>zidovudine</i>	10		
ZILBRYSQ	31		
<i>ziprasidone hcl</i>	31		
<i>ziprasidone mesylate</i>	31		
ZIRGAN	56		
ZOKINVY	41		
ZOLINZA	18		

**Section:
Prior Authorization**

Acitretin

Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Actimmune

Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ADHD Drugs

Products Affected

- *guanfacine oral tablet extended release 24 hr*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Afinitor

Products Affected

- *everolimus (antineoplastic) oral tablet*
- *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>For everolimus only, documentation of advanced, hormone receptor-positive, HER2-negative breast cancer -AND- all of the following (1-3): 1) member is a postmenopausal woman 2) using in combination with exemestane 3) therapeutic failure or intolerance to prior treatment with letrozole or anastrozole. For everolimus only, documentation of non-functional neuroendocrine tumors -AND- all of the following (1-2): 1) disease is classified as progressive, well-differentiated, non-functional 2) disease is of gastrointestinal or lung origin that are unresectable, locally advanced or metastatic. For everolimus only, documentation of advanced renal cell carcinoma -AND- therapeutic failure or intolerance to prior treatment with sunitinib or sorafenib. For everolimus only, documentation of renal angiomyolipoma and tuberous sclerosis complex (TSC), member does not require immediate surgery. For everolimus and everolimus tablets for oral suspension, documentation of TSC with Subependymal Giant Cell Astrocytoma -AND- member is not a candidate for curative surgical resection. For everolimus only, documentation of progressive neuroendocrine tumors of pancreatic origin -AND- disease is unresectable, locally advanced or metastatic. For everolimus tablets for oral suspension only, documentation of use for adjunctive treatment of TSC-associated partial-onset seizures.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Aimovig

Products Affected

- **AIMOVIG AUTOINJECTOR
SUBCUTANEOUS AUTO-INJECTOR
140 MG/ML, 70 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-2): 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Akeega

Products Affected

- **AKEEGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. -AND- Concomitant therapy with prednisone. -AND- Concomitant therapy with a gonadotropin-releasing hormone analog or a bilateral orchiectomy, -AND- For metastatic castration-sensitive prostate cancer, disease harbors a deleterious or suspected deleterious BRCA2 mutation based on an FDA-approved test. -AND- For metastatic castration-resistant prostate cancer, BRCA mutations.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alecensa

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test - AND- One of the following (1-2): 1) metastatic disease, 2) will be used as adjuvant treatment following tumor resection of node positive or greater than or equal to 4 cm tumor(s).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alpha1-Proteinase Inhibitors

Products Affected

- **PROLASTIN-C INTRAVENOUS SOLUTION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of panacinar emphysema AND documentation of a decline in forced expiratory volume in 1 second (fev1) despite medical therapy (bronchodilators, corticosteroids) AND documentation of phenotype (pi*zz, pi*znull or pi*nullnull) associated with causing serum alpha 1-antitrypsin of less than 80 mg/dl AND documentation of an alpha 1-antitrypsin serum level below the value of 35% of normal (less than 80 mg/dl).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when furnished incident to a physician service and is not self-administered.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alunbrig

Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ampyra

Products Affected

- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure disorder, Cr Cl less than 50ml/min
Required Medical Information	Documentation of diagnosis -AND- documentation that the patient is ambulatory and has walking impairment as evidenced by one of the following. 1. Functional status score (EDSS score). 2. Timed 25-foot Walk Test (T25W).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, documentation supporting improvement in walking impairment from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Anticholinergic Meds

Products Affected

- *paroxetine hcl oral suspension*
- *paroxetine hcl oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Attestation of counseling on risk-benefit profile and potential anticholinergic adverse effects -AND- Therapeutic failure, contraindication, or intolerance to 2 other antidepressants (i.e., SSRI, SNRI).
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Anzupgo

Products Affected

- ANZUPGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic hand eczema (CHE) -AND- symptoms persisting for more than 3 months or recurring at least twice within a 12-month timeframe -AND- therapeutic failure, contraindication, or intolerance to 1 generic, on-formulary, medium/high/super-high potency topical corticosteroid.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Arikayce

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Mycobacterium avium complex lung disease -AND- Attestation of not achieving negative sputum cultures despite at least 6 months with a multidrug background regimen containing 2 of the following: 1) macrolide 2) rifamycin or 3) ethambutal -AND- Arikayce will be used in conjunction with a background multidrug regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive sputum cultures -OR- negative sputum cultures for insufficient period of time (e.g. less than 12 months).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ATTR-CM drugs

Products Affected

- ATTRUBY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitantly with transthyretin-lowering agents
Required Medical Information	Documentation of cardiomyopathy of transthyretin-mediated amyloidosis (ATTR-CM) with amyloid deposits on cardiac biopsy or scintigraphy with heart contralateral greater than 1.5 (Grade III) or visual grade 2 to 3 - AND- Cardiac involvement supported by cardiac magnetic resonance, echocardiography or serum cardiac biomarker (e.g. B-type natriuretic peptide, cardiac troponin) -AND- ATTR-CM type is wild-type or variant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement or delayed disease progression from baseline demonstrated by 6-minute walk test, cardiac function (e.g. LVEF, NYHA class), Kansas City Cardiomyopathy Questionnaire-Overall Summary, number of cardiovascular-related hospitalizations or serum cardiac biomarkers (e.g. B-type natriuretic peptide, cardiac troponin)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Atypical Antipsychotics

Products Affected

- *aripiprazole oral solution*
- *aripiprazole oral tablet, disintegrating*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Aubagio

Products Affected

- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as fingolimod, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Augtyro

Products Affected

- AUGTYRO ORAL CAPSULE 160 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive. Documentation of solid tumors - AND- disease harbors a NTRK gene fusion -AND- one of the following (1-2): 1) disease is locally advanced or metastatic, or 2) surgical resection is likely to result in severe morbidity -AND- one of the following (3-4): 3) disease has progressed following treatment, or 4) the member has no satisfactory alternative therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Austedo

Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG **ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG**
- AUSTEDO XR
- AUSTEDO XR TITRATION KT(WK1-4)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3) 1) Chorea associated with Huntington's disease 2) In patients with comorbid depression, attestation of adequate treatment for depression is required. 3) Attestation that patient is not actively suicidal. -OR- 4) Tardive Dyskinesia
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Auvelity

Products Affected

- AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Avmapki-Fakzynja

Products Affected

- **AVMAPKI-FAKZYNJA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent low-grade serous ovarian cancer (LGSOC), disease is KRAS-mutated -AND- member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ayvakit

Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable or metastatic gastrointestinal stromal tumor -AND- tumors harbor a PDGFRA exon 18 mutation. Documentation of aggressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm, mast cell leukemia, or indolent systemic mastocytosis -AND- platelet count greater than or equal to $50 \times 10^9/L$.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Bafiertam

Products Affected

- **BAFIERTAM**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Balversa

Products Affected

- **BALVERSA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) FGFR3 mutation status as detected by an FDA approved test 2) disease progression on or after at least on prior line of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Banzel

Products Affected

- *rufinamide*

PA Criteria	Criteria Details
Exclusion Criteria	Patients with familial short QT syndrome
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Benlysta

Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>For diagnosis of active systemic lupus erythematosus (SLE), both of the following (1-2): 1) Documentation of positive anti-nuclear antibody (ANA) titer (greater than or equal to 1:80) or anti-double-stranded DNA antibody (anti-dsDNA) greater than or equal to 30IU/mL, 2) member will continue to receive concomitant standard of care treatment with use of at least one of the following (alone or in combination) (a-c): a) corticosteroids (e.g. prednisone), b) antimalarials (e.g. hydroxychloroquine), c) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate). For diagnosis of active lupus nephritis (LN), one of the following (3 or 4): 3) Diagnosis confirmed by renal biopsy, 4) Contraindication to renal biopsy, laboratory findings specific to LN (e.g., elevated serum creatinine, abnormal urine analysis, decreased eGFR). -AND- Member will continue to receive concomitant standard of care treatment which includes corticosteroids (e.g. prednisone) with at least one of the following (d or e): d) mycophenolate for induction followed by mycophenolate for maintenance, e) cyclophosphamide for induction followed by azathioprine for maintenance.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For SLE reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with corticosteroids, antimalarials, or immunosuppressives. For active LN reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with mycophenolate or azathioprine. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Besremi

Products Affected

- **BESREMI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Bosulif

Products Affected

- **BOSULIF ORAL CAPSULE 100 MG, 50 MG**
- **BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy. For pediatric patients 1 year of age and older, one of the following (3, 4): 3) newly diagnosed PH-positive CML in the chronic phase 4) diagnosis of Ph-positive CML in the chronic phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Bosutinib capsules and 18 years of age or older, inability to swallow tablets is required. For Bosutinib 100mg capsules and pediatric 1 year of age or older, inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Braftovi

Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	Use in wild-type BRAF melanoma, wild-type BRAF CRC, or wild-type BRAF NSCLC.
Required Medical Information	Documentation of diagnosis. For metastatic colorectal cancer (mCRC) and using in combination with cetuximab, all of the following (1-3): 1) BRAF V600E mutation status as detected by an FDA-authorized test 2) using in combination with cetuximab 3) member has received prior therapy for CRC. For mCRC and using in combination with cetuximab and fluorouracil-based chemotherapy, all of the following (4-5): 4) BRAF V600E mutation status, as detected by an FDA-authorized test 5) using in combination with cetuximab and fluorouracil-based chemotherapy. For unresectable or metastatic melanoma, all of the following (6-7): 6) BRAF V600E or V600K mutation status, as detected by an FDA-authorized test 7) using in combination with binimetinib. For metastatic non-small cell lung cancer, all of the following (8-9): 8) BRAF V600E mutation status, as detected by an FDA-authorized test 9) using in combination with binimetinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Brinsupri

Products Affected

- **BRINSUPRI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of non-cystic fibrosis bronchiectasis (NCFB) confirmed by computed tomography (CT) -AND- at least 1 of the following symptoms consistent with bronchiectasis (1-8): 1) cough on most days of the week, 2) chronic sputum production, 3) history of recurrent respiratory infections, 4) dyspnea, 5) wheezing, 6) rhinosinusitis, 7) hemoptysis, 8) recurrent pleurisy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Brukinsa

Products Affected

- BRUKINSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma (MCL), previous treatment with at least 1 prior therapy. For marginal zone lymphoma (MZL), previous treatment with at least 1 anti-CD20-based regimen. For follicular lymphoma (FL), using in combination with obinutuzumab and previous treatment with at least 2 prior lines of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Buphenyl

Products Affected

- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cabometyx

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-6): 1) diagnosis of advanced renal cell carcinoma (RCC), 2) diagnosis of advanced RCC and using as a first-line treatment in combination with nivolumab, 3) member has previously been treated with sorafenib for hepatocellular carcinoma, 4) member has experienced disease progression following prior VEGFR-targeted therapy and is either radioactive iodine-refractory or is ineligible for radioactive iodine therapy for locally advanced or metastatic differentiated thyroid cancer, 5) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated pancreatic neuroendocrine tumors (pNET), 6) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated extra-pancreatic neuroendocrine tumors (epNET)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Calquence

Products Affected

- CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma, member has received at least one prior therapy -OR- all of the following (1-3): 1) member has not received prior therapy for MCL, 2) member is ineligible for autologous hematopoietic stem cell transplantation (HSCT), and 3) using in combination with bendamustine and rituximab.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Caplyta

Products Affected

- CAPLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For schizophrenia, bipolar I disorder, and bipolar II disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Caprelsa

Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic or progressive medullary thyroid cancer in patients with unresectable locally advanced or metastatic disease.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Carbaglu

Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use as an adjunct therapy for acute hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA) -OR- maintenance therapy for chronic hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cayston

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cerdelga

Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of type 1 Gaucher disease confirmed by the following A. or B. A) With one of the following symptoms (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm ³ . 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B) Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles -AND- Documentation of CYP2D6 metabolizer status (e.g. intermediate metabolizer).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CF drugs

Products Affected

- **TOBI PODHALER**
- *tobramycin in 0.225 % nacl*
- *tobramycin inhalation*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization of tobramycin products, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cialis

Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of benign prostatic hyperplasia (BPH) and trial/failure of at least two alternative medications in the following classes (alpha-1 adrenergic blockers and/or 5-alpha reductase inhibitors)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cinryze

Products Affected

- CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	<p>For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) due to deficiency of C1INH (type I & II) with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) with normal C1INH (type III) with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE, FXII mutation, angiotensin-1 mutation, plasminogen mutation, kininogen-1 mutation, myoferlin mutation, or heparin sulfate-glucosamine 3-O-sulfotransferase 6 mutation. 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cobenfy

Products Affected

- **COBENFY**
- **COBENFY STARTER PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cometriq

Products Affected

- **COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressive, metastatic medullary thyroid cancer
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Copiktra

Products Affected

- **COPIKTRA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is no longer responding or is intolerant to at least 2 prior therapies for chronic lymphocytic leukemia and small lymphocytic leukemia.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Corlanor

Products Affected

- *ivabradine oral tablet 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) Normal sinus rhythm, 2) Resting heart rate greater than or equal to 70 beats per minute, 3) For adults, left ventricular ejection fraction less than or equal to 35 percent, 4) For adults, concurrent use of 1 beta-blocker used for treatment of heart failure (i.e. bisoprolol, carvedilol, metoprolol succinate) or therapeutic failure, intolerance, or contraindication to the maximum tolerated dose of 2 beta-blockers used for treatment of heart failure.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cosentyx

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs or all are contraindicated.
Age Restrictions	Pending CMS Review
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication. For hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cotellic

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following for unresectable or metastatic melanoma (1-2): 1) BRAF V600E or V600K mutation status 2) Concomitant therapy with vemurafenib. For cobimetinib monotherapy, documentation of histiocytic neoplasms.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cysteamine Ophthalmic Drops

Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystinosis -AND- Attestation of accumulation of corneal cystine crystals
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Danziten

Products Affected

- DANZITEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib -OR- member is newly diagnosed in the chronic phase.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Daraprim

Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For primary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 100 cells/mm ³ -AND- Toxoplasma IgG positive -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For secondary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 200 cells/mm ³ . For secondary prophylaxis of cystoisosporiasis with CD4 count less than 200 cells/mm ³ or acute cystoisosporiasis infection: failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For primary prophylaxis of Pneumocystis jirovecii pneumonia: diagnosis of HIV - AND- CD4 count less than 200 cells/mm ³ -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Daurismo

Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of newly diagnosed Acute Myeloid Leukemia -AND- Used in combination with cytarabine -AND- At least one comorbidity that preclude use of intensive induction chemotherapy defined as one of the following: 1) Age greater than or equal to 75 2) Severe cardiac or pulmonary comorbidity 3) Reduced renal function 4) Hepatic impairment 5.) Physician attests patient is not a candidate for intensive induction therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Deferasirox

Products Affected

- *deferasirox oral tablet, dispersible*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic iron overload due to blood transfusions, transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -And- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw). For Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia (NTDT) Syndrome, LIC of at least 5 mg Fe/g dw -AND- serum ferritin greater than 300 mcg/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of chronic iron overload due to blood transfusion, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw. For reauthorization of chronic iron overload in NTDT syndrome, LIC greater than or equal to 3 mg Fe/g dw.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Diacomit

Products Affected

- **DIACOMIT ORAL CAPSULE 250 MG, 500 MG**
- **DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravets syndrome - AND- Used in combination with clobazam
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Dihydroergotamine

Products Affected

- *dihydroergotamine nasal*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- requires non-oral route of administration -AND- therapeutic failure or intolerance to generic sumatriptan nasal spray.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Drizalma

Products Affected

- **DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG, 30 MG, 40 MG, 60 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow tablets/capsules. For fibromyalgia, members must also have widespread bilateral pain above and below the waist for greater than 3 months duration -AND- At least 1 fibromyalgia-related symptom (e.g., cognitive impairment, fatigue, sleep disturbance, neurologic symptoms, exercise intolerance).
Age Restrictions	Deny if less than 18 years of age in the treatment of fibromyalgia, major depressive disorder, diabetic peripheral neuropathy and chronic musculoskeletal pain -OR- if less than 7 years of age in generalized anxiety disorder
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Dupixent

Products Affected

- **DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML**
- **DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Mod/Sev atopic dermatitis (AD): failure/intolerance to 1 topical corticosteroid (TCS) or, if 2+ yrs, topical calcineurin inhibitor (e.g. tacrolimus, etc) OR, if sev AD, topical tx are precluded d/t body surface area involvement or severely damaged skin. TCS in combo w/Dupixent for mod/sev AD is permitted & allows coverage. Mod/sev asthma: hx of 2+ exacerbations req oral/inj corticosteroids in past 12mos or 1+ exacerbation req hospitalization in past 12mos AND blood eosinophils of at least 150cells/uL or current daily/alternate-day oral corticosteroid (OCS) AND inadequate control despite regular med/high-dose inhaled corticosteroids (ICS) & 1 or more add'l asthma controller (e.g. LABA, leukotriene receptor antagonist [LTRA], theophylline) +/- OCS, unless intolerant or contraindicated (c/i) to all AND will continue med/high dose ICS & 1 or more add'l asthma controller +/- OCS. Chronic rhinosinusitis w/ nasal polyposis (CRSwNP): failure, intolerance, or c/i to intranasal corticosteroid. Eosinophilic esophagitis: esophageal eosinophils of 15eos/hpf or higher on esophageal biopsy AND clinical esophageal dysfxn symptoms. Prurigo nodularis. COPD: blood eosinophils of at least 300cells/uL or current daily/alternate-day OCS AND inadequate control despite 3+ mos of regular LAMA, LABA, & ICS tx, unless intolerant or c/i to all. Chronic Spontaneous Urticaria: failure, intolerance, or c/i to 1 2nd-gen H1 antihistamine at max rec dose (e.g. cetirizine, etc). Bullous pemphigoid: failure, intolerance, or c/i on high/super high potency TCS or OCS. Allergic fungal rhinosinusitis: hx of sinonasal surgery AND diagnosis confirmed by at least 2 of the following: Type 1 hypersensitivity to fungus (history, in vitro IgE, or skin testing), positive fungal stain from surgical sinus contents, presence of eosinophilic mucin w/o sinus tissue invasion, presence of CRSwNP, or characteristic CT signs</p>
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For atopic dermatitis reauthorization, attestation of positive clinical response to therapy. For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For EoE reauthorization, attestation of histological remission (less than 15 eos/hpf) on esophageal biopsy or reduced severity or frequency of clinical symptoms of esophageal dysfunction. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline. For COPD reauthorization, attestation of one of the following is required (1-4): 1) reduction in COPD symptoms, 2) improvement in exercise tolerance, 3) delayed disease progression, or 4) reduction in the number of COPD exacerbations. For CSU reauthorization, improved CSU symptoms. For BP reauthorization, 1 of the following is required (1-4): 1) attestation of disease control, 2) reduction in number of relapses, 3) improvement in BP symptoms, or 4) reduction in oral corticosteroid use. For allergic fungal rhinosinusitis reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Duvyzat

Products Affected

- DUVYZAT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy with pathogenic mutation in the dystrophin gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

EGFR Tyrosine Kinase Inhibitors

Products Affected

- *erlotinib*
- **GILOTRIF**

PA Criteria	Criteria Details
Exclusion Criteria	Afatinib products: tumors with resistant EGFR mutations. Erlotinib products: use in NSCLC tumors with mutations other than those in FDA-approved indications. Use in combination with platinum based chemotherapy.
Required Medical Information	For afatinib, documentation of metastatic non-small cell lung cancer (NSCLC) -AND- one of the following, as detected by an FDA-approved test (1-3): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutation 3) disease harbors non-resistant EGFR mutation (i.e., S768I, L861Q, G719X) -OR- documentation of squamous metastatic NSCLC and member has experienced progression on platinum-based chemotherapy. For erlotinib, documentation of metastatic NSCLC -AND- one of the following, as detected by an FDA-approved test (1-2): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutations -OR- documentation of locally advanced, unresectable or metastatic pancreatic cancer -AND- all of the following (1-2): 1) using erlotinib as first-line therapy 2) using in combination with gemcitabine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Emgality

Products Affected

- **EMGALITY PEN**
- **EMGALITY SYRINGE
SUBCUTANEOUS SYRINGE 120
MG/ML, 300 MG/3 ML (100 MG/ML X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). For episodic cluster headaches, characterized by severe or very severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes when left untreated -AND- Attack frequency of at least one attack every other day during the cluster period.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency is required. For reauthorization of cluster headache, attestation of reduction in the number of mean weekly cluster headaches from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Ensacove

Products Affected

- ENSACOVE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally advanced or metastatic non-small cell lung cancer (NSCLC) -AND- Disease is ALK-positive, as detected by an FDA-approved test -AND- Not previously received an ALK inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Entresto Sprinkle

Products Affected

- ENTRESTO SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of symptomatic heart failure with left ventricular systolic dysfunction -AND- weight less than 40kg, or weight greater than or equal to 40kg and less than 50kg and one of the following (1 or 2): 1) therapeutic failure or intolerance to Entresto (sacubitril/valsartan) tablets, or 2) inability to swallow tablets.
Age Restrictions	Deny if greater than 17 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, all of the following are required (1-3): 1) attestation of positive clinical response to therapy, 2) member is younger than 18 years of age, 3) weight less than 40kg, or weight greater than or equal to 40kg and less than 50kg and one of the following (4 or 5): 4) intolerance to Entresto (sacubitril/valsartan) tablets, or 5) inability to swallow tablets.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Eohilia

Products Affected

- EOHILIA

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 12 weeks
Required Medical Information	Diagnosis of eosinophilic esophagitis -AND- esophageal eosinophil count greater than or equal to 15 eos/hpf on esophageal biopsy -AND- clinical symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, gastroesophageal reflux)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Epclusa

Products Affected

- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance. For Brand Epclusa, the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Epidiolex

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Lennox-Gastaut, Dravet syndromes or Tuberous Sclerosis Complex. For Lennox-Gastaut, trial and failure or intolerance of at least two standard of care treatments (e.g. lamotrigine, clobazam). For Lennox-Gastaut and Dravet syndromes, treatment is in combination with other conventional agents.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Eprontia

Products Affected

- *topiramate oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow pills or food. For migraine, therapeutic failure, intolerance, or contraindication to two generic preventive migraine therapies. For partial onset seizures, primary generalized tonic-clonic seizures, or adjunctive treatment of Lennox-Gastaut Syndrome, therapeutic failure, contraindication, or intolerance to two generic anti-epileptic drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ergotamine

Products Affected

- *ergotamine-caffeine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use to abort a vascular headache -AND- therapeutic failure, intolerance, or contraindication to a generic triptan -OR- documentation of use to prevent a vascular headache -AND- therapeutic failure, intolerance, or contraindication to generic prophylactic migraine medication (e.g., topiramate, propranolol, timolol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Erivedge

Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if disease is locally advanced all of the following: 1) disease has recurred following surgery, or is not a candidate for surgery, 2) is not a candidate for radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Erleada

Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member meets one of the following (1 or 2) 1. Documentation of use in combination with a GnRH analog -OR- 2. The member has had a bilateral orchiectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Escitalopram Capsule

Products Affected

- *escitalopram oxalate oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of major depressive disorder or generalized anxiety -AND- Therapy has been initiated with escitalopram 10mg or member has been receiving and is experiencing unfavorable tolerability to escitalopram 20mg -AND- Intolerance to generic escitalopram tablets -AND- Therapeutic failure, intolerance or contraindication to at least one other antidepressant (e.g., SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Etanercept

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE
25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Ankylosing Spondylitis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Psoriatic Arthritis or Less than 4 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	Therapeutic failure or intolerance to 1 of the following preferred adalimumab products is required, when being utilized for the same medically accepted indication and age: Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Exxua

Products Affected

- **EXXUA ORAL TABLET EXTENDED RELEASE 24 HR**
- **EXXUA ORAL TABLET, EXT REL 24HR DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fabhalta

Products Affected

- FABHALTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis. For paroxysmal nocturnal hemoglobinuria (PNH), meets one of the following (1 or 2): 1) PNH mutant clones confirmed by flow cytometry, or 2) glycosylphosphatidylinositol-anchored proteins (GPI-AP)-deficient polymorphonuclear cells (PMNs) confirmed by flow cytometry -AND- meets one of the following (3-6): 3) anemia secondary to PNH (e.g. hemoglobin less than 10.5 g/dL with symptoms of anemia), 4) elevated lactate dehydrogenase (LDH) greater than or equal to 1.5 times the upper limit of normal, 5) history of a thromboembolic event, or 6) clinical findings of systemic complications (e.g. fatigue, hemoglobinuria, abdominal pain, dyspnea, dysphagia, erectile dysfunction, history of blood cell transfusion due to PNH) -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)) unless initially cross-titrating. For diagnosis of primary immunoglobulin A nephropathy (IgAN) confirmed by biopsy, member is at risk for rapid disease progression evidenced by one of the following (7 or 8): 7) Urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g or 8) Proteinuria greater than or equal to 1 g/day -AND- has experienced therapeutic failure, contraindication, or intolerance to a maximally tolerated dose of one of the following (a or b): a) Angiotensin converting enzyme (ACE) inhibitor, b) Angiotensin receptor blocker (ARB) -AND- experienced therapeutic failure, contraindication, or intolerance to one of the following (c or d): c) Filspari (sparsentan) or d) Tarpeyo (budesonide). For Complement 3 Glomerulopathy (C3G) confirmed by biopsy, meets all of the following (13-14): 13) UPCR greater than or equal to 1.0 g/g, 14) currently therapy of the member should be on the maximally tolerated dose of one of the following (e or f): e) ACE-inhibitor, f) ARB.</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of PNH, attestation of positive clinical response defined as one of the following (1-3): 1) hemoglobin stabilization or increase from baseline, 2) decrease in transfusions from baseline, or 3) decrease in LDH levels from baseline or reduction of hemolysis -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)). For reauthorization of IgAN, reduction in urine protein-to-creatinine ratio (UPCR) or proteinuria from baseline. For reauthorization of C3G, reduction in UPCR from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fanapt

Products Affected

- FANAPT
- FANAPT TITRATION PACK A

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fasenra

Products Affected

- **FASENRA PEN**
- **FASENRA SUBCUTANEOUS SYRINGE**
10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe asthma and all of the following (1-4): 1) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos 2) blood eosinophils greater than or equal to 150cells/uL within the past 6 weeks or greater than or equal to 300cells/uL within the past 12 mos in without other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) 3) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), with or without oral corticosteroids (OCS), unless intolerant or contraindicated to all 4) will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, with or without OCS -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) and all of the following (5-6): 5) history of relapsing or refractory disease 6) will be receiving standard of care while on Fasenra therapy with glucocorticoid treatment (e.g. prednisone or prednisolone), with or without immunosuppressive therapy (e.g. cyclosporine, leflunomide, azathioprine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For severe asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EPGA reauthorization, attestation of one of the following is required (5-8): 5) reduction in the frequency and/or severity of relapses, 6) reduction or discontinuation of doses of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fecal Microbiota Products

Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a recent diagnosis of Clostridioides difficile infection (CDI) confirmed by a positive stool test -AND- Classified as recurrent (e.g., at least 2 CDI episodes) -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	For reauthorization, attestation of recurrent CDI episodes after administration of the initial fecal microbiota product -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Fetzima

Products Affected

- **FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)** **CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG**
- **FETZIMA ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder and trial and failure of two other generic antidepressants.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Filsuvez

Products Affected

- **FILSUVEZ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB) -AND- at least one open wound.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement in targeted wound(s) - AND- member requires additional courses of treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Fintepla

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravet syndrome or Lennox-Gastaut syndrome. For Lennox-Gastaut syndrome, therapeutic failure, contraindication, or intolerance to at least 2 standard of care treatments (e.g. lamotrigine, clobazam).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Firazyr

Products Affected

- *icatibant*
- SAJAZIR

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) due to deficiency of C1INH (type I & II) with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) with normal C1INH (type III) with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE, FXII mutation, angiotensin-converting enzyme mutation, plasminogen mutation, kininogen-1 mutation, myoferlin mutation, or heparin sulfate-glucosamine 3-O-sulfotransferase 6 mutation. 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Firmagon

Products Affected

- **FIRMAGON KIT W DILUENT SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Forteo

Products Affected

- *teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of teriparatide will be limited to a coverage duration of 24 months in the absence of provider attestation that the member remains at or has returned to having a high risk for fracture
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Fotivda

Products Affected

- FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received at least two prior systemic therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fruzaqla

Products Affected

- **FRUZAQLA ORAL CAPSULE 1 MG, 5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received previous threatment with a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy -AND- member has received previous treatment with an anti-VEGF therapy -AND- one of the following, if member is RAS wild-type (1-2): 1) member has received previous therapy with an anti-EGFR therapy 2) prescriber attests that treatment with an anti-EGFR therapy would not be medically appropriate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gattex

Products Affected

- **GATTEX 30-VIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of short bowel syndrome (SBS) having less than 200 cm of functional small bowel -AND- Dependence on parenteral/intravenous nutrition -AND- weight of at least 10 kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline - AND- 1 of the following (1-2): 1) continued dependence on parenteral nutrition/intravenous nutritional support, 2) not receiving parenteral nutrition/intravenous support and experienced therapeutic failure upon trial of discontinuation of parenteral nutrition/intravenous support.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gavreto

Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, disease is RET fusion-positive as detected by an FDA approved test. For advanced or metastatic thyroid cancer, all of the following (1-2): 1) disease is RET fusion-positive 2) if radioactive iodine is appropriate, the member is radioactive iodine-refractory.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gilenya

Products Affected

- *fingolimod*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Glatiramer

Products Affected

- **COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**
 - *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- **GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gleevec

Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult aggressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvant treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Gleostine

Products Affected

- *lomustine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or metastatic brain tumor(s) -AND- member has previously received surgical and/or radiotherapeutic procedure(s). Documentation of Hodgkin's lymphoma -AND- all of the following (1-2): 1) using lomustine in combination with other chemotherapies 2) member has experienced disease progression with initial chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GLP1

Products Affected

- *liraglutide*
 - **MOUNJARO**
 - **OZEMPIC SUBCUTANEOUS**
 - **RYBELSUS**
- **TRULICITY**

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gomekli

Products Affected

- GOMEKLI ORAL CAPSULE 1 MG, 2 MG
- GOMEKLI ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) -AND- presence of symptomatic plexiform neurofibromas (PN) not amenable to complete resection.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Growth Hormone

Products Affected

- **NORDITROPIN FLEXPRO
SUBCUTANEOUS PEN INJECTOR 10
MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML
(10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, member has open epiphyses, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria. Additionally for growth failure due to chronic kidney disease, glomerular filtration rate is less than 89ml/min per 1.73m ² . For HIV wasting and cachexia, Concurrent use of antiretroviral therapy -AND- weight loss from baseline of at least 10% in the past 12 months or at least 5% in the past 6 months. For short bowel syndrome, receiving management for short bowel syndrome, including specialized nutritional support -AND- less than 200 cm of functional small bowel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and member has open epiphyses, as applicable to meet standard continuation of therapy guidelines. For reauthorization of HIV wasting and cachexia, attestation of increase in weight from start of therapy. For reauthorization of short bowel syndrome, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Hernexeos

Products Affected

- **HERNEXEOS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For non-squamous non-small cell lung cancer (NSCLC), all of the following (1-2): 1) disease is unresectable or metastatic, 2) disease harbors HER2 (ERBB2) tyrosine kinase domain activating mutations, as detected by an FDA-authorized test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

High-risk meds

Products Affected

- *amitriptyline*
- *benztropine oral*
- *clomipramine*
- *cyproheptadine oral tablet*
- *doxepin oral capsule*
- *doxepin oral concentrate*
- *doxepin oral tablet*
- *hydroxyzine hcl oral tablet*
- *imipramine hcl*
- *promethazine oral tablet*
- *trimipramine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For all medications subject to this PA group, the following information (1 through 3) is required: 1. Documentation of diagnosis 2. Explanation of risk-benefit profile favoring use of the high-risk medication 3. Attestation of an intent to monitor and address treatment-related adverse events. For target tricyclic antidepressants (TCAs), in addition to criteria 1 through 3 above, trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs for the same indication, if available, is required (e.g. SSRIs and SNRIs). If using a TCA for a medically-accepted indication not shared by the safer alternatives listed, then no trial of alternatives is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Doxepin doses less than or equal to 6 mg per day will receive automatic approval.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

High-risk meds phenobarbital

Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia
Required Medical Information	For use in seizures the following are required: 1. Explanation of risk-benefit profile favoring use of the high-risk medication 2. Attestation of an intent to monitor and address treatment-related adverse events. 3. For new starts, the trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs used for seizures (e.g. carbamazepine, lamotrigine) is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Seizure disorders
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Hyruno

Products Affected

- **HYRNUO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For non-squamous non-small cell lung cancer (NSCLC), all of the following (1-3): 1) disease is locally advanced or metastatic, 2) disease harbors HER2 (ERBB2) tyrosine kinase domain activating mutations, as detected by an FDA-approved test, and 3) member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ibrance

Products Affected

- **IBRANCE ORAL CAPSULE 125 MG**
- **IBRANCE ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of HR-positive, HER2-negative advanced or metastatic breast cancer -AND- meets one of the following (1. or 2.): 1) documentation of use with an aromatase inhibitor as initial endocrine-based therapy -OR- 2) documentation of use with fulvestrant in patients with disease progression following endocrine therapy. Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (3-6): 3) disease is HR-positive, HER2-negative 4) disease is PIK3CA-mutated, as detected by an FDA-approved test 5) the member is using in combination with inavolisib and fulvestrant 6) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ibuprofen

Products Affected

- **IBTROI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iclusig

Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of newly-diagnosed chronic phase CML
Required Medical Information	Documentation of T3151+ chronic myeloid leukemia (CML) -OR- documentation of chronic phase CML and member has experienced resistance or intolerance to at least two prior kinase inhibitors -OR- documentation of accelerated phase or blast phase CML and no other kinase inhibitor is indicated -OR- member is using Iclusig as monotherapy and meets one of the following (1-2): 1) documentation of T3151+ acute lymphoblastic leukemia (ALL) 2) documentation of Ph+ ALL and no other tyrosine kinase inhibitor therapy is indicated -OR- member has newly diagnosed Ph+ ALL and is using Iclusig in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Idhifa

Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IG

Products Affected

- **GAMMAGARD LIQUID**
- **GAMMAGARD LIQUID ERC**
- **GAMMAGARD S-D (IGA)**
- **PRIVIGEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis. For select diagnoses the following apply 1) For Myasthenia Gravis Syndrome, documentation that the patient is refractory to at least one other standard therapy (e.g., cholinesterase inhibitors, corticosteroids, azathioprine) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 2) For Multiple Sclerosis, patient is refractory to at least one other standard therapy (e.g., interferons) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 3) For B-cell CLL, associated with recurrent bacterial infections OR with Associated Hypogammaglobulinemia defined as IgG level less than 600mg/dL or evidence of a specific antibody deficiency. 4) For Bone Marrow Transplantation, the member is 20 years of age or older and within the first 100 days after transplantation. 5) For Dermatomyositis/Polymyositis, trial and failure, intolerance, or contraindication to at least one standard first line therapy (i.e. corticosteroids or immunosuppressants). 6) For Pediatric HIV, the patient is less than 13 y.o. who have entry CD4 lymphocyte count greater than or equal to 200/mcl and IgG less than 400mg/dL OR recurrent bacterial infections. 7) For Guillain-Barre syndrome, impaired function by objective assessment and/or objective findings on physical exam at the time of initial therapy and IVIG therapy must be initiated within 2 weeks of symptom onset. 8) For Autoimmune Mucocutaneous Blistering Diseases (e.g. Stevens-Johnson Syndrome), trial and failure, intolerance, or contraindication to at least one conventional therapy (e.g. corticosteroids) or the patient has rapidly progressive disease in which a clinical response could not be affected quickly enough using conventional agents.</p>
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	Covered under Part B when administered in the home to a member with a diagnosis of primary immunodeficiency disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Myasthenia Gravis syndrome, Multiple Sclerosis, Polymyositis, Bone Marrow Transplant, Pediatric HIV, Guillain-Barre syndrome, Autoimmune Mucocutaneous Blistering Diseases
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Imbruvica

Products Affected

- **IMBRUVICA ORAL CAPSULE 140 MG, 70 MG**
- **IMBRUVICA ORAL SUSPENSION**
- **IMBRUVICA ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic graft versus host disease, previous treatment with at least 1 prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For suspension, inability to swallow oral tablets or oral capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Imkeldi

Products Affected

- **IMKELDI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult aggressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvant treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of an inability to swallow oral tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Increlex

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe primary IGF-1 deficiency and all of the following: 1) Normal or elevated response (greater than 10 ng/ml) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 2) Serum IGF-1 concentration that is less than or equal to three (3) standard deviations below the normal value based on laboratory reference range. 3) Height less than or equal to three (3) standard deviations below normal (at or below the third percentile for gender and age). 4) Member has open epiphyses. -OR- Documentation of diagnosis of growth hormone deficiency caused by gene deletion and all of the following: 1) Growth velocity at least 2 standard deviations below the age-appropriate mean or height at least 2.25 standard deviations below the age-appropriate mean. 2) Subnormal response (less than 10 ng/mL) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 3) Development of neutralizing antibodies to growth hormone product(s). 4) Member has open epiphyses
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For pediatric patients, reauthorization requires open epiphyses.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Inluriyo

Products Affected

- INLURIYO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) disease is ER-positive, HER2-negative, 2) disease harbors an ESR1 mutation, as detected by an FDA-approved test, 3) member has experienced disease progression following at least one line of endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inlyta

Products Affected

- INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced renal cell carcinoma -AND- one of the following (1-2): 1) if using axitinib as first line therapy, member is using axitinib in combination with avelumab or pembrolizumab 2) if using axitinib as a single-agent, member has been treated with at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inqovi

Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation chronic myelomonocytic leukemia. Documentation of myelodysplastic syndrome -AND- One of the following (1 or 2): 1) French American-British MDS subtypes of refractory anemia, refractory anemia with ringed sideroblasts or refractory anemia with excess blasts. 2) International Prognostic Scoring System group of intermediate-1, intermediate-2 or high-risk.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Inrebic

Products Affected

- **INREBIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate-2 or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) -AND- If a new start, baseline platelet count of greater than or equal to $50 \times 10^9/L$
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Insulin Supplies

Products Affected

- ALCOHOL PADS
- GAUZE PAD TOPICAL BANDAGE 2 X 2 "

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation the product is being used for the delivery of insulin into the body.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Interferon Alfa

Products Affected

- PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Interferon Beta

Products Affected

- **AVONEX INTRAMUSCULAR PEN INJECTOR KIT**
- **AVONEX INTRAMUSCULAR SYRINGE KIT**
- **BETASERON SUBCUTANEOUS KIT**
- **PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML**
- **PLEGRIDY SUBCUTANEOUS**

SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IPF Agents

Products Affected

- **OFEV**
- *pirfenidone oral capsule*
- *pirfenidone oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 50% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30%. For Ofev only, documentation of systemic sclerosis-associated interstitial lung disease -AND- baseline forced vital capacity (FVC) of at least 40% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- documentation of a chest computed tomography (CT) scan demonstrating greater than or equal to 10% pulmonary fibrosis. For Ofev only, documentation of chronic fibrosing interstitial lung disease with progressive phenotype -AND- chest computing tomography (CT) scan demonstrating greater than 10% fibrosing disease -AND- baseline forced vital capacity (FVC) of at least 45% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- disease progression in previous 24 months shown by one of the following : 1. Relative decline in FVC greater than or equal to 10% predicted 2. Relative decline in FVC greater than or equal to 5% but less than 10% predicted and either worsening of respiratory symptoms or increased extent of fibrotic changes on CT 3. Worsening of respiratory symptoms and increasing extent of fibrotic changes on CT
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iressa

Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	Use in tumors with EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.
Required Medical Information	Documentation of diagnosis -AND- the following: 1) EGFR exon 19 deletion mutations or exon 21 (L858R) mutations as detected by an FDA-approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Itovebi

Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (1-4): 1) disease is HR-positive, HER2-negative 2) disease is PIK3CA-mutated, as detected by an FDA-approved test 3) the member is using in combination with palbociclib and fulvestrant 4) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Itraconazole

Products Affected

- *itraconazole oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If using for diagnosis of onychomycosis, confirmation through positive laboratory testing (e.g. KOH preparation, fungal culture, or nail biopsy) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Onychomycosis: 3 months. All other indications: 3 months initial, 12 months reauth
Other Criteria	Documentation of trial/failure or intolerance of amphotericin b must be provided for approval in patients with aspergillosis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ivermectin Oral

Products Affected

- *ivermectin oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of strongyloidiasis of the intestinal tract (non-disseminated disease) or onchocerciasis -AND- Member weighs greater than or equal to 15kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iwilfin

Products Affected

- IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of high-risk neuroblastoma (HRNB) -AND- partial response to anti-glycolipid disialoganglioside (GD2) immunotherapy (e.g., dinutuximab, naxitamab).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jakafi

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) and if a new start, baseline platelet count of greater than or equal to $50 \times 10^9/L$ -OR- documentation of polycythemia vera and inadequate response or intolerance to hydroxyurea -OR- Documentation of steroid refractory acute graft-versus-host disease and prior therapy with at least one systemic corticosteroid -OR- Documentation of chronic graft-versus-host disease with prior failure of at least one systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jascayd

Products Affected

- JASCAVD

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 45% -AND- a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 25% -AND- 1 of the following (1-3): 1) Concomitant use of a stable dose of Esbriet (pirfenidone) or Ofev (nintedanib). For concomitant Esbriet (pirfenidone) therapy, Jascayd dose will remain 18 mg orally twice daily. 2) Receiving Jascayd as monotherapy and intolerance to Esbriet (pirfenidone) and Ofev (nintedanib), 3) Receiving Jascayd as monotherapy and contraindication to Esbriet (pirfenidone) and Ofev (nintedanib). Diagnosis of progressive pulmonary fibrosis (PPF) -AND- documentation of a chest computed tomography (CT) scan demonstrating at least 10% pulmonary fibrosis -AND- a baseline FVC of at least 45% -AND- a percent predicted DLCO of at least 25% -AND- at least 2 of the following (4-6): 4) Worsening of respiratory symptoms 5) Physiological evidence of disease progression defined as 1 of the following (a or b): a) Decline in FVC of at least 5% (relative from baseline). b) Decline in percent predicted DLCO of at least 10% (relative from baseline). 6) Radiological evidence of disease progression defined as increased extent of fibrotic changes on imaging (e.g., chest computed tomography scan) -AND- One of the following (7-9): 7) Concomitant use of a stable dose of Ofev (nintedanib), 8) Treatment naive to antifibrotic treatment, or 9) Previously discontinued Ofev (nintedanib) with no plans to initiate or resume background PPF treatment.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jaypirca

Products Affected

- **JAYPIRCA ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory mantle cell lymphoma, member has received at least two (2) previous lines of systemic therapy, at least one (1) of which was a BTK inhibitor. For relapsed or refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia, member has previously been treated with a covalent BTK inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Joenja

Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of activated phosphoinositide 3-kinase delta syndrome (APDS) with genetic confirmation of variant in PIK3CD or PIK3R1 gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Jynarque

Products Affected

- *tolvaptan (polycys kidney dis) oral tablet*
- *tolvaptan (polycys kidney dis) oral tablets, sequential*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of rapidly progressing autosomal dominant polycystic kidney disease defined by one of the following: 1.) Historical decline in eGFR greater than or equal to 3mL/min/1.73 m*2 within a 12 month period, or 2.) Mayo imaging classification of 1C, 1D, or 1E.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, prescriber attestation of slowed decline of kidney function
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kalydeco

Products Affected

- **KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG**
- **KALYDECO ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to ivacaftor based on clinical and or in vitro assay (e.g. G551D, G1244E, G1349D)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kerendia

Products Affected

- **KERENDIA**

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than 25 mL/min/1.73 m*2, serum potassium greater than 5.5 mEq/L
Required Medical Information	Documentation of type 2 diabetes mellitus with chronic kidney disease (T2DM+CKD) or heart failure (HF) with left ventricular ejection fraction (LVEF) of greater than or equal to 40% -AND- one of the following (1. 2. or 3.): 1) concomitant use of a sodium-glucose Cotransporter-2 (SGLT2) inhibitor 2) therapeutic failure to at least one SGLT2 inhibitor or 3) contraindication or intolerance to at least one SGLT2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that signs or symptoms of hyperkalemia are not present.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kesimpta

Products Affected

- **KESIMPTA PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kineret

Products Affected

- KINERET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), therapeutic failure or intolerance to at least one (1) corticosteroid, or all corticosteroids are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kisqali

Products Affected

- **KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative -AND- one of the following (1-3): 1) member is using ribociclib in combination with an aromatase inhibitor as initial endocrine-based therapy 2) member is using ribociclib in combination with fulvestrant and member is using fulvestrant as initial endocrine-based therapy or member has experienced disease progression on endocrine therapy. 3) disease is classified as stage II or stage III early breast cancer at high risk of recurrence, ribociclib is being used in combination with an aromatase inhibitor as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kisqali Femara

Products Affected

- **KISQALI FEMARA CO-PACK ORAL
TABLET 400 MG/DAY(200 MG X 2)-2.5
MG, 600 MG/DAY(200 MG X 3)-2.5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative - AND- one of the following (1-2): 1) disease is classified as stage II or stage III early breast cancer at high risk of recurrence and used as adjuvant treatment 2) used as initial endocrine-based therapy for advanced or metastatic breast cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Korlym

Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endogenous Cushing's syndrome -AND- Patient is not a candidate for surgery or where surgery has failed -AND- one of the following (1 or 2): 1) Diagnosis of diabetes, 2) Glucose intolerance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Koselugo

Products Affected

- **KOSELUGO ORAL CAPSULE 10 MG, 25 MG**
- **KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- for neurofibromatosis type 1 (NF1), documentation of symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Koselugo granules, 1 of the following is required (1-2): 1) inability to swallow oral capsules, 2) body surface area of less than 0.55 m*2
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Krazati

Products Affected

- **KRAZATI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of locally advanced or metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA-approved test -AND- using in combination with cetuximab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kuvan

Products Affected

- **JAVYGTOR**
- *sapropterin*
- **ZELVYSIA ORAL POWDER IN PACKET 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of PKU -AND- documented baseline Phe level greater than 6 mg/dL -AND- clinical documentation of current weight -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, initial therapy has resulted in decrease in phenylalanine levels from baseline or current phenylalanine levels within the range of 120-360 micromol/L -AND- clinical documentation of current weight is required -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kygevvi

Products Affected

- **KYGEVVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Thymidine Kinase 2 Deficiency (TK2d) -AND- Pathogenic mutation in TK2 gene, as confirmed by genetic testing -AND- symptom onset on or before 12 years of age
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Latuda

Products Affected

- *lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for bipolar 1 disorder, documentation of trial and failure or intolerance to one other formulary medication indicated in bipolar 1 disorder (e.g. quetiapine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lazcluze

Products Affected

- **LAZCLUZE ORAL TABLET 240 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) -AND- all of the following (1-4): 1) disease harbors EGFR exon 19 deletions or EGFR exon 21 (L858R) substitution mutation, as detected by an FDA-approved test 2) disease is locally advanced or metastatic, 3) member is treatment naive for advanced disease, 4) using in combination with amivantinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lenvima

Products Affected

- LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of differentiated thyroid cancer -AND- meets all of the following (1-3): 1) disease is locally recurrent or metastatic 2) disease is progressive 3) disease is radioactive iodine refractory. Documentation of advanced renal cell carcinoma -AND- one of the following (4-5): 4) member is using lenvatinib in combination with pembrolizumab and is using lenvatinib and pembrolizumab as first-line treatment 5) member is using lenvatinib in combination with everolimus and has experienced therapeutic failure or intolerance to one prior anti-angiogenic therapy. Documentation of unresectable hepatocellular carcinoma -AND- member is using lenvatinib as first-line treatment. Documentation of endometrial cancer -AND- meets all of the following (6-10): 6) member is using lenvatinib in combination with pembrolizumab 7) disease is advanced 8) disease is not classified as microsatellite instability-high or disease is classified as mismatch repair proficient as determined by an FDA-approved test 9) member has experienced disease progression following prior systemic therapy 10) member is not a candidate for curative surgery or radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Leukine

Products Affected

- LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis: following induction chemotherapy in patients who are 55 years or older with acute myelogenous leukemia (AML) -OR- mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation -OR- acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation -OR- acceleration of myeloid reconstitution following allogeneic BMT -OR- treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic BMT -OR- following acute exposure to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lidoderm

Products Affected

- *lidocaine topical adhesive patch, medicated 5 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) or diabetic peripheral neuropathy (DPN) -AND- One of the following (1-3): 1) trial and failure of 1 other agent used to treat diagnosis (e.g. gabapentin for PHN, duloxetine for DPN), 2) inability to swallow oral medication, 3) unable to take an oral medication due to potential adverse events (e.g. sedation).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Diabetic peripheral neuropathy
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Litfulo

Products Affected

- LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For severe alopecia areata, therapeutic failure/intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Livtency

Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of CMV prophylaxis.
Required Medical Information	Documentation of refractory cytomegalovirus infection or disease as evidenced by antigenemia or polymerase chain reaction (PCR) test -AND- all of the following (1-3): 1) member weighs at least 35 kg, 2) member is a recipient of hematopoietic stem cell transplant -OR- solid organ transplant. 3) member has experienced therapeutic failure to one of the following: ganciclovir, valganciclovir, cidofovir, or foscarnet.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of a previous reduction in CMV DNA level -AND- documentation of one of the following (1-3): 1) new onset symptomatic CMV infection, 2) virologic relapse with treatment-emergent maribavir resistance or 3) continued antiviral treatment is required to achieve virologic clearance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Loargys

Products Affected

- LOARGYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Arginase 1 Deficiency (ARG1-D) -AND- Used in conjunction with dietary protein restriction -AND- Diagnosis is confirmed by 1 of the following (1-3): 1) genetic testing, 2) newborn screening, or 3) elevated arginine levels in plasma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lokelma

Products Affected

- **LOKELMA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 7.4 mmol/L on at least two (2) screenings -AND- Modification of medications to reduce serum potassium levels were not successful, when applicable
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in serum potassium levels following Lokelma administration and continued treatment for hyperkalemia is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lonsurf

Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer as a single agent or in combination with bevacizumab in patients who have previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy -OR- documentation of metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/neu-targeted therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lorbrena

Products Affected

- **LORBRENA ORAL TABLET 100 MG,
25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ALK-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lotronex

Products Affected

- *alosetron oral tablet 0.5 mg, 1 mg*

PA Criteria	Criteria Details
Exclusion Criteria	For irritable bowel syndrome (IBS): Exclude if male gender
Required Medical Information	Documentation of chronic severe diarrhea-predominant IBS -AND- trial and failure or intolerance to one anti-diarrheal (e.g. loperamide), anti-spasmodic, or tricyclic antidepressant, or contraindication to all
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that symptoms of IBS continue to persist AND positive clinical response.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lumakras

Products Affected

- LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- using in combination with panitumumab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lynparza

Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) in complete or partial response to first-line platinum-based chemotherapy 2) diagnosis of deleterious or suspected deleterious BRCA mutated disease or disease is associated with homologous recombination deficiency (HRD) positive status with a deleterious or suspected deleterious BRCA mutation or genomic instability and will be using in combination with bevacizumab. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) deleterious or suspected deleterious germline or somatic BRCA mutation 2) in complete or partial response to platinum-based chemotherapy. For deleterious or suspected deleterious gBRCAm, HER2-negative breast cancer, 1 of the following (1-2): 1) classified as high-risk, early breast cancer and has been treated with neoadjuvant or adjuvant chemotherapy 2) has been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting and if hormone receptor (HR)-positive, has been previously treated with or considered inappropriate for treatment with endocrine therapy. For metastatic pancreatic adenocarcinoma, all of the following (1-2): 1) a deleterious or suspected deleterious gBRCA mutation 2) did not progress on at least 16 weeks of a first-line platinum-based chemotherapy regimen. For deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene mutation metastatic castration-resistant prostate cancer, all of the following (1-2): 1) progressed following prior treatment with enzalutamide or abiraterone 2) concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy. For deleterious or suspected deleterious BRCA mutation metastatic castration-resistant prostate cancer, using in combination with all of the following (1-2): 1) abiraterone 2) prednisone or prednisolone.</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lyrica

Products Affected

- *pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg*
- *pregabalin oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. When using pregabalin products concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lytgobi

Products Affected

- **LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-2): 1) disease harbors FGFR2 fusions or other rearrangements 2) member has experienced therapeutic failure or intolerance to at least one prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Mavyret

Products Affected

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Megace

Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Mekinist

Products Affected

- **MEKINIST ORAL RECON SOLN**
- **MEKINIST ORAL TABLET 0.5 MG, 2 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For trametinib monotherapy, documentation of unresectable or metastatic melanoma -AND- meets all of the following (1-2): 1) member has a BRAF V600E or V600K mutation 2) member is BRAF inhibitor treatment naive. For use in combination with dabrafenib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with dabrafenib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA approved test when FDA indicated. For use in combination with dabrafenib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Mekinist powder for oral solution, attestation of inability to swallow Mekinist (trametinib) tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Mektovi

Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, all of the following (1-2): 1) BRAF V600E or V600K mutation status, as detected by an FDA-authorized test 2) using in combination with encorafenib. For metastatic non-small cell lung cancer, all of the following (1-2): 1) BRAF V600E mutation status, as detected by an FDA-authorized test 2) using in combination with encorafenib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Metyrosine

Products Affected

- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pheochromocytoma defined by 1 of the following (1 or 2): 1) Elevated metanephrines in plasma or urine, or 2) Tumor evidence from CT scan or MRI -AND- Documentation of 1 of the following (3., 4., or 5.): 3) Planned resection surgery, 4) Resection surgery is contraindicated, or 5) malignant pheochromocytoma. -AND- therapeutic failure, contraindication, or intolerance to a selective alpha-blocker (e.g., doxazosin, prazosin, terazosin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Modeyso

Products Affected

- **MODEYSO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of diffuse midline glioma -AND- disease harbors an H3 K27M mutation -AND- member has experienced progressive disease following prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Mulpleta

Products Affected

- MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -AND- trial and failure or intolerance to Doptelet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Myasthenia Gravis

Products Affected

- **ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML, 23 MG/0.574 ML, 32.4 MG/0.81 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized myasthenia gravis (gMG) -AND- Anti-acetylcholine receptor (AChR) antibody-positive -AND- Therapeutic failure, contraindication, or intolerance to generic pyridostigmine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of gMG signs/symptoms improvement (e.g., speech, swallowing, mobility, and/or respiratory function) -OR- decreased gMG exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Namzarin

Products Affected

- *memantine-donepezil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and intolerance to generic memantine and generic donepezil
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nayzilam

Products Affected

- NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nemluvio

Products Affected

- NEMLUVIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of prurigo nodularis -OR- Documentation of moderate to severe atopic dermatitis and one of the following (1-3): 1) trial and failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus), 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement, or 3) severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For prurigo nodularis, must follow recommended dosing guidelines based upon weight. For atopic dermatitis, must follow recommended dosing based on FDA approved dosing guidelines. For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Nerlynx

Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of early-stage, HER2-positive breast cancer -AND- meets all of the following: 1)member has received adjuvant trastuzumab-based therapy 2) member is using neratinib as a single agent. Documentation of advanced HER-2 positive, or metastatic HER2-positive breast cancer and meets all of the following 1) using neratinib in combination with capecitabine 2) member has received two or more prior anti-HER2 based regimens in the metastatic setting.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nexavar

Products Affected

- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally recurrent or metastatic, progressive, differentiated thyroid carcinoma, refractory to radioactive iodine treatment
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nexletol

Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>1. HeFH supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C at least 190mg/dL or untreated LDL-C at least 160mg/dL before 20 yrs of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite Simon Broome register criteria, or definite on the Make Early Diagnosis to Prevent Early Deaths tool -AND- LDL-C at least 70mg OR non-HDL at least 100mg/dL or less than 50% reduction in baseline LDL-C despite use with maximally tolerated statin therapy, or if very high risk, LDL-C at least 55mg/dL or non-HDL at least 85mg/dL -AND- Attestation of combination use with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documented statin intolerance -AND- Therapeutic failure/intolerance/contraindication to ezetimibe</p> <p>2. ASCVD or attestation of high risk for CVD, -AND- Therapeutic failure to a maximally tolerated statin or documented statin intolerance -AND- Therapeutic failure/intolerance/contraindication to ezetimibe.</p> <p>3. Primary Hyperlipidemia not associated w/ ASCVD or HeFH -AND- LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% reduction in baseline LDL-C despite use with maximally tolerated statin therapy -AND- Attestation of combination use with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documented statin intolerance -AND- Therapeutic failure/intolerance/contraindication to ezetimibe.</p>
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of ASCVD or High ASCVD Risk, attestation of need for continued therapy is required. For reauthorization associated with all other diagnoses, documentation showing an LDL-C reduction from baseline -AND- attestation of continued use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statin which resolved upon discontinuation of statin or documentation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ninlaro

Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- previous treatment with at least 1 prior therapy -AND- used in combination with lenalidomide and dexamethasone
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nitisinone

Products Affected

- *nitisinone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary tyrosinemia type 1 (HT-1) confirmed by biochemical or genetic testing -AND- dietary restriction of tyrosine and phenylalanine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Northera

Products Affected

- *droxidopa oral capsule 100 mg, 200 mg, 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of neurogenic orthostatic hypotension caused by primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency or non-diabetic autonomic neuropathy -AND- Documentation of inadequate response, intolerance or contraindication to preferred generic alternative midodrine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of increase from baseline of systolic or diastolic blood pressure upon standing -OR- attestation of decrease from baseline of neurogenic orthostatic hypotension symptoms upon standing (e.g., dizziness, feeling faint, etc.).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nubeqa

Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of non-metastatic castration-resistant prostate cancer or metastatic castration-sensitive prostate cancer -AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nuedexta

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting improvement in symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nuplazid

Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hallucinations and delusions associated with Parkinson's disease psychosis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nurtec

Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine Prevention, defined as 4-14 migraine days per month, the following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). For acute treatment of migraine with or without aura, trial and failure of one generic triptan or all triptans are contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency. For reauthorization of acute treatment of migraine, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nuvigil

Products Affected

- *armodafinil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Octreotide

Products Affected

- *octreotide acetate injection solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- therapeutic failure or cannot be treated with surgical resection, pituitary irradiation or bromocriptine mesylate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Odomzo

Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced basal cell carcinoma (laBCC) that has recurred following surgery or radiation therapy or for use in patients who are not candidates for surgery or radiation therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ogsiveo

Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressing desmoid tumor(s) requiring systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ojemda

Products Affected

- **OJEMDA ORAL SUSPENSION FOR RECONSTITUTION** **MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)**
- **OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory pediatric low-grade glioma - AND- one of the following (1-2): 1) BRAF fusion or rearrangement, or 2) BRAF V600K mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ojjaara

Products Affected

- OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of anemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Onfi

Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Onureg

Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute myeloid leukemia that has achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy -AND- Inability to complete intensive curative therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Opipza

Products Affected

- OPIPZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- 1 of the following (1-2): 1) intolerance to generic aripiprazole tablets, 2) inability to swallow oral tablets. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Orencia

Products Affected

- **ORENCIA CLICKJECT**
- **ORENCIA SUBCUTANEOUS SYRINGE**
125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	<p>For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients over 18 years of age must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Cosentyx, Otezla/Otezla XR, Skyrizi SC, Rinvoq, and Xeljanz/Xeljanz XR. For polyarticular juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq/Rinvoq LQ, and Xeljanz/Xeljanz solution. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Orgovyx

Products Affected

- **ORGOVYX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Orkambi

Products Affected

- **ORKAMBI ORAL GRANULES IN PACKET**
- **ORKAMBI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis and homozygous F508del mutation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Orserdu

Products Affected

- **ORSERDU ORAL TABLET 345 MG, 86 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-4): 1) member is male or a postmenopausal female 2) disease is ER-positive, HER2-negative 3) disease harbors an ESR1 mutation, as detected by an FDA-approved test 4) member has experienced disease progression on or after an endocrine based regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Otezla

Products Affected

- OTEZLA (47)
- OTEZLA STARTER ORAL TABLETS, DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG
- OTEZLA XR
- OTEZLA XR INITIATION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For plaque psoriasis, all of the following (1-2): 1) member is greater than or equal to 18 years of age with mild-to-severe disease -OR- if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg for Otezla or 50 kg for Otezla XR, and member has moderate-to-severe disease -AND- 2) inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For oral ulcers associated with Behcet's Disease, inadequate response or intolerance to one systemic therapy for prevention of recurrent oral ulcers. For psoriatic arthritis, if the member is greater than or equal to 6 years of age and less than 18 years and weight is greater than or equal to 20 kg for Otezla or 50 kg for Otezla XR.
Age Restrictions	Deny if less than 6 years of age for plaque psoriasis or psoriatic arthritis. Deny if less than 18 years of age for Behcet's Disease.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Panretin

Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions in patients with AIDS-related Kaposi Sarcoma (KS) who are not receiving systemic therapy for KS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pemazyre

Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable locally advanced cholangiocarcinoma or metastatic cholangiocarcinoma -AND- all of the following (1-2): 1) disease harbors FGFR2 fusion or other rearrangement as detected by an FDA-approved test 2) member has experienced therapeutic failure or intolerance to at least on prior therapy. Documentation of relapsed or refractory myeloid/lymphoid neoplasms -AND- disease harbors an FGFR1 rearrangement.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Pheburane

Products Affected

- PHEBURANE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- intolerance to generic sodium phenylbutyrate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Piqray

Products Affected

- **PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR-positive, HER2-negative tumor status, 2) PIK3CA mutation positive as detected by an FDA-approved test, 3) disease progression on or after an endocrine-based regimen, 4) Concomitant therapy with fulvestrant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pomalyst

Products Affected

- *pomalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma, and combination use with dexamethasone, and previous trial of at least 2 therapies including lenalidomide and a proteasome inhibitor, and disease progression on or within 60 days of completion of the last therapy -OR- Documentation of AIDS-related Kaposi sarcoma (KS) after failure of highly active antiretroviral therapy (HAART) or in patients with KS who are HIV negative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Posaconazole Tablet

Products Affected

- *posaconazole oral tablet, delayed release (dr/ec)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight greater than 40 kg. For invasive aspergillosis infection, trial/failure or contraindication to voriconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Preferred Adalimumab

Products Affected

- **HADLIMA** **SYRINGE KIT 20 MG/0.2 ML, 40**
- **HADLIMA PUSH TOUCH** **MG/0.4 ML**
- **HADLIMA(CF)** • **YUFLYMA(CF)**
- **HADLIMA(CF) PUSH TOUCH** • **YUFLYMA(CF) AI CROHN'S-UC-HS**
- **SIMLANDI(CF) AUTOINJECTOR** • **YUFLYMA(CF) AUTOINJECTOR**
- **SIMLANDI(CF) SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 1 immunosuppressant or corticosteroid, or all are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. These criteria are applicable to Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Preferred Ustekinumab

Products Affected

- **STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**
- **YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**
- **YESINTEK SUBCUTANEOUS SOLUTION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease or Ulcerative Colitis, attestation of receiving or currently undergoing a single induction dose of IV ustekinumab within 2 months of initiating therapy with ustekinumab SC.
Age Restrictions	Deny if less than 18 years of age for Crohn's Disease and Ulcerative Colitis. Deny if less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	Must follow recommended dosing guidelines based upon weight. Induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Prenatal Vitamins

Products Affected

- **PRENATAL VITAMIN PLUS LOW IRON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of nutritional supplementation required in a female of child-bearing potential during pre-conception, pregnancy, or lactation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Prescription Drug Combo

Products Affected

- acetaminophen-codeine oral solution 120-12 mg/5 ml
- acetaminophen-codeine oral tablet
- alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg
- buprenorphine
- clonazepam oral tablet 0.5 mg, 1 mg, 2 mg
- clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg
- clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg
- **DIAZEPAM INTENSOL**
- diazepam oral solution 5 mg/5 ml (1 mg/ml)
- diazepam oral tablet
- fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr
- hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg
- hydromorphone oral liquid
- hydromorphone oral tablet 2 mg, 4 mg, 8 mg
- **LORAZEPAM INTENSOL**
- lorazepam oral tablet 0.5 mg, 1 mg, 2 mg
- methadone oral solution 10 mg/5 ml, 5 mg/5 ml
- methadone oral tablet 10 mg, 5 mg
- morphine concentrate oral solution
- morphine oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)
- morphine oral tablet
- morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg
- oxycodone oral capsule
- oxycodone oral concentrate
- oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg
- oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg
- tramadol oral tablet 50 mg
- tramadol-acetaminophen
- zaleplon oral capsule 10 mg, 5 mg
- zolpidem oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	<p>For concomitant use of opiate agonist and substance abuse therapy, documentation of acute pain condition (eg, acute traumatic injury) in which treatment with other agents would cause insufficient pain control or treatment for terminal illness related pain. For concomitant use of an opiate agonist, benzodiazepine, and centrally acting skeletal muscle relaxant, trial/failure of at least 2 other skeletal muscle relaxants (eg, methocarbamol, metaxalone), understanding these skeletal muscle relaxants are high-risk medications in geriatric patients, attestations that non-opiate (eg, NSAIDs) and non-benzodiazepine therapies (eg, SSRI, SNRI) have been considered, and attestation of intent to monitor & address concomitant drug-drug interaction adverse events. For concomitant use of opiate agonist and other opiate potentiators (eg, gabapentinoids, sedative-hypnotics), attestation of intent to monitor & address concomitant drug-drug interaction adverse events. For long acting opioid medications, the following apply (1-3): 1)Severe pain requiring daily, around-the-clock, long-term opioid treatment. 2)Not opioid naive. 3)Attestations that non-opiate alternative therapies have been explored (eg, NSAIDs), controlled substance history was reviewed in the state Prescription Drug Monitoring Program (PDMP), counseling on potential adverse effects of opioid analgesics, including risk of misuse, abuse, addiction.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	<p>Opiate tx for pain+subs. abuse, approve opiate x 1mo. All other combos and dx approve x 12mo.</p>
Other Criteria	<p>Opiate agonists will receive automatic approval if no recent claims for a substance abuse therapy (e.g. buprenorphine-naloxone) OR a benzodiazepine with a centrally acting skeletal muscle relaxant (e.g., carisoprodol) OR a gabapentinoid OR a sedative-hypnotic. Benzodiazepines (e.g. triazolam, alprazolam) will receive automatic approval if no recent claims for an opiate agonist with a centrally acting skeletal muscle relaxant (e.g. carisoprodol). Sedative-hypnotics (e.g. zolpidem) will receive automatic approval if no recent claims for an opiate agonist.</p>
Indications	<p>All FDA-approved Indications.</p>
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Prevymis Oral Pellets

Products Affected

- **PREVYMIS ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 6 months of age with HSCT. Deny if less than 12 years of age with kidney transplant.
Prescriber Restrictions	
Coverage Duration	7 months
Other Criteria	One of the following is required (1-2): 1) inability to swallow tablets, 2) unable to use Prevymis (letermovir) tablets due to body weight dosing limitations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Prolia

Products Affected

- PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -And- For osteoporosis at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 (i.e. osteopenia) -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For osteoporosis and osteopenia, documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Provigil

Products Affected

- *modafinil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications. Diagnosis of fatigue associated with Multiple Sclerosis (MS)</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Fatigue associated with Multiple Sclerosis (MS)
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pulmonary Arterial Hypertension

Products Affected

- **ADEMPAS**
- *ambrisentan*
- **OPSUMIT**
- **OPSYNVI**
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group. For sildenafil in pediatric individuals, an exception to RHC will be allowed when the risk of RHC outweighs the benefit -AND- prescriber attests alternative studies have been completed (i.e. CT, MRI or specified test ruling out other causes of pulmonary hypertension). For Adempas, additional diagnosis of CTEPH as documented by right heart catheterization and V/Q scan substantiating mPAP greater than 20 mmHg at rest and PWP less than or equal to 15 mmHg and documented presence of occlusive thrombi within the pulmonary arteries will be approved.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Pulmozyme

Products Affected

- PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis -AND- Used in conjunction with standard therapies for management of cystic fibrosis to improve pulmonary function.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization, attestation of increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Qinlock

Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced gastrointestinal stromal tumor -AND- Prior treatment with imatinib and 2 other kinase inhibitors.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Quinine

Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment or prevention of leg cramps
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	Doses for duration greater than 10 days will not be approved
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Qulipta

Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, of which 8 or more are migraine days. The following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ravicti

Products Affected

- *glycerol phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Urea cycle disorders due to N-acetylglutamate synthetase deficiency, Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of a urea cycle disorders (UCDs)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Repatha

Products Affected

- **REPATHA SURECLICK**
- **REPATHA SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
Required Medical Information	<p>1.HoFH supported by 1 of the following (a-d): a) bi-allelic causal mutations of FH by genetic testing, b) LDL-C over 560mg/dL, c) LDL-C over 400mg/dL w/ aortic valve disease or xanthomata at less than 20 yrs old, d) LDL-C over 400mg/dL and 1 or both parents w/ clinically diagnosed FH, positive genetic testing for a known LDL-C-raising (LDLR, Apo[b], PCSK9) gene defect, or autosomal-recessive FH AND current LDL-C over 115mg/dL (if 17 yrs or younger) or over 70mg/dL (18 yrs or older) or over 55mg/dL w/ ASCVD or major ASCVD risk factors (18 yrs or older) AND Therapeutic failure to a max tolerated statin or documented statin intolerance. 2.HeFH supported by 1 of the following (a-d): a) causal mutation of FH by genetic testing, b) untreated LDL-C at least 190mg/dL or untreated LDL-C at least 160mg/dL before 20 yrs old, c) physical signs of FH (e.g. xanthomas, xanthelasma), d) diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria w/ score over 8 points, definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool. AND Therapeutic failure to a max tolerated statin or documented statin intolerance AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C, or if very high risk, LDL-C at least 55mg/dL or non-HDL-C at least 85mg/dL. 3.Hypercholesterolemia w/ ASCVD or attestation of major CV risk AND LDL-C at least 55mg/dL or non-HDL-C at least 85mg/dL or, if ASCVD, less than 50% decrease in baseline LDL-C AND Therapeutic failure to a max tolerated statin or documented statin intolerance. 4.Primary Hyperlipidemia/Hypercholesterolemia (w/o ASCVD, HeFH, or HoFH) AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C AND Therapeutic failure to a max tolerated statin or documented statin intolerance.</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization associated with Major Cardiovascular Risk, attestation of need for continued Repatha therapy is required. For reauthorization associated with all other diagnoses, documentation showing an LDL-C reduction on Repatha therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Retevmo

Products Affected

- **RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is classified as RET gene fusion as detected by an FDA approved test. Documentation of advanced or metastatic medullary thyroid cancer -AND- disease is classified as RET mutation as detected by an FDA approved test. Documentation of advanced or metastatic thyroid cancer -AND- all of the following (1-2): 1) disease is classified as RET gene fusion as detected by an FDA approved test 2) if radioactive iodine is appropriate for the member, the member is radioactive iodine-refractory. Documentation of locally advanced or metastatic solid tumor(s) -AND- disease harbors a RET gene fusion, as detected by an FDA-approved test -AND- one of the following (1-2): 1) the member has no satisfactory alternative treatments 2) the member's tumors have progressed following prior systemic treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Revlimid

Products Affected

- *lenalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of chronic lymphocytic leukemia outside of a controlled clinical trial
Required Medical Information	Diagnosis of multiple myeloma in combination with dexamethasone -OR- diagnosis of multiple myeloma, as maintenance following autologous hematopoietic stem cell transplant (auto-HSCT) -OR- diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities -OR- diagnosis of mantle cell lymphoma (MCL) in which disease has relapsed or progressed after two prior therapies, one of which included bortezomib -OR- diagnosis of follicular lymphoma in combination with a rituximab product after previous treatment -OR- diagnosis of marginal zone lymphoma in combination with a rituximab product after previous treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Revuforj

Products Affected

- **REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory acute leukemia, member has a lysine methyltransferase 2A gene (KMT2A) translocation, as determined by an FDA-authorized test. For relapsed or refractory acute myeloid leukemia, the member has a susceptible NPM1 mutation -AND- the member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rexulti

Products Affected

- REXULTI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI). For schizophrenia, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rezdiffra

Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in conjunction with diet and exercise. Liver biopsy (within the past 3 years) or noninvasive tests (NITs) (performed within the past 6 months) indicating stage F2 or F3 fibrosis -AND- attestation of 1 cardiometabolic risk factor (e.g. obesity, hyperglycemia, hypertriglyceridemia, hyperlipidemia, hypertension) -AND- attestation that member does not have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma -AND- attestation that member's alcohol consumption is not defined as excess alcohol use (greater than 20g/day in females or greater than 30g/day in males)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of continued use in conjunction with diet and exercise -AND- member has experienced improvement or stabilization of fibrosis as demonstrated by NIT -AND- member has not progressed to stage F4 fibrosis -AND- attestation that member does not have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma -AND- attestation that member's alcohol consumption is not defined as excess alcohol use (greater than 20g/day in females or greater than 30g/day in males)
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rezlidhia

Products Affected

- **REZLIDHIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rinvoq

Products Affected

- **RINVOQ ORAL TABLET EXTENDED
RELEASE 24 HR 15 MG, 30 MG, 45 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate) or all non-biologic DMARDs are contraindicated. For moderate to severe refractory atopic dermatitis whose disease is not adequately controlled with other systemic drug products, documentation of one of the following (1 or 2): 1) trial & failure, or intolerance to at least one topical corticosteroid -OR- topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) 2) The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For Non-radiographic Axial Spondyloarthritis, trial & failure or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or contraindication to all. For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	For diagnoses in which tumor necrosis factor (TNF) blockers are also indicated (e.g., Rheumatoid Arthritis, Psoriatic Arthritis), the member has experienced therapeutic failure or intolerance to at least 1 TNF blocker. For ulcerative colitis or Crohns disease, the member has experienced therapeutic failure or intolerance to at least 1 systemic therapy if TNF blockers are not advisable.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rinvoq LQ

Products Affected

- RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	For PsA, deny if 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rivfloza

Products Affected

- **RIVFLOZA SUBCUTANEOUS SOLUTION**
- **RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-4): 1) diagnosis of primary hyperoxaluria type 1 (PH1) confirmed by one of the following (a or b): a) Genetic testing (i.e., AGXT gene mutation), b) Liver biopsy (i.e., absence or significantly reduced AGT activity), 2) relatively preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m ²), 3) at least two elevated urinary oxalate levels greater than 1.5 times the upper reference limit, and 4) The member has not received a liver transplant.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, continued preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m ²) -AND- reduction in urinary oxalate levels from baseline -AND- has not received a liver transplant.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Romvimza

Products Affected

- ROMVIMZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor -AND- attestation that surgical resection may cause one of the following (1-2): 1) worsening functional limitation, 2) severe morbidity
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rozlytrek

Products Affected

- **ROZLYTREK ORAL CAPSULE 100 MG, 200 MG**
- **ROZLYTREK ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, the tumor status is ROS1-positive. For solid tumors with NTRK gene fusion without a known acquired resistance mutation, the tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or the tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rubraca

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) Disease harbors a deleterious BRCA mutation (germline or somatic) 2) member is in complete or partial response to platinum-based chemotherapy. For metastatic castration-resistant prostate cancer, all of the following (1-3): 1) disease harbors a deleterious BRCA mutation (germline and/or somatic) 2) member has been treated with androgen receptor-directed therapy 3) member is concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rydapt

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Use as single agent induction therapy for AML
Required Medical Information	Documentation of diagnosis. For a new diagnosis of acute myeloid leukemia, member is using in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation chemotherapy regimens and member is FLT3 mutation positive as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Sabril

Products Affected

- *vigabatrin*
- **VIGADRONE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of refractory complex partial seizures -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance to at least two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, tiagabine) -OR- documentation of use as monotherapy in treatment of infantile spasms
Age Restrictions	Deny if less than 2 years of age in treatment of refractory complex partial seizures -OR- if less than 1 month old and greater than 2 years of age in treatment of infantile spasms
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Samsca

Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic hypervolemic or euvolemic hyponatremia evidenced by (1. or 2.): 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	Doses must be initiated in the hospital setting to closely monitor serum sodium. Treatment should be limited to 30 days to minimize risk of liver injury. For reauthorization, treatment is for a new episode of a clinically significant euvolemic or hypervolemic hyponatremia -AND- on of the following (1. or 2.) 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Saphris

Products Affected

- *asenapine maleate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Scemblix

Products Affected

- **SCSEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-3): 1) the member is newly-diagnosed, 2) member has been previously treated for Ph+ CML in chronic phase, 3) disease is positive for the T3151 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Secuado

Products Affected

- **SECUADO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- intolerance to generic asenapine sublingual tablets -AND- trial/failure or intolerance to 1 of the following or all are contraindicated (1-3): 1) olanzapine, 2) quetiapine, 3) risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Siliq

Products Affected

- SILIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For psoriasis, patients must have therapeutic failure or intolerance to 2 preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Cosentyx, Otezla/Otezla XR, and Skyrizi SC. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Sirturo

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Criteria will be applied consistent with the current ATS/CDC/ERS/IDSA Clinical Practice Guideline for the Treatment of Drug-Susceptible and Drug-Resistant Tuberculosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of disease improvement -AND- member requires additional antimicrobial therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Skyclarys

Products Affected

- SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Friedreichs ataxia confirmed by genetic testing (i.e., FXN gene mutation).
Age Restrictions	Deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Skyrizi

Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)
- SKYRIZI SUBCUTANEOUS SYRINGE
- SKYRIZI SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohns disease, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC. For moderate to severe ulcerative colitis, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sohonos

Products Affected

- SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 2.5 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	Deny if female and less than 8 years of age -OR- if male and less than 10 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in volume of new heterotopic ossification from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Solaraze

Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure, intolerance, or contraindication to one of the following (1 or 2): 1) topical fluorouracil solution or fluorouracil 5% cream 2) topical imiquimod 5% cream
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of 30 day washout period since optimal therapeutic effect may not be evident until 30 days following cessation of therapy AND attestation of previous response to diclofenac sodium 3% topical gel therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Somavert

Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- inadequate or partial response to surgery or radiotherapy or not a candidate for surgery or radiotherapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Sprycel

Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adults with Ph+ chronic myeloid leukemia, the member is newly diagnosed in the chronic phase -OR- the member is in chronic, accelerated, or myeloid or lymphoid blast phase and has resistance or intolerance to prior therapy including imatinib. For adults with Ph+ acute lymphocytic leukemia, member has had resistance or intolerance to prior therapy. For pediatric patients with Ph+ CML, the member is in the chronic phase. For pediatric patients with Ph+ acute lymphoblastic leukemia, the member is newly diagnosed and will be using in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Steroidogenesis Inhibitors

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in urinary free cortisol levels from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Stivarga

Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer and trial of a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy, AND an anti-VEGF therapy AND if RAS wild-type, an anti-EGFR therapy -OR- documentation of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with both imatinib and sunitinib -OR- documentation of hepatocellular cancer AND previous treatment with sorafenib
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sutent

Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For gastrointestinal stromal tumor (GIST), the member has experienced therapeutic failure, intolerance, or contraindication to imatinib. For a high risk of recurrent renal cell carcinoma, member has had a nephrectomy and sunitinib is to be used as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sympazan

Products Affected

- SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy -AND- unable to tolerate generic clobazam
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Synarel

Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For central precocious puberty (CPP), advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L. For female with endometriosis, attestation of not pregnant if of childbearing age -AND- Therapeutic failure, contraindication or intolerance to 2 of the following standard of care treatments: NSAIDs, combination hormonal contraceptive, progestin (i.e. medroxyprogesterone injection), GnRH agonist (i.e. Leuprolide) or danazol
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty. Deny if less than 18 years of age for endometriosis.
Prescriber Restrictions	
Coverage Duration	Endometriosis: 6 months, CPP: 6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization for CPP, attestation of pubertal development slowing from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tabrecta

Products Affected

- **TABRECTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with MET exon 14 skipping mutation as detected by an FDA approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tafinlar

Products Affected

- **TAFINLAR ORAL CAPSULE**
- **TAFINLAR ORAL TABLET FOR SUSPENSION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For dabrafenib monotherapy, documentation of unresectable or metastatic melanoma -AND- documentation of a BRAF V600E mutation. For use in combination with trametinib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with trametinib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA-approved test when FDA indicated. For use in combination with trametinib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Tafinlar tablets for oral suspension, attestation of inability to swallow Tafinlar (dabrafenib) capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Tagrisso

Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of non-small cell lung cancer (NSCLC) and one of the following (1-5): 1) Adjuvant therapy after tumor resection -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test 2) Locally advanced disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used in combination with pemetrexed and platinum-based chemotherapy, 3) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used with or without combination therapy of pemetrexed and platinum-based chemotherapy, 4) Metastatic disease -AND- disease harbors EGFR T790M mutations, as detected by an FDA-approved test -AND- has progressed on or after EGFR TKI therapy, 5) Locally advanced, unresectable (stage III) disease -AND- disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy -AND- disease harbors EGFR exon 19 deletions or EGFR exon 21 L858R mutations, as detected by an FDA-approved test.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Talzenna

Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of deleterious or suspected deleterious gBRCAm, HER2-negative locally advanced or metastatic breast cancer as a single agent - OR- Documentation of HRR gene-mutated metastatic castration-resistant prostate cancer in combination with enzalutamide
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Targretin

Products Affected

- *bexarotene oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations of cutaneous T-cell lymphoma in patients who are refractory to at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Targretin Gel

Products Affected

- *bexarotene topical*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions associated with one (1) of the following stages of cutaneous T-cell lymphoma (CTCL) (1. or 2.): 1) Stage IA or 2) Stage IB - AND- the member has experience therapeutic failure, contraindication, or intolerance to at least one other therapy for CTCL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tascenso ODT

Products Affected

- TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Inability to swallow capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tasigna

Products Affected

- *nilotinib hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adult patients with Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib - OR- member is newly diagnosed in the chronic phase. For pediatric patients, one of the following (1-2): 1) member has chronic phase or accelerated phase Ph+ CML and is is resistant or intolerant to prior tyrosine kinase inhibitor therapy 2) member is newly diagnosed with Ph+ CML in the chronic phase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tasimelteon

Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tazorac

Products Affected

- *tazarotene topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 of the following (A or B). A) Documentation of plaque psoriasis -AND- trial and failure or intolerance to at least one topical corticosteroid (e.g. fluocinonide, mometasone, triamcinolone, betamethasone). B) Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tecfidera

Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tepmetko

Products Affected

- **TEPMETKO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with a MET exon 14 skipping alteration
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Testosterone (androgens)

Products Affected

- *testosterone cypionate*
- *testosterone enanthate*
- *testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or secondary hypogonadism in males -OR- documentation of chronic steroid treatment in men. In all previously noted indications, members must also have documented low total testosterone level below the normal range for the laboratory -OR- a total testosterone level near the lower limit of the normal range with a low free testosterone level which is less than normal based upon the laboratory reference range -OR- the member is not producing any testosterone. Additional approvable indications include female patients with metastatic breast cancer (testosterone enanthate only), primary or secondary hypogonadism in males with testicular failure due to double orchidectomy, and delayed puberty in males (testosterone enanthate only).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Gender Dysphoria, Gender Identity Disorder
Part B Prerequisite	No
Prerequisite Therapy Required	No

Thalomid

Products Affected

- **THALOMID ORAL CAPSULE 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Use as monotherapy for ENL treatment in the presence of moderate to severe neuritis
Required Medical Information	Documentation of multiple myeloma in combination with dexamethasone -OR- documentation for use in the treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) -OR- documentation of therapy for prevention and suppression of the cutaneous manifestations of ENL recurrence
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Thrombopoiesis Stimulating Agents

Products Affected

- *eltrombopag olamine oral powder in packet*
12.5 mg, 25 mg
- *eltrombopag olamine oral tablet* 12.5 mg, 25 mg, 50 mg, 75 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of persistent or chronic immune idiopathic thrombocytopenia purpura and trial and failure of corticosteroid or immunoglobulin therapy or splenectomy -OR- documentation of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy -OR- severe aplastic anemia who have had an insufficient response to immunosuppressive therapy -OR- documentation of first line treatment for severe aplastic anemia and used in combination with at least two immunosuppressive therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Platelet count to be provided
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tibsovo

Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is IDH1 mutation positive as detected by an FDA-approved test. For newly-diagnosed acute myeloid leukemia, member is using as monotherapy or in combination with azacitidine -AND- member meets one of the following (1-5): 1) age is greater than or equal to 75 years of age 2) severe cardiac or pulmonary comorbidity 3) reduced renal function 4) hepatic impairment 5) or prescriber attestation that member is not a candidate for intensive induction therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tocilizumab

Products Affected

- **ACTEMRA ACTPEN**
- **ACTEMRA SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide). For polyarticular juvenile idiopathic arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For systemic sclerosis-associated interstitial lung disease (SSc-ILD), patients must have therapeutic failure or intolerance to one immunosuppressant (e.g., mycophenolate mofetil, corticosteroids, cyclophosphamide). Documentation of systemic juvenile idiopathic arthritis.
Age Restrictions	Deny if less than 18 years of age for systemic sclerosis-associated interstitial lung disease (SSc-ILD), Rheumatoid Arthritis, and Giant Cell Arteritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. For polyarticular juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq/Rinvoq LQ, and Xeljanz/Xeljanz solution. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Topical Lidocaine

Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical ointment*
- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tretinoin

Products Affected

- *tretinoin topical cream*
- *tretinoin topical gel 0.01 %, 0.025 %*

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two generic topical non-retinoid acne medications (e.g. clindamycin-benzoyl peroxide)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Trikafta

Products Affected

- **TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL**
- **TRIKAFTA ORAL TABLETS, SEQUENTIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have at least one F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to elexacaftor/tezacaftor/ivacaftor based on in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Trintellix

Products Affected

- **TRINTELLIX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Truqap

Products Affected

- TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) HR mutation status, HER2 mutation status, and PIK3CA/AKT1/PTEN status 2) concomitant therapy with fulvestrant 3) disease progression on at least one endocrine-based regimen in the metastatic setting -OR- recurrence on or within 12 months of completing adjuvant therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tukysa

Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced unresectable HER2-positive breast cancer or metastatic HER2-positive breast cancer, member will be using in combination with trastuzumab and capecitabine -AND- member has received one or more prior anti-HER2 based regimens in the metastatic setting. For RAS wild-type HER2-positive unresectable or metastatic colorectal cancer, member will be using in combination with trastuzumab -AND- member has experienced disease progression following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Turalio

Products Affected

- TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor associated with severe morbidity and functional limitations -AND- patient is not amenable to improvement with surgery or not a candidate for surgery
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tykerb

Products Affected

- *lapatinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced HER2-positive, or metastatic HER2-positive breast cancer, the member has received prior therapy with an anthracycline, a taxane, and trastuzumab -AND- will be using in combination with capecitabine. For HR+, metastatic breast cancer, the member is post-menopausal -AND- the member's cancer over expresses the HER2 receptor -AND- the member will be using lapatinib in combination with letrozole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tymlos

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 24 months.
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Additional documentation of trial/failure, intolerance or contraindication to preferred parathyroid hormone analog teriparatide. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Tymlos will be limited to a coverage duration of 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ubrelvy

Products Affected

- **UBRELVY ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of acute treatment of migraine with or without aura -AND- trial and failure of one generic triptan or all triptans are contraindicated
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	Patients must have therapeutic failure, intolerance, or contraindication to Nurtec ODT. For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Uptravi

Products Affected

- **UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG** **PACK**
- **UPTRAVI ORAL TABLETS,DOSE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Trial and failure of Opsumit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ustekinumab

Products Affected

- **STELARA SUBCUTANEOUS SOLUTION**
- **STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**
- *ustekinumab subcutaneous solution*
- *ustekinumab subcutaneous syringe 45 mg/0.5 ml, 90 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease or Ulcerative Colitis, attestation of receiving or currently undergoing a single induction dose of IV ustekinumab within 2 months of initiating therapy with ustekinumab SC.
Age Restrictions	Deny if less than 18 years of age for Crohn's Disease and Ulcerative Colitis. Deny if less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For ustekinumab products except Steqeyma and Yesintek, intolerance to both preferred ustekinumab products is required: Steqeyma with NDC starting with 72606 and Yesintek with NDC starting with 83257. Must follow recommended dosing guidelines based upon weight. Subcutaneous induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Valchlor

Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received at least one prior skin-directed therapy (e.g. topical corticosteroids, topical chemotherapy, local radiation and topical retinoids).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Valtoco

Products Affected

- VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vanflyta

Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is FLT3-ITD-positive as detected by an FDA-approved test -AND- one of the following (1-3): 1) member is receiving induction therapy and is using Vanflyta in combination with standard cytarabine and anthracycline induction therapy 2) member is receiving consolidation therapy and is using Vanflyta in combination with standard cytarabine consolidation therapy 3) member is receiving maintenance therapy and is using Vanflyta as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Venclexta

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For newly-diagnosed AML, member is using in combination with azacitidine, decitabine, or cytarabine -AND- age greater than or equal to 75 years or presence of at least one comorbidity that precludes use of intensive induction chemotherapy (i.e. severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Verquvo

Products Affected

- VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of heart failure (NYHA Class II to IV) -AND- Left ventricular ejection fraction less than 45% -AND- Hospitalization for heart failure or received outpatient IV diuretics for heart failure -AND- Used in combination with a angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker or Entresto -AND- Used in combination with bisoprolol, carvedilol IR/ER or metoprolol succinate ER.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Verzenio

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative. For early breast cancer that is at high risk of recurrence and is node-positive, all of the following (1-2): 1) used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) 2) used as adjuvant treatment. For advanced or metastatic breast cancer, used as initial endocrine-based therapy and used in combination with an aromatase inhibitor -OR- used after documented disease progression following endocrine therapy and used in combination with fulvestrant -OR- used after documented disease progression and used following endocrine therapy and prior chemotherapy in the metastatic setting and will be used as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Viibryd

Products Affected

- *vilazodone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vioice

Products Affected

- **VIJOICE ORAL GRANULES IN PACKET** **MG**
- **VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe manifestations of PIK3CA Related Overgrowth Spectrum (PROS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vittrakvi

Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation -AND- Tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vivjoa

Products Affected

- VIVJOA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of recurrent vulvovaginal candidiasis defined as at least 3 episodes of vulvovaginal candidiasis in less than one year -AND Documentation the member is NOT of reproductive potential defined as postmenopausal or another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy) -AND- the member has experienced therapeutic failure, contraindication, or intolerance to a six-month maintenance course of oral fluconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vizimpro

Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer -AND- one of the following, as detected by an FDA-approved test (1 or 2): 1) Epidermal growth factor (EGFR) exon 19 deletions, 2) Epidermal growth factor receptor (EGFR) exon 21 L858R substitution mutations.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vonjo

Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of a platelet count of less than $50 \times 10^9/L$.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Voranigo

Products Affected

- **VORANIGO ORAL TABLET 10 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) grade 2 astrocytoma or grade 2 oligodendroglioma, 2) disease harbors a susceptible isocitrate dehydrogenase (IDH)-1 or IDH-2 mutation, as detected by an FDA-approved test, 3) will be used following surgery including biopsy, sub-total resection, or gross total resection.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Voriconazole

Products Affected

- *voriconazole intravenous recon soln*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary cannot take oral voriconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of continued indicators of active disease (e.g. histopathology, positive cultures) is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vosevi

Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Votrient

Products Affected

- *pazopanib oral tablet 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of adipocytic soft tissue sarcoma or gastrointestinal stromal tumor
Required Medical Information	Documentation of diagnosis. For advanced soft-tissue sarcoma, trial/failure of at least one prior chemotherapy regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vraylar

Products Affected

- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For a diagnosis of schizophrenia and bipolar I disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vumerity

Products Affected

- VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Welireg

Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For von Hippel Lindau (VHL) syndrome, one of the following diagnoses not requiring immediate surgery (1, 2, or 3): 1) Renal cell carcinoma (RCC) 2) CNS hemangioblastoma 3) Pancreatic neuroendocrine tumor. For advanced RCC with a clear cell component, prior treatment with a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) -AND- a programmed death receptor-1 (PD-1) or a programmed death-ligand (PD-L1) inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Winrevair

Products Affected

- WINREVAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group -AND- Concomitant use of at least one of the following (1-4): 1) generic endothelin-1 receptor antagonist, 2) phosphodiesterase type 5 inhibitor, 3) soluble guanylate cyclase stimulator, 4) generic prostacyclin agent -AND- Background therapy with PAH-specific therapies will be continued while being treated with Winrevair.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xalkori

Products Affected

- **XALKORI ORAL CAPSULE**
- **XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer (NSCLC), disease is ALK-positive as detected by an FDA-approved test or ROS1-positive as detected by an FDA-approved test. For relapsed or refractory anaplastic large cell lymphoma (ALCL), disease is ALK-positive. For unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT), disease is ALK-positive.
Age Restrictions	Deny if less than 1 year of age or greater than 21 years of age for ALCL.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For crizotinib oral pellets and NSCLC, inability to swallow capsules is required. For crizotinib oral pellets and ALCL / IMT, inability to swallow oral capsules -OR- body surface area less than 1.34 m ² is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xcopri

Products Affected

- **XCOPRI**
- **XCOPRI MAINTENANCE PACK**
- **XCOPRI TITRATION PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Therapeutic failure, intolerance or contraindication to 1 other anti-epileptic drug (e.g. carbamazepine, levetiracetam)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xdemvy

Products Affected

- XDEMVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis confirmed by identification of Demodex infection via microscopic examination of pulled eyelashes -OR- identification of collarettes via slit-lamp evaluation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xeljanz

Products Affected

- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate). Xeljanz immediate release for polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID).
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis, ulcerative colitis, ankylosing spondylitis. For Xeljanz XR, deny if less than 18 years of age for psoriatic arthritis. For Xeljanz regular release tablet, deny if less than 2 years of age for polyarticular juvenile idiopathic arthritis and psoriatic arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced inadequate response or intolerance to at least 1 tumor necrosis factor (TNF) blocker. Doses greater than 10 mg per day for Xeljanz and 11 mg per day for Xeljanz XR will not be approved for rheumatoid arthritis and psoriatic arthritis. Doses greater than 20mg per day for Xeljanz and 22 mg per day for Xeljanz XR will not be approved for ulcerative colitis.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xeljanz Solution

Products Affected

- XELJANZ ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Documentation of polyarticular juvenile idiopathic arthritis -AND- Inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age for polyarticular juvenile idiopathic arthritis and psoriatic arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xenazine

Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary is not actively suicidal
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	In patients with comorbid depression, attestation of adequate treatment for depression is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xermelo

Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of carcinoid syndrome diarrhea AND used in combination with a somatostatin analog AND trial and failure of somatostatin analog monotherapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in average number of daily bowel movements -AND- the member will continue to use in combination with a somatostatin analog.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xgeva

Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For hypercalcemia of malignancy, refractory to bisphosphonates. For giant cell tumor of bone, unresectable or surgical resection is likely to result in severe morbidity -AND- one of the following (1. or 2.)- 1.) the member is 18 years old or older -OR- 2.) the member is a skeletally mature adolescent (e.g. has at least one mature long bone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xifaxan

Products Affected

- XIFAXAN ORAL TABLET 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 or 2. 1) Diagnosis of hepatic encephalopathy AND trial/failure, intolerance, or contraindication to lactulose. 2) Diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D) AND trial/failure, intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Hepatic encephalopathy: 1 year. IBS-D: 14 days.
Other Criteria	No more than three courses of rifaximin for the treatment of IBS-D will be approved per lifetime.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xolair

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of the following (1-2): 1) Chronic Spontaneous Urticaria, 2) trial/failure or intolerance of a second-generation non-sedating H1 antihistamine at the maximum recommended doses (e.g. cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine). -OR- Documentation of the following (3-9): 3) moderate to severe persistent asthma, 4) a positive skin test or in vitro reactivity to a perennial aeroallergen, 5) Baseline IgE titer greater than or equal to 30 IU/mL, 6) documented pretreatment FEV1 less than 80% predicted in adults or less than 90% predicted in children and adolescents or FEV1 reversibility of at least 12% and 200mL after albuterol administration, 7) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12mos or at least 1 asthma exacerbation requiring hospitalization in past 12mos, 8) inadequate symptom control despite regular treatment w/ medium- or high-dose inhaled corticosteroids (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline) w/ or w/o OCS, unless intolerant or contraindicated to all, 9) will continue treatment with a medium- or high-dose ICS and at least 1 add'l asthma controller medication w/ or w/o OCS -OR- Documentation of the following (10-11): 10) chronic rhinosinusitis with nasal polyps (CRSwNP), 11) will use concomitantly with nasal corticosteroid maintenance treatment, -OR- Documentation of the following (12-17): 12) IgE mediated food allergy, 13) diagnosis confirmed by skin prick test or food-specific antibodies, 14) previous allergic reaction to food, 15) using for the reduction of allergic reactions (type 1), including anaphylaxis, 16) will be used in conjunction with food allergen avoidance, 17) member has a documented prescription for epinephrine.</p>
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CSU reauthorization, improved CSU symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For IgE-mediated food allergy reauthorization, member requires continuation of therapy and will continue food allergen avoidance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xolremdi

Products Affected

- XOLREMDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of WHIM syndrome (warts, hypogammaglobulinemia, infections, myelokathexis).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in incidence of infections is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xospata

Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is FLT3 mutation-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xpovio

Products Affected

- **XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80 MG/WEEK (80 MG X 1), 80MG TWICE WEEK (160 MG/WEEK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use in combination with dexamethasone for relapse or refractory multiple myeloma with failure, intolerance or contraindication to 5 therapies (e.g. bortezomib, carfilzomib, lenalidomide, pomalidomide and daratumumab) -OR- Documentation of use in combination with both bortezomib and dexamethasone for multiple myeloma after receiving 1 prior multiple myeloma therapy -OR- Documentation of relapsing or refractory diffuse large B-cell lymphoma with failure, intolerance or contraindication to at least 2 lines of systemic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xtandi

Products Affected

- **XTANDI ORAL CAPSULE**
- **XTANDI ORAL TABLET 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For castration-resistant prostate cancer and metastatic castration sensitive-prostate cancer, the member is using in combination with a GnRH analog or the member has had a bilateral orchiectomy. For non-metastatic castration-sensitive prostate cancer, the member has biochemical recurrence at high risk for metastasis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xyrem

Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	<p>If 18 years or older and no diagnosis of cataplexy, trial and failure, intolerance, or contraindication to generic modafinil -AND- a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.</p>
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Yonsa

Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with methylprednisolone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Yorvipath

Products Affected

- **YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML, 294 MCG/0.98 ML, 420 MCG/1.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hypoparathyroidism -AND- albumin-corrected serum calcium greater than or equal to 7.8 mg/dL -AND- trial and failure of an active form of vitamin D (e.g. calcitriol, alfacalcidol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of normalization or improvement in total serum calcium from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zavesca

Products Affected

- *miglustat*
- YARGESA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of mild to moderate type 1 Gaucher disease confirmed by the following (A. or B.) A. (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm ³ -OR- 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B. Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation enzyme replacement therapy (e.g. Cerezyme, Elelyso, or VPRIV) is not a therapeutic option
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zavzpret

Products Affected

- ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- Therapeutic failure, contraindication or intolerance to one generic triptan - AND- Inability to swallow capsules/tablets
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zejula

Products Affected

- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer, member is in complete or partial response to first-line platinum-based therapy -AND- disease is associated with homologous recombination deficiency-positive status, defined by at least one of the following (1 or 2): 1) a deleterious or suspected deleterious BRCA mutation or 2) genomic instability. For recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, all of the following (1-2): 1) disease harbors a deleterious or suspected deleterious germline BRCA mutation 2) member is in a complete or partial response to platinum-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zelboraf

Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, member has a BRAF V600E mutation. For Erdheim-Chester Disease, member has a BRAF V600 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zelsuvmi

Products Affected

- ZELSUVMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of molluscum contagiosum
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	For reauthorization, attestation that the member has previously experienced complete or partial clearance of molluscum contagiosum lesions with Zelsuvmi -AND- additional course of therapy is required for recurrence of molluscum contagiosum
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zeposia

Products Affected

- ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Zeposia and other disease modifying agents such as interferons, Copaxone, Tysabri.
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Ulcerative Colitis: Plan Year, Multiple Sclerosis: 24 months
Other Criteria	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Skyrizi SC, Rinvoq, and Xeljanz/Xeljanz XR. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zokinvy

Products Affected

- ZOKINVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- BSA of at least 0.39 m ² . For Hutchinson-Gilford Progeria Syndrome (HGPS), a mutation in the LMNA gene is required. For Processing-Deficient Progeroid Lamionpathy (PL), 1 of the following is required (1-3): 1) heterozygous LMNA mutation and progerin-like protein accumulation, 2) homozygous ZMPSTE24 mutations, or 3) compound heterozygous ZMPSTE24 mutations.
Age Restrictions	Deny if less than 12 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zolinza

Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations in patients with cutaneous T-cell lymphoma (CTCL) who have progressive, persistent, or recurrent disease on or following 2 systemic therapies. Systemic therapies include bexarotene, interferon alpha, extracorporeal photochemotherapy, PUVA, single agent or combination chemotherapies (e.g. cyclophosphamide, vinblastine, romidepsin)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zonisade

Products Affected

- ZONISADE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Documentation of adjunctive therapy -AND- Inability to swallow capsules -AND- Therapeutic failure/intolerance to 2 or contraindication to all of the following (1-6): 1) generic carbamazepine suspension/chewable tablet/extended-release capsule, 2) generic gabapentin capsules/solution, 3) generic lacosamide solution, 4) generic levetiracetam solution, 5) generic oxcarbazepine suspension, 6) generic pregabalin capsules/solution.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zoryve 0.15 Cream

Products Affected

- ZORYVE TOPICAL CREAM 0.15 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3). 1) Documentation of mild to moderate atopic dermatitis -AND- 2) therapeutic failure, contraindication or intolerance to one generic formulary topical corticosteroid -OR- documentation of facial or anogenital involvement -AND- 3) therapeutic failure, contraindication or intolerance to topical generic tacrolimus or topical generic pimecrolimus.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ztalmy

Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures associated with CDKL5 deficiency confirmed by genetic testing.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zurzuvaе

Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of postpartum depression -AND- attestation of symptom onset began during the third trimester of pregnancy or up to 4 weeks post-delivery -AND- less than or equal to 12 months postpartum.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zykadia

Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- ALK mutations, as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zytiga

Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*
- **ABIRTEGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with prednisone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Index of Drugs

<i>abiraterone oral tablet 250 mg, 500 mg</i>	362	AVONEX INTRAMUSCULAR	
ABIRTEGA	362	SYRINGE KIT	129
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i>	224	AYVAKIT	22
<i>acetaminophen-codeine oral tablet</i>	224	BAFIERTAM	23
<i>acitretin</i>	1	BALVERSA	24
ACTEMRA ACTPEN	294	BENLYSTA SUBCUTANEOUS	26
ACTEMRA SUBCUTANEOUS	294	<i>benztropine oral</i>	109
ACTIMMUNE	2	BESREMI	28
ADEMPAS	232	BETASERON SUBCUTANEOUS KIT	129
AIMOVIG AUTOINJECTOR		<i>bexarotene oral</i>	282
SUBCUTANEOUS AUTO-INJECTOR		<i>bexarotene topical</i>	283
140 MG/ML, 70 MG/ML	6	BOSULIF ORAL CAPSULE 100 MG, 50 MG	29
AKEEGA	7	BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG	29
ALCOHOL PADS	127	BRAFTOVI	30
ALECENSA	8	BRINSUPRI	32
<i>alosetron oral tablet 0.5 mg, 1 mg</i>	167	BRUKINSA ORAL TABLET	33
<i>alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	224	<i>buprenorphine</i>	224
ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG	10	CABOMETYX	35
ALUNBRIG ORAL TABLETS,DOSE PACK	10	CALQUENCE (ACALABRUTINIB MAL)	36
<i>ambrisentan</i>	232	CAPLYTA	37
<i>amitriptyline</i>	109	CAPRELSA ORAL TABLET 100 MG, 300 MG	38
ANZUPGO	13	<i>carglumic acid</i>	39
ARIKAYCE	14	CAYSTON	40
<i>aripiprazole oral solution</i>	16	CERDELGA	41
<i>aripiprazole oral tablet,disintegrating</i>	16	CINRYZE	44
<i>armodafinil</i>	197	<i>clobazam oral suspension</i>	204
<i>asenapine maleate</i>	259	<i>clobazam oral tablet</i>	204
ATTRUBY	15	<i>clomipramine</i>	109
AUGTYRO ORAL CAPSULE 160 MG, 40 MG	18	<i>clonazepam oral tablet 0.5 mg, 1 mg, 2 mg</i>	224
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG	19	<i>clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	224
AUSTEDO XR	19	<i>clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg</i>	224
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG	19	COBENFY	46
AUVELITY	20	COBENFY STARTER PACK	46
AVMAPKI-FAKZYNJA	21	COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)	47
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	129	COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML	100

COPIKTRA	48	<i>eltrombopag olamine oral powder in</i>	
COSENTYX (2 SYRINGES)	50	<i>packet 12.5 mg, 25 mg</i>	292
COSENTYX PEN (2 PENS)	50	<i>eltrombopag olamine oral tablet 12.5 mg,</i>	
COSENTYX SUBCUTANEOUS		<i>25 mg, 50 mg, 75 mg</i>	292
SYRINGE 75 MG/0.5 ML	50	EMGALITY PEN	66
COSENTYX UNOREADY PEN	50	EMGALITY SYRINGE	
COTELLIC	51	SUBCUTANEOUS SYRINGE 120	
<i>cyproheptadine oral tablet</i>	109	MG/ML, 300 MG/3 ML (100 MG/ML X	
CYSTARAN	52	3)	66
<i>dalfampridine</i>	11	ENBREL MINI	78
DANZITEN	53	ENBREL SUBCUTANEOUS	
<i>dasatinib oral tablet 100 mg, 140 mg, 20</i>		SOLUTION	78
<i>mg, 50 mg, 70 mg, 80 mg</i>	270	ENBREL SUBCUTANEOUS SYRINGE	
DAURISMO ORAL TABLET 100 MG,		25 MG/0.5 ML (0.5), 50 MG/ML (1 ML) ..	78
25 MG	55	ENBREL SURECLICK	78
<i>deferasirox oral tablet, dispersible</i>	56	ENSACOVE	68
DIACOMIT ORAL CAPSULE 250 MG,		ENTRESTO SPRINKLE	69
500 MG	58	EOHILIA	70
DIACOMIT ORAL POWDER IN		EPIDIOLEX	72
PACKET 250 MG, 500 MG	58	<i>ergotamine-caffeine</i>	74
DIAZEPAM INTENSOL	224	ERIVEDGE	75
<i>diazepam oral solution 5 mg/5 ml (1</i>		ERLEADA ORAL TABLET 240 MG,	
<i>mg/ml)</i>	224	60 MG	76
<i>diazepam oral tablet</i>	224	<i>erlotinib</i>	64
<i>diclofenac sodium topical gel 3 %</i>	268	<i>escitalopram oxalate oral capsule</i>	77
<i>dihydroergotamine nasal</i>	59	<i>everolimus (antineoplastic) oral tablet</i>	4
<i>dimethyl fumarate oral capsule, delayed</i>		<i>everolimus (antineoplastic) oral tablet for</i>	
<i>release(dr/ec) 120 mg, 120 mg (14)- 240</i>		<i>suspension 2 mg, 3 mg, 5 mg</i>	4
<i>mg (46), 240 mg</i>	288	EXXUA ORAL TABLET EXTENDED	
<i>doxepin oral capsule</i>	109	RELEASE 24 HR	80
<i>doxepin oral concentrate</i>	109	EXXUA ORAL TABLET, EXT REL	
<i>doxepin oral tablet</i>	109	24HR DOSE PACK	80
DRIZALMA SPRINKLE ORAL		FABHALTA	81
CAPSULE, DELAYED REL		FANAPT	83
SPRINKLE 20 MG, 30 MG, 40 MG, 60		FANAPT TITRATION PACK A	83
MG	60	FASENRA PEN	84
<i>droxidopa oral capsule 100 mg, 200 mg,</i>		FASENRA SUBCUTANEOUS	
<i>300 mg</i>	192	SYRINGE 10 MG/0.5 ML, 30 MG/ML	84
DUPIXENT PEN SUBCUTANEOUS		<i>fentanyl transdermal patch 72 hour 100</i>	
PEN INJECTOR 200 MG/1.14 ML, 300		<i>mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr,</i>	
MG/2 ML	61	<i>75 mcg/hr</i>	224
DUPIXENT SYRINGE		FETZIMA ORAL CAPSULE,EXT REL	
SUBCUTANEOUS SYRINGE 200		24HR DOSE PACK 20 MG (2)- 40 MG	
MG/1.14 ML, 300 MG/2 ML	61	(26)	87
DUVYZAT	63		

FETZIMA ORAL	
CAPSULE,EXTENDED RELEASE 24	
HR 120 MG, 20 MG, 40 MG, 80 MG	87
FILSUVEZ	88
<i> fingolimod</i>	99
FINTEPLA	89
FIRMAGON KIT W DILUENT	
SYRINGE	92
FOTIVDA	95
FRUZAQLA ORAL CAPSULE 1 MG, 5	
MG	96
GAMMAGARD LIQUID	116
GAMMAGARD LIQUID ERC	116
GAMMAGARD S-D (IGA < 1	
MCG/ML)	116
GATTEX 30-VIAL	97
GAUZE PAD TOPICAL BANDAGE 2	
X 2 "	127
GAVRETO	98
<i> gefitinib</i>	133
GILOTRIF	64
<i> glatiramer subcutaneous syringe 20 mg/ml,</i>	
<i> 40 mg/ml</i>	100
GLATOPA SUBCUTANEOUS	
SYRINGE 20 MG/ML, 40 MG/ML	100
<i> glycerol phenylbutyrate</i>	238
GOMEKLI ORAL CAPSULE 1 MG, 2	
MG	105
GOMEKLI ORAL TABLET FOR	
SUSPENSION	105
<i> guanfacine oral tablet extended release 24</i>	
<i> hr</i>	3
HADLIMA	220
HADLIMA PUSHTOUCH	220
HADLIMA(CF)	220
HADLIMA(CF) PUSHTOUCH	220
HERNEXEOS	108
<i> hydrocodone-acetaminophen oral tablet</i>	
<i> 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325</i>	
<i> mg</i>	224
<i> hydromorphone oral liquid</i>	224
<i> hydromorphone oral tablet 2 mg, 4 mg, 8</i>	
<i> mg</i>	224
<i> hydroxyzine hcl oral tablet</i>	109
HYRNUO	111
IBRANCE ORAL CAPSULE 125 MG	112
IBRANCE ORAL TABLET	112
IBTROZI	113
<i> icatibant</i>	90
ICLUSIG	114
IDHIFA	115
<i> imatinib oral tablet 100 mg, 400 mg</i>	101
IMBRUVICA ORAL CAPSULE 140	
MG, 70 MG	118
IMBRUVICA ORAL SUSPENSION	118
IMBRUVICA ORAL TABLET	118
<i> imipramine hcl</i>	109
IMKELDI	119
INCRELEX	121
INLURIYO	123
INLYTA	124
INQOVI	125
INREBIC	126
ITOVEBI ORAL TABLET 3 MG, 9 MG	
.....	134
<i> itraconazole oral capsule</i>	135
<i> ivabradine oral tablet 5 mg, 7.5 mg</i>	49
<i> ivermectin oral</i>	136
IWILFIN	137
JAKAFI	138
JASCAYD	139
JAVYGTOR	153
JAYPIRCA ORAL TABLET 100 MG,	
50 MG	141
JOENJA	142
KALYDECO ORAL GRANULES IN	
PACKET 13.4 MG, 25 MG, 5.8 MG, 50	
MG, 75 MG	144
KALYDECO ORAL TABLET	144
KERENDIA	145
KESIMPTA PEN	146
KINERET	147
KISQALI FEMARA CO-PACK ORAL	
TABLET 400 MG/DAY(200 MG X 2)-	
2.5 MG, 600 MG/DAY(200 MG X 3)-2.5	
MG	149
KISQALI ORAL TABLET 200	
MG/DAY (200 MG X 1), 400 MG/DAY	
(200 MG X 2), 600 MG/DAY (200 MG X	
3)	148
KOSELUGO ORAL CAPSULE 10 MG,	
25 MG	151

KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG	151	<i>memantine-donepezil</i>	182
KRAZATI	152	<i>methadone oral solution 10 mg/5 ml, 5 mg/5 ml</i>	224
KYGEVVI	154	<i>methadone oral tablet 10 mg, 5 mg</i>	224
<i>lapatinib</i>	303	<i>metirosine</i>	178
LAZCLUZE ORAL TABLET 240 MG, 80 MG	156	<i>mifepristone oral tablet 300 mg</i>	150
<i>lenalidomide</i>	243	<i>miglustat</i>	349
LENVIMA	157	<i>modafinil</i>	230
LEUKINE INJECTION RECON SOLN	159	MODEYSO	179
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i>	296	<i>morphine concentrate oral solution</i>	224
<i>lidocaine topical adhesive patch,medicated 5 %</i>	160	<i>morphine oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)</i>	224
<i>lidocaine topical ointment</i>	296	<i>morphine oral tablet</i>	224
<i>lidocaine-prilocaine topical cream</i>	296	<i>morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg</i>	224
<i>liraglutide</i>	104	MOUNJARO	104
LITFULO	161	MULPLETA	180
LIVTENCITY	162	NAYZILAM	183
LOARGYS	163	NEMLUVIO	184
LOKELMA	164	NERLYNX	186
<i>lomustine</i>	103	NEXLETOL	188
LONSURF	165	NEXLIZET	188
LORAZEPAM INTENSOL	224	<i>nilotinib hcl</i>	285
<i>lorazepam oral tablet 0.5 mg, 1 mg, 2 mg</i> ..	224	NINLARO	190
LORBRENA ORAL TABLET 100 MG, 25 MG	166	<i>nitisinone</i>	191
LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG	168	NORDITROPIN FLEXPRO SUBCUTANEOUS PEN INJECTOR 10 MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML (10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML)	106
<i>lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg</i>	155	NUBEQA	193
LYNPARZA	169	NUDEXTA	194
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)	172	NUPLAZID	195
MAVYRET ORAL PELLETS IN PACKET	173	NURTEC ODT	196
MAVYRET ORAL TABLET	173	<i>octreotide acetate injection solution</i>	199
<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)</i>	174	ODOMZO	200
<i>megestrol oral tablet</i>	174	OFEV	131
MEKINIST ORAL RECON SOLN	175	OGSIVEO ORAL TABLET 100 MG, 150 MG	201
MEKINIST ORAL TABLET 0.5 MG, 2 MG	175	OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	202
MEKTOVI	177	OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500 MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)	202
		OJJAARA	203

ONUREG	205	PLEGRIDY SUBCUTANEOUS	
OPIPZA	206	SYRINGE 125 MCG/0.5 ML	129
OPSUMIT	232	<i>pomalidomide</i>	218
OPSYNVI	232	<i>posaconazole oral tablet, delayed release</i>	
ORENCIA CLICKJECT	207	<i>(dr/ec)</i>	219
ORENCIA SUBCUTANEOUS		<i>pregabalin oral capsule 100 mg, 150 mg,</i>	
SYRINGE 125 MG/ML, 50 MG/0.4 ML,		<i>200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75</i>	
87.5 MG/0.7 ML	207	<i>mg</i>	171
ORGOVYX	209	<i>pregabalin oral solution</i>	171
ORKAMBI ORAL GRANULES IN		PRENATAL VITAMIN PLUS LOW	
PACKET	210	IRON	223
ORKAMBI ORAL TABLET	210	PREVYMIS ORAL PELLETS IN	
ORSERDU ORAL TABLET 345 MG,		PACKET	227
86 MG	211	PRIVIGEN	116
OTEZLA	212	PROLASTIN-C INTRAVENOUS	
OTEZLA STARTER ORAL		SOLUTION	9
TABLETS, DOSE PACK 10 MG (4)- 20		PROLIA	228
MG (51), 10 MG (4)-20 MG (4)-30 MG		<i>promethazine oral tablet</i>	109
(47)	212	PULMOZYME	234
OTEZLA XR	212	<i>pyrimethamine</i>	54
OTEZLA XR INITIATION	212	QINLOCK	235
<i>oxycodone oral capsule</i>	224	<i>quinine sulfate</i>	236
<i>oxycodone oral concentrate</i>	224	QULIPTA	237
<i>oxycodone oral tablet 10 mg, 15 mg, 20</i>		REPATHA SURECLICK	239
<i>mg, 30 mg, 5 mg</i>	224	REPATHA SYRINGE	239
<i>oxycodone-acetaminophen oral tablet 10-</i>		RETEVMO ORAL TABLET 120 MG,	
<i>325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i>		160 MG, 40 MG, 80 MG	241
.....	224	REVUFORJ ORAL TABLET 110 MG,	
OZEMPIC SUBCUTANEOUS	104	160 MG, 25 MG	244
PANRETIN	214	REXULTI ORAL TABLET	245
<i>paroxetine hcl oral suspension</i>	12	REZDIFFRA	246
<i>paroxetine hcl oral tablet</i>	12	REZLIDHIA	248
<i>pazopanib oral tablet 200 mg</i>	324	RINVOQ LQ	251
PEGASYS	128	RINVOQ ORAL TABLET EXTENDED	
PEMAZYRE	215	RELEASE 24 HR 15 MG, 30 MG, 45	
PHEBURANE	216	MG	249
<i>phenobarbital</i>	110	RIVFLOZA SUBCUTANEOUS	
PIQRAY ORAL TABLET 200		SOLUTION	252
MG/DAY (200 MG X 1), 250 MG/DAY		RIVFLOZA SUBCUTANEOUS	
(200 MG X1-50 MG X1), 300 MG/DAY		SYRINGE 128 MG/0.8 ML, 160	
(150 MG X 2)	217	MG/ML	252
<i>pirfenidone oral capsule</i>	131	ROMVIMZA	253
<i>pirfenidone oral tablet</i>	131	ROZLYTREK ORAL CAPSULE 100	
PLEGRIDY SUBCUTANEOUS PEN		MG, 200 MG	254
INJECTOR 125 MCG/0.5 ML	129	ROZLYTREK ORAL PELLETS IN	
		PACKET	254

RUBRACA	255	TAFINLAR ORAL CAPSULE	277
<i>rufinamide</i>	25	TAFINLAR ORAL TABLET FOR	
RYBELSUS	104	SUSPENSION	277
RYDAPT	256	TAGRISSE	279
SAJAZIR	90	TALZENNA	281
<i>sapropterin</i>	153	TASCENSO ODT	284
SCEMBLIX ORAL TABLET 100 MG,		<i>tasimelteon</i>	286
20 MG, 40 MG	260	<i>tazarotene topical cream</i>	287
SECUADO	261	TEPMETKO	289
SIGNIFOR	271	<i>teriflunomide</i>	17
<i>sildenafil (pulm.hypertension) oral tablet..</i>	232	<i>teriparatide subcutaneous pen injector 20</i>	
SILIQ	262	<i>mcg/dose (560mcg/2.24ml)</i>	93
SIMLANDI(CF) AUTOINJECTOR	220	<i>testosterone cypionate</i>	290
SIMLANDI(CF) SUBCUTANEOUS		<i>testosterone enanthate</i>	290
SYRINGE KIT 20 MG/0.2 ML, 40		<i>testosterone transdermal gel in metered-</i>	
MG/0.4 ML	220	<i>dose pump 20.25 mg/1.25 gram (1.62 %)</i> ...290	
SIRTURO	264	<i>tetrabenazine oral tablet 12.5 mg, 25 mg</i> ...335	
SKYCLARYS	265	THALOMID ORAL CAPSULE 100	
SKYRIZI SUBCUTANEOUS PEN		MG, 50 MG	291
INJECTOR	266	TIBSOVO	293
SKYRIZI SUBCUTANEOUS		TOBI PODHALER	42
SYRINGE	266	<i>tobramycin in 0.225 % nacl</i>	42
SKYRIZI SUBCUTANEOUS		<i>tobramycin inhalation</i>	42
WEARABLE INJECTOR 180 MG/1.2		<i>tolvaptan</i>	258
ML (150 MG/ML), 360 MG/2.4 ML (150		<i>tolvaptan (polycys kidney dis) oral tablet</i> ...143	
MG/ML)	266	<i>tolvaptan (polycys kidney dis) oral tablets,</i>	
<i>sodium oxybate</i>	345	<i>sequential</i>	143
<i>sodium phenylbutyrate</i>	34	<i>topiramate oral solution</i>	73
<i>sofosbuvir-velpatasvir</i>	71	<i>tramadol oral tablet 50 mg</i>	224
SOHONOS ORAL CAPSULE 1 MG, 1.5		<i>tramadol-acetaminophen</i>	224
MG, 10 MG, 2.5 MG, 5 MG	267	<i>tretinoin topical cream</i>	297
SOMAVERT	269	<i>tretinoin topical gel 0.01 %, 0.025 %</i>	297
<i>sorafenib</i>	187	TRIKAFTA ORAL GRANULES IN	
STELARA SUBCUTANEOUS		PACKET, SEQUENTIAL	298
SOLUTION	307	TRIKAFTA ORAL TABLETS,	
STELARA SUBCUTANEOUS		SEQUENTIAL	298
SYRINGE 45 MG/0.5 ML, 90 MG/ML ..	307	<i>trimipramine</i>	109
STEQEYMA SUBCUTANEOUS		TRINTELLIX	299
SYRINGE 45 MG/0.5 ML, 90 MG/ML ..	222	TRULICITY	104
STIVARGA	272	TRUQAP	300
<i>sunitinib malate</i>	273	TUKYSA ORAL TABLET 150 MG, 50	
SYMPAZAN	274	MG	301
SYNAREL	275	TURALIO	302
TABRECTA	276	TYMLOS	304
<i>tadalafil (pulm. hypertension)</i>	232	UBRELVY ORAL TABLET 100 MG,	
<i>tadalafil oral tablet 2.5 mg, 5 mg</i>	43	50 MG	305

UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG	306	XDEMVI	331
UPTRAVI ORAL TABLETS,DOSE PACK	306	XELJANZ ORAL SOLUTION	334
<i>ustekinumab subcutaneous solution</i>	307	XELJANZ ORAL TABLET	332
<i>ustekinumab subcutaneous syringe 45 mg/0.5 ml, 90 mg/ml</i>	307	XELJANZ XR	332
VALCHLOR	309	XERMELO	336
VALTOCO	310	XGEVA	337
VANFLYTA	311	XIFAXAN ORAL TABLET 550 MG	338
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG	312	XOLAIR	339
VENCLEXTA STARTING PACK	312	XOLREMDI	341
VERQUVO	313	XOSPATA	342
VERZENIO	314	XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80 MG/WEEK (80 MG X 1), 80MG TWICE WEEK (160 MG/WEEK)	343
<i>vigabatrin</i>	257	XTANDI ORAL CAPSULE	344
VIGADRONE	257	XTANDI ORAL TABLET 40 MG, 80 MG	344
VIJOICE ORAL GRANULES IN PACKET	316	YARGESA	349
VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50 MG	316	YESINTEK SUBCUTANEOUS SOLUTION	222
<i>vilazodone</i>	315	YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML ..	222
VITRAKVI ORAL CAPSULE 100 MG, 25 MG	317	YONSA	347
VITRAKVI ORAL SOLUTION	317	YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML, 294 MCG/0.98 ML, 420 MCG/1.4 ML	348
VIVJOA	318	YUFLYMA(CF)	220
VIZIMPRO	319	YUFLYMA(CF) AI CROHN'S-UC-HS ..	220
VONJO	320	YUFLYMA(CF) AUTOINJECTOR	220
VORANIGO ORAL TABLET 10 MG, 40 MG	321	<i>zaleplon oral capsule 10 mg, 5 mg</i>	224
<i>voriconazole intravenous recon soln</i>	322	ZAVZPRET	350
VOSEVI	323	ZEJULA ORAL TABLET	351
VOWST	86	ZELBORAF	352
VRAYLAR ORAL CAPSULE	325	ZELSUVMI	353
VUMERITY	326	ZELVYSIA ORAL POWDER IN PACKET 500 MG	153
WELIREG	327	ZEPOSIA	354
WINREVAIR	328	ZEPOSIA STARTER KIT (28-DAY)	354
XALKORI ORAL CAPSULE	329	ZEPOSIA STARTER PACK (7-DAY) ...354	
XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG	329		
XCOPRI	330		
XCOPRI MAINTENANCE PACK	330		
XCOPRI TITRATION PACK	330		

ZILBRYSQ SUBCUTANEOUS	
SYRINGE 16.6 MG/0.416 ML, 23	
MG/0.574 ML, 32.4 MG/0.81 ML.....	181
ZOKINVY	355
ZOLINZA	356
<i>zolpidem oral tablet.....</i>	224
ZONISADE.....	357
ZORYVE TOPICAL CREAM 0.15 %...358	
ZTALMY	359
ZURZUVAE ORAL CAPSULE 20 MG,	
25 MG, 30 MG.....	360
ZYKADIA	361

Section: Step Therapy

Entresto Tablet

Products Affected

- ENTRESTO 24 MG-26 MG TABLET
- ENTRESTO 49 MG-51 MG TABLET
- ENTRESTO 97 MG-103 MG TABLET

Details

Criteria	Require a 1 month trial of generic sacubitril/valsartan tablets (Step 1 drug) in the last 90 days
-----------------	---

GnRH Agonists

Products Affected

- **ELIGARD 22.5 MG (3 MONTH)
SUBCUTANEOUS SYRINGE**
- **ELIGARD 30 MG (4 MONTH)
SUBCUTANEOUS SYRINGE**
- **ELIGARD 45 MG (6 MONTH)
SUBCUTANEOUS SYRINGE**
- **ELIGARD 7.5 MG (1 MONTH)
SUBCUTANEOUS SYRINGE**
- **TRELSTAR 11.25 MG IM SUSPENSION**
- **TRELSTAR 22.5 MG IM SUSPENSION**
- **TRELSTAR 3.75 MG IM SUSPENSION**

Details

Criteria	Require a trial of Leuprolide 22.5mg (Step 1 drug) in the last 180 days when being utilized for the same medically accepted indication
-----------------	--

Herpetic Keratitis

Products Affected

- ZIRGAN 0.15 % EYE GEL

Details

Criteria	Require a 1 month trial of generic trifluridine eye drops (Step 1 drug) in the last 90 days when utilized for the same medically accepted indication and age
-----------------	--

Pulmonary Antiinflammatory

Products Affected

- *fluticasone propionate 100 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 110 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 220 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 250 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 44 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 50 mcg/actuation blister powder for inhalation*

Details

Criteria	Require a 1 month trial of Qvar and Asmanex/Asmanex HFA (Step 1 drugs) in the last 180 days
-----------------	---

Rho Kinase Inhibitors

Products Affected

- **RHOPRESSA 0.02 % EYE DROPS** **DROPS**
- **ROCKLATAN 0.02 %-0.005 % EYE**

Details

Criteria	Require a 1 month trial of one preferred glaucoma drug (Step 1 drug) in the last 120 days
-----------------	---

Rytary

Products Affected

- **RYTARY 23.75 MG-95 MG
CAPSULE,EXTENDED RELEASE**
- **RYTARY 36.25 MG-145 MG
CAPSULE,EXTENDED RELEASE**
- **RYTARY 48.75 MG-195 MG
CAPSULE,EXTENDED RELEASE**
- **RYTARY 61.25 MG-245 MG
CAPSULE,EXTENDED RELEASE**

Details

Criteria	Require a trial of generic carbidopa/levodopa product (Step 1 drug) in the last 90 days
-----------------	---

Index of Drugs

**ELIGARD 22.5 MG (3 MONTH)
SUBCUTANEOUS SYRINGE..... 2**

**ELIGARD 30 MG (4 MONTH)
SUBCUTANEOUS SYRINGE..... 2**

**ELIGARD 45 MG (6 MONTH)
SUBCUTANEOUS SYRINGE..... 2**

**ELIGARD 7.5 MG (1 MONTH)
SUBCUTANEOUS SYRINGE..... 2**

ENTRESTO 24 MG-26 MG TABLET 1

ENTRESTO 49 MG-51 MG TABLET 1

ENTRESTO 97 MG-103 MG TABLET 1

*fluticasone propionate 100 mcg/actuation
blister powder for inhalation..... 4*

*fluticasone propionate 110 mcg/actuation
hfa aerosol inhaler..... 4*

*fluticasone propionate 220 mcg/actuation
hfa aerosol inhaler..... 4*

*fluticasone propionate 250 mcg/actuation
blister powder for inhalation..... 4*

*fluticasone propionate 44 mcg/actuation
hfa aerosol inhaler..... 4*

*fluticasone propionate 50 mcg/actuation
blister powder for inhalation..... 4*

RHOPRESSA 0.02 % EYE DROPS..... 5

**ROCKLATAN 0.02 %-0.005 % EYE
DROPS..... 5**

**RYTARY 23.75 MG-95 MG
CAPSULE,EXTENDED RELEASE..... 6**

**RYTARY 36.25 MG-145 MG
CAPSULE,EXTENDED RELEASE..... 6**

**RYTARY 48.75 MG-195 MG
CAPSULE,EXTENDED RELEASE..... 6**

**RYTARY 61.25 MG-245 MG
CAPSULE,EXTENDED RELEASE..... 6**

**TRELSTAR 11.25 MG IM
SUSPENSION..... 2**

TRELSTAR 22.5 MG IM SUSPENSION.. 2

TRELSTAR 3.75 MG IM SUSPENSION.. 2

ZIRGAN 0.15 % EYE GEL..... 3