

Prior Authorization Criteria
Lantidra (donislecel-juju)

All requests for Lantidra (donislecel-juju) require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Coverage may be provided with a diagnosis of **Type I Diabetes** and the following criteria is met:

- Documentation the member has had Type I diabetes for more than 5 years accompanied by all the following despite intensive insulin management efforts (including insulin, devices, and education):
 - Unable to reach target HbA1C due to repeated episodes of severe hypoglycemia
 - At least one episode of severe hypoglycemia in the past 3 years defined as an event with symptoms compatible with hypoglycemia in which the member required the assistance of another person, and which was associated with either a blood glucose level < 50 mg/dL (2.8 mmol/L) or prompt recovery after oral carbohydrate, intravenous glucose, or glucagon administration
 - Prescriber attestation that the member reports reduced awareness of hypoglycemia as defined by the absence of adequate autonomic symptoms at capillary glucose levels of < 54 mg/dL (3 mmol/L) AND the member is unable to prevent repeated severe hypoglycemia events using intensive diabetes management (including insulin, devices, and education)
- Prescriber by or in consultation with an endocrinologist
- The member must not have any of the following:
 - Co-existing cardiac disease defined by one of the following:
 - Recent (within past 6 months) myocardial infarction
 - Angiographic evidence of non-correctable coronary artery disease
 - Evidence of ischemia on functional cardiac exam (with a stress echo test recommended for members with a history of ischemic disease).
 - Heart failure > New York Heart Association (NYHA) II
 - History of stroke within the past 6 months
 - Active alcohol or substance abuse-includes cigarette smoking (must be abstinent for six months). Active alcohol abuse should be considered using the current National Institute on Alcohol Abuse and Alcoholism (NIAAA) definitions.
 - Psychiatric disorder making the member not a suitable candidate for transplantation, e.g., schizophrenia, bipolar disorder, or major depression that is unstable or uncontrolled on current medication.
 - History of non-adherence to prescribed regimens
 - Active infection including hepatitis C, hepatitis B, HIV
 - TB (by history or currently infected as evidenced by a positive QuantiFERON® - TB Gold test or under treatment for suspected TB)
 - Any history of malignancies except squamous or basal skin cancer. Any member found to have squamous or basal cancer is required to have it removed prior to transplant.
 - Body Mass Index (BMI) > 27 kg/m².

- C-peptide response to glucagon stimulation (1 mg IV) (any C-peptide \geq 0.3 ng/mL)
- Creatinine clearance $<$ 80 mL/min/1.73 m² by 24-hour urine collection. If corrected creatinine clearance is $<$ 80 and serum creatinine is $<$ 1.2 mg/dL, then a nuclear renal scan is required to determine glomerular filtration rate.
- Serum creatinine consistently $>$ 1.5 mg/dL
- Macroalbuminuria (urinary albumin excretion rate $>$ 300 mg/24h)
- Baseline Hb $<$ 12 gm/dL in women or $<$ 13 gm/dL in men
- Baseline liver function tests (LFT) outside of normal range (An initial LFT test panel with any values $>$ 1.5 times normal upper limits will exclude a member without a re-test. A re-test for any values between normal and 1.5 times normal should be made, and if the values remain elevated above normal limits, the member will be excluded.)
- Untreated proliferative retinopathy
- Positive pregnancy test, or presently breast-feeding
- Insulin requirement $>$ 0.7 IU/kg/day
- HbA1c $>$ 12%
- Hyperlipidemia (fasting LDL cholesterol $>$ 130 mg/dL, treated or untreated; and/or fasting triglycerides $>$ 200 mg/dL)
- Under treatment for a medical condition requiring chronic use of steroids other than a previous organ transplant
- Use of coumadin or other antiplatelet or anticoagulant therapy, or member with PT INR $>$ 1.5. Low dose aspirin is allowed after transplantation.
- History of Factor V deficiency
- Addison's disease
- Allergy to radiographic contrast material
- Symptomatic cholecystolithiasis
- Acute or chronic pancreatitis
- Symptomatic peptic ulcer disease
- Severe unrelenting diarrhea, vomiting, or other gastrointestinal disorders that could interfere with the ability to absorb oral medications
- Received live attenuated vaccine(s) within 2 months
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
- Must be age-appropriate according to FDA-approved labeling, nationally recognized compendia, or evidence-based practice guidelines
- Must be used in conjunction with concomitant immunosuppression
- **Initial Duration of Approval:** 12 months
- **Reauthorization criteria**
 - Documentation the member has not achieved independence from exogenous insulin within one year of infusion OR within one year after losing independence from exogenous insulin after a previous infusion
- **Reauthorization Duration of Approval:** 12 months (maximum 3 infusions per lifetime)



Updated: 09/2024
PARP Approved: 09/2024

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.

**LANTIDRA (DONISLECEL-JUJU)
PRIOR AUTHORIZATION FORM**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

PROVIDER INFORMATION

Requesting Provider:	NPI:
Provider Specialty:	Office Contact:
Office Address:	Office Phone:
	Office Fax:

MEMBER INFORMATION

Member Name:	DOB:	
Member ID:	Member weight:	Height:

REQUESTED DRUG INFORMATION

Medication:	Strength:	
Directions:	Quantity:	Refills:

Is the member currently receiving requested medication? Yes No Date Medication Initiated:

Is this medication being used for a chronic or long-term condition for which the medication may be necessary for the life of the patient? Yes No

Billing Information

This medication will be billed: at a pharmacy **OR** medically, JCODE:

Place of Service: Hospital Provider's office Member's home Other

Place of Service Information

Name:	NPI:
Address:	Phone:

MEDICAL HISTORY (Complete for ALL requests)

Diagnosis:	ICD Code:
------------	-----------

How long has the member been diagnosed with Type I diabetes?

Does the member have any of the following (please check all that apply):

- the inability to reach target HbA1C due to repeated episodes of severe hypoglycemia
- History of at least one episode of severe hypoglycemia in the past 3 years defined as an event with symptoms compatible with hypoglycemia in which the member required the assistance of another person, and which was associated with either a blood glucose level < 50 mg/dL (2.8 mmol/L) or prompt recovery after oral carbohydrate, intravenous glucose, or glucagon administration
- reduced awareness of hypoglycemia as defined by the absence of adequate autonomic symptoms at capillary glucose levels of < 54 mg/dL (3 mmol/L) AND the member is unable to prevent repeated severe hypoglycemia events using intensive diabetes management (including insulin, devices, and education)
- Recent (within past 6 months) myocardial infarction
- Angiographic evidence of non-correctable coronary artery disease
- Evidence of ischemia on functional cardiac exam (with a stress echo test recommended for members with a history of ischemic disease).
- Heart failure > New York Heart Association (NYHA) II
- History of stroke within the past 6 months
- Active alcohol or substance abuse-includes cigarette smoking (must be abstinent for six months). Active alcohol abuse should be considered using the current National Institute on Alcohol Abuse and Alcoholism (NIAAA) definitions.
- Psychiatric disorder making the member not a suitable candidate for transplantation, e.g., schizophrenia, bipolar disorder, or major depression that is unstable or uncontrolled on current medication.
- History of non-adherence to prescribed regimens
- Active infection including hepatitis C, hepatitis B, HIV
- TB (by history or currently infected as evidenced by a positive QuantiFERON® -TB Gold test or under treatment for suspected TB)
- Any history of malignancies except squamous or basal skin cancer. Any member found to have squamous or basal cancer is required to have it removed prior to transplant.