



Updated: 02/2025
DMMA Approved: 02/2025

Request for Prior Authorization for Complement Inhibitors
Website Form – www.highmarkhealthoptions.com
Submit request via: Fax - 1-855-476-4158

All requests for Complement inhibitors require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Complement inhibitors include Soliris (eculizumab), Ultomiris (ravulizumab-cwvz), Empaveli (pegcetacoplan), Fabhalta (iptacopan), Enjaymo (sutimlimab-jome), Tavneos (avacopan) and Veopoz (pozelimab-bbfg), PiaSky (crovalimab-akkz), Voydeya (danicopan) and Zilbrysq (zilucoplan). New products with this classification will require the same documentation.

**** For all requests for complement inhibitors for the treatment of myasthenia gravis please refer to the Myasthenia Gravis Medications policy ****

For all requests for Complement Inhibitors all of the following criteria must be met:

- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
- Is age-appropriate according to FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature
- The member has received appropriate vaccinations as recommended in the FDA-approved package labeling unless contraindicated

Coverage may be provided with a diagnosis of **paroxysmal nocturnal hemoglobinuria (PNH)** and the following criteria are met:

- Medication is prescribed by, or in consultation with, a hematologist, oncologist, immunologist, or genetic specialist
- Member has a diagnosis of PNH confirmed by flow cytometry testing. Flow Cytometry pathology report must be supplied and demonstrate at least 2 different GPI protein deficiencies within 2 different cell lines from granulocytes, monocytes, or erythrocytes.
- Member is transfusion dependent as defined by having a transfusion within the last 12 months and one of the following:
 - Member's hemoglobin is less than or equal to 7 g/dL
 - Member has symptoms of anemia and the hemoglobin is less than or equal to 9 g/dL (**Soliris only**) or 10.5g/dL (**Empaveli only**)
 - Member has symptoms of anemia and the hemoglobin is less than or equal to 10 g/dL (**Ultomiris and Fabhalta only**)
- Must have a Lactate dehydrogenase (LDH) level at least 1.5 times the upper limit of the normal range (laboratory results with reference range must be submitted)
- If requesting Soliris, must have documentation of inadequate response, contraindication or intolerance to Ultomiris.
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria:**
 - Documentation of each of the following:
 - Documentation of a recent (within 3 months) LDH level that shows a reduction from baseline

- Documentation that hemoglobin has not dropped by more than 2 g/dL from baseline.
 - If baseline hemoglobin was less than 9g/dL, then the most recent hemoglobin must remain above 7g/dL
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **atypical hemolytic uremic syndrome (aHUS)** and the following criteria are met:

- The member weighs at least 5 kilograms
- Medication is prescribed by, or in consultation with, a hematologist, oncologist, immunologist, genetic specialist, or nephrologist
- Must provide documentation of hemolysis such as an elevation in serum LDH and serum creatinine above the upper limits of normal or required dialysis.
- The diagnosis of aHUS is supported by the absence of Shiga toxin-producing *E.coli* infection
- Must provide documentation member does not have a disintegrin and metalloproteinase with thrombospondin type 1 motif member 13 (ADAMTS13) deficiency
- If requesting Soliris, must have documentation of inadequate response, contraindication or intolerance to Ultomiris.
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Documentation from the provider that the member had a positive clinical response as evidenced by any of the following:
 - An increase in platelet count from baseline
 - Maintenance of normal platelet counts and LDH levels for at least four weeks
 - A 25% reduction in serum creatinine for a minimum of four weeks
 - The member has not experienced one of the following for at least 12 weeks after initiation of treatment:
 - A decrease in platelet count of >25% from baseline
 - Plasma exchange or plasma infusion
 - New dialysis requirement
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Neuromyelitis Optica Spectrum Disorder (NMOSD) (Soliris (eculizumab) and Ultomiris (ravulizumab-cwvz) ONLY)** and the following criteria are met:

- Medication is prescribed by, or in consultation with a neurologist
- Documentation of a positive test for AQP4-IgG antibodies
- The prescriber submits documentation of baseline number of relapse(s), which occurred over the last year.
- Documentation of an Expanded Disability Status Scale (EDSS) score of ≤ 7
- If using concurrent corticosteroids, dose is less than or equal to the equivalent of prednisone 20 mg per day
- Must have documentation of inadequate response, contraindication or intolerance to one (1) immunosuppressant (e.g., mycophenolate mofetil, azathioprine, methotrexate) or an inadequate response, contraindication or intolerance to rituximab or any of its biosimilars

- **Initial Duration of Approval:** 12 months
- **Reauthorization criteria**
 - Documentation the member has experienced a decrease from baseline in the number of NMOSD relapse(s).
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **cold agglutinin disease (CAD)** and the following criteria are met:

- Documentation of the diagnosis of CAD confirmed by:
 - Evidence of hemolysis indicated by both of the following:
 - Lactate dehydrogenase (LDH) level above the upper limit of normal AND
 - Haptoglobin level below the lower limit of normal
- Positive direct antiglobulin (Coombs) test for C3d only
- Cold agglutinin titer of ≥ 64 at 4 degrees Celsius
- Lack of overt malignant disease
- Documented history of at least one blood transfusion in the past 6 months
- Hemoglobin level ≤ 10.0 g/dL
- Bilirubin level above normal reference range, including patients with Gilbert's syndrome
- Presence of one or more symptoms associated with CAD including:
 - Symptomatic anemia
 - Acrocyanosis
 - Raynaud's phenomenon
 - Hemoglobinuria
 - Disabling circulatory symptoms
 - Major adverse vascular event
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Documentation of benefit from therapy including one of the following:
 - Increase in Hgb from baseline by ≥ 2 g/dL or achieving Hgb level of ≥ 12 g/dL
 - Normalization of LDH and/or bilirubin levels
 - Decrease in transfusion burden
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Complement Hyperactivation, Angiopathic Thrombosis, and Protein-Losing Enteropathy (CHAPLE)** and the following criteria are met:

- Must be prescribed by or in consultation with a provider who specializes in the treatment of CHAPLE disease
- Member has a diagnosis of CD55-deficient protein-losing enteropathy (PLE), also known as CHAPLE disease confirmed by biallelic CD55 loss-of-function mutation detected by genetic testing
- Member has active disease defined as hypoalbuminemia (serum albumin concentration ≤ 3.2 g/dL) with one or more of the following in the last 6 months:
 - Abdominal pain
 - Vomiting

- Diarrhea
- Peripheral or facial edema
- Infection with concomitant hypogammaglobinemia
- New thromboembolic event
- Member will not receive the requested drug in combination with another complement inhibitor
- Member has adequate titers or will receive meningococcal vaccines at least two weeks prior to the first dose of Veopoz
- **Initial Duration of Approval: 6 months**
- **Reauthorization criteria**
 - No evidence of unacceptable toxicity or disease progression while on the requested medication
 - Demonstrates a positive response to therapy as defined by ALL of the following:
 - Improvement or stabilization in disease activity (e.g., improvement of clinical symptoms [abdominal pain, diarrhea, and/or edema], increase in or stabilization of IgG concentrations, increase in growth percentiles, reduction in hospitalizations)
 - Normalization of serum albumin levels
- **Reauthorization Duration of Approval: 12 months**

Coverage may be provided with a diagnosis of **Anti-Neutrophil Cytoplasmic autoantibody (ANCA)-Associated Vasculitis** and the following criteria are met:

- Member must have severe and active ANCA- associated vasculitis
- Disease is one of the following types:
 - Granulomatosis with polyangiitis (GPA)
 - Microscopic polyangiitis (MPA)
- Member is positive for proteinase 3 antibodies, myeloperoxidase antibodies, or anti-neutrophil cytoplasmic autoantibody (ANCA)
- Must be prescribed by or in consultation with a rheumatologist,
- nephrologist, or immunologist
- A baseline Birmingham Vasculitis Activity Score (BVAS) has been performed
- Must be used as adjunctive treatment in combination with standard therapy (e.g., prednisone, azathioprine, mycophenolate, methotrexate, rituximab, cyclophosphamide)
- **Initial Duration of Approval: 6 months**
- **Reauthorization criteria**
 - Member experienced a beneficial clinical response from baseline exhibited by improvement in estimated glomerular filtration rate, decrease in urinary albumin creatinine ratio, or improvement in the BVAS from baseline
 - Member has experienced an improvement in at least one symptom, such as joint pain, ulcers, myalgia, persistent cough, skin rash, abdominal pain, or improvement in function or activities of daily living
- **Reauthorization Duration of Approval: 12 months**

Coverage may be provided with a diagnosis of **primary immunoglobulin A nephropathy (IgAN)** and the following criteria are met (**Fabhalta only**):

- The member has a diagnosis of IgAN confirmed by:
 - biopsy-proven IgAN
 - $\text{eGFR} \geq 20 \text{ mL/min/1.73 m}^2$
 - urine protein-to-creatinine ratio (UPCR) $\geq 1 \text{ g/g}$ on a stable dose of maximally-tolerated renin-angiotensin system (RAS) inhibitor therapy with or without a stable dose of an SGLT2 inhibitor
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria**
 - Member experienced a beneficial clinical response from baseline exhibited by improvement or reduction in UPCR from baseline
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

Drugs are authorized in generic form unless the branded product is on the preferred drug list or the prescriber has indicated in writing that the branded product is medically necessary. If only the branded product is on the preferred drug list, the generic form will be considered non-preferred and shall not require the prescriber to indicate in writing that the branded product is medically necessary.

Attachments

Attachment 1. Expanded Disability Status Scale (EDSS)

Score	Description
1.0	No disability, minimal signs in one functional system (FS)
1.5	No disability, minimal signs in more than one FS
2.0	Minimal disability in one FS
2.5	Mild disability in one FS or minimal disability in two FS
3.0	Moderate disability in one FS, or mild disability in three or four FS. No impairment to walking
3.5	Moderate disability in one FS and more than minimal disability in several others. No impairment to walking
4.0	Significant disability but self-sufficient and up and about some 12 hours a day. Able to walk without aid or rest for 500m
4.5	Significant disability but up and about much of the day, able to work a full day, may otherwise have some limitation of full activity or require minimal assistance. Able to walk without aid or rest for 300m
5.0	Disability severe enough to impair full daily activities and ability to work a full day without special provisions. Able to walk without aid or rest for 200m
5.5	Disability severe enough to preclude full daily activities. Able to walk without aid or rest for 100m
6.0	Requires a walking aid – cane, crutch, etc. – to walk about 100m with or without resting
6.5	Requires two walking aids – pair of canes, crutches, etc. – to walk about 20m without resting
7.0	Unable to walk beyond approximately 5m even with aid. Essentially restricted to wheelchair; though wheels self in standard wheelchair and transfers alone. Up and about in wheelchair some 12 hours a day
7.5	Unable to take more than a few steps. Restricted to wheelchair and may need aid in transferring. Can wheel self but cannot carry on in standard wheelchair for a full day and may require a motorised wheelchair
8.0	Essentially restricted to bed or chair or pushed in wheelchair. May be out of bed itself much of the day. Retains many self-care functions. Generally has effective use of arms
8.5	Essentially restricted to bed much of day. Has some effective use of arms retains some self-care functions
9.0	Confined to bed. Can still communicate and eat
9.5	Confined to bed and totally dependent. Unable to communicate effectively or eat/swallow
10.0	Death

COMPLEMENT INHIBITORS PRIOR AUTHORIZATION FORM-PAGE 1 of 3

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Health Options Pharmacy Services. **FAX:** (855) 476-4158

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (844) 325-6251 Mon-Fri 8 am to 7 pm

PROVIDER INFORMATION

Requesting Provider:	NPI:
Provider Specialty:	Office Contact:
Office Address:	Office Phone:
	Office Fax:

MEMBER INFORMATION

Member Name:	DOB:
Member ID:	Member weight: Height:

REQUESTED DRUG INFORMATION

Medication:	Strength:
Directions:	Quantity: Refills:
Is the member currently receiving requested medication? <input type="checkbox"/> Yes <input type="checkbox"/> No Date Medication Initiated:	
Is this medication being used for a chronic or long-term condition for which the medication may be necessary for the life of the patient? <input type="checkbox"/> Yes <input type="checkbox"/> No	

Billing Information

This medication will be billed: <input type="checkbox"/> at a pharmacy OR <input type="checkbox"/> medically, JCODE:
Place of Service: <input type="checkbox"/> Hospital <input type="checkbox"/> Provider's office <input type="checkbox"/> Member's home <input type="checkbox"/> Other

Place of Service Information


Name:	NPI:
Address:	Phone:

REFERENCE VALUES

Lab	Initial (Pre-Treatment) Value	Reference Range	Date	Post-Therapy Value (Reauthorization only)	Reference Range	Date
Hemoglobin (Hgb)						
Lactate dehydrogenase (LDH)						
Platelet count						
Serum Creatinine						
UPCR						

MEDICAL HISTORY (Complete for ALL requests)

Diagnosis:	ICD Code:
For Paroxysmal Nocturnal Hemoglobinuria (PNH) only: Does the member's flow cytometry pathology report demonstrate at least 2 different GPI protein deficiencies within 2 different cell lines from granulocytes, monocytes, or erythrocytes? Please include a copy <input type="checkbox"/> Yes <input type="checkbox"/> No Has the patient had a blood transfusion within the last 12 months? <input type="checkbox"/> Yes <input type="checkbox"/> No Does the patient have symptoms of anemia? <input type="checkbox"/> Yes <input type="checkbox"/> No	
For Atypical Hemolytic Uremic Syndrome only: Has the absence of Shiga toxin-producing <i>E.coli</i> been confirmed? <input type="checkbox"/> Yes <input type="checkbox"/> No Does the member have an ADAMTS13 deficiency? <input type="checkbox"/> Yes <input type="checkbox"/> No Is the member currently on dialysis? <input type="checkbox"/> Yes <input type="checkbox"/> No	



Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Health Options Pharmacy Services. **FAX:** (855) 476-4158

MEMBER INFORMATION

MEDICAL HISTORY (Complete for ALL requests)

What is the baseline number of relapse(s) which has occurred over the last year?_____

Does the member have one or more symptoms associated with CAD? ☐ Yes ☐ No

Has the member had adequate titers or will receive meningococcal vaccines at least two weeks prior to the first dose of Veopoz? ☐ Yes ☐ No

Does the member have a urine protein-to-creatinine ratio (UPCR) ≥ 1 g/g on a stable dose of maximally-tolerated renin-angiotensin system (RAS) inhibitor therapy with or without a stable dose of an SGLT2 inhibitor? ☐ Yes ☐ No

[illegible]

**CYTOKINE AND CAM ANTAGONISTS
PRIOR AUTHORIZATION FORM (CONTINUED) – PAGE 3 OF 3**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Health Options Pharmacy Services. **FAX:** (855) 476-4158

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (844) 325-6251 Mon – Fri 8 am to 7 pm

MEMBER INFORMATION

Member Name:	DOB:	
Health Options ID:	Member weight:	Height:

**MEDICAL HISTORY (Complete for ALL requests)
REAUTHORIZATION**

Has the member experienced a significant improvement with treatment? ☐ Yes (documentation required) ☐ No

For Paroxysmal Nocturnal Hemoglobinuria (PNH) only:

Has the patient had a blood transfusion since taking Soliris? ☐ Yes ☐ No

For Atypical Hemolytic Uremic Syndrome only:

Has the patient been able to maintain a normal platelet or LDH level for at least four weeks? ☐ Yes ☐ No

Has the patient experienced a 25% serum creatinine reduction for at least four weeks? ☐ Yes ☐ No

In the past 12 weeks, has the patient had any of the following?

A decrease in platelet count of >25% from baseline ☐ Yes ☐ No

Increased need for plasma exchange or plasma infusion ☐ Yes ☐ No

New dialysis requirement ☐ Yes ☐ No

For Neuromyelitis Optica Spectrum Disorder:

Has the member experienced a decrease from baseline in the number of NMOSD relapse(s)? ☐ Yes ☐ No

For Cold Agglutinin Disease (CAD):

Is there documentation of benefit from therapy including one of the following:

- Increase in Hgb from baseline by ≥ 2 g/dL or achieving Hgb level of ≥ 12 g/dL ☐ Yes ☐ No
- Normalization of LDH and/or bilirubin levels ☐ Yes ☐ No
- Decrease in transfusion burden ☐ Yes ☐ No

For CHAPLE disease:

Is there evidence of unacceptable toxicity or disease progression while on the requested medication? ☐ Yes ☐ No

Has the member demonstrated a positive response defined by ALL of the following? ☐ Yes ☐ No

- Improvement or stabilization in disease activity (e.g., improvement of clinical symptoms [abdominal pain, diarrhea, and/or edema], increase in or stabilization of IgG concentrations, increase in growth percentiles, reduction in hospitalizations)
- Normalization of serum albumin levels

For Primary immunoglobulin A nephropathy (IgAN):

Has the member experienced a beneficial clinical response from baseline exhibited by improvement or reduction in UPCR from baseline? ☐ Yes ☐ No

SUPPORTING INFORMATION or CLINICAL RATIONALE

Prescribing Provider Signature	Date