

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS:

ABRILADA™ (adalimumab-afzb) SQ

Adalimumab-aacf SQ

Adalimumab-aaty SQ

Adalimumab-adaz SQ

Adalimumab-adbm SQ

Adalimumab-fkjp SQ

Adalimumab-ryvk SQ

AMJEVITA™ (adalimumab-atto) SQ by Amgen

AMJEVITA™ (adalimumab-atto) SQ by Optum Health Solutions Limited

CYLTEZO® (adalimumab-adbm) SQ

HADLIMA™ (adalimumab-bwwd) SQ

HULIO® (adalimumab-fkjp) SQ

HUMIRA® (adalimumab) SQ

HYRIMOZ® (adalimumab-adaz) SQ

IDACIO® (adalimumab-aacf) SQ

SIMLANDI® (adalimumab-ryvk) SQ

YUFLYMA® (adalimumab-aaty) SQ

YUSIMRY™ (adalimumab-aqvh) SQ

This Pharmacy Coverage Guideline (PCG):

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

Scope

- This PCG applies to Commercial and Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

Instructions & Guidance

- To determine whether a member is eligible for the Service, read the entire PCG.
- This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
- Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
- The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
- The “Description” section describes the Service.
- The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
- The “Resources” section lists the information and materials we considered in developing this PCG
- **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
- Information about medications that require prior authorization is available at www.azblue.com/pharmacy. You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 6/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to Pharmacyprecert@azblue.com.

Criteria:

Section A. Ankylosing Spondylitis (AS):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderately to severely active ankylosing spondylitis:
 1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Rheumatologist
 3. Individual is 18 years of age or older
 4. Clinical and diagnostic imaging evidence of ankylosing spondylitis as indicated by **ALL** of the following:
 - a. Back pain of 3 months or more duration with an age of onset of 45 years or younger
 - b. Sacroiliitis on x-ray imaging **showing** definitive radiographic evidence of **structural damage of sacroiliac joints**
 - c. Spondyloarthritis signs or symptoms as indicated by **ONE or more** of the following:
 - i. Arthritis
 - ii. Elevated serum C-reactive protein
 - iii. Enthesitis (e.g., inflammation of Achilles tendon insertion)
 - iv. HLA-B27
 - v. Limited chest expansion
 - vi. Morning stiffness for one hour or more
 - d. A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 or more, and spinal pain rated as at least 4 or more on a 0 to 10 numerical rating scale
 5. Disease activity and treatment scenario as indicated by **ONE or more** of the following:
 - a. Axial (spinal) disease
 - b. Peripheral arthritis without axial involvement, and failure, contraindication per FDA label, or intolerance to 4 or more months of therapy with sulfasalazine
 6. Individual has documented failure, contraindication per FDA label, intolerance, or is not a candidate for **TWO or more** different NSAIDs (at maximum recommended doses) over a total period of at least 4 or more weeks of therapy
 7. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited,

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- c. Hadlima
 - d. Simlandi
8. Individual does **NOT** have **ANY** of the following:
- a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
9. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):
- 1. Individual continues to be seen by a physician specializing in or is in consultation with a Rheumatologist
 - 2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 20% improvement in BASDAI ([see Definitions section](#))
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
 - 3. Individual has been adherent with the medication
 - 4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
 - 5. Individual has not developed any significant adverse drug effects that may exclude continued use
 - 6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section B. Crohn's Disease (CD):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderately to severely active Crohn's disease:
 1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Gastroenterologist
 3. Individual is 6 years of age or older
 4. Individual has a confirmed diagnosis of moderate to severe active Crohn's disease as indicated by **ONE** of the following:
 - a. Crohn's disease activity index (CDAI) greater than 220 in adults
 - b. Pediatric Crohn's disease activity index (PCDAI) greater than 30
 - c. **At least 5** of the following signs and symptoms:
 - i. Anemia
 - ii. Chronic intermittent diarrhea (with or without food)
 - iii. Crampy abdominal pain
 - iv. Elevated serum C-reactive protein level and/or fecal calprotectin
 - v. Extraintestinal manifestations such as arthritis or arthropathy, eye and skin disorders, biliary tract involvement, and kidney stones
 - vi. Fatigue
 - vii. Fistulas
 - viii. Perianal disease (e.g., anal fissures, anorectal abscess)
 - ix. Weight loss or growth failure in children
 5. Individual has documented failure (used for ≥ 3 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **ONE or MORE** of the following [Note this criterion is waived if the individual already has tried an FDA-approved Crohn's disease biologic]:
 - a. 6-mercaptopurine
 - b. Azathioprine
 - c. Methotrexate
 - d. Oral corticosteroids
 6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
7. Individual does **NOT** have **ANY** of the following:
- a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in or is in consultation with a Gastroenterologist
2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation ONE of the following:**
 - i. AT LEAST a 20% improvement in the signs and symptoms of Crohn's disease
 - ii. Decrease in Crohn's disease activity index of more than 70 from baseline or a Crohn's disease activity index of < 150 (in remission) in adults
 - iii. Pediatric Crohn disease activity index (PCDAI) ≤ 30 in children indicating mild disease or disease remission
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
3. Individual has been adherent with the medication
4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section C. Juvenile Idiopathic Arthritis (JIA) subtype: Polyarticular (pJIA):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for juvenile idiopathic arthritis subtype polyarticular JIA:
1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Rheumatologist
 3. Individual is 2 years of age or older
 4. Individual has arthritis in **five** or more joints during the first six months of disease and **NONE** of the following:
 - a. Fever, rash, lymphadenopathy, hepatosplenomegaly
 - b. Arthritis starting after 6 years of age in male individual who is positive for HLA-B27
 - c. Personal history or first degree relative with psoriasis, ERA, ankylosing spondylitis, sacroiliitis with IBD, reactive arthritis, anterior uveitis
 5. Individual has documented failure, contraindication per FDA label, intolerance, or is not a candidate for methotrexate
 6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
 7. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
- b. Concurrent use of live vaccines
8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in or is in consultation with a Rheumatologist
2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 30% improvement in JIA Core Set ([see Definitions section](#))
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
3. Individual has been adherent with the medication
4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Section D. Plaque Psoriasis (Ps also as PsO):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderate to severe plaque psoriasis:
1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Dermatologist
 3. Individual is 18 years of age or older
 4. Individual has a diagnosis of moderate to severe plaque psoriasis, as indicated by **ALL** of the following:
 - a. Individual is a candidate for photochemotherapy or phototherapy
 - b. Plaque psoriasis involves $\geq 10\%$ body surface area (BSA) **or** plaque psoriasis involves $< 10\%$ BSA but includes sensitive areas or areas that significantly impact daily function (e.g., palms, soles of feet, head/neck, or genitalia)
 - c. A Psoriasis Area and Index (PASI) of at least 10
 5. Individual has documented failure (used for ≥ 3 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for a treatment regimen that includes **ALL** of the following:
 - a. A trial of least **TWO** topical agents (e.g., anthralin, calcipotriene, coal tars, corticosteroids, tazarotene)
 - b. A trial of **ONE** immunosuppressive treatment (e.g., cyclosporine, methotrexate)
 - c. A trial of Ultraviolet Light therapy (e.g., Photochemotherapy (i.e., psoralen plus ultraviolet A therapy), Phototherapy (i.e., ultraviolet light therapy), or Excimer laser)
 6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
 7. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
 8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):
1. Individual continues to be seen by a physician specializing in or is in consultation with a Dermatologist
 2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 20% improvement in PASI ([see Definitions section](#))
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
 3. Individual has been adherent with the medication
 4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
 5. Individual has not developed any significant adverse drug effects that may exclude continued use
 6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
 7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section E. Psoriatic Arthritis (PsA):

- **Criteria for initial therapy:** Humira or Humira Biosimilars considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderately to severely active psoriatic arthritis:

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
2. Prescriber is a Rheumatologist or Dermatologist
3. Individual is 18 years of age or older
4. Individual has a confirmed diagnosis of moderate to severe active psoriatic arthritis is identified by **ONE or more** of the following:
 - a. Predominantly axial disease (i.e., sacroiliitis or spondylitis) as indicated by **ALL** of the following:
 - i. Radiographic evidence of axial disease (e.g., sacroiliac joint space narrowing or erosions, vertebral syndesmophytes)
 - ii. Symptoms (e.g., limited spinal range of motion, spinal morning stiffness more than 30 minutes) present for more than 3 months duration
 - iii. Failure, contraindication per FDA label, or intolerance to 1 or more different NSAIDs (at maximum recommended doses) over total period of at least 4 or more weeks of therapy
 - b. Predominantly non-axial disease, and failure (used for ≥ 3 consecutive months), intolerance, or contraindication per FDA label to methotrexate or NSAIDs
5. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):
1. Individual continues to be seen by a physician specializing in or is in consultation with a Rheumatologist or Dermatologist
 2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 20% improvement in any of the following: ACR, CDAI, DAS28, PAS, PASII, RAPID-3, SDAI ([see Definitions section](#))
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
 3. Individual has been adherent with the medication
 4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
 5. Individual has not developed any significant adverse drug effects that may exclude continued use
 6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
 7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section F. Rheumatoid Arthritis (RA):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderately to severely active rheumatoid arthritis:
1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Rheumatologist
 3. Individual is 18 years of age or older

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

4. Individual has a confirmed diagnosis of rheumatoid arthritis identified by **ONE** of the following:
 - a. Clinical Disease Activity Index (CDAI) score greater than 10
 - b. Disease Activity Score 28 (DAS28) of greater than 3.2
 - c. Patient Activity Scale (PAS) of greater than 3.7
 - d. Patient Activity Scale II (PASII) of greater than 3.7
 - e. Routine Assessment of Patient Index Data 3 (RAPID-3) score greater than 2
 - f. Simplified Disease Activity Index (SDAI) score greater than 11
5. Individual has documented failure (used for ≥ 3 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **methotrexate**
6. Individual has documented failure (used for ≥ 3 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **ONE** of the following: [Note this criterion is waived if the individual already has tried an FDA-approved Rheumatoid Arthritis biologic]
 - a. Leflunomide
 - b. Sulfasalazine
7. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
8. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
9. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in or is in consultation with a Rheumatologist
2. Individual's condition has responded while on therapy with response defined as the following:

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- a. **With first request for continuation:** AT LEAST a 20% improvement in any of the following: ACR, CDAI, DAS28, PAS, PASII, RAPID-3, SDAI ([see Definitions section](#))
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
3. Individual has been adherent with the medication
4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section G. Ulcerative Colitis (UC):

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderately to severely active ulcerative colitis (UC):
1. Request is for **ONE** of the following: Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, Yusimry
 2. Prescriber is a Gastroenterologist
 3. Individual is 5 years of age or older
 4. Individual has a confirmed diagnosis of moderate to severe active ulcerative colitis, as indicated by **ONE** of the following:
 - a. American College of Gastroenterology Ulcerative Colitis activity index rating of moderate to severe disease in adults
 - b. Pediatric ulcerative colitis activity index (PUCAI) greater than or equal to 35
 - c. **At least 5** of the following signs and symptoms:

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- i. Anemia
 - ii. Bloody diarrhea or visible blood in stool
 - iii. Bowel movements 4-6 or more times per day
 - iv. Colicky abdominal pain
 - v. Elevated fecal calprotectin
 - vi. Elevated serum C-reactive protein or erythrocyte sedimentation rate
 - vii. Fatigue
 - viii. Fever
 - ix. Tenesmus
 - x. Urgency
 - xi. Weight loss or delayed growth in children
5. Individual has documented failure (used for ≥ 3 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **ONE or more** of the following: [Note: this criterion is waived if the individual already has tried an FDA-approved Ulcerative Colitis biologic]
 - a. 6-mercaptopurine
 - b. Azathioprine
 - c. Oral corticosteroids
 - d. Salicylates (such as mesalamine, sulfasalazine, balsalazide, olsalazine)
6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
7. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

1. Individual continues to be seen by a physician specializing in or is in consultation with a Gastroenterologist
2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation ONE of the following:**
 - i. AT LEAST a 20% improvement in signs and symptoms of ulcerative colitis
 - ii. American College of Gastroenterology Ulcerative Colitis activity index rating of mild disease or disease in remission in adults
 - iii. Pediatric ulcerative colitis activity index (PUCAI) of ≤ 34 in children indicating mild disease or disease remission
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
3. Individual has been adherent with the medication
4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section H. Hidradenitis Suppurativa:

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderate to severe hidradenitis suppurativa:
1. Request is for Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, or Yusimry
 2. Prescriber is a Dermatologist

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

3. Individual is 12 years of age or older
4. Diagnosis of moderate to severe disease as indicated by **ONE or more** of the following:
 - a. Multiple interconnected tracts and abscesses in single anatomic area
 - b. Widely separated and recurrent abscesses with sinus tracts and scarring
5. Individual has documented failure, contraindication per FDA label, intolerance, or is not a candidate for oral antibiotics (at maximum recommended doses) for at least 3 consecutive months (i.e., tetracycline, clindamycin plus rifampin, minocycline, doxycycline)
6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, or Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
7. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
 - b. Concurrent use of live vaccines
8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):
1. Individual continues to be seen by a physician specializing in or is in consultation with a Dermatologist
 2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 20% improvement in the signs and symptoms of hidradenitis suppurativa
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
 3. Individual has been adherent with the medication

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, or Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:
 - a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section I. Uveitis:

- **Criteria for initial therapy:** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met for moderate non-infectious intermediate uveitis, non-infectious posterior uveitis or non-infectious panuveitis:
 1. Request is for Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-adbm, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Amjevita by Optum Health Solutions Limited, Cyltezo, Hadlima, Hulio, Humira, Hyrimoz, Idacio, Simlandi, Yuflyma, or Yusimry
 2. Prescriber is an Ophthalmologist
 3. Individual is 2 years of age or older
 4. Individual has a confirmed diagnosis of non-infectious intermediate, posterior, or panuveitis
 5. Individual has documented failure, contraindication per FDA label, intolerance, or is not a candidate for **ONE** agent for **BOTH** categories:
 - a. Corticosteroids (> 2-week trial at up to maximally indicated doses)
 - b. Systemic immunosuppressant (i.e., methotrexate, cyclosporine, azathioprine, mycophenolate, cyclophosphamide, leflunomide, hydroxychloroquine, sulfasalazine, tacrolimus, sirolimus, or chlorambucil)
 6. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira, Hyrimoz, Idacio, Yuflyma, or**

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Yusimry: Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:

- a. Adalimumab-adbm
- b. Amjevita by Optum Health Solutions Limited
- c. Hadlima
- d. Simlandi

7. Individual does **NOT** have **ANY** of the following:

- a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - i. Serologic tests for hepatitis B and C (HB surface Ag, anti-HB surface Ab, anti-HB core Ab, and hepatitis C antibody tests) have been done within the previous 12 months
 - ii. Screening for latent tuberculosis infection with a tuberculin skin test or blood test has been done and if positive, treatment has been initiated
- b. Concurrent use of live vaccines

8. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Approval Duration: 6 months

➤ **Criteria for continuation of coverage (renewal request):** Humira or Humira Biosimilars is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in or is in consultation with an Ophthalmologist
2. Individual's condition has responded while on therapy with response defined as the following:
 - a. **With first request for continuation:** AT LEAST a 20% improvement in the signs and symptoms of uveitis or panuveitis
 - b. **With subsequent request for continuation:** Documented evidence of disease stability and/or improvement with no evidence of disease progression
3. Individual has been adherent with the medication
4. **For Abrilada, adalimumab-aacf, adalimumab-aaty, adalimumab-adaz, adalimumab-fkjp, adalimumab-ryvk, Amjevita by Amgen, Cyltezo, Hulio, Humira (effective 7/1/2025), Hyrimoz, Idacio, Yuflyma, Yusimry:** Individual has documented failure (used for ≥ 6 consecutive months), contraindication per FDA label, intolerance, or is not a candidate for **THREE** of the following:
 - a. Adalimumab-adbm
 - b. Amjevita by Optum Health Solutions Limited
 - c. Hadlima
 - d. Simlandi
5. Individual has not developed any significant adverse drug effects that may exclude continued use
6. Individual does **NOT** have **ANY** of the following:

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

- a. Evidence of active serious infections including, opportunistic infections, fungal infections, tuberculosis, clinically important localized infections, sepsis, Hepatitis B, or Hepatitis C
 - b. Concurrent use of live vaccines
7. There is no concomitant use with biologic immunomodulators or other potent immunosuppressants (e.g., Adbry, azathioprine, cyclosporine, Dupixent, rituximab, infliximab, Enbrel, Otezla, Vtama, Xolair, or JAK inhibitors (Cibinqo, Olumiant, Opzelura, Rinvoq, Rinvoq LQ, Xeljanz IR, XR, solution), etc.)

Renewal Duration: 12 months

Section J. Measurement of Antibodies to Biologic/Immunologic Agents:

- Measurement of antibodies for biologic or immunologic agents in an individual receiving treatment, either alone or as a combination test, which includes the measurement of serum levels for the biologic or immunologic agents is considered **experimental or investigational** when any **ONE** or more of the following criteria are met:
1. Lack of final approval from the appropriate governmental regulatory bodies (e.g., Food and Drug Administration); or
 2. Insufficient scientific evidence to permit conclusions concerning the effect on health outcomes; or
 3. Insufficient evidence to support improvement of the net health outcome; or
 4. Insufficient evidence to support improvement of the net health outcome as much as, or more than, established alternatives; or
 5. Insufficient evidence to support improvement outside the investigational setting.

These measurements include, *but are not limited to*:

- Anser™ ADA

Section K. Other:

- Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:
1. **Off-Label Use of Non-Cancer Medications**
 2. **Off-Label Use of Cancer Medications**
-

Definitions:

Adult: Age 18 years and older.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Bath Ankylosing Spondylitis Disease Activity Index (BASDAI):

1. How would you describe the overall level of fatigue/tiredness you have experienced?	None 0 1 2 3 4 5 6 7 8 9 10 Very Severe
2. How would you describe the overall level of ankylosing spondylitis neck, back or hip pain you have had?	None 0 1 2 3 4 5 6 7 8 9 10 Very Severe
3. How would you describe the overall level of pain/swelling you have had in joints other than neck, back and hips?	None 0 1 2 3 4 5 6 7 8 9 10 Very Severe
4. How would you describe the level of discomfort you have had from an area tender to touch or pressure?	None 0 1 2 3 4 5 6 7 8 9 10 Very Severe
5. How would you describe the level of morning stiffness you have had from the time you wake up?	None 0 1 2 3 4 5 6 7 8 9 10 Very Severe
6. How long does your morning stiffness last from the time you wake up?	0 hours 0 1 2 3 4 5 6 7 8 9 10 2 or more hours

Calculation of BASDAI:

Compute the mean of questions 5 and 6

Calculate the sum of the values of question 1-4 and add the result to the mean of questions 5 and 6

© 2018 UpToDate, Inc. Originally published in: Garrett S, Jenkinson T, Kennedy LG, et al. A new approach to defining disease status in ankylosing spondylitis: the Bath Ankylosing Spondylitis Disease Activity Index. J Rheumatol 1994; 21:2286. Reproduced with permission from: the Royal National Hospital for Rheumatic Diseases NHS Foundation Trust, Bath. www.rnhrd.nhs.uk. Copyright ©

Crohn's Disease Activity Index:

Sum each factor after adjustment with a weighting factor

Clinical or laboratory variable	Weighting factor	Factor Sum
Number of liquid or soft stools each day for seven days	x 2	
Abdominal pain (graded 0 = none, 1 = mild, 2 = moderate, 3 = severe) each day for 7 days	x 5	
General well-being (assessed from 0 = well, 1 = slightly under par, 2 = poor, 3 = very poor, 4 = terrible) each day for 7 days	x 7	
Presence of complications†	x 20	
Taking Lomotil (diphenoxylate/atropine) or opiates for diarrhea (0 = No, 1 = Yes)	x 30	
Presence of an abdominal mass (0 = none, 2 = questionable, 5 = definite)	x 10	
Hematocrit of < 0.47 in men and < 0.42 in women	x 6	
Percentage deviation from standard weight [1 – (ideal/observed)] x 100	x 1	

† **Complications:** one point each is added for each:

- the presence of joint pains (arthralgia) or frank arthritis
- inflammation of the iris or uveitis
- presence of erythema nodosum, pyoderma gangrenosum, or aphthous ulcers
- anal fissures, fistulae or abscesses
- other fistulae
- fever during the previous week

Total CDAI

Remission of CD: CDAI < 150

Severe CD: CDAI > 450

CD response: decrease in CDAI of > 70

Pediatric Crohn disease activity index (PCDAI):

HISTORY: Recall from previous week		
Abdominal Pain	None	0 points

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

	Mild – Brief, does not interfere with activities	5 points
	Moderate or severe – Daily, longer lasting, affects activities, nocturnal	10 points
Stools (per day)	0-1 liquid stools, no blood	0 points
	Up to 2 semi-formed stools with small blood, or 2-5 liquid stools without blood	5 points
	Gross bleeding, or ≥ 6 liquid stools, or nocturnal diarrhea	10 points
Patient functioning, general well-being	No limitations of activities, well	0 points
	Occasional difficulty in maintaining age-appropriate activities, below par	5 points
	Frequent limitation of activity, very poor	10 points
Laboratory		
Hematocrit (%) <10 years	>33	0 points
	28 to 32	2.5 points
	<28	5 points
Hematocrit (%) 11-19 years (females)	≥ 34	0 points
	29 to 33	2.5 points
	<29	5 points
Hematocrit (%) 11-14 years (males)	≥ 35	0 points
	30 to 34	2.5 points
	<30	5 points
Hematocrit (%) 15 to 19 years (male)	≥ 37	0 points
	32 to 36	2.5 points
	<32	5 points
ESR (mm/hour)	<20	0 points
	20 to 50	2.5 points
	>50	5 points
Albumin (g/dl)	≥ 3.5	0 points
	3.1 to 3.4	5 points
	≤ 3	10 points
Examination		
Weight	Weight gain, weight stable, or voluntary weight loss	0 points
	Involuntary weight stable, or weight loss 1 to 9%	5 points
	Weight loss $\geq 10\%$	10 points
Height (at diagnosis)	<1 channel decrease*	0 points
	1 to 2 channel decrease	5 points
	≥ 2 channel decrease	10 points
Height (at follow-up)	High velocity ≥ -1 SD	0 points
	High velocity between -1 and -2 SD	5 points
	High velocity ≤ -2 SD	10 points
Abdomen	No tenderness, no mass	0 points
	Tenderness, or mass without tenderness	5 points
	Tenderness, involuntary guarding, definite mass	10 points
Perirectal disease	None, asymptomatic tags	0 points
	1 to 2 indolent fistula(e), scant drainage, no tenderness	5 points
	Active fistula, drainage, tenderness, or abscess	10 points

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Extraintestinal manifestations (Fever $\geq 38.5^{\circ}\text{C}$ for 3 days over past week, definite arthritis, uveitis, erythema nodosum, pyoderma gangrenosum)	None	0 points
	1	5 points
	≥ 2	10 points
<p>The PCDAI is interpreted as follows: a score of 0 to 10 indicates inactive disease, 11 to 30 indicates mild disease activity, and >30 indicates moderate to severe disease activity. A decrease in PCDAI of ≥ 12.5 points reflects a clinical response (improvement from moderate/severe to mild/inactive disease)</p> <p>ESR: erythrocyte sedimentation rate; SD: standard deviation.</p> <p>* A "channel decrease" refers to serial height measurements that deviate across the width of a major curve on a standard height-for-age chart. For example, decreasing from the 40th to 20th percentile is a 1-channel decrease.</p>		

Psoriasis Area and Severity Index (PASI):

	Head	Upper Extremities	Trunk	Lower extremities
1. Redness ¹				
2. Thickness ¹				
3. Scale ¹				
4. Sum of rows 1,2 and 3				
5. Area score ²				
6. Score of row 4 x row 5 x the area multiplier	row 4 x row 5 x 0.1	row 4 x row 5 x 0.2	Row 4 x row 5 x 0.3	Row 4 x row 5 x 0.4
7. Sum row 6 for each column for PASI score				

Steps in generating PASI score:

- Divide body into four areas: head, arms, trunk to groin, and legs to top of buttocks.
- Generate an average score for the erythema, thickness, and scale for each of the 4 areas (0 = clear; 1–4 = increasing severity)¹.
- Sum scores of erythema, thickness, and scale for each area.
- Generate a percentage for skin covered with psoriasis for each area and convert that to a 0–6 scale (0 = 0%; 1 = $<10\%$; 2 = $10\text{--}<30\%$; 3 = $30\text{--}<50\%$; 4 = $50\text{--}<70\%$; 5 = $70\text{--}<90\%$; 6 = $90\text{--}100\%$).
- Multiply score of item (c) above times item (d) above for each area and multiply that by 0.1, 0.2, 0.3, and 0.4 for head, arms, trunk, and legs, respectively.
- Add these scores to get the PASI score.

¹ Erythema, induration and scale are measured on a 0–4 scale (none, slight, mild, moderate, severe)

² Area scoring criteria (score: % involvement)

- 0: 0 (clear)
 1: $<10\%$
 2: $10\text{--}<30\%$
 3: $30\text{--}<50\%$
 4: $50\text{--}<70\%$
 5: $70\text{--}<90\%$
 6: $90\text{--}<100\%$

Feldman, SR and Krueger, GG. Psoriasis assessment tools in clinical trials. *Ann Rheum Dis* 2005; 64 (Suppl III): ii65-ii68.

JIA Core Set 30%:

At least 30 percent improvement in at least 3 of the 6 core set variables with no more than 1 remaining variable worsening by $> 30\%$	
1.	Physician's global assessment of overall disease activity measured on a visual analog scale (VAS)
2.	Parent or patient global assessment of overall well-being measured on VAS
3.	Functional ability
4.	Number of joints with active arthritis
5.	Number of joints with limited range of motion

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

6. Erythrocyte sedimentation rate (ESR)

Giannini, EH, Ruperto, N, Ravelli A, et al. Preliminary Definition of Improvement in Juvenile Arthritis. *Arthritis & Rheumatism* 1997

Rheumatoid Arthritis Disease Activity Measurement Instruments:

Instrument	Threshold of Disease Activity
Clinical Disease Activity Index (CDAI)	Range: 0 to 76 Remission: ≤ 2.8 Low activity: >2.8 to ≤ 10 Moderate activity: >10 to ≤ 22 High activity: >22
Disease Activity Score 28 (DAS28)	Range: 0.5 to 9 Remission: < 2.6 Low activity: > 2.6 to ≤ 3.2 Moderate activity: > 3.2 to ≤ 5.1 High activity: > 5.1
Patient Activity Scale (PAS) Patient Activity Scale II (PASII)	Range 0 to 10 Remission: 0 to 0.25 Low activity: >0.25 to 3.7 Moderate activity: > 3.7 to < 8.0 High activity: ≥ 8.0
Routine Assessment of Patient Index Data 3 (RAPID-3)	Range: 0 to 10 Remission: 0 to 1.0 Low activity: > 1.0 to 2.0 Moderate activity: > 2.0 to 4.0 High activity: > 4.0 to 10
Simplified Disease Activity Index (SDAI)	Range: 0 to 90 Remission: ≤ 3.3 Low activity: > 3.3 to ≤ 11.0 Moderate activity: > 11.0 to ≤ 26 High activity: > 26

American College of Rheumatology 20 Percent Improvement Criteria (ACR20):

At least 20 percent improvement in the following:
1. Swollen joint count
2. Tender joint count
And three of the following five variables:
3. Patient-assessed global disease activity (e.g., by VAS)
4. Evaluator-assessed global disease activity (e.g., by VAS)
5. Patient pain assessment (e.g., by VAS)
6. Functional disability (e.g., by HAQ)
7. Acute phase response (ESR or CRP)
A 50 and 70 percent ACR response (ACR50 and ACR70, respectively) represents respective improvement of at least 50 or 70 percent ¹ .
© 2018 UpToDate, Inc.
1. Felson DT, Anderson JJ, Lange ML, et al. Should improvement in rheumatoid arthritis clinical trials be defined as fifty percent or seventy percent improvement in core set measures, rather than twenty percent?. <i>Arthritis Rheum</i> 1998; 41:1564.
2. Felson DT, Anderson JJ, Boers M, et al. American College of Rheumatology preliminary definition of improvement in rheumatoid arthritis. <i>Arthritis Rheum</i> 1995; 38:727.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Ulcerative Colitis Activity (Adults):

American College of Gastroenterology Ulcerative Colitis Activity Index				
	Remission	Mild	Moderate-severe	Fulminant
Stools (no./d)	Formed	< 4	> 6	> 10
Blood in stools	None	Intermittent	Frequent	Continuous
Urgency	None	Mild, occasional	Often	Continuous
Hemoglobin	Normal	Normal	< 75% of normal	Transfusion needed
ESR	< 30	< 30	> 30	> 30
CRP (mg/L)	Normal	Elevated	Elevated	Elevated
Fecal calprotectin (mg/g)	< 150-200	> 150-200	> 150-200	> 150-200
Endoscopy (Mayo sub-score)	0-1	1	2-3	3
UCEIS	0-1	2-4	5-8	7-8
The above factors are general guides for disease activity. With the exception of remission, a patient does not need to have all the factors to be considered in a specific category. CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; UCEIS, Ulcerative Colitis Endoscopic Index of Severity.				
Endoscopic Assessment of Disease Activity				
Endoscopic Features	UCEIS Score		Mayo Score	
Normal	0		0	
Erythema, decreased vascular pattern, mild friability	1-3		1	
Marked erythema, absent vascular pattern, friability, erosions	4-6		2	
Spontaneous bleeding, ulceration	7-8		3	

Pediatric ulcerative colitis activity index (PUCAI)

Abdominal pain	No pain	0 points
	Pain can be ignored	5 points
	Pain cannot be ignored	10 points
Rectal Bleeding	None	0 points
	Small amount only, in <50% of stools	10 points
	Small amount with most stools	20 points
	Large amount (>50% of the stool content)	30 points
Stool consistency of most stools	Formed	0 points
	Partially formed	5 points
	Completely unformed	10 points
Number of stools er 24 hours	0 to 2	0 points
	3 to 5	5 points
	6 to 8	10 points
	>8	15 points
Nocturnal stools (any episode causing wakening)	No	0 points
	Yes	10 points
Activity level	No limitation of activity	0 points
	Occasional limitation of activity	5 points
	Severe restricted activity	10 points
Sum (0-85) PUCAI scores are interpreted as follows: 0 to 9 – Remission 10 to 34 – Mild disease 35 to 64 – Moderate disease 65 to 85 – Severe disease		

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Uveitis:

Uveitis is characterized by inflammation of the uvea, which is the middle portion of the eye made up of the iris, ciliary body and choroid. The anterior portion of the uvea includes the iris and ciliary body, the posterior portion of the uvea is known as the choroid. There are several types of uveitis, defined by the part of the eye where it occurs:

- Iritis also called anterior uveitis, is the most common type of uveitis
- Intermediate uveitis or pars planitis is inflammation of the uvea in the middle or intermediate region of the eye
- Posterior uveitis affects the back parts of your eye
- Panuveitis occurs when all layers of the uvea are inflamed

Resources:

Abrilada (adalimumab-afzb) product information, revised by Pfizer Laboratories Div Pfizer, Inc. 04-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-aacf injection product information, revised by Fresenius Kabi USA, LLC. 06-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-aaty injection product information, revised by Celltrion USA, Inc 12-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-adaz product information, revised by Sandoz, Inc. 06-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-adbm injection product information, revised by Boehringer Ingelheim Pharmaceuticals, Inc. 04-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-fkjp injection product information, revised by Biocon Biologics, Inc. 12-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Adalimumab-ryvk injection product information, revised by Quallent Pharmaceuticals Health LLC. 07-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Amjevita (adalimumab-atto) injection product information, revised by Amgen, Inc. 08-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed January 16, 2025.

Amjevita (adalimumab-atto) injection product information, revised by Optum Health Solutions (Ireland) Limited. 08-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Cyltezo (adalimumab-adbm) injection product information, revised by Boehringer Ingelheim Pharmaceuticals, Inc. 04-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Hadlima (adalimumab-bwwd) injection product information, revised by Organon LLC. 06-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Hulio (adalimumab-fkjp) injection product information, revised by Biocon Biologics, Inc. 12-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Humira (adalimumab) injection product information, revised by AbbVie Inc. 02-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed January 16, 2025.

Hyrimoz (adalimumab-adaz) injection product information, revised by Cordavis Limited. 04-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Idacio (adalimumab-aacf) injection product information, revised by Fresenius Kabi USA, LLC. 12-2022. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Simlandi (adalimumab-ryvk) product info, revised by Teva Pharmaceuticals USA, Inc. 07-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Yuflyma (adalimumab-aaty) injection product information, revised by Celltrion USA, Inc. 01-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

Yusimry (adalimumab-aqvh) injection product information, revised by Meitheal Pharmaceuticals, Inc. 10-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed December 03, 2024.

van Tubergen A. Treatment of axial spondyloarthritis (ankylosing spondylitis and non-radiographic axial spondyloarthritis) in adults. In: UpToDate, Sieper J, Seo P (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated on January 22, 2024. Accessed January 09, 2025.

Reguerio M, Al Hashash J. Overview of the medical management of mild (low risk) Crohn disease in adults. In: UpToDate, Kane SV, Robson KM (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated June 12, 2024. Accessed January 09, 2025.

Al Hashash J, Reguerio M. Medical management of moderate to severe Crohn disease in adults. In: UpToDate, Kane SV, Robson KM (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated September 09, 2024. Accessed January 09, 2025.

Zitomersky N, Bousvaros A. Overview of the management of Crohn disease in children and adolescents. In: UpToDate, Heyman MB, Hoppin AG (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated August 01, 2022. Accessed January 17, 2025.

Martini A, Ravelli A, Avcin T, et al.: Towards a new classification criteria for juvenile idiopathic arthritis: First steps, Pediatric International Trials Organization International Consensus. *J Rheumatol* 2019;46(2):190-197. Re-evaluated January 17, 2025.

Weiss PF. Polyarticular juvenile idiopathic arthritis: Treatment and prognosis. In: UpToDate, Klein-Gitelman M, Case SM (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated September 25, 2024. Accessed January 09, 2025.

Weiss PF. Polyarticular juvenile idiopathic arthritis: Clinical manifestations, diagnosis, and complications. In: UpToDate, Klein-Gitelman M, Case SM (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated December 12, 2023. Accessed January 17, 2025.

Feldman SR, Soung J. Chronic plaque psoriasis in adults: Treatment of disease requiring phototherapy or systemic therapy. In: UpToDate, Dellavalle RP, Duffin KC, Ofori, AO (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated November 21, 2024. Accessed January 09, 2025.

Gladman DD, Orbai AM. Treatment of psoriatic arthritis. In UpToDate, Sieper J, Seo P (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated on September 29, 2023. Accessed January 09, 2025.

Moreland LW, Cannella A. General principles and overview of management of rheumatoid arthritis in adults. In: UpToDate, O'Dell JR, Seo P (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated May 14, 2024. Accessed January 09, 2025.

Cohen S, Mikuls TR. Initial treatment of rheumatoid arthritis in adults. In: UpToDate, O'Dell JR, Seo P (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated January 05, 2024. Accessed January 17, 2025.

Cohen S, Cannella A. Treatment of rheumatoid arthritis in adults resistant to initial conventional nonbiologic DMARD therapy. In: UpToDate, O'Dell JR, Seo P (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated June 10, 2024. Accessed January 17, 2025.

Al Hashash J, Reguerio M. Medical management of low-risk adult patients with mild to moderate ulcerative colitis. In: UpToDate, Kane SV, Robson KM (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through December 2024. Topic last updated March 22, 2023. Accessed January 09, 2025.

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINE

HUMIRA AND HUMIRA BIOSIMILARS

Cohen RD, Stein AC. Management of moderate to severe ulcerative colitis in adults. In: UpToDate, Kane SV, Robson KM (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated December 03, 2024. Accessed January 09, 2025.

Bousvaros A, Kaplan JL. Management of mild to moderate ulcerative colitis in children and adolescents. In: UpToDate, Heyman MB, Hoppin AG (Eds), UpToDate, Waltham, MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated July 16, 2024. Accessed January 17, 2025.

Ingram JR. Hidradenitis suppurativa: Management. In: UpToDate, Dellavalle RP, Owen C, Ofori OA (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated October 31, 2024. Accessed January 09, 2025.

Papaliodis GN. Uveitis: Treatment. In: UpToDate, Thorne JE, Case SM (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature review through December 2024. Topic last updated December 06, 2024. Accessed January 09, 2025.

ORIGINAL EFFECTIVE DATE: 04/01/2019 | ARCHIVE DATE: | LAST REVIEW DATE: 02/20/2025 | LAST CRITERIA REVISION DATE: 06/23/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.