

Policy and Procedure	
PHARMACY PRIOR AUTHORIZATION POLICY AND CRITERIA ORPTCEND039.0425	ENDOCRINE & METABOLIC DRUGS MEDICAL HORMONE THERAPY See Table 1 for Medications
Effective Date: 6/1/2025	Review/Revised Date: 08/15, 03/16, 07/16, 03/17, 3/18, 05/18, 07/18, 01/19, 02/19, 03/19, 02/20, 03/21, 03/22, 03/23, 05/23, 02/24, 09/24, 12/24, 02/25 (MTW)
Original Effective Date: 09/15	P&T Committee Meeting Date: 02/15, 08/15, 04/16, 08/16, 04/17, 04/18, 06/18, 07/18 (cv), 02/19, 02/19 (cv), 04/19, 04/20, 04/21, 04/22, 04/23, 06/23, 04/24, 10/24, 02/25, 04/25
Approved by: Oregon Region Pharmacy and Therapeutics Committee	

SCOPE:

Providence Health Plan and Providence Health Assurance as applicable (referred to individually as “Company” and collectively as “Companies”).

APPLIES TO:

Commercial
Medicaid

POLICY CRITERIA:

COVERED USES:

All Food and Drug Administration (FDA). Some medically accepted indications not otherwise excluded from the benefit.

Coverage for Medicaid is limited to a condition that has been designated a covered line-item number by the Oregon Health Services Commission listed on the Prioritized List of Health Care Services when all applicable indication-specific criteria below are met. The Early and Periodic Screening, Diagnostic and Treatment (EPSDT) benefit provides comprehensive and preventive health care services for children and adolescents up to their 21st birthday who are enrolled in Medicaid. Management of unfunded conditions falls under this benefit when they impact the ability to grow, develop or participate in school and the applicable indication-specific criteria below are met.

REQUIRED MEDICAL INFORMATION:

Hormone replacement therapy is considered medically necessary if all the following criteria are met:

1. One of the following diagnoses:
 - a. Gender dysphoria or gender incongruence (diagnosis codes F64.0, F64.2, F64.8, or F64.9)

OR

 - b. Diagnosis of primary or secondary (hypogonadotropic) hypogonadism
 - c. Delayed puberty (Testopel® only)

2. For testosterone replacement: Documented inability to reach therapeutic levels or experiencing fluctuations in levels (resulting in symptoms) of the following:
 - a. **For topical products:** Generic formulary topical testosterone (such as generic topical testosterone 1% or generic topical testosterone 1.62% pump)
 - b. **For injectable products:** Generic injectable testosterone cypionate
 - c. **For all other products including oral, pellet, and nasal products:** One of the following:
 - i. For Testopel® for the treatment of delayed puberty: generic injectable testosterone enanthate
OR
 - ii. Must try both of the following:
 - 1) Generic formulary topical testosterone (such as generic topical testosterone 1% or generic topical testosterone 1.62% pump)
 - 2) Generic injectable testosterone cypionate
3. For estrogen replacement: The use of a subcutaneous pellet formulation of estrogen is considered investigational for all indications.
4. For reauthorization for delayed puberty must meet both of the following:
 - a. Documentation of continued need for therapy
 - b. Documentation of monitoring for adverse effects, including an x-ray of hand and wrist to determine bone age

EXCLUSION CRITERIA:

- Use for improvement of sexual signs and symptoms (such as decreased libido, sexual dysfunction)
- Compounded hormones for pellet insertion that are not FDA approved

AGE RESTRICTIONS: N/A

PRESCRIBER RESTRICTIONS: N/A

COVERAGE DURATION:

Delayed puberty: Initial authorization and reauthorization will be approved for six months.

All other indications: Authorization will be approved until no longer eligible with the plan, subject to formulary or benefit changes

Requests for indications that were approved by the FDA within the previous six (6) months may not have been reviewed by the health plan for safety and effectiveness and inclusion on this policy document. These requests will be reviewed using the New Drug and or Indication Awaiting P&T Review; Prior Authorization Request ORPTCOPS047.

Requests for a non-FDA approved (off-label) indication requires the proposed indication be listed in either the American Hospital Formulary System (AHFS), Drugdex, or the National Comprehensive Cancer Network (NCCN) and is considered subject to evaluation of the prescriber's medical rationale, formulary alternatives, the available published evidence-based research and whether the proposed use is determined to be experimental/investigational.

Coverage for Medicaid is limited to a condition that has been designated a covered line item number by the Oregon Health Services Commission listed on the Prioritized List of Health Care Services.

Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case.

INTRODUCTION:

Each testosterone replacement therapy (TRT) product is approved for use in adult males for conditions associated with a deficiency or absence of endogenous testosterone, commonly referred to as hypogonadism. The condition is further classified into primary hypogonadism, failure of the testicles to produce testosterone, and secondary (hypogonadotropic) hypogonadism, central defects in the hypothalamus or pituitary gland, both leading to low testosterone levels. In primary hypogonadism, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) are generally at normal or elevated levels; where as in secondary hypogonadisms levels are decreased. There are various administration routes available for TRT, including transdermal patch, topical gels, buccal formulations, implantable pellets and intramuscular injections (short & long acting) of which all have been proven effective in increasing testosterone levels.

Note: All TRT products are Drug Enforcement Administration (DEA) Controlled Substance Class III

FDA APPROVED INDICATIONS:

Replacement therapy in congenital or acquired conditions associated with a deficiency or absence of endogenous testosterone, such as:

- Primary hypogonadism: testicular failure from conditions such as cryptorchidism, bilateral torsions, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter syndrome, chemotherapy, or toxic damage from alcohol or heavy metals.
- Secondary (hypogonadotropic) hypogonadism: idiopathic gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency, or pituitary-hypothalamic injury from tumors, trauma, or radiation.
- Delayed puberty (Testopel® only): Androgens may be used to stimulate puberty in carefully selected males with clearly delayed puberty. These patients usually have a familial pattern of delayed puberty that is not secondary to a pathological disorder; puberty is expected to occur spontaneously at a relatively late date. Brief treatment with conservative doses may occasionally be justified in these

patients if they do not respond to psychological support. The potential adverse effect on bone maturation should be discussed with the patient and parents prior to androgen administration. An x-ray of the hand and wrist to determine bone age should be taken every 6 months to assess the effect of treatment on epiphyseal centers

POSITION STATEMENT:

Testosterone Replacement Therapy (TRT)

- There is no evidence demonstrating any one TRT product is safer or more effective than other therapeutic alternatives. All products were proven effective based on pharmacokinetic data.
- Total serum testosterone level must be measured to confirm diagnosis of hypogonadism.
- According to the 2018 Endocrine Society Guidelines: For {total serum testosterone} laboratories that are not CDC certified and do not participate in an accuracy-based quality control program, the reference range may vary considerably depending on the assay and reference population used. Using the lower limit of the range established in local laboratories may not accurately identify men with hypogonadism.
- A harmonized reference range for free testosterone (FT) has not been established, so reference ranges may vary considerably depending on the specific equilibrium dialysis method or the algorithm used to calculate FT. Therefore, until a harmonized reference range is established, the lower limits established by the laboratory may be used.
- Testosterone exhibit diurnal variation levels, with peak levels in the morning and varying levels throughout the day. A confirmed diagnosis of hypogonadism must be measured by morning serum testosterone levels before 10 AM on at least two occasions.
- Normal ranges for testosterone vary among laboratories and assays. The Endocrine Society and Center for Disease Control recommends a lower limit of normal testosterone of 264ng/dL for total serum testosterone. The American Urological Association recommends a lower limit of normal total serum testosterone of less than 300ng/dL.
- TRT is contraindicated in men with known or suspected prostate or breast cancer.
- The FDA requires manufacturer labeling to reflect the appropriate use of TRT, specifically as replacement therapy only for men who have low testosterone levels due to disorders of the testicles, pituitary gland, or brain that cause a condition called hypogonadism.
- In 2018, both the Endocrine Society and American Urological Association concluded that the evidence is inconclusive with regards to TRT and concerns for increased cardiovascular risk in hypogonadal men.

- According to the National Institute for Health and Care Excellence (NICE), testosterone supplementation may be considered for people with low sexual desire associated with menopause if hormone replacement therapy (HRT) alone is not effective. None of the androgen therapies attached to this policy are approved by the FDA for treating sexual dysfunction due to limited clinical trial data, limited efficacy compared with placebo, and concerns about long-term safety. Drugs or medicines used to treat sexual dysfunction are a benefit exclusion for many groups.

Estrogen replacement therapy (ERT)

- Used by biological females typically for the treatment of symptoms of menopause
- Also used for gender-affirming care in biological males
- ERT is available in several different formulations, including estradiol tablets, transdermal patches, injectable, and vaginal applications.
- There are no FDA approved estrogen formulations for pellet insertion; these products are often compounded

REFERENCE/RESOURCES:

1. Bhasin, S., Cunningham, G.R., Hayes, F.J., et al. Testosterone Therapy in men with androgen deficiency syndromes: An endocrine society clinical practice guideline. *J. Clin Endocrinol and Metab.*, Jun 2010;95,2536-2559.
2. Relevant package inserts
3. Lunenfeld B, Mskhalaya G, Zitzmann M, et al. (2021). Recommendations on the diagnosis, treatment and monitoring of testosterone deficiency in men. *The Aging Male*, 24(1), 119–138.
<https://doi.org/10.1080/13685538.2021.1962840>
4. Shalender Bhasin, Juan P Brito, et al, Testosterone Therapy in Men With Hypogonadism: An Endocrine Society Clinical Practice Guideline, *The Journal of Clinical Endocrinology & Metabolism*, Volume 103, Issue 5, May 2018, Pages 1715–1744, <https://doi.org/10.1210/jc.2018-00229>
5. Morales A, Bebb RA, Manjoo P et al. Diagnosis and management of testosterone deficiency syndrome in men: clinical practice guideline. *CMAJ*. 2015;187(18):1369-1377.
6. Travison TG, Vesper HW, Orwoll E, Wu F, Kaufman JM, Wang Y, et al. Harmonized Reference Ranges for Circulating Testosterone Levels in Men of Four Cohort Studies in the United States and Europe. *J Clin Endocrinol Metab*. 2017;102(4):1161-73.
7. Mulhall JP, Trost LW, Brannigan RE et al: Evaluation and management of testosterone deficiency: AUA guideline. *J Urol* 2018; **200**: 423.
8. National Institute for Health and Care Excellence. (2024). Menopause: identification and management [NICE Guideline No. 23].

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCEND039**

**ENDOCRINE & METABOLIC DRUGS
MEDICAL HORMONE THERAPY**
See [Table 1](#) for Medications

<https://www.nice.org.uk/guidance/ng23/resources/menopause-identification-and-management-pdf-1837330217413>

Table 1: Medications covered by this policy

Brand Name	Benefit
Androderm [®] patch	Pharmacy
Androgel [®] 1.62 % packet	Pharmacy
Aveed [®] testosterone vial	Medical
Fortesta [®] pump	Pharmacy
Jatenzo [®] capsule	Pharmacy
Natesto [®] nasal pump	Pharmacy
Testopel [®] pellet	Medical
Tlando [®] capsule	Pharmacy
Xyosted [®] auto-injector	Pharmacy
Compounded estradiol and/or testosterone for use in pellet implantation	N/A - not covered as these are not FDA approved

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCEND039**

**ENDOCRINE & METABOLIC DRUGS
MEDICAL HORMONE THERAPY**
See [Table 1](#) for Medications

Appendix 1. Available TRT products

Testosterone Product	Standard Daily Dose	Max Daily Dose	How supplied (per package)	Site of Administration
<i>Androgel®</i>				
1%	5 g (4 pumps) applied topically once daily	10 g (8 pumps)	<ul style="list-style-type: none"> # 30 – 5 g tubes or packets 2 x 75 g pumps (120 metered pumps per package) 	Applied to right and left upper arms /shoulders and/or right/left abdomen
1.62%	Initial: 40.5 mg (2 pumps) applied topically once daily	81 mg (4 pumps)	<ul style="list-style-type: none"> # 30 – 5 g tubes or packets 2 x 75 g pumps (120 metered pumps per package) 	
<i>Androderm®</i> 2 mg/day & 4 mg/day	Apply topically once daily	1 patch per application	<ul style="list-style-type: none"> 2 mg: 60 patches 4 mg: 30 patches 	Applied to the back, abdomen, upper arms, or thighs
<i>Axiron®</i>	Initial: 60 mg (1 pump actuation to each axilla) applied topically once daily	120 mg (2 pumps actuations to each axilla)	<ul style="list-style-type: none"> 110 mL (60 metered pumps per package) 	Applied to the axilla only.
<i>Fortesta®</i>	40 mg (4 pumps) applied topically once daily	70 mg (7 pumps)	<ul style="list-style-type: none"> 60 g canister (120 metered pumps per package) 	Applied to front and inner thighs only.
<i>Vogelxo®</i>	50 mg (1 tube or packet or 4 pumps) once daily	100 mg (2 tubes or packets or 8 pumps)	<ul style="list-style-type: none"> # 30 – 5 g tubes or packets 2 x 75 g pumps (120 metered pumps per package) <p>* MAX 2 packages/month</p>	Applied to the shoulders and/or upper arms only.
<i>Testim®</i>	5 g (1 tube) applied topically once daily	10 g (2 tubes)	<ul style="list-style-type: none"> #30 – 5 g tubes 	Applied to the shoulders and/or upper arms only.
<i>Tlando®</i>	225 mg orally twice daily	N/A	112.5 mg capsule	Oral
<i>Jatenzo®</i>	237 mg orally twice daily	396 mg twice daily	<ul style="list-style-type: none"> Capsules are available in three strengths of 158 mg, 198 mg, and 237 	Oral

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCEND039**

**ENDOCRINE & METABOLIC DRUGS
MEDICAL HORMONE THERAPY**
See [Table 1](#) for Medications

Testosterone Product	Standard Daily Dose	Max Daily Dose	How supplied (per package)	Site of Administration
			mg. Capsules are packaged as 120 units per bottle.	
<i>Kyzatrex®</i> Note: this product is only available direct from the manufacturer ("cash" pay) and cannot be billed through insurance	Recommended starting dose of 200 mg twice daily. Minimum recommended dose is 100 mg once daily.	400 mg twice daily	100 mg, 150 mg, and 200 mg capsules	Oral
<i>Natesto™</i>	1 actuation per nostril (5.5.mg per actuation) INTRANASALLY 3 times a day 6 to 8 hours apart for total daily dose of 33 mg	33 mg/ day – no dose adjustments are recommended. If testosterone levels remain below 300 ng/dL, treatment should be discontinued	Metered dose pump, containing 60 actuations (10day supply)	Nasal
<i>Xyosted®</i>	Initial: 75mg (1 autoinjector) subcut once weekly, titrate to trough concentration between 350 ng/dL and 650 ng/dL		Autoinjector (carton containing 4 autoinjectors) available as 50mg/0.5ml; 75/0.5ml; or 100mg/0.5ml injectors	Administer subcutaneously in the abdominal region
<i>Testopel®</i>	The number of pellets to be implanted depends upon the minimal daily requirements of testosterone propionate determined by a gradual reduction of the amount administered parenterally. The usual dosage is as follows: implant two 75mg pellets for each 25mg testosterone propionate required weekly.	The suggested dosage for androgens varies depending on the age, and diagnosis of the individual patient. 150mg to 450mg subcutaneously every 3 to 6 months	Each TESTOPEL® (testosterone pellets) contains 75mg testosterone	Implanted subcutaneously

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCEND039**

**ENDOCRINE & METABOLIC DRUGS
MEDICAL HORMONE THERAPY**
See [Table 1](#) for Medications

Testosterone Product	Standard Daily Dose	Max Daily Dose	How supplied (per package)	Site of Administration
<i>Compounded testosterone (not FDA approved)</i>	Not determined – not FDA approved	Not determined – not FDA approved	Compounded for implantation	Implanted subcutaneously

Appendix 2. Available ERT products

Route of Administration	FDA-Approved Products
Oral	Estradiol tablets (Estrace®) Menest® (esterified estrogens) tablets Premarin® (conjugated estrogens) tablets
Injectable	Estradiol valerate (Delestrogen®) Estradiol cypionate
Implant insertion	Compounded estradiol for pellet insertion (not FDA approved)
Transdermal	Estradiol gel (Divigel®) Estradiol patch (Dotti®, Climara®, Vivelle-DOT®) Estradiol cream (Estrace®)
Vaginal	Estradiol (Vagifem®) vaginal tablet Estradiol acetate (Femring®) vaginal ring Premarin® (conjugated estrogens) cream with applicator
Combination Products	Premphase®/Prempro® (conjugated estrogens/medroxyprogesterone) Esterified estrogens/methyltestosterone tablets

BILLING GUIDELINES AND CODING

Per Coding Policy 22.0 HCPCS S-Codes and H-Codes, S-codes are not accepted as billable codes; testosterone pellet (Testopel®) must be billed with the unclassified code.

INJECTABLE MEDICATIONS & ADMINISTRATION:	
DRUG CODE(S)	Description
J3145	Injection, testosterone undecanoate, 1 mg
96372	Ther/proph/diag inj sc/im
HORMONE PELLET & ADMINISTRATION	
DRUG CODE(S)	Description
J3490	Unclassified drugs (Testopel®)
11980*	Subcutaneous hormone pellet implantation (implantation of estradiol and/or testosterone pellets beneath the skin)
<i>*11980 may be covered when billed with J3490 for Testopel® for which there is an approved prior authorization on file. 11980 is not covered when billed separately and without J3490 for Testopel®</i>	
EXCLUSIONS	
J7999	Compounded drug, not otherwise classified
S0189	Testosterone pellet, 75 mg