

# Retisert

## (fluocinolone acetonide intravitreal implant)

Override(s)	Approval Duration
Prior Authorization	One time

Medications	Dosing Limit
Retisert (fluocinolone acetonide) 0.59 mg intravitreal implant	One intravitreal implant (0.59 mg) per eye; each implant may be replaced following depletion of fluocinolone acetonide as evidenced by recurrence of uveitis

### **APPROVAL CRITERIA**

Requests for Retisert (fluocinolone acetonide intravitreal implant) may be approved if the following criterion is met:

- I. Individual has a diagnosis of chronic (duration of 1 year or more) non-infectious uveitis affecting the posterior segment of the eye.

Requests for Retisert (fluocinolone acetonide intravitreal implant) may **not** be approved for the following criteria:

- I. All other indications not included above; **OR**
- II. Individual has active viral diseases of cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella; **OR**
- III. Individual has active bacterial, mycobacterial or fungal infections of the eye.

### **Key References:**

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2019. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: July 8, 2019.
3. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
4. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2019; Updated periodically.