

PHARMACY COVERAGE GUIDELINE

EOHILIA™ (budesonide) oral suspension Generic Equivalent (if available)

This Pharmacy Coverage Guideline (PCG):

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

Scope

- This PCG applies to Commercial and/or Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

Instructions & Guidance

- To determine whether a member is eligible for the Service, read the entire PCG.
 - This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
 - Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
 - The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
 - The “Description” section describes the Service.
 - The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
 - The “Resources” section lists the information and materials we considered in developing this PCG
 - **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
 - Information about medications that require prior authorization is available at www.azblue.com/pharmacy. You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to Pharmacyprecert@azblue.com.
-

Medical Necessity Requirements for EOHILIA (budesonide) oral suspension

Criteria for Initial Therapy:

Prescriber Qualifications

- Prescribed by or in consultation with an allergist or gastroenterologist

Indication

- Diagnosis of eosinophilic esophagitis (EoE)

PHARMACY COVERAGE GUIDELINE

EOHILIA™ (budesonide) oral suspension Generic Equivalent (if available)

Age Requirement

- 11 years of age or older

Baseline Clinical Evaluation

- History of esophageal dysfunction symptoms (eating problems, abdominal pain, heartburn, dysphagia, vomiting, food impaction, weight loss) intermittently or continuously
- Endoscopic biopsy shows 15 or greater intraepithelial eosinophils per high power field or 60 eosinophils per mm² (must submit copy of endoscopy report)
- Screening for hepatitis B infection

Alternative Therapies

- Failure (trial for at least two months duration), contraindication, intolerance from **EACH** of the following treatment classes:
 - Proton pump inhibitor (e.g., pantoprazole, omeprazole)
 - Topical corticosteroid using swallowed (not inhaled) fluticasone

Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

Safety

- No concomitant use with CYP3A4 inhibitors (ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, cyclosporine, grapefruit juice)
- Does not have hypersensitivity to any budesonide product
- Does not have fungal infection, Strongyloides infestation, cerebral malaria, or active ocular herpes simplex
- Does not have severe hepatic impairment (Child-Pugh Class C)

Documentation Requirements

- A completed request form must be submitted, including:
 - Chart notes
 - Lab results (eosinophil count, hepatitis B screening)
 - Supporting clinical documentation

Initial Therapy Criteria Approval Duration

- 3 months (12 weeks) maximum duration of treatment OR end of plan year

Criteria for Continuation of Therapy (renewal therapy)

Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy

PHARMACY COVERAGE GUIDELINE

EOHILIA™ (budesonide) oral suspension Generic Equivalent (if available)

Prescriber Qualifications

- Continues to be seen by or in consultation with an allergist or gastroenterologist

Clinical Response

- Achieved and maintains **TWO** of the following:
 - Significant reduction in dysphagia
 - Improvement in abdominal pain, reflux or heartburn, abdominal pain or vomiting
 - Endoscopic biopsy shows less than 7 eosinophils per high power field (eos/hpf) or greater than 50 percent reduction from baseline

Adherence

- Adherence to the prescribed therapy regimen has been documented

Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

Safety

- No concomitant use with CYP3A4 inhibitors (ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, cyclosporine, grapefruit juice)
- Does not have hypersensitivity to any budesonide product
- Does not have new or worsening localized or systemic infection (bacterial, fungal, viral, protozoal, helminthic), including oropharyngeal and esophageal candidiasis
- Does not have severe hepatic impairment (Child-Pugh Class C)

Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement in eosinophilic esophagitis (eosinophilic esophagitis)
- Lab values confirming safe use (eosinophil count)

Continuation Therapy Criteria Approval Duration

- Duration will be adjusted to meet the recommended maximum duration of 12 weeks OR end of plan year

Criteria for Off-Label Use Requests:

Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. Off-Label Use of Non-Cancer Medications

ORIGINAL EFFECTIVE DATE: 05/16/2024 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

EOHILIA™ (budesonide) oral suspension Generic Equivalent (if available)

2. Off-Label Use of Cancer Medications

Description:

Eohilia (budesonide oral suspension) is a corticosteroid indicated for 12 weeks of treatment in adult and pediatric patients 11 years of age and older with eosinophilic esophagitis (EoE). Eohilia (budesonide oral suspension) has not been shown to be safe and effective for the treatment of EoE for longer than 12 weeks.

Eosinophilic esophagitis (EoE) is a chronic, immune-mediated, esophageal disease characterized by symptoms related to esophageal dysfunction and histologically by eosinophil-predominant inflammation. Diagnostic criteria include symptoms related to esophageal dysfunction, esophageal biopsy, characteristically consisting of ≥ 15 eosinophils per high power field (HPF) (or 60 eosinophils per mm), and exclusion of other causes responsible for or contributing to symptoms and eosinophilia.

Established treatments for EoE include dietary therapy, proton pump inhibitors (PPIs), topical corticosteroids, and dupilumab. While PPIs and corticosteroids are not FDA approved treatment options, both are recognized as standard treatment per American College of Gastroenterology guidelines.

Definitions:

U.S. Food and Drug Administration (FDA) MedWatch Forms for FDA Safety Reporting
[MedWatch Forms for FDA Safety Reporting | FDA](#)

Resources:

Eohilia (budesonide oral suspension) product information, revised by Takeda Pharmaceuticals America, Inc. 01-2025. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed February 19, 2025.

Gonsalves N. Eosinophilic gastrointestinal diseases. In: UpToDate, Friedman LS, Robson KM (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated May 09, 2024. Accessed March 17, 2025.

Bonis PAL, Gupta SK. Clinical manifestations and diagnosis of eosinophilic esophagitis (EoE). In: UpToDate, Talley NJ, Meyer C (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated January 15, 2025. Accessed March 17, 2025.

Rothenberg ME. Eosinophilic esophagitis (EoE): Genetics and immunopathogenesis. In: UpToDate, Sicherer SH, TePas E (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated January 10, 2023. Accessed March 17, 2025.

Bonis PAL, Sandeep SK. Treatment of eosinophilic esophagitis (EoE). In: UpToDate, Talley NJ, Meyer C (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated February 05, 2025. Accessed March 17, 2025.

ClinicalTrials.gov Bethesda (MD): National Library of Medicine (US). Identifier NCT02605837: Oral Budesonide Suspension (OBS) in Adolescent and Adult Subjects (11 to 55 Years of Age, Inclusive) With Eosinophilic Esophagitis: A Phase 3 Randomized, Double-blind, Placebo-controlled Study. Available from: <http://clinicaltrials.gov>. Last update posted June 08, 2021. Last verified May 2021. Accessed March 19, 2024. Re-evaluated March 17, 2025.

ORIGINAL EFFECTIVE DATE: 05/16/2024 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.



An Independent Licensee of the Blue Cross Blue Shield Association

PHARMACY COVERAGE GUIDELINE

EOHILIA™ (budesonide) oral suspension **Generic Equivalent (if available)**

ORIGINAL EFFECTIVE DATE: 05/16/2024 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.