

Prior Authorization Criteria
Chimeric Antigen Receptor T cell (CAR-T) Immunotherapy

All requests for Chimeric Antigen Receptor T cell (CAR-T)* Immunotherapy require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

*CAR-T Immunotherapy medications include Kymriah (tisagenlecleucel), Yescarta (axicabtagene ciloleucel), Tecartus (brexucabtagene autoleucel), Breyanzi (lisocabtagene maraleucel), Abecma (idecabtagene vicleucel) and Carvykti (ciltacabtagene autoleucel). New products with this classification will require the same documentation.

For all requests for CAR-T Immunotherapy, all of the following criteria must be met:

- Must have documentation of CD19 tumor expression (excluding Abecma and Carvykti)
- Must be prescribed by an Oncologist or Hematologist
- Must be given as a one-time, single administration treatment
- The member has received or will receive lymphodepleting chemotherapy within two weeks preceding infusion unless the member's WBC count is less than or equal to $1 \times 10^9/L$ within 1 week prior to infusion
- Documentation screening for HBV, HCV, and HIV in accordance with clinical guidelines before collection of cells for manufacturing must be performed due to risk of viral reactivation
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
- Must be age-appropriate according to FDA-approved labeling, nationally recognized compendia, or evidence-based practice guidelines
- Exclusion criteria:
 - Will not be used as first-line therapy;
 - Will not be used in combination with other chemotherapy agents;
 - Will not be given as repeat treatment in individuals who have received CAR-T treatment previously.
 - Will not be given if the member has primary central nervous system (CNS) lymphoma (excluding Abecma and Carvykti)

KYMRIAH

Coverage may be provided with a diagnosis of B-cell acute lymphoblastic leukemia (ALL) and the following criteria is met:

- Disease is considered refractory, or in second or later relapse, in *any* of the following scenarios:
 - Second or later bone marrow relapse;
 - Bone marrow relapse after allogeneic stem cell transplant;
 - Primary refractory or chemo-refractory after relapse;
 - Presence of > 5% blasts at screening
- For members with Ph+ ALL only:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - At least two tyrosine kinase inhibitors (TKIs)
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory (r/r) large B-cell lymphoma and the following criteria is met:

- The member is diagnosed with any of the following large B-cell lymphomas:
 - Diffuse large B-cell lymphoma (DLBCL) not otherwise specified,
 - High grade B-cell lymphoma
 - DLBCL arising from follicular lymphoma.
- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory (r/r) follicular lymphoma (FL) and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

YESCARTA

Coverage may be provided with a diagnosis of relapsed or refractory large B-cell lymphoma and the following criteria is met:

- Members with large B-cell lymphoma that is refractory to first-line chemoimmunotherapy or that relapses within 12 months of first-line chemoimmunotherapy **OR**
- The member is diagnosed with any of the following large B-cell lymphomas:
 - Diffuse large B-cell lymphoma (DLBCL) not otherwise specified,
 - Primary mediastinal large B-cell lymphoma,
 - High grade B-cell lymphoma,
 - DLBCL arising from follicular lymphoma **AND**
 - Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory follicular lymphoma (FL) and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

TECARTUS

Coverage may be provided with a diagnosis of relapsed or refractory mantle cell lymphoma and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - A covalent Bruton tyrosine kinase inhibitor (BTKi; ibrutinib, acalabrutinib, or zanubrutinib) during the following scenarios:
 - no response or progressive disease following second-line therapy with covalent BTKi or other continuous treatment regimens i.e. lenalidomide and rituximab
 - partial response, no response, or progressive disease following second-line therapy with fixed-duration regimens
 - relapsed or progressive disease (relapse #2 or greater) if not previously given
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL) and the following criteria is met:

- Member has Philadelphia chromosome-negative disease that is relapsed or refractory as defined as one of the following:
 - Primary refractory disease
 - First relapse with remission of 12 months or less
 - Relapsed or refractory disease after at least 2 previous lines of systemic therapy
 - Relapsed or refractory disease after allogeneic stem cell transplant (allo-SCT)

For members with Ph+ ALL only:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - At least two tyrosine kinase inhibitors (TKIs)
- **Duration of Approval:** 1 treatment

BREYANZI

Coverage may be provided with a diagnosis of relapsed or refractory large B-cell lymphoma and the following criteria is met:

- The member is diagnosed with any of the following large B-cell lymphomas:

- Diffuse large B-cell lymphoma (DLBCL) not otherwise specified (including DLBCL arising from indolent lymphoma),
- Primary mediastinal large B-cell lymphoma,
- High grade B-cell lymphoma,
- Follicular lymphoma grade 3B
- Documentation the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to **one** of the following:
 - refractory disease to first-line chemoimmunotherapy or relapse within 12 months of first-line chemoimmunotherapy
 - refractory disease to first-line chemoimmunotherapy or relapse after first-line chemoimmunotherapy and are not eligible for hematopoietic stem cell transplantation (HSCT) due to comorbidities or age
 - two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to at least 2 prior lines of therapy including:
 - a Bruton tyrosine kinase (BTK) inhibitor and
 - a B-cell lymphoma 2 (BCL-2) inhibitor
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory follicular lymphoma (FL) and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more lines of systemic therapy
- **Duration of Approval:** 1 treatment

Coverage may be provided with a diagnosis of relapsed or refractory mantle cell lymphoma (MCL) and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to at least 2 prior lines of systemic therapy including:
 - a Bruton tyrosine kinase (BTK) inhibitor
- **Duration of Approval:** 1 treatment

ABECMA

Coverage may be provided with a diagnosis of relapsed or refractory multiple myeloma and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - Two or more prior lines of therapy including:
 - an immunomodulatory agent
 - a proteasome inhibitor
 - an anti-CD38 monoclonal antibody
- **Duration of Approval:** 1 treatment

CARVYKTI

Coverage may be provided with a diagnosis of relapsed or refractory multiple myeloma and the following criteria is met:

- Must provide documentation showing the member has tried and failed (which will be verified via pharmacy claims or submitted documentation) or had an intolerance or contraindication to the following:
 - At least 1 prior line of therapy including:
 - an immunomodulatory agent
 - a proteasome inhibitor
 - refractory to lenalidomide
- **Duration of Approval:** 1 treatment

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.



Updated: 06/2025
PARP Approved: 07/2025

**CHIMERIC ANTIGEN RECEPTOR T-CELL (CAR-T) IMMUNOTHERAPY
PRIOR AUTHORIZATION FORM**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049
If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

PROVIDER INFORMATION

Requesting Provider:	Provider NPI:
Provider Specialty:	Office Contact:
State license #:	Office NPI:
Office Address:	Office Phone:
	Office Fax:

MEMBER INFORMATION

Member Name:	DOB:	
Member ID:	Member weight:	Height:

REQUESTED DRUG INFORMATION

Medication:	Strength:	
Directions:	Quantity:	Refills:
Is the member currently receiving requested medication? <input type="checkbox"/> Yes <input type="checkbox"/> No		Date Medication Initiated:

Billing Information

This medication will be billed: <input type="checkbox"/> at a pharmacy OR <input type="checkbox"/> medically, JCODE: _____
Place of Service: <input type="checkbox"/> Hospital <input type="checkbox"/> Provider's office <input type="checkbox"/> Member's home <input type="checkbox"/> Other

Place of Service Information

Name:	NPI:
Address:	Phone:

MEDICAL HISTORY (Complete for ALL requests)

Diagnosis:	ICD Code:
------------	-----------

For all CAR-T Therapy:

Does the member have CD19 tumor expression documentation (excluding Abecma and Carvykti)? Yes No

Has the member received or will receive lymphodepleting chemotherapy within two weeks preceding the infusion?

Yes No

Is there documentation of screening for HBV, HCV, and HIV in accordance with clinical guidelines before collection of cells for manufacturing must be performed due to risk of viral reactivation? Yes No

Does the member have any of the following exclusions? Please mark if any apply. If NONE, leave blank.

Medication will be used as first-line therapy

Medication will be used in combination with other chemotherapy agents

Medication will be given as repeat treatment in individuals who have received CAR-T treatment previously

Medication will be given if the member has primary central nervous system (CNS) lymphoma (excluding Abecma & Carvykti)

Kymriah only:

Does the member have a diagnosis of B-cell acute lymphoblastic leukemia (ALL)? Yes No

Is the disease considered refractory, or in second or later relapse, in *any* of the following scenarios? Please mark which applies:

Second or later bone marrow relapse

Bone marrow relapse after allogeneic stem cell transplant

Primary refractory or chemo-refractory after relapse

Presence of > 5% blasts at screening

For members with Ph+ ALL only: Has the member tried and failed or had an intolerance or contraindication to at least two (2) tyrosine kinase inhibitors (TKIs)? Yes No

**CHIMERIC ANTIGEN RECEPTOR T-CELL (CAR-T) IMMUNOTHERAPY
PRIOR AUTHORIZATION FORM (CONTINUED)– PAGE 2 of 3**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

MEMBER INFORMATION

Member Name:	DOB:
Member ID:	Member weight: Height:

MEDICAL HISTORY (Complete for ALL requests)

Kymriah only:

Has the member been diagnosed with relapsed or refractory (r/r) large B-cell lymphoma? Yes No

Has the member been diagnosed with one of the following large B-cell lymphomas? Please mark which applies.

- Diffuse large B-cell lymphoma (DLBCL) not otherwise specified
- High grade B-cell lymphoma
- DLBCL arising from follicular lymphoma

If the member is Ph+ ALL: has the member tried and failed or had an intolerance or contraindication to two (2) or more lines of systemic therapy?

Yes No

Does the member have a diagnosis of relapsed or refractory follicular lymphoma (FL)? Yes No

Has the member tried and failed or had an intolerance or contraindication to two (2) or more lines of systemic therapy?

Yes No

Yescarta only:

Does the member have a diagnosis of relapsed or refractory large B-cell lymphoma? Yes No

Does the member have large B-cell lymphoma that is refractory to first line chemoimmunotherapy or that relapses within 12 months of first line chemoimmunotherapy? Yes No

Has the member been diagnosed with one of the following large B-cell lymphomas? Please mark which applies:

- Diffuse large B-cell lymphoma (DLBCL) not otherwise specified
- Primary mediastinal large B-cell lymphoma
- High grade B-cell lymphoma
- DLBCL arising from follicular lymphoma

Has the member tried and failed or had an intolerance or contraindication to two (2) or more lines of systemic therapy?

Yes No

Does the member have a diagnosis of relapsed or refractory follicular lymphoma (FL)? Yes No

Has the member tried and failed or had an intolerance or contraindication to two (2) or more lines of systemic therapy?

Yes No

Tecartus only:

Does the member have a diagnosis of relapsed or refractory mantle cell lymphoma? Yes No

Has the member tried and failed or had an intolerance or contraindication to all of the following: anthracycline or bendamustine containing chemotherapy, an anti-CD20 antibody and a Bruton tyrosine kinase inhibitor (BTKi; ibrutinib or acalabrutinib)?

Yes No

Does the member have a diagnosis of relapsed or refractory B-cell precursor acute lymphoblastic leukemia (ALL)? Yes No

Does the member have Philadelphia chromosome-negative disease that is relapsed or refractory as defined as one of the following:

- Primary refractory disease
- First relapse with remission of 12 months or less
- Relapsed or refractory disease after at least 2 previous lines of systemic therapy
- Relapsed or refractory disease after allogeneic stem cell transplant (allo-SCT)

If the member is Ph+ ALL: Has the member tried and failed or had an intolerance or contraindication to two (2) or more lines of systemic therapy? Yes No

**CHIMERIC ANTIGEN RECEPTOR T-CELL (CAR-T) IMMUNOTHERAPY
PRIOR AUTHORIZATION FORM (CONTINUED)– PAGE 3 of 3**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Highmark Wholecare Pharmacy Services. **FAX:** (888) 245-2049

If needed, you may call to speak to a Pharmacy Services Representative. **PHONE:** (800) 392-1147 Mon – Fri 8:30am to 5:00pm

MEMBER INFORMATION

Member Name:	DOB:
Member ID:	Member weight: Height:

MEDICAL HISTORY (Complete for ALL requests)

Brevanzi only:

Does the member have a diagnosis of relapsed or refractory large B-cell lymphoma? Yes No
 Has the member been diagnosed with any of the following large B-cell lymphomas? Please mark which applies:
 Diffuse large B-cell lymphoma (DLBCL) not otherwise specified (including DLBCL arising from indolent lymphoma)
 Primary mediastinal large B-cell lymphoma
 High grade B-cell lymphoma
 Follicular lymphoma grade 3B
 Has the member tried and failed or had an intolerance or contraindication to **one** of the following? Please mark which applies:
 refractory disease to first-line chemoimmunotherapy or relapse within 12 months of first-line chemoimmunotherapy
 refractory disease to first-line chemoimmunotherapy or relapse after first-line chemoimmunotherapy and are not eligible for hematopoietic stem cell transplantation (HSCT) due to comorbidities or age
 two or more lines of systemic therapy

Abecma and Carvykti only:

Does the member have a diagnosis of relapsed or refractory multiple myeloma? Yes No
 Has the member tried and failed or had an intolerance or contraindication to four or more prior lines of therapy including an immunomodulatory agent, a proteasome inhibitor and an anti-CD38 monoclonal antibody?
 Yes No

CURRENT or PREVIOUS THERAPY

Medication Name	Strength/ Frequency	Dates of Therapy	Status (Discontinued & Why/Current)

SUPPORTING INFORMATION or CLINICAL RATIONALE

Prescribing Provider Signature

Date

--	--