

Aranesp (darbopoetin alfa)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	Dialysis-dependent use: 1 year All other: 6 months
Medications	Quantity Limit
Aranesp (darbopoetin alfa)	May be subject to quantity limit

APPROVAL CRITERIA

Initial requests for Aranesp (darbopoetin alfa) may be approved if the following criteria are met:

- I. Individual has a baseline hemoglobin (Hgb) level less than 10 g/dL; **AND**
- II. Baseline iron status is adequate as defined by one of the following:
 - A. Transferrin saturation is at least 20%; **OR**
 - B. Ferritin is at least 80ng/mL; **OR**
 - C. Bone marrow demonstrates adequate iron stores;

AND

- III. Individual is using for one of the following:
 - A. Anemia associated with chronic kidney disease (CKD), for individuals on dialysis, to achieve and maintain Hgb levels within the range of 10 to 11 g/dL; **OR**
 - B. Anemia associated with CKD for individuals not on dialysis, to achieve and maintain Hgb levels of 10 g/dL; **OR**
 - C. Anemia associated with myelosuppressive chemotherapy when the following are met:
 1. The individual has a diagnosis of non-myeloid cancer and the anticipated outcome is not cure; **OR**
 2. Individual refuses blood transfusion (NCCN 2A);
- OR**
- D. Myelodysplastic syndrome with endogenous erythropoietin level less than or equal to 500 mU/mL (NCCN 2A); **OR**
 - E. Myelofibrosis-associated anemia with an endogenous erythropoietin level less than 500 mU/mL (NCCN 2A).

Continuation requests for Aranesp (darbopoetin alfa) may be approved if the following criteria are met:

- I. Individual demonstrates continued need for ESA treatment and has confirmation of response to treatment as evidenced by an increase in hemoglobin levels from baseline (i.e., increase of approximately 1 g/dL or greater from baseline); **AND**

- II. Individual is using the lowest ESA dose necessary to avoid transfusions; **AND**
- III. Individual meets one of the following criteria:
 - A. Hemoglobin level is not greater than 11 g/dL for CKD individuals on dialysis, or greater than 10 g/dL for CKD non-dialysis, unless otherwise specified (for example, pediatric individuals with CKD where target Hgb levels is within the range of 10 to 12 g/dL); **OR**
 - B. Hemoglobin level is not greater than 11 g/dL for individuals using for myelosuppressive chemotherapy related anemia or myelodysplastic syndrome ; **AND**
- IV. If using for myelosuppressive chemotherapy-related anemia, individual is not using beyond 6 weeks after chemotherapy has completed.

Aranesp (darbepoetin alfa) may **not** be approved for all of the following:

- I. Continued use when the hemoglobin level exceeds 11 g/dL unless otherwise specified (for example, pediatric individuals with CKD where target Hgb levels within the range of 10 to 12 g/dL); **OR**
- II. Use beyond 12 weeks of continuous treatment at therapeutic doses in the absence of response in individuals with chronic kidney disease; **OR**
- III. Use beyond 8 weeks of continuous treatment at therapeutic doses in the absence of response in individuals with myelodysplastic syndrome (MDS); **OR**
- IV. Use beyond 8 weeks of continuous treatment at therapeutic doses in the absence of response or if transfusions are still required in individuals with metastatic, non-myeloid cancer being treated with myelosuppressive chemotherapy agents known to produce anemia; **OR**
- V. As treatment in the presence of a sudden loss of response with severe anemia and low reticulocyte count; **OR**
- VI. Continued use beyond 6 weeks after therapy with myelosuppressive chemotherapy known to produce anemia is completed.

Note:

ESAs are contraindicated in individuals with uncontrolled hypertension. Blood pressure should be adequately controlled prior to initiation and during treatment with ESAs.

Key References:

1. Bohlius J, Bohlke K, Castelli R, et al. Management of cancer-associated anemia with erythropoiesis-stimulating agents: ASCO/ASH clinical practice guideline update. *J Clin Oncol.* 2019;37(15):1336-1351.
2. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2025. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
3. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Updated periodically.
4. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
5. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Inter.* 2012; Suppl 2: 279–335. Available from: https://www.kidney.org/professionals/guidelines/guidelines_commentaries/anemia. Accessed on July 14, 2023.

6. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2024; Updated periodically.
7. NCCN Clinical Practice Guidelines in Oncology™. © 2024 National Comprehensive Cancer Network, Inc. For additional information visit the NCCN website: <http://www.nccn.org/index.asp>. Accessed on June 17, 2025.
 - a. Hematopoietic Growth Factors. Version 1.2025. Revised October 11, 2024.
 - b. Myelodysplastic Syndromes. Version 2.2025. Revised January 17, 2025.
 - c. Myeloproliferative Neoplasms. Version 1.2025. Revised February 21, 2025.

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.