

PHARMACY COVERAGE GUIDELINE

IBRANCE® (palbociclib) Generic Equivalent (if available)

This Pharmacy Coverage Guideline (PCG):

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

Scope

- This PCG applies to Commercial and Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

Instructions & Guidance

- To determine whether a member is eligible for the Service, read the entire PCG.
- This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
- Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
- The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
- The “Description” section describes the Service.
- The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
- The “Resources” section lists the information and materials we considered in developing this PCG
- **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
- Information about medications that require prior authorization is available at www.azblue.com/pharmacy. You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to Pharmacyprecert@azblue.com.

Criteria:

- **Criteria for initial therapy:** Ibrance (palbociclib) and/or generic equivalent (if available) is considered ***medically necessary*** and will be approved when **ALL** the following criteria are met:
1. Prescriber is a physician specializing in the patient’s diagnosis or is in consultation with an Oncologist
 2. Individual is 18 years of age or older
 3. Individual has a confirmed diagnosis of **ONE** of the following:
 - a. Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer in combination with **one** of the following:
 - i. An aromatase inhibitor as initial endocrine-based therapy ([see Definitions section](#))
 - ii. Fulvestrant in individuals with disease progression following endocrine-based therapy

ORIGINAL EFFECTIVE DATE: 03/17/2016 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/16/2024

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

IBRANCE® (palbociclib) Generic Equivalent (if available)

- b. Other request for a specific oncologic direct treatment use that is found and listed in the National Comprehensive Cancer Network (NCCN) Guidelines with Categories of Evidence and Consensus of 1 and 2A
4. Individual has a baseline Eastern Cooperative Oncology Group (ECOG) Performance Status of 0-1
5. **If available:** Individual has failure after adequate trial, contraindication per FDA label, intolerance, or is not a candidate for a **generic equivalent** [Note: Failure, contraindication or intolerance to the generic should be reported to the FDA] ([see Definitions section](#))
6. Individual is not currently taking any other drugs which cause severe adverse reactions or any drug interactions requiring discontinuation such as use with strong CYP3A inducers such (e.g., rifampin, rifabutin, phenobarbital, carbamazepine, others)

Initial approval duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Ibrance (palbociclib) and/or generic equivalent (if available) is considered **medically necessary** and will be approved when **ALL** the following criteria are met (**samples are not considered for continuation of therapy**):

1. Individual continues to be seen by a physician specializing in the patient's diagnosis or is in consultation with an Oncologist
2. Individual's condition has responded while on therapy with response defined as there is no evidence of disease progression or unacceptable toxicity
3. Individual has been adherent with the medication
4. **If available:** Individual has failure after adequate trial, contraindication per FDA label, intolerance, or is not a candidate for a **generic equivalent** [Note: Failure, contraindication or intolerance to the generic should be reported to the FDA] ([see Definitions section](#))
5. Individual has not developed any significant adverse drug effects that may exclude continued use such as:
 - a. Severe or life-threatening neutropenia or febrile neutropenia
 - b. Severe or life-threatening interstitial lung disease or pneumonitis
6. Individual is not currently taking any other drugs which cause severe adverse reactions or any drug interactions requiring discontinuation such as use with strong CYP3A inducers such (e.g., rifampin, rifabutin, phenobarbital, carbamazepine, others)

Renewal duration: 6 months

- Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

ORIGINAL EFFECTIVE DATE: 03/17/2016 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/16/2024

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

IBRANCE® (palbociclib) Generic Equivalent (if available)

1. Off-Label Use of Non-Cancer Medications
2. Off-Label Use of Cancer Medications

Description:

Ibrance (palbociclib) is a kinase inhibitor indicated for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer in combination with letrozole as initial endocrine based therapy in postmenopausal women, or fulvestrant in women with disease progression following endocrine therapy.

Palbociclib is the first oral cyclin-dependent kinase (CDK) inhibitor that works by blocking the action of enzymes called kinases. Kinases are involved in numerous cellular functions, including cell signaling, growth, and division. The majority of breast cancers are hormone receptor positive. They are stimulated to grow by the circulating female hormones estrogen and/or progesterone. Treatment of hormone receptor-positive breast cancer often involves hormonal therapies that suppress or block the action of estrogen. Growth of hormone-receptor-positive breast cancer is also dependent on the cyclin-dependent kinases 4 and 6 (CDK4 and CDK6), which promote progression through the various phases of the cell cycle that result in cell division. Specifically, palbociclib inhibits CDK4 and CDK6.

Letrozole is a nonsteroidal competitive inhibitor of the aromatase enzyme system; it inhibits the conversion of androgens to estrogens. Fulvestrant is an estrogen receptor antagonist that binds to the estrogen receptor (ER) and downregulates the ER protein in human breast cancer cells, blocking the actions of estrogen.

Definitions:

U.S. Food and Drug Administration (FDA) MedWatch Forms for FDA Safety Reporting
[MedWatch Forms for FDA Safety Reporting | FDA](#)

Aromatase Inhibitors:

Arimidex (anastrozole)
Femara (letrozole)
Aromasin (exemestane)

Antiestrogens:

Faslodex (fulvestrant)
Tamoxifen
Fareston (toremifene)

Gonadotropin-Releasing Hormone Analog – for men with breast cancer along with aromatase inhibitors:

Zoladex (goserelin)
Vantas (histrelin)
Eligard, Lupron (leuprolide)
Trelstar (triptorelin)
Progestin Combination

ORIGINAL EFFECTIVE DATE: 03/17/2016 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/16/2024

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

IBRANCE® (palbociclib) Generic Equivalent (if available)

Antiandrogens:

Zytiga, Yonsa (abiraterone)
Erleada (apalutamide)
Casodex (bicalutamide)
Xtandi (enzalutamide)
Flutamide
Nilandron (nilutamide)

ECOG Performance status:

Eastern Co-operative Oncology Group (ECOG) Performance Status	
Grade	ECOG description
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g., light housework, office work
2	Ambulatory and capable of all selfcare but unable to carry out any work activities. Up and about more than 50% of waking hours
3	Capable of only limited selfcare, confined to bed or chair more than 50% of waking hours
4	Completely disabled. Cannot carry on any self-care. Totally confined to bed or chair
5	Dead

Oken, M.M., Creech, R.H., Tormey, D.C., Horton, J., Davis, T.E., McFadden, E.T., Carbone, P.P.: Toxicity And Response Criteria Of The Eastern Cooperative Oncology Group. Am J Clin Oncol 5:649-655, 1982

NCCN recommendation definitions:

Category 1:

Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A:

Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B:

Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3:

Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate

Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0:

Grade 1	Mild; asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated
Grade 2	Moderate; minimal, local or noninvasive intervention indicated; limiting age-appropriate instrumental ADL*
Grade 3	Severe or medically significant but not immediately life-threatening; hospitalization or prolongation of hospitalization indicated; disabling; limiting self-care ADL**
Grade 4	Life-threatening consequences; urgent intervention indicated
Grade 5	Death related to AE

U.S. department of Health and Human Services, National Institutes of Health, and National Cancer Institute

ORIGINAL EFFECTIVE DATE: 03/17/2016 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/16/2024

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

IBRANCE® (palbociclib) Generic Equivalent (if available)

Resources:

Ibrance (palbociclib) capsule product information, revised by Pfizer Laboratories Div Pfizer Inc. 12-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed February 21, 2025.

Ibrance (palbociclib) tablet product information, revised by Pfizer Laboratories Div Pfizer Inc. 12-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed February 21, 2025.

Ma CX, Sparano JA. Treatment for hormone-receptor positive, HER2-negative advanced breast cancer. In: UpToDate, Burnstein HJ, Vora SR (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through March 2025. Topic last updated March 20, 2025. Accessed April 17, 2025.

Gradishar WJ, Ruddy KJ. Breast cancer in men. In: UpToDate, Chagpar AB, Isaacs C, Vora SR (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through March 2024. Topic last updated March 03, 2025. Accessed April 17, 2025.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Breast Cancer Version 4.2025 – Updated April 17, 2025. Available at <https://www.nccn.org>. Accessed April 17, 2025.

Off Label Use of Cancer Medications: A.R.S. §§ 20-826(R) & (S). Subscription contracts; definitions.

Off Label Use of Cancer Medications: A.R.S. §§ 20-1057(V) & (W). Evidence of coverage by health care service organizations; renewability; definitions.