

Prior Authorization Criteria

Pulmonary Arterial Hypertension (PAH), Injectable Agents

All requests for injectable Pulmonary Arterial Hypertension (PAH) agents require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Pulmonary Arterial Hypertension (PAH) injectable agents Prior Authorization Criteria:

For all requests for injectable Pulmonary Arterial Hypertension (PAH) agents all of the following criteria must be met:

- Treatment is prescribed by, or in consultation with, a cardiologist or pulmonologist
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines. If a requested dose is above these recommendations, medical rationale must be submitted. For infused products, must provide member's weight, dose, frequency and titration schedule.
- Request meets diagnostic and drug criteria outlined in sections A and B
- If member is new to the plan and requests a continuation of therapy, must provide chart documentation indicating member is currently on requested therapy

A. Diagnostic Criteria

Coverage may be provided with a diagnosis of Pulmonary Arterial Hypertension (PAH) WHO Group I and the following criteria is met:

- Member has a diagnosis of PAH WHO Group I (refer below to Appendix I) confirmed by chart documentation of right-heart catheterization (RHC) or echocardiography if the provider indicates RHC is not recommended. RHC documentation must contain the following hemodynamic values:
 - Mean pulmonary arterial pressure \geq to 20 mmHg
 - Pulmonary capillary wedge pressure **OR** left atrial pressure **OR** left ventricular end-diastolic pressure \leq to 15 mmHg
 - Pulmonary vascular resistance greater than 3 Wood units.
- Documentation of member's vasoreactivity test and one of the following, unless member has a contraindication to vasoreactivity testing (e.g. low systemic blood pressure, low cardiac index, or the presence of severe (functional class IV) symptoms):
 - Member had a positive response (pulmonary artery pressure decreases at least 10 mmHg and to a value less than 40 mmHg, with an increased or unchanged cardiac output, and a minimally reduced or unchanged systemic blood pressure) and had inadequate response, contraindication or intolerance to calcium channel blocker therapy with diltiazem or a dihydropyridine
 - Member did not have a positive response to the vasoreactivity test
- Member has functional class II, III or IV symptoms (refer below to Appendix II)

- One of the following:
 - The requested drug will be used as monotherapy
 - The requested drug will be used for add-on therapy to existing monotherapy or dual therapy in addition to **both** of the following:
 - The medications must be from different therapeutic classes
 - The member must have unresponsive or progressive disease despite established PAH-specific therapies

B. Drug Criteria

- **Prostanoids/prostacyclin therapies**
 - Infused agents: epoprostenol, Flolan, Veletri, Remodulin
 - For Flolan, Veletri, or Remodulin requests in members with functional class III or IV symptoms, must provide documentation of inadequate response, contraindication or intolerance to generic epoprostenol or clinical rationale for why generic epoprostenol cannot be used
 - Must meet **one** of the following criteria:
 - Documentation of WHO functional class IV symptoms or functional class III symptoms with **any** of the following:
 - Evidence of progression of their disease
 - Any marker of poor clinical prognosis defined as:
 - Clinical signs of right heart failure
 - Repeated episodes of syncope, even with little or regular physical activity
 - <165 meter 6-minute walking distance (6MWD)
 - Peak oxygen consumption (VO₂) <11ml/min/kg (<35% predicted)
 - Ventilatory equivalents of CO₂ (VE/VCO₂) slope ≥45
 - BNP >300 ng/l
 - NT-proBNP >1400ng/l
 - Right atrium area ≥26 cm²
 - Presence of pericardial effusion
 - Right atrial pressure >14 mmHg
 - Cardiac index <2.0 l/min/m²
 - Mixed venous oxygen saturation (SvO₂) <60%
 - Documentation of WHO functional class II symptoms and all of the following:
 - Request is for Remodulin
 - Inadequate response, contraindication or intolerance to combination therapy of generic ambrisentan and tadalafil
 - Inadequate response, contraindication or intolerance to one additional PAH medication therapy (could have been in combination or monotherapy)

Initial Duration of Approval: 3 months

Reauthorization criteria:

- Must provide documentation that demonstrates member is tolerating and receiving clinical benefit from treatment.

Reauthorization Duration of Approval: 12 months

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.

Appendix I: World Health Organization (WHO) Clinical Classification of Pulmonary Hypertension (PH)

Group 1	Pulmonary Arterial Hypertension (PAH) <ul style="list-style-type: none"> • Idiopathic • Heritable • Drug/toxin-induced • Associated with: <ul style="list-style-type: none"> ○ Connective tissue disease ○ HIV infection ○ Portal hypertension ○ Congenital heart disease ○ Schistosomiasis
Group 1'	Pulmonary veno-occlusive disease (PVOD) and/or pulmonary capillary hemangiomatosis (PCH)
Group 1''	Persistent pulmonary hypertension of the newborn (PPHN)
Group 2	PH due to left heart disease
Group 3	PH due to lung diseases and/or hypoxia
Group 4	Chronic thromboembolic PH (CTEPH) and other pulmonary artery obstructions
Group 5	PH with unclear and/or multifactorial mechanisms

Appendix II: World Health Organization (WHO) Functional Classification of Members with PH

Classification	Physical Activity	Symptoms (Dyspnea, fatigue, chest pain, syncope)
Class I	No limitation	None upon ordinary physical activity
Class II	Slight limitation	Symptoms appear upon ordinary physical activity
Class III	Marked limitation	Symptoms appear upon less than ordinary activity
Class IV	Severe limitation	Symptoms appear upon any physical activity or may even be present at rest; signs of right heart failure present



It's
Wholecare.

Updated: 04/2020
PARP Approved: 04/2020

**Pulmonary Arterial Hypertension (PAH) Agents (Oral and Inhaled)
PRIOR AUTHORIZATION FORM (CONTINUED) – PAGE 2 OF 2**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Gateway HealthSM Pharmacy Services. **FAX:** (888) 245-2049
If needed, you may call to speak to a Pharmacy Services Representative.
PHONE: (800) 392-1147 Monday through Friday 8:30am to 5:00pm

MEMBER INFORMATION

Member Name:	DOB:
Gateway ID:	Member weight: Height:

MEDICAL HISTORY (Complete for ALL requests)

Please check any boxes applicable to the member:

- | | |
|--|--|
| <input type="checkbox"/> Clinical signs of right heart failure | <input type="checkbox"/> Repeated episodes of syncope, even with little or regular physical activity |
| <input type="checkbox"/> Presence of pericardial effusion | <input type="checkbox"/> Peak oxygen consumption (VO ₂) <11ml/min/kg (<35% predicted) |
| <input type="checkbox"/> Ventilatory equivalents of CO ₂ (VE/VCO ₂) slope ≥45 | <input type="checkbox"/> <165 meter 6-minute walking distance (6MWD) |
| <input type="checkbox"/> Mixed venous oxygen saturation (SvO ₂) <60% | <input type="checkbox"/> Cardiac index <2.0 l/min/m ² |
| <input type="checkbox"/> BNP >300 ng/l <input type="checkbox"/> NT-proBNP >1400ng/l | <input type="checkbox"/> Right atrium area ≥26 cm ² <input type="checkbox"/> Right atrial pressure >14 mmHg |

PREVIOUS THERAPY

Medication Name	Strength/ Frequency	Dates of Therapy	Status (Discontinued & Why/Current)

REAUTHORIZATION

Has the member experienced a significant improvement with treatment? Yes No
Please describe:

SUPPORTING INFORMATION or CLINICAL RATIONALE

Prescribing Provider Signature

Date

--	--