

PHARMACY COVERAGE GUIDELINE

VALCHLOR™ (mechlorethamine) Generic Equivalent (if available)

This Pharmacy Coverage Guideline (PCG):

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

Scope

- This PCG applies to Commercial and Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

Instructions & Guidance

- To determine whether a member is eligible for the Service, read the entire PCG.
- This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
- Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
- The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
- The “Description” section describes the Service.
- The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
- The “Resources” section lists the information and materials we considered in developing this PCG
- **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
- Information about medications that require prior authorization is available at www.azblue.com/pharmacy. You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to Pharmacyprecert@azblue.com.

Criteria:

- **Criteria for initial therapy:** Valchlor (mechlorethamine) gel and/or generic equivalent (if available) is considered **medically necessary** when **ALL** of the following criteria are met:
1. Prescriber is a physician specializing in the patient’s diagnosis or is in consultation with an Oncologist or Dermatologist
 2. Individuals 18 years of age or older
 3. Individual has a confirmed diagnosis of **ONE** of the following:
 - a. Stage IA and IB mycosis fungoides-type cutaneous T-cell Lymphoma who have received prior skin-directed therapy

ORIGINAL EFFECTIVE DATE: 02/28/2014 | ARCHIVE DATE: | LAST REVIEW DATE: 11/21/2024 | LAST CRITERIA REVISION DATE: 11/16/2023

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

VALCHLOR™ (mechlorethamine) Generic Equivalent (if available)

- b. Other request for a specific oncologic direct treatment use that is found and listed in the National Comprehensive Cancer Network (NCCN) Guidelines with Categories of Evidence and Consensus of 1 or 2A
4. **If available:** Individual has failure after adequate trial, contraindication per FDA label, intolerance, or is not a candidate for a **generic equivalent** [Note: Failure, contraindication or intolerance to the generic should be reported to the FDA] ([see Definitions section](#))
5. Individual has documented failure, contraindication per FDA label, intolerance or is not a candidate for at least **TWO** of the following prior skin-directed therapy:
 - a. Topical corticosteroid
 - b. Topical retinoid
 - c. Topical imiquimod
 - d. Local superficial radiation
 - e. Phototherapy (such as PUVA for thicker plaques, UVB, and NB-UVB for patch/thin plaques)
 - f. Total skin electron beam radiation (TSEBT)

Initial approval duration: 6 months

- **Criteria for continuation of coverage (renewal request):** Valchlor (mechlorethamine) gel and/or generic equivalent (if available) is considered **medically necessary** and will be approved when **ALL** of the following criteria are met (**samples are not considered for continuation of therapy**):
1. Individual continues to be seen by a physician specializing in the patient's diagnosis or is in consultation with an Oncologist or Dermatologist
 2. Individual's condition has responded while on therapy with response defined as **ONE** of the following:
 - a. No evidence of disease progression defined as worsening of index lesion(s) or development of new cutaneous tumor lesions or development of non-cutaneous manifestations of disease
 - b. Documented evidence of efficacy, disease stability and/or improvement defined as at least a 50% improvement or complete disappearance of the index lesion(s)
 3. Individual has been adherent with the medication
 4. **If available:** Individual has failure after adequate trial, contraindication per FDA label, intolerance, or is not a candidate for a **generic equivalent** [Note: Failure, contraindication or intolerance to the generic should be reported to the FDA] ([see Definitions section](#))
 5. Individual has not developed any significant **adverse drug effects** that may exclude continued use such as skin ulceration, blistering, or moderately severe or severe dermatitis (i.e., marked skin redness with edema)

Renewal duration: 6 months

PHARMACY COVERAGE GUIDELINE

VALCHLOR™ (mechlorethamine) Generic Equivalent (if available)

- Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. **Off-Label Use of Non-Cancer Medications**
2. **Off-Label Use of Cancer Medications**

Description:

Valchlor (mechlorethamine) is indicated for the topical treatment of Stage IA and IB mycosis fungoides type cutaneous T-cell lymphoma in patients who have received prior skin-directed therapy.

Mechlorethamine, also known as nitrogen mustard, is an alkylating agent that inhibits rapidly proliferating cells, is indicated for the topical treatment of Stage IA and IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received prior skin-directed therapy.

The efficacy of Valchlor (mechlorethamine) was assessed in a randomized, multicenter, observer-blind, active-controlled, non-inferiority clinical trial of 260 patients with Stage IA, IB, and IIA mycosis fungoides-type cutaneous T-cell lymphoma who had received at least one prior skin-directed therapy. Ninety-eight percent (256 study subjects) of enrolled patients were Stage IA and IB. There were too few Stage IIA patients to evaluate. Qualifying prior therapies included topical corticosteroids, phototherapy, bexarotene gel, and topical nitrogen mustard.

Mycosis fungoides (MF also known as Alibert-Bazin syndrome or granuloma fungoides), is the most common form of cutaneous T-cell lymphoma (CTCL). It is a rare indolent form of non-Hodgkin's lymphoma that affects approximately 1,400 individuals yearly in the US and occurs more commonly in men than in women. In MF, malignant T-cells migrate and accumulate in the skin, initially resulting in dry skin and red rash that may or may not itch, eventually other skin lesions form. The malignant T-cells may also involve lymph nodes and spread to other areas such as liver, spleen, and lungs.

Sézary syndrome is a more aggressive leukemic form of CTCL with widespread skin involvement, enlarged lymph nodes and malignant lymphocytes (Sézary cells) in the skin, lymph nodes, and blood. It is a leukemic form of CTCL in which there is significant blood involvement with Sézary cells, lymphadenopathy, and erythrodermic skin. It is an advanced variant form of MF.

MF may be classified into various stages depending upon skin (T), node (N), metastasis (M), and blood (B) involvement. Stages IA, IB, and IIA are considered early-stage MF. Prognosis and survival depend on the stage at diagnosis. In the management of early-stage MF, skin-directed therapies may be categorized in two ways: "skin-limited/local therapies" for limited or localized disease and "skin-generalized therapies" for generalized skin involvement

Skin-limited therapies include topical corticosteroids, topical chemotherapy (such as nitrogen mustard), local superficial radiation (8-36 gray or Gy), topical retinoids (such as bexarotene and tazarotene), phototherapy (such as PUVA for thicker plaques, UVB, and NB-UVB for patch/thin plaques), and topical imiquimod.

ORIGINAL EFFECTIVE DATE: 02/28/2014 | ARCHIVE DATE: | LAST REVIEW DATE: 11/21/2024 | LAST CRITERIA REVISION DATE: 11/16/2023

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

VALCHLOR™ (mechlorethamine) Generic Equivalent (if available)

Skin-generalized therapies include topical corticosteroids, topical chemotherapy (such as nitrogen mustard), imiquimod, retinoids, phototherapy (such as PUVA for thicker plaques, UVB, and NB-UVB for patch/thin plaques), and total skin electron beam radiation (TSEBT [12-36 Gy]).

Systemic therapies may include (alphabetical order): acitretin, alemtuzumab, bexarotene, brentuximab vedotin, chlorambucil, cladribine, cyclophosphamide, etoposide, extracorporeal photopheresis— especially if have some blood involvement (B1 or B2), fludarabine, folotyn (pralatrexate), gemcitabine, interferons (alpha-interferon, gamma-interferon), isotretinoin, liposomal doxorubicin, methotrexate, mogamulizumab, pembrolizumab, romidepsin.

Definitions:

U.S. Food and Drug Administration (FDA) MedWatch Forms for FDA Safety Reporting
[MedWatch Forms for FDA Safety Reporting | FDA](#)

Staging of Mycosis fungoides:

In **Stage IA**, less than 10% of the skin is covered with patches, papules, and/or plaques, lymph nodes are not enlarged or abnormal, there is no visceral involvement, and the blood does not contain or has a low burden of circulating Sézary cells, defined as < 5% of peripheral blood. With **Stage IB**, 10% or more of the skin is covered with patches, papules, and/or plaques.

In **Stage IIA**, any amount of skin may be covered with patches, papules and/or plaques, lymph nodes are enlarged and may or may not have abnormal cells, there is still no visceral involvement, and the blood does not contain or has a low burden of circulating Sézary cells. **Stage IIB** has the same characteristics except now there are one or more tumorous skin lesions.

With **Stage III**, there is erythrodermic skin (greater than 80% of body surface with red patches, papules, or plaques), the lymph nodes may or may not be enlarged, when enlarged the nodes may or may not contain abnormal cells, and there is no visceral involvement. With **Stage IIIA** there are no circulating Sézary cells in the blood, with **Stage IIIB** there is a low burden of circulating Sézary cells.

In **Stages IVA and IVB**, patches, papules, plaques or tumors involve any amount of the skin surface. The lymph nodes tend to be enlarged and contain atypical cells and there is a significant level of Sézary cells in the blood. Patients with visceral involvement are classified as Stage IVB.

Clinical staging system for mycosis fungoides and Sézary syndrome:

Clinical stage	TNMB classification			
	Skin	Node	Visceral	Blood
IA – limited skin involvement	T ₁ (patches, papules, &/or plaques covering < 10% BSA)	N ₀	M ₀	B ₀ or B ₁
IB – skin only disease	T ₂ (patches, papules, &/or plaques covering ≥ 10% BSA)	N ₀	M ₀	B ₀ or B ₁
IIA	T ₁ or T ₂	N ₁ or N ₂	M ₀	B ₀ or B ₁

ORIGINAL EFFECTIVE DATE: 02/28/2014 | ARCHIVE DATE: | LAST REVIEW DATE: 11/21/2024 | LAST CRITERIA REVISION DATE: 11/16/2023

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

PHARMACY COVERAGE GUIDELINE

VALCHLOR™ (mechlorethamine) Generic Equivalent (if available)

IIB – tumor stage disease	T ₃ (one or more tumors: ≥ 1 cm in diameter)	N ₀ to N ₂	M ₀	B ₀ or B ₁
IIIA – erythrodermic disease	T ₄ (confluence of erythema $\geq 80\%$ BSA)	N ₀ to N ₂	M ₀	B ₀
IIIB – erythrodermic disease	T ₄ (confluence of erythema $\geq 80\%$ BSA)	N ₀ to N ₂	M ₀	B ₁
IVA1 - Sézary syndrome	T ₁ to T ₄	N ₀ to N ₂	M ₀	B ₂
IVA2 – Sézary syndrome or non- Sézary syndrome	T ₁ to T ₄	N ₃	M ₀	B ₀ to B ₂
IVB – visceral disease	T ₁ to T ₄	N ₀ to N ₃	M ₁	B ₀ to B ₂
	Large-cell transformation (LCT)			

To be used in conjunction with the TNMB classification system for mycosis fungoides
Skin (T), node (N), metastasis (M), and blood (B) involvement

Resources:

Valchlor (mechlorethamine) product information, revised by Helsinn Therapeutics, Inc. 05-2024. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 22, 2024.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Primary Cutaneous Lymphomas Version 3.2024 – Updated August 22, 2024. Available at <https://www.nccn.org>. Accessed October 21, 2024.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology (NCCN Guidelines®): T-Cell Lymphomas Version 4.2024 – Updated May 28, 2024. Available at <https://www.nccn.org>. Accessed October 21, 2024.

Hoppe RT, Kim YH. Clinical manifestations, pathologic features, and diagnosis of mycosis fungoides. In: UpToDate, Kuzel TM, Zic JA, Rosmarin AG, Corona R (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated September 23, 2022. Accessed October 21, 2024.

Hoppe RT, Kim YH, Horwitz S. Treatment of early stage (IA to IIA) mycosis fungoides. In: UpToDate, Kuzel TM, Zic JA, Corona R, Rosmarin AG, (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated October 24, 2023. Accessed October 21, 2024.

Hoppe RT, Kim YH, Horwitz S. Treatment of advanced stage (IIB to IV) mycosis fungoides. In: UpToDate, Kuzel TM, Zic JA, Rosmarin AG, Corona R (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated September 20, 2024. Accessed October 21, 2024.

Rook AH, Olsen EA. Clinical manifestations, pathologic features, and diagnosis of Sézary syndrome. In: UpToDate, Kuzel TM, Zic JA, Rosmarin AG (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated June 28, 2023. Accessed October 21, 2024.

Hoppe RT, Kim YH. Staging and prognosis of mycosis fungoides and Sézary syndrome. In: UpToDate, Kuzel TM, Zic JA, Rosmarin AG, Corona R (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated March 25, 2024. Accessed October 21, 2024.

Kim EJ, Rook AH. Sézary syndrome: Treatment and prognosis. In: UpToDate, Kuzel TM, Zic JA, Rosmarin AG (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through September 2024. Topic last updated July 03, 2024. Accessed October 21, 2024.

Off Label Use of Cancer Medications: A.R.S. §§ 20-826(R) & (S). Subscription contracts; definitions.

Off Label Use of Cancer Medications: A.R.S. §§ 20-1057(V) & (W). Evidence of coverage by health care service organizations; renewability; definitions.

ORIGINAL EFFECTIVE DATE: 02/28/2014 | ARCHIVE DATE: | LAST REVIEW DATE: 11/21/2024 | LAST CRITERIA REVISION DATE: 11/16/2023

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.