

2024 Retiree Pharmacy Prescription Drug Plan (PDP) Formulary

Medicare Part D: Compass Formulary 2024

Please click here.

For Medicare Part D: Prior Authorization Criteria

Please click here.

For Medicare Part D: Step Therapy Criteria

Please click here.

For more recent information or other questions, please contact:

Retiree Pharmacy PDP

Pharmacy Service at 1-877-461-9218.

For TTY users, 711 National Relay Service, Oct. 1 – March 31, 8 a.m. – 8 p.m. ET, seven days a week, and April 1 – Aug. 30, 8 a.m. – 8 p.m. ET, Monday – Sunday.

Visit [medicare.highmark.com](https://www.medicare.highmark.com).

Notice: Oral Antivirals for COVID-19 Coverage

Paxlovid is covered with \$0 copay when dispensed at a preferred or non-preferred pharmacy.

Formulary ID: 24018 Version: 17

Updated: 9/2024

Note to existing members: This formulary has changed since last year. Please review this document to make sure that it still contains the drugs you take.

When this drug list (formulary) refers to “we,” “us,” or “our,” it means Highmark Blue Cross Blue Shield.

When it refers to “plan” or “our plan,” it means 2024 Retiree Pharmacy PDP.

This document includes a list of the drugs (formulary) for our plan which is current as of January 1, 2024. For an updated formulary, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2025, and from time to time during the year.

What is the 2024 Retiree Pharmacy PDP Formulary?

A formulary is a list of covered drugs selected by our plan in consultation with a team of health care providers, which represents the prescription therapies believed to be a necessary part of a quality treatment program. Our plan will generally cover the drugs listed in our formulary as long as the drug is medically necessary, the prescription is filled at one of our plan’s network pharmacies, and other plan rules are followed. For more information on how to fill your prescriptions, please review your Evidence of Coverage.

Can the 2024 Retiree Pharmacy PDP Formulary (drug list) change?

Most changes in drug coverage happen on January 1, but we may add or remove drugs on the Drug List during the year, move them to different cost-sharing tiers, or add new restrictions. We must follow Medicare rules in making these changes.

Changes that can affect you this year: In the below cases, you will be affected by coverage changes during the year:

- **New generic drugs.** We may immediately remove a brand name drug on our Drug List if we are replacing it with a new generic drug that will appear on the same or lower cost sharing tier and with the same or fewer restrictions. Also, when adding the new generic drug, we may decide to keep the brand name drug on our Drug List, but immediately move it to a different cost-sharing tier or add new restrictions. If you are currently taking that brand name drug, we may not tell you in advance before we make that change, but we will later provide you with information about the specific change(s) we have made.
- If we make such a change, you or your prescriber can ask us to make an exception and continue to cover the brand name drug for you. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled “How do I request an exception to the Retiree Pharmacy PDP Formulary?”

- o **Drugs removed from the market.** If the Food and Drug Administration deems a drug on our formulary to be unsafe or the drug’s manufacturer removes the drug from the market, we will immediately remove the drug from our formulary and provide notice to members who take the drug.
- **Other changes.** We may make other changes that affect members currently taking a drug. For instance, we may add a generic drug that is not new to market to replace a brand name drug currently on the formulary or add new restrictions to the brand name drug or move it to a different cost-sharing tier or both. Or we may make changes based on new clinical guidelines. If we remove drugs from our formulary, add prior authorization, quantity limits and/or step therapy restrictions on a drug, or move a drug to a higher cost-sharing tier, we must notify affected members of the change at least 30 days before the change becomes effective, or at the time the member requests a refill of the drug, at which time the member will receive a 31-day supply of the drug.
 - o If we make these other changes, you or your prescriber can ask us to make an exception and continue to cover the brand name drug for you. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled “How do I request an exception to the Retiree Pharmacy PDP Formulary?”

Changes that will not affect you if you are currently taking the drug. Generally, if you are taking a drug on our 2024 formulary that was covered at the beginning of the year, we will not discontinue or reduce coverage of the drug during the 2024 coverage year except as described above. This means these drugs will remain available at the same cost-sharing and with no new restrictions for those members taking them for the remainder of the coverage year. You will not get direct notice this year about changes that do not affect you. However, on January 1 of the next year, such changes would affect you, and it is important to check the Drug List for the new benefit year for any changes to drugs.

The enclosed formulary is current as of January 1, 2024. To get updated information about the drugs covered by our plan, please contact us. Our contact information appears on the front and back cover pages. In the event of mid-year non-maintenance formulary changes, members will be notified by mail and prospective members will receive an update with this formulary. The most up-to-date formulary is available on our website, [medicare.highmark.com](https://www.medicare.highmark.com).

How do I use the 2024 Retiree Pharmacy PDP Formulary?

There are two ways to find your drug within the formulary:

Medical Condition

The formulary begins on page 9. The drugs in this formulary are grouped into categories depending on the type of medical conditions that they are used to treat. For example, drugs used to treat a heart condition are listed under the category, “Cardiovascular – Hypertension & Lipids.” If you know what your drug is used for, look for the category name in the list that begins on page 9. Then look under the category name for your drug.

Alphabetical Listing

If you are not sure what category to look under, you should look for your drug in the Index that begins at the end of this document. The Index provides an alphabetical list of all of the drugs included in this document. Both brand name drugs and generic drugs are listed in the Index. Look in the Index and find your drug. Next to your drug, you will see the page number where you can find coverage information. Turn to the page listed in the Index and find the name of your drug in the first column of the list.

What are generic drugs?

Our plan covers both brand name drugs and generic drugs. A generic drug is approved by the FDA as having the same active ingredient as the brand name drug. Generally, generic drugs cost less than brand name drugs.

Are there any restrictions on my coverage?

Some covered drugs may have additional requirements or limits on coverage. These requirements and limits may include:

- **Prior Authorization:** Our plan requires you or your physician to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don't get approval, our plan may not cover the drug.
- **Quantity Limits:** For certain drugs, our plan limits the amount of the drug we will cover. For example, our plan provides 31 tablets, per 31 days, per prescription for 100mg losartan. This may be in addition to a standard one-month or three-month supply.
- **Step Therapy:** In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. For example, if Drug A and Drug B both treat your medical condition, our plan may not cover Drug B unless you try Drug A first. If Drug A does not work for you, our plan will then cover Drug B.

You can find out if your drug has any additional requirements or limits by looking in the formulary that begins on page 9. You can also get more information about the restrictions applied to specific covered drugs by visiting our website. We have posted online document(s) that explain(s) our prior authorization and step therapy restrictions. You may also ask us to send you a copy. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You can ask our plan to make an exception to these restrictions or limits or for a list of other, similar drugs that may treat your health condition. See the section, "How do I request an exception to the Retiree Pharmacy PDP Formulary?" on page 5 for information about how to request an exception.

What if my drug is not on the 2024 Retiree Pharmacy PDP Formulary?

If your drug is not included in this formulary (list of covered drugs), you should first contact Pharmacy Service and ask if your drug is covered.

If you learn that our plan does not cover your drug, you have two options:

- You can ask Pharmacy Service for a list of similar drugs that are covered by our plan. When you receive the list, show it to your doctor and ask them to prescribe a similar drug that is covered by our plan.
- You can ask our plan to make an exception and cover your drug. See below for information about how to request an exception.

How do I request an exception to the 2024 Retiree Pharmacy PDP Formulary?

You can ask our plan to make an exception to our coverage rules. There are several types of exceptions that you can ask us to make.

- You can ask us to cover a drug even if it is not on our formulary. If approved, this drug will be covered at a pre-determined cost-sharing level, and you would not be able to ask us to provide the drug at a lower cost-sharing level.
- You can ask us to cover a formulary drug at a lower cost-sharing level unless the drug is on the specialty tier. If approved, this would lower the amount you must pay for your drug.
- You can ask us to waive coverage restrictions or limits on your drug. For example, for certain drugs, our plan limits the amount of the drug that we will cover. If your drug has a quantity limit, you can ask us to waive the limit and cover a greater amount.

Generally, our plan will only approve your request for an exception if the alternative drugs included on the plan's formulary, the lower cost-sharing drug or additional utilization restrictions, would not be as effective in treating your condition and/or would cause you to have adverse medical effects.

You should contact us to ask us for an initial coverage decision for a formulary, or utilization restriction exception. **When you request a formulary or utilization restriction exception, you should submit a statement from your prescriber or physician supporting your request.** Generally, we must make our decision within 72 hours of getting your prescriber's supporting statement. You can request an expedited (fast) exception if you or your doctor believe that your health could be seriously harmed by waiting up to 72 hours for a decision. If your request to expedite is granted, we must give you a decision no later than 24 hours after we get a supporting statement from your doctor or other prescriber.

What do I do before I can talk to my doctor about changing my drugs or requesting an exception?

As a new or continuing member in our plan, you may be taking drugs that are not on our formulary. Or you may be taking a drug that is on our formulary but your ability to get it is limited. For example, you may need a prior authorization from us before you can fill your prescription. You should talk to your doctor to decide if you should switch to an appropriate drug that we cover or request a formulary exception so that we will cover the drug you take. While you talk to your doctor to determine the right course of action for you, we may cover your drug in certain cases during the first 90 days you are a member of our plan.

For each of your drugs that is not on our formulary or if your ability to get your drugs is limited, we will cover a temporary 31-day supply. If your prescription is written for fewer days, we'll allow refills to provide up to a maximum 31-day supply of medication. After your first 31-day supply, we will not pay for these drugs, even if you have been a member of the plan less than 90 days.

If you are a resident of a long-term care facility and you need a drug that is not on our formulary or if your ability to get your drugs is limited, but you are past the first 90 days of membership in our plan, we will cover a 31-day emergency supply of that drug while you pursue a formulary exception.

The above transition process will be implemented to accommodate you if you have an immediate need for a non-formulary drug or a drug that requires prior authorization due to a change in your level of care while you are waiting for an exception request to be processed.

For more information

For more detailed information about your plan's prescription drug coverage, please review your Evidence of Coverage and other plan materials.

If you have questions about your plan, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

If you have general questions about Medicare prescription drug coverage, please call Medicare at 1-800-MEDICARE (1-800-633-4227) 24 hours a day / 7 days a week. TTY users should call 1-877-486-2048. Or visit <http://www.medicare.gov>.

2024 Retiree Pharmacy PDP Formulary

The formulary that begins on the next page provides coverage information about the drugs covered by your plan. If you have trouble finding your drug in the list, turn to the Index at the end of this document.

The first column of the chart lists the drug name. Brand name drugs are capitalized (e.g., ABELCET) and generic drugs are listed in lowercase italics (e.g., *abacavir*).

The information in the Requirements/Limits column tells you if your plan has any special requirements for coverage of your drug.

The following is a Formulary Format Example Only:

Drug Name	Compass Drug Tier	Requirements/ Limits
Anti - Infectives		
XYZ DRUG	NF	QL- 28

Table of Contents

Anti - Infectives.....	3
Antineoplastic / Immunosuppressant Drugs.....	16
Autonomic / Cns Drugs, Neurology / Psych.....	25
Cardiovascular, Hypertension / Lipids.....	58
Dermatologicals/Topical Therapy.....	70
Diagnostics / Miscellaneous Agents.....	79
Ear, Nose / Throat Medications.....	82
Endocrine/Diabetes.....	83
Gastroenterology.....	91
Immunology, Vaccines / Biotechnology.....	98
Miscellaneous Supplies.....	102
Musculoskeletal / Rheumatology.....	102
Obstetrics / Gynecology.....	106
Ophthalmology.....	113
Respiratory And Allergy.....	116
Urologicals.....	122
Vitamins, Hematinics / Electrolytes.....	124

Requirements/Limits

LA = Limited access

PA = Prior authorization required

PA-BvD = This drug may be covered under Medicare part B or D depending on the circumstance.

Information may need to be submitted describing the use and setting of the drug to make the determination.

PA-NS = Prior authorization required for new starts only

QL = Quantity limit applies. The quantity limit is noted for each drug. For example, if the quantity limit is QL (90 EA per 180 days), the quantity limit would be 90 units per 180-day supply.

ST = Step therapy applies

ST-NS = Step therapy applies to new starts only

Drug Tier**T1** = Cost-Sharing Tier 1 includes preferred generic drugs. This is the lowest cost-sharing tier.**T2** = Cost-Sharing Tier 2 includes generic drugs.**T3** = Cost-Sharing Tier 3 includes preferred brand name drugs and may include some single-sourced drugs (those generic drugs made by a single manufacturer).**T4** = Cost-Sharing Tier 4 includes non-preferred brand name drugs and may include some single-sourced generic drugs (those generic drugs made by a single manufacturer).**T5** = Cost-Sharing Tier 5 includes specialty drugs. This is the highest cost-sharing tier.**lowercase italics** = Generic drugs**UPPERCASE BOLD** = Brand name drugs

Drug Name	Drug Tier	Requirements/Limits
Anti - Infectives		
<i>abacavir</i>	T2	
<i>abacavir-lamivudine</i>	T3	
ABELCET	T4	PA-BvD
<i>acyclovir oral capsule</i>	T2	
<i>acyclovir oral suspension 200 mg/5 ml</i>	T2	
<i>acyclovir oral tablet</i>	T2	
<i>acyclovir sodium intravenous solution</i>	T4	PA-BvD
<i>adefovir</i>	T4	
AEMCOLO	T4	QL (12 EA per 3 days)
<i>albendazole</i>	T4	
<i>amantadine hcl oral capsule</i>	T2	QL (124 EA per 31 days)
<i>amantadine hcl oral solution</i>	T2	
<i>amantadine hcl oral tablet</i>	T2	
AMBISOME	T5	PA-BvD
<i>amikacin injection solution 500 mg/2 ml</i>	T4	
<i>amoxicillin oral capsule</i>	T2	
<i>amoxicillin oral suspension for reconstitution</i>	T2	
<i>amoxicillin oral tablet</i>	T2	
<i>amoxicillin oral tablet, chewable 125 mg, 250 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>amoxicillin-pot clavulanate oral suspension for reconstitution</i>	T2	
<i>amoxicillin-pot clavulanate oral tablet</i>	T2	
<i>amoxicillin-pot clavulanate oral tablet extended release 12 hr</i>	T4	
<i>amoxicillin-pot clavulanate oral tablet, chewable</i>	T2	
<i>amphotericin b</i>	T4	PA-BvD
<i>amphotericin b liposome</i>	T5	PA-BvD
<i>ampicillin oral capsule 500 mg</i>	T2	
<i>ampicillin sodium injection recon soln 1 gram, 10 gram, 125 mg</i>	T4	
<i>ampicillin-sulbactam injection</i>	T4	
ANCOBON	T5	
APTIVUS	T5	
ARIKAYCE	T5	PA
<i>atazanavir</i>	T4	
<i>atovaquone</i>	T4	
<i>atovaquone-proguanil</i>	T2	
AUGMENTIN ES-600	T4	
AUGMENTIN ORAL SUSPENSION FOR RECONSTITUTION 125-31.25 MG/5 ML	T4	
AVYCAZ	T5	
AZACTAM	T4	
<i>azithromycin intravenous</i>	T4	
<i>azithromycin oral</i>	T2	
<i>aztreonam</i>	T4	
BACTRIM	T4	
BACTRIM DS	T4	
BARACLUDE	T5	
BAXDELA	T5	
<i>benznidazole</i>	T3	PA
BETHKIS	T5	PA
BICILLIN C-R	T3	
BICILLIN L-A INTRAMUSCULAR SYRINGE 1,200,000 UNIT/2 ML, 600,000 UNIT/ML	T4	
BICILLIN L-A INTRAMUSCULAR SYRINGE 2,400,000 UNIT/4 ML	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
BIKTARVY	T5	QL (31 EA per 31 days)
BILTRICIDE	T4	
CANCIDAS	T5	
<i>caspofungin intravenous recon soln 50 mg</i>	T5	
<i>caspofungin intravenous recon soln 70 mg</i>	T4	
CAYSTON	T5	PA
<i>cefaclor oral capsule</i>	T2	
<i>cefaclor oral suspension for reconstitution 250 mg/5 ml</i>	T2	
<i>cefaclor oral tablet extended release 12 hr</i>	T4	
<i>cefadroxil oral capsule</i>	T2	
<i>cefadroxil oral suspension for reconstitution 250 mg/5 ml, 500 mg/5 ml</i>	T2	
<i>cefadroxil oral tablet</i>	T2	
<i>cefazolin injection recon soln 1 gram, 10 gram, 500 mg</i>	T4	
<i>cefdinir</i>	T2	
<i>cefepime injection</i>	T4	
<i>cefixime</i>	T4	
<i>cefotetan injection</i>	T4	
<i>cefoxitin</i>	T4	
<i>cefpodoxime</i>	T2	
<i>cefprozil</i>	T2	
<i>ceftazidime</i>	T4	
<i>ceftriaxone injection recon soln 1 gram, 10 gram, 2 gram, 250 mg, 500 mg</i>	T4	
<i>cefuroxime axetil oral tablet</i>	T2	
<i>cefuroxime sodium injection recon soln 750 mg</i>	T4	
<i>cefuroxime sodium intravenous recon soln 1.5 gram</i>	T4	
<i>cephalexin oral capsule 250 mg, 500 mg</i>	T2	
<i>cephalexin oral capsule 750 mg</i>	T4	
<i>cephalexin oral suspension for reconstitution</i>	T2	
<i>cephalexin oral tablet</i>	T4	
<i>chloroquine phosphate oral tablet 250 mg</i>	T2	QL (50 EA per 30 days)
<i>chloroquine phosphate oral tablet 500 mg</i>	T2	QL (25 EA per 30 days)
CIMDUO	T5	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
CIPRO ORAL SUSPENSION, MICROCAPSULE RECON	T4	
CIPRO ORAL TABLET 250 MG, 500 MG	T4	
<i>ciprofloxacin hcl oral tablet 250 mg, 500 mg</i>	T1	
<i>ciprofloxacin hcl oral tablet 750 mg</i>	T2	
<i>ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml</i>	T4	
<i>clarithromycin</i>	T2	
CLEOCIN HCL	T4	
CLEOCIN INJECTION	T4	
CLEOCIN PEDIATRIC	T4	
<i>clindamycin hcl</i>	T2	
<i>clindamycin in 5 % dextrose</i>	T4	
CLINDAMYCIN PEDIATRIC	T2	
<i>clindamycin phosphate injection</i>	T4	
<i>clotrimazole mucous membrane</i>	T2	
COARTEM	T4	
<i>colistin (colistimethate na)</i>	T4	
COMBIVIR	T5	
COMPLERA	T5	
CRESEMBA ORAL	T5	PA
CUBICIN RF	T5	
<i>cycloserine</i>	T4	
DALVANCE	T5	
<i>dapsone oral</i>	T3	
<i>daptomycin intravenous recon soln 350 mg</i>	T5	
<i>daptomycin intravenous recon soln 500 mg</i>	T4	
DARAPRIM	T5	PA
<i>darunavir</i>	T5	
DELSTRIGO	T5	QL (31 EA per 31 days)
<i>demeclocycline</i>	T4	
DESCOVY	T5	QL (31 EA per 31 days)
<i>dicloxacillin</i>	T2	
DIFICID ORAL SUSPENSION FOR RECONSTITUTION	T5	QL (136 ML per 12 days)
DIFICID ORAL TABLET	T5	QL (20 EA per 10 days)
DIFLUCAN ORAL SUSPENSION FOR RECONSTITUTION 40 MG/ML	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
DIFLUCAN ORAL TABLET 100 MG, 200 MG	T4	
DORYX MPC ORAL TABLET,DELAYED RELEASE (DR/EC) 60 MG	T5	
DOVATO	T5	QL (31 EA per 31 days)
DOXY-100	T4	
<i>doxycycline hyclate oral capsule</i>	T2	
<i>doxycycline hyclate oral tablet 100 mg, 150 mg, 50 mg, 75 mg</i>	T4	
<i>doxycycline hyclate oral tablet 20 mg</i>	T2	
<i>doxycycline hyclate oral tablet, delayed release (dr/ec)</i>	T4	
<i>doxycycline monohydrate oral capsule 100 mg, 50 mg</i>	T2	
<i>doxycycline monohydrate oral capsule 150 mg, 75 mg</i>	T4	
<i>doxycycline monohydrate oral capsule, ir - delay rel, biphase</i>	T4	
<i>doxycycline monohydrate oral suspension for reconstitution</i>	T4	
<i>doxycycline monohydrate oral tablet 100 mg, 50 mg, 75 mg</i>	T2	
<i>doxycycline monohydrate oral tablet 150 mg</i>	T4	
E.E.S. 400 ORAL TABLET	T4	
E.E.S. GRANULES	T4	
EDURANT	T5	
<i>efavirenz</i>	T4	
<i>efavirenz-emtricitabin-tenofovir</i>	T5	
<i>efavirenz-lamivudine-tenofovir disoproxil fumarate</i>	T5	QL (31 EA per 31 days)
<i>emtricitabine</i>	T2	
<i>emtricitabine-tenofovir (tdf) oral tablet 100-150 mg, 133-200 mg, 167-250 mg</i>	T5	
<i>emtricitabine-tenofovir (tdf) oral tablet 200-300 mg</i>	T4	
EMTRIVA ORAL CAPSULE	T4	
EMTRIVA ORAL SOLUTION	T3	
EMVERM	T5	
<i>entecavir</i>	T4	
EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG	T5	PA; QL (28 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
EPCLUSA ORAL PELLETS IN PACKET 200-50 MG	T5	PA; QL (56 EA per 28 days)
EPCLUSA ORAL TABLET	T5	PA; QL (28 EA per 28 days)
EPIVIR	T4	
EPZICOM	T5	
ERAXIS(WATER DILUENT) INTRAVENOUS RECON SOLN 100 MG	T5	
ERAXIS(WATER DILUENT) INTRAVENOUS RECON SOLN 50 MG	T4	
<i>ertapenem</i>	T4	
ERYPED 200	T4	
ERYPED 400	T4	
ERY-TAB	T4	
ERYTHROCIN (AS STEARATE) ORAL TABLET 250 MG	T4	
ERYTHROCIN INTRAVENOUS RECON SOLN 500 MG	T4	
<i>erythromycin ethylsuccinate oral suspension for reconstitution 200 mg/5 ml</i>	T4	
<i>erythromycin ethylsuccinate oral suspension for reconstitution 400 mg/5 ml</i>	T5	
<i>erythromycin ethylsuccinate oral tablet</i>	T4	
<i>erythromycin oral</i>	T4	
<i>ethambutol</i>	T2	
<i>etravirine</i>	T5	
EVOTAZ	T5	
<i>famciclovir</i>	T2	
FIRVANQ	T4	
FLAGYL ORAL CAPSULE	T4	
<i>fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml</i>	T4	
<i>fluconazole oral suspension for reconstitution</i>	T3	
<i>fluconazole oral tablet</i>	T2	
<i>flucytosine</i>	T5	
<i>fosamprenavir</i>	T5	
<i>fosfomycin tromethamine</i>	T4	
FUZEON SUBCUTANEOUS RECON SOLN	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml</i>	T4	
<i>gentamicin injection solution 40 mg/ml</i>	T4	
GENVOYA	T5	
<i>griseofulvin microsize</i>	T4	
<i>griseofulvin ultramicrosize</i>	T4	
HARVONI ORAL PELLETS IN PACKET	T5	PA; QL (28 EA per 28 days)
HARVONI ORAL TABLET 90-400 MG	T5	PA; QL (28 EA per 28 days)
HIPREX	T4	
HUMATIN	T4	
<i>hydroxychloroquine oral tablet 100 mg</i>	T4	QL (93 EA per 31 days)
<i>hydroxychloroquine oral tablet 200 mg</i>	T2	QL (93 EA per 31 days)
<i>hydroxychloroquine oral tablet 300 mg</i>	T4	QL (62 EA per 31 days)
<i>hydroxychloroquine oral tablet 400 mg</i>	T4	QL (31 EA per 31 days)
<i>imipenem-cilastatin</i>	T4	
IMPAVIDO	T5	
INTELENCE ORAL TABLET 100 MG, 200 MG	T5	
INTELENCE ORAL TABLET 25 MG	T4	
INVANZ INJECTION	T4	
ISENTRESS HD	T5	
ISENTRESS ORAL POWDER IN PACKET	T5	
ISENTRESS ORAL TABLET	T5	
ISENTRESS ORAL TABLET,CHEWABLE 100 MG	T5	
ISENTRESS ORAL TABLET,CHEWABLE 25 MG	T3	
<i>isoniazid oral</i>	T2	
<i>itraconazole</i>	T4	PA
<i>ivermectin oral</i>	T2	PA
JULUCA	T5	
KALETRA ORAL SOLUTION	T5	
KALETRA ORAL TABLET	T4	
<i>ketoconazole oral</i>	T2	
KITABIS PAK	T4	PA
KRINTAFEL	T4	
LAGEVRIO (EUA)	T3	QL (360 EA per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>lamivudine</i>	T3	
<i>lamivudine-zidovudine</i>	T3	
LAMPIT	T4	PA
<i>ledipasvir-sofosbuvir</i>	T5	PA; QL (28 EA per 28 days)
<i>levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml</i>	T3	
<i>levofloxacin oral</i>	T2	
LEXIVA ORAL SUSPENSION	T4	
LEXIVA ORAL TABLET	T5	
<i>linezolid</i>	T4	
<i>linezolid in dextrose 5%</i>	T4	
LIVTENCITY	T5	PA; QL (372 EA per 31 days)
<i>lopinavir-ritonavir oral solution</i>	T4	
<i>lopinavir-ritonavir oral tablet 100-25 mg</i>	T3	
<i>lopinavir-ritonavir oral tablet 200-50 mg</i>	T4	
MACROBID	T4	QL (90 EA per 365 days)
MACRODANTIN ORAL CAPSULE 100 MG	T4	QL (90 EA per 365 days)
MACRODANTIN ORAL CAPSULE 25 MG	T4	QL (360 EA per 365 days)
MACRODANTIN ORAL CAPSULE 50 MG	T4	QL (180 EA per 365 days)
MALARONE	T4	
MALARONE PEDIATRIC	T4	
<i>maraviroc oral tablet 150 mg</i>	T5	
<i>maraviroc oral tablet 300 mg</i>	T4	
MAVYRET ORAL PELLETS IN PACKET	T5	PA; QL (140 EA per 28 days)
MAVYRET ORAL TABLET	T5	PA; QL (84 EA per 28 days)
<i>mefloquine</i>	T2	
MEPRON	T4	
<i>meropenem intravenous recon soln 1 gram, 500 mg</i>	T4	
<i>methenamine hippurate</i>	T2	
<i>metronidazole in nacl (iso-os)</i>	T4	
<i>metronidazole oral capsule</i>	T4	
<i>metronidazole oral tablet</i>	T2	
<i>micafungin intravenous recon soln 100 mg</i>	T4	
<i>micafungin intravenous recon soln 50 mg</i>	T5	
<i>minocycline oral capsule</i>	T2	
<i>minocycline oral tablet</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>minocycline oral tablet extended release 24 hr</i>	T4	
<i>moxifloxacin oral</i>	T2	
<i>moxifloxacin-sod.chloride(iso)</i>	T4	
MYAMBUTOL ORAL TABLET 400 MG	T4	
MYCAMINE INTRAVENOUS RECON SOLN 50 MG	T5	
MYCOBUTIN	T4	
<i>nafcillin injection recon soln 1 gram, 2 gram</i>	T4	
<i>nafcillin injection recon soln 10 gram</i>	T5	
NEBUPENT	T4	PA-BvD
<i>neomycin</i>	T2	
<i>nevirapine oral suspension</i>	T4	
<i>nevirapine oral tablet</i>	T3	
<i>nevirapine oral tablet extended release 24 hr 400 mg</i>	T4	
<i>nitazoxanide</i>	T4	
<i>nitrofurantoin macrocrystal oral capsule 100 mg</i>	T3	QL (90 EA per 365 days)
<i>nitrofurantoin macrocrystal oral capsule 25 mg</i>	T4	QL (360 EA per 365 days)
<i>nitrofurantoin macrocrystal oral capsule 50 mg</i>	T3	QL (180 EA per 365 days)
<i>nitrofurantoin monohyd/m-cryst</i>	T3	QL (90 EA per 365 days)
<i>nitrofurantoin oral suspension 25 mg/5 ml</i>	T5	QL (1800 ML per 365 days)
<i>nitrofurantoin oral suspension 50 mg/5 ml</i>	T5	QL (900 ML per 365 days)
NORVIR ORAL POWDER IN PACKET	T4	
NORVIR ORAL TABLET	T4	
NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON	T5	PA; QL (32 EA per 31 days)
NOXAFIL ORAL SUSPENSION	T5	PA
NOXAFIL ORAL TABLET,DELAYED RELEASE (DR/EC)	T5	PA
NUZYRA	T5	
<i>nystatin oral</i>	T2	
ODEFSEY	T5	QL (31 EA per 31 days)
<i>ofloxacin oral tablet 300 mg, 400 mg</i>	T4	
ORACEA	T4	
<i>oseltamivir oral capsule 30 mg</i>	T2	QL (170 EA per 365 days)
<i>oseltamivir oral capsule 45 mg, 75 mg</i>	T2	QL (90 EA per 365 days)
<i>oseltamivir oral suspension for reconstitution</i>	T3	QL (1080 ML per 365 days)
<i>oxacillin</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>oxacillin in dextrose(iso-osm)</i>	T4	
PAXLOVID ORAL TABLETS,DOSE PACK 150-100 MG	T3	QL (180 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 300 MG (150 MG X 2)-100 MG	T3	QL (270 EA per 365 days)
<i>penicillin g pot in dextrose intravenous piggyback 2 million unit/50 ml, 3 million unit/50 ml</i>	T4	
<i>penicillin g potassium injection recon soln 20 million unit</i>	T4	
<i>penicillin g sodium</i>	T5	
<i>penicillin v potassium</i>	T2	
PENTAM	T4	
<i>pentamidine inhalation</i>	T4	PA-BvD
<i>pentamidine injection</i>	T4	
PIFELTRO	T5	QL (62 EA per 31 days)
<i>piperacillin-tazobactam intravenous recon soln 2.25 gram, 3.375 gram, 4.5 gram, 40.5 gram</i>	T4	
PLAQUENIL	T4	QL (93 EA per 31 days)
<i>polymyxin b sulfate</i>	T4	
<i>posaconazole oral</i>	T5	PA
<i>praziquantel</i>	T4	
<i>pretomanid</i>	T4	PA; QL (31 EA per 31 days)
PREVYMIS ORAL	T5	QL (31 EA per 31 days)
PREZCOBIX	T5	
PREZISTA ORAL SUSPENSION	T5	
PREZISTA ORAL TABLET 150 MG, 600 MG, 75 MG, 800 MG	T5	
PRIFTIN	T3	
<i>primaquine</i>	T3	
PRIMAXIN IV INTRAVENOUS RECON SOLN 500 MG	T4	
<i>pyrazinamide</i>	T4	
<i>pyrimethamine</i>	T5	PA
QUALAQUIN	T4	PA; QL (42 EA per 28 days)
<i>quinine sulfate</i>	T4	PA; QL (42 EA per 28 days)
RELENZA DISKHALER	T4	
RETROVIR ORAL CAPSULE	T4	
RETROVIR ORAL SYRUP	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
REYATAZ ORAL CAPSULE 200 MG, 300 MG	T5	
REYATAZ ORAL POWDER IN PACKET	T5	
<i>ribavirin oral capsule</i>	T3	
<i>ribavirin oral tablet 200 mg</i>	T3	
<i>rifabutin</i>	T4	
<i>rifampin intravenous</i>	T5	
<i>rifampin oral</i>	T3	
<i>rimantadine</i>	T2	
<i>ritonavir</i>	T3	
RUKOBIA	T5	QL (62 EA per 31 days)
SELZENTRY ORAL SOLUTION	T5	
SELZENTRY ORAL TABLET 150 MG, 300 MG, 75 MG	T5	
SELZENTRY ORAL TABLET 25 MG	T4	
SEYSARA	T5	PA
SIRTURO	T5	PA
SIVEXTRO INTRAVENOUS	T5	
SIVEXTRO ORAL	T5	QL (6 EA per 31 days)
<i>sofosbuvir-velpatasvir</i>	T5	PA; QL (28 EA per 28 days)
SOLODYN ORAL TABLET EXTENDED RELEASE 24 HR 80 MG	T4	
SOLOSEC	T4	
SOVALDI	T5	PA; QL (28 EA per 28 days)
SOVUNA ORAL TABLET 200 MG	T4	QL (93 EA per 31 days)
SOVUNA ORAL TABLET 300 MG	T4	QL (62 EA per 31 days)
SPORANOX ORAL CAPSULE	T5	PA
SPORANOX ORAL SOLUTION	T4	PA
<i>streptomycin</i>	T5	
STRIBILD	T5	
STROMEKTOL	T4	PA
<i>sulfadiazine</i>	T4	
<i>sulfamethoxazole-trimethoprim oral suspension</i>	T2	
<i>sulfamethoxazole-trimethoprim oral tablet</i>	T1	
SUNLENCA ORAL	T5	
SYMFI	T5	QL (31 EA per 31 days)
SYMFI LO	T5	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SYMTUZA	T5	QL (31 EA per 31 days)
TAMIFLU ORAL CAPSULE 30 MG	T4	QL (170 EA per 365 days)
TAMIFLU ORAL CAPSULE 45 MG, 75 MG	T4	QL (90 EA per 365 days)
TAMIFLU ORAL SUSPENSION FOR RECONSTITUTION	T4	QL (1080 ML per 365 days)
TARGADOX	T4	
TAZICEF INJECTION	T4	
TEFLARO INTRAVENOUS RECON SOLN 400 MG	T4	
TEFLARO INTRAVENOUS RECON SOLN 600 MG	T5	
<i>tenofovir disoproxil fumarate</i>	T3	
<i>terbinafine hcl oral</i>	T2	QL (90 EA per 180 days)
<i>tetracycline oral capsule</i>	T4	
<i>tigecycline</i>	T5	
<i>tinidazole</i>	T2	
TIVICAY ORAL TABLET 10 MG	T4	
TIVICAY ORAL TABLET 25 MG, 50 MG	T5	
TIVICAY PD	T5	
TOBI	T5	PA
TOBI PODHALER	T5	PA; QL (224 EA per 56 days)
<i>tobramycin in 0.225 % nacl</i>	T5	PA
<i>tobramycin inhalation</i>	T5	PA
<i>tobramycin sulfate injection solution</i>	T4	
TOLSURA	T5	PA; QL (130 EA per 31 days)
TRECTOR	T4	
<i>trimethoprim</i>	T2	
TRIUMEQ	T5	
TRIUMEQ PD	T5	QL (186 EA per 31 days)
TRIZIVIR	T5	
TRUVADA	T5	
TYBOST	T3	
TYGACIL	T4	
UNASYN INJECTION RECON SOLN 15 GRAM, 3 GRAM	T4	
VABOMERE	T4	
<i>valacyclovir</i>	T2	
VALCYTE	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>valganciclovir oral recon soln</i>	T5	
<i>valganciclovir oral tablet</i>	T3	
VALTREX	T4	
VANCOCIN ORAL CAPSULE 125 MG	T4	PA; QL (124 EA per 31 days)
VANCOCIN ORAL CAPSULE 250 MG	T5	PA; QL (248 EA per 31 days)
<i>vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg</i>	T4	
<i>vancomycin oral capsule 125 mg</i>	T4	PA; QL (124 EA per 31 days)
<i>vancomycin oral capsule 250 mg</i>	T4	PA; QL (248 EA per 31 days)
<i>vancomycin oral recon soln</i>	T4	
VEMLIDY	T5	QL (31 EA per 31 days)
VFEND IV	T4	PA
VFEND ORAL SUSPENSION FOR RECONSTITUTION	T5	
VFEND ORAL TABLET	T4	
VIBRAMYCIN ORAL CAPSULE 100 MG	T4	
VIRACEPT ORAL TABLET	T5	
VIREAD	T5	
VIVJOA	T4	PA; QL (18 EA per 84 days)
<i>voriconazole intravenous</i>	T5	PA
<i>voriconazole oral suspension for reconstitution</i>	T5	
<i>voriconazole oral tablet</i>	T4	
VOSEVI	T5	PA; QL (28 EA per 28 days)
XIFAXAN ORAL TABLET 200 MG	T4	QL (27 EA per 365 days)
XIFAXAN ORAL TABLET 550 MG	T5	PA; QL (62 EA per 31 days)
XOFLUZA ORAL TABLET 40 MG, 80 MG	T3	QL (9 EA per 365 days)
ZEMDRI	T5	
ZEPATIER	T5	PA; QL (28 EA per 28 days)
ZERBAXA	T4	
ZIAGEN ORAL SOLUTION	T4	
<i>zidovudine</i>	T2	
ZITHROMAX INTRAVENOUS	T4	
ZITHROMAX ORAL PACKET	T4	
ZITHROMAX ORAL SUSPENSION FOR RECONSTITUTION	T4	
ZITHROMAX ORAL TABLET 250 MG, 500 MG	T4	
ZITHROMAX TRI-PAK	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZITHROMAX Z-PAK	T4	
ZOSYN IN DEXTROSE (ISO-OSM) INTRAVENOUS PIGGYBACK 2.25 GRAM/50 ML	T4	
ZYVOX INTRAVENOUS PIGGYBACK 600 MG/300 ML	T4	
ZYVOX ORAL	T5	
Antineoplastic / Immunosuppressant Drugs		
<i>abiraterone oral tablet 250 mg</i>	T5	PA-NS; QL (124 EA per 31 days)
<i>abiraterone oral tablet 500 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
AFINITOR	T5	PA-NS; QL (31 EA per 31 days)
AFINITOR DISPERZ ORAL TABLET FOR SUSPENSION 2 MG, 5 MG	T5	PA-NS; QL (62 EA per 31 days)
AFINITOR DISPERZ ORAL TABLET FOR SUSPENSION 3 MG	T5	PA-NS; QL (93 EA per 31 days)
AKEEGA	T5	PA-NS; QL (62 EA per 31 days)
ALECENSA	T5	PA-NS; QL (248 EA per 31 days)
ALUNBRIG ORAL TABLET 180 MG, 90 MG	T5	PA-NS; QL (31 EA per 31 days)
ALUNBRIG ORAL TABLET 30 MG	T5	PA-NS; QL (186 EA per 31 days)
ALUNBRIG ORAL TABLETS,DOSE PACK	T5	PA-NS; QL (60 EA per 365 days)
<i>anastrozole</i>	T2	
ARIMIDEX	T5	
AROMASIN	T5	
ASTAGRAF XL ORAL CAPSULE,EXTENDED RELEASE 24HR 0.5 MG, 1 MG	T4	PA-BvD
ASTAGRAF XL ORAL CAPSULE,EXTENDED RELEASE 24HR 5 MG	T5	PA-BvD
AUGTYRO	T5	PA-NS; QL (248 EA per 31 days)
AYVAKIT	T5	PA-NS; QL (31 EA per 31 days)
AZASAN	T4	PA-BvD
<i>azathioprine oral tablet 100 mg, 75 mg</i>	T4	PA-BvD
<i>azathioprine oral tablet 50 mg</i>	T2	PA-BvD
BALVERSA	T5	PA-NS
<i>bexarotene oral</i>	T5	PA-NS
<i>bexarotene topical</i>	T5	PA-NS; QL (60 GM per 28 days)
<i>bicalutamide</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
BOSULIF ORAL CAPSULE 100 MG	T5	PA-NS; QL (186 EA per 31 days)
BOSULIF ORAL CAPSULE 50 MG	T5	PA-NS; QL (341 EA per 31 days)
BOSULIF ORAL TABLET 100 MG	T5	PA-NS; QL (93 EA per 31 days)
BOSULIF ORAL TABLET 400 MG, 500 MG	T5	PA-NS; QL (31 EA per 31 days)
BRAFTOVI	T5	PA-NS; QL (186 EA per 31 days)
BRUKINSA	T5	PA-NS; QL (124 EA per 31 days)
CABOMETYX	T5	PA-NS; QL (31 EA per 31 days)
CALQUENCE	T5	PA-NS; QL (62 EA per 31 days)
CALQUENCE (ACALABRUTINIB MAL)	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 300 MG	T5	PA-NS; QL (31 EA per 31 days)
CASODEX	T4	
CELLCEPT	T5	PA-BvD
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
COMETRIQ ORAL CAPSULE 140 MG/DAY(80 MG X1-20 MG X3)	T5	PA-NS; QL (112 EA per 28 days)
COMETRIQ ORAL CAPSULE 60 MG/DAY (20 MG X 3/DAY)	T5	PA-NS; QL (84 EA per 28 days)
COPIKTRA	T5	PA-NS; QL (62 EA per 31 days)
COTELLIC	T5	PA-NS; LA; QL (63 EA per 28 days)
<i>cyclophosphamide oral</i>	T3	PA-BvD
<i>cyclosporine modified oral capsule</i>	T2	PA-BvD
<i>cyclosporine modified oral solution</i>	T4	PA-BvD
<i>cyclosporine oral capsule</i>	T2	PA-BvD
DAURISMO ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
DAURISMO ORAL TABLET 25 MG	T5	PA-NS; QL (62 EA per 31 days)
DROXIA	T4	
ELIGARD	T4	
ELIGARD (3 MONTH)	T4	
ELIGARD (4 MONTH)	T4	
ELIGARD (6 MONTH)	T4	
ENSPRYNG	T5	PA; QL (1 ML per 28 days)
ENVARBUS XR	T4	PA-BvD
ERIVEDGE	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 240 MG	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 60 MG	T5	PA-NS; QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>erlotinib</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 7.5 mg</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet 5 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 2 mg, 5 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 3 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>everolimus (immunosuppressive)</i>	T5	PA-BvD
<i>exemestane</i>	T4	
FARESTON	T5	
FEMARA	T4	
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 120 MG	T5	
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 80 MG	T4	
FOTIVDA	T5	PA-NS; QL (21 EA per 28 days)
FRUZAQLA ORAL CAPSULE 1 MG	T5	PA-NS; QL (84 EA per 28 days)
FRUZAQLA ORAL CAPSULE 5 MG	T5	PA-NS; QL (21 EA per 28 days)
GAVRETO	T5	PA-NS; QL (124 EA per 31 days)
<i>gefitinib</i>	T5	PA-NS; QL (31 EA per 31 days)
GENGRAF	T2	PA-BvD
GILOTRIF	T5	PA-NS; QL (31 EA per 31 days)
GLEEVEC ORAL TABLET 100 MG	T5	PA-NS; QL (93 EA per 31 days)
GLEEVEC ORAL TABLET 400 MG	T5	PA-NS; QL (62 EA per 31 days)
GLEOSTINE ORAL CAPSULE 10 MG, 40 MG	T4	PA-NS
GLEOSTINE ORAL CAPSULE 100 MG	T5	PA-NS
HYDREA	T4	
<i>hydroxyurea</i>	T2	
IBRANCE	T5	PA-NS; QL (21 EA per 28 days)
ICLUSIG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA ORAL TABLET 50 MG	T5	PA-NS; QL (62 EA per 31 days)
<i>imatinib oral tablet 100 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>imatinib oral tablet 400 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
IMBRUVICA ORAL CAPSULE 140 MG	T5	PA-NS; QL (124 EA per 31 days)
IMBRUVICA ORAL CAPSULE 70 MG	T5	PA-NS; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
IMBRUVICA ORAL SUSPENSION	T5	PA-NS; QL (216 ML per 25 days)
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG	T5	PA-NS; QL (31 EA per 31 days)
IMURAN	T4	PA-BvD
INLYTA	T5	PA-NS; QL (124 EA per 31 days)
INQOVI	T5	PA-NS; QL (5 EA per 28 days)
INREBIC	T5	PA-NS; QL (124 EA per 31 days)
IRESSA	T5	PA-NS; QL (31 EA per 31 days)
IWILFIN	T5	PA-NS; QL (248 EA per 31 days)
JAKAFI	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
JYLAMVO	T4	PA-BvD
KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG	T5	PA-NS; QL (49 EA per 28 days)
KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG	T5	PA-NS; QL (70 EA per 28 days)
KISQALI FEMARA CO-PACK ORAL TABLET 600 MG/DAY(200 MG X 3)-2.5 MG	T5	PA-NS; QL (91 EA per 28 days)
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (21 EA per 28 days)
KISQALI ORAL TABLET 400 MG/DAY (200 MG X 2)	T5	PA-NS; QL (42 EA per 28 days)
KISQALI ORAL TABLET 600 MG/DAY (200 MG X 3)	T5	PA-NS; QL (63 EA per 28 days)
KLISYRI	T5	PA
KOSELUGO ORAL CAPSULE 10 MG	T5	PA; QL (279 EA per 31 days)
KOSELUGO ORAL CAPSULE 25 MG	T5	PA; QL (124 EA per 31 days)
KRAZATI	T5	PA-NS; QL (186 EA per 31 days)
<i>lapatinib</i>	T5	PA-NS; QL (186 EA per 31 days)
<i>lenalidomide</i>	T5	PA-NS; QL (21 EA per 28 days)
LENVIMA	T5	PA-NS
<i>letrozole</i>	T2	
<i>leucovorin calcium oral</i>	T3	
LEUKERAN	T5	
<i>leuprolide (3 month)</i>	T4	ST
<i>leuprolide subcutaneous kit</i>	T3	
LONSURF	T5	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
LORBRENA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
LORBRENA ORAL TABLET 25 MG	T5	PA-NS; QL (93 EA per 31 days)
LUMAKRAS ORAL TABLET 120 MG	T5	PA-NS; QL (124 EA per 31 days)
LUMAKRAS ORAL TABLET 320 MG	T5	PA-NS; QL (93 EA per 31 days)
LUPKYNIS	T5	PA; QL (186 EA per 31 days)
LUPRON DEPOT	T5	ST
LUPRON DEPOT (3 MONTH)	T5	ST
LUPRON DEPOT (4 MONTH)	T5	ST
LUPRON DEPOT (6 MONTH)	T5	ST
LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	T5	PA
LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)	T5	PA
LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT	T5	PA
LYNPARZA	T5	PA-NS; QL (124 EA per 31 days)
LYSODREN	T5	
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3)	T5	PA-NS; QL (93 EA per 31 days)
LYTGOBI ORAL TABLET 16 MG/DAY (4 MG X 4)	T5	PA-NS; QL (124 EA per 31 days)
LYTGOBI ORAL TABLET 20 MG/DAY (4 MG X 5)	T5	PA-NS; QL (155 EA per 31 days)
MATULANE	T5	
<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml)</i>	T3	PA
<i>megestrol oral suspension 625 mg/5 ml (125 mg/ml)</i>	T4	PA
<i>megestrol oral tablet</i>	T3	PA-NS
MEKINIST ORAL RECON SOLN	T5	PA-NS; QL (1260 ML per 31 days)
MEKINIST ORAL TABLET 0.5 MG	T5	PA-NS; QL (93 EA per 31 days)
MEKINIST ORAL TABLET 2 MG	T5	PA-NS; QL (31 EA per 31 days)
MEKTOVI	T5	PA-NS; QL (186 EA per 31 days)
<i>mercaptopurine</i>	T2	
MESNEX ORAL	T4	
<i>methotrexate sodium</i>	T2	PA-BvD
<i>methotrexate sodium (pf) injection solution</i>	T2	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
MYCAPSSA	T5	PA; QL (124 EA per 31 days)
<i>mycophenolate mofetil oral capsule</i>	T3	PA-BvD
<i>mycophenolate mofetil oral suspension for reconstitution</i>	T5	PA-BvD
<i>mycophenolate mofetil oral tablet</i>	T3	PA-BvD
<i>mycophenolate sodium</i>	T4	PA-BvD
MYFORTIC ORAL TABLET,DELAYED RELEASE (DR/EC) 180 MG	T4	PA-BvD
MYFORTIC ORAL TABLET,DELAYED RELEASE (DR/EC) 360 MG	T5	PA-BvD
MYHIBBIN	T5	PA-BvD
NEORAL	T4	PA-BvD
NERLYNX	T5	PA-NS; QL (186 EA per 31 days)
NEXAVAR	T5	PA-NS; QL (124 EA per 31 days)
NILANDRON	T5	
<i>nilutamide</i>	T5	
NINLARO	T5	PA-NS; QL (3 EA per 28 days)
NUBEQA	T5	PA-NS; QL (124 EA per 31 days)
<i>octreotide acetate injection solution 1,000 mcg/ml, 100 mcg/ml, 200 mcg/ml, 50 mcg/ml</i>	T4	PA
<i>octreotide acetate injection solution 500 mcg/ml</i>	T5	PA
ODOMZO	T5	PA-NS; LA; QL (31 EA per 31 days)
OGSIVEO ORAL TABLET 100 MG, 150 MG	T5	PA-NS; QL (62 EA per 31 days)
OGSIVEO ORAL TABLET 50 MG	T5	PA-NS; QL (186 EA per 31 days)
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	T5	PA-NS; QL (96 ML per 28 days)
OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)	T5	PA-NS; QL (20 EA per 28 days)
OJJAARA	T5	PA-NS; QL (31 EA per 31 days)
ONUREG	T5	PA-NS; QL (14 EA per 28 days)
ORGOVYX	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 345 MG	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 86 MG	T5	PA-NS; QL (93 EA per 31 days)
<i>pazopanib</i>	T5	PA-NS; QL (124 EA per 31 days)
PEMAZYRE	T5	PA-NS; QL (14 EA per 21 days)
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (28 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
PIQRAY ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	T5	PA-NS; QL (56 EA per 28 days)
POMALYST	T5	PA-NS; QL (21 EA per 28 days)
PROGRAF ORAL CAPSULE 0.5 MG, 1 MG	T4	PA-BvD
PROGRAF ORAL CAPSULE 5 MG	T5	PA-BvD
PROGRAF ORAL GRANULES IN PACKET	T4	PA-BvD
PURIXAN	T5	
QINLOCK	T5	PA-NS; QL (93 EA per 31 days)
RAPAMUNE ORAL SOLUTION	T5	PA-BvD
RAPAMUNE ORAL TABLET 0.5 MG	T4	PA-BvD
RAPAMUNE ORAL TABLET 1 MG, 2 MG	T5	PA-BvD
RETEVMO ORAL CAPSULE 40 MG	T5	PA-NS; QL (186 EA per 31 days)
RETEVMO ORAL CAPSULE 80 MG	T5	PA-NS; QL (124 EA per 31 days)
REVLIMID	T5	PA-NS; QL (21 EA per 28 days)
REZLIDHIA	T5	PA-NS; QL (62 EA per 31 days)
REZUROCK	T5	PA; QL (62 EA per 31 days)
ROZLYTREK ORAL CAPSULE 100 MG	T5	PA-NS; QL (155 EA per 31 days)
ROZLYTREK ORAL CAPSULE 200 MG	T5	PA-NS; QL (93 EA per 31 days)
ROZLYTREK ORAL PELLETS IN PACKET	T5	PA-NS; QL (372 EA per 31 days)
RUBRACA	T5	PA-NS; QL (124 EA per 31 days)
RYDAPT	T5	PA-NS; QL (248 EA per 31 days)
SANDIMMUNE ORAL	T4	PA-BvD
SANDOSTATIN INJECTION SOLUTION 100 MCG/ML	T5	PA
SANDOSTATIN INJECTION SOLUTION 50 MCG/ML, 500 MCG/ML	T4	PA
SCEMBLIX ORAL TABLET 100 MG	T5	PA-NS; QL (124 EA per 31 days)
SCEMBLIX ORAL TABLET 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SCEMBLIX ORAL TABLET 40 MG	T5	PA-NS; QL (310 EA per 31 days)
SIGNIFOR	T5	PA
SIKLOS ORAL TABLET 1,000 MG	T5	
SIKLOS ORAL TABLET 100 MG	T4	
<i>sirolimus oral solution</i>	T5	PA-BvD
<i>sirolimus oral tablet</i>	T4	PA-BvD
SOLTAMOX	T5	
<i>sorafenib</i>	T5	PA-NS; QL (124 EA per 31 days)
SPRYCEL	T5	PA-NS; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
STIVARGA	T5	PA-NS; QL (84 EA per 28 days)
<i>sunitinib malate</i>	T5	PA-NS; QL (31 EA per 31 days)
SUTENT	T5	PA-NS; QL (31 EA per 31 days)
TABLOID	T4	
TABRECTA	T5	PA-NS; QL (124 EA per 31 days)
<i>tacrolimus oral capsule</i>	T2	PA-BvD
TAFINLAR ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
TAFINLAR ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (930 EA per 31 days)
TAGRISSE	T5	PA-NS; LA; QL (31 EA per 31 days)
TALZENNA	T5	PA-NS; QL (31 EA per 31 days)
<i>tamoxifen</i>	T2	
TARGRETIN ORAL	T5	PA-NS
TARGRETIN TOPICAL	T5	PA-NS; QL (60 GM per 28 days)
TASIGNA	T5	PA-NS; QL (124 EA per 31 days)
TAZVERIK	T5	PA-NS; QL (248 EA per 31 days)
TEPMETKO	T5	PA-NS; QL (62 EA per 31 days)
THALOMID ORAL CAPSULE 100 MG, 150 MG, 50 MG	T5	PA-NS; QL (28 EA per 28 days)
THALOMID ORAL CAPSULE 200 MG	T5	PA-NS; QL (56 EA per 28 days)
TIBSOVO	T5	PA-NS; QL (62 EA per 31 days)
<i>toremifene</i>	T4	
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION	T4	ST
<i>tretinoin (antineoplastic)</i>	T5	
TREXALL	T4	PA-BvD
TRUQAP	T5	PA-NS; QL (64 EA per 28 days)
TUKYSA ORAL TABLET 150 MG	T5	PA-NS; QL (124 EA per 31 days)
TUKYSA ORAL TABLET 50 MG	T5	PA-NS; QL (248 EA per 31 days)
TURALIO ORAL CAPSULE 125 MG	T5	PA-NS; QL (124 EA per 31 days)
TYKERB	T5	PA-NS; QL (186 EA per 31 days)
VANFLYTA	T5	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 10 MG	T3	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 100 MG	T5	PA-NS; QL (186 EA per 31 days)
VENCLEXTA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
VENCLEXTA STARTING PACK	T5	PA-NS; QL (84 EA per 365 days)
VERZENIO	T5	PA-NS; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
VIJOICE ORAL GRANULES IN PACKET	T5	PA-NS; QL (31 EA per 31 days)
VIJOICE ORAL TABLET 125 MG, 50 MG	T5	PA-NS; QL (28 EA per 28 days)
VIJOICE ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
VITRAKVI ORAL CAPSULE 100 MG	T5	PA-NS; QL (62 EA per 31 days)
VITRAKVI ORAL CAPSULE 25 MG	T5	PA-NS; QL (186 EA per 31 days)
VITRAKVI ORAL SOLUTION	T5	PA-NS; QL (310 ML per 31 days)
VIZIMPRO	T5	PA-NS; QL (31 EA per 31 days)
VONJO	T5	PA-NS; QL (124 EA per 31 days)
VOTRIENT	T5	PA-NS; QL (124 EA per 31 days)
WELIREG	T5	PA-NS; QL (93 EA per 31 days)
XALKORI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XALKORI ORAL PELLETT 150 MG	T5	PA-NS; QL (186 EA per 31 days)
XALKORI ORAL PELLETT 20 MG, 50 MG	T5	PA-NS; QL (124 EA per 31 days)
XATMEP	T4	PA-BvD
XERMELO	T5	PA; QL (93 EA per 31 days)
XGEVA	T5	PA-NS
XOSPATA	T5	PA-NS; QL (124 EA per 31 days)
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40MG TWICE WEEK (40 MG X 2), 80 MG/WEEK (40 MG X 2)	T5	PA-NS; QL (8 EA per 28 days)
XPOVIO ORAL TABLET 40 MG/WEEK (40 MG X 1), 60 MG/WEEK (60 MG X 1)	T5	PA-NS; QL (4 EA per 28 days)
XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK)	T5	PA-NS; QL (24 EA per 28 days)
XPOVIO ORAL TABLET 80MG TWICE WEEK (160 MG/WEEK)	T5	PA-NS; QL (32 EA per 28 days)
XTANDI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 40 MG	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 80 MG	T5	PA-NS; QL (62 EA per 31 days)
YONSA	T5	PA-NS; QL (124 EA per 31 days)
ZEJULA ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
ZELBORAF	T5	PA-NS; QL (248 EA per 31 days)
ZOLINZA	T5	PA-NS
ZORTRESS	T5	PA-BvD
ZYDELIG	T5	PA-NS; QL (62 EA per 31 days)
ZYKADIA	T5	PA-NS; QL (93 EA per 31 days)
ZYTIGA ORAL TABLET 250 MG	T5	PA-NS; QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZYTIGA ORAL TABLET 500 MG	T5	PA-NS; QL (62 EA per 31 days)
Autonomic / Cns Drugs, Neurology / Psych		
ABILIFY ASIMTUFII INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 720 MG/2.4 ML	T5	QL (2.4 ML per 56 days)
ABILIFY ASIMTUFII INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 960 MG/3.2 ML	T5	QL (3.2 ML per 56 days)
ABILIFY MAINTENA	T5	QL (1 EA per 28 days)
ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET WITH SENSOR AND STRIP 15 MG, 2 MG, 20 MG, 30 MG, 5 MG	T5	PA-NS
ABILIFY MYCITE STARTER KIT ORAL TABLET WITH SENSOR, STRIP, POD 10 MG	T5	PA-NS
ABILIFY ORAL TABLET	T4	PA-NS
<i>acetaminophen-caff-dihydrocod</i>	T2	PA; QL (372 EA per 31 days)
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i>	T2	PA; QL (5167 ML per 31 days)
<i>acetaminophen-codeine oral tablet</i>	T2	PA; QL (403 EA per 31 days)
ADDERALL ORAL TABLET 20 MG	T4	ST; QL (93 EA per 31 days)
ADDERALL ORAL TABLET 5 MG, 7.5 MG	T4	ST; QL (62 EA per 31 days)
ADDERALL XR	T4	ST; QL (31 EA per 31 days)
ADLARITY	T4	PA; QL (4 EA per 28 days)
ADZENYS XR-ODT	T4	ST; QL (31 EA per 31 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML	T3	PA; QL (1 ML per 28 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 70 MG/ML	T3	PA; QL (2 ML per 28 days)
AJOVY AUTOINJECTOR	T3	PA; QL (1.5 ML per 28 days)
AJOVY SYRINGE	T3	PA; QL (1.5 ML per 28 days)
ALLZITAL	T4	QL (372 EA per 31 days)
<i>almotriptan malate oral tablet 12.5 mg</i>	T4	QL (8 EA per 28 days)
<i>almotriptan malate oral tablet 6.25 mg</i>	T4	QL (16 EA per 28 days)
ALPRAZOLAM INTENSOL	T4	PA
<i>alprazolam oral tablet 0.25 mg, 0.5 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>alprazolam oral tablet 1 mg, 2 mg</i>	T4	PA; QL (155 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg</i>	T4	PA; QL (31 EA per 31 days)
<i>alprazolam oral tablet extended release 24 hr 2 mg</i>	T4	PA; QL (155 EA per 31 days)
<i>alprazolam oral tablet extended release 24 hr 3 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>alprazolam oral tablet,disintegrating 0.25 mg, 0.5 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>alprazolam oral tablet,disintegrating 1 mg, 2 mg</i>	T4	PA; QL (155 EA per 31 days)
AMBIEN	T4	PA; QL (31 EA per 31 days)
AMBIEN CR	T4	PA; QL (31 EA per 31 days)
<i>amitriptyline</i>	T2	PA-NS
<i>amitriptyline-chlordiazepoxide</i>	T4	PA-NS
<i>amoxapine</i>	T3	
<i>amphetamine sulfate</i>	T4	PA
AMPYRA	T5	PA; QL (62 EA per 31 days)
AMRIX	T5	QL (31 EA per 31 days)
ANAFRANIL	T4	PA-NS
ALENZIN	T5	QL (31 EA per 31 days)
APOKYN	T5	PA; QL (60 ML per 30 days)
<i>apomorphine</i>	T5	PA; QL (60 ML per 30 days)
APTENSIO XR	T4	ST; QL (31 EA per 31 days)
APTIOM ORAL TABLET 200 MG	T5	QL (186 EA per 31 days)
APTIOM ORAL TABLET 400 MG	T5	QL (93 EA per 31 days)
APTIOM ORAL TABLET 600 MG, 800 MG	T5	QL (62 EA per 31 days)
ARICEPT	T4	
<i>aripiprazole oral solution</i>	T4	PA-NS
<i>aripiprazole oral tablet</i>	T2	PA-NS
<i>aripiprazole oral tablet,disintegrating</i>	T4	PA-NS
ARISTADA INITIO	T5	QL (4.8 ML per 365 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 1,064 MG/3.9 ML	T5	QL (3.9 ML per 28 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 441 MG/1.6 ML	T5	QL (1.6 ML per 28 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 662 MG/2.4 ML	T5	QL (2.4 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 882 MG/3.2 ML	T5	QL (3.2 ML per 28 days)
<i>armodafinil</i>	T4	PA; QL (31 EA per 31 days)
ARTHROTEC 50	T4	
ARTHROTEC 75	T4	
ASCOMP WITH CODEINE	T4	PA; QL (372 EA per 31 days)
<i>asenapine maleate</i>	T4	PA-NS; QL (62 EA per 31 days)
ATIVAN ORAL TABLET 0.5 MG	T5	PA; QL (124 EA per 31 days)
ATIVAN ORAL TABLET 1 MG	T5	PA; QL (186 EA per 31 days)
ATIVAN ORAL TABLET 2 MG	T5	PA; QL (155 EA per 31 days)
<i>atomoxetine oral capsule 10 mg, 25 mg, 40 mg</i>	T4	QL (62 EA per 31 days)
<i>atomoxetine oral capsule 100 mg, 60 mg, 80 mg</i>	T4	QL (31 EA per 31 days)
<i>atomoxetine oral capsule 18 mg</i>	T4	QL (124 EA per 31 days)
AUBAGIO	T5	PA; QL (31 EA per 31 days)
AUSTEDO ORAL TABLET 12 MG, 6 MG	T5	PA; QL (124 EA per 31 days)
AUSTEDO ORAL TABLET 9 MG	T5	PA; QL (155 EA per 31 days)
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG	T5	PA; QL (93 EA per 31 days)
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 24 MG	T5	PA; QL (62 EA per 31 days)
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 30 MG, 36 MG, 42 MG, 48 MG	T5	PA; QL (31 EA per 31 days)
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 6 MG	T5	PA; QL (217 EA per 31 days)
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 6 MG (14)-12 MG (14)-24 MG (14)	T5	PA; QL (84 EA per 365 days)
AUVELITY	T4	PA-NS; QL (62 EA per 31 days)
AZILECT	T4	
AZSTARYS	T4	ST; QL (31 EA per 31 days)
<i>baclofen oral solution 10 mg/5 ml (2 mg/ml)</i>	T4	PA; QL (1240 ML per 31 days)
<i>baclofen oral suspension</i>	T4	PA; QL (496 ML per 31 days)
<i>baclofen oral tablet 10 mg, 20 mg, 5 mg</i>	T2	
<i>baclofen oral tablet 15 mg</i>	T4	
BAFIERTAM	T5	PA; QL (124 EA per 31 days)
BANZEL	T5	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
BELBUCA BUCCAL FILM 150 MCG, 300 MCG, 450 MCG, 600 MCG, 75 MCG	T3	PA; QL (62 EA per 31 days)
BELBUCA BUCCAL FILM 750 MCG, 900 MCG	T4	PA; QL (62 EA per 31 days)
BELSOMRA	T4	ST; QL (31 EA per 31 days)
<i>benztropine oral</i>	T1	PA
BRIVIACT ORAL SOLUTION	T5	QL (620 ML per 31 days)
BRIVIACT ORAL TABLET 10 MG, 100 MG, 25 MG, 75 MG	T5	QL (62 EA per 31 days)
BRIVIACT ORAL TABLET 50 MG	T4	QL (62 EA per 31 days)
<i>bromocriptine</i>	T4	
BUPAP	T4	QL (403 EA per 31 days)
<i>buprenorphine</i>	T4	PA; QL (4 EA per 28 days)
<i>buprenorphine hcl sublingual tablet 2 mg</i>	T2	QL (93 EA per 31 days)
<i>buprenorphine hcl sublingual tablet 8 mg</i>	T2	QL (62 EA per 31 days)
<i>buprenorphine-naloxone sublingual film 12-3 mg, 4-1 mg, 8-2 mg</i>	T3	QL (62 EA per 31 days)
<i>buprenorphine-naloxone sublingual film 2-0.5 mg</i>	T3	QL (93 EA per 31 days)
<i>buprenorphine-naloxone sublingual tablet</i>	T2	QL (93 EA per 31 days)
<i>bupropion hcl oral tablet</i>	T1	
<i>bupropion hcl oral tablet extended release 24 hr 150 mg</i>	T2	QL (93 EA per 31 days)
<i>bupropion hcl oral tablet extended release 24 hr 300 mg</i>	T2	QL (31 EA per 31 days)
<i>bupropion hcl oral tablet extended release 24 hr 450 mg</i>	T4	QL (31 EA per 31 days)
<i>bupropion hcl oral tablet sustained-release 12 hr</i>	T2	QL (62 EA per 31 days)
<i>buspirone</i>	T2	
<i>butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg</i>	T4	PA; QL (403 EA per 31 days)
<i>butalbital-acetaminop-caf-cod oral capsule 50-325-40-30 mg</i>	T4	PA; QL (372 EA per 31 days)
<i>butalbital-acetaminophen oral capsule</i>	T4	QL (403 EA per 31 days)
<i>butalbital-acetaminophen oral tablet 50-300 mg</i>	T4	QL (403 EA per 31 days)
<i>butalbital-acetaminophen oral tablet 50-325 mg</i>	T4	QL (372 EA per 31 days)
<i>butalbital-acetaminophen-caff oral capsule 50-300-40 mg</i>	T4	QL (403 EA per 31 days)
<i>butalbital-acetaminophen-caff oral capsule 50-325-40 mg</i>	T4	QL (372 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>butalbital-acetaminophen-caff oral tablet</i>	T4	QL (372 EA per 31 days)
<i>butalbital-aspirin-caffeine oral capsule</i>	T4	
<i>butorphanol nasal</i>	T2	QL (5 ML per 28 days)
BUTRANS	T4	PA; QL (4 EA per 28 days)
CAMBIA	T4	
CAPLYTA	T5	PA-NS; QL (31 EA per 31 days)
<i>carbamazepine oral capsule, er multiphase 12 hr</i>	T2	
<i>carbamazepine oral suspension 100 mg/5 ml</i>	T2	
<i>carbamazepine oral tablet</i>	T2	
<i>carbamazepine oral tablet extended release 12 hr</i>	T2	
<i>carbamazepine oral tablet,chewable</i>	T2	
CARBATROL	T4	
<i>carbidopa</i>	T2	
<i>carbidopa-levodopa</i>	T2	
<i>carbidopa-levodopa-entacapone</i>	T4	
<i>carisoprodol</i>	T4	PA
CELEBREX	T4	ST; QL (62 EA per 31 days)
<i>celecoxib</i>	T2	ST; QL (62 EA per 31 days)
CELEXA ORAL TABLET	T4	
CELONTIN ORAL CAPSULE 300 MG	T4	
<i>chlordiazepoxide hcl</i>	T4	PA
<i>chlorpromazine oral</i>	T4	
<i>chlorzoxazone oral tablet 250 mg</i>	T5	PA
<i>chlorzoxazone oral tablet 375 mg, 500 mg, 750 mg</i>	T4	PA
<i>citalopram oral capsule</i>	T4	PA-NS; QL (31 EA per 31 days)
<i>citalopram oral solution</i>	T3	
<i>citalopram oral tablet</i>	T1	
<i>clobazam oral suspension</i>	T4	PA-NS; QL (496 ML per 31 days)
<i>clobazam oral tablet</i>	T3	PA-NS; QL (62 EA per 31 days)
<i>clomipramine</i>	T4	PA-NS
<i>clonazepam oral tablet 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>clonazepam oral tablet 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet,disintegrating 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>clonazepam oral tablet,disintegrating 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clonidine hcl oral tablet extended release 12 hr</i>	T4	PA
<i>clorazepate dipotassium oral tablet 15 mg</i>	T2	PA-NS; QL (186 EA per 31 days)
<i>clorazepate dipotassium oral tablet 3.75 mg, 7.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clozapine oral tablet 100 mg, 25 mg</i>	T3	QL (279 EA per 31 days)
<i>clozapine oral tablet 200 mg</i>	T3	QL (124 EA per 31 days)
<i>clozapine oral tablet 50 mg</i>	T3	QL (93 EA per 31 days)
<i>clozapine oral tablet,disintegrating 100 mg, 25 mg</i>	T4	QL (279 EA per 31 days)
<i>clozapine oral tablet,disintegrating 12.5 mg</i>	T4	QL (93 EA per 31 days)
<i>clozapine oral tablet,disintegrating 150 mg</i>	T4	QL (186 EA per 31 days)
<i>clozapine oral tablet,disintegrating 200 mg</i>	T4	QL (124 EA per 31 days)
CLOZARIL ORAL TABLET 100 MG	T5	QL (279 EA per 31 days)
CLOZARIL ORAL TABLET 200 MG	T4	QL (124 EA per 31 days)
CLOZARIL ORAL TABLET 25 MG	T4	QL (279 EA per 31 days)
CLOZARIL ORAL TABLET 50 MG	T4	QL (93 EA per 31 days)
<i>codeine sulfate</i>	T4	PA; QL (186 EA per 31 days)
<i>codeine-butalbital-asa-caff</i>	T4	PA; QL (372 EA per 31 days)
COMTAN	T4	
CONCERTA	T4	ST; QL (31 EA per 31 days)
CONZIP	T4	PA; QL (30 EA per 30 days)
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
COPAXONE SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
COTEMPLA XR-ODT	T4	ST; QL (62 EA per 31 days)
<i>cyclobenzaprine oral capsule,extended release 24hr</i>	T4	QL (31 EA per 31 days)
<i>cyclobenzaprine oral tablet 10 mg, 7.5 mg</i>	T4	QL (93 EA per 31 days)
<i>cyclobenzaprine oral tablet 5 mg</i>	T4	QL (155 EA per 31 days)
CYMBALTA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 20 MG, 60 MG	T4	QL (62 EA per 31 days)
CYMBALTA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 30 MG	T4	QL (31 EA per 31 days)
<i>dalfampridine</i>	T5	PA; QL (62 EA per 31 days)
DANTRIUM ORAL CAPSULE 25 MG	T4	
<i>dantrolene oral</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
DAYBUE	T5	PA; QL (3600 ML per 30 days)
DAYPRO	T4	
DAYTRANA	T4	PA; QL (30 EA per 30 days)
DAYVIGO	T4	ST; QL (31 EA per 31 days)
DEMEROL (PF) INJECTION SYRINGE 25 MG/ML	T4	PA; QL (824 ML per 31 days)
DEMEROL INJECTION	T4	PA; QL (412 ML per 31 days)
DEPAKOTE	T4	
DEPAKOTE ER	T4	
DEPAKOTE SPRINKLES	T4	
<i>desipramine</i>	T2	
<i>desvenlafaxine oral tablet extended release 24 hr 100 mg</i>	T4	QL (124 EA per 31 days)
<i>desvenlafaxine oral tablet extended release 24 hr 50 mg</i>	T4	QL (31 EA per 31 days)
<i>desvenlafaxine succinate</i>	T2	QL (31 EA per 31 days)
DEXEDRINE SPANSULE ORAL CAPSULE, EXTENDED RELEASE 10 MG	T4	ST; QL (155 EA per 31 days)
<i>dexmethylphenidate oral capsule,er biphasic 50-50</i>	T2	QL (31 EA per 31 days)
<i>dexmethylphenidate oral tablet 10 mg</i>	T2	QL (62 EA per 31 days)
<i>dexmethylphenidate oral tablet 2.5 mg, 5 mg</i>	T2	QL (93 EA per 31 days)
<i>dextroamphetamine sulfate oral capsule, extended release 10 mg</i>	T4	QL (155 EA per 31 days)
<i>dextroamphetamine sulfate oral capsule, extended release 15 mg</i>	T4	QL (124 EA per 31 days)
<i>dextroamphetamine sulfate oral capsule, extended release 5 mg</i>	T4	QL (186 EA per 31 days)
<i>dextroamphetamine sulfate oral solution</i>	T4	
<i>dextroamphetamine sulfate oral tablet 10 mg</i>	T4	QL (186 EA per 31 days)
<i>dextroamphetamine sulfate oral tablet 15 mg, 20 mg, 30 mg</i>	T4	QL (62 EA per 31 days)
<i>dextroamphetamine sulfate oral tablet 5 mg</i>	T4	QL (341 EA per 31 days)
<i>dextroamphetamine-amphetamine oral capsule, er triphasic 24 hr</i>	T4	ST; QL (31 EA per 31 days)
<i>dextroamphetamine-amphetamine oral capsule,extended release 24hr</i>	T4	QL (31 EA per 31 days)
<i>dextroamphetamine-amphetamine oral tablet 10 mg, 12.5 mg, 15 mg, 30 mg, 5 mg, 7.5 mg</i>	T3	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>dextroamphetamine-amphetamine oral tablet 20 mg</i>	T3	QL (93 EA per 31 days)
DHIVY	T4	ST
DIACOMIT ORAL CAPSULE 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL CAPSULE 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIAZEPAM INTENSOL	T2	PA-NS; QL (248 ML per 31 days)
<i>diazepam oral solution 5 mg/5 ml (1 mg/ml)</i>	T2	PA-NS; QL (1500 ML per 31 days)
<i>diazepam oral tablet</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>diazepam rectal</i>	T4	
<i>diclofenac epolamine</i>	T4	PA; QL (62 EA per 31 days)
<i>diclofenac potassium oral capsule</i>	T4	
<i>diclofenac potassium oral powder in packet</i>	T4	
<i>diclofenac potassium oral tablet 25 mg</i>	T5	
<i>diclofenac potassium oral tablet 50 mg</i>	T2	
<i>diclofenac sodium oral</i>	T2	
<i>diclofenac sodium topical drops</i>	T4	QL (450 ML per 28 days)
<i>diclofenac sodium topical solution in metered-dose pump</i>	T5	ST; QL (224 GM per 28 days)
<i>diclofenac-misoprostol</i>	T4	
<i>diflunisal</i>	T2	
<i>dihydroergotamine nasal</i>	T4	PA; QL (8 ML per 31 days)
DILANTIN	T3	
DILANTIN EXTENDED	T4	
DILANTIN INFATABS	T4	
DILANTIN-125	T4	
DILAUDID ORAL LIQUID	T4	PA; QL (1550 ML per 31 days)
DILAUDID ORAL TABLET 2 MG	T4	PA; QL (186 EA per 31 days)
DILAUDID ORAL TABLET 4 MG, 8 MG	T5	PA; QL (186 EA per 31 days)
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg (14)- 240 mg (46)</i>	T5	PA; QL (120 EA per 365 days)
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 240 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>divalproex</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>donepezil oral tablet 10 mg, 5 mg</i>	T1	
<i>donepezil oral tablet 23 mg</i>	T4	QL (31 EA per 31 days)
<i>donepezil oral tablet, disintegrating</i>	T2	
<i>doxepin oral capsule</i>	T2	PA-NS
<i>doxepin oral concentrate</i>	T2	PA-NS
<i>doxepin oral tablet</i>	T3	PA
<i>duloxetine oral capsule, delayed release(dr/ec) 20 mg, 60 mg</i>	T2	QL (62 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 30 mg</i>	T2	QL (31 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 40 mg</i>	T4	QL (31 EA per 31 days)
DUOPA	T5	PA-BvD
DYANAVEL XR ORAL SUSPEN, IR - ER, BIPHASIC 24HR	T4	ST; QL (248 ML per 31 days)
DYANAVEL XR ORAL TABLET, IR - ER, BIPHASIC 24HR	T4	ST; QL (31 EA per 31 days)
EDLUAR	T4	PA; QL (31 EA per 31 days)
EFFEXOR XR ORAL CAPSULE, EXTENDED RELEASE 24HR 150 MG, 37.5 MG	T4	QL (31 EA per 31 days)
EFFEXOR XR ORAL CAPSULE, EXTENDED RELEASE 24HR 75 MG	T4	QL (93 EA per 31 days)
<i>eletriptan oral tablet 20 mg</i>	T4	QL (12 EA per 28 days)
<i>eletriptan oral tablet 40 mg</i>	T4	QL (6 EA per 28 days)
ELYXYB	T4	PA
EMGALITY PEN	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 300 MG/3 ML (100 MG/ML X 3)	T5	PA; QL (3 ML per 28 days)
EMSAM	T5	QL (30 EA per 30 days)
ENDOCET	T3	PA; QL (372 EA per 31 days)
<i>entacapone</i>	T3	
EPIDIOLEX	T5	PA-NS
EPITOL	T2	
EPRONTIA	T4	PA-NS; QL (496 ML per 31 days)
EQUETRO	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>ergoloid</i>	T4	PA
<i>ergotamine-caffeine</i>	T3	PA
<i>escitalopram oxalate oral solution</i>	T2	QL (620 ML per 31 days)
<i>escitalopram oxalate oral tablet 10 mg</i>	T1	QL (45 EA per 30 days)
<i>escitalopram oxalate oral tablet 20 mg, 5 mg</i>	T1	QL (30 EA per 30 days)
ESGIC ORAL TABLET	T4	QL (372 EA per 31 days)
<i>estazolam</i>	T4	PA
<i>eszopiclone</i>	T4	PA; QL (31 EA per 31 days)
<i>ethosuximide</i>	T2	
<i>etodolac</i>	T2	
EVEKEO	T4	PA
EVRYSDI	T5	PA; QL (240 ML per 31 days)
EXELON PATCH	T4	QL (30 EA per 30 days)
FANAPT ORAL TABLET 1 MG	T4	QL (62 EA per 31 days)
FANAPT ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG	T5	QL (62 EA per 31 days)
FANAPT ORAL TABLETS,DOSE PACK	T4	QL (16 EA per 365 days)
<i>felbamate oral suspension</i>	T5	
<i>felbamate oral tablet</i>	T4	
FELBATOL ORAL TABLET	T5	
<i>fenoprofen oral capsule 400 mg</i>	T4	
<i>fenoprofen oral tablet</i>	T4	
<i>fentanyl citrate buccal lozenge on a handle 1,200 mcg</i>	T5	PA; QL (40 EA per 31 days)
<i>fentanyl citrate buccal lozenge on a handle 1,600 mcg</i>	T5	PA; QL (30 EA per 31 days)
<i>fentanyl citrate buccal lozenge on a handle 200 mcg</i>	T5	PA; QL (124 EA per 31 days)
<i>fentanyl citrate buccal lozenge on a handle 400 mcg</i>	T5	PA; QL (119 EA per 31 days)
<i>fentanyl citrate buccal lozenge on a handle 600 mcg</i>	T5	PA; QL (79 EA per 31 days)
<i>fentanyl citrate buccal lozenge on a handle 800 mcg</i>	T5	PA; QL (59 EA per 31 days)
<i>fentanyl citrate buccal tablet, effervescent 100 mcg, 200 mcg</i>	T5	PA; QL (124 EA per 31 days)
<i>fentanyl citrate buccal tablet, effervescent 400 mcg</i>	T5	PA; QL (119 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>fentanyl citrate buccal tablet, effervescent 600 mcg</i>	T5	PA; QL (79 EA per 31 days)
<i>fentanyl citrate buccal tablet, effervescent 800 mcg</i>	T5	PA; QL (59 EA per 31 days)
<i>fentanyl transdermal patch 72 hour 100 mcg/hr</i>	T4	PA; QL (10 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 12 mcg/hr, 25 mcg/hr, 37.5 mcg/hour</i>	T4	PA; QL (20 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 50 mcg/hr</i>	T4	PA; QL (17 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 62.5 mcg/hour</i>	T4	PA; QL (15 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 75 mcg/hr</i>	T4	PA; QL (12 EA per 30 days)
<i>fentanyl transdermal patch 72 hour 87.5 mcg/hour</i>	T5	PA; QL (11 EA per 30 days)
FENTORA BUCCAL TABLET, EFFERVESCENT 100 MCG, 200 MCG	T5	PA; QL (124 EA per 31 days)
FENTORA BUCCAL TABLET, EFFERVESCENT 400 MCG	T5	PA; QL (119 EA per 31 days)
FENTORA BUCCAL TABLET, EFFERVESCENT 600 MCG	T5	PA; QL (79 EA per 31 days)
FENTORA BUCCAL TABLET, EFFERVESCENT 800 MCG	T5	PA; QL (59 EA per 31 days)
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)	T3	PA-NS; QL (56 EA per 365 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 40 MG, 80 MG	T3	PA-NS; QL (31 EA per 31 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 20 MG	T3	PA-NS; QL (93 EA per 31 days)
FEXMID	T4	QL (124 EA per 31 days)
<i> fingolimod</i>	T5	PA; QL (31 EA per 31 days)
FINTEPLA	T5	PA-NS; QL (360 ML per 30 days)
FIORICET	T4	QL (403 EA per 31 days)
FIORICET WITH CODEINE	T4	PA; QL (403 EA per 31 days)
FIRDAPSE	T5	PA; QL (248 EA per 31 days)
FLECTOR	T4	PA; QL (62 EA per 31 days)
FLEQSUVY	T4	PA; QL (496 ML per 31 days)
<i>fluoxetine (pmdd)</i>	T2	
<i>fluoxetine oral capsule</i>	T1	
<i>fluoxetine oral capsule,delayed release(dr/ec)</i>	T2	
<i>fluoxetine oral solution</i>	T2	
<i>fluoxetine oral tablet 10 mg, 20 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>fluoxetine oral tablet 60 mg</i>	T4	
<i>fluphenazine decanoate</i>	T4	
<i>fluphenazine hcl</i>	T4	
<i>flurazepam</i>	T4	PA
<i>flurbiprofen oral tablet 100 mg</i>	T2	
<i>fluvoxamine oral capsule,extended release 24hr</i>	T4	
<i>fluvoxamine oral tablet</i>	T2	
FOCALIN ORAL TABLET 10 MG	T4	ST; QL (62 EA per 31 days)
FOCALIN ORAL TABLET 2.5 MG, 5 MG	T4	ST; QL (93 EA per 31 days)
FOCALIN XR	T4	ST; QL (31 EA per 31 days)
FORFIVO XL	T4	QL (31 EA per 31 days)
FROVA	T4	QL (12 EA per 28 days)
<i>frovatriptan</i>	T4	QL (12 EA per 28 days)
FYCOMPA ORAL SUSPENSION	T5	QL (744 ML per 31 days)
FYCOMPA ORAL TABLET 10 MG, 12 MG, 4 MG, 6 MG, 8 MG	T5	QL (31 EA per 31 days)
FYCOMPA ORAL TABLET 2 MG	T4	QL (31 EA per 31 days)
<i>gabapentin oral capsule 100 mg, 400 mg</i>	T1	PA-NS; QL (270 EA per 30 days)
<i>gabapentin oral capsule 300 mg</i>	T1	PA-NS; QL (360 EA per 30 days)
<i>gabapentin oral solution 250 mg/5 ml</i>	T2	PA-NS; QL (2160 ML per 30 days)
<i>gabapentin oral tablet 600 mg</i>	T1	PA-NS; QL (180 EA per 30 days)
<i>gabapentin oral tablet 800 mg</i>	T1	PA-NS; QL (120 EA per 30 days)
<i>gabapentin oral tablet extended release 24 hr 300 mg</i>	T3	PA; QL (155 EA per 31 days)
<i>gabapentin oral tablet extended release 24 hr 600 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>galantamine</i>	T3	
GEODON INTRAMUSCULAR	T4	
GEODON ORAL	T5	QL (62 EA per 31 days)
GILENYA	T5	PA; QL (31 EA per 31 days)
<i>glatiramer subcutaneous syringe 20 mg/ml</i>	T5	PA; QL (31 ML per 31 days)
<i>glatiramer subcutaneous syringe 40 mg/ml</i>	T5	PA; QL (12 ML per 28 days)
GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
GLATOPA SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
GOCOVRI ORAL CAPSULE,EXTENDED RELEASE 24HR 137 MG	T5	PA; QL (62 EA per 31 days)
GOCOVRI ORAL CAPSULE,EXTENDED RELEASE 24HR 68.5 MG	T5	PA; QL (124 EA per 31 days)
GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 300 MG	T4	PA; QL (155 EA per 31 days)
GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 450 MG	T4	PA; QL (31 EA per 31 days)
GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 600 MG	T4	PA; QL (93 EA per 31 days)
GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 750 MG, 900 MG	T4	PA; QL (62 EA per 31 days)
<i>guanfacine oral tablet extended release 24 hr</i>	T2	PA
HALCION ORAL TABLET 0.25 MG	T4	PA
HALDOL DECANOATE INTRAMUSCULAR SOLUTION 100 MG/ML	T4	
<i>haloperidol</i>	T1	
<i>haloperidol decanoate</i>	T2	
<i>haloperidol lactate injection</i>	T2	
<i>haloperidol lactate oral</i>	T2	
HETLIOZ	T5	PA; QL (31 EA per 31 days)
HETLIOZ LQ	T5	PA; QL (158 ML per 31 days)
HORIZANT ORAL TABLET EXTENDED RELEASE 300 MG	T4	PA; QL (90 EA per 30 days)
HORIZANT ORAL TABLET EXTENDED RELEASE 600 MG	T4	PA; QL (60 EA per 30 days)
<i>hydrocodone bitartrate oral capsule, oral only, er 12hr</i>	T4	PA; QL (100 EA per 31 days)
<i>hydrocodone bitartrate oral tablet,oral only,ext.rel.24 hr 100 mg, 120 mg</i>	T5	PA; QL (31 EA per 31 days)
<i>hydrocodone bitartrate oral tablet,oral only,ext.rel.24 hr 20 mg, 30 mg, 40 mg, 60 mg, 80 mg</i>	T4	PA; QL (31 EA per 31 days)
<i>hydrocodone-acetaminophen oral solution 7.5-325 mg/15 ml</i>	T3	PA; QL (5723 ML per 31 days)
<i>hydrocodone-acetaminophen oral tablet 10-300 mg, 5-300 mg, 7.5-300 mg</i>	T3	PA; QL (403 EA per 31 days)
<i>hydrocodone-acetaminophen oral tablet 10-325 mg, 5-325 mg, 7.5-325 mg</i>	T3	PA; QL (372 EA per 31 days)
<i>hydrocodone-ibuprofen</i>	T3	PA; QL (155 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>hydromorphone (pf) injection solution 10 (mg/ml) (5 ml), 10 mg/ml</i>	T4	PA; QL (124 ML per 31 days)
<i>hydromorphone oral liquid</i>	T4	PA; QL (1550 ML per 31 days)
<i>hydromorphone oral tablet</i>	T3	PA; QL (186 EA per 31 days)
<i>hydromorphone oral tablet extended release 24 hr 12 mg, 16 mg, 8 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>hydromorphone oral tablet extended release 24 hr 32 mg</i>	T4	PA; QL (48 EA per 31 days)
HYSINGLA ER ORAL TABLET,ORAL ONLY,EXT.REL.24 HR 100 MG, 120 MG, 80 MG	T5	PA; QL (31 EA per 31 days)
HYSINGLA ER ORAL TABLET,ORAL ONLY,EXT.REL.24 HR 20 MG, 30 MG, 40 MG, 60 MG	T4	PA; QL (31 EA per 31 days)
IBU ORAL TABLET 600 MG, 800 MG	T1	
<i>ibuprofen oral suspension</i>	T2	
<i>ibuprofen oral tablet 400 mg, 600 mg, 800 mg</i>	T1	
<i>ibuprofen-famotidine</i>	T4	PA; QL (93 EA per 31 days)
<i>imipramine hcl</i>	T4	PA-NS
<i>imipramine pamoate</i>	T4	PA-NS
IMITREX ORAL TABLET 100 MG	T4	QL (9 EA per 28 days)
IMITREX ORAL TABLET 25 MG	T4	QL (36 EA per 28 days)
IMITREX ORAL TABLET 50 MG	T4	QL (18 EA per 28 days)
IMITREX STATDOSE PEN SUBCUTANEOUS PEN INJECTOR 4 MG/0.5 ML	T4	QL (6 ML per 28 days)
IMITREX STATDOSE REFILL SUBCUTANEOUS CARTRIDGE 6 MG/0.5 ML	T5	QL (4 ML per 28 days)
INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE	T5	PA; QL (300 EA per 30 days)
INDOCIN ORAL	T4	
INDOCIN RECTAL	T5	
<i>indomethacin oral capsule</i>	T1	
<i>indomethacin oral capsule, extended release</i>	T2	
<i>indomethacin oral suspension</i>	T4	
<i>indomethacin rectal suppository 50 mg</i>	T5	
INGREZZA INITIATION PK(TARDIV)	T5	PA; QL (56 EA per 365 days)
INGREZZA ORAL CAPSULE 40 MG	T5	PA; QL (62 EA per 31 days)
INGREZZA ORAL CAPSULE 60 MG, 80 MG	T5	PA; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
INGREZZA SPRINKLE	T5	PA; QL (31 EA per 31 days)
INTUNIV ER	T4	PA
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,092 MG/3.5 ML	T5	QL (3.5 ML per 180 days)
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,560 MG/5 ML	T5	QL (5 ML per 180 days)
INVEGA ORAL TABLET EXTENDED RELEASE 24HR 3 MG, 9 MG	T4	QL (31 EA per 31 days)
INVEGA ORAL TABLET EXTENDED RELEASE 24HR 6 MG	T4	QL (62 EA per 31 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 117 MG/0.75 ML	T5	QL (0.75 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 156 MG/ML	T5	QL (1 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 234 MG/1.5 ML	T5	QL (1.5 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 39 MG/0.25 ML	T3	QL (0.25 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 78 MG/0.5 ML	T5	QL (0.5 ML per 28 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 273 MG/0.88 ML	T5	QL (0.88 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 410 MG/1.32 ML	T5	QL (1.32 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 546 MG/1.75 ML	T5	QL (1.75 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 819 MG/2.63 ML	T5	QL (2.63 ML per 84 days)
JORNAY PM	T4	ST; QL (31 EA per 31 days)
KEPPRA ORAL SOLUTION	T5	
KEPPRA ORAL TABLET 1,000 MG	T5	
KEPPRA ORAL TABLET 250 MG, 500 MG, 750 MG	T4	
KEPPRA XR ORAL TABLET EXTENDED RELEASE 24 HR 500 MG	T5	
KEPPRA XR ORAL TABLET EXTENDED RELEASE 24 HR 750 MG	T4	
KESIMPTA PEN	T5	PA; QL (0.4 ML per 28 days)
<i>ketoprofen oral capsule 25 mg, 50 mg</i>	T4	
<i>ketoprofen oral capsule,ext rel. pellets 24 hr 200 mg</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>ketorolac oral</i>	T4	
KEVEYIS	T5	PA; QL (124 EA per 31 days)
KIPROFEN	T4	
KLONOPIN ORAL TABLET 0.5 MG	T4	PA-NS; QL (93 EA per 31 days)
KLONOPIN ORAL TABLET 1 MG	T4	PA-NS; QL (124 EA per 31 days)
KLONOPIN ORAL TABLET 2 MG	T4	PA-NS; QL (310 EA per 31 days)
KLOXXADO	T3	
<i>lacosamide oral</i>	T4	
LAMICTAL ODT ORAL TABLET,DISINTEGRATING 100 MG	T5	
LAMICTAL ODT ORAL TABLET,DISINTEGRATING 200 MG, 25 MG, 50 MG	T4	
LAMICTAL ORAL TABLET 100 MG, 150 MG, 200 MG	T5	
LAMICTAL ORAL TABLET 25 MG	T4	
LAMICTAL ORAL TABLET, CHEWABLE DISPERSIBLE 25 MG, 5 MG	T4	
LAMICTAL STARTER (BLUE) KIT	T4	
LAMICTAL STARTER (GREEN) KIT	T4	
LAMICTAL STARTER (ORANGE) KIT	T4	
LAMICTAL XR ORAL TABLET EXTENDED RELEASE 24HR 100 MG, 250 MG, 300 MG, 50 MG	T4	
LAMICTAL XR ORAL TABLET EXTENDED RELEASE 24HR 200 MG, 25 MG	T5	
LAMICTAL XR STARTER (BLUE)	T4	
LAMICTAL XR STARTER (GREEN)	T4	
LAMICTAL XR STARTER (ORANGE)	T4	
<i>lamotrigine oral tablet</i>	T1	
<i>lamotrigine oral tablet disintegrating, dose pk</i>	T4	
<i>lamotrigine oral tablet extended release 24hr</i>	T4	
<i>lamotrigine oral tablet, chewable dispersible</i>	T2	
<i>lamotrigine oral tablet,disintegrating</i>	T4	
<i>lamotrigine oral tablets,dose pack</i>	T4	
LATUDA ORAL TABLET 120 MG, 20 MG, 40 MG, 60 MG	T5	PA-NS; QL (31 EA per 31 days)
LATUDA ORAL TABLET 80 MG	T5	PA-NS; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>levetiracetam oral solution 100 mg/ml</i>	T2	
<i>levetiracetam oral tablet</i>	T2	
<i>levetiracetam oral tablet extended release 24 hr</i>	T2	
<i>levorphanol tartrate</i>	T5	PA; QL (186 EA per 31 days)
LEXAPRO ORAL TABLET 10 MG	T4	QL (45 EA per 30 days)
LEXAPRO ORAL TABLET 20 MG, 5 MG	T4	QL (30 EA per 30 days)
LIBERVANT	T5	PA-NS; QL (10 EA per 30 days)
LICART	T4	PA; QL (31 EA per 31 days)
<i>lisdexamfetamine</i>	T4	ST; QL (31 EA per 31 days)
<i>lithium carbonate</i>	T1	
<i>lithium citrate</i>	T1	
LITHOBID	T4	
LODINE ORAL TABLET	T4	
LODOSYN	T4	
LOFENA	T5	
LORAZEPAM INTENSOL	T2	PA; QL (155 ML per 31 days)
<i>lorazepam oral tablet 0.5 mg</i>	T2	PA; QL (124 EA per 31 days)
<i>lorazepam oral tablet 1 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>lorazepam oral tablet 2 mg</i>	T2	PA; QL (155 EA per 31 days)
LOREEV XR ORAL CAPSULE,EXTENDED RELEASE 24HR 1 MG, 3 MG	T4	PA; QL (93 EA per 31 days)
LOREEV XR ORAL CAPSULE,EXTENDED RELEASE 24HR 1.5 MG, 2 MG	T4	PA; QL (155 EA per 31 days)
LORZONE	T4	PA
<i>loxapine succinate</i>	T2	
LUCEMYRA	T5	
LUMRYZ	T5	PA; QL (31 EA per 31 days)
LUNESTA	T4	PA; QL (31 EA per 31 days)
<i>lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>lurasidone oral tablet 80 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
LYBALVI	T5	PA-NS; QL (31 EA per 31 days)
LYRICA CR	T4	PA; QL (31 EA per 31 days)
LYRICA ORAL CAPSULE 100 MG, 150 MG, 200 MG, 25 MG, 50 MG, 75 MG	T4	PA-NS; QL (93 EA per 31 days)
LYRICA ORAL CAPSULE 225 MG, 300 MG	T4	PA-NS; QL (62 EA per 31 days)
LYRICA ORAL SOLUTION	T4	PA-NS; QL (930 ML per 31 days)
LYVISPAH	T4	PA; QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
MARPLAN	T4	
MAVENCLAD (10 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (4 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (5 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (6 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (7 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (8 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAVENCLAD (9 TABLET PACK)	T5	PA; QL (40 EA per 365 days)
MAXALT ORAL TABLET 10 MG	T4	QL (12 EA per 28 days)
MAXALT-MLT ORAL TABLET,DISINTEGRATING 10 MG	T4	QL (12 EA per 28 days)
MAYZENT ORAL TABLET 0.25 MG	T5	PA; QL (155 EA per 31 days)
MAYZENT ORAL TABLET 1 MG, 2 MG	T5	PA; QL (31 EA per 31 days)
MAYZENT STARTER(FOR 1MG MAINT)	T4	PA; QL (14 EA per 365 days)
MAYZENT STARTER(FOR 2MG MAINT)	T4	PA; QL (24 EA per 365 days)
<i>meclofenamate</i>	T4	
<i>mefenamic acid</i>	T4	
<i>meloxicam oral tablet</i>	T1	
<i>meloxicam submicronized</i>	T4	PA; QL (31 EA per 31 days)
<i>memantine oral capsule,sprinkle,er 24hr</i>	T4	
<i>memantine oral solution</i>	T3	
<i>memantine oral tablet</i>	T2	
<i>memantine oral tablets,dose pack</i>	T4	
<i>meperidine (pf) injection solution 100 mg/ml</i>	T4	PA; QL (200 ML per 31 days)
<i>meperidine (pf) injection solution 25 mg/ml</i>	T4	PA; QL (800 ML per 31 days)
<i>meperidine (pf) injection solution 50 mg/ml</i>	T4	PA; QL (400 ML per 31 days)
<i>meperidine oral solution</i>	T4	PA; QL (6200 ML per 31 days)
<i>meperidine oral tablet 50 mg</i>	T4	PA; QL (1240 EA per 31 days)
<i>meprobamate oral tablet 200 mg</i>	T4	QL (341 EA per 31 days)
<i>meprobamate oral tablet 400 mg</i>	T4	QL (186 EA per 31 days)
MESTINON ORAL	T5	
MESTINON TIMESPAN	T5	
METADATE CD	T4	ST; QL (31 EA per 31 days)
<i>metaxalone</i>	T4	PA
<i>methadone oral solution 10 mg/5 ml</i>	T3	PA; QL (1033 ML per 31 days)
<i>methadone oral solution 5 mg/5 ml</i>	T3	PA; QL (2066 ML per 31 days)
<i>methadone oral tablet 10 mg</i>	T3	PA; QL (206 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>methadone oral tablet 5 mg</i>	T3	PA; QL (248 EA per 31 days)
<i>methamphetamine</i>	T5	PA
<i>methocarbamol oral tablet 500 mg, 750 mg</i>	T4	
<i>methsuximide</i>	T4	
METHYLIN ORAL SOLUTION	T4	ST
<i>methylphenidate</i>	T4	PA; QL (30 EA per 30 days)
<i>methylphenidate hcl oral cap,er sprinkle,biphasic 40-60</i>	T4	QL (31 EA per 31 days)
<i>methylphenidate hcl oral capsule, er biphasic 30-70</i>	T4	QL (31 EA per 31 days)
<i>methylphenidate hcl oral capsule,er biphasic 50-50 10 mg</i>	T4	QL (186 EA per 31 days)
<i>methylphenidate hcl oral capsule,er biphasic 50-50 20 mg</i>	T4	QL (93 EA per 31 days)
<i>methylphenidate hcl oral capsule,er biphasic 50-50 30 mg, 40 mg</i>	T4	QL (62 EA per 31 days)
<i>methylphenidate hcl oral capsule,er biphasic 50-50 60 mg</i>	T4	QL (31 EA per 31 days)
<i>methylphenidate hcl oral solution</i>	T4	
<i>methylphenidate hcl oral tablet</i>	T3	QL (93 EA per 31 days)
<i>methylphenidate hcl oral tablet extended release 10 mg</i>	T4	QL (186 EA per 31 days)
<i>methylphenidate hcl oral tablet extended release 20 mg</i>	T4	QL (93 EA per 31 days)
<i>methylphenidate hcl oral tablet extended release 24hr 18 mg, 18 mg (bx rating), 27 mg, 27 mg (bx rating), 36 mg, 36 mg (bx rating), 54 mg, 54 mg (bx rating)</i>	T4	QL (31 EA per 31 days)
<i>methylphenidate hcl oral tablet extended release 24hr 45 mg, 63 mg, 72 mg</i>	T4	ST; QL (31 EA per 31 days)
<i>methylphenidate hcl oral tablet,chewable 10 mg</i>	T4	QL (186 EA per 31 days)
<i>methylphenidate hcl oral tablet,chewable 2.5 mg, 5 mg</i>	T4	QL (93 EA per 31 days)
MIGERGOT	T5	
MIGRANAL	T5	PA; QL (8 ML per 31 days)
<i>mirtazapine oral tablet</i>	T1	
<i>mirtazapine oral tablet,disintegrating</i>	T2	
<i>modafinil</i>	T2	PA; QL (31 EA per 31 days)
<i>molindone</i>	T2	
<i>morphine concentrate oral solution</i>	T3	PA; QL (310 ML per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>morphine oral capsule, er multiphase 24 hr 120 mg</i>	T4	PA; QL (51 EA per 31 days)
<i>morphine oral capsule, er multiphase 24 hr 30 mg, 45 mg, 60 mg, 75 mg, 90 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>morphine oral capsule, extend. release pellets 10 mg, 100 mg, 20 mg, 30 mg, 50 mg, 60 mg, 80 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>morphine oral solution 10 mg/5 ml</i>	T3	PA; QL (2800 ML per 31 days)
<i>morphine oral solution 20 mg/5 ml (4 mg/ml)</i>	T3	PA; QL (1400 ML per 31 days)
<i>morphine oral tablet</i>	T3	PA; QL (186 EA per 31 days)
<i>morphine oral tablet extended release 100 mg</i>	T3	PA; QL (62 EA per 31 days)
<i>morphine oral tablet extended release 15 mg, 30 mg, 60 mg</i>	T3	PA; QL (100 EA per 31 days)
<i>morphine oral tablet extended release 200 mg</i>	T3	PA; QL (31 EA per 31 days)
MOTPOLY XR ORAL CAPSULE, EXTENDED RELEASE 24HR 100 MG	T4	PA-NS; QL (31 EA per 31 days)
MOTPOLY XR ORAL CAPSULE, EXTENDED RELEASE 24HR 150 MG, 200 MG	T5	PA-NS; QL (62 EA per 31 days)
MS CONTIN ORAL TABLET EXTENDED RELEASE 100 MG	T5	PA; QL (62 EA per 31 days)
MS CONTIN ORAL TABLET EXTENDED RELEASE 15 MG, 30 MG, 60 MG	T4	PA; QL (100 EA per 31 days)
MS CONTIN ORAL TABLET EXTENDED RELEASE 200 MG	T5	PA; QL (31 EA per 31 days)
MYDAYIS	T4	ST; QL (31 EA per 31 days)
MYSOLINE	T5	
<i>nabumetone</i>	T2	
NALFON ORAL CAPSULE 400 MG	T4	
NALFON ORAL TABLET	T4	
NALOCET	T4	PA; QL (403 EA per 31 days)
<i>naloxone injection solution</i>	T2	
<i>naloxone injection syringe 0.4 mg/ml, 1 mg/ml</i>	T2	
<i>naloxone nasal</i>	T4	
<i>naltrexone</i>	T2	
NAMENDA TITRATION PAK	T4	PA
NAMENDA XR ORAL CAPSULE, SPRINKLE, ER 24HR 14 MG, 21 MG, 28 MG	T4	PA
NAMZARIC	T3	PA

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
NAPRELAN CR	T4	
NAPROSYN ORAL SUSPENSION	T4	
<i>naproxen oral suspension</i>	T2	
<i>naproxen oral tablet</i>	T1	
<i>naproxen oral tablet, delayed release (dr/ec)</i>	T2	
<i>naproxen sodium oral tablet 275 mg, 550 mg</i>	T2	
<i>naproxen sodium oral tablet, er multiphase 24 hr 375 mg, 750 mg</i>	T4	
<i>naproxen sodium oral tablet, er multiphase 24 hr 500 mg</i>	T5	
<i>naproxen-esomeprazole</i>	T5	PA; QL (62 EA per 31 days)
<i>naratriptan oral tablet 1 mg</i>	T3	QL (20 EA per 28 days)
<i>naratriptan oral tablet 2.5 mg</i>	T3	QL (9 EA per 28 days)
NARDIL	T4	
NAYZILAM	T4	PA-NS; QL (10 EA per 30 days)
<i>nefazodone</i>	T2	
NEUPRO	T4	
NEURONTIN ORAL CAPSULE 100 MG, 400 MG	T4	PA-NS; QL (270 EA per 30 days)
NEURONTIN ORAL CAPSULE 300 MG	T4	PA-NS; QL (360 EA per 30 days)
NEURONTIN ORAL SOLUTION	T4	PA-NS; QL (2160 ML per 30 days)
NEURONTIN ORAL TABLET 600 MG	T5	PA-NS; QL (180 EA per 30 days)
NEURONTIN ORAL TABLET 800 MG	T4	PA-NS; QL (120 EA per 30 days)
NORGESIC	T4	PA; QL (248 EA per 31 days)
NORGESIC FORTE	T4	PA; QL (124 EA per 31 days)
NORPRAMIN ORAL TABLET 10 MG, 25 MG	T4	
<i>nortriptyline</i>	T2	
NOURIANZ	T5	PA; QL (31 EA per 31 days)
NUCYNTA ER ORAL TABLET EXTENDED RELEASE 12 HR 100 MG, 150 MG, 200 MG, 250 MG	T5	PA; QL (62 EA per 31 days)
NUCYNTA ER ORAL TABLET EXTENDED RELEASE 12 HR 50 MG	T4	PA; QL (62 EA per 31 days)
NUCYNTA ORAL TABLET 100 MG	T5	PA; QL (186 EA per 31 days)
NUCYNTA ORAL TABLET 50 MG, 75 MG	T4	PA; QL (186 EA per 31 days)
NUEDEXTA	T5	PA; QL (62 EA per 31 days)
NUPLAZID	T5	PA-NS; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
NURTEC ODT	T5	PA; QL (18 EA per 28 days)
NUVIGIL ORAL TABLET 150 MG, 200 MG, 250 MG	T5	PA; QL (31 EA per 31 days)
NUVIGIL ORAL TABLET 50 MG	T4	PA; QL (31 EA per 31 days)
<i>olanzapine intramuscular</i>	T4	
<i>olanzapine oral</i>	T2	QL (31 EA per 31 days)
<i>olanzapine-fluoxetine</i>	T4	
ONFI ORAL SUSPENSION	T5	PA-NS; QL (496 ML per 31 days)
ONFI ORAL TABLET	T5	PA-NS; QL (62 EA per 31 days)
ONGENTYS	T4	PA; QL (31 EA per 31 days)
ONZETRA XSAIL	T4	QL (16 EA per 28 days)
OPVEE	T4	
ORMALVI	T5	PA; QL (124 EA per 31 days)
<i>orphenadrine citrate oral</i>	T4	PA
<i>orphenadrine-asa-caffeine oral tablet 25-385-30 mg</i>	T4	PA; QL (248 EA per 31 days)
OSMOLEX ER ORAL TABLET, IR - ER, BIPHASIC 24HR 129 MG	T4	PA; QL (31 EA per 31 days)
<i>oxaprozin oral tablet</i>	T4	
<i>oxazepam</i>	T4	PA
<i>oxcarbazepine oral suspension</i>	T4	
<i>oxcarbazepine oral tablet</i>	T3	
OXTELLAR XR ORAL TABLET EXTENDED RELEASE 24 HR 150 MG, 300 MG	T4	
OXTELLAR XR ORAL TABLET EXTENDED RELEASE 24 HR 600 MG	T5	
<i>oxycodone oral capsule</i>	T3	PA; QL (186 EA per 31 days)
<i>oxycodone oral concentrate</i>	T4	PA; QL (180 ML per 31 days)
<i>oxycodone oral solution</i>	T3	PA; QL (4133 ML per 31 days)
<i>oxycodone oral tablet 10 mg, 15 mg, 20 mg, 5 mg</i>	T3	PA; QL (186 EA per 31 days)
<i>oxycodone oral tablet 30 mg</i>	T3	PA; QL (138 EA per 31 days)
<i>oxycodone oral tablet,oral only,ext.rel.12 hr 10 mg, 20 mg</i>	T4	PA; QL (100 EA per 31 days)
<i>oxycodone-acetaminophen oral solution 5-325 mg/5 ml</i>	T4	PA; QL (1907 ML per 31 days)
<i>oxycodone-acetaminophen oral tablet 10-300 mg, 5-300 mg, 7.5-300 mg</i>	T5	PA; QL (403 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i>	T3	PA; QL (372 EA per 31 days)
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 10 MG, 15 MG, 20 MG, 30 MG	T3	PA; QL (100 EA per 31 days)
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 40 MG	T4	PA; QL (100 EA per 31 days)
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 60 MG	T5	PA; QL (69 EA per 31 days)
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 80 MG	T5	PA; QL (62 EA per 31 days)
<i>oxymorphone oral tablet</i>	T4	PA; QL (186 EA per 31 days)
<i>oxymorphone oral tablet extended release 12 hr 10 mg, 15 mg, 20 mg, 5 mg, 7.5 mg</i>	T4	PA; QL (100 EA per 31 days)
<i>oxymorphone oral tablet extended release 12 hr 30 mg</i>	T4	PA; QL (69 EA per 31 days)
<i>oxymorphone oral tablet extended release 12 hr 40 mg</i>	T4	PA; QL (51 EA per 31 days)
OZOBAX DS	T4	PA; QL (1240 ML per 31 days)
<i>paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 9 mg</i>	T4	QL (31 EA per 31 days)
<i>paliperidone oral tablet extended release 24hr 6 mg</i>	T4	QL (62 EA per 31 days)
PAMELOR ORAL CAPSULE 10 MG, 50 MG, 75 MG	T4	
PAMELOR ORAL CAPSULE 25 MG	T5	
PARLODEL	T4	
PARNATE	T4	
<i>paroxetine hcl oral suspension</i>	T4	
<i>paroxetine hcl oral tablet</i>	T1	
<i>paroxetine hcl oral tablet extended release 24 hr</i>	T4	
<i>paroxetine mesylate(menop.sym)</i>	T4	
PAXIL	T4	
PAXIL CR	T4	
PENNSAID TOPICAL SOLUTION IN METERED-DOSE PUMP	T5	ST; QL (224 GM per 28 days)
<i>pentazocine-naloxone</i>	T4	QL (335 EA per 31 days)
PERCOCET ORAL TABLET 10-325 MG, 5-325 MG, 7.5-325 MG	T5	PA; QL (372 EA per 31 days)
PERCOCET ORAL TABLET 2.5-325 MG	T4	PA; QL (372 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>perphenazine</i>	T2	
<i>perphenazine-amitriptyline</i>	T4	PA-NS
PERSERIS	T5	QL (1 EA per 28 days)
<i>phenelzine</i>	T3	
<i>phenobarbital</i>	T2	PA-NS
PHENYTEK	T4	
<i>phenytoin oral suspension 125 mg/5 ml</i>	T2	
<i>phenytoin oral tablet, chewable</i>	T2	
<i>phenytoin sodium extended</i>	T2	
<i>pimozide</i>	T4	
<i>piroxicam</i>	T2	
PONVORY	T5	PA; QL (31 EA per 31 days)
PONVORY 14-DAY STARTER PACK	T5	PA; QL (28 EA per 365 days)
<i>pramipexole oral tablet</i>	T2	
<i>pramipexole oral tablet extended release 24 hr</i>	T4	
<i>pregabalin oral capsule 100 mg, 150 mg, 200 mg, 25 mg, 50 mg, 75 mg</i>	T3	PA-NS; QL (93 EA per 31 days)
<i>pregabalin oral capsule 225 mg, 300 mg</i>	T3	PA-NS; QL (62 EA per 31 days)
<i>pregabalin oral solution</i>	T3	PA-NS; QL (930 ML per 31 days)
<i>pregabalin oral tablet extended release 24 hr</i>	T4	PA; QL (31 EA per 31 days)
<i>primidone oral tablet 125 mg</i>	T4	
<i>primidone oral tablet 250 mg, 50 mg</i>	T2	
PRISTIQ	T4	QL (31 EA per 31 days)
PROCENTRA	T4	
PROLATE ORAL SOLUTION	T4	PA; QL (5167 ML per 31 days)
PROLATE ORAL TABLET	T4	PA; QL (403 EA per 31 days)
<i>protriptyline</i>	T4	
PROVIGIL	T5	PA; QL (31 EA per 31 days)
PROZAC ORAL CAPSULE 10 MG, 20 MG	T4	
PROZAC ORAL CAPSULE 40 MG	T5	
<i>pyridostigmine bromide oral syrup</i>	T2	
<i>pyridostigmine bromide oral tablet 30 mg</i>	T4	
<i>pyridostigmine bromide oral tablet 60 mg</i>	T3	
<i>pyridostigmine bromide oral tablet extended release</i>	T3	
QDOLO	T5	PA; QL (2400 ML per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
QELBREE ORAL CAPSULE,EXTENDED RELEASE 24HR 100 MG, 200 MG	T4	PA; QL (93 EA per 31 days)
QELBREE ORAL CAPSULE,EXTENDED RELEASE 24HR 150 MG	T4	PA; QL (62 EA per 31 days)
QUDEXY XR ORAL CAPSULE,SPRINKLE,ER 24HR 100 MG, 150 MG, 25 MG, 50 MG	T4	
QUDEXY XR ORAL CAPSULE,SPRINKLE,ER 24HR 200 MG	T5	
<i>quetiapine oral tablet 100 mg, 200 mg, 25 mg, 300 mg, 400 mg, 50 mg</i>	T2	QL (62 EA per 31 days)
<i>quetiapine oral tablet 150 mg</i>	T3	QL (62 EA per 31 days)
<i>quetiapine oral tablet extended release 24 hr</i>	T2	QL (62 EA per 31 days)
QUILLICHEW ER ORAL TABLET,CHEW,IR-ER.BIPHASIC24HR 20 MG, 40 MG	T4	ST; QL (31 EA per 31 days)
QUILLICHEW ER ORAL TABLET,CHEW,IR-ER.BIPHASIC24HR 30 MG	T4	ST; QL (62 EA per 31 days)
QUILLIVANT XR	T4	ST; QL (360 ML per 30 days)
QULIPTA	T5	PA; QL (31 EA per 31 days)
QUVIVIQ	T4	ST; QL (31 EA per 31 days)
RADICAVA ORS STARTER KIT SUSP	T5	PA; QL (70 ML per 28 days)
<i>ramelteon</i>	T3	QL (31 EA per 31 days)
<i>rasagiline</i>	T4	
RELAFEN DS	T5	
RELEXXII ORAL TABLET EXTENDED RELEASE 24HR 18 MG, 27 MG, 36 MG, 45 MG, 63 MG	T4	ST; QL (31 EA per 31 days)
RELPAK ORAL TABLET 20 MG	T4	QL (12 EA per 28 days)
RELPAK ORAL TABLET 40 MG	T4	QL (6 EA per 28 days)
REMERON ORAL TABLET 15 MG, 30 MG	T4	
REMERON SOLTAB	T4	
RESTORIL	T4	PA; QL (31 EA per 31 days)
REXULTI ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
REYVOW ORAL TABLET 100 MG	T4	QL (8 EA per 28 days)
REYVOW ORAL TABLET 50 MG	T4	QL (4 EA per 28 days)
RISPERDAL CONSTA INTRAMUSCULAR SUSPENSION,EXTENDED REL RECON 12.5 MG/2 ML, 25 MG/2 ML, 37.5 MG/2 ML	T4	QL (2 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
RISPERDAL CONSTA INTRAMUSCULAR SUSPENSION,EXTENDED REL RECON 50 MG/2 ML	T5	QL (2 EA per 28 days)
RISPERDAL ORAL SOLUTION	T4	QL (496 ML per 31 days)
RISPERDAL ORAL TABLET 0.5 MG, 1 MG, 2 MG	T4	QL (31 EA per 31 days)
RISPERDAL ORAL TABLET 3 MG	T4	QL (93 EA per 31 days)
RISPERDAL ORAL TABLET 4 MG	T4	QL (124 EA per 31 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 12.5 mg/2 ml, 25 mg/2 ml</i>	T3	QL (2 EA per 28 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 37.5 mg/2 ml</i>	T4	QL (2 EA per 28 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 50 mg/2 ml</i>	T5	QL (2 EA per 28 days)
<i>risperidone oral solution</i>	T2	QL (496 ML per 31 days)
<i>risperidone oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T1	QL (31 EA per 31 days)
<i>risperidone oral tablet 3 mg</i>	T1	QL (93 EA per 31 days)
<i>risperidone oral tablet 4 mg</i>	T1	QL (124 EA per 31 days)
<i>risperidone oral tablet,disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T4	QL (31 EA per 31 days)
<i>risperidone oral tablet,disintegrating 3 mg</i>	T4	QL (93 EA per 31 days)
<i>risperidone oral tablet,disintegrating 4 mg</i>	T4	QL (124 EA per 31 days)
RITALIN	T4	ST; QL (93 EA per 31 days)
RITALIN LA ORAL CAPSULE,ER BIPHASIC 50-50 10 MG	T4	ST; QL (186 EA per 31 days)
RITALIN LA ORAL CAPSULE,ER BIPHASIC 50-50 20 MG, 40 MG	T4	ST; QL (31 EA per 31 days)
RITALIN LA ORAL CAPSULE,ER BIPHASIC 50-50 30 MG	T4	ST; QL (62 EA per 31 days)
<i>rivastigmine</i>	T4	QL (30 EA per 30 days)
<i>rivastigmine tartrate</i>	T3	
<i>rizatriptan oral tablet 10 mg</i>	T2	QL (12 EA per 28 days)
<i>rizatriptan oral tablet 5 mg</i>	T2	QL (24 EA per 28 days)
<i>rizatriptan oral tablet,disintegrating 10 mg</i>	T3	QL (12 EA per 28 days)
<i>rizatriptan oral tablet,disintegrating 5 mg</i>	T3	QL (24 EA per 28 days)
<i>ropinirole oral tablet</i>	T2	
<i>ropinirole oral tablet extended release 24 hr</i>	T4	
ROWEEPRA ORAL TABLET 500 MG	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ROXICODONE ORAL TABLET 15 MG	T4	PA; QL (186 EA per 31 days)
ROXICODONE ORAL TABLET 30 MG	T5	PA; QL (138 EA per 31 days)
ROXYBOND	T5	PA; QL (186 EA per 31 days)
ROZEREM	T4	QL (31 EA per 31 days)
<i>rufinamide oral suspension</i>	T5	PA-NS
<i>rufinamide oral tablet 200 mg</i>	T4	PA-NS
<i>rufinamide oral tablet 400 mg</i>	T5	PA-NS
RYTARY	T3	ST
SABRIL	T5	PA-NS
SAPHRIS	T4	PA-NS; QL (62 EA per 31 days)
SECUADO	T5	PA-NS; QL (31 EA per 31 days)
SEGLENTIS	T4	PA; QL (124 EA per 31 days)
<i>selegiline hcl</i>	T2	
SEROQUEL ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG	T4	QL (62 EA per 31 days)
SEROQUEL ORAL TABLET 400 MG	T5	QL (62 EA per 31 days)
SEROQUEL XR ORAL TABLET EXTENDED RELEASE 24 HR 150 MG, 200 MG, 50 MG	T4	QL (62 EA per 31 days)
SEROQUEL XR ORAL TABLET EXTENDED RELEASE 24 HR 300 MG, 400 MG	T5	QL (62 EA per 31 days)
<i>sertraline oral capsule</i>	T4	PA-NS; QL (31 EA per 31 days)
<i>sertraline oral concentrate</i>	T2	
<i>sertraline oral tablet</i>	T1	
SILENOR	T4	PA
SINEMET ORAL TABLET 10-100 MG, 25-100 MG	T4	
SKYCLARYS	T5	PA; QL (93 EA per 31 days)
<i>sodium oxybate</i>	T5	PA; QL (540 ML per 30 days)
SOMA	T4	PA
SPRITAM	T4	
SPRIX	T5	QL (5 EA per 31 days)
STALEVO 100	T4	
STALEVO 125	T4	
STALEVO 150	T4	
STALEVO 200	T4	
STALEVO 50	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
STALEVO 75	T4	
STRATTERA ORAL CAPSULE 10 MG, 25 MG, 40 MG	T4	ST; QL (62 EA per 31 days)
STRATTERA ORAL CAPSULE 100 MG, 60 MG, 80 MG	T4	ST; QL (31 EA per 31 days)
STRATTERA ORAL CAPSULE 18 MG	T4	ST; QL (124 EA per 31 days)
SUBOXONE SUBLINGUAL FILM 12-3 MG, 4-1 MG, 8-2 MG	T4	ST; QL (62 EA per 31 days)
SUBOXONE SUBLINGUAL FILM 2-0.5 MG	T4	ST; QL (93 EA per 31 days)
SUBVENITE	T2	
SUBVENITE STARTER (BLUE) KIT	T4	
SUBVENITE STARTER (GREEN) KIT	T4	
SUBVENITE STARTER (ORANGE) KIT	T4	
<i>sulindac</i>	T2	
<i>sumatriptan nasal spray,non-aerosol 20 mg/actuation</i>	T4	QL (8 EA per 28 days)
<i>sumatriptan nasal spray,non-aerosol 5 mg/actuation</i>	T4	QL (32 EA per 28 days)
<i>sumatriptan succinate oral tablet 100 mg</i>	T2	QL (9 EA per 28 days)
<i>sumatriptan succinate oral tablet 25 mg</i>	T2	QL (36 EA per 28 days)
<i>sumatriptan succinate oral tablet 50 mg</i>	T2	QL (18 EA per 28 days)
<i>sumatriptan succinate subcutaneous cartridge 6 mg/0.5 ml</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan succinate subcutaneous pen injector 4 mg/0.5 ml</i>	T4	QL (6 ML per 28 days)
<i>sumatriptan succinate subcutaneous pen injector 6 mg/0.5 ml</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan succinate subcutaneous solution</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan-naproxen</i>	T4	QL (9 EA per 28 days)
SUNOSI	T4	PA; QL (31 EA per 31 days)
SYMBYAX ORAL CAPSULE 3-25 MG, 6-25 MG	T4	
SYMPAZAN ORAL FILM 10 MG, 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SYMPAZAN ORAL FILM 5 MG	T4	PA-NS; QL (62 EA per 31 days)
TASCENSO ODT	T5	PA; QL (31 EA per 31 days)
<i>tasimelteon</i>	T5	PA; QL (31 EA per 31 days)
TASMAR ORAL TABLET 100 MG	T5	
TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG (14)- 240 MG (46)	T5	PA; QL (120 EA per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TECFIDERA ORAL CAPSULE,DELAYED RELEASE(DR/EC) 120 MG, 240 MG	T5	PA; QL (62 EA per 31 days)
TEGRETOL ORAL SUSPENSION	T4	
TEGRETOL ORAL TABLET	T4	
TEGRETOL XR	T4	
TEGSEDI	T5	PA; QL (6 ML per 28 days)
<i>temazepam</i>	T2	PA; QL (31 EA per 31 days)
TENCON	T4	QL (372 EA per 31 days)
<i>teriflunomide</i>	T5	PA; QL (31 EA per 31 days)
<i>tetrabenazine oral tablet 12.5 mg</i>	T5	PA; QL (93 EA per 31 days)
<i>tetrabenazine oral tablet 25 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>thioridazine</i>	T3	
<i>thiothixene</i>	T2	
<i>tiagabine</i>	T4	
<i>tizanidine oral capsule</i>	T4	
<i>tizanidine oral tablet</i>	T2	
<i>tolcapone</i>	T5	
TOLECTIN 600	T5	
<i>tolmetin oral capsule</i>	T2	
TOPAMAX ORAL CAPSULE, SPRINKLE	T4	
TOPAMAX ORAL TABLET 100 MG, 200 MG	T5	
TOPAMAX ORAL TABLET 25 MG, 50 MG	T4	
<i>topiramate oral capsule, sprinkle</i>	T2	
<i>topiramate oral capsule,extended release 24hr</i>	T4	
<i>topiramate oral capsule,sprinkle,er 24hr 100 mg, 150 mg, 25 mg, 50 mg</i>	T4	
<i>topiramate oral capsule,sprinkle,er 24hr 200 mg</i>	T5	
<i>topiramate oral tablet</i>	T1	
TOSYMRA	T5	QL (12 EA per 28 days)
<i>tramadol oral capsule,er biphase 24 hr 17-83</i>	T4	PA; QL (30 EA per 30 days)
<i>tramadol oral capsule,er biphase 24 hr 25-75 100 mg, 200 mg</i>	T4	PA; QL (30 EA per 30 days)
<i>tramadol oral solution</i>	T5	PA; QL (2400 ML per 30 days)
<i>tramadol oral tablet 100 mg</i>	T4	PA; QL (124 EA per 31 days)
<i>tramadol oral tablet 25 mg</i>	T4	PA; QL (496 EA per 31 days)
<i>tramadol oral tablet 50 mg</i>	T2	PA; QL (240 EA per 30 days)
<i>tramadol oral tablet extended release 24 hr</i>	T4	PA; QL (30 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>tramadol oral tablet, er multiphase 24 hr</i>	T4	PA; QL (30 EA per 30 days)
<i>tramadol-acetaminophen</i>	T2	PA; QL (372 EA per 31 days)
<i>tranylcypromine</i>	T4	
<i>trazodone</i>	T1	
TREXIMET	T5	QL (9 EA per 28 days)
TREZIX	T4	PA; QL (372 EA per 31 days)
<i>triazolam</i>	T4	PA
<i>trifluoperazine</i>	T2	
<i>trihexyphenidyl</i>	T4	
TRILEPTAL	T4	
<i>trimipramine</i>	T4	PA-NS
TRINTELLIX	T3	
TROKENDI XR ORAL CAPSULE,EXTENDED RELEASE 24HR 100 MG, 25 MG, 50 MG	T4	
TROKENDI XR ORAL CAPSULE,EXTENDED RELEASE 24HR 200 MG	T5	
TRUDHESA	T5	PA; QL (12 ML per 28 days)
UBRELVY ORAL TABLET 100 MG	T5	PA; QL (17 EA per 28 days)
UBRELVY ORAL TABLET 50 MG	T5	PA; QL (34 EA per 28 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 100 MG/0.28 ML	T5	QL (0.28 ML per 30 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 125 MG/0.35 ML	T5	QL (0.35 ML per 30 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 150 MG/0.42 ML	T5	QL (0.42 ML per 60 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 200 MG/0.56 ML	T5	QL (0.56 ML per 60 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 250 MG/0.7 ML	T5	QL (0.7 ML per 60 days)
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 50 MG/0.14 ML	T5	QL (0.14 ML per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
UZEDY SUBCUTANEOUS SUSPENSION,EXTENDED REL SYRING 75 MG/0.21 ML	T5	QL (0.21 ML per 30 days)
VALIUM	T4	PA-NS; QL (124 EA per 31 days)
<i>valproic acid</i>	T2	
<i>valproic acid (as sodium salt) oral solution 250 mg/5 ml</i>	T2	
VALTOCO NASAL SPRAY,NON-AEROSOL 10 MG/SPRAY (0.1 ML), 5 MG/SPRAY (0.1 ML)	T4	PA-NS; QL (10 EA per 30 days)
VALTOCO NASAL SPRAY,NON-AEROSOL 15 MG/2 SPRAY (7.5/0.1ML X 2), 20 MG/2 SPRAY (10MG/0.1ML X2)	T5	PA-NS; QL (10 EA per 30 days)
<i>venlafaxine besylate</i>	T4	PA-NS; QL (62 EA per 31 days)
<i>venlafaxine oral capsule,extended release 24hr 150 mg, 37.5 mg</i>	T2	QL (31 EA per 31 days)
<i>venlafaxine oral capsule,extended release 24hr 75 mg</i>	T2	QL (93 EA per 31 days)
<i>venlafaxine oral tablet</i>	T2	
<i>venlafaxine oral tablet extended release 24hr</i>	T2	QL (31 EA per 31 days)
VERSACLOZ	T5	QL (558 ML per 31 days)
<i>vigabatrin</i>	T5	PA-NS
VIGADRONE	T5	PA-NS
VIGPODER	T5	PA-NS
VIIBRYD ORAL TABLET	T4	QL (31 EA per 31 days)
<i>vilazodone</i>	T3	QL (31 EA per 31 days)
VIMOVO	T5	PA; QL (62 EA per 31 days)
VIMPAT ORAL SOLUTION	T5	
VIMPAT ORAL TABLET 100 MG, 150 MG, 200 MG	T5	
VIMPAT ORAL TABLET 50 MG	T3	
VIVITROL	T5	
VRAYLAR ORAL CAPSULE	T5	PA-NS; QL (31 EA per 31 days)
VUMERITY	T5	PA; QL (124 EA per 31 days)
VYVANSE	T4	ST; QL (31 EA per 31 days)
WAINUA	T5	PA; QL (0.8 ML per 30 days)
WAKIX	T5	PA; QL (62 EA per 31 days)
WELLBUTRIN SR	T4	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
WELLBUTRIN XL ORAL TABLET EXTENDED RELEASE 24 HR 150 MG	T5	QL (93 EA per 31 days)
WELLBUTRIN XL ORAL TABLET EXTENDED RELEASE 24 HR 300 MG	T5	QL (31 EA per 31 days)
XANAX ORAL TABLET 0.25 MG, 0.5 MG	T4	PA; QL (93 EA per 31 days)
XANAX ORAL TABLET 1 MG, 2 MG	T4	PA; QL (155 EA per 31 days)
XANAX XR ORAL TABLET EXTENDED RELEASE 24 HR 0.5 MG, 1 MG	T4	PA; QL (31 EA per 31 days)
XANAX XR ORAL TABLET EXTENDED RELEASE 24 HR 2 MG	T4	PA; QL (155 EA per 31 days)
XANAX XR ORAL TABLET EXTENDED RELEASE 24 HR 3 MG	T4	PA; QL (93 EA per 31 days)
XCOPRI	T5	PA-NS
XCOPRI MAINTENANCE PACK ORAL TABLET 250MG/DAY(150 MG X1-100MG X1)	T4	PA-NS
XCOPRI MAINTENANCE PACK ORAL TABLET 350 MG/DAY (200 MG X1-150MG X1)	T5	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 12.5 MG (14)- 25 MG (14)	T4	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 150 MG (14)- 200 MG (14), 50 MG (14)- 100 MG (14)	T5	PA-NS
XELSTRYM	T4	ST; QL (30 EA per 30 days)
XENAZINE ORAL TABLET 12.5 MG	T5	PA; QL (93 EA per 31 days)
XENAZINE ORAL TABLET 25 MG	T5	PA; QL (124 EA per 31 days)
XTAMPZA ER ORAL CAP,SPRINKL,ER12HR(DONT CRUSH) 13.5 MG, 18 MG, 27 MG, 9 MG	T4	PA; QL (62 EA per 31 days)
XTAMPZA ER ORAL CAP,SPRINKL,ER12HR(DONT CRUSH) 36 MG	T5	PA; QL (62 EA per 31 days)
XYREM	T5	PA; QL (540 ML per 30 days)
XYWAV	T5	PA; QL (540 ML per 30 days)
<i>zaleplon oral capsule 10 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>zaleplon oral capsule 5 mg</i>	T4	PA; QL (93 EA per 31 days)
ZANAFLEX	T4	
ZARONTIN	T4	
ZAVZPRET	T5	PA; QL (8 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZELAPAR	T5	
ZEMBRACE SYMTOUCH	T5	QL (8 ML per 28 days)
ZENZEDI	T4	QL (62 EA per 31 days)
ZEPOSIA	T5	PA; QL (31 EA per 31 days)
ZEPOSIA STARTER KIT (28-DAY)	T5	PA; QL (56 EA per 365 days)
ZEPOSIA STARTER PACK (7-DAY)	T5	PA; QL (14 EA per 365 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML	T5	PA; QL (11.648 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 23 MG/0.574 ML	T5	PA; QL (16 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 32.4 MG/0.81 ML	T5	PA; QL (22.68 ML per 28 days)
ZIMHI	T4	
<i>ziprasidone hcl</i>	T2	QL (62 EA per 31 days)
<i>ziprasidone mesylate</i>	T4	
ZIPSOR	T4	
<i>zolmitriptan nasal spray,non-aerosol 5 mg</i>	T4	QL (8 EA per 28 days)
<i>zolmitriptan oral tablet 2.5 mg</i>	T4	QL (16 EA per 28 days)
<i>zolmitriptan oral tablet 5 mg</i>	T4	QL (8 EA per 28 days)
<i>zolmitriptan oral tablet,disintegrating 2.5 mg</i>	T4	QL (16 EA per 28 days)
<i>zolmitriptan oral tablet,disintegrating 5 mg</i>	T4	QL (8 EA per 28 days)
ZOLOFT	T4	
<i>zolpidem oral capsule</i>	T4	PA; QL (31 EA per 31 days)
<i>zolpidem oral tablet</i>	T2	PA; QL (31 EA per 31 days)
<i>zolpidem oral tablet,ext release multiphase</i>	T4	PA; QL (31 EA per 31 days)
<i>zolpidem sublingual</i>	T4	PA; QL (31 EA per 31 days)
ZOMIG NASAL SPRAY,NON-AEROSOL 5 MG	T4	QL (8 EA per 28 days)
ZONEGRAN ORAL CAPSULE 100 MG, 25 MG	T5	
ZONISADE	T5	PA-NS; QL (930 ML per 31 days)
<i>zonisamide</i>	T2	
ZTALMY	T5	PA-NS; QL (1100 ML per 30 days)
ZUBSOLV SUBLINGUAL TABLET 0.7-0.18 MG, 2.9-0.71 MG, 8.6-2.1 MG	T3	QL (62 EA per 31 days)
ZUBSOLV SUBLINGUAL TABLET 1.4-0.36 MG	T3	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZUBSOLV SUBLINGUAL TABLET 11.4-2.9 MG, 5.7-1.4 MG	T3	QL (31 EA per 31 days)
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG	T5	PA-NS; QL (28 EA per 180 days)
ZURZUVAE ORAL CAPSULE 30 MG	T5	PA-NS; QL (14 EA per 180 days)
ZYPREXA INTRAMUSCULAR	T4	
ZYPREXA ORAL TABLET 10 MG, 2.5 MG, 5 MG, 7.5 MG	T4	QL (31 EA per 31 days)
ZYPREXA ORAL TABLET 15 MG, 20 MG	T5	QL (31 EA per 31 days)
ZYPREXA RELPREVV INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 210 MG	T5	QL (2 EA per 28 days)
ZYPREXA ZYDIS ORAL TABLET,DISINTEGRATING 10 MG, 5 MG	T4	QL (31 EA per 31 days)
ZYPREXA ZYDIS ORAL TABLET,DISINTEGRATING 15 MG, 20 MG	T5	QL (31 EA per 31 days)
Cardiovascular, Hypertension / Lipids		
<i>acebutolol</i>	T2	
ALDACTONE	T4	
<i>aliskiren</i>	T4	
ALTACE ORAL CAPSULE 1.25 MG	T4	QL (62 EA per 31 days)
ALTACE ORAL CAPSULE 10 MG	T4	QL (93 EA per 31 days)
ALTACE ORAL CAPSULE 2.5 MG, 5 MG	T4	
ALTOPREV ORAL TABLET EXTENDED RELEASE 24 HR 20 MG, 60 MG	T5	
ALTOPREV ORAL TABLET EXTENDED RELEASE 24 HR 40 MG	T4	
ALVAIZ ORAL TABLET 18 MG, 9 MG	T5	PA; QL (31 EA per 31 days)
ALVAIZ ORAL TABLET 36 MG, 54 MG	T5	PA; QL (62 EA per 31 days)
<i>amiloride</i>	T2	
<i>amiloride-hydrochlorothiazide</i>	T2	
<i>amiodarone oral</i>	T2	
<i>amlodipine</i>	T1	
<i>amlodipine-atorvastatin</i>	T2	
<i>amlodipine-benazepril</i>	T1	
<i>amlodipine-olmesartan</i>	T2	QL (31 EA per 31 days)
<i>amlodipine-valsartan</i>	T1	
<i>amlodipine-valsartan-hcthiazyd</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ARIXTRA SUBCUTANEOUS SYRINGE 10 MG/0.8 ML, 5 MG/0.4 ML, 7.5 MG/0.6 ML	T5	
ARIXTRA SUBCUTANEOUS SYRINGE 2.5 MG/0.5 ML	T4	
<i>aspirin-dipyridamole</i>	T4	
ASPRUZYO SPRINKLE	T4	PA; QL (60 EA per 30 days)
ATACAND	T4	
ATACAND HCT	T4	
<i>atenolol</i>	T1	
<i>atenolol-chlorthalidone</i>	T2	
ATORVALIQ	T4	PA; QL (600 ML per 30 days)
<i>atorvastatin</i>	T1	
AVALIDE	T4	QL (31 EA per 31 days)
AVAPRO	T4	QL (31 EA per 31 days)
AZOR	T4	QL (31 EA per 31 days)
<i>benazepril</i>	T1	
<i>benazepril-hydrochlorothiazide</i>	T1	
BENICAR HCT	T4	QL (31 EA per 31 days)
BENICAR ORAL TABLET 20 MG, 40 MG	T4	QL (31 EA per 31 days)
BENICAR ORAL TABLET 5 MG	T4	QL (93 EA per 31 days)
BETAPACE AF	T4	
<i>betaxolol oral</i>	T3	
BIDIL	T4	
<i>bisoprolol fumarate</i>	T2	
<i>bisoprolol-hydrochlorothiazide</i>	T1	
BRILINTA	T3	
<i>bumetanide</i>	T2	
BYSTOLIC ORAL TABLET 10 MG, 2.5 MG	T4	QL (93 EA per 31 days)
BYSTOLIC ORAL TABLET 20 MG	T4	QL (62 EA per 31 days)
BYSTOLIC ORAL TABLET 5 MG	T4	QL (217 EA per 31 days)
CABLIVI INJECTION KIT	T5	PA; QL (31 EA per 31 days)
CADUET	T4	
CAMZYOS	T5	PA; QL (31 EA per 31 days)
<i>candesartan</i>	T2	
<i>candesartan-hydrochlorothiazid</i>	T2	
<i>captopril</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
CARDIZEM CD ORAL CAPSULE,EXTENDED RELEASE 24HR 120 MG, 180 MG, 240 MG, 300 MG	T4	
CARDIZEM CD ORAL CAPSULE,EXTENDED RELEASE 24HR 360 MG	T5	
CARDIZEM LA	T4	
CARDIZEM ORAL TABLET 120 MG, 30 MG	T4	
CARDIZEM ORAL TABLET 60 MG	T5	
CARDURA	T4	
CARDURA XL	T4	
CAROSPIR	T4	
CARTIA XT	T2	
<i>carvedilol</i>	T1	
<i>carvedilol phosphate</i>	T4	
<i>chlorthalidone oral tablet 25 mg, 50 mg</i>	T2	
<i>cholestyramine (with sugar) oral powder in packet</i>	T2	
CHOLESTYRAMINE LIGHT ORAL POWDER IN PACKET	T2	
<i>cilostazol</i>	T2	
<i>clonidine</i>	T4	
<i>clonidine hcl oral tablet</i>	T1	
<i>clonidine hcl oral tablet extended release 24 hr</i>	T4	ST
<i>clopidogrel oral tablet 75 mg</i>	T1	
<i>colesevelam</i>	T4	
COLESTID ORAL TABLET	T4	
<i>colestipol oral packet</i>	T4	
<i>colestipol oral tablet</i>	T4	
COREG	T4	
COREG CR	T4	
CORGARD ORAL TABLET 20 MG, 40 MG	T4	
CORLANOR ORAL SOLUTION	T3	PA; QL (420 ML per 28 days)
CORLANOR ORAL TABLET 5 MG	T3	PA; QL (93 EA per 31 days)
CORLANOR ORAL TABLET 7.5 MG	T3	PA; QL (62 EA per 31 days)
COZAAR ORAL TABLET 100 MG	T4	QL (31 EA per 31 days)
COZAAR ORAL TABLET 25 MG	T4	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
COZAAR ORAL TABLET 50 MG	T4	QL (62 EA per 31 days)
CRESTOR	T4	
<i>dabigatran etexilate oral capsule 110 mg</i>	T4	QL (124 EA per 31 days)
<i>dabigatran etexilate oral capsule 150 mg, 75 mg</i>	T4	QL (62 EA per 31 days)
DEMSER	T5	PA
DIBENZYLINE	T5	PA
<i>digoxin oral solution</i>	T3	QL (155 ML per 31 days)
<i>digoxin oral tablet 125 mcg (0.125 mg)</i>	T2	QL (62 EA per 31 days)
<i>digoxin oral tablet 250 mcg (0.25 mg)</i>	T2	QL (31 EA per 31 days)
<i>digoxin oral tablet 62.5 mcg (0.0625 mg)</i>	T3	QL (124 EA per 31 days)
<i>diltiazem hcl oral capsule,extended release 12 hr</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24 hr 360 mg, 420 mg</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24hr 120 mg, 180 mg, 240 mg, 300 mg</i>	T2	
<i>diltiazem hcl oral tablet</i>	T2	
<i>diltiazem hcl oral tablet extended release 24 hr</i>	T2	
DILT-XR	T2	
DIOVAN HCT	T4	QL (31 EA per 31 days)
DIOVAN ORAL TABLET 160 MG, 40 MG, 80 MG	T4	QL (62 EA per 31 days)
DIOVAN ORAL TABLET 320 MG	T4	QL (31 EA per 31 days)
<i>dipyridamole oral</i>	T4	
<i>disopyramide phosphate oral capsule</i>	T4	
DIURIL	T4	
<i>dofetilide</i>	T4	
DOPTELET (10 TAB PACK)	T5	PA
DOPTELET (15 TAB PACK)	T5	PA
DOPTELET (30 TAB PACK)	T5	PA
<i>doxazosin</i>	T1	
DYRENIUM	T4	
EDARBI	T3	
EDARBYCLOR	T3	
EDECRIN	T5	
EFFIENT	T4	
ELIQUIS DVT-PE TREAT 30D START	T3	QL (74 EA per 30 days)
ELIQUIS ORAL TABLET 2.5 MG	T3	QL (60 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ELIQUIS ORAL TABLET 5 MG	T3	QL (74 EA per 30 days)
<i>enalapril maleate oral solution</i>	T4	
<i>enalapril maleate oral tablet</i>	T1	
<i>enalapril-hydrochlorothiazide</i>	T1	
<i>enoxaparin subcutaneous syringe</i>	T4	
ENTRESTO ORAL TABLET 24-26 MG	T3	QL (186 EA per 31 days)
ENTRESTO ORAL TABLET 49-51 MG	T3	QL (93 EA per 31 days)
ENTRESTO ORAL TABLET 97-103 MG	T3	QL (62 EA per 31 days)
<i>eplerenone</i>	T2	
<i>ethacrynic acid</i>	T4	
EXFORGE	T4	
EXFORGE HCT	T4	
EZALLOR SPRINKLE	T4	
<i>ezetimibe</i>	T2	
<i>ezetimibe-simvastatin</i>	T2	QL (31 EA per 31 days)
<i>felodipine</i>	T2	
<i>fenofibrate micronized oral capsule 130 mg</i>	T4	
<i>fenofibrate micronized oral capsule 134 mg, 200 mg, 43 mg, 67 mg</i>	T2	
<i>fenofibrate nanocrystallized</i>	T2	
<i>fenofibrate oral capsule</i>	T4	
<i>fenofibrate oral tablet 120 mg, 40 mg</i>	T4	
<i>fenofibrate oral tablet 160 mg, 54 mg</i>	T2	
<i>fenofibric acid (choline)</i>	T4	
FENOGLIDE	T4	
FILSPARI	T5	PA; QL (31 EA per 31 days)
<i>flecainide</i>	T2	
FLOLIPID	T4	PA
<i>fluvastatin oral capsule</i>	T2	
<i>fluvastatin oral tablet extended release 24 hr</i>	T4	
<i>fondaparinux subcutaneous syringe 10 mg/0.8 ml, 5 mg/0.4 ml, 7.5 mg/0.6 ml</i>	T5	
<i>fondaparinux subcutaneous syringe 2.5 mg/0.5 ml</i>	T4	
<i>fosinopril</i>	T1	
<i>fosinopril-hydrochlorothiazide</i>	T2	
FRAGMIN SUBCUTANEOUS SOLUTION 25,000 ANTI-XA UNIT/ML	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
FRAGMIN SUBCUTANEOUS SYRINGE 10,000 ANTI-XA UNIT/ML, 12,500 ANTI-XA UNIT/0.5 ML, 15,000 ANTI-XA UNIT/0.6 ML, 18,000 ANTI-XA UNIT/0.72 ML, 7,500 ANTI- XA UNIT/0.3 ML	T5	
FRAGMIN SUBCUTANEOUS SYRINGE 2,500 ANTI-XA UNIT/0.2 ML, 5,000 ANTI- XA UNIT/0.2 ML	T4	
FUROSCIX	T5	PA; QL (8 EA per 30 days)
<i>furosemide injection solution</i>	T2	
<i>furosemide oral solution 10 mg/ml, 40 mg/5 ml (8 mg/ml)</i>	T2	
<i>furosemide oral tablet</i>	T1	
<i>gemfibrozil</i>	T1	
<i>guanfacine oral tablet</i>	T4	
<i>heparin (porcine) injection solution</i>	T3	
<i>hydralazine oral</i>	T2	
<i>hydrochlorothiazide</i>	T1	
HYZAAR	T4	
<i>icosapent ethyl oral capsule 0.5 gram</i>	T2	QL (248 EA per 31 days)
<i>icosapent ethyl oral capsule 1 gram</i>	T2	QL (124 EA per 31 days)
<i>indapamide</i>	T1	
INDERAL LA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 60 MG	T4	
INDERAL LA ORAL CAPSULE,EXTENDED RELEASE 24 HR 160 MG, 80 MG	T5	
INNOPRAN XL	T5	
INSPRA	T4	
<i>irbesartan</i>	T1	QL (31 EA per 31 days)
<i>irbesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
ISORDIL	T5	
ISORDIL TITRADOSE ORAL TABLET 5 MG	T4	
<i>isosorbide dinitrate oral tablet 10 mg, 20 mg, 30 mg, 5 mg</i>	T2	
<i>isosorbide dinitrate oral tablet 40 mg</i>	T4	
<i>isosorbide mononitrate</i>	T1	
<i>isosorbide-hydralazine</i>	T3	
<i>isradipine</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
JANTOVEN	T1	
JUXTAPID	T5	PA
KATERZIA	T4	PA
KERENDIA	T4	PA; QL (31 EA per 31 days)
<i>labetalol oral</i>	T2	
LANOXIN ORAL TABLET 125 MCG (0.125 MG)	T4	QL (62 EA per 31 days)
LANOXIN ORAL TABLET 250 MCG (0.25 MG)	T4	QL (31 EA per 31 days)
LANOXIN ORAL TABLET 62.5 MCG (0.0625 MG)	T4	QL (124 EA per 31 days)
LASIX	T4	
LESCOL XL	T4	
<i>levamlodipine</i>	T4	ST; QL (31 EA per 31 days)
LIPITOR	T4	
LIPOFEN	T4	
<i>lisinopril</i>	T1	
<i>lisinopril-hydrochlorothiazide</i>	T1	
LIVALO	T3	
LODOCO	T4	PA; QL (31 EA per 31 days)
LOPID	T4	
LOPRESSOR ORAL	T4	
<i>losartan oral tablet 100 mg</i>	T1	QL (31 EA per 31 days)
<i>losartan oral tablet 25 mg</i>	T1	QL (93 EA per 31 days)
<i>losartan oral tablet 50 mg</i>	T1	QL (62 EA per 31 days)
<i>losartan-hydrochlorothiazide</i>	T1	
LOTENSIN ORAL TABLET 10 MG, 20 MG, 40 MG	T4	
LOTREL	T4	
<i>lovastatin</i>	T1	
LOVAZA	T4	PA; QL (124 EA per 31 days)
LOVENOX SUBCUTANEOUS SYRINGE 100 MG/ML	T5	
LOVENOX SUBCUTANEOUS SYRINGE 120 MG/0.8 ML, 150 MG/ML, 30 MG/0.3 ML, 40 MG/0.4 ML, 60 MG/0.6 ML, 80 MG/0.8 ML	T4	
MATZIM LA	T2	
<i>metolazone</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>metoprolol succinate</i>	T1	
<i>metoprolol ta-hydrochlorothiaz</i>	T2	
<i>metoprolol tartrate oral</i>	T1	
<i>metyrosine</i>	T5	PA
<i>mexiletine</i>	T2	
MICARDIS	T4	
MICARDIS HCT	T4	
<i>minoxidil oral</i>	T2	
<i>moexipril</i>	T1	
MULPLETA	T5	PA
MULTAQ	T3	
<i>nadolol</i>	T2	
<i>nebivolol oral tablet 10 mg, 2.5 mg</i>	T2	QL (93 EA per 31 days)
<i>nebivolol oral tablet 20 mg</i>	T2	QL (62 EA per 31 days)
<i>nebivolol oral tablet 5 mg</i>	T2	QL (217 EA per 31 days)
NEXICLON XR	T4	ST
NEXLETOL	T3	PA; QL (31 EA per 31 days)
NEXLIZET	T4	PA; QL (31 EA per 31 days)
<i>niacin oral tablet 500 mg</i>	T4	
<i>niacin oral tablet extended release 24 hr 1,000 mg, 750 mg</i>	T4	
<i>niacin oral tablet extended release 24 hr 500 mg</i>	T4	QL (31 EA per 31 days)
NIACOR	T4	
<i>nicardipine oral</i>	T4	
<i>nifedipine oral capsule</i>	T4	
<i>nifedipine oral tablet extended release</i>	T2	
<i>nifedipine oral tablet extended release 24hr</i>	T2	
<i>nimodipine</i>	T4	
<i>nisoldipine</i>	T4	
NITRO-BID	T2	
NITRO-DUR TRANSDERMAL PATCH 24 HOUR 0.1 MG/HR, 0.2 MG/HR, 0.4 MG/HR, 0.6 MG/HR	T4	
NITRO-DUR TRANSDERMAL PATCH 24 HOUR 0.3 MG/HR, 0.8 MG/HR	T5	
<i>nitroglycerin sublingual</i>	T2	
<i>nitroglycerin transdermal patch 24 hour</i>	T2	
<i>nitroglycerin translingual</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
NITROLINGUAL	T4	
NITROSTAT	T4	
NORLIQVA	T4	PA; QL (300 ML per 30 days)
NORPACE	T4	
NORPACE CR	T4	
NORVASC	T4	
NYMALIZE ORAL SYRINGE 60 MG/10 ML	T5	
<i>olmesartan oral tablet 20 mg, 40 mg</i>	T1	QL (31 EA per 31 days)
<i>olmesartan oral tablet 5 mg</i>	T1	QL (93 EA per 31 days)
<i>olmesartan-amlodipin-hcthiazyd</i>	T3	
<i>olmesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>omega-3 acid ethyl esters</i>	T2	QL (124 EA per 31 days)
ORENITRAM MONTH 1 TITRATION KT	T5	PA; QL (336 EA per 365 days)
ORENITRAM MONTH 2 TITRATION KT	T5	PA; QL (672 EA per 365 days)
ORENITRAM MONTH 3 TITRATION KT	T5	PA; QL (504 EA per 365 days)
ORENITRAM ORAL TABLET EXTENDED RELEASE 0.125 MG	T4	PA; QL (93 EA per 31 days)
ORENITRAM ORAL TABLET EXTENDED RELEASE 0.25 MG, 1 MG	T5	PA; QL (186 EA per 31 days)
ORENITRAM ORAL TABLET EXTENDED RELEASE 2.5 MG	T5	PA; QL (521 EA per 31 days)
ORENITRAM ORAL TABLET EXTENDED RELEASE 5 MG	T5	PA; QL (261 EA per 31 days)
PACERONE ORAL TABLET 100 MG, 200 MG, 400 MG	T2	
<i>pentoxifylline</i>	T2	
<i>perindopril erbumine</i>	T1	
<i>phenoxybenzamine</i>	T5	PA
<i>pindolol</i>	T3	
<i>pitavastatin calcium</i>	T3	
PLAVIX ORAL TABLET 75 MG	T4	
PRADAXA ORAL CAPSULE 110 MG	T4	QL (124 EA per 31 days)
PRADAXA ORAL CAPSULE 150 MG, 75 MG	T4	QL (62 EA per 31 days)
PRADAXA ORAL PELLETS IN PACKET 110 MG, 30 MG, 40 MG, 50 MG	T5	QL (120 EA per 30 days)
PRADAXA ORAL PELLETS IN PACKET 150 MG, 20 MG	T5	QL (60 EA per 30 days)
PRALUENT PEN	T4	PA; QL (2 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>prasugrel</i>	T2	
<i>pravastatin</i>	T1	
<i>prazosin</i>	T2	
PREVALITE ORAL POWDER IN PACKET	T2	
PROCARDIA XL	T4	
PROMACTA ORAL POWDER IN PACKET 12.5 MG	T5	PA; QL (372 EA per 31 days)
PROMACTA ORAL POWDER IN PACKET 25 MG	T5	PA; QL (31 EA per 31 days)
PROMACTA ORAL TABLET 12.5 MG, 25 MG	T5	PA; QL (31 EA per 31 days)
PROMACTA ORAL TABLET 50 MG, 75 MG	T5	PA; QL (62 EA per 31 days)
<i>propafenone oral capsule, extended release 12 hr</i>	T4	
<i>propafenone oral tablet</i>	T2	
<i>propranolol oral capsule, extended release 24 hr</i>	T2	
<i>propranolol oral solution</i>	T2	
<i>propranolol oral tablet</i>	T1	
QBRELIS	T5	
QUESTRAN LIGHT	T4	
QUESTRAN ORAL POWDER	T4	
<i>quinapril</i>	T1	
<i>quinidine gluconate oral</i>	T4	
<i>quinidine sulfate oral tablet</i>	T2	
<i>ramipril</i>	T1	
<i>ranolazine</i>	T2	QL (62 EA per 31 days)
REPATHA PUSHTRONEX	T3	PA; QL (7 ML per 28 days)
REPATHA SURECLICK	T3	PA; QL (3 ML per 28 days)
REPATHA SYRINGE	T3	PA; QL (3 ML per 28 days)
<i>rosuvastatin</i>	T1	
RYTHMOL SR	T4	
SAVAYSA	T4	QL (31 EA per 31 days)
<i>simvastatin</i>	T1	
SOAANZ	T4	ST
SORINE ORAL TABLET 120 MG, 160 MG	T2	
SOTALOL AF	T2	
<i>sotalol oral</i>	T2	
SOTYLIZE	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>spironolactone oral suspension</i>	T4	
<i>spironolactone oral tablet</i>	T1	
<i>spironolacton-hydrochlorothiaz</i>	T2	
SULAR ORAL TABLET EXTENDED RELEASE 24 HR 17 MG, 34 MG, 8.5 MG	T4	
TAVALISSE	T5	PA; QL (62 EA per 31 days)
TEKTURNA	T4	
<i>telmisartan</i>	T2	
<i>telmisartan-amlodipine</i>	T2	
<i>telmisartan-hydrochlorothiazid</i>	T2	
TENORETIC 100	T4	
TENORETIC 50	T4	
TENORMIN	T4	
<i>terazosin</i>	T1	
THALITONE	T4	
TIADYLT ER	T2	
TIAZAC	T4	
TIKOSYN	T4	
<i>timolol maleate oral</i>	T2	
TOPROL XL	T4	
<i>toremide oral</i>	T2	
<i>trandolapril</i>	T1	
<i>trandolapril-verapamil</i>	T2	
<i>triamterene</i>	T4	
<i>triamterene-hydrochlorothiazid</i>	T1	
TRIBENZOR	T4	
TRICOR	T4	
TRILIPIX	T4	
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 400 MCG, 600 MCG, 800 MCG	T5	PA; QL (62 EA per 31 days)
UPTRAVI ORAL TABLET 200 MCG	T5	PA; QL (224 EA per 28 days)
UPTRAVI ORAL TABLETS,DOSE PACK	T5	PA; QL (400 EA per 365 days)
<i>valsartan oral solution</i>	T5	QL (2480 ML per 31 days)
<i>valsartan oral tablet 160 mg, 40 mg, 80 mg</i>	T1	QL (62 EA per 31 days)
<i>valsartan oral tablet 320 mg</i>	T1	QL (31 EA per 31 days)
<i>valsartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
VASCEPA ORAL CAPSULE 0.5 GRAM	T3	QL (248 EA per 31 days)
VASCEPA ORAL CAPSULE 1 GRAM	T3	QL (124 EA per 31 days)
VASERETIC	T4	
VASOTEC ORAL TABLET 10 MG	T5	
VASOTEC ORAL TABLET 2.5 MG, 20 MG, 5 MG	T4	
VECAMYL	T5	
<i>verapamil oral capsule, 24 hr er pellet ct</i>	T2	
<i>verapamil oral capsule,ext rel. pellets 24 hr</i>	T2	
<i>verapamil oral tablet</i>	T1	
<i>verapamil oral tablet extended release</i>	T2	
VERELAN	T4	
VERELAN PM	T4	
VERQUVO	T3	PA; QL (31 EA per 31 days)
VYNDAMAX	T5	PA; QL (31 EA per 31 days)
VYNDAQEL	T5	PA; QL (124 EA per 31 days)
VYTORIN 10-10	T4	QL (31 EA per 31 days)
VYTORIN 10-20	T4	QL (31 EA per 31 days)
VYTORIN 10-40	T4	QL (31 EA per 31 days)
VYTORIN 10-80	T4	QL (31 EA per 31 days)
<i>warfarin</i>	T1	
WELCHOL	T4	
XARELTO DVT-PE TREAT 30D START	T3	QL (51 EA per 30 days)
XARELTO ORAL SUSPENSION FOR RECONSTITUTION	T3	QL (930 ML per 31 days)
XARELTO ORAL TABLET 10 MG, 20 MG	T3	QL (31 EA per 31 days)
XARELTO ORAL TABLET 15 MG	T3	QL (52 EA per 31 days)
XARELTO ORAL TABLET 2.5 MG	T3	QL (62 EA per 31 days)
ZESTORETIC	T4	
ZESTRIL	T4	
ZETIA	T4	
ZIAC	T4	
ZOCOR ORAL TABLET 10 MG, 20 MG, 40 MG	T4	
ZONTIVITY	T4	
ZYPITAMAG	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
Dermatologicals/Topical Therapy		
ABSORICA	T5	
ABSORICA LD	T5	
ACANYA TOPICAL GEL WITH PUMP	T4	
ACCUTANE ORAL CAPSULE 10 MG, 20 MG, 40 MG	T4	
<i>acitretin</i>	T4	PA
<i>acyclovir topical cream</i>	T4	QL (5 GM per 28 days)
<i>acyclovir topical ointment</i>	T4	QL (30 GM per 30 days)
ACZONE	T4	QL (90 GM per 28 days)
<i>adapalene topical cream</i>	T4	PA
<i>adapalene topical gel 0.3 %</i>	T4	PA
<i>adapalene topical swab</i>	T4	PA
<i>adapalene-benzoyl peroxide</i>	T4	PA
ADBRY SUBCUTANEOUS SYRINGE	T5	PA; QL (4 ML per 28 days)
AKLIEF	T4	PA
ALA-CORT TOPICAL CREAM 1 %	T2	
ALA-CORT TOPICAL CREAM 2.5 %	T2	QL (30 GM per 28 days)
ALA-SCALP	T4	
<i>alclometasone</i>	T2	
ALTABAX	T4	QL (30 GM per 28 days)
ALTRENO	T4	PA; QL (45 GM per 28 days)
<i>amcinonide topical cream</i>	T2	
<i>amcinonide topical ointment</i>	T2	
<i>ammonium lactate</i>	T2	
AMNESTEEM	T4	
AMZEEQ	T4	
APEXICON E	T4	QL (60 GM per 28 days)
ARAZLO	T4	PA; QL (45 GM per 28 days)
ATRALIN	T4	PA; QL (45 GM per 28 days)
<i>azelaic acid</i>	T4	QL (50 GM per 28 days)
AZELEX	T4	QL (50 GM per 28 days)
BENZAMYCIN	T4	
<i>betamethasone dipropionate</i>	T2	
<i>betamethasone valerate topical cream</i>	T2	
<i>betamethasone valerate topical foam</i>	T4	
<i>betamethasone valerate topical lotion</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>betamethasone valerate topical ointment</i>	T2	
<i>betamethasone, augmented</i>	T2	
BIMZELX	T5	PA; QL (2 ML per 28 days)
BIMZELX AUTOINJECTOR	T5	PA; QL (2 ML per 28 days)
<i>brimonidine topical</i>	T4	
BRYHALI	T4	QL (100 GM per 28 days)
CABTREO	T4	PA; QL (50 GM per 28 days)
<i>calcipotriene scalp</i>	T3	QL (60 ML per 28 days)
<i>calcipotriene topical cream</i>	T4	QL (60 GM per 28 days)
<i>calcipotriene topical foam</i>	T5	ST; QL (120 GM per 28 days)
<i>calcipotriene topical ointment</i>	T4	QL (60 GM per 28 days)
<i>calcipotriene-betamethasone</i>	T4	ST; QL (400 GM per 28 days)
<i>calcitriol topical</i>	T4	ST
CARAC	T5	PA
CIBINQO	T5	PA; QL (31 EA per 31 days)
<i>ciclopirox topical cream</i>	T2	QL (90 GM per 28 days)
<i>ciclopirox topical gel</i>	T2	QL (45 GM per 28 days)
<i>ciclopirox topical shampoo</i>	T2	QL (120 ML per 28 days)
<i>ciclopirox topical solution</i>	T2	
<i>ciclopirox topical suspension</i>	T2	QL (60 ML per 28 days)
CLARAVIS	T4	
CLEOCIN T TOPICAL LOTION	T4	QL (60 ML per 28 days)
CLINDACIN	T2	QL (100 GM per 28 days)
CLINDACIN ETZ TOPICAL SWAB	T4	
CLINDAGEL	T5	QL (75 ML per 28 days)
<i>clindamycin phosphate topical foam</i>	T4	QL (100 GM per 28 days)
<i>clindamycin phosphate topical gel</i>	T2	QL (60 GM per 28 days)
<i>clindamycin phosphate topical gel, once daily</i>	T2	QL (75 ML per 28 days)
<i>clindamycin phosphate topical lotion</i>	T3	QL (60 ML per 28 days)
<i>clindamycin phosphate topical solution</i>	T3	QL (60 ML per 28 days)
<i>clindamycin phosphate topical swab</i>	T4	
<i>clindamycin-benzoyl peroxide topical gel</i>	T4	
<i>clindamycin-benzoyl peroxide topical gel with pump 1.2 %(1 % base) -3.75 %, 1.2-2.5 %</i>	T4	
<i>clindamycin-tretinoin</i>	T4	PA; QL (60 GM per 28 days)
<i>clobetasol scalp</i>	T4	QL (50 ML per 28 days)
<i>clobetasol topical cream</i>	T4	QL (60 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>clobetasol topical foam</i>	T4	QL (100 GM per 28 days)
<i>clobetasol topical gel</i>	T4	QL (60 GM per 28 days)
<i>clobetasol topical lotion</i>	T4	QL (118 ML per 28 days)
<i>clobetasol topical ointment</i>	T4	QL (60 GM per 28 days)
<i>clobetasol topical shampoo</i>	T4	QL (118 ML per 28 days)
<i>clobetasol topical spray,non-aerosol</i>	T4	QL (125 ML per 28 days)
<i>clobetasol-emollient topical cream</i>	T4	QL (60 GM per 28 days)
<i>clobetasol-emollient topical foam</i>	T4	QL (100 GM per 28 days)
CLOBEX TOPICAL LOTION	T4	QL (118 ML per 28 days)
CLOBEX TOPICAL SHAMPOO	T4	QL (118 ML per 28 days)
CLOBEX TOPICAL SPRAY,NON-AEROSOL	T4	QL (125 ML per 28 days)
<i>clocortolone pivalate</i>	T4	
CLODAN	T4	QL (118 ML per 28 days)
<i>clotrimazole topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole topical solution</i>	T2	QL (30 ML per 28 days)
<i>clotrimazole-betamethasone topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole-betamethasone topical lotion</i>	T2	QL (60 ML per 28 days)
CONDYLOX TOPICAL GEL	T4	
CORDRAN TAPE LARGE ROLL	T4	
CORDRAN TOPICAL CREAM 0.05 %	T4	QL (120 GM per 28 days)
CORDRAN TOPICAL LOTION	T4	QL (120 ML per 28 days)
COSENTYX (2 SYRINGES)	T5	PA; QL (2 ML per 28 days)
COSENTYX PEN (2 PENS)	T5	PA; QL (2 ML per 28 days)
COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
COSENTYX UNOREADY PEN	T5	PA; QL (2 ML per 28 days)
CROTAN	T4	
<i>dapsone topical</i>	T4	QL (90 GM per 28 days)
DENAVIR	T4	QL (5 GM per 28 days)
DERMA-SMOOTH/FS SCALP OIL	T4	QL (118.28 ML per 28 days)
<i>desonide topical cream</i>	T4	QL (60 GM per 28 days)
<i>desonide topical gel</i>	T4	QL (60 GM per 28 days)
<i>desonide topical lotion</i>	T4	QL (118 ML per 28 days)
<i>desonide topical ointment</i>	T4	QL (60 GM per 28 days)
DESOWEN TOPICAL CREAM	T4	QL (60 GM per 28 days)
<i>desoximetasone topical cream</i>	T4	QL (100 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>desoximetasone topical gel</i>	T4	QL (60 GM per 28 days)
<i>desoximetasone topical ointment</i>	T4	QL (100 GM per 28 days)
<i>desoximetasone topical spray,non-aerosol</i>	T4	QL (100 ML per 28 days)
<i>diclofenac sodium topical gel 3 %</i>	T4	PA; QL (100 GM per 28 days)
DIFFERIN TOPICAL CREAM	T4	PA
DIFFERIN TOPICAL GEL WITH PUMP	T4	PA
DIFFERIN TOPICAL LOTION	T4	PA
<i>diflorasone</i>	T4	QL (60 GM per 28 days)
DIPROLENE (AUGMENTED) TOPICAL OINTMENT	T4	
<i>doxepin topical</i>	T4	PA; QL (45 GM per 28 days)
DUOBRII	T4	PA; QL (200 GM per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 100 MG/0.67 ML	T5	PA; QL (1.34 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
<i>econazole</i>	T4	QL (85 GM per 28 days)
EFUDEX TOPICAL CREAM	T4	
ELIDEL	T4	QL (100 GM per 28 days)
ENSTILAR	T5	ST; QL (60 GM per 28 days)
EPIDUO FORTE	T4	PA
EPIDUO TOPICAL GEL WITH PUMP	T4	PA
EPSOLAY	T4	QL (30 GM per 28 days)
ERTACZO	T4	ST; QL (60 GM per 28 days)
ERY PADS	T2	
ERYGEL	T4	QL (60 GM per 28 days)
<i>erythromycin with ethanol topical gel</i>	T4	QL (60 GM per 28 days)
<i>erythromycin with ethanol topical solution</i>	T2	QL (60 ML per 28 days)
<i>erythromycin-benzoyl peroxide</i>	T4	
EUCRISA	T4	PA; QL (60 GM per 30 days)
EXELDERM TOPICAL CREAM	T4	ST; QL (60 GM per 28 days)
EXELDERM TOPICAL SOLUTION	T4	ST; QL (30 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
FABIOR	T4	PA; QL (100 GM per 28 days)
FILSUVEZ	T5	PA
FINACEA	T4	QL (50 GM per 28 days)
<i>fluocinolone and shower cap</i>	T4	QL (118.28 ML per 28 days)
<i>fluocinolone topical cream 0.01 %</i>	T4	QL (60 GM per 28 days)
<i>fluocinolone topical cream 0.025 %</i>	T4	QL (120 GM per 28 days)
<i>fluocinolone topical ointment</i>	T4	QL (120 GM per 28 days)
<i>fluocinolone topical solution</i>	T4	QL (90 ML per 28 days)
<i>fluocinonide topical cream 0.05 %</i>	T4	QL (60 GM per 28 days)
<i>fluocinonide topical cream 0.1 %</i>	T4	QL (120 GM per 28 days)
<i>fluocinonide topical gel</i>	T4	QL (60 GM per 28 days)
<i>fluocinonide topical ointment</i>	T4	QL (60 GM per 28 days)
<i>fluocinonide topical solution</i>	T4	QL (60 ML per 28 days)
<i>fluocinonide-emollient</i>	T4	QL (60 GM per 28 days)
<i>fluorouracil topical cream 0.5 %</i>	T5	PA
<i>fluorouracil topical cream 5 %</i>	T3	
<i>fluorouracil topical solution</i>	T3	
<i>flurandrenolide topical cream</i>	T4	QL (120 GM per 28 days)
<i>flurandrenolide topical lotion</i>	T4	QL (120 ML per 28 days)
<i>fluticasone propionate topical cream</i>	T4	
<i>fluticasone propionate topical lotion</i>	T4	QL (120 ML per 28 days)
<i>fluticasone propionate topical ointment</i>	T4	
<i>gentamicin topical</i>	T2	QL (60 GM per 28 days)
<i>halcinonide topical cream</i>	T4	QL (60 GM per 28 days)
<i>halobetasol propionate topical cream</i>	T4	QL (50 GM per 28 days)
<i>halobetasol propionate topical foam</i>	T4	QL (120 GM per 28 days)
<i>halobetasol propionate topical ointment</i>	T4	QL (50 GM per 28 days)
HALOG TOPICAL CREAM	T4	QL (60 GM per 28 days)
HALOG TOPICAL OINTMENT	T4	QL (60 GM per 28 days)
HALOG TOPICAL SOLUTION	T4	QL (120 ML per 28 days)
<i>hydrocortisone butyrate topical cream</i>	T4	QL (45 GM per 28 days)
<i>hydrocortisone butyrate topical lotion</i>	T4	QL (118 ML per 28 days)
<i>hydrocortisone butyrate topical ointment</i>	T4	QL (45 GM per 28 days)
<i>hydrocortisone butyrate topical solution</i>	T4	QL (60 ML per 28 days)
<i>hydrocortisone topical cream 1 %</i>	T2	
<i>hydrocortisone topical lotion 2.5 %</i>	T2	QL (118 ML per 28 days)
<i>hydrocortisone topical ointment 1 %, 2.5 %</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>hydrocortisone valerate</i>	T4	QL (60 GM per 28 days)
HYFTOR	T5	PA; QL (30 GM per 30 days)
ILUMYA	T5	PA; QL (1 ML per 84 days)
<i>imiquimod topical cream in metered-dose pump</i>	T5	
<i>imiquimod topical cream in packet 5 %</i>	T2	
<i>isotretinoin</i>	T4	
<i>ivermectin topical cream</i>	T2	QL (45 GM per 28 days)
JUBLIA	T4	QL (8 ML per 28 days)
KENALOG TOPICAL	T4	QL (100 GM per 28 days)
<i>ketoconazole topical cream</i>	T2	QL (60 GM per 28 days)
<i>ketoconazole topical foam</i>	T4	QL (100 GM per 28 days)
<i>ketoconazole topical shampoo</i>	T2	QL (120 ML per 28 days)
KETODAN	T4	ST; QL (100 GM per 28 days)
KLARON	T4	
LEXETTE	T5	QL (200 GM per 28 days)
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i>	T2	PA; QL (50 ML per 28 days)
<i>lidocaine topical adhesive patch,medicated 5 %</i>	T2	PA; QL (93 EA per 31 days)
<i>lidocaine topical ointment</i>	T4	PA; QL (50 GM per 28 days)
LIDOCAINE VISCOUS	T2	
<i>lidocaine-prilocaine topical cream</i>	T2	PA; QL (30 GM per 28 days)
LIDOCAN III	T4	PA; QL (93 EA per 31 days)
LIDODERM	T5	PA; QL (93 EA per 31 days)
LOCOID LIPOCREAM	T4	QL (60 GM per 28 days)
LOCOID TOPICAL LOTION	T4	QL (118 ML per 28 days)
LOPROX TOPICAL SHAMPOO	T4	QL (120 ML per 28 days)
<i>luliconazole</i>	T4	QL (60 GM per 28 days)
LUZU	T4	QL (60 GM per 28 days)
<i>mafenide acetate</i>	T2	
<i>malathion</i>	T2	
<i>methoxsalen</i>	T5	
METROCREAM	T4	
METROGEL TOPICAL GEL 1 %	T4	
METROLOTION	T4	
<i>metronidazole topical cream</i>	T4	
<i>metronidazole topical gel</i>	T4	
<i>metronidazole topical lotion</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
MIRVASO	T4	
<i>mometasone topical</i>	T2	
<i>mupirocin</i>	T2	
<i>mupirocin calcium</i>	T4	ST
<i>naftifine topical cream 1 %</i>	T4	ST; QL (90 GM per 28 days)
<i>naftifine topical cream 2 %</i>	T4	ST; QL (60 GM per 28 days)
<i>naftifine topical gel 2 %</i>	T4	ST; QL (60 GM per 28 days)
NAFTIN TOPICAL GEL 1 %	T4	ST; QL (90 GM per 28 days)
NAFTIN TOPICAL GEL 2 %	T4	ST; QL (60 GM per 28 days)
NATROBA	T4	
NEO-SYNALAR	T4	
NEUAC	T4	
NORITATE	T5	
NYAMYC	T2	QL (60 GM per 28 days)
<i>nystatin topical cream</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical ointment</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical powder</i>	T2	QL (60 GM per 28 days)
<i>nystatin-triamcinolone</i>	T3	QL (60 GM per 28 days)
NYSTOP	T2	QL (60 GM per 28 days)
ONEXTON TOPICAL GEL WITH PUMP	T4	
OPZELURA	T5	PA; QL (240 GM per 28 days)
OVIDE	T4	
<i>oxiconazole</i>	T4	ST; QL (90 GM per 28 days)
OXISTAT TOPICAL CREAM	T4	ST; QL (90 GM per 28 days)
OXISTAT TOPICAL LOTION	T4	ST; QL (60 ML per 28 days)
PANDEL	T5	QL (80 GM per 28 days)
PANRETIN	T5	PA-NS
<i>penciclovir</i>	T3	QL (5 GM per 28 days)
<i>permethrin</i>	T2	
<i>pimecrolimus</i>	T4	QL (100 GM per 28 days)
PLIAGLIS	T4	PA; QL (100 GM per 28 days)
<i>podofilox topical gel</i>	T4	
<i>podofilox topical solution</i>	T2	
PRUDOXIN	T4	PA; QL (45 GM per 28 days)
QBREXZA	T4	QL (30 EA per 30 days)
REGRANEX	T5	PA
RETIN-A	T4	PA; QL (45 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
RETIN-A MICRO	T4	PA; QL (45 GM per 28 days)
RETIN-A MICRO PUMP TOPICAL GEL WITH PUMP 0.06 %	T4	PA; QL (50 GM per 28 days)
RETIN-A MICRO PUMP TOPICAL GEL WITH PUMP 0.08 %	T5	PA; QL (50 GM per 28 days)
SANTYL	T4	QL (180 GM per 30 days)
<i>selenium sulfide topical lotion</i>	T2	
SILIQ	T5	PA; QL (6 ML per 28 days)
SILVADENE	T4	
<i>silver sulfadiazine</i>	T2	
SKYRIZI SUBCUTANEOUS PEN INJECTOR	T5	PA; QL (1 ML per 84 days)
SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML	T5	PA; QL (1 ML per 84 days)
SOOLANTRA	T4	QL (45 GM per 28 days)
SORILUX	T5	ST; QL (120 GM per 28 days)
SOTYKTU	T5	PA; QL (31 EA per 31 days)
SPEVIGO SUBCUTANEOUS	T5	PA; QL (2 ML per 28 days)
<i>spinosad</i>	T4	
SSD	T4	
STELARA SUBCUTANEOUS SOLUTION	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
<i>sulfacetamide sodium (acne)</i>	T2	
SULFAMYLON TOPICAL CREAM	T3	
SYNALAR TOPICAL CREAM	T4	QL (120 GM per 28 days)
SYNALAR TOPICAL OINTMENT	T4	QL (120 GM per 28 days)
TACLONEX TOPICAL OINTMENT	T4	ST; QL (400 GM per 28 days)
TACLONEX TOPICAL SUSPENSION	T5	ST; QL (400 GM per 28 days)
<i>tacrolimus topical</i>	T4	QL (100 GM per 28 days)
TALTZ AUTOINJECTOR	T5	PA; QL (1 ML per 28 days)
TALTZ SYRINGE SUBCUTANEOUS SYRINGE 80 MG/ML	T5	PA; QL (1 ML per 28 days)
<i>tavaborole</i>	T4	QL (10 ML per 28 days)
<i>tazarotene topical cream</i>	T4	PA; QL (60 GM per 28 days)
<i>tazarotene topical foam</i>	T4	PA; QL (100 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>tazarotene topical gel</i>	T4	PA; QL (100 GM per 28 days)
TAZORAC TOPICAL CREAM	T4	PA; QL (60 GM per 28 days)
TAZORAC TOPICAL GEL	T4	PA; QL (100 GM per 28 days)
TEXACORT	T4	
TOPICORT TOPICAL CREAM	T4	QL (100 GM per 28 days)
TOPICORT TOPICAL GEL	T4	QL (60 GM per 28 days)
TOPICORT TOPICAL OINTMENT 0.05 %	T4	QL (100 GM per 28 days)
TOPICORT TOPICAL SPRAY, NON-AEROSOL	T4	QL (100 ML per 28 days)
TOVET EMOLLIENT	T4	QL (100 GM per 28 days)
TREMFYA	T5	PA; QL (1 ML per 56 days)
<i>tretinoin microspheres topical gel</i>	T4	PA; QL (50 GM per 28 days)
<i>tretinoin microspheres topical gel with pump 0.08 %</i>	T4	PA; QL (50 GM per 28 days)
<i>tretinoin topical cream</i>	T4	PA; QL (45 GM per 28 days)
<i>tretinoin topical gel</i>	T3	PA; QL (45 GM per 28 days)
<i>triamcinolone acetonide topical cream</i>	T2	
<i>triamcinolone acetonide topical lotion</i>	T2	
<i>triamcinolone acetonide topical ointment 0.025 %, 0.1 %, 0.5 %</i>	T2	
<i>triamcinolone acetonide topical ointment 0.05 %</i>	T4	
TRIDERM TOPICAL CREAM	T2	
TWYNEO	T4	PA; QL (30 GM per 28 days)
ULTRAVATE TOPICAL LOTION	T4	QL (120 ML per 28 days)
VALCHLOR	T5	PA-NS
VANOS	T5	QL (120 GM per 28 days)
VECTICAL	T4	ST
VELTIN	T4	PA; QL (60 GM per 28 days)
VERDESO	T4	QL (100 GM per 28 days)
VEREGEN	T5	QL (30 GM per 28 days)
VTAMA	T5	PA; QL (60 GM per 28 days)
WINLEVI	T4	PA; QL (60 GM per 28 days)
XERESE	T5	
ZENATANE	T4	
ZIANA	T4	PA; QL (60 GM per 28 days)
ZILXI	T4	
ZONALON	T4	PA; QL (30 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZORYVE TOPICAL CREAM 0.3 %	T4	PA; QL (60 GM per 28 days)
ZORYVE TOPICAL FOAM	T4	PA; QL (60 GM per 28 days)
ZOVIRAX TOPICAL CREAM	T4	QL (5 GM per 28 days)
ZOVIRAX TOPICAL OINTMENT	T4	QL (30 GM per 30 days)
ZTLIDO	T4	PA; QL (93 EA per 31 days)
ZYCLARA TOPICAL CREAM IN METERED-DOSE PUMP	T5	
Diagnostics / Miscellaneous Agents		
<i>acamprosate</i>	T4	
AGRYLIN	T4	
<i>anagrelide</i>	T2	
ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG	T5	PA
AURYXIA	T5	PA; QL (372 EA per 31 days)
BUPHENYL	T5	PA
<i>bupropion hcl (smoking deter)</i>	T2	QL (62 EA per 31 days)
CARBAGLU	T5	PA
<i>carglumic acid</i>	T5	PA
CARNITOR ORAL	T4	PA-BvD
<i>cevimeline</i>	T4	
CHEMET	T4	
CLINIMIX 4.25%/D5W SULFIT FREE	T4	PA-BvD
CLINIMIX E 2.75%/D5W SULF FREE	T4	PA-BvD
CUVRIOR	T5	PA; QL (310 EA per 31 days)
<i>d10 %-0.45 % sodium chloride</i>	T2	
<i>d2.5 %-0.45 % sodium chloride</i>	T2	
<i>d5 % and 0.9 % sodium chloride</i>	T2	
<i>d5 %-0.45 % sodium chloride</i>	T2	
<i>deferasirox oral granules in packet</i>	T5	PA
<i>deferasirox oral tablet 180 mg, 360 mg</i>	T5	PA
<i>deferasirox oral tablet 90 mg</i>	T4	PA
<i>deferasirox oral tablet, dispersible 125 mg</i>	T4	PA
<i>deferasirox oral tablet, dispersible 250 mg, 500 mg</i>	T5	PA
<i>deferiprone</i>	T5	PA
<i>dextrose 10 % and 0.2 % nacl</i>	T2	
<i>dextrose 10 % in water (d10w)</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>dextrose 5 % in water (d5w) intravenous piggyback</i>	T2	
<i>dextrose 5%-0.2 % sod chloride</i>	T2	
<i>disulfiram</i>	T2	
<i>droxidopa oral capsule 100 mg</i>	T5	PA; QL (465 EA per 31 days)
<i>droxidopa oral capsule 200 mg, 300 mg</i>	T5	PA; QL (186 EA per 31 days)
ENDARI	T5	PA; QL (180 EA per 30 days)
EVOXAC	T4	
EXJADE	T5	PA
EXSERVAN	T5	PA; QL (62 EA per 31 days)
FABHALTA	T5	PA; QL (62 EA per 31 days)
FERRIPROX	T5	PA
FERRIPROX (2 TIMES A DAY)	T5	PA
FOSRENOL ORAL POWDER IN PACKET	T5	
FOSRENOL ORAL TABLET,CHEWABLE 1,000 MG, 750 MG	T5	
FOSRENOL ORAL TABLET,CHEWABLE 500 MG	T4	
GLASSIA	T5	PA
INCRELEX	T5	PA
JADENU	T5	PA
JADENU SPRINKLE	T5	PA
JOENJA	T5	PA; QL (60 EA per 30 days)
KIONEX (WITH SORBITOL)	T2	
<i>lanthanum oral tablet,chewable 1,000 mg</i>	T5	
<i>lanthanum oral tablet,chewable 500 mg, 750 mg</i>	T4	
<i>levocarnitine (with sugar)</i>	T2	PA-BvD
<i>levocarnitine oral tablet</i>	T2	PA-BvD
LITFULO	T5	PA; QL (28 EA per 28 days)
LITHOSTAT	T4	
LOKELMA	T3	PA; QL (93 EA per 31 days)
<i>midodrine</i>	T2	
NICOTROL	T4	
NICOTROL NS	T5	
<i>nitisinone</i>	T5	PA
NITYR	T5	PA
NORTHERA ORAL CAPSULE 100 MG	T5	PA; QL (465 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
NORTHERA ORAL CAPSULE 200 MG, 300 MG	T5	PA; QL (186 EA per 31 days)
OLPRUVA ORAL PELLETS IN PACKET 2 GRAM, 3 GRAM	T5	PA; QL (186 EA per 31 days)
OLPRUVA ORAL PELLETS IN PACKET 4 GRAM	T5	PA; QL (155 EA per 31 days)
OLPRUVA ORAL PELLETS IN PACKET 5 GRAM	T5	PA; QL (124 EA per 31 days)
OLPRUVA ORAL PELLETS IN PACKET 6 GRAM	T5	PA; QL (93 EA per 31 days)
OLPRUVA ORAL PELLETS IN PACKET 6.67 GRAM	T5	PA; QL (62 EA per 31 days)
ORFADIN	T5	PA
OXBRYTA ORAL TABLET 300 MG	T5	PA; QL (248 EA per 31 days)
OXBRYTA ORAL TABLET 500 MG	T5	PA; QL (155 EA per 31 days)
OXBRYTA ORAL TABLET FOR SUSPENSION	T5	PA; QL (248 EA per 31 days)
PHEBURANE	T5	PA; QL (620 GM per 31 days)
<i>pilocarpine hcl oral</i>	T2	
PROLASTIN-C INTRAVENOUS SOLUTION	T5	PA
PYRUKYND ORAL TABLET 20 MG, 5 MG (4-WEEK PACK), 50 MG	T5	PA; QL (56 EA per 28 days)
PYRUKYND ORAL TABLET 5 MG	T5	PA; QL (14 EA per 365 days)
PYRUKYND ORAL TABLETS,DOSE PACK	T5	PA; QL (28 EA per 365 days)
RAVICTI	T5	PA
REVELA	T4	
REVCIVI	T5	
REZDIFFRA	T5	PA; QL (31 EA per 31 days)
<i>riluzole</i>	T3	
<i>risedronate oral tablet 30 mg</i>	T4	
SALAGEN (PILOCARPINE)	T4	
<i>sevelamer carbonate oral powder in packet</i>	T4	
<i>sevelamer carbonate oral tablet</i>	T3	
<i>sevelamer hcl</i>	T4	
<i>sodium chloride 0.9 % intravenous parenteral solution</i>	T2	
<i>sodium chloride irrigation</i>	T2	
<i>sodium phenylbutyrate</i>	T5	PA
<i>sodium polystyrene sulfonate oral powder</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 2.5 MG, 5 MG	T5	PA; QL (31 EA per 31 days)
SOHONOS ORAL CAPSULE 10 MG	T5	PA; QL (62 EA per 31 days)
SPS (WITH SORBITOL) ORAL	T2	
SYPRINE	T5	QL (248 EA per 31 days)
TAVNEOS	T5	PA; QL (186 EA per 31 days)
TEGLUTIK	T5	PA
THIOLA	T5	PA
THIOLA EC	T5	PA
<i>tiopronin oral tablet</i>	T5	PA
<i>trientine oral capsule 250 mg</i>	T5	QL (248 EA per 31 days)
<i>trientine oral capsule 500 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>varenicline oral tablet</i>	T4	QL (60 EA per 30 days)
<i>varenicline oral tablets,dose pack</i>	T4	QL (106 EA per 365 days)
VELPHORO	T5	
VELTASSA	T5	PA; QL (30 EA per 30 days)
WEGOVY SUBCUTANEOUS PEN INJECTOR 0.25 MG/0.5 ML, 0.5 MG/0.5 ML, 1 MG/0.5 ML	T5	PA; QL (2 ML per 28 days)
WEGOVY SUBCUTANEOUS PEN INJECTOR 1.7 MG/0.75 ML, 2.4 MG/0.75 ML	T5	PA; QL (3 ML per 28 days)
XURIDEN	T5	PA; QL (124 EA per 31 days)
ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG	T5	PA
ZOKINVY	T5	PA
Ear, Nose / Throat Medications		
<i>acetic acid otic (ear)</i>	T2	
<i>azelastine nasal spray,non-aerosol 137 mcg (0.1 %)</i>	T3	QL (30 ML per 25 days)
CETRAXAL	T4	
<i>chlorhexidine gluconate mucous membrane</i>	T1	
CIPRO HC	T4	
<i>ciprofloxacin hcl otic (ear)</i>	T4	
<i>ciprofloxacin-dexamethasone</i>	T3	
<i>ciprofloxacin-fluocinolone</i>	T4	
DERMOTIC OIL	T4	
FLAC OTIC OIL	T4	
<i>fluocinolone acetonide oil</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>hydrocortisone-acetic acid</i>	T2	
<i>ipratropium bromide nasal spray,non-aerosol 21 mcg (0.03 %)</i>	T2	QL (30 ML per 28 days)
<i>ipratropium bromide nasal spray,non-aerosol 42 mcg (0.06 %)</i>	T2	QL (15 ML per 28 days)
KOURZEQ	T2	
<i>neomycin-polymyxin-hc otic (ear)</i>	T2	
<i>ofloxacin otic (ear)</i>	T2	
<i>olopatadine nasal</i>	T4	QL (30.5 GM per 30 days)
OTOVEL	T4	
PERIOGARD	T1	
<i>triamcinolone acetonide dental</i>	T2	
Endocrine/Diabetes		
<i>acarbose</i>	T2	QL (93 EA per 31 days)
ACTHAR	T5	PA
ACTOPLUS MET ORAL TABLET 15-850 MG	T4	QL (93 EA per 31 days)
ACTOS	T4	QL (31 EA per 31 days)
ADMELOG SOLOSTAR U-100 INSULIN	T4	ST
ADMELOG U-100 INSULIN LISPRO	T4	ST
AFREZZA INHALATION CARTRIDGE WITH INHALER 12 UNIT, 4 UNIT (90)/ 8 UNIT (90), 4 UNIT/8 UNIT/ 12 UNIT (60), 8 UNIT (90)/ 12 UNIT (90)	T5	
AFREZZA INHALATION CARTRIDGE WITH INHALER 4 UNIT, 8 UNIT	T4	
AGAMREE	T5	PA; QL (300 ML per 40 days)
ALCOHOL PADS	T2	
ALKINDI SPRINKLE ORAL CAPSULE, SPRINKLE 0.5 MG	T4	PA
ALKINDI SPRINKLE ORAL CAPSULE, SPRINKLE 1 MG, 2 MG, 5 MG	T5	PA
<i>alogliptin</i>	T4	ST; QL (31 EA per 31 days)
<i>alogliptin-metformin</i>	T4	ST; QL (62 EA per 31 days)
<i>alogliptin-pioglitazone oral tablet 12.5-30 mg, 25-15 mg, 25-30 mg, 25-45 mg</i>	T4	ST; QL (31 EA per 31 days)
ANDROGEL TRANSDERMAL GEL IN METERED-DOSE PUMP	T4	PA
APIDRA SOLOSTAR U-100 INSULIN	T4	ST

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
APIDRA U-100 INSULIN	T4	ST
AVEED	T4	PA
BAQSIMI	T3	
BASAGLAR KWIKPEN U-100 INSULIN	T4	
BASAGLAR TEMPO PEN(U-100)INSLN	T4	
BYDUREON BCISE	T4	PA; QL (3.4 ML per 28 days)
BYETTA SUBCUTANEOUS PEN INJECTOR 10 MCG/DOSE(250 MCG/ML) 2.4 ML	T4	PA; QL (2.4 ML per 30 days)
BYETTA SUBCUTANEOUS PEN INJECTOR 5 MCG/DOSE (250 MCG/ML) 1.2 ML	T4	PA; QL (1.2 ML per 30 days)
<i>cabergoline</i>	T3	
<i>calcitonin (salmon) nasal</i>	T2	PA-BvD
<i>calcitriol oral</i>	T2	PA-BvD
CERDELGA	T5	PA; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 30 mg, 60 mg</i>	T4	PA-BvD; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 90 mg</i>	T4	PA-BvD; QL (124 EA per 31 days)
CORTEF	T4	
CORTROPHIN GEL	T5	PA
CYCLOSET	T4	
CYTOMEL	T4	
<i>danazol</i>	T4	
<i>dapaglifloz propaned-metformin oral tablet, ir - er, biphasic 24hr 10-1,000 mg</i>	T4	QL (31 EA per 31 days)
<i>dapaglifloz propaned-metformin oral tablet, ir - er, biphasic 24hr 5-1,000 mg</i>	T4	QL (62 EA per 31 days)
<i>dapagliflozin propanediol</i>	T4	QL (31 EA per 31 days)
DDAVP ORAL	T4	
<i>deflazacort</i>	T5	PA
DEPO-TESTOSTERONE	T4	PA
<i>desmopressin nasal spray,non-aerosol 10 mcg/spray (0.1 ml)</i>	T3	
<i>desmopressin oral</i>	T3	
DEXABLISS	T4	
<i>dexamethasone oral solution</i>	T2	
<i>dexamethasone oral tablet</i>	T1	
<i>dexamethasone oral tablets,dose pack</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>diazoxide</i>	T4	
<i>doxercalciferol oral</i>	T4	PA-BvD
DUETACT	T4	QL (31 EA per 31 days)
EMFLAZA	T5	PA
ERMEZA	T4	
EUTHYROX	T3	
FARXIGA	T3	QL (31 EA per 31 days)
FIASP FLEXTOUCH U-100 INSULIN	T4	ST
FIASP PENFILL U-100 INSULIN	T4	ST
FIASP U-100 INSULIN	T4	ST
<i>fludrocortisone</i>	T1	
GALAFOLD	T5	PA; QL (14 EA per 28 days)
<i>glimepiride</i>	T1	
<i>glipizide oral tablet 10 mg, 5 mg</i>	T1	
<i>glipizide oral tablet 2.5 mg</i>	T3	
<i>glipizide oral tablet extended release 24hr</i>	T1	
<i>glipizide-metformin</i>	T1	
GLUCAGON EMERGENCY KIT (HUMAN)	T4	
GLUCOTROL XL ORAL TABLET EXTENDED RELEASE 24HR 10 MG, 5 MG	T4	
GLUMETZA ORAL TABLET,ER GAST.RETENTION 24 HR 1,000 MG	T5	ST; QL (62 EA per 31 days)
GLUMETZA ORAL TABLET,ER GAST.RETENTION 24 HR 500 MG	T5	ST; QL (31 EA per 31 days)
<i>glyburide</i>	T4	
<i>glyburide micronized</i>	T4	
<i>glyburide-metformin</i>	T4	
GLYXAMBI	T3	QL (31 EA per 31 days)
GVOKE	T3	
GVOKE HYPOPEN 2-PACK	T3	
GVOKE PFS 1-PACK SYRINGE SUBCUTANEOUS SYRINGE 1 MG/0.2 ML	T3	
HEMADY	T4	PA-NS
HUMALOG JUNIOR KWIKPEN U-100	T3	
HUMALOG KWIKPEN INSULIN	T3	
HUMALOG MIX 50-50 KWIKPEN	T3	
HUMALOG MIX 75-25 KWIKPEN	T3	
HUMALOG MIX 75-25(U-100)INSULN	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
HUMALOG TEMPO PEN(U-100)INSULN	T3	
HUMALOG U-100 INSULIN	T3	
HUMULIN 70/30 U-100 INSULIN	T3	
HUMULIN 70/30 U-100 KWIKPEN	T3	
HUMULIN N NPH INSULIN KWIKPEN	T3	
HUMULIN N NPH U-100 INSULIN	T3	
HUMULIN R REGULAR U-100 INSULN	T3	
HUMULIN R U-500 (CONC) INSULIN	T3	
HUMULIN R U-500 (CONC) KWIKPEN	T3	
<i>hydrocortisone oral</i>	T2	
INPEFA ORAL TABLET 200 MG	T4	QL (62 EA per 31 days)
INPEFA ORAL TABLET 400 MG	T4	QL (31 EA per 31 days)
<i>insulin asp prt-insulin aspart</i>	T4	ST
<i>insulin aspart u-100</i>	T4	ST
<i>insulin degludec</i>	T4	
<i>insulin glargine u-300 conc</i>	T4	
<i>insulin glargine-yfgn</i>	T4	
<i>insulin lispro</i>	T3	
<i>insulin lispro protamin-lispro</i>	T3	
INVOKAMET	T4	QL (62 EA per 31 days)
INVOKAMET XR	T4	QL (62 EA per 31 days)
INVOKANA ORAL TABLET 100 MG	T4	QL (62 EA per 31 days)
INVOKANA ORAL TABLET 300 MG	T4	QL (31 EA per 31 days)
ISTURISA ORAL TABLET 1 MG	T5	PA; QL (248 EA per 31 days)
ISTURISA ORAL TABLET 5 MG	T5	PA; QL (62 EA per 31 days)
JANUMET	T3	QL (62 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 100-1,000 MG, 50-500 MG	T3	QL (31 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 50-1,000 MG	T3	QL (62 EA per 31 days)
JANUVIA ORAL TABLET 100 MG, 50 MG	T3	QL (31 EA per 31 days)
JANUVIA ORAL TABLET 25 MG	T3	QL (93 EA per 31 days)
JARDIANCE ORAL TABLET 10 MG	T3	QL (62 EA per 31 days)
JARDIANCE ORAL TABLET 25 MG	T3	QL (31 EA per 31 days)
JATENZO ORAL CAPSULE 158 MG, 198 MG	T4	PA; QL (124 EA per 31 days)
JATENZO ORAL CAPSULE 237 MG	T4	PA; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
JAVYGTOR	T5	PA
JENTADUETO ORAL TABLET 2.5-1,000 MG, 2.5-500 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 5-1,000 MG	T3	QL (31 EA per 31 days)
JYNARQUE ORAL TABLET	T5	PA; QL (112 EA per 28 days)
JYNARQUE ORAL TABLETS, SEQUENTIAL	T5	PA; QL (56 EA per 28 days)
KAZANO	T4	ST; QL (62 EA per 31 days)
KORLYM	T5	PA; QL (124 EA per 31 days)
KUVAN	T5	PA
LANTUS SOLOSTAR U-100 INSULIN	T3	
LANTUS U-100 INSULIN	T3	
LEVEMIR FLEXPEN	T4	
LEVEMIR U-100 INSULIN	T4	
<i>levothyroxine oral capsule</i>	T4	
<i>levothyroxine oral tablet</i>	T1	
LEVOXYL ORAL TABLET 100 MCG, 112 MCG, 125 MCG, 137 MCG, 150 MCG, 175 MCG, 200 MCG, 25 MCG, 50 MCG, 75 MCG, 88 MCG	T3	
<i>liothyronine oral</i>	T2	
LYUMJEV KWIKPEN U-100 INSULIN	T4	ST
LYUMJEV KWIKPEN U-200 INSULIN	T4	ST
LYUMJEV TEMPO PEN(U-100)INSULN	T4	ST
LYUMJEV U-100 INSULIN	T4	ST
MEDROL (PAK)	T4	
MEDROL ORAL TABLET 16 MG, 2 MG, 4 MG, 8 MG	T4	
<i>metformin oral solution</i>	T4	ST; QL (791 ML per 31 days)
<i>metformin oral tablet 1,000 mg, 500 mg, 850 mg</i>	T1	
<i>metformin oral tablet 625 mg</i>	T4	QL (124 EA per 31 days)
<i>metformin oral tablet extended release 24 hr</i>	T1	
<i>metformin oral tablet extended release 24hr 1,000 mg</i>	T4	ST; QL (62 EA per 31 days)
<i>metformin oral tablet extended release 24hr 500 mg</i>	T4	ST; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>metformin oral tablet,er gast.retention 24 hr 1,000 mg</i>	T5	ST; QL (62 EA per 31 days)
<i>metformin oral tablet,er gast.retention 24 hr 500 mg</i>	T4	ST; QL (31 EA per 31 days)
<i>methimazole oral tablet 10 mg, 5 mg</i>	T1	
METHITEST	T5	PA
<i>methylprednisolone</i>	T2	
<i>methyltestosterone oral capsule</i>	T5	PA
<i>mifepristone oral tablet 300 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>miglitol</i>	T4	
<i>miglustat</i>	T5	PA; QL (93 EA per 31 days)
MOUNJARO	T3	PA; QL (2 ML per 28 days)
MYALEPT	T5	PA
<i>nateglinide</i>	T2	QL (93 EA per 31 days)
NATESTO	T4	PA
NESINA	T4	ST; QL (31 EA per 31 days)
NOVOLIN 70/30 U-100 INSULIN	T4	ST
NOVOLIN 70-30 FLEXPEN U-100	T4	ST
NOVOLIN N FLEXPEN	T4	ST
NOVOLIN N NPH U-100 INSULIN	T4	ST
NOVOLIN R FLEXPEN	T4	ST
NOVOLIN R REGULAR U100 INSULIN	T4	ST
NOVOLOG FLEXPEN U-100 INSULIN	T4	ST
NOVOLOG MIX 70-30 U-100 INSULN	T4	ST
NOVOLOG MIX 70-30FLEXPEN U-100	T4	ST
NOVOLOG PENFILL U-100 INSULIN	T4	ST
NOVOLOG U-100 INSULIN ASPART	T4	ST
ORAPRED ODT	T4	
ORILISSA ORAL TABLET 150 MG	T5	PA; QL (31 EA per 31 days)
ORILISSA ORAL TABLET 200 MG	T5	PA; QL (62 EA per 31 days)
OSENI ORAL TABLET 12.5-30 MG, 25-15 MG, 25-30 MG, 25-45 MG	T4	ST; QL (31 EA per 31 days)
OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)	T3	PA; QL (3 ML per 28 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (15 ML per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
PALYNZIQ SUBCUTANEOUS SYRINGE 2.5 MG/0.5 ML	T5	PA; QL (4 ML per 30 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (90 ML per 30 days)
<i>paricalcitol oral</i>	T4	PA-BvD
<i>pioglitazone</i>	T1	QL (31 EA per 31 days)
<i>pioglitazone-glimepiride</i>	T4	QL (31 EA per 31 days)
<i>pioglitazone-metformin</i>	T4	QL (93 EA per 31 days)
<i>prednisolone oral solution</i>	T2	
<i>prednisolone oral tablet</i>	T4	
<i>prednisolone sodium phosphate oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)</i>	T4	
<i>prednisolone sodium phosphate oral solution 25 mg/5 ml (5 mg/ml), 5 mg base/5 ml (6.7 mg/5 ml)</i>	T2	
<i>prednisolone sodium phosphate oral tablet, disintegrating</i>	T4	
PREDNISON INTENSOL	T2	
<i>prednisone oral solution</i>	T2	
<i>prednisone oral tablet</i>	T1	
<i>prednisone oral tablets, dose pack</i>	T1	
PROGLYCEM	T4	
<i>propylthiouracil</i>	T2	
QTERN	T3	QL (31 EA per 31 days)
RAYALDEE	T5	QL (62 EA per 31 days)
RAYOS	T5	
RECORLEV	T5	PA; QL (248 EA per 31 days)
<i>repaglinide oral tablet 0.5 mg, 1 mg</i>	T2	QL (124 EA per 31 days)
<i>repaglinide oral tablet 2 mg</i>	T2	QL (248 EA per 31 days)
REZVOGLAR KWIKPEN	T4	
ROCALTROL	T4	PA-BvD
RYBELSUS	T3	PA; QL (31 EA per 31 days)
SAMSCA	T5	PA
<i>sapropterin</i>	T5	PA
<i>saxagliptin</i>	T4	ST; QL (31 EA per 31 days)
<i>saxagliptin-metformin oral tablet, er multiphase 24 hr 2.5-1,000 mg</i>	T4	ST; QL (62 EA per 31 days)
<i>saxagliptin-metformin oral tablet, er multiphase 24 hr 5-1,000 mg, 5-500 mg</i>	T4	ST; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SEGLUROMET	T4	QL (62 EA per 31 days)
SEMGLEE(INSULIN GLARGINE-YFGN)	T4	
SEMGLEE(INSULIN GLARG-YFGN)PEN	T4	
SENSIPAR ORAL TABLET 30 MG, 60 MG	T4	PA-BvD; QL (62 EA per 31 days)
SENSIPAR ORAL TABLET 90 MG	T5	PA-BvD; QL (124 EA per 31 days)
<i>sitagliptin oral tablet 100 mg, 50 mg</i>	T4	ST; QL (31 EA per 31 days)
<i>sitagliptin oral tablet 25 mg</i>	T4	ST; QL (93 EA per 31 days)
<i>sitagliptin-metformin</i>	T4	ST; QL (62 EA per 31 days)
SOLQUA 100/33	T3	QL (18 ML per 30 days)
SOMAVERT	T5	PA
STEGLATRO	T3	QL (31 EA per 31 days)
STEGLUJAN	T4	ST; QL (31 EA per 31 days)
SYMLINPEN 120	T5	QL (10.8 ML per 28 days)
SYMLINPEN 60	T5	QL (6 ML per 28 days)
SYNAREL	T5	PA
SYNJARDY	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 12.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 25-1,000 MG	T3	QL (31 EA per 31 days)
SYNTHROID	T4	
TAPERDEX	T4	
TARPEYO	T5	PA; QL (124 EA per 31 days)
TESTIM	T4	PA
<i>testosterone cypionate</i>	T3	PA
<i>testosterone enanthate</i>	T3	PA
<i>testosterone transdermal gel in metered-dose pump 10 mg/0.5 gram /actuation, 20.25 mg/1.25 gram (1.62 %)</i>	T3	PA
<i>testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %)</i>	T4	PA
<i>testosterone transdermal gel in packet</i>	T3	PA
<i>testosterone transdermal solution in metered pump w/app</i>	T3	PA
THYQUIDITY	T4	
TIROSINT	T4	
TIROSINT-SOL	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TLANDO	T4	PA; QL (124 EA per 31 days)
<i>tolvaptan</i>	T5	PA
TOUJEO MAX U-300 SOLOSTAR	T3	
TOUJEO SOLOSTAR U-300 INSULIN	T3	
TRADJENTA	T3	QL (31 EA per 31 days)
TRESIBA FLEXTOUCH U-100	T4	
TRESIBA FLEXTOUCH U-200	T4	
TRESIBA U-100 INSULIN	T4	
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-5-1,000 MG, 25-5-1,000 MG	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 12.5-2.5-1,000 MG, 5-2.5-1,000 MG	T3	QL (62 EA per 31 days)
TRULICITY	T3	PA; QL (2 ML per 28 days)
UNITHROID	T3	
VICTOZA 3-PAK	T3	PA; QL (9 ML per 30 days)
VOGELXO TRANSDERMAL GEL	T4	PA
VOGELXO TRANSDERMAL GEL IN METERED-DOSE PUMP	T4	PA
VOXZOGO	T5	PA; QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 10-500 MG, 5-500 MG	T3	QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
XULTOPHY 100/3.6	T3	QL (15 ML per 30 days)
XYOSTED	T4	PA
YARGESA	T5	PA; QL (93 EA per 31 days)
ZAVESCA	T5	PA; QL (93 EA per 31 days)
ZEGALOGUE AUTOINJECTOR	T4	
ZEGALOGUE SYRINGE	T4	
ZEMPLAR ORAL CAPSULE 1 MCG, 2 MCG	T4	PA-BvD
ZITUVIO ORAL TABLET 100 MG, 50 MG	T4	ST; QL (31 EA per 31 days)
ZITUVIO ORAL TABLET 25 MG	T4	ST; QL (93 EA per 31 days)
Gastroenterology		
ACIPHEX	T4	QL (62 EA per 31 days)
<i>alosetron oral tablet 0.5 mg</i>	T5	PA; QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>alosetron oral tablet 1 mg</i>	T5	PA; QL (62 EA per 31 days)
AMITIZA	T4	QL (62 EA per 31 days)
<i>amoxicil-clarithromy-lansopraz</i>	T4	
ANTIVERT ORAL TABLET 50 MG	T4	
ANTIVERT ORAL TABLET,CHEWABLE	T4	
ANUSOL-HC TOPICAL	T4	
ANZEMET ORAL TABLET 50 MG	T4	PA-BvD
<i>aprepitant</i>	T4	PA-BvD
APRISO	T4	QL (124 EA per 31 days)
AZULFIDINE	T4	
AZULFIDINE EN-TABS	T4	
<i>balsalazide</i>	T2	
<i>betaine</i>	T5	
<i>bismuth subcit k-metronidz-tcn</i>	T4	
BONJESTA	T4	PA; QL (62 EA per 31 days)
<i>budesonide oral capsule,delayed,extend.release</i>	T4	
<i>budesonide oral tablet,delayed and ext.release</i>	T5	
<i>budesonide rectal</i>	T4	
BYLVAY ORAL CAPSULE 1,200 MCG	T5	PA; QL (186 EA per 31 days)
BYLVAY ORAL CAPSULE 400 MCG	T5	PA; QL (558 EA per 31 days)
BYLVAY ORAL PELLETT 200 MCG	T5	PA; QL (1116 EA per 31 days)
BYLVAY ORAL PELLETT 600 MCG	T5	PA; QL (372 EA per 31 days)
CANASA	T5	QL (31 EA per 31 days)
CARAFATE	T4	
CHENODAL	T5	PA
<i>chlordiazepoxide-clidinium</i>	T4	
CHOLBAM	T5	PA
<i>cimetidine</i>	T2	
CIMZIA	T5	PA; QL (2 EA per 28 days)
CIMZIA POWDER FOR RECONST	T5	PA; QL (2 EA per 28 days)
CLENPIQ	T4	
COLAZAL	T5	
COMPRO	T2	
CONSTULOSE	T2	
CREON	T3	
<i>cromolyn oral</i>	T4	
CUVPOSA	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
CYSTADANE	T5	
CYTOTEC	T4	
DELZICOL	T4	QL (186 EA per 31 days)
DEXILANT	T4	QL (31 EA per 31 days)
<i>dexlansoprazole</i>	T4	QL (31 EA per 31 days)
DICLEGIS	T4	PA; QL (124 EA per 31 days)
<i>dicyclomine oral capsule</i>	T2	
<i>dicyclomine oral solution</i>	T2	
<i>dicyclomine oral tablet</i>	T2	
DIPENTUM	T5	
<i>diphenoxylate-atropine</i>	T2	
<i>doxylamine-pyridoxine (vit b6)</i>	T4	PA; QL (124 EA per 31 days)
<i>dronabinol</i>	T4	PA-BvD
EMEND ORAL CAPSULE 80 MG	T4	PA-BvD
EMEND ORAL CAPSULE,DOSE PACK	T4	PA-BvD
EMEND ORAL SUSPENSION FOR RECONSTITUTION	T4	PA-BvD
ENTYVIO PEN	T5	PA; QL (1.36 ML per 28 days)
ENULOSE	T2	
<i>esomeprazole magnesium oral capsule, delayed release(dr/ec)</i>	T2	QL (31 EA per 31 days)
<i>esomeprazole magnesium oral granules dr for susp in packet</i>	T4	QL (31 EA per 31 days)
<i>famotidine oral suspension for reconstitution</i>	T2	
<i>famotidine oral tablet 20 mg, 40 mg</i>	T1	
GASTROCROM	T4	
GATTEX 30-VIAL	T5	PA
GAVILYTE-C	T2	
GAVILYTE-G	T2	
GENERLAC	T2	
GIMOTI	T5	PA; QL (9.8 ML per 28 days)
GLYCATE	T4	PA; QL (155 EA per 31 days)
<i>glycopyrrolate oral solution</i>	T4	
<i>glycopyrrolate oral tablet 1 mg, 2 mg</i>	T3	
<i>glycopyrrolate oral tablet 1.5 mg</i>	T4	PA; QL (155 EA per 31 days)
GOLYTELY	T4	
<i>granisetron hcl oral</i>	T2	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
HELIDAC	T4	
<i>hydrocortisone rectal</i>	T4	
<i>hydrocortisone topical cream with perineal applicator 2.5 %</i>	T2	
<i>hydrocortisone-pramoxine rectal cream 1-1 %</i>	T4	
IBSRELA	T5	PA; QL (62 EA per 31 days)
KONVOMEF	T4	
KRISTALOSE	T4	ST
<i>lactulose oral packet</i>	T5	ST
<i>lactulose oral solution 10 gram/15 ml</i>	T2	
<i>lansoprazole oral capsule, delayed release(dr/ec) 15 mg</i>	T2	QL (31 EA per 31 days)
<i>lansoprazole oral capsule, delayed release(dr/ec) 30 mg</i>	T2	QL (62 EA per 31 days)
<i>lansoprazole oral tablet, disintegrat, delay rel 15 mg</i>	T4	QL (31 EA per 31 days)
<i>lansoprazole oral tablet, disintegrat, delay rel 30 mg</i>	T4	QL (62 EA per 31 days)
LIALDA	T4	QL (124 EA per 31 days)
LIBRAX (WITH CLIDINIUM)	T4	
LINZESS	T3	QL (31 EA per 31 days)
LIVMARLI ORAL SOLUTION 9.5 MG/ML	T5	PA; QL (93 ML per 31 days)
LOMOTIL	T4	
<i>loperamide oral capsule</i>	T2	
LOTRONEX ORAL TABLET 0.5 MG	T5	PA; QL (93 EA per 31 days)
LOTRONEX ORAL TABLET 1 MG	T5	PA; QL (62 EA per 31 days)
<i>lubiprostone</i>	T4	QL (62 EA per 31 days)
MARINOL ORAL CAPSULE 10 MG, 5 MG	T5	PA-BvD
MARINOL ORAL CAPSULE 2.5 MG	T4	PA-BvD
<i>meclizine oral tablet 12.5 mg, 25 mg</i>	T2	
<i>mesalamine oral capsule (with del rel tablets)</i>	T4	QL (186 EA per 31 days)
<i>mesalamine oral capsule, extended release</i>	T4	QL (248 EA per 31 days)
<i>mesalamine oral capsule, extended release 24hr</i>	T4	QL (124 EA per 31 days)
<i>mesalamine oral tablet, delayed release (dr/ec) 1.2 gram</i>	T4	QL (124 EA per 31 days)
<i>mesalamine oral tablet, delayed release (dr/ec) 800 mg</i>	T4	QL (186 EA per 31 days)
<i>mesalamine rectal enema</i>	T4	QL (1860 ML per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>mesalamine rectal suppository</i>	T4	QL (31 EA per 31 days)
<i>methscopolamine</i>	T4	
<i>metoclopramide hcl oral solution</i>	T2	
<i>metoclopramide hcl oral tablet</i>	T1	
<i>metoclopramide hcl oral tablet, disintegrating 5 mg</i>	T4	
<i>misoprostol</i>	T3	
MOTEGRITY	T4	PA; QL (31 EA per 31 days)
MOVANTI	T3	QL (31 EA per 31 days)
MOVIPREP	T4	
MYTESI	T5	QL (62 EA per 31 days)
NEXIUM	T4	QL (31 EA per 31 days)
NEXIUM PACKET	T4	QL (31 EA per 31 days)
<i>nitroglycerin rectal</i>	T3	
<i>nizatidine oral capsule</i>	T2	
OCALIVA	T5	PA; QL (31 EA per 31 days)
<i>omeprazole oral capsule, delayed release (dr/ec)</i>	T1	
<i>omeprazole-sodium bicarbonate oral capsule</i>	T4	
<i>omeprazole-sodium bicarbonate oral packet</i>	T5	
OMVOH PEN	T5	PA; QL (2 ML per 28 days)
OMVOH SUBCUTANEOUS	T5	PA; QL (2 ML per 28 days)
<i>ondansetron hcl oral solution</i>	T2	PA-BvD
<i>ondansetron hcl oral tablet 4 mg, 8 mg</i>	T2	PA-BvD
<i>ondansetron oral tablet, disintegrating 4 mg, 8 mg</i>	T2	PA-BvD
PANCREAZE ORAL CAPSULE, DELAYED RELEASE (DR/EC) 10,500-35,500- 61,500 UNIT, 16,800-56,800- 98,400 UNIT, 2,600-8,800- 15,200 UNIT, 21,000-54,700- 83,900 UNIT, 37,000-97,300- 149,900 UNIT, 4,200-14,200- 24,600 UNIT	T4	
<i>pantoprazole oral granules dr for susp in packet</i>	T4	
<i>pantoprazole oral tablet, delayed release (dr/ec)</i>	T1	
<i>peg 3350-electrolytes</i>	T2	
<i>peg3350-sod sul-nacl-kcl-asb-c</i>	T4	
<i>peg-electrolyte soln</i>	T2	
PENTASA ORAL CAPSULE, EXTENDED RELEASE 250 MG	T3	QL (496 EA per 31 days)
PENTASA ORAL CAPSULE, EXTENDED RELEASE 500 MG	T5	QL (248 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
PEPCID ORAL TABLET	T4	
PERTZYE	T4	
PLENVU	T4	
PREVACID	T4	QL (62 EA per 31 days)
PREVACID SOLUTAB ORAL TABLET,DISINTEGRAT, DELAY REL 15 MG	T4	QL (31 EA per 31 days)
PREVACID SOLUTAB ORAL TABLET,DISINTEGRAT, DELAY REL 30 MG	T4	QL (62 EA per 31 days)
PRILOSEC ORAL SUSP,DELAYED RELEASE FOR RECON	T4	
<i>prochlorperazine</i>	T2	
<i>prochlorperazine maleate</i>	T2	
PROCTOFOAM HC	T4	
PROCTO-MED HC	T2	
PROCTOSOL HC TOPICAL	T2	
PROCTOZONE-HC	T2	
PROTONIX ORAL	T4	
PYLERA	T4	
<i>rabeprazole oral tablet, delayed release (dr/ec)</i>	T4	QL (62 EA per 31 days)
RECTIV	T3	
REGLAN ORAL	T4	
RELISTOR ORAL	T5	PA; QL (93 EA per 31 days)
RELISTOR SUBCUTANEOUS SOLUTION	T5	PA; QL (18.6 ML per 31 days)
RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML	T5	PA; QL (18.6 ML per 31 days)
RELISTOR SUBCUTANEOUS SYRINGE 8 MG/0.4 ML	T5	PA; QL (12.4 ML per 31 days)
RELTONE	T5	ST
ROBINUL FORTE	T4	
ROBINUL ORAL	T4	
ROWASA RECTAL ENEMA KIT	T4	QL (31 EA per 31 days)
<i>scopolamine base</i>	T3	QL (10 EA per 30 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML)	T5	PA; QL (1.2 ML per 56 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 360 MG/2.4 ML (150 MG/ML)	T5	PA; QL (2.4 ML per 56 days)
<i>sodium,potassium,mag sulfates</i>	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SUCRAID	T5	PA
<i>sucralfate oral suspension</i>	T4	
<i>sucralfate oral tablet</i>	T2	
SUFLAVE	T4	
<i>sulfasalazine</i>	T2	
SUPREP BOWEL PREP KIT	T4	
SUTAB	T4	
SYMPROIC	T4	PA; QL (31 EA per 31 days)
SYNDROS	T5	PA
TALICIA	T4	
TRANSDERM-SCOP	T4	QL (10 EA per 30 days)
<i>trimethobenzamide oral</i>	T4	PA
TRULANCE	T3	QL (31 EA per 31 days)
UCERIS ORAL	T5	
UCERIS RECTAL	T4	
URSO 250	T4	ST
URSO FORTE	T4	ST
<i>ursodiol oral capsule 200 mg, 400 mg</i>	T4	ST
<i>ursodiol oral capsule 300 mg</i>	T3	
<i>ursodiol oral tablet</i>	T3	
VARUBI	T4	PA-BvD
VELSIPITY	T5	PA; QL (31 EA per 31 days)
VIBERZI	T5	PA; QL (62 EA per 31 days)
VIOKACE ORAL TABLET 10,440-39,150-39,150 UNIT	T4	
VIOKACE ORAL TABLET 20,880-78,300-78,300 UNIT	T5	
VOQUEZNA DUAL PAK	T4	PA; QL (112 EA per 14 days)
VOQUEZNA ORAL TABLET 10 MG	T4	PA; QL (31 EA per 31 days)
VOQUEZNA ORAL TABLET 20 MG	T4	PA; QL (62 EA per 31 days)
VOQUEZNA TRIPLE PAK	T4	PA; QL (112 EA per 14 days)
VOWST	T5	PA; QL (12 EA per 14 days)
ZEGERID ORAL CAPSULE	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZENPEP ORAL CAPSULE,DELAYED RELEASE(DR/EC) 10,000-32,000 -42,000 UNIT, 15,000-47,000 -63,000 UNIT, 20,000-63,000- 84,000 UNIT, 25,000-79,000- 105,000 UNIT, 3,000-10,000 -14,000-UNIT, 40,000-126,000- 168,000 UNIT, 5,000-17,000- 24,000 UNIT, 60,000-189,600- 252,600 UNIT	T3	
ZYMFENTRA	T5	PA; QL (1 EA per 28 days)
Immunology, Vaccines / Biotechnology		
ABRYSVO (PF)	T3	QL (2 EA per 365 days)
ACTHIB (PF)	T3	
ACTIMMUNE	T5	PA
ADACEL(TDAP ADOLESN/ADULT)(PF)	T3	
ARANESP (IN POLYSORBATE) INJECTION SOLUTION 100 MCG/ML, 200 MCG/ML	T5	PA-BvD; ST
ARANESP (IN POLYSORBATE) INJECTION SOLUTION 25 MCG/ML, 40 MCG/ML, 60 MCG/ML	T4	PA-BvD; ST
ARANESP (IN POLYSORBATE) INJECTION SYRINGE 10 MCG/0.4 ML, 25 MCG/0.42 ML, 40 MCG/0.4 ML	T4	PA-BvD; ST
ARANESP (IN POLYSORBATE) INJECTION SYRINGE 100 MCG/0.5 ML, 150 MCG/0.3 ML, 200 MCG/0.4 ML, 300 MCG/0.6 ML, 500 MCG/ML, 60 MCG/0.3 ML	T5	PA-BvD; ST
ARCALYST	T5	PA
AREXVY (PF)	T3	QL (2 EA per 365 days)
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	T5	PA; QL (4 EA per 28 days)
AVONEX INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (4 EA per 28 days)
<i>bcg vaccine, live (pf)</i>	T3	
BESREMI	T5	PA-NS; QL (2 ML per 28 days)
BETASERON SUBCUTANEOUS KIT	T5	PA; QL (14 EA per 28 days)
BEXSERO	T3	
BIVIGAM	T5	PA
BOOSTRIX TDAP	T3	
DAPTACEL (DTAP PEDIATRIC) (PF)	T3	
EGRIFTA SV	T5	PA
ENGERIX-B (PF)	T3	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ENGERIX-B PEDIATRIC (PF)	T3	PA-BvD
EPOGEN INJECTION SOLUTION 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML	T4	PA-BvD; ST
FULPHILA	T5	
FYLNETRA	T5	ST
GAMMAGARD LIQUID	T5	PA
GAMMAGARD S-D (IGA < 1 MCG/ML)	T5	PA
GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GAMMAPLEX	T5	PA
GAMMAPLEX (WITH SORBITOL)	T5	PA
GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GARDASIL 9 (PF)	T3	
GENOTROPIN	T5	PA
GENOTROPIN MINIQUICK SUBCUTANEOUS SYRINGE 0.2 MG/0.25 ML	T4	PA
GENOTROPIN MINIQUICK SUBCUTANEOUS SYRINGE 0.4 MG/0.25 ML, 0.6 MG/0.25 ML, 0.8 MG/0.25 ML, 1 MG/0.25 ML, 1.2 MG/0.25 ML, 1.4 MG/0.25 ML, 1.6 MG/0.25 ML, 1.8 MG/0.25 ML, 2 MG/0.25 ML	T5	PA
GRANIX	T5	ST
GRASTEK	T4	PA
HAVRIX (PF)	T3	
HEPLISA V-B (PF)	T3	PA-BvD
HIBERIX (PF)	T3	
HUMATROPE INJECTION CARTRIDGE 12 MG (36 UNIT), 24 MG (72 UNIT)	T5	PA
HUMATROPE INJECTION CARTRIDGE 6 MG (18 UNIT)	T4	PA
IMOVAX RABIES VACCINE (PF)	T3	PA-BvD
INFANRIX (DTAP) (PF)	T3	
IPOL	T3	
IXCHIQ (PF)	T3	
IXIARO (PF)	T3	
JYNNEOS (PF)	T3	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
KINRIX (PF)	T3	
LEUKINE INJECTION RECON SOLN	T5	PA
MENACTRA (PF) INTRAMUSCULAR SOLUTION	T3	
MENQUADFI (PF)	T3	
MENVEO A-C-Y-W-135-DIP (PF) INTRAMUSCULAR KIT	T3	
M-M-R II (PF)	T3	
NEULASTA	T5	
NEUPOGEN	T5	ST
NGENLA	T5	PA
NIVESTYM	T5	
NORDITROPIN FLEXPRO	T5	PA
NUTROPIN AQ NUSPIN	T5	PA
NYVEPRIA	T5	ST
OCTAGAM	T5	PA
ODACTRA	T4	PA
OMNITROPE SUBCUTANEOUS CARTRIDGE 10 MG/1.5 ML (6.7 MG/ML)	T5	PA
OMNITROPE SUBCUTANEOUS CARTRIDGE 5 MG/1.5 ML (3.3 MG/ML)	T4	PA
OMNITROPE SUBCUTANEOUS RECON SOLN	T5	PA
ORALAIR SUBLINGUAL TABLET 300 INDX REACTIVITY	T4	PA
PANZYGA	T5	PA
PEDIARIX (PF)	T3	PA-BvD
PEDVAX HIB (PF)	T3	
PEGASYS	T5	PA
PENBRAYA (PF)	T3	
PENTACEL (PF) INTRAMUSCULAR KIT 15LF-48MCG-62DU -10 MCG/0.5ML	T3	
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PREHEVBRIO (PF)	T3	PA-BvD
PRIORIX (PF)	T3	
PRIVIGEN	T5	PA

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML	T3	PA-BvD
PROCRIT INJECTION SOLUTION 20,000 UNIT/ML, 40,000 UNIT/ML	T5	PA-BvD
PROQUAD (PF)	T3	
QUADRACEL (PF)	T3	
RABAVERT (PF)	T3	PA-BvD
REBIF (WITH ALBUMIN)	T5	PA; QL (6 ML per 28 days)
REBIF REBIDOSE SUBCUTANEOUS PEN INJECTOR 22 MCG/0.5 ML, 44 MCG/0.5 ML	T5	PA; QL (6 ML per 28 days)
REBIF REBIDOSE SUBCUTANEOUS PEN INJECTOR 8.8MCG/0.2ML-22 MCG/0.5ML (6)	T5	PA; QL (4.2 ML per 365 days)
REBIF TITRATION PACK	T5	PA; QL (8.4 ML per 365 days)
RECOMBIVAX HB (PF)	T3	PA-BvD
RELEUKO SUBCUTANEOUS	T5	ST
RETACRIT	T3	PA-BvD
ROTARIX	T3	
ROTATEQ VACCINE	T3	
SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG	T5	PA
SHINGRIX (PF)	T3	QL (2 EA per 999 days)
SKYTROFA	T5	PA
SOGROYA	T5	PA
STIMUFEND	T5	ST
TDVAX	T3	
TENIVAC (PF)	T3	
<i>tetanus, diphtheria tox ped(pf)</i>	T3	
TICOVAC	T3	
TRUMENBA	T3	
TWINRIX (PF)	T3	
TYPHIM VI	T3	
UDENYCA	T5	ST
UDENYCA AUTOINJECTOR	T5	ST
VAQTA (PF)	T3	
VARIVAX (PF)	T3	
XOLREMDI	T5	PA; QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
YF-VAX (PF)	T3	
ZARXIO	T5	
ZIEXTENZO	T5	
ZOMACTON	T4	PA
Miscellaneous Supplies		
ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"	T3	
GAUZE PAD TOPICAL BANDAGE 2 X 2 "	T3	
<i>insulin syringe-needle u-100 syringe 0.3 ml 29 gauge, 1 ml 29 gauge x 1/2", 1/2 ml 28 gauge</i>	T3	
<i>pen needle, diabetic needle 29 gauge x 1/2"</i>	T3	
Musculoskeletal / Rheumatology		
ABRILADA(CF)	T5	PA; QL (2 EA per 28 days)
ABRILADA(CF) PEN	T5	PA; QL (2 EA per 28 days)
ACTEMRA ACTPEN	T5	PA; QL (3.6 ML per 28 days)
ACTEMRA SUBCUTANEOUS	T5	PA; QL (3.6 ML per 28 days)
ACTONEL ORAL TABLET 150 MG, 35 MG	T4	
<i>adalimumab-aacf subcutaneous pen injector kit</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-aaty</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-adaz</i>	T5	PA; QL (0.8 ML per 28 days)
<i>adalimumab-adbm subcutaneous pen injector kit 40 mg/0.4 ml</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-adbm subcutaneous pen injector kit 40 mg/0.8 ml</i>	T5	PA; QL (1.6 EA per 28 days)
<i>adalimumab-adbm subcutaneous syringe kit 10 mg/0.2 ml</i>	T5	PA; QL (0.4 EA per 28 days)
<i>adalimumab-adbm subcutaneous syringe kit 20 mg/0.4 ml</i>	T5	PA; QL (0.8 EA per 28 days)
<i>adalimumab-adbm subcutaneous syringe kit 40 mg/0.4 ml</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-adbm subcutaneous syringe kit 40 mg/0.8 ml</i>	T5	PA; QL (1.6 EA per 28 days)
ADALIMUMAB-ADBM(CF) PEN CROHNS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML	T5	PA; QL (12 EA per 365 days)
ADALIMUMAB-ADBM(CF) PEN CROHNS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML	T5	PA; QL (9.6 EA per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ADALIMUMAB-ADBM(CF) PEN PS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML	T5	PA; QL (8 EA per 365 days)
ADALIMUMAB-ADBM(CF) PEN PS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML	T5	PA; QL (6.4 EA per 365 days)
<i>adalimumab-fkjp subcutaneous pen injector kit</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-fkjp subcutaneous syringe kit</i>	T5	PA; QL (2 EA per 28 days)
<i>adalimumab-ryvk subcutaneous auto-injector, kit</i>	T5	PA; QL (2 EA per 28 days)
<i>alendronate oral solution</i>	T2	
<i>alendronate oral tablet 10 mg, 35 mg, 70 mg</i>	T1	
<i>allopurinol oral tablet 100 mg, 300 mg</i>	T1	
<i>allopurinol oral tablet 200 mg</i>	T4	
AMJEVITA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 40 MG/0.4 ML	T5	PA; QL (0.8 ML per 28 days)
AMJEVITA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 40 MG/0.8 ML	T5	PA; QL (1.6 EA per 28 days)
AMJEVITA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 80 MG/0.8 ML	T5	PA; QL (1.6 ML per 28 days)
AMJEVITA(CF) SUBCUTANEOUS SYRINGE 10 MG/0.2 ML, 20 MG/0.2 ML	T5	PA; QL (0.4 ML per 28 days)
AMJEVITA(CF) SUBCUTANEOUS SYRINGE 20 MG/0.4 ML	T5	PA; QL (0.8 EA per 28 days)
AMJEVITA(CF) SUBCUTANEOUS SYRINGE 40 MG/0.4 ML	T5	PA; QL (0.8 ML per 28 days)
AMJEVITA(CF) SUBCUTANEOUS SYRINGE 40 MG/0.8 ML	T5	PA; QL (1.6 EA per 28 days)
ARAVA	T5	
AELVIA	T4	
BENLYSTA SUBCUTANEOUS	T5	PA; QL (4 ML per 28 days)
BINOSTO	T4	
<i>colchicine oral capsule</i>	T3	QL (62 EA per 31 days)
<i>colchicine oral tablet</i>	T2	QL (62 EA per 31 days)
COLCRYS	T4	QL (124 EA per 31 days)
CUPRIMINE	T5	
CYLTEZO(CF)	T5	PA; QL (2 EA per 28 days)
CYLTEZO(CF) PEN	T5	PA; QL (2 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
CYLTEZO(CF) PEN CROHN'S-UC-HS	T5	PA; QL (12 EA per 365 days)
CYLTEZO(CF) PEN PSORIASIS-UV	T5	PA; QL (8 EA per 365 days)
DEPEN TITRATABS	T5	
ENBREL MINI	T5	PA; QL (8 ML per 28 days)
ENBREL SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5)	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 50 MG/ML (1 ML)	T5	PA; QL (8 ML per 28 days)
ENBREL SURECLICK	T5	PA; QL (8 ML per 28 days)
EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML (105MG/1.17MLX2)	T5	PA; QL (2.34 ML per 28 days)
EVISTA	T4	
<i>febuxostat</i>	T3	PA
FORTEO	T5	PA; QL (2.4 ML per 28 days)
FOSAMAX ORAL TABLET 70 MG	T4	
FOSAMAX PLUS D	T4	
GLOPERBA	T4	QL (300 ML per 30 days)
HADLIMA	T5	PA; QL (1.6 ML per 28 days)
HADLIMA PUSHTOUCH	T5	PA; QL (1.6 ML per 28 days)
HADLIMA(CF)	T5	PA; QL (0.8 ML per 28 days)
HADLIMA(CF) PUSHTOUCH	T5	PA; QL (0.8 ML per 28 days)
HULIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT	T5	PA; QL (2 EA per 28 days)
HULIO(CF) SUBCUTANEOUS SYRINGE KIT	T5	PA; QL (2 EA per 28 days)
HUMIRA PEN	T5	PA; QL (2 EA per 28 days)
HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF)	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF) PEN	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF) PEN CROHNS-UC-HS	T5	PA; QL (3 EA per 28 days)
HUMIRA(CF) PEN PEDIATRIC UC	T5	PA; QL (4 EA per 28 days)
HUMIRA(CF) PEN PSOR-UV-ADOL HS	T5	PA; QL (3 EA per 28 days)
HYRIMOZ PEN CROHN'S-UC STARTER	T5	PA; QL (4.8 ML per 365 days)
HYRIMOZ PEN PSORIASIS STARTER	T5	PA; QL (3.2 ML per 365 days)
HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML	T5	PA; QL (4.8 ML per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML-40 MG/0.4 ML	T5	PA; QL (2.4 ML per 365 days)
HYRIMOZ(CF) PEN SUBCUTANEOUS PEN INJECTOR 40 MG/0.4 ML	T5	PA; QL (0.8 ML per 28 days)
HYRIMOZ(CF) PEN SUBCUTANEOUS PEN INJECTOR 80 MG/0.8 ML	T5	PA; QL (1.6 ML per 28 days)
HYRIMOZ(CF) SUBCUTANEOUS SYRINGE 10 MG/0.1 ML	T5	PA; QL (0.2 ML per 28 days)
HYRIMOZ(CF) SUBCUTANEOUS SYRINGE 20 MG/0.2 ML	T5	PA; QL (0.4 ML per 28 days)
HYRIMOZ(CF) SUBCUTANEOUS SYRINGE 40 MG/0.4 ML	T5	PA; QL (0.8 ML per 28 days)
<i>ibandronate oral</i>	T2	
IDACIO(CF)	T5	PA; QL (2 EA per 28 days)
IDACIO(CF) PEN CROHN-UC STARTR	T5	PA; QL (12 EA per 365 days)
IDACIO(CF) PEN PSORIASIS START	T5	PA; QL (8 EA per 365 days)
IDACIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT	T5	PA; QL (2 EA per 28 days)
KEVZARA	T5	PA; QL (2.28 ML per 28 days)
KINERET	T5	PA; QL (18.76 ML per 28 days)
<i>leflunomide</i>	T2	
MITIGARE	T4	QL (62 EA per 31 days)
OLUMIANT	T5	PA; QL (31 EA per 31 days)
ORENCIA CLICKJECT	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 50 MG/0.4 ML	T5	PA; QL (1.6 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 87.5 MG/0.7 ML	T5	PA; QL (2.8 ML per 28 days)
OTEZLA ORAL TABLET 30 MG	T5	PA; QL (62 EA per 31 days)
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)	T5	PA; QL (55 EA per 28 days)
OTREXUP (PF)	T4	PA
<i>penicillamine</i>	T5	
<i>probenecid</i>	T2	
<i>probenecid-colchicine</i>	T2	
PROLIA	T3	PA; QL (1 ML per 180 days)
<i>raloxifene</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
RASUVO (PF)	T4	PA
RIDAURA	T5	
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG	T5	PA; QL (31 EA per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 45 MG	T5	PA; QL (168 EA per 365 days)
<i>risedronate oral tablet 150 mg, 35 mg, 35 mg (12 pack), 35 mg (4 pack), 5 mg</i>	T4	
<i>risedronate oral tablet, delayed release (dr/ec)</i>	T4	
SAVELLA	T3	PA
SIMLANDI(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML	T5	PA; QL (1 ML per 28 days)
SIMPONI SUBCUTANEOUS PEN INJECTOR 50 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML	T5	PA; QL (1 ML per 28 days)
SIMPONI SUBCUTANEOUS SYRINGE 50 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
<i>teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)</i>	T5	PA; QL (2.48 ML per 28 days)
TYMLOS	T5	PA; QL (1.56 ML per 30 days)
ULORIC	T4	PA
XELJANZ ORAL SOLUTION	T5	PA; QL (310 ML per 31 days)
XELJANZ ORAL TABLET	T5	PA; QL (62 EA per 31 days)
XELJANZ XR	T5	PA; QL (31 EA per 31 days)
YUFLYMA(CF)	T5	PA; QL (2 EA per 28 days)
YUFLYMA(CF) AI CROHN'S-UC-HS	T5	PA; QL (6 EA per 365 days)
YUFLYMA(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
YUSIMRY(CF) PEN	T5	PA; QL (1.6 ML per 28 days)
Obstetrics / Gynecology		
ACTIVELLA	T4	
ALTAVERA (28)	T2	
ALYACEN 1/35 (28)	T2	
AMETHIA	T4	
ANGELIQ	T4	
ANNOVERA	T4	
APRI	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ARANELLE (28)	T2	
ASHLYNA	T4	
AUBRA EQ	T2	
AVIANE	T2	
BALCOLTRA	T4	
BALZIVA (28)	T4	
BEYAZ	T4	
BIJUVA	T4	
BLISOVI 24 FE	T4	
BLISOVI FE 1.5/30 (28)	T4	
BRIELLYN	T4	
CAMILA	T2	
CAMRESE LO	T4	
CLEOCIN VAGINAL	T4	
CLIMARA	T4	
CLIMARA PRO	T4	
<i>clindamycin phosphate vaginal</i>	T2	
CLINDESSE	T4	
COMBIPATCH	T4	
CRINONE	T4	PA
CRYSELLE (28)	T2	
CYRED EQ	T2	
DEBLITANE	T2	
DELESTROGEN INTRAMUSCULAR OIL 10 MG/ML, 20 MG/ML	T4	
DEPO-ESTRADIOL	T4	
DEPO-PROVERA INTRAMUSCULAR SUSPENSION 150 MG/ML	T4	
DEPO-PROVERA INTRAMUSCULAR SYRINGE	T4	
DEPO-SUBQ PROVERA 104	T4	
<i>desog-e.estradiol/e.estradiol</i>	T2	
<i>desogestrel-ethinyl estradiol</i>	T2	
DIVIGEL	T4	
DOLISHALE	T4	
DOTTI	T3	
<i>drospirenone-e.estradiol-lm.fa oral tablet 3-0.02- 0.451 mg (24) (4)</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>drospirenone-ethinyl estradiol</i>	T2	
DUAVEE	T3	
ELESTRIN	T4	
ELURYNG	T4	
ENILLORING	T4	
ENPRESSE	T2	
ENSKYCE	T2	
ERRIN	T2	
ESTARYLLA	T2	
ESTRACE	T4	
<i>estradiol oral</i>	T4	
<i>estradiol transdermal gel in metered-dose pump</i>	T4	
<i>estradiol transdermal gel in packet</i>	T4	
<i>estradiol transdermal patch semiweekly</i>	T3	
<i>estradiol transdermal patch weekly</i>	T3	
<i>estradiol vaginal</i>	T4	
<i>estradiol valerate</i>	T4	
<i>estradiol-norethindrone acet</i>	T3	
ESTRING	T4	
ESTROGEL	T4	
<i>ethynodiol diac-eth estradiol</i>	T2	
<i>etonogestrel-ethinyl estradiol</i>	T4	
EVAMIST	T4	
FALMINA (28)	T2	
FEMRING	T4	
FINZALA	T4	
FYAVOLV	T4	
GEMMILY	T4	
GYNAZOLE-1	T4	
HAILEY 24 FE	T4	
HALOETTE	T4	
HEATHER	T2	
ICLEVIA	T4	
IMVEXXY MAINTENANCE PACK	T3	
IMVEXXY STARTER PACK	T3	
INCASSIA	T2	
INTRAROSA	T4	PA; QL (28 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
INTROVALE	T2	
ISIBLOOM	T2	
JASMIEL (28)	T2	
JINTELI	T4	
JOYEAUX	T4	
JULEBER	T2	
JUNEL 1.5/30 (21)	T4	
JUNEL 1/20 (21)	T4	
JUNEL FE 1.5/30 (28)	T4	
JUNEL FE 1/20 (28)	T4	
JUNEL FE 24	T4	
KAITLIB FE	T4	
KARIVA (28)	T2	
KELNOR 1/35 (28)	T2	
KELNOR 1/50 (28)	T2	
KURVELO (28)	T2	
KYLEENA	T4	
<i>l norgest/e.estradiol-e.estrad</i>	T2	
LARIN 1.5/30 (21)	T2	
LARIN 1/20 (21)	T2	
LARIN FE 1.5/30 (28)	T2	
LARIN FE 1/20 (28)	T2	
LAYOLIS FE	T4	
LEENA 28	T4	
LESSINA	T2	
LEVONEST (28)	T2	
<i>levonorgestrel-ethinyl estrad</i>	T2	
<i>levonorg-eth estrad triphasic</i>	T2	
LEVORA-28	T2	
LILETTA	T4	
LO LOESTRIN FE	T4	
LOESTRIN 1.5/30 (21)	T4	
LOESTRIN 1/20 (21)	T4	
LOESTRIN FE 1.5/30 (28-DAY)	T4	
LOESTRIN FE 1/20 (28-DAY)	T4	
LORYNA (28)	T2	
LOW-OGESTREL (28)	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
LUTERA (28)	T2	
LYLEQ	T2	
LYLLANA	T3	
LYZA	T2	
MARLISSA (28)	T2	
<i>medroxyprogesterone</i>	T2	
MENEST	T3	
MENOSTAR	T4	
MERZEE	T4	
<i>metronidazole vaginal gel 0.75 % (37.5mg/5 gram)</i>	T3	
MIBELAS 24 FE	T4	
MICONAZOLE-3 VAGINAL SUPPOSITORY	T4	
MICROGESTIN 1.5/30 (21)	T2	
MICROGESTIN 1/20 (21)	T2	
MICROGESTIN 24 FE	T4	
MICROGESTIN FE 1.5/30 (28)	T2	
MICROGESTIN FE 1/20 (28)	T2	
MILI	T2	
MIMVEY	T3	
MINIVELLE	T4	
MIRENA	T4	
MYFEMBREE	T5	PA; QL (31 EA per 31 days)
NATAZIA	T4	
NECON 0.5/35 (28)	T4	
NEXPLANON	T4	
NEXTSTELLIS	T4	
NIKKI (28)	T2	
NORA-BE	T2	
<i>norelgestromin-ethin.estradiol</i>	T2	
<i>noreth-ethinyl estradiol-iron</i>	T4	
<i>norethindrone (contraceptive)</i>	T2	
<i>norethindrone acetate</i>	T2	
<i>norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg</i>	T4	
<i>norethindrone ac-eth estradiol oral tablet 1-20 mg-mcg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>norethindrone-e.estradiol-iron oral capsule</i>	T4	
<i>norethindrone-e.estradiol-iron oral tablet 1 mg-20 mcg (21)/75 mg (7), 1-20(5)/1-30(7) /1mg-35mcg (9)</i>	T2	
<i>norethindrone-e.estradiol-iron oral tablet,chewable</i>	T4	
<i>norgestimate-ethinyl estradiol</i>	T2	
NORTREL 0.5/35 (28)	T2	
NORTREL 1/35 (21)	T2	
NORTREL 1/35 (28)	T2	
NORTREL 7/7/7 (28)	T2	
NUVARING	T4	
NUVESSA	T4	
NYLIA 1/35 (28)	T4	
NYLIA 7/7/7 (28)	T4	
NYMYO	T4	
OCELLA	T4	
ORIAHNN	T5	PA; QL (56 EA per 28 days)
OSPHENA	T4	PA; QL (31 EA per 31 days)
PHEXXI	T4	
PIMTREA (28)	T2	
PORTIA 28	T2	
PREMARIN ORAL	T3	
PREMARIN VAGINAL	T3	
PREMPHASE	T3	
PREMPRO	T3	
<i>progesterone micronized</i>	T2	
PROMETRIUM	T4	
PROVERA	T4	
QUARTETTE	T4	
RECLIPSEN (28)	T2	
RIVELSA	T4	
SAFYRAL	T4	
SETLAKIN	T2	
SHAROBEL	T2	
SKYLA	T4	
SLYND	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SPRINTEC (28)	T2	
SRONYX	T2	
SYEDA	T2	
TARINA 24 FE	T2	
TARINA FE 1-20 EQ (28)	T2	
<i>terconazole</i>	T3	
TILIA FE	T2	
<i>tranexamic acid oral</i>	T3	
TRI-ESTARYLLA	T2	
TRI-LEGEST FE	T2	
TRI-LO-ESTARYLLA	T2	
TRI-LO-SPRINTEC	T2	
TRI-MILI	T4	
TRI-NYMYO	T4	
TRI-SPRINTEC (28)	T2	
TRIVORA (28)	T2	
TRI-VYLIBRA	T4	
TRI-VYLIBRA LO	T4	
TURQOZ (28)	T2	
TYBLUME	T4	
TYDEMY	T4	
VAGIFEM	T4	
VANDAZOLE	T3	
VELIVET TRIPHASIC REGIMEN (28)	T2	
VEOZAH	T4	PA; QL (31 EA per 31 days)
VESTURA (28)	T2	
VIENVA	T2	
VIVELLE-DOT	T4	
VYFEMLA (28)	T4	
VYLIBRA	T4	
WYMZYA FE	T4	
XACIATO	T4	
XULANE	T4	
YASMIN (28)	T4	
YAZ (28)	T4	
YUVAFEM	T4	
ZAFEMY	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZOVIA 1-35 (28)	T2	
Ophthalmology		
<i>acetazolamide</i>	T3	
ACULAR	T4	
ACULAR LS	T4	
ACUVAIL (PF)	T4	
ALOMIDE	T4	
ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.1 %	T3	
ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.15 %	T4	
ALREX	T3	
<i>apraclonidine</i>	T3	
<i>atropine ophthalmic (eye) drops 1 %</i>	T2	
AZASITE	T3	
<i>azelastine ophthalmic (eye)</i>	T2	
AZOPT	T4	
<i>bacitracin ophthalmic (eye)</i>	T2	
<i>bacitracin-polymyxin b</i>	T2	
<i>bepotastine besilate</i>	T3	
BEPREVE	T4	
BESIVANCE	T3	
<i>betaxolol ophthalmic (eye)</i>	T3	
BETIMOL	T4	
BETOPTIC S	T4	
<i>bimatoprost ophthalmic (eye)</i>	T4	
<i>brimonidine ophthalmic (eye) drops 0.1 %</i>	T3	
<i>brimonidine ophthalmic (eye) drops 0.15 %, 0.2 %</i>	T2	
<i>brimonidine-timolol</i>	T3	
<i>brinzolamide</i>	T4	
<i>bromfenac</i>	T3	
BROMSITE	T4	
<i>carteolol</i>	T2	
CEQUA	T4	ST; QL (60 EA per 30 days)
CILOXAN OPHTHALMIC (EYE) OINTMENT	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>ciprofloxacin hcl ophthalmic (eye)</i>	T2	
COMBIGAN	T3	
COSOPT	T4	
COSOPT (PF)	T4	
<i>cromolyn ophthalmic (eye)</i>	T2	
<i>cyclosporine ophthalmic (eye)</i>	T3	QL (60 EA per 30 days)
CYSTADROPS	T5	PA; QL (20 ML per 28 days)
CYSTARAN	T5	PA; QL (60 ML per 28 days)
<i>dexamethasone sodium phosphate ophthalmic (eye)</i>	T2	
<i>diclofenac sodium ophthalmic (eye)</i>	T2	
<i>difluprednate</i>	T4	
<i>dorzolamide</i>	T2	
<i>dorzolamide-timolol</i>	T2	
<i>dorzolamide-timolol (pf) ophthalmic (eye) dropperette</i>	T4	
DUREZOL	T4	
<i>epinastine</i>	T3	
<i>erythromycin ophthalmic (eye)</i>	T2	
EYSUVIS	T4	QL (8.3 ML per 30 days)
FLAREX	T4	
<i>fluorometholone</i>	T3	
<i>flurbiprofen sodium</i>	T2	
FML FORTE	T4	
FML LIQUIFILM	T4	
<i>gatifloxacin</i>	T2	
<i>gentamicin ophthalmic (eye) drops</i>	T2	
ILEVRO	T4	
INVELTYS	T3	
IOPIDINE OPHTHALMIC (EYE) DROPPERETTE	T4	
ISTALOL	T4	
IYUZEH (PF)	T4	ST; QL (30 EA per 30 days)
<i>ketorolac ophthalmic (eye)</i>	T2	
LACRISERT	T4	
<i>latanoprost</i>	T1	
<i>levobunolol ophthalmic (eye) drops 0.5 %</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>levofloxacin ophthalmic (eye) drops 0.5 %</i>	T3	
LOTEMAX	T4	
LOTEMAX SM	T4	
<i>loteprednol etabonate</i>	T3	
LUMIGAN OPHTHALMIC (EYE) DROPS 0.01 %	T3	QL (5 ML per 31 days)
MAXIDEX	T4	
MAXITROL	T4	
<i>methazolamide</i>	T4	
MIEBO (PF)	T5	ST; QL (9 ML per 30 days)
<i>moxifloxacin ophthalmic (eye) drops</i>	T3	
NATACYN	T4	
<i>neomycin-bacitracin-poly-hc</i>	T2	
<i>neomycin-bacitracin-polymyxin</i>	T2	
<i>neomycin-polymyxin b-dexameth</i>	T2	
<i>neomycin-polymyxin-gramicidin</i>	T2	
<i>neomycin-polymyxin-hc ophthalmic (eye)</i>	T3	
NEO-POLYCIN	T2	
NEO-POLYCIN HC	T2	
NEVANAC	T4	
OCUFLOX	T4	
<i>ofloxacin ophthalmic (eye)</i>	T2	
OXERVATE	T5	PA; QL (112 ML per 56 days)
PHOSPHOLINE IODIDE	T5	PA; QL (5 ML per 25 days)
<i>pilocarpine hcl ophthalmic (eye) drops 1 %, 2 %, 4 %</i>	T2	
POLYCIN	T2	
<i>polymyxin b sulf-trimethoprim</i>	T2	
PRED FORTE	T4	
PRED MILD	T4	
<i>prednisolone acetate</i>	T2	
<i>prednisolone sodium phosphate ophthalmic (eye)</i>	T2	
PROLENSA	T4	
RESTASIS	T3	QL (60 EA per 30 days)
RESTASIS MULTIDOSE	T3	QL (5.5 ML per 27 days)
RHOPRESSA	T3	ST
ROCKLATAN	T3	ST

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SIMBRINZA	T4	
<i>sulfacetamide sodium ophthalmic (eye)</i>	T2	
<i>sulfacetamide-prednisolone</i>	T2	
<i>tafluprost (pf)</i>	T4	
<i>timolol maleate (pf)</i>	T4	
<i>timolol maleate ophthalmic (eye) drops</i>	T1	
<i>timolol maleate ophthalmic (eye) drops, once daily</i>	T3	
<i>timolol maleate ophthalmic (eye) gel forming solution</i>	T3	
TIMOPTIC OCUDOSE (PF)	T4	
TOBRADEX OPHTHALMIC (EYE) OINTMENT	T3	
TOBRADEX ST	T3	
<i>tobramycin ophthalmic (eye)</i>	T2	
<i>tobramycin-dexamethasone</i>	T3	
TOBEX OPHTHALMIC (EYE) OINTMENT	T4	
TRAVATAN Z	T4	
<i>travoprost</i>	T3	
<i>trifluridine</i>	T3	
TYRVAYA	T4	ST; QL (8.4 ML per 30 days)
VERKAZIA	T5	PA; QL (120 EA per 30 days)
VEVYE	T4	ST; QL (2 ML per 50 days)
VIGAMOX	T4	
VUITY	T4	PA; QL (5 ML per 25 days)
VYZULTA	T4	ST; QL (5 ML per 31 days)
XALATAN	T4	
XDEMVY	T5	PA; QL (10 ML per 42 days)
XELPROS	T4	
XIIDRA	T3	QL (60 EA per 30 days)
ZERVIAE	T4	
ZIOPTAN (PF)	T4	
ZIRGAN	T4	ST
ZYLET	T4	
Respiratory And Allergy		
<i>acetylcysteine</i>	T3	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ADCIRCA	T5	PA; QL (62 EA per 31 days)
ADEMPAS	T5	PA; QL (93 EA per 31 days)
ADVAIR DISKUS	T4	QL (60 EA per 30 days)
ADVAIR HFA	T4	QL (12 GM per 30 days)
AIRDUO RESPICLICK	T4	QL (1 EA per 30 days)
AIRSUPRA	T4	ST; QL (32.1 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation</i>	T2	QL (17 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020503)</i>	T2	QL (13.4 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020983)</i>	T4	ST; QL (36 GM per 30 days)
<i>albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml</i>	T2	PA-BvD
<i>albuterol sulfate oral syrup</i>	T2	
<i>albuterol sulfate oral tablet</i>	T4	
ALVESCO	T4	QL (12.2 GM per 30 days)
ALYQ	T5	PA; QL (62 EA per 31 days)
<i>ambrisentan</i>	T5	PA; QL (31 EA per 31 days)
ANORO ELLIPTA	T4	QL (60 EA per 30 days)
<i>arformoterol</i>	T3	PA-BvD
ARNUTY ELLIPTA	T4	QL (30 EA per 30 days)
ASMANEX HFA	T3	QL (13 GM per 30 days)
ASMANEX TWISTHALER INHALATION AEROSOL POWDR BREATH ACTIVATED 110 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (120), 220 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (60)	T3	QL (1 EA per 30 days)
ATROVENT HFA	T3	QL (25.8 GM per 30 days)
AUVI-Q	T4	ST
<i>azelastine-fluticasone</i>	T4	QL (23 GM per 30 days)
BERINERT INTRAVENOUS KIT	T5	PA
BEVESPI AEROSPHERE	T4	QL (10.7 GM per 30 days)
<i>bosentan</i>	T5	PA; QL (62 EA per 31 days)
BREO ELLIPTA	T3	QL (60 EA per 30 days)
BREYNA	T3	QL (10.3 GM per 30 days)
BREZTRI AEROSPHERE	T3	QL (10.7 GM per 30 days)
BRONCHITOL	T5	PA; QL (560 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
BROVANA	T4	PA-BvD
<i>budesonide inhalation</i>	T4	PA-BvD
<i>budesonide-formoterol</i>	T3	QL (10.2 GM per 30 days)
<i>carbinoxamine maleate oral liquid</i>	T4	PA
<i>carbinoxamine maleate oral tablet 4 mg</i>	T4	PA
<i>cetirizine oral solution 1 mg/ml</i>	T2	QL (310 ML per 31 days)
CINRYZE	T5	PA; QL (20 EA per 28 days)
CLARINEX ORAL TABLET	T4	QL (31 EA per 31 days)
CLARINEX-D 12 HOUR	T4	
<i>clemastine oral syrup</i>	T4	PA
<i>clemastine oral tablet</i>	T4	
COMBIVENT RESPIMAT	T3	QL (4 GM per 30 days)
<i>cromolyn inhalation</i>	T5	PA-BvD
<i>cyproheptadine</i>	T4	PA
DALIRESP	T4	QL (31 EA per 31 days)
<i>desloratadine</i>	T4	QL (31 EA per 31 days)
DUAKLIR PRESSAIR	T5	QL (1 EA per 30 days)
DULERA	T4	QL (13 GM per 30 days)
DYMISTA	T4	ST; QL (23 GM per 30 days)
<i>epinephrine injection auto-injector 0.15 mg/0.15 ml, 0.3 mg/0.3 ml</i>	T4	
<i>epinephrine injection auto-injector 0.15 mg/0.3 ml, 0.3 mg/0.3 ml</i>	T3	
EPIPEN 2-PAK	T4	
EPIPEN JR 2-PAK	T4	
ESBRIET ORAL CAPSULE	T5	PA; QL (279 EA per 31 days)
ESBRIET ORAL TABLET	T5	PA; QL (93 EA per 31 days)
FASENRA PEN	T5	PA; QL (1 ML per 56 days)
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (0.5 ML per 56 days)
FASENRA SUBCUTANEOUS SYRINGE 30 MG/ML	T5	PA; QL (1 ML per 56 days)
FIRAZYR	T5	PA; QL (18 ML per 30 days)
<i>flunisolide</i>	T2	QL (50 ML per 25 days)
<i>fluticasone furoate-vilanterol</i>	T4	ST; QL (60 EA per 30 days)
<i>fluticasone propionate inhalation blister with device</i>	T3	QL (60 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>fluticasone propionate inhalation hfa aerosol inhaler 110 mcg/actuation</i>	T3	QL (12 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 220 mcg/actuation</i>	T3	QL (24 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 44 mcg/actuation</i>	T3	QL (10.6 GM per 30 days)
<i>fluticasone propionate nasal</i>	T2	QL (16 GM per 30 days)
<i>fluticasone propion-salmeterol inhalation aerosol powdr breath activated</i>	T3	QL (1 EA per 30 days)
<i>fluticasone propion-salmeterol inhalation blister with device</i>	T3	QL (60 EA per 30 days)
<i>fluticasone propion-salmeterol inhalation hfa aerosol inhaler</i>	T4	QL (12 GM per 30 days)
<i>formoterol fumarate</i>	T3	PA-BvD
HAEGARDA	T5	PA
<i>hydroxyzine hcl oral solution 10 mg/5 ml</i>	T4	PA
<i>hydroxyzine hcl oral tablet</i>	T2	PA
<i>hydroxyzine pamoate</i>	T4	PA
<i>icatibant</i>	T5	PA; QL (18 ML per 30 days)
INCRUSE ELLIPTA	T4	ST; QL (30 EA per 30 days)
<i>ipratropium bromide inhalation</i>	T2	PA-BvD
<i>ipratropium-albuterol</i>	T2	PA-BvD
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 5.8 MG, 50 MG, 75 MG	T5	PA; QL (56 EA per 28 days)
KALYDECO ORAL GRANULES IN PACKET 25 MG	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL TABLET	T5	PA; QL (62 EA per 31 days)
LETAIRIS	T5	PA; QL (31 EA per 31 days)
<i>levalbuterol hcl inhalation solution for nebulization 0.31 mg/3 ml, 0.63 mg/3 ml</i>	T2	PA-BvD
<i>levalbuterol hcl inhalation solution for nebulization 1.25 mg/0.5 ml</i>	T4	PA-BvD
<i>levalbuterol hcl inhalation solution for nebulization 1.25 mg/3 ml</i>	T3	PA-BvD
<i>levalbuterol tartrate</i>	T4	QL (30 GM per 30 days)
<i>levocetirizine oral solution</i>	T2	QL (310 ML per 31 days)
<i>levocetirizine oral tablet</i>	T2	QL (31 EA per 31 days)
LIQREV	T5	PA; QL (744 ML per 31 days)
<i>mometasone nasal</i>	T2	QL (34 GM per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>montelukast</i>	T2	QL (31 EA per 31 days)
NUCALA SUBCUTANEOUS AUTO-INJECTOR	T5	PA; QL (3 ML per 28 days)
NUCALA SUBCUTANEOUS RECON SOLN	T5	PA; QL (3 EA per 28 days)
NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML	T5	PA; QL (3 ML per 28 days)
NUCALA SUBCUTANEOUS SYRINGE 40 MG/0.4 ML	T5	PA; QL (0.4 ML per 28 days)
OFEV	T5	PA; QL (62 EA per 31 days)
OMNARIS	T4	QL (12.5 GM per 30 days)
OPSUMIT	T5	PA; QL (31 EA per 31 days)
OPSYNVI	T5	PA; QL (31 EA per 31 days)
ORKAMBI ORAL GRANULES IN PACKET	T5	PA; QL (62 EA per 31 days)
ORKAMBI ORAL TABLET	T5	PA; QL (124 EA per 31 days)
ORLADEYO	T5	PA; QL (31 EA per 31 days)
PERFOROMIST	T4	PA-BvD
<i>pirfenidone oral capsule</i>	T5	PA; QL (279 EA per 31 days)
<i>pirfenidone oral tablet</i>	T5	PA; QL (93 EA per 31 days)
PROAIR RESPICLICK	T4	ST; QL (2 EA per 30 days)
<i>promethazine oral</i>	T4	PA
<i>promethazine rectal suppository 12.5 mg, 25 mg</i>	T4	
PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG	T4	
PULMICORT	T4	PA-BvD
PULMICORT FLEXHALER	T4	QL (1 EA per 30 days)
PULMOZYME	T5	PA
QNASL NASAL HFA AEROSOL INHALER 40 MCG/ACTUATION	T4	QL (6.8 GM per 30 days)
QNASL NASAL HFA AEROSOL INHALER 80 MCG/ACTUATION	T4	QL (10.6 GM per 30 days)
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION	T3	QL (10.6 GM per 30 days)
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION	T3	QL (21.2 GM per 30 days)
REVATIO ORAL SUSPENSION FOR RECONSTITUTION	T5	PA; QL (784 ML per 31 days)
REVATIO ORAL TABLET	T5	PA; QL (372 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>roflumilast</i>	T4	QL (31 EA per 31 days)
RUCONEST	T5	PA
RYALTRIS	T4	ST; QL (29 GM per 30 days)
RYCLORA	T4	
RYVENT	T4	PA
SAJAZIR	T5	PA; QL (18 ML per 30 days)
SEREVENT DISKUS	T3	QL (60 EA per 30 days)
<i>sildenafil (pulm.hypertension) oral suspension for reconstitution</i>	T5	PA; QL (784 ML per 31 days)
<i>sildenafil (pulm.hypertension) oral tablet</i>	T3	PA; QL (372 EA per 31 days)
SINGULAIR	T4	QL (31 EA per 31 days)
SPIRIVA RESPIMAT	T3	QL (4 GM per 30 days)
SPIRIVA WITH HANDIHALER	T3	QL (30 EA per 30 days)
STIOLTO RESPIMAT	T3	QL (4 GM per 30 days)
STRIVERDI RESPIMAT	T3	QL (4 GM per 30 days)
SYMBICORT	T4	ST; QL (10.2 GM per 30 days)
SYMDEKO	T5	PA; QL (56 EA per 28 days)
<i>tadalafil (pulm. hypertension)</i>	T5	PA; QL (62 EA per 31 days)
TADLIQ	T5	PA; QL (310 ML per 31 days)
TAKHZYRO SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)
TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML	T5	PA; QL (2 ML per 28 days)
TAKHZYRO SUBCUTANEOUS SYRINGE 300 MG/2 ML (150 MG/ML)	T5	PA; QL (4 ML per 28 days)
<i>terbutaline oral</i>	T4	
THEO-24	T3	
<i>theophylline oral solution</i>	T2	
<i>theophylline oral tablet extended release 12 hr</i>	T2	
<i>theophylline oral tablet extended release 24 hr</i>	T2	
<i>tiotropium bromide</i>	T3	QL (30 EA per 30 days)
TRACLEER ORAL TABLET	T5	PA; QL (62 EA per 31 days)
TRACLEER ORAL TABLET FOR SUSPENSION	T5	PA; QL (124 EA per 31 days)
TRELEGY ELLIPTA	T3	QL (60 EA per 30 days)
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	T5	PA; QL (56 EA per 28 days)
TRIKAFTA ORAL TABLETS, SEQUENTIAL	T5	PA; QL (84 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TUDORZA PRESSAIR	T4	ST; QL (1 EA per 30 days)
TYVASO DPI INHALATION CARTRIDGE WITH INHALER 16 MCG, 32 MCG, 32-48 MCG, 48 MCG, 64 MCG	T5	PA
TYVASO DPI INHALATION CARTRIDGE WITH INHALER 16(112)-32(112) -48(28) MCG	T5	PA; QL (504 EA per 365 days)
VENTOLIN HFA	T3	QL (36 GM per 30 days)
VISTARIL ORAL CAPSULE 25 MG	T4	PA
WINREVAIR	T5	PA
WIXELA INHUB	T3	QL (60 EA per 30 days)
XHANCE	T4	QL (32 ML per 30 days)
XOLAIR	T5	PA
XOPENEX HFA	T4	ST; QL (30 GM per 30 days)
YUPELRI	T5	PA-BvD
<i>zafirlukast oral tablet 10 mg</i>	T2	QL (93 EA per 31 days)
<i>zafirlukast oral tablet 20 mg</i>	T2	QL (62 EA per 31 days)
ZETONNA	T4	QL (6.1 GM per 30 days)
<i>zileuton</i>	T5	PA; QL (124 EA per 31 days)
ZYFLO	T5	PA; QL (124 EA per 31 days)
Urologicals		
<i>alfuzosin</i>	T2	QL (31 EA per 31 days)
AVODART	T4	QL (31 EA per 31 days)
<i>bethanechol chloride</i>	T2	
CIALIS ORAL TABLET 2.5 MG	T4	PA; QL (62 EA per 31 days)
CIALIS ORAL TABLET 5 MG	T4	PA; QL (31 EA per 31 days)
CYSTAGON	T4	
<i>darifenacin</i>	T4	QL (31 EA per 31 days)
DETROL	T4	QL (62 EA per 31 days)
DETROL LA	T4	QL (31 EA per 31 days)
<i>dutasteride</i>	T2	QL (31 EA per 31 days)
<i>dutasteride-tamsulosin</i>	T4	QL (31 EA per 31 days)
ELMIRON	T4	
ENTADFI	T4	PA; QL (31 EA per 31 days)
<i>fesoterodine</i>	T3	QL (31 EA per 31 days)
<i>finasteride oral tablet 5 mg</i>	T2	
<i>flavoxate</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
FLOMAX	T4	
GEMTESA	T4	QL (31 EA per 31 days)
<i>mirabegron</i>	T4	ST; QL (31 EA per 31 days)
MYRBETRIQ ORAL SUSPENSION,EXTENDED REL RECON	T3	QL (300 ML per 30 days)
MYRBETRIQ ORAL TABLET EXTENDED RELEASE 24 HR	T3	QL (31 EA per 31 days)
<i>oxybutynin chloride oral syrup</i>	T3	
<i>oxybutynin chloride oral tablet 5 mg</i>	T3	
<i>oxybutynin chloride oral tablet extended release 24hr 10 mg, 5 mg</i>	T3	QL (31 EA per 31 days)
<i>oxybutynin chloride oral tablet extended release 24hr 15 mg</i>	T3	QL (62 EA per 31 days)
OXYTROL	T4	QL (8 EA per 28 days)
<i>potassium citrate oral tablet extended release</i>	T2	
PROCYSBI ORAL GRANULES DEL RELEASE IN PACKET	T5	PA
PROSCAR	T4	
RAPAFLO	T4	
RIVFLOZA SUBCUTANEOUS SOLUTION	T5	PA; QL (1 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML	T5	PA; QL (0.8 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 160 MG/ML	T5	PA; QL (1 ML per 28 days)
<i>silodosin</i>	T2	
<i>solifenacin</i>	T4	QL (31 EA per 31 days)
<i>tadalafil oral tablet 2.5 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>tadalafil oral tablet 5 mg</i>	T4	PA; QL (31 EA per 31 days)
<i>tamsulosin</i>	T1	
<i>tolterodine oral capsule,extended release 24hr</i>	T3	QL (31 EA per 31 days)
<i>tolterodine oral tablet</i>	T3	QL (62 EA per 31 days)
TOVIAZ	T4	QL (31 EA per 31 days)
<i>trospium oral capsule,extended release 24hr</i>	T4	QL (31 EA per 31 days)
<i>trospium oral tablet</i>	T2	QL (93 EA per 31 days)
UROCIT-K 10	T4	
UROCIT-K 15	T4	
UROCIT-K 5	T4	
UROXATRAL	T4	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
VESICARE	T4	QL (31 EA per 31 days)
VESICARE LS	T4	QL (310 ML per 31 days)
Vitamins, Hematinics / Electrolytes		
<i>calcium acetate(phosphat bind)</i>	T2	
CLINIMIX 5%/D15W SULFITE FREE	T4	PA-BvD
CLINIMIX 4.25%/D10W SULF FREE	T4	PA-BvD
CLINIMIX 5%-D20W(SULFITE-FREE)	T4	PA-BvD
CLINIMIX E 4.25%/D10W SUL FREE	T4	PA-BvD
CLINIMIX E 4.25%/D5W SULF FREE	T4	PA-BvD
CLINIMIX E 5%/D15W SULFIT FREE	T4	PA-BvD
CLINIMIX E 5%/D20W SULFIT FREE	T4	PA-BvD
CLINISOL SF 15 %	T4	PA-BvD
DOJOLVI	T5	PA
<i>electrolyte-148</i>	T4	PA-BvD
<i>fluoride (sodium) oral tablet</i>	T2	
INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %	T4	PA-BvD
ISOLYTE S PH 7.4	T3	PA-BvD
ISOLYTE-P IN 5 % DEXTROSE	T4	PA-BvD
KLOR-CON	T4	
KLOR-CON 10	T3	
KLOR-CON 8	T3	
KLOR-CON M10	T2	
KLOR-CON M15	T2	
KLOR-CON M20	T2	
<i>magnesium sulfate injection</i>	T2	
NUTRILIPID	T4	PA-BvD
PLASMA-LYTE 148	T4	PA-BvD
PLASMA-LYTE A	T4	PA-BvD
PLENAMINE	T4	PA-BvD
<i>potassium chlorid-d5-0.45%nacl</i>	T2	
<i>potassium chloride in 0.9%nacl intravenous parenteral solution 20 meq/l, 40 meq/l</i>	T2	
<i>potassium chloride in 5 % dex intravenous parenteral solution 20 meq/l</i>	T2	
<i>potassium chloride in lr-d5 intravenous parenteral solution 20 meq/l</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>potassium chloride in water intravenous piggyback 10 meq/100 ml, 20 meq/100 ml, 40 meq/100 ml</i>	T2	
<i>potassium chloride intravenous</i>	T2	
<i>potassium chloride oral capsule, extended release</i>	T2	
<i>potassium chloride oral liquid</i>	T4	
<i>potassium chloride oral packet</i>	T4	
<i>potassium chloride oral tablet extended release</i>	T2	
<i>potassium chloride oral tablet, er particles/crystals</i>	T2	
<i>potassium chloride-0.45 % nacl</i>	T2	
<i>potassium chloride-d5-0.2%nacl intravenous parenteral solution 20 meq/l</i>	T2	
<i>potassium chloride-d5-0.9%nacl</i>	T2	
PREMASOL 10 %	T4	PA-BvD
PRENATAL VITAMIN PLUS LOW IRON	T2	PA
PROSOL 20 %	T4	PA-BvD
<i>sodium chloride 0.45 % intravenous</i>	T2	
<i>sodium chloride 3 % hypertonic</i>	T2	
<i>sodium chloride 5 % hypertonic</i>	T2	
TPN ELECTROLYTES	T4	
TRAVASOL 10 %	T4	PA-BvD
TROPHAMINE 10 %	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Index of Drugs

<i>abacavir</i>	3	ADALIMUMAB-ADBM(CF)	ALOMIDE	113
<i>abacavir-lamivudine</i>	3	PEN PS-UV	<i>alosepron</i>	91, 92
ABELCET	3	<i>adalimumab-fkjp</i>	ALPHAGAN P	113
ABILIFY	25	<i>adalimumab-ryvk</i>	<i>alprazolam</i>	25, 26
ABILIFY ASIMTUFII	25	<i>adapalene</i>	ALPRAZOLAM INTENSOL ..	25
ABILIFY MAINTENA	25	<i>adapalene-benzoyl peroxide</i>	ALREX	113
ABILIFY MYCITE		ADBRY	ALTABAX	70
MAINTENANCE KIT	25	ADCIRCA	ALTACE	58
ABILIFY MYCITE		ADDERALL	ALTAVERA (28)	106
STARTER KIT	25	ADDERALL XR	ALTOPREV	58
<i>abiraterone</i>	16	<i>adefovir</i>	ALTRENO	70
ABRILADA(CF)	102	ADEMPAS	ALUNBRIG	16
ABRILADA(CF) PEN	102	ADLARITY	ALVAIZ	58
ABRYSVO (PF)	98	ADMELOG SOLOSTAR U-	ALVESCO	117
ABSORICA	70	100 INSULIN	ALYACEN 1/35 (28)	106
ABSORICA LD	70	ADMELOG U-100 INSULIN	ALYQ	117
<i>acamprosate</i>	79	LISPRO	<i>amantadine hcl</i>	3
ACANYA	70	ADVAIR DISKUS	AMBIEN	26
<i>acarbose</i>	83	ADVAIR HFA	AMBIEN CR	26
ACCUTANE	70	ADZENYS XR-ODT	AMBISOME	3
<i>acebutolol</i>	58	AEMCOLO	<i>ambrisentan</i>	117
<i>acetaminophen-caff-dihydrocod</i> ..	25	AFINITOR	<i>amcinonide</i>	70
<i>acetaminophen-codeine</i>	25	AFINITOR DISPERZ	AMETHIA	106
<i>acetazolamide</i>	113	AFREZZA	<i>amikacin</i>	3
<i>acetic acid</i>	82	AGAMREE	<i>amiloride</i>	58
<i>acetylcysteine</i>	116	AGRYLIN	<i>amiloride-hydrochlorothiazide</i> ..	58
ACIPHEX	91	AIMOVIG AUTOINJECTOR ..	<i>amiodarone</i>	58
<i>acitretin</i>	70	AIRDUO RESPICLICK	AMITIZA	92
ACTEMRA	102	AIRSUPRA	<i>amitriptyline</i>	26
ACTEMRA ACTPEN	102	AJOVY AUTOINJECTOR	<i>amitriptyline-chlordiazepoxide</i> ..	26
ACTHAR	83	AJOVY SYRINGE	AMJEVITA(CF)	103
ACTHIB (PF)	98	AKEEGA	AMJEVITA(CF)	
ACTIMMUNE	98	AKLIEF	AUTOINJECTOR	103
ACTIVELLA	106	ALA-CORT	<i>amlodipine</i>	58
ACTONEL	102	ALA-SCALP	<i>amlodipine-atorvastatin</i>	58
ACTOPLUS MET	83	<i>albendazole</i>	<i>amlodipine-benazepril</i>	58
ACTOS	83	<i>albuterol sulfate</i>	<i>amlodipine-olmesartan</i>	58
ACULAR	113	<i>alclometasone</i>	<i>amlodipine-valsartan</i>	58
ACULAR LS	113	ALCOHOL PADS	<i>amlodipine-valsartan-hcthiazid</i> ..	58
ACUVAIL (PF)	113	ALDACTONE	<i>ammonium lactate</i>	70
<i>acyclovir</i>	3, 70	ALECENSA	AMNESTEEM	70
<i>acyclovir sodium</i>	3	<i>alendronate</i>	<i>amoxapine</i>	26
ACZONE	70	<i>alfuzosin</i>	<i>amoxicil-clarithromy-lansopraz</i> ..	92
ADACEL(TDAP		<i>aliskiren</i>	<i>amoxicillin</i>	3
ADOLESN/ADULT)(PF)	98	ALKINDI SPRINKLE	<i>amoxicillin-pot clavulanate</i>	4
<i>adalimumab-aacf</i>	102	<i>allopurinol</i>	<i>amphetamine sulfate</i>	26
<i>adalimumab-aaty</i>	102	ALLZITAL	<i>amphotericin b</i>	4
<i>adalimumab-adaz</i>	102	<i>almotriptan malate</i>	<i>amphotericin b liposome</i>	4
<i>adalimumab-adbm</i>	102	<i>alogliptin</i>	<i>ampicillin</i>	4
ADALIMUMAB-ADBM(CF)		<i>alogliptin-metformin</i>	<i>ampicillin sodium</i>	4
PEN CROHNS	102	<i>alogliptin-pioglitazone</i>	<i>ampicillin-sulbactam</i>	4

AMPYRA	26	ASMANEX HFA	117	AZOPT	113
AMRIX	26	ASMANEX TWISTHALER	117	AZOR	59
AMZEEQ	70	<i>aspirin-dipyridamole</i>	59	AZSTARYS	27
ANAFRANIL	26	ASPRUZYO SPRINKLE	59	<i>aztreonam</i>	4
<i>anagrelide</i>	79	ASSURE ID INSULIN		AZULFIDINE	92
<i>anastrozole</i>	16	SAFETY	102	AZULFIDINE EN-TABS	92
ANCOBON	4	ASTAGRAF XL	16	<i>bacitracin</i>	113
ANDROGEL	83	ATACAND	59	<i>bacitracin-polymyxin b</i>	113
ANGELIQ	106	ATACAND HCT	59	<i>baclofen</i>	27
ANNOVERA	106	<i>atazanavir</i>	4	BACTRIM	4
ANORO ELLIPTA	117	ATELVIA	103	BACTRIM DS	4
ANTIVERT	92	<i>atenolol</i>	59	BAFIERTAM	27
ANUSOL-HC	92	<i>atenolol-chlorthalidone</i>	59	BALCOLTRA	107
ANZEMET	92	ATIVAN	27	<i>balsalazide</i>	92
APEXICON E	70	<i>atomoxetine</i>	27	BALVERSA	16
APIDRA SOLOSTAR U-100		ATORVALIQ	59	BALZIVA (28)	107
INSULIN	83	<i>atorvastatin</i>	59	BANZEL	27
APIDRA U-100 INSULIN	84	<i>atovaquone</i>	4	BAQSIMI	84
APLENZIN	26	<i>atovaquone-proguanil</i>	4	BARACLUDGE	4
APOKYN	26	ATRALIN	70	BASAGLAR KWIKPEN U-	
<i>apomorphine</i>	26	<i>atropine</i>	113	100 INSULIN	84
<i>apraclonidine</i>	113	ATROVENT HFA	117	BASAGLAR TEMPO	
<i>aprepitant</i>	92	AUBAGIO	27	PEN(U-100)INSLN	84
APRI	106	AUBRA EQ	107	BAXDELA	4
APRISO	92	AUGMENTIN	4	<i>bcg vaccine, live (pf)</i>	98
APTENSIO XR	26	AUGMENTIN ES-600	4	BELBUCA	28
APTIOM	26	AUGTYRO	16	BELSOMRA	28
APTIVUS	4	AURYXIA	79	<i>benazepril</i>	59
ARALAST NP	79	AUSTEDO	27	<i>benazepril-hydrochlorothiazide</i>	59
ARANELLE (28)	107	AUSTEDO XR	27	BENICAR	59
ARANESP (IN		AUSTEDO XR TITRATION		BENICAR HCT	59
POLYSORBATE)	98	KT(WK1-4)	27	BENLYSTA	103
ARAVA	103	AUVELITY	27	BENZAMYCIN	70
ARAZLO	70	AUVI-Q	117	<i>benznidazole</i>	4
ARCALYST	98	AVALIDE	59	<i>benztropine</i>	28
AREXVY (PF)	98	AVAPRO	59	<i>bepotastine besilate</i>	113
<i>arformoterol</i>	117	AVEED	84	BEPREVE	113
ARICEPT	26	AVIANE	107	BERINERT	117
ARIKAYCE	4	AVODART	122	BESIVANCE	113
ARIMIDEX	16	AVONEX	98	BESREMI	98
<i>aripiprazole</i>	26	AVYCAZ	4	<i>betaine</i>	92
ARISTADA	26, 27	AYVAKIT	16	<i>betamethasone dipropionate</i>	70
ARISTADA INITIO	26	AZACTAM	4	<i>betamethasone valerate</i>	70, 71
ARIXTRA	59	AZASAN	16	<i>betamethasone, augmented</i>	71
<i>armodafinil</i>	27	AZASITE	113	BETAPACE AF	59
ARNUITY ELLIPTA	117	<i>azathioprine</i>	16	BETASERON	98
AROMASIN	16	<i>azelaic acid</i>	70	<i>betaxolol</i>	59, 113
ARTHROTEC 50	27	<i>azelastine</i>	82, 113	<i>bethanechol chloride</i>	122
ARTHROTEC 75	27	<i>azelastine-fluticasone</i>	117	BETHKIS	4
ASCOMP WITH CODEINE	27	AZELEX	70	BETIMOL	113
<i>asenapine maleate</i>	27	AZILECT	27	BETOPTIC S	113
ASHLYNA	107	<i>azithromycin</i>	4	BEVESPI AEROSPHERE	117

<i>bexarotene</i>	16	<i>butalbital-acetaminop-caf-cod</i>	28	CAROSPIR	60
BEXSERO	98	<i>butalbital-acetaminophen</i>	28	<i>carteolol</i>	113
BEYAZ	107	<i>butalbital-acetaminophen-caff</i>	28, 29	CARTIA XT	60
<i>bicalutamide</i>	16	<i>butalbital-aspirin-caffeine</i>	29	<i>carvedilol</i>	60
BICILLIN C-R	4	<i>butorphanol</i>	29	<i>carvedilol phosphate</i>	60
BICILLIN L-A	4	BUTRANS	29	CASODEX	17
BIDIL	59	BYDUREON BCISE	84	<i>casprofungin</i>	5
BIJUVA	107	BYETTA	84	CAYSTON	5
BIKTARVY	5	BYLVAY	92	<i>cefaclor</i>	5
BILTRICIDE	5	BYSTOLIC	59	<i>cefadroxil</i>	5
<i>bimatoprost</i>	113	<i>cabergoline</i>	84	<i>cefazolin</i>	5
BIMZELX	71	CABLIVI	59	<i>cefdinir</i>	5
BIMZELX AUTOINJECTOR	71	CABOMETYX	17	<i>cefepime</i>	5
BINOSTO	103	CABTREO	71	<i>cefixime</i>	5
<i>bismuth subcit k-metronidz-tcn</i> ...	92	CADUET	59	<i>cefotetan</i>	5
<i>bisoprolol fumarate</i>	59	<i>calcipotriene</i>	71	<i>cefoxitin</i>	5
<i>bisoprolol-hydrochlorothiazide</i> ..	59	<i>calcipotriene-betamethasone</i>	71	<i>cefpodoxime</i>	5
BIVIGAM	98	<i>calcitonin (salmon)</i>	84	<i>cefprozil</i>	5
BLISOVI 24 FE	107	<i>calcitriol</i>	71, 84	<i>ceftazidime</i>	5
BLISOVI FE 1.5/30 (28)	107	<i>calcium acetate(phosphat bind)</i>	124	<i>ceftriaxone</i>	5
BONJESTA	92	CALQUENCE	17	<i>cefuroxime axetil</i>	5
BOOSTRIX TDAP	98	CALQUENCE		<i>cefuroxime sodium</i>	5
<i>bosentan</i>	117	(ACALABRUTINIB MAL)	17	CELEBREX	29
BOSULIF	17	CAMBIA	29	<i>celecoxib</i>	29
BRAFTOVI	17	CAMILA	107	CELEXA	29
BREO ELLIPTA	117	CAMRESE LO	107	CELLCEPT	17
BREYNA	117	CAMZYOS	59	CELONTIN	29
BREZTRI AEROSPHERE	117	CANASA	92	<i>cephalexin</i>	5
BRIELLYN	107	CANCIDAS	5	CEQUA	113
BRILINTA	59	<i>candesartan</i>	59	CERDELGA	84
<i>brimonidine</i>	71, 113	<i>candesartan-hydrochlorothiazid</i> ..	59	<i>cetirizine</i>	118
<i>brimonidine-timolol</i>	113	CAPLYTA	29	CETRAXAL	82
<i>brinzolamide</i>	113	CAPRELSA	17	<i>cevimeline</i>	79
BRIVIACT	28	<i>captopril</i>	59	CHEMET	79
<i>bromfenac</i>	113	CARAC	71	CHENODAL	92
<i>bromocriptine</i>	28	CARAFATE	92	<i>chlordiazepoxide hcl</i>	29
BROMSITE	113	CARBAGLU	79	<i>chlordiazepoxide-clidinium</i>	92
BRONCHITOL	117	<i>carbamazepine</i>	29	<i>chlorhexidine gluconate</i>	82
BROVANA	118	CARBATROL	29	<i>chloroquine phosphate</i>	5
BRUKINSA	17	<i>carbidopa</i>	29	<i>chlorpromazine</i>	29
BRYHALI	71	<i>carbidopa-levodopa</i>	29	<i>chlorthalidone</i>	60
<i>budesonide</i>	92, 118	<i>carbidopa-levodopa-entacapone</i>	29	<i>chlorzoxazone</i>	29
<i>budesonide-formoterol</i>	118	<i>carbinoxamine maleate</i>	118	CHOLBAM	92
<i>bumetanide</i>	59	CARDIZEM	60	<i>cholestyramine (with sugar)</i>	60
BUPAP	28	CARDIZEM CD	60	CHOLESTYRAMINE	
BUPHENYL	79	CARDIZEM LA	60	LIGHT	60
<i>buprenorphine</i>	28	CARDURA	60	CIALIS	122
<i>buprenorphine hcl</i>	28	CARDURA XL	60	CIBINQO	71
<i>buprenorphine-naloxone</i>	28	<i>carglumic acid</i>	79	<i>ciclopirox</i>	71
<i>bupropion hcl</i>	28	<i>carisoprodol</i>	29	<i>cilostazol</i>	60
<i>bupropion hcl (smoking deter)</i>	79	CARNITOR	79	CILOXAN	113
<i>buspirona</i>	28			CIMDUO	5

<i>cimetidine</i>	92	CLINIMIX E 5%/D20W		COSENTYX	72
CIMZIA	92	SULFIT FREE	124	COSENTYX (2 SYRINGES) ...	72
CIMZIA POWDER FOR RECONST	92	CLINISOL SF 15 %	124	COSENTYX PEN (2 PENS)	72
<i>cinacalcet</i>	84	<i>clobazam</i>	29	COSENTYX UNOREADY PEN	72
CINRYZE	118	<i>clobetasol</i>	71, 72	COSOPT	114
CIPRO	6	<i>clobetasol-emollient</i>	72	COSOPT (PF)	114
CIPRO HC	82	CLOBEX	72	COTELLIC	17
<i>ciprofloxacin hcl</i>	6, 82, 114	<i>clocortolone pivalate</i>	72	COTEMPLA XR-ODT	30
<i>ciprofloxacin in 5 % dextrose</i>	6	CLODAN	72	COZAAR	60, 61
<i>ciprofloxacin-dexamethasone</i>	82	<i>clomipramine</i>	29	CREON	92
<i>ciprofloxacin-fluocinolone</i>	82	<i>clonazepam</i>	29, 30	CRESEMBA	6
<i>citalopram</i>	29	<i>clonidine</i>	60	CRESTOR	61
CLARAVIS	71	<i>clonidine hcl</i>	30, 60	CRINONE	107
CLARINEX	118	<i>clopidogrel</i>	60	<i>cromolyn</i>	92, 114, 118
CLARINEX-D 12 HOUR	118	<i>clorazepate dipotassium</i>	30	CROTAN	72
<i>clarithromycin</i>	6	<i>clotrimazole</i>	6, 72	CRYSELLE (28)	107
<i>clemastine</i>	118	<i>clotrimazole-betamethasone</i>	72	CUBICIN RF	6
CLENPIQ	92	<i>clozapine</i>	30	CUPRIMINE	103
CLEOCIN	6, 107	CLOZARIL	30	CUVPOSA	92
CLEOCIN HCL	6	COARTEM	6	CUVRIOR	79
CLEOCIN PEDIATRIC	6	<i>codeine sulfate</i>	30	<i>cyclobenzaprine</i>	30
CLEOCIN T	71	<i>codeine-butalbital-asa-caff</i>	30	<i>cyclophosphamide</i>	17
CLIMARA	107	COLAZAL	92	<i>cycloserine</i>	6
CLIMARA PRO	107	<i>colchicine</i>	103	CYCLOSET	84
CLINDACIN	71	COLCRYS	103	<i>cyclosporine</i>	17, 114
CLINDACIN ETZ	71	<i>colesevelam</i>	60	<i>cyclosporine modified</i>	17
CLINDAGEL	71	COLESTID	60	CYLTEZO(CF)	103
<i>clindamycin hcl</i>	6	<i>colestipol</i>	60	CYLTEZO(CF) PEN	103
<i>clindamycin in 5 % dextrose</i>	6	<i>colistin (colistimethate na)</i>	6	CYLTEZO(CF) PEN	
CLINDAMYCIN PEDIATRIC	6	COMBIGAN	114	CROHN'S-UC-HS	104
<i>clindamycin phosphate</i> ... 6, 71, 107		COMBIPATCH	107	CYLTEZO(CF) PEN	
<i>clindamycin-benzoyl peroxide</i>	71	COMBIVENT RESPIMAT ..	118	PSORIASIS-UV	104
<i>clindamycin-tretinoin</i>	71	COMBIVIR	6	CYMBALTA	30
CLINDESSE	107	COMETRIQ	17	<i>cyproheptadine</i>	118
CLINIMIX 5%/D15W SULFITE FREE	124	COMPLERA	6	CYRED EQ	107
CLINIMIX 4.25%/D10W SULF FREE	124	COMPRO	92	CYSTADANE	93
CLINIMIX 4.25%/D5W SULFIT FREE	79	COMTAN	30	CYSTADROPS	114
CLINIMIX 5%-D20W(SULFITE-FREE)	124	CONCERTA	30	CYSTAGON	122
CLINIMIX E 2.75%/D5W SULF FREE	79	CONDYLOX	72	CYSTARAN	114
CLINIMIX E 4.25%/D10W SUL FREE	124	CONSTULOSE	92	CYTOMEL	84
CLINIMIX E 4.25%/D5W SULF FREE	124	CONZIP	30	CYTOTEC	93
CLINIMIX E 5%/D15W SULFIT FREE	124	COPAXONE	30	<i>d10 %-0.45 % sodium chloride</i> ..	79
		COPIKTRA	17	<i>d2.5 %-0.45 % sodium chloride</i> ..	79
		CORDRAN	72	<i>d5 % and 0.9 % sodium chloride</i>	79
		CORDRAN TAPE LARGE ROLL	72	<i>d5 %-0.45 % sodium chloride</i>	79
		COREG	60	<i>dabigatran etexilate</i>	61
		COREG CR	60	<i>dalfampridine</i>	30
		CORGARD	60	DALIRESP	118
		CORLANOR	60	DALVANCE	6
		CORTEF	84	<i>danazol</i>	84
		CORTROPHIN GEL	84		

DANTRIUM	30	DETROL LA	122	<i>disopyramide phosphate</i>	61
<i>danrolene</i>	30	DEXABLISS	84	<i>disulfiram</i>	80
<i>dapaglifloz propaned-metformin</i>	84	<i>dexamethasone</i>	84	DIURIL	61
<i>dapagliflozin propanediol</i>	84	<i>dexamethasone sodium</i>		<i>divalproex</i>	32
<i>dapsone</i>	6, 72	<i>phosphate</i>	114	DIVIGEL	107
DAPTACEL (DTAP		DEXEDRINE SPANSULE	31	<i>dofetilide</i>	61
PEDIATRIC) (PF)	98	DEXILANT	93	DOJOLVI	124
<i>daptomycin</i>	6	<i>dexlansoprazole</i>	93	DOLISHALE	107
DARAPRIM	6	<i>dexmethylphenidate</i>	31	<i>donepezil</i>	33
<i>darifenacin</i>	122	<i>dextroamphetamine sulfate</i>	31	DOPTELET (10 TAB PACK) .	61
<i>darunavir</i>	6	<i>dextroamphetamine-</i>		DOPTELET (15 TAB PACK) .	61
DAURISMO	17	<i>amphetamine</i>	31, 32	DOPTELET (30 TAB PACK) .	61
DAYBUE	31	<i>dextrose 10 % and 0.2 % nacl</i> ...	79	DORYX MPC	7
DAYPRO	31	<i>dextrose 10 % in water (d10w)</i> ..	79	<i>dorzolamide</i>	114
DAYTRANA	31	<i>dextrose 5 % in water (d5w)</i>	80	<i>dorzolamide-timolol</i>	114
DAYVIGO	31	<i>dextrose 5%-0.2 % sod chloride</i> .	80	<i>dorzolamide-timolol (pf)</i>	114
DDAVP	84	DHIVY	32	DOTTI	107
DEBLITANE	107	DIACOMIT	32	DOVATO	7
<i>deferasirox</i>	79	<i>diazepam</i>	32	<i>doxazosin</i>	61
<i>deferiprone</i>	79	DIAZEPAM INTENSOL	32	<i>doxepin</i>	33, 73
<i>deflazacort</i>	84	<i>diazoxide</i>	85	<i>doxercalciferol</i>	85
DELESTROGEN	107	DIBENZYLINE	61	DOXY-100	7
DELSTRIGO	6	DICLEGIS	93	<i>doxycycline hyclate</i>	7
DELZICOL	93	<i>diclofenac epolamine</i>	32	<i>doxycycline monohydrate</i>	7
<i>demeclocycline</i>	6	<i>diclofenac potassium</i>	32	<i>doxylamine-pyridoxine (vit b6)</i> ..	93
DEMEROL	31	<i>diclofenac sodium</i>	32, 73, 114	<i>dronabinol</i>	93
DEMEROL (PF)	31	<i>diclofenac-misoprostol</i>	32	<i>drospirenone-e.estradiol-lm.fa</i> .	107
DEMSEER	61	<i>dicloxacillin</i>	6	<i>drospirenone-ethinyl estradiol</i> ..	108
DENAVIR	72	<i>dicyclomine</i>	93	DROXIA	17
DEPAKOTE	31	DIFFERIN	73	<i>droxidopa</i>	80
DEPAKOTE ER	31	DIFICID	6	DUAKLIR PRESSAIR	118
DEPAKOTE SPRINKLES	31	<i>diflorasone</i>	73	DUAVEE	108
DEPEN TITRATABS	104	DIFLUCAN	6, 7	DUETACT	85
DEPO-ESTRADIOL	107	<i>diflunisal</i>	32	DULERA	118
DEPO-PROVERA	107	<i>difluprednate</i>	114	<i>duloxetine</i>	33
DEPO-SUBQ PROVERA	104 107	<i>digoxin</i>	61	DUOBRII	73
DEPO-TESTOSTERONE	84	<i>dihydroergotamine</i>	32	DUOPA	33
DERMA-SMOOTHIE/FS		DILANTIN	32	DUPIXENT PEN	73
SCALP OIL	72	DILANTIN EXTENDED	32	DUPIXENT SYRINGE	73
DERMOTIC OIL	82	DILANTIN INFATABS	32	DUREZOL	114
DESCOVY	6	DILANTIN-125	32	<i>dutasteride</i>	122
<i>desipramine</i>	31	DILAUDID	32	<i>dutasteride-tamsulosin</i>	122
<i>desloratadine</i>	118	<i>diltiazem hcl</i>	61	DYANAVEL XR	33
<i>desmopressin</i>	84	DILT-XR	61	DYMISTA	118
<i>desog-e.estradiol/e.estradiol</i>	107	<i>dimethyl fumarate</i>	32	DYRENIUM	61
<i>desogestrel-ethinyl estradiol</i>	107	DIOVAN	61	E.E.S. 400	7
<i>desonide</i>	72	DIOVAN HCT	61	E.E.S. GRANULES	7
DESOWEN	72	DIPENTUM	93	<i>econazole</i>	73
<i>desoximetasone</i>	72, 73	<i>diphenoxylate-atropine</i>	93	EDARBI	61
<i>desvenlafaxine</i>	31	DIPROLENE		EDARBYCLOR	61
<i>desvenlafaxine succinate</i>	31	(AUGMENTED)	73	EDECRIIN	61
DETROL	122	<i>dipyridamole</i>	61	EDLUAR	33

EDURANT	7	ENULOSE	93	<i>eszopiclone</i>	34
<i>efavirenz</i>	7	ENVARUS XR	17	<i>ethacrynic acid</i>	62
<i>efavirenz-emtricitabin-tenofov</i>	7	EPCLUSA	7, 8	<i>ethambutol</i>	8
<i>efavirenz-lamivu-tenofov disop</i>	7	EPIDIOLEX	33	<i>ethosuximide</i>	34
EFFEXOR XR	33	EPIDUO	73	<i>ethynodiol diac-eth estradiol</i>	108
EFFIENT	61	EPIDUO FORTE	73	<i>etodolac</i>	34
EFUDEX	73	<i>epinastine</i>	114	<i>etonogestrel-ethinyl estradiol</i> ...	108
EGRIFTA SV	98	<i>epinephrine</i>	118	<i>etravirine</i>	8
<i>electrolyte-148</i>	124	EIPEN 2-PAK	118	EUCRISA	73
ELESTRIN	108	EIPEN JR 2-PAK	118	EUTHYROX	85
<i>eletriptan</i>	33	EPITOL	33	EVAMIST	108
ELIDEL	73	EPIVIR	8	EVEKEO	34
ELIGARD	17	<i>eplerenone</i>	62	EVENITY	104
ELIGARD (3 MONTH)	17	EPOGEN	99	<i>everolimus (antineoplastic)</i>	18
ELIGARD (4 MONTH)	17	EPRONTIA	33	<i>everolimus</i>	
ELIGARD (6 MONTH)	17	EPSOLAY	73	<i>(immunosuppressive)</i>	18
ELIQUIS	61, 62	EPZICOM	8	EVISTA	104
ELIQUIS DVT-PE TREAT		EQUETRO	33	EVOTAZ	8
30D START	61	ERAXIS(WATER DILUENT) ..	8	EVOXAC	80
ELMIRON	122	<i>ergoloid</i>	34	EVRYSDI	34
ELURYNG	108	<i>ergotamine-caffeine</i>	34	EXELDERM	73
ELYXYB	33	ERIVEDGE	17	EXELON PATCH	34
EMEND	93	ERLEADA	17	<i>exemestane</i>	18
EMFLAZA	85	<i>erlotinib</i>	18	EXFORGE	62
EMGALITY PEN	33	ERMEZA	85	EXFORGE HCT	62
EMGALITY SYRINGE	33	ERRIN	108	EXJADE	80
EMSAM	33	ERTACZO	73	EXSERVAN	80
<i>emtricitabine</i>	7	<i>ertapenem</i>	8	EYSUVIS	114
<i>emtricitabine-tenofovir (tdf)</i>	7	ERY PADS	73	EZALLOR SPRINKLE	62
EMTRIVA	7	ERYGEL	73	<i>ezetimibe</i>	62
EMVERM	7	ERYPED 200	8	<i>ezetimibe-simvastatin</i>	62
<i>enalapril maleate</i>	62	ERYPED 400	8	FABHALTA	80
<i>enalapril-hydrochlorothiazide</i>	62	ERY-TAB	8	FABIOR	74
ENBREL	104	ERYTHROCIN	8	FALMINA (28)	108
ENBREL MINI	104	ERYTHROCIN (AS		<i>famciclovir</i>	8
ENBREL SURECLICK	104	STEARATE)	8	<i>famotidine</i>	93
ENDARI	80	<i>erythromycin</i>	8, 114	FANAPT	34
ENDOCET	33	<i>erythromycin ethylsuccinate</i>	8	FARESTON	18
ENGERIX-B (PF)	98	<i>erythromycin with ethanol</i>	73	FARXIGA	85
ENGERIX-B PEDIATRIC		<i>erythromycin-benzoyl peroxide</i> ...73		FASENRA	118
(PF)	99	ESBRIET	118	FASENRA PEN	118
ENILLORING	108	<i>escitalopram oxalate</i>	34	<i>febuxostat</i>	104
<i>enoxaparin</i>	62	ESGIC	34	<i>felbamate</i>	34
ENPRESSE	108	<i>esomeprazole magnesium</i>	93	FELBATOL	34
ENSKYCE	108	ESTARYLLA	108	<i>felodipine</i>	62
ENSPRYNG	17	<i>estazolam</i>	34	FEMARA	18
ENSTILAR	73	ESTRACE	108	FEMRING	108
<i>entacapone</i>	33	<i>estradiol</i>	108	<i>fenofibrate</i>	62
ENTADFI	122	<i>estradiol valerate</i>	108	<i>fenofibrate micronized</i>	62
<i>entecavir</i>	7	<i>estradiol-norethindrone acet</i>	108	<i>fenofibrate nanocrystallized</i>	62
ENTRESTO	62	ESTRING	108	<i>fenofibric acid (choline)</i>	62
ENTYVIO PEN	93	ESTROGEL	108	FENOGLIDE	62

<i>fenoprofen</i>	34	<i>fluoxetine (pmd)</i>	35	<i>gatifloxacin</i>	114
<i>fentanyl</i>	35	<i>fluphenazine decanoate</i>	36	GATTEX 30-VIAL	93
<i>fentanyl citrate</i>	34, 35	<i>fluphenazine hcl</i>	36	GAUZE PAD	102
FENTORA	35	<i>flurandrenolide</i>	74	GAVILYTE-C	93
FERRIPROX	80	<i>flurazepam</i>	36	GAVILYTE-G	93
FERRIPROX (2 TIMES A DAY)	80	<i>flurbiprofen</i>	36	GAVRETO	18
<i>fesoterodine</i>	122	<i>flurbiprofen sodium</i>	114	<i>gefitinib</i>	18
FETZIMA	35	<i>fluticasone furoate-vilanterol</i> ... 118		<i>gemfibrozil</i>	63
FEXMID	35	<i>fluticasone propionate</i> 74, 118, 119		GEMMILY	108
FIASP FLEXTOUCH U-100 INSULIN	85	<i>fluticasone propion-salmeterol</i> . 119		GEMTESA	123
FIASP PENFILL U-100 INSULIN	85	<i>fluvastatin</i>	62	GENERLAC	93
FIASP U-100 INSULIN	85	<i>fluvoxamine</i>	36	GENGRAF	18
FILSPARI	62	FML FORTE	114	GENOTROPIN	99
FILSUVEZ	74	FML LIQUIFILM	114	GENOTROPIN MINIQUICK . 99	
FINACEA	74	FOCALIN	36	<i>gentamicin</i>9, 74, 114	
<i>finasteride</i>	122	FOCALIN XR	36	<i>gentamicin in nacl (iso-osm)</i>9	
<i>finingolimod</i>	35	<i>fondaparinux</i>	62	GENVOYA	9
FINTEPLA	35	FORFIVO XL	36	GEODON	36
FINZALA	108	<i>formoterol fumarate</i>	119	GILENYA	36
FIORICET	35	FORTEO	104	GILOTRIF	18
FIORICET WITH CODEINE . 35		FOSAMAX	104	GIMOTI	93
FIRAZYR	118	FOSAMAX PLUS D	104	GLASSIA	80
FIRDAPSE	35	<i>fosamprenavir</i>	8	<i>glatiramer</i>	36
FIRMAGON KIT W DILUENT SYRINGE	18	<i>fosfomycin tromethamine</i>	8	GLATOPA	36
FIRVANQ	8	<i>fosinopril</i>	62	GLEEVEC	18
FLAC OTIC OIL	82	<i>fosinopril-hydrochlorothiazide</i> ... 62		GLEOSTINE	18
FLAGYL	8	FOSRENOL	80	<i>glimepiride</i>	85
FLAREX	114	FOTIVDA	18	<i>glipizide</i>	85
<i>flavoxate</i>	122	FRAGMIN	62, 63	<i>glipizide-metformin</i>	85
<i>flecainide</i>	62	FROVA	36	GLOPERBA	104
FLECTOR	35	<i>frovatriptan</i>	36	GLUCAGON EMERGENCY KIT (HUMAN)	85
FLEQSUVY	35	FRUZAQLA	18	GLUCOTROL XL	85
FLOLIPID	62	FULPHILA	99	GLUMETZA	85
FLOMAX	123	FUROSCIX	63	<i>glyburide</i>	85
<i>fluconazole</i>	8	<i>furosemide</i>	63	<i>glyburide micronized</i>	85
<i>fluconazole in nacl (iso-osm)</i>	8	FUZEON	8	<i>glyburide-metformin</i>	85
<i>flucytosine</i>	8	FYAVOLV	108	GLYCATE	93
<i>fludrocortisone</i>	85	FYCOMPA	36	<i>glycopyrrolate</i>	93
<i>flunisolide</i>	118	FYLNETRA	99	GLYXAMBI	85
<i>fluocinolone</i>	74	<i>gabapentin</i>	36	GOCOVRI	37
<i>fluocinolone acetonide oil</i>	82	GALAFOLD	85	GOLYTELY	93
<i>fluocinolone and shower cap</i>	74	<i>galantamine</i>	36	GRALISE	37
<i>fluocinonide</i>	74	GAMMAGARD LIQUID	99	<i>granisetron hcl</i>	93
<i>fluocinonide-emollient</i>	74	GAMMAGARD S-D (IGA < 1 MCG/ML)	99	GRANIX	99
<i>fluoride (sodium)</i>	124	GAMMAKED	99	GRASTEK	99
<i>fluorometholone</i>	114	GAMMAPLEX	99	<i>griseofulvin microsize</i>	9
<i>fluorouracil</i>	74	GAMMAPLEX (WITH SORBITOL)	99	<i>griseofulvin ultramicrosize</i>	9
<i>fluoxetine</i>	35, 36	GAMUNEX-C	99	<i>guanfacine</i>	37, 63
		GARDASIL 9 (PF)	99	GVOKE	85
		GASTROCROM	93	GVOKE HYPOPEN 2-PACK . 85	

GVOKE PFS 1-PACK			
SYRINGE	85	HUMIRA(CF) PEN	
GYNAZOLE-1	108	CROHNS-UC-HS	104
HADLIMA	104	HUMIRA(CF) PEN	
HADLIMA PUSH TOUCH ...	104	PEDIATRIC UC	104
HADLIMA(CF)	104	HUMIRA(CF) PEN PSOR-	
HADLIMA(CF)		UV-ADOL HS	104
PUSH TOUCH	104	HUMULIN 70/30 U-100	
HAEGARDA	119	INSULIN	86
HAILEY 24 FE	108	HUMULIN 70/30 U-100	
<i>halcinonide</i>	74	KWIKPEN	86
HALCION	37	HUMULIN N NPH INSULIN	
HALDOL DECANOATE	37	KWIKPEN	86
<i>halobetasol propionate</i>	74	HUMULIN N NPH U-100	
HALOETTE	108	INSULIN	86
HALOG	74	HUMULIN R REGULAR U-	
<i>haloperidol</i>	37	100 INSULN	86
<i>haloperidol decanoate</i>	37	HUMULIN R U-500 (CONC)	
<i>haloperidol lactate</i>	37	INSULIN	86
HARVONI	9	HUMULIN R U-500 (CONC)	
HAVRIX (PF)	99	KWIKPEN	86
HEATHER	108	<i>hydralazine</i>	63
HELIDAC	94	HYDREA	18
HEMADY	85	<i>hydrochlorothiazide</i>	63
<i>heparin (porcine)</i>	63	<i>hydrocodone bitartrate</i>	37
HEPLISAV-B (PF)	99	<i>hydrocodone-acetaminophen</i>	37
HETLIOZ	37	<i>hydrocodone-ibuprofen</i>	37
HETLIOZ LQ	37	<i>hydrocortisone</i>	74, 86, 94
HIBERIX (PF)	99	<i>hydrocortisone butyrate</i>	74
HIPREX	9	<i>hydrocortisone valerate</i>	75
HORIZANT	37	<i>hydrocortisone-acetic acid</i>	83
HULIO(CF)	104	<i>hydrocortisone-pramoxine</i>	94
HULIO(CF) PEN	104	<i>hydromorphone</i>	38
HUMALOG JUNIOR		<i>hydromorphone (pf)</i>	38
KWIKPEN U-100	85	<i>hydroxychloroquine</i>	9
HUMALOG KWIKPEN		<i>hydroxyurea</i>	18
INSULIN	85	<i>hydroxyzine hcl</i>	119
HUMALOG MIX 50-50		<i>hydroxyzine pamoate</i>	119
KWIKPEN	85	HYFTOR	75
HUMALOG MIX 75-25		HYRIMOZ PEN CROHN'S-	
KWIKPEN	85	UC STARTER	104
HUMALOG MIX 75-25(U-		HYRIMOZ PEN PSORIASIS	
100)INSULN	85	STARTER	104
HUMALOG TEMPO PEN(U-		HYRIMOZ(CF)	105
100)INSULN	86	HYRIMOZ(CF) PEDI	
HUMALOG U-100 INSULIN ..	86	CROHN STARTER	104, 105
HUMATIN	9	HYRIMOZ(CF) PEN	105
HUMATROPE	99	HYSINGLA ER	38
HUMIRA	104	HYZAAR	63
HUMIRA PEN	104	<i>ibandronate</i>	105
HUMIRA(CF)	104	IBRANCE	18
HUMIRA(CF) PEN	104	IBSRELA	94
		IBU	38
		<i>ibuprofen</i>	38
		<i>ibuprofen-famotidine</i>	38
		<i>icatibant</i>	119
		ICLEVIA	108
		ICLUSIG	18
		<i>icosapent ethyl</i>	63
		IDACIO(CF)	105
		IDACIO(CF) PEN	105
		IDACIO(CF) PEN CROHN-	
		UC STARTR	105
		IDACIO(CF) PEN	
		PSORIASIS START	105
		IDHIFA	18
		ILEVRO	114
		ILUMYA	75
		<i>imatinib</i>	18
		IMBRUVICA	18, 19
		<i>imipenem-cilastatin</i>	9
		<i>imipramine hcl</i>	38
		<i>imipramine pamoate</i>	38
		<i>imiquimod</i>	75
		IMITREX	38
		IMITREX STATDOSE PEN ..	38
		IMITREX STATDOSE	
		REFILL	38
		IMOVAX RABIES	
		VACCINE (PF)	99
		IMPAVIDO	9
		IMURAN	19
		IMVEXXY MAINTENANCE	
		PACK	108
		IMVEXXY STARTER PACK	
		108
		INBRIJA	38
		INCASSIA	108
		INCRELEX	80
		INCRUSE ELLIPTA	119
		<i>indapamide</i>	63
		INDERAL LA	63
		INDOCIN	38
		<i>indomethacin</i>	38
		INFANRIX (DTAP) (PF)	99
		INGREZZA	38
		INGREZZA INITIATION	
		PK(TARDIV)	38
		INGREZZA SPRINKLE	39
		INLYTA	19
		INNOPRAN XL	63
		INPEFA	86
		INQOVI	19
		INREBIC	19
		INSPIRA	63

<i>insulin asp prt-insulin aspart</i>	86	JADENU SPRINKLE	80	KIPROFEN	40
<i>insulin aspart u-100</i>	86	JAKAFI	19	KISQALI	19
<i>insulin degludec</i>	86	JANTOVEN	64	KISQALI FEMARA CO-	
<i>insulin glargine u-300 conc</i>	86	JANUMET	86	PACK	19
<i>insulin glargine-yfgn</i>	86	JANUMET XR	86	KITABIS PAK	9
<i>insulin lispro</i>	86	JANUVIA	86	KLARON	75
<i>insulin lispro protamin-lispro</i>	86	JARDIANCE	86	KLISYRI	19
<i>insulin syringe-needle u-100</i>	102	JASMIEL (28)	109	KLONOPIN	40
INTELENCE	9	JATENZO	86	KLOR-CON	124
INTRALIPID	124	JAVYGTOR	87	KLOR-CON 10	124
INTRAROSA	108	JAYPIRCA	19	KLOR-CON 8	124
INTROVALE	109	JENTADUETO	87	KLOR-CON M10	124
INTUNIV ER	39	JENTADUETO XR	87	KLOR-CON M15	124
INVANZ	9	JINTELI	109	KLOR-CON M20	124
INVEGA	39	JOENJA	80	KLOXXADO	40
INVEGA HAFYERA	39	JORNAY PM	39	KONVOMEF	94
INVEGA SUSTENNA	39	JOYEAUX	109	KORLYM	87
INVEGA TRINZA	39	JUBLIA	75	KOSELUGO	19
INVELTYS	114	JULEBER	109	KOURZEQ	83
INVOKAMET	86	JULUCA	9	KRAZATI	19
INVOKAMET XR	86	JUNEL 1.5/30 (21)	109	KRINTAFEL	9
INVOKANA	86	JUNEL 1/20 (21)	109	KRISTALOSE	94
IOPIDINE	114	JUNEL FE 1.5/30 (28)	109	KURVELO (28)	109
IPOL	99	JUNEL FE 1/20 (28)	109	KUVAN	87
<i>ipratropium bromide</i>	83, 119	JUNEL FE 24	109	KYLEENA	109
<i>ipratropium-albuterol</i>	119	JUXTAPID	64	<i>l norgest/e.estradiol-e.estrad</i>	109
<i>irbesartan</i>	63	JYLAMVO	19	<i>labetalol</i>	64
<i>irbesartan-hydrochlorothiazide</i> ..	63	JYNARQUE	87	<i>lacosamide</i>	40
IRESSA	19	JYNNEOS (PF)	99	LACRISERT	114
ISENTRESS	9	KAITLIB FE	109	<i>lactulose</i>	94
ISENTRESS HD	9	KALETRA	9	LAGEVRIO (EUA)	9
ISIBLOOM	109	KALYDECO	119	LAMICTAL	40
ISOLYTE S PH 7.4	124	KARIVA (28)	109	LAMICTAL ODT	40
ISOLYTE-P IN 5 %		KATERZIA	64	LAMICTAL STARTER	
DEXTROSE	124	KAZANO	87	(BLUE) KIT	40
<i>isoniazid</i>	9	KELNOR 1/35 (28)	109	LAMICTAL STARTER	
ISORDIL	63	KELNOR 1/50 (28)	109	(GREEN) KIT	40
ISORDIL TITRADOSE	63	KENALOG	75	LAMICTAL STARTER	
<i>isosorbide dinitrate</i>	63	KEPPRA	39	(ORANGE) KIT	40
<i>isosorbide mononitrate</i>	63	KEPPRA XR	39	LAMICTAL XR	40
<i>isosorbide-hydralazine</i>	63	KERENDIA	64	LAMICTAL XR STARTER	
<i>isotretinoin</i>	75	KESIMPTA PEN	39	(BLUE)	40
<i>isradipine</i>	63	<i>ketoconazole</i>	9, 75	LAMICTAL XR STARTER	
ISTALOL	114	KETODAN	75	(GREEN)	40
ISTURISA	86	<i>ketoprofen</i>	39	LAMICTAL XR STARTER	
<i>itraconazole</i>	9	<i>ketorolac</i>	40, 114	(ORANGE)	40
<i>ivermectin</i>	9, 75	KEVEYIS	40	<i>lamivudine</i>	10
IWILFIN	19	KEVZARA	105	<i>lamivudine-zidovudine</i>	10
IXCHIQ (PF)	99	KINERET	105	<i>lamotrigine</i>	40
IXIARO (PF)	99	KINRIX (PF)	100	LAMPIT	10
IYUZEH (PF)	114	KIONEX (WITH		LANOXIN	64
JADENU	80	SORBITOL)	80	<i>lansoprazole</i>	94

<i>lanthanum</i>	80	LIBRAX (WITH CLIDINIUM)	94	LOREEV XR	41
LANTUS SOLOSTAR U-100 INSULIN	87	LICART	41	LORYNA (28)	109
LANTUS U-100 INSULIN	87	<i>lidocaine</i>	75	LORZONE	41
<i>lapatinib</i>	19	<i>lidocaine hcl</i>	75	<i>losartan</i>	64
LARIN 1.5/30 (21)	109	LIDOCAINE VISCOUS	75	<i>losartan-hydrochlorothiazide</i>	64
LARIN 1/20 (21)	109	<i>lidocaine-prilocaine</i>	75	LOTEMAX	115
LARIN FE 1.5/30 (28)	109	LIDOCAN III	75	LOTEMAX SM	115
LARIN FE 1/20 (28)	109	LIDODERM	75	LOTENSIN	64
LASIX	64	LILETTA	109	<i>loteprednol etabonate</i>	115
<i>latanoprost</i>	114	<i>linezolid</i>	10	LOTREL	64
LATUDA	40	<i>linezolid in dextrose 5%</i>	10	LOTRONEX	94
LAYOLIS FE	109	LINZESS	94	<i>lovastatin</i>	64
<i>ledipasvir-sofosbuvir</i>	10	<i>liothyronine</i>	87	LOVAZA	64
LEENA 28	109	LIPITOR	64	LOVENOX	64
<i>leflunomide</i>	105	LIPOFEN	64	LOW-OGESTREL (28)	109
<i>lenalidomide</i>	19	LIQREV	119	<i>loxapine succinate</i>	41
LENVIMA	19	<i>lisdexamfetamine</i>	41	<i>lubiprostone</i>	94
LESCOL XL	64	<i>lisinopril</i>	64	LUCEMYRA	41
LESSINA	109	<i>lisinopril-hydrochlorothiazide</i>	64	<i>luliconazole</i>	75
LETAIRIS	119	LITFULO	80	LUMAKRAS	20
<i>letrozole</i>	19	<i>lithium carbonate</i>	41	LUMIGAN	115
<i>leucovorin calcium</i>	19	<i>lithium citrate</i>	41	LUMRYZ	41
LEUKERAN	19	LITHOBID	41	LUNESTA	41
LEUKINE	100	LITHOSTAT	80	LUPKYNIS	20
<i>leuprolide</i>	19	LIVALO	64	LUPRON DEPOT	20
<i>leuprolide (3 month)</i>	19	LIVMARLI	94	LUPRON DEPOT (3 MONTH)	20
<i>levabuterol hcl</i>	119	LIVTENCITY	10	LUPRON DEPOT (4 MONTH)	20
<i>levabuterol tartrate</i>	119	LO LOESTRIN FE	109	LUPRON DEPOT (6 MONTH)	20
<i>levamlodipine</i>	64	LOCOID	75	LUPRON DEPOT (6 MONTH)	20
LEVEMIR FLEXPEN	87	LOCOID LIPOCREAM	75	LUPRON DEPOT-PED	20
LEVEMIR U-100 INSULIN	87	LODINE	41	LUPRON DEPOT-PED (3 MONTH)	20
<i>levetiracetam</i>	41	LODOCO	64	<i>lurasidone</i>	41
<i>levobunolol</i>	114	LODOSYN	41	LUTERA (28)	110
<i>levocarnitine</i>	80	LOESTRIN 1.5/30 (21)	109	LUZU	75
<i>levocarnitine (with sugar)</i>	80	LOESTRIN 1/20 (21)	109	LYBALVI	41
<i>levocetirizine</i>	119	LOESTRIN FE 1.5/30 (28-DAY)	109	LYLEQ	110
<i>levofloxacin</i>	10, 115	LOESTRIN FE 1/20 (28-DAY)	109	LYLLANA	110
<i>levofloxacin in d5w</i>	10	LOFENA	41	LYNPARZA	20
LEVONEST (28)	109	LOKELMA	80	LYRICA	41
<i>levonorgestrel-ethinyl estrad</i>	109	LOMOTIL	94	LYRICA CR	41
<i>levonorg-eth estrad triphasic</i>	109	LONSURF	19	LYSODREN	20
LEVORA-28	109	<i>loperamide</i>	94	LYTGOBI	20
<i>levorphanol tartrate</i>	41	LOPID	64	LYUMJEV KWIKPEN U-100 INSULIN	87
<i>levothyroxine</i>	87	<i>lopinavir-ritonavir</i>	10	LYUMJEV KWIKPEN U-200 INSULIN	87
LEVOXYL	87	LOPRESSOR	64	LYUMJEV TEMPO PEN(U-100)INSULN	87
LEXAPRO	41	LOPROX	75	LYUMJEV U-100 INSULIN	87
LEXETTE	75	<i>lorazepam</i>	41		
LEXIVA	10	LORAZEPAM INTENSOL	41		
LIALDA	94	LORBRENA	20		
LIBERVANT	41				

LYVISPAH	41	MENACTRA (PF)	100	MICARDIS HCT	65
LYZA	110	MENEST	110	MICONAZOLE-3	110
MACROBID	10	MENOSTAR	110	MICROGESTIN 1.5/30 (21) ..	110
MACRODANTIN	10	MENQUADFI (PF)	100	MICROGESTIN 1/20 (21)	110
<i>mafenide acetate</i>	75	MENVEO A-C-Y-W-135-DIP		MICROGESTIN 24 FE	110
<i>magnesium sulfate</i>	124	(PF)	100	MICROGESTIN FE 1.5/30	
MALARONE	10	<i>meperidine</i>	42	(28)	110
MALARONE PEDIATRIC	10	<i>meperidine (pf)</i>	42	MICROGESTIN FE 1/20 (28)	
<i>malathion</i>	75	<i>meprobamate</i>	42	110
<i>maraviroc</i>	10	MEPRON	10	<i>midodrine</i>	80
MARINOL	94	<i>mercaptapurine</i>	20	MIEBO (PF)	115
MARLISSA (28)	110	<i>meropenem</i>	10	<i>mifepristone</i>	88
MARPLAN	42	MERZEE	110	MIGERGOT	43
MATULANE	20	<i>mesalamine</i>	94, 95	<i>miglitol</i>	88
MATZIM LA	64	MESNEX	20	<i>miglustat</i>	88
MAVENCLAD (10 TABLET		MESTINON	42	MIGRANAL	43
PACK)	42	MESTINON TIMESPAN	42	MILI	110
MAVENCLAD (4 TABLET		METADATE CD	42	MIMVEY	110
PACK)	42	<i>metaxalone</i>	42	MINIVELLE	110
MAVENCLAD (5 TABLET		<i>metformin</i>	87, 88	<i>minocycline</i>	10, 11
PACK)	42	<i>methadone</i>	42, 43	<i>minoxidil</i>	65
MAVENCLAD (6 TABLET		<i>methamphetamine</i>	43	<i>mirabegron</i>	123
PACK)	42	<i>methazolamide</i>	115	MIRENA	110
MAVENCLAD (7 TABLET		<i>methenamine hippurate</i>	10	<i>mirtazapine</i>	43
PACK)	42	<i>methimazole</i>	88	MIRVASO	76
MAVENCLAD (8 TABLET		METHITEST	88	<i>misoprostol</i>	95
PACK)	42	<i>methocarbamol</i>	43	MITIGARE	105
MAVENCLAD (9 TABLET		<i>methotrexate sodium</i>	20	M-M-R II (PF)	100
PACK)	42	<i>methotrexate sodium (pf)</i>	20	<i>modafinil</i>	43
MAVYRET	10	<i>methoxsalen</i>	75	<i>moexipril</i>	65
MAXALT	42	<i>methscopolamine</i>	95	<i>molindone</i>	43
MAXALT-MLT	42	<i>methsuximide</i>	43	<i>mometasone</i>	76, 119
MAXIDEX	115	METHYLIN	43	<i>montelukast</i>	120
MAXITROL	115	<i>methylphenidate</i>	43	<i>morphine</i>	44
MAYZENT	42	<i>methylphenidate hcl</i>	43	<i>morphine concentrate</i>	43
MAYZENT STARTER(FOR		<i>methylprednisolone</i>	88	MOTEGRITY	95
1MG MAINT)	42	<i>methyltestosterone</i>	88	MOTPOLY XR	44
MAYZENT STARTER(FOR		<i>metoclopramide hcl</i>	95	MOUNJARO	88
2MG MAINT)	42	<i>metolazone</i>	64	MOVANTIK	95
<i>meclizine</i>	94	<i>metoprolol succinate</i>	65	MOVIPREP	95
<i>meclofenamate</i>	42	<i>metoprolol ta-hydrochlorothiaz</i> ..	65	<i>moxifloxacin</i>	11, 115
MEDROL	87	<i>metoprolol tartrate</i>	65	<i>moxifloxacin-sod.chloride(iso)</i> ...	11
MEDROL (PAK)	87	METROCREAM	75	MS CONTIN	44
<i>medroxyprogesterone</i>	110	METROGEL	75	MULPLETA	65
<i>mefenamic acid</i>	42	METROLOTION	75	MULTAQ	65
<i>mefloquine</i>	10	<i>metronidazole</i>	10, 75, 110	<i>mupirocin</i>	76
<i>megestrol</i>	20	<i>metronidazole in nacl (iso-os)</i>	10	<i>mupirocin calcium</i>	76
MEKINIST	20	<i>metryrosine</i>	65	MYALEPT	88
MEKTOVI	20	<i>mexiletine</i>	65	MYAMBUTOL	11
<i>meloxicam</i>	42	MIBELAS 24 FE	110	MYCAMINE	11
<i>meloxicam submicronized</i>	42	<i>micafungin</i>	10	MYCAPSSA	21
<i>memantine</i>	42	MICARDIS	65	MYCOBUTIN	11

<i>mycophenolate mofetil</i>	21	NESINA	88	<i>norgestimate-ethinyl estradiol</i> ..	111
<i>mycophenolate sodium</i>	21	NEUAC	76	NORITATE	76
MYDAYIS	44	NEULASTA	100	NORLIQVA	66
MYFEMBREE	110	NEUPOGEN	100	NORPACE	66
MYFORTIC	21	NEUPRO	45	NORPACE CR	66
MYHIBBIN	21	NEURONTIN	45	NORPRAMIN	45
MYRBETRIQ	123	NEVANAC	115	NORTHERA	80, 81
MYSOLINE	44	<i>nevirapine</i>	11	NORTREL 0.5/35 (28)	111
MYTESI	95	NEXAVAR	21	NORTREL 1/35 (21)	111
<i>nabumetone</i>	44	NEXICLON XR	65	NORTREL 1/35 (28)	111
<i>nadolol</i>	65	NEXIUM	95	NORTREL 7/7/7 (28)	111
<i>nafticillin</i>	11	NEXIUM PACKET	95	<i>nortriptyline</i>	45
<i>naftifine</i>	76	NEXLETOL	65	NORVASC	66
NAFTIN	76	NEXLIZET	65	NORVIR	11
NALFON	44	NEXPLANON	110	NOURIANZ	45
NALOCET	44	NEXTSTELLIS	110	NOVOLIN 70/30 U-100	
<i>naloxone</i>	44	NGENLA	100	INSULIN	88
<i>naltrexone</i>	44	<i>niacin</i>	65	NOVOLIN 70-30 FLEXPEN	
NAMENDA TITRATION		NIACOR	65	U-100	88
PAK	44	<i>nicardipine</i>	65	NOVOLIN N FLEXPEN	88
NAMENDA XR	44	NICOTROL	80	NOVOLIN N NPH U-100	
NAMZARIC	44	NICOTROL NS	80	INSULIN	88
NAPRELAN CR	45	<i>nifedipine</i>	65	NOVOLIN R FLEXPEN	88
NAPROSYN	45	NIKKI (28)	110	NOVOLIN R REGULAR	
<i>naproxen</i>	45	NILANDRON	21	U100 INSULIN	88
<i>naproxen sodium</i>	45	<i>nilutamide</i>	21	NOVOLOG FLEXPEN U-100	
<i>naproxen-esomeprazole</i>	45	<i>nimodipine</i>	65	INSULIN	88
<i>naratriptan</i>	45	NINLARO	21	NOVOLOG MIX 70-30 U-100	
NARDIL	45	<i>nisoldipine</i>	65	INSULN	88
NATACYN	115	<i>nitazoxanide</i>	11	NOVOLOG MIX 70-	
NATAZIA	110	<i>nitisinone</i>	80	30FLEXPEN U-100	88
<i>nateglinide</i>	88	NITRO-BID	65	NOVOLOG PENFILL U-100	
NATESTO	88	NITRO-DUR	65	INSULIN	88
NATROBA	76	<i>nitrofurantoin</i>	11	NOVOLOG U-100 INSULIN	
NAYZILAM	45	<i>nitrofurantoin macrocrystal</i>	11	ASPART	88
<i>nebivolol</i>	65	<i>nitrofurantoin monohyd/m-cryst.</i>	11	NOXAFIL	11
NEBUPENT	11	<i>nitroglycerin</i>	65, 95	NUBEQA	21
NECON 0.5/35 (28)	110	NITROLINGUAL	66	NUCALA	120
<i>nefazodone</i>	45	NITROSTAT	66	NUCYNTA	45
<i>neomycin</i>	11	NITYR	80	NUCYNTA ER	45
<i>neomycin-bacitracin-poly-hc</i>	115	NIVESTYM	100	NUEDEXTA	45
<i>neomycin-bacitracin-polymyxin</i>	115	<i>nizatidine</i>	95	NUPLAZID	45
<i>neomycin-polymyxin b-</i>		NORA-BE	110	NURTEC ODT	46
<i>dexameth</i>	115	NORDITROPIN FLEXPEN	100	NUTRILIPID	124
<i>neomycin-polymyxin-gramicidin</i>		<i>norelgestromin-ethin.estradiol.</i>	110	NUTROPIN AQ NUSPIN	100
.....	115	<i>noreth-ethinyl estradiol-iron</i>	110	NUVARING	111
<i>neomycin-polymyxin-hc</i>	83, 115	<i>norethindrone (contraceptive)</i> ..	110	NUVESSA	111
NEO-POLYCIN	115	<i>norethindrone acetate</i>	110	NUVIGIL	46
NEO-POLYCIN HC	115	<i>norethindrone ac-eth estradiol.</i>	110	NUZYRA	11
NEORAL	21	<i>norethindrone-e.estradiol-iron.</i>	111	NYAMYC	76
NEO-SYNALAR	76	NORGESIC	45	NYLIA 1/35 (28)	111
NERLYNX	21	NORGESIC FORTE	45	NYLIA 7/7/7 (28)	111

NYMALIZE	66	ORENITRAM MONTH 1	<i>paricalcitol</i>	89	
NYMYO	111	TITRATION KT	66	PARLODEL	47
<i>nystatin</i>	11, 76	ORENITRAM MONTH 2	PARNATE	47	
<i>nystatin-triamcinolone</i>	76	TITRATION KT	66	<i>paroxetine hcl</i>	47
NYSTOP	76	ORENITRAM MONTH 3	<i>paroxetine</i>		
NYVEPRIA	100	TITRATION KT	66	<i>mesylate(menop.sym)</i>	47
OALIVA	95	ORFADIN	81	PAXIL	47
OCELLA	111	ORGOVYX	21	PAXIL CR	47
OCTAGAM	100	ORIAHNN	111	PAXLOVID	12
<i>octreotide acetate</i>	21	ORILISSA	88	<i>pazopanib</i>	21
OCUFLOX	115	ORKAMBI	120	PEDIARIX (PF)	100
ODACTRA	100	ORLADEYO	120	PEDVAX HIB (PF)	100
ODEFSEY	11	ORMALVI	46	<i>peg 3350-electrolytes</i>	95
ODOMZO	21	<i>orphenadrine citrate</i>	46	<i>peg3350-sod sul-nacl-kcl-asb-c.</i>	95
OFEV	120	<i>orphenadrine-asa-caffeine</i>	46	PEGASYS	100
<i>ofloxacin</i>	11, 83, 115	ORSERDU	21	<i>peg-electrolyte soln</i>	95
OGSIVEO	21	<i>oseltamivir</i>	11	PEMAZYRE	21
OJEMDA	21	OSENI	88	<i>pen needle, diabetic</i>	102
OJJAARA	21	OSMOLEX ER	46	PENBRAYA (PF)	100
<i>olanzapine</i>	46	OSPHENA	111	<i>penciclovir</i>	76
<i>olanzapine-fluoxetine</i>	46	OTEZLA	105	<i>penicillamine</i>	105
<i>olmesartan</i>	66	OTEZLA STARTER	105	<i>penicillin g pot in dextrose</i>	12
<i>olmesartan-amlodipin-hcthiazyd</i>	66	OTOVEL	83	<i>penicillin g potassium</i>	12
<i>olmesartan-hydrochlorothiazide</i>	66	OTREXUP (PF)	105	<i>penicillin g sodium</i>	12
<i>olopatadine</i>	83	OVIDE	76	<i>penicillin v potassium</i>	12
OLPRUVA	81	<i>oxacillin</i>	11	PENNSAID	47
OLUMIANT	105	<i>oxacillin in dextrose(iso-osm)</i>	12	PENTACEL (PF)	100
<i>omega-3 acid ethyl esters</i>	66	<i>oxaprozin</i>	46	PENTAM	12
<i>omeprazole</i>	95	<i>oxazepam</i>	46	<i>pentamidine</i>	12
<i>omeprazole-sodium bicarbonate</i>	95	OXBRYTA	81	PENTASA	95
OMNARIS	120	<i>oxcarbazepine</i>	46	<i>pentazocine-naloxone</i>	47
OMNITROPE	100	OXERVATE	115	<i>pentoxifylline</i>	66
OMVOH	95	<i>oxiconazole</i>	76	PEPCID	96
OMVOH PEN	95	OXISTAT	76	PERCOCET	47
<i>ondansetron</i>	95	OXTELLAR XR	46	PERFOROMIST	120
<i>ondansetron hcl</i>	95	<i>oxybutynin chloride</i>	123	<i>perindopril erbumine</i>	66
ONEXTON	76	<i>oxycodone</i>	46	PERIOGARD	83
ONFI	46	<i>oxycodone-acetaminophen</i>	46, 47	<i>permethrin</i>	76
ONGENTYS	46	OXYCONTIN	47	<i>perphenazine</i>	48
ONUREG	21	<i>oxymorphone</i>	47	<i>perphenazine-amitriptyline</i>	48
ONZETRA XSAIL	46	OXYTROL	123	PERSERIS	48
OPSUMIT	120	OZEMPIC	88	PERTZYE	96
OPSYNVI	120	OZOBAX DS	47	PHEBURANE	81
OPVEE	46	PACERONE	66	<i>phenelzine</i>	48
OPZELURA	76	<i>paliperidone</i>	47	<i>phenobarbital</i>	48
ORACEA	11	PALYNZIQ	88, 89	<i>phenoxybenzamine</i>	66
ORALAIR	100	PAMELOR	47	PHENYTEK	48
ORAPRED ODT	88	PANCREAZE	95	<i>phenytoin</i>	48
ORENCIA	105	PANDEL	76	<i>phenytoin sodium extended</i>	48
ORENCIA CLICKJECT	105	PANRETIN	76	PHEXXI	111
ORENITRAM	66	<i>pantoprazole</i>	95	PHOSPHOLINE IODIDE	115
		PANZYGA	100	PIFELTRO	12

<i>pilocarpine hcl</i>	81, 115	PRED MILD	115	PROMETHEGAN	120
<i>pimecrolimus</i>	76	<i>prednisolone</i>	89	PROMETRIUM	111
<i>pimozide</i>	48	<i>prednisolone acetate</i>	115	<i>propafenone</i>	67
PIMTREA (28)	111	<i>prednisolone sodium phosphate</i>	89, 115	<i>propranolol</i>	67
<i>pindolol</i>	66	<i>prednisone</i>	89	<i>propylthiouracil</i>	89
<i>pioglitazone</i>	89	PREDNISON	89	PROQUAD (PF)	101
<i>pioglitazone-glimepiride</i>	89	<i>pregabalin</i>	48	PROSCAR	123
<i>pioglitazone-metformin</i>	89	PREHEVBRI (PF).....	100	PROSOL 20 %	125
<i>piperacillin-tazobactam</i>	12	PREMARIN	111	PROTONIX	96
PIQRAY	21, 22	PREMASOL 10 %	125	<i>protriptyline</i>	48
<i>pirfenidone</i>	120	PREMPHASE	111	PROVERA	111
<i>piroxicam</i>	48	PREMPRO	111	PROVIGIL	48
<i>pitavastatin calcium</i>	66	PRENATAL VITAMIN PLUS LOW IRON	125	PROZAC	48
PLAQUENIL	12	<i>pretomanid</i>	12	PRUDOXIN	76
PLASMA-LYTE 148	124	PREVACID	96	PULMICORT	120
PLASMA-LYTE A	124	PREVACID SOLUTAB	96	PULMICORT FLEXHALER	120
PLAVIX	66	PREVALITE	67	PULMOZYME	120
PLEGRIDY	100	PREVYMIS	12	PURIXAN	22
PLENAMINE	124	PREZCOBIX	12	PYLERA	96
PLENVU	96	PREZISTA	12	<i>pyrazinamide</i>	12
PLIAGLIS	76	PRIFTIN	12	<i>pyridostigmine bromide</i>	48
<i>podofilox</i>	76	PRIOSEC	96	<i>pyrimethamine</i>	12
POLYCIN	115	<i>primaquine</i>	12	PYRUKYND	81
<i>polymyxin b sulfate</i>	12	PRIMAXIN IV	12	QBRELIS	67
<i>polymyxin b sulf-trimethoprim</i>	115	<i>primidone</i>	48	QBREXZA	76
POMALYST	22	PRIORIX (PF)	100	QDOLO	48
PONVORY	48	PRISTIQ	48	QELBREE	49
PONVORY 14-DAY STARTER PACK	48	PRIVIGEN	100	QINLOCK	22
PORTIA 28	111	PROAIR RESPICLICK	120	QNASL	120
<i>posaconazole</i>	12	<i>probenecid</i>	105	QTERN	89
<i>potassium chlorid-d5-</i> <i>0.45%nacl</i>	124	<i>probenecid-colchicine</i>	105	QUADRACEL (PF)	101
<i>potassium chloride</i>	125	PROCARDIA XL	67	QUALAQUIN	12
<i>potassium chloride in 0.9%nacl</i>	124	PROCENTRA	48	QUARTETTE	111
<i>potassium chloride in 5 % dex.</i>	124	<i>prochlorperazine</i>	96	QUDEXY XR	49
<i>potassium chloride in lr-d5</i>	124	<i>prochlorperazine maleate</i>	96	QUESTRAN	67
<i>potassium chloride in water</i>	125	PROCRIT	101	QUESTRAN LIGHT	67
<i>potassium chloride-0.45 % nacl</i>	125	PROCTOFOAM HC	96	<i>quetiapine</i>	49
<i>potassium chloride-d5-</i> <i>0.2%nacl</i>	125	PROCTO-MED HC	96	QUILLICHEW ER	49
<i>potassium chloride-d5-</i> <i>0.9%nacl</i>	125	PROCTOSOL HC	96	QUILLIVANT XR	49
<i>potassium citrate</i>	123	PROCTOZONE-HC	96	<i>quinapril</i>	67
PRADAXA	66	PROCYSBI	123	<i>quinidine gluconate</i>	67
PRALUENT PEN	66	<i>progesterone micronized</i>	111	<i>quinidine sulfate</i>	67
<i>pramipexole</i>	48	PROGLYCEM	89	<i>quinine sulfate</i>	12
<i>prasugrel</i>	67	PROGRAF	22	QULIPTA	49
<i>pravastatin</i>	67	PROLASTIN-C	81	QUVIVIQ	49
<i>praziquantel</i>	12	PROLATE	48	QVAR REDHALER	120
<i>prazosin</i>	67	PROLENSA	115	RABAVERT (PF)	101
PRED FORTE	115	PROLIA	105	<i>rabeprazole</i>	96
		PROMACTA	67	RADICAVA ORS STARTER KIT SUSP	49
		<i>promethazine</i>	120	<i>raloxifene</i>	105
				<i>ramelteon</i>	49

<i>ramipril</i>	67	<i>ribavirin</i>	13	SANDOSTATIN	22
<i>ranolazine</i>	67	RIDAURA	106	SANTYL	77
RAPAFLO	123	<i>rifabutin</i>	13	SAPHRIS	51
RAPAMUNE	22	<i>rifampin</i>	13	<i>sapropterin</i>	89
<i>rasagiline</i>	49	<i>riluzole</i>	81	SAVAYSA	67
RASUVO (PF)	106	<i>rimantadine</i>	13	SAVELLA	106
RAVICTI	81	RINVOQ	106	<i>saxagliptin</i>	89
RAYALDEE	89	<i>risedronate</i>	81, 106	<i>saxagliptin-metformin</i>	89
RAYOS	89	RISPERDAL	50	SCSEMBLIX	22
REBIF (WITH ALBUMIN) ... 101		RISPERDAL CONSTA	49, 50	<i>scopolamine base</i>	96
REBIF REBIDOSE	101	<i>risperidone</i>	50	SECUADO	51
REBIF TITRATION PACK .. 101		<i>risperidone microspheres</i>	50	SEGLENTIS	51
RECLIPSEN (28)	111	RITALIN	50	SEGLUROMET	90
RECOMBIVAX HB (PF) 101		RITALIN LA	50	<i>selegiline hcl</i>	51
RECORLEV	89	<i>ritonavir</i>	13	<i>selenium sulfide</i>	77
RECTIV	96	<i>rivastigmine</i>	50	SELZENTRY	13
REGLAN	96	<i>rivastigmine tartrate</i>	50	SEMGLEE(INSULIN	
REGRANEX	76	RIVELSA	111	GLARGINE-YFGN)	90
RELAFEN DS	49	RIVFLOZA	123	SEMGLEE(INSULIN	
RELENZA DISKHALER 12		<i>rizatriptan</i>	50	GLARG-YFGN)PEN	90
RELEUKO	101	ROBINUL	96	SENSIPAR	90
RELEXXII	49	ROBINUL FORTE	96	SEREVENT DISKUS	121
RELISTOR	96	ROCALTROL	89	SEROQUEL	51
RELPAK	49	ROCKLATAN	115	SEROQUEL XR	51
RELTONE	96	<i>roflumilast</i>	121	SEROSTIM	101
REMERON	49	<i>ropinirole</i>	50	<i>sertraline</i>	51
REMERON SOLTAB	49	<i>rosuvastatin</i>	67	SETLAKIN	111
REVELA	81	ROTARIX	101	<i>sevelamer carbonate</i>	81
<i>repaglinide</i>	89	ROTATEQ VACCINE	101	<i>sevelamer hcl</i>	81
REPATHA PUSHTRONEX ... 67		ROWASA	96	SEYSARA	13
REPATHA SURECLICK 67		ROWEEPRA	50	SHAROBEL	111
REPATHA SYRINGE	67	ROXICODONE	51	SHINGRIX (PF)	101
RESTASIS	115	ROXYBOND	51	SIGNIFOR	22
RESTASIS MULTIDOSE 115		ROZEREM	51	SIKLOS	22
RESTORIL	49	ROZLYTREK	22	<i>sildenafil (pulm.hypertension)</i> .. 121	
RETACRIT	101	RUBRACA	22	SILENOR	51
RETEVMO	22	RUCONEST	121	SILIQ	77
RETIN-A	76	<i>rufinamide</i>	51	<i>silodosin</i>	123
RETIN-A MICRO	77	RUKOBIA	13	SILVADENE	77
RETIN-A MICRO PUMP 77		RYALTRIS	121	<i>silver sulfadiazine</i>	77
RETROVIR	12	RYBELSUS	89	SIMBRINZA	116
REVATIO	120	RYCLORA	121	SIMLANDI(CF)	
REVCovi	81	RYDAPT	22	AUTOINJECTOR	106
REVLIMID	22	RYTARY	51	SIMPONI	106
REXULTI	49	RYTHMOL SR	67	<i>simvastatin</i>	67
REYATAZ	13	RYVENT	121	SINEMET	51
REYVOW	49	SABRIL	51	SINGULAIR	121
REZDIFFRA	81	SAFYRAL	111	<i>sirolimus</i>	22
REZLIDHIA	22	SAJAZIR	121	SIRTURO	13
REZUROCK	22	SALAGEN (PILOCARPINE) . 81		<i>sitagliptin</i>	90
REZVOGLAR KWIKPEN 89		SAMSCA	89	<i>sitagliptin-metformin</i>	90
RHOPRESSA	115	SANDIMMUNE	22	SIVEXTRO	13

SKYCLARYS	51	STALEVO 100	51	SYMLINPEN 120	90
SKYLA	111	STALEVO 125	51	SYMLINPEN 60	90
SKYRIZI	77, 96	STALEVO 150	51	SYMPAZAN	52
SKYTROFA	101	STALEVO 200	51	SYMPROIC	97
SLYND	111	STALEVO 50	51	SYMTUZA	14
SOAAZ	67	STALEVO 75	52	SYNALAR	77
<i>sodium chloride</i>	81	STEGLATRO	90	SYNAREL	90
<i>sodium chloride 0.45 %</i>	125	STEGLUJAN	90	SYNDROS	97
<i>sodium chloride 0.9 %</i>	81	STELARA	77	SYNJARDY	90
<i>sodium chloride 3 % hypertonic</i>	125	STIMUFEND	101	SYNJARDY XR	90
<i>sodium chloride 5 % hypertonic</i>	125	STIOLTO RESPIMAT	121	SYNTHROID	90
<i>sodium oxybate</i>	51	STIVARGA	23	SYPRINE	82
<i>sodium phenylbutyrate</i>	81	STRATTERA	52	TABLOID	23
<i>sodium polystyrene sulfonate</i>	81	<i>streptomycin</i>	13	TABRECTA	23
<i>sodium,potassium,mag sulfates</i> ..	96	STRIBILD	13	TACLONEX	77
<i>sofosbuvir-velpatasvir</i>	13	STRIVERDI RESPIMAT	121	<i>tacrolimus</i>	23, 77
SOGROYA	101	STROMECTOL	13	<i>tadalafil</i>	123
SOHONOS	82	SUBOXONE	52	<i>tadalafil (pulm. hypertension)</i> ..	121
<i>solifenacin</i>	123	SUBVENITE	52	TADLIQ	121
SOLQUA 100/33	90	SUBVENITE STARTER (BLUE) KIT	52	TAFINLAR	23
SOLODYN	13	SUBVENITE STARTER (GREEN) KIT	52	<i>tafluprost (pf)</i>	116
SOLOSEC	13	SUBVENITE STARTER (ORANGE) KIT	52	TAGRISO	23
SOLTAMOX	22	SUCRAID	97	TAKHZYRO	121
SOMA	51	<i>sucralfate</i>	97	TALICIA	97
SOMAVERT	90	SUFLAVE	97	TALTZ AUTOINJECTOR	77
SOOLANTRA	77	SULAR	68	TALTZ SYRINGE	77
<i>sorafenib</i>	22	<i>sulfacetamide sodium</i>	116	TALZENNA	23
SORILUX	77	<i>sulfacetamide sodium (acne)</i>	77	TAMIFLU	14
SORINE	67	<i>sulfacetamide-prednisolone</i>	116	<i>tamoxifen</i>	23
<i>sotalol</i>	67	<i>sulfadiazine</i>	13	<i>tamsulosin</i>	123
SOTALOL AF	67	<i>sulfamethoxazole-trimethoprim</i> ..	13	TAPERDEX	90
SOTYKTU	77	SULFAMYLON	77	TARGADOX	14
SOTYLIZE	67	<i>sulfasalazine</i>	97	TARGRETIN	23
SOVALDI	13	<i>sulindac</i>	52	TARINA 24 FE	112
SOVUNA	13	<i>sumatriptan</i>	52	TARINA FE 1-20 EQ (28)	112
SPEVIGO	77	<i>sumatriptan succinate</i>	52	TARPEYO	90
<i>spinosad</i>	77	<i>sumatriptan-naproxen</i>	52	TASCENSO ODT	52
SPIRIVA RESPIMAT	121	<i>sunitinib malate</i>	23	TASIGNA	23
SPIRIVA WITH HANDIHALER	121	SUNLENCA	13	<i>tasimelteon</i>	52
<i>spironolactone</i>	68	SUNOSI	52	TASMAR	52
<i>spironolacton-hydrochlorothiaz.</i>	68	SUPREP BOWEL PREP KIT	97	<i>tavorole</i>	77
SPORANOX	13	SUTAB	97	TAVALISSE	68
SPRINTEC (28)	112	SUTENT	23	TAVNEOS	82
SPRITAM	51	SYEDA	112	<i>tazarotene</i>	77, 78
SPRIX	51	SYMBICORT	121	TAZICEF	14
SPRYCEL	22	SYMBYAX	52	TAZORAC	78
SPS (WITH SORBITOL)	82	SYMDEKO	121	TAZVERIK	23
SRONYX	112	SYMFI	13	TDVAX	101
SSD	77	SYMFI LO	13	TECFIDERA	52, 53
				TEFLARO	14
				TEGLUTIK	82
				TEGRETOL	53

TEGRETOL XR	53	TIROSINT-SOL	90	TRESIBA FLEXTOUCH U-	
TEGSEDI	53	TIVICAY	14	100	91
TEKTURNA	68	TIVICAY PD	14	TRESIBA FLEXTOUCH U-	
<i>telmisartan</i>	68	<i>tizanidine</i>	53	200	91
<i>telmisartan-amlodipine</i>	68	TLANDO	91	TRESIBA U-100 INSULIN	91
<i>telmisartan-hydrochlorothiazid</i> ..	68	TOBI	14	<i>tretinoin</i>	78
<i>temazepam</i>	53	TOBI PODHALER	14	<i>tretinoin (antineoplastic)</i>	23
TENCON	53	TOBRADEX	116	<i>tretinoin microspheres</i>	78
TENIVAC (PF)	101	TOBRADEX ST	116	TREXALL	23
<i>tenofovir disoproxil fumarate</i>	14	<i>tobramycin</i>	14, 116	TREXIMET	54
TENORETIC 100	68	<i>tobramycin in 0.225 % nacl</i>	14	TREZIX	54
TENORETIC 50	68	<i>tobramycin sulfate</i>	14	<i>triamcinolone acetonide</i>	78, 83
TENORMIN	68	<i>tobramycin-dexamethasone</i>	116	<i>triamterene</i>	68
TEPMETKO	23	TOBREX	116	<i>triamterene-hydrochlorothiazid</i> ..	68
<i>terazosin</i>	68	<i>tolcapone</i>	53	<i>triazolam</i>	54
<i>terbinafine hcl</i>	14	TOLECTIN 600	53	TRIBENZOR	68
<i>terbutaline</i>	121	<i>tolmetin</i>	53	TRICOR	68
<i>terconazole</i>	112	TOLSURA	14	TRIDERM	78
<i>teriflunomide</i>	53	<i>tolterodine</i>	123	<i>trientine</i>	82
<i>teriparatide</i>	106	<i>tolvaptan</i>	91	TRI-ESTARYLLA	112
TESTIM	90	TOPAMAX	53	<i>trifluoperazine</i>	54
<i>testosterone</i>	90	TOPICORT	78	<i>trifluridine</i>	116
<i>testosterone cypionate</i>	90	<i>topiramate</i>	53	<i>trihexyphenidyl</i>	54
<i>testosterone enanthate</i>	90	TOPROL XL	68	TRIJARDY XR	91
<i>tetanus,diphtheria tox ped(pf)</i> ... 101		<i>toremifene</i>	23	TRIKAFTA	121
<i>tetrabenazine</i>	53	<i>torseamide</i>	68	TRI-LEGEST FE	112
<i>tetracycline</i>	14	TOSYMRA	53	TRILEPTAL	54
TEXACORT	78	TOUJEO MAX U-300		TRILIPIX	68
THALITONE	68	SOLOSTAR	91	TRI-LO-ESTARYLLA	112
THALOMID	23	TOUJEO SOLOSTAR U-300		TRI-LO-SPRINTEC	112
THEO-24	121	INSULIN	91	<i>trimethobenzamide</i>	97
<i>theophylline</i>	121	TOVET EMOLLIENT	78	<i>trimethoprim</i>	14
THIOLA	82	TOVIAZ	123	TRI-MILI	112
THIOLA EC	82	TPN ELECTROLYTES	125	<i>trimipramine</i>	54
<i>thioridazine</i>	53	TRACLEER	121	TRINTELLIX	54
<i>thiothixene</i>	53	TRADJENTA	91	TRI-NYMYO	112
THYQUIDITY	90	<i>tramadol</i>	53, 54	TRI-SPRINTEC (28)	112
TIADYLT ER	68	<i>tramadol-acetaminophen</i>	54	TRIUMEQ	14
<i>tiagabine</i>	53	<i>trandolapril</i>	68	TRIUMEQ PD	14
TIAZAC	68	<i>trandolapril-verapamil</i>	68	TRIVORA (28)	112
TIBSOVO	23	<i>tranexamic acid</i>	112	TRI-VYLIBRA	112
TICOVAC	101	TRANSDERM-SCOP	97	TRI-VYLIBRA LO	112
<i>tigecycline</i>	14	<i>tranylcypromine</i>	54	TRIZIVIR	14
TIKOSYN	68	TRAVASOL 10 %	125	TROKENDI XR	54
TILIA FE	112	TRAVATAN Z	116	TROPHAMINE 10 %	125
<i>timolol maleate</i>	68, 116	<i>travoprost</i>	116	<i>tropium</i>	123
<i>timolol maleate (pf)</i>	116	<i>trazodone</i>	54	TRUDHESA	54
TIMOPTIC OCUDOSE (PF) 116		TRECATOR	14	TRULANCE	97
<i>tinidazole</i>	14	TRELEGY ELLIPTA	121	TRULICITY	91
<i>tiopronin</i>	82	TRELSTAR	23	TRUMENBA	101
<i>tiotropium bromide</i>	121	TREMFYA	78	TRUQAP	23
TIROSINT	90			TRUVADA	14

TUDORZA PRESSAIR	122	<i>varenicline</i>	82	VIREAD	15
TUKYSA	23	VARIVAX (PF)	101	VISTARIL	122
TURALIO	23	VARUBI	97	VITRAKVI	24
TURQOZ (28)	112	VASCEPA	69	VIVELLE-DOT	112
TWINRIX (PF)	101	VASERETIC	69	VIVITROL	55
TWYNEO	78	VASOTEC	69	VIVJOA	15
TYBLUME	112	VECAMYL	69	VIZIMPRO	24
TYBOST	14	VECTICAL	78	VOGELXO	91
TYDEMY	112	VELIVET TRIPHASIC		VONJO	24
TYGACIL	14	REGIMEN (28)	112	VOQUEZNA	97
TYKERB	23	VELPHORO	82	VOQUEZNA DUAL PAK	97
TYMLOS	106	VELSIPITY	97	VOQUEZNA TRIPLE PAK	97
TYPHIM VI	101	VELTASSA	82	<i>voriconazole</i>	15
TYRVAYA	116	VELTIN	78	VOSEVI	15
TYVASO DPI	122	VEMLIDY	15	VOTRIENT	24
UBRELVY	54	VENCLEXTA	23	VOWST	97
UCERIS	97	VENCLEXTA STARTING		VOXZOGO	91
UDENYCA	101	PACK	23	VRAYLAR	55
UDENYCA		<i>venlafaxine</i>	55	VTAMA	78
AUTOINJECTOR	101	<i>venlafaxine besylate</i>	55	VUITY	116
ULORIC	106	VENTOLIN HFA	122	VUMERITY	55
ULTRAVATE	78	VEOZAH	112	VYFEMLA (28)	112
UNASYN	14	<i>verapamil</i>	69	VYLIBRA	112
UNITHROID	91	VERDESO	78	VYNDAMAX	69
UPTRAVI	68	VEREGEN	78	VYNDAQEL	69
UROCIT-K 10	123	VERELAN	69	VYTORIN 10-10	69
UROCIT-K 15	123	VERELAN PM	69	VYTORIN 10-20	69
UROCIT-K 5	123	VERKAZIA	116	VYTORIN 10-40	69
UROXATRAL	123	VERQUVO	69	VYTORIN 10-80	69
URSO 250	97	VERSACLOZ	55	VYVANSE	55
URSO FORTE	97	VERZENIO	23	VYZULTA	116
<i>ursodiol</i>	97	VESICARE	124	WAINUA	55
UZEDY	54, 55	VESICARE LS	124	WAKIX	55
VABOMERE	14	VESTURA (28)	112	<i>warfarin</i>	69
VAGIFEM	112	VEVYE	116	WEGOVY	82
<i>valacyclovir</i>	14	VFEND	15	WELCHOL	69
VALCHLOR	78	VFEND IV	15	WELIREG	24
VALCYTE	14	VIBERZI	97	WELLBUTRIN SR	55
<i>valganciclovir</i>	15	VIBRAMYCIN	15	WELLBUTRIN XL	56
VALIUM	55	VICTOZA 3-PAK	91	WINLEVI	78
<i>valproic acid</i>	55	VIENVA	112	WINREVAIR	122
<i>valproic acid (as sodium salt)</i>	55	<i>vigabatrin</i>	55	WIXELA INHUB	122
<i>valsartan</i>	68	VIGADRONE	55	WYMZYA FE	112
<i>valsartan-hydrochlorothiazide</i>	68	VIGAMOX	116	XACIATO	112
VALTOCO	55	VIGPODER	55	XALATAN	116
VALTREX	15	VIIBRYD	55	XALKORI	24
VANCOGIN	15	VIJOICE	24	XANAX	56
<i>vancomycin</i>	15	<i>vilazodone</i>	55	XANAX XR	56
VANDAZOLE	112	VIMOVO	55	XARELTO	69
VANFLYTA	23	VIMPAT	55	XARELTO DVT-PE TREAT	
VANOS	78	VIOKACE	97	30D START	69
VAQTA (PF)	101	VIRACEPT	15	XATMEP	24

XCOPRI	56	ZEGALOGUE		ZONALON	78
XCOPRI MAINTENANCE		AUTOINJECTOR	91	ZONEGRAN	57
PACK	56	ZEGALOGUE SYRINGE	91	ZONISADE	57
XCOPRI TITRATION PACK	56	ZEGERID	97	<i>zonisamide</i>	57
XDEMVY	116	ZEJULA	24	ZONTIVITY	69
XELJANZ	106	ZELAPAR	57	ZORTRESS	24
XELJANZ XR	106	ZELBORAF	24	ZORYVE	79
XELPROS	116	ZEMAIRA	82	ZOSYN IN DEXTROSE	
XELSTRYM	56	ZEMBRACE SYMTOUCH	57	(ISO-OSM)	16
XENAZINE	56	ZEMDRI	15	ZOVIA 1-35 (28)	113
XERESE	78	ZEMPLAR	91	ZOVIRAX	79
XERMELO	24	ZENATANE	78	ZTALMY	57
XGEVA	24	ZENPEP	98	ZTLIDO	79
XHANCE	122	ZENZEDI	57	ZUBSOLV	57, 58
XIFAXAN	15	ZEPATIER	15	ZURZUVAE	58
XIGDUO XR	91	ZEPOSIA	57	ZYCLARA	79
XIIDRA	116	ZEPOSIA STARTER KIT		ZYDELIG	24
XOFLUZA	15	(28-DAY)	57	ZYFLO	122
XOLAIR	122	ZEPOSIA STARTER PACK		ZYKADIA	24
XOLREMDI	101	(7-DAY)	57	ZYLET	116
XOPENEX HFA	122	ZERBAXA	15	ZYMFENTRA	98
XOSPATA	24	ZERVIAE	116	ZYPITAMAG	69
XPOVIO	24	ZESTORETIC	69	ZYPREXA	58
XTAMPZA ER	56	ZESTRIL	69	ZYPREXA RELPREVV	58
XTANDI	24	ZETIA	69	ZYPREXA ZYDIS	58
XULANE	112	ZETONNA	122	ZYTIGA	24, 25
XULTOPHY 100/3.6	91	ZIAC	69	ZYVOX	16
XURIDEN	82	ZIAGEN	15		
XYOSTED	91	ZIANA	78		
XYREM	56	<i>zidovudine</i>	15		
XYWAV	56	ZIEXTENZO	102		
YARGESA	91	ZILBRYSQ	57		
YASMIN (28)	112	<i>zileuton</i>	122		
YAZ (28)	112	ZILXI	78		
YF-VAX (PF)	102	ZIMHI	57		
YONSA	24	ZIOPTAN (PF)	116		
YUFLYMA(CF)	106	<i>ziprasidone hcl</i>	57		
YUFLYMA(CF) AI		<i>ziprasidone mesylate</i>	57		
CROHN'S-UC-HS	106	ZIPSOR	57		
YUFLYMA(CF)		ZIRGAN	116		
AUTOINJECTOR	106	ZITHROMAX	15		
YUPELRI	122	ZITHROMAX TRI-PAK	15		
YUSIMRY(CF) PEN	106	ZITHROMAX Z-PAK	16		
YUVAFEM	112	ZITUVIO	91		
ZAFEMY	112	ZOCOR	69		
<i>zafirlukast</i>	122	ZOKINVY	82		
<i>zaleplon</i>	56	ZOLINZA	24		
ZANAFLEX	56	<i>zolmitriptan</i>	57		
ZARONTIN	56	ZOLOFT	57		
ZARXIO	102	<i>zolpidem</i>	57		
ZAVESCA	91	ZOMACTON	102		
ZAVZPRET	56	ZOMIG	57		

Acitretin

Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Actemra

Products Affected

- **ACTEMRA ACTPEN**
- **ACTEMRA SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide). For giant cell arteritis, patients must have therapeutic failure or intolerance to one systemic corticosteroid (e.g., prednisone). For polyarticular juvenile idiopathic arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For systemic sclerosis-associated interstitial lung disease (SSc-ILD), patients must have therapeutic failure or intolerance to one immunosuppressant (e.g., mycophenolate mofetil, corticosteroids, cyclophosphamide). Documentation of systemic juvenile idiopathic arthritis.
Age Restrictions	Deny if less than 18 years of age for systemic sclerosis-associated interstitial lung disease (SSc-ILD), Rheumatoid Arthritis, and Giant Cell Arteritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Acthar H.P.

Products Affected

- ACTHAR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Covered for the following indications: 1. Infantile spasms (West syndrome) in children less than 2 years of age. 2. Acute exacerbations of multiple sclerosis (MS) for patients receiving concurrent immunomodulator therapy (e.g., interferon beta, glatiramer acetate, dimethyl fumarate, fingolimod, teriflunomide) 3. Rheumatic disorders for patients receiving maintenance therapy with at least one NSAID, DMARD (e.g. leflunomide) or biologic (e.g. adalimumab) 4. Collagen diseases for members receiving maintenance therapy with at least one antimalarial (e.g. hydroxychloroquine) or immunosuppressant (e.g. azathioprine) 5. Dermatologic diseases 6. Allergic states (i.e. serum sickness and transfusion reaction due to serum protein reaction 7. Ophthalmic diseases 8. Respiratory diseases 9. Gout and unable to take first-line therapies. 10. Pediatric acquired epileptic aphasia. 11. Proteinuria in nephrotic syndrome and trial/failure or contraindication to two therapies from any of the following different classes: corticosteroids (e.g., cortisone or dexamethasone), calcineurin inhibitors (e.g., cyclosporine or tacrolimus, per DRUGDEX). For covered indications 2 through 10, limited/unsatisfactory response or intolerance (i.e. severe anaphylaxis) to two corticosteroids (i.e. IV methylprednisolone, IV dexamethasone, or high dose oral steroids) must be documented.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month

PA Criteria	Criteria Details
Other Criteria	For reauthorization, the following (1. 2. and 3.) must be met. 1) Prescriber attestation that the member cannot use corticosteroids (e.g. IV methylprednisolone, high dose oral corticosteroids) due to unsatisfactory response, intolerance (e.g. severe anaphylaxis) or experienced a severe adverse event to corticosteroids (e.g. psychosis). 2) If the reauthorization is for the treatment of multiple sclerosis, a rheumatic disorder, dermatologic disease, or nephrotic syndrome, the prescriber attests that H.P. Acthar is being used for a new acute exacerbation and not on a routine basis to prevent an exacerbation as supported by Compendia. 3) If the reauthorization is for treatment of multiple sclerosis, a rheumatic disorder, or collagen disease, the member continues to receive maintenance therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Gout, Epileptic Aphasia
Part B Prerequisite	No

Actimmune

Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adalimumab

Products Affected

- **ABRILADA(CF)**
- **ABRILADA(CF) PEN**
- *adalimumab-aacf subcutaneous pen injector kit*
- *adalimumab-aaty*
- *adalimumab-fkjp subcutaneous pen injector kit*
- *adalimumab-fkjp subcutaneous syringe kit*
- *adalimumab-ryvk subcutaneous auto-injector, kit*
- **AMJEVITA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 40 MG/0.4 ML, 40 MG/0.8 ML, 80 MG/0.8 ML**
- **AMJEVITA(CF) SUBCUTANEOUS SYRINGE 10 MG/0.2 ML, 20 MG/0.2 ML, 20 MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8 ML**
- **HADLIMA**
- **HADLIMA PUSHTOUCH**
- **HADLIMA(CF)**
- **HADLIMA(CF) PUSHTOUCH**
- **HULIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT**
- **HULIO(CF) SUBCUTANEOUS SYRINGE KIT**
- **IDACIO(CF)**
- **IDACIO(CF) PEN CROHN-UC STARTR**
- **IDACIO(CF) PEN PSORIASIS START**
- **IDACIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT**
- **SIMLANDI(CF) AUTOINJECTOR**
- **YUFLYMA(CF)**
- **YUFLYMA(CF) AI CROHN'S-UC-HS**
- **YUFLYMA(CF) AUTOINJECTOR**
- **YUSIMRY(CF) PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 1 immunosuppressant or corticosteroid, or all are contraindicated.

PA Criteria	Criteria Details
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Therapeutic failure or intolerance to 2 of the following preferred products is required: Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm. For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. Please Note for New Starts Only: preferred adalimumab products include Humira with NDC starting 00074, Cyltezo with NDC starting 00597, Hyrimoz with NDC starting 61314, adalimumab-adaz with NDC starting 61314, and adalimumab-adbm with NDC starting 00597.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adbry

Products Affected

- **ADBRY SUBCUTANEOUS SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: 1) Dupixent and 2) If 18 years or older, Rinvoq. For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADHD Drugs

Products Affected

- *clonidine hcl oral tablet extended release 12 hr*
- *guanfacine oral tablet extended release 24 hr*
- **INTUNIV ER**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adlarity

Products Affected

- ADLARITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1 or 2): 1) therapeutic failure or intolerance to donepezil tablets, 2) Unable to take daily oral donepezil due to impaired memory.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Afinitor

Products Affected

- **AFINITOR** *mg, 2.5 mg, 5 mg, 7.5 mg*
- **AFINITOR DISPERZ ORAL TABLET FOR SUSPENSION 2 MG, 3 MG, 5 MG** • *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*
- *everolimus (antineoplastic) oral tablet 10*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) HR mutation status and HER2 mutation status 2) Alternatives tried/failed 3) Concomitant therapy 4) Candidacy for surgical resection
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Afinitor, documentation of failure on generic everolimus tablets is required. For brand Afinitor Disperz, documentation of failure on generic everolimus tablets for suspension is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Agamree

Products Affected

- AGAMREE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy (DMD) with mutation of dystrophin gene -AND- One of the following: 1) Therapeutic failure, intolerance, or contraindication to prednisone, or 2) Growth delay or decline while on prednisone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Aimovig

Products Affected

- **AIMOVIG AUTOINJECTOR
SUBCUTANEOUS AUTO-INJECTOR
140 MG/ML, 70 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ajovy

Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Akeega

Products Affected

- **AKEEGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) BRCA mutations, 2) Comcomitant therapy with prednisone, 3) Concomitant therapy with a gonadotropin-releasing hormone analog or a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alecensa

Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive -AND- One of the following (1 or 2): 1) metastatic disease, or 2) will be used as adjuvant treatment following tumor resection of node positive or greater than or equal to 4 cm tumor(s).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alkindi

Products Affected

- **ALKINDI SPRINKLE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of adrenocortical insufficiency -AND- Therapeutic failure or intolerance to oral generic hydrocortisone tablets.
Age Restrictions	Deny if greater than 17 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alpha1-Proteinase Inhibitors

Products Affected

- **ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG**
- **GLASSIA**
- **PROLASTIN-C INTRAVENOUS SOLUTION**
- **ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of panacinar emphysema AND documentation of a decline in forced expiratory volume in 1 second (fev1) despite medical therapy (bronchodilators, corticosteroids) AND documentation of phenotype (pi*zz, pi*znull or pi*nullnull) associated with causing serum alpha 1-antitrypsin of less than 80 mg/dl AND documentation of an alpha 1-antitrypsin serum level below the value of 35% of normal (less than 80 mg/dl).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when furnished incident to a physician service and is not self-administered.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alunbrig

Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Amlodipine Liquid

Products Affected

- KATERZIA
- NORLIQVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Inability to swallow amlodipine tablets
Age Restrictions	Deny if less than 6 years of age for Hypertension
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ampyra

Products Affected

- AMPYRA
- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure disorder, Cr Cl less than 50ml/min
Required Medical Information	Documentation of diagnosis -AND- documentation that the patient is ambulatory and has walking impairment as evidenced by one of the following. 1. Functional status score (EDSS score). 2. Timed 25-foot Walk Test (T25W).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For brand Ampyra, documentation of failure on generic dalfampridine. For reauthorization, documentation supporting improvement in walking impairment from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Anabolic Steroids

Products Affected

- **METHITEST**
- *methyltestosterone oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Apokyn

Products Affected

- **APOKYN**
- *apomorphine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease -AND- for use in acute, intermittent treatment of hypomobility off episodes -AND- experiencing off episodes despite the use of oral carbidopa/levodopa -AND- Therapeutic failure, intolerance, or contraindication to one of the following generic products: pramipexole, ropinirole, entacapone, selegiline or rasagiline
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Arikayce

Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Mycobacterium avium complex lung disease -AND- Attestation of not achieving negative sputum cultures despite at least 6 months with a multidrug background regimen containing 2 of the following: 1) macrolide 2) rifamycin or 3) ethambutal -AND- Arikayce will be used in conjunction with a background multidrug regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive sputum cultures -OR- negative sputum cultures for insufficient period of time (e.g. less than 12 months).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Aspruzyo Sprinkle

Products Affected

- ASPRUZYO SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic angina -AND- Therapeutic failure, contraindication, or intolerance to one beta-blocker (e.g., propranolol oral solution, metoprolol tartrate, propranolol immediate-release) -AND- Inability to swallow generic ranolazine tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ATTR-CM drugs

Products Affected

- VYNDAMAX
- VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	Concomitantly with transthyretin-lowering agents
Required Medical Information	Documentation of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) with amyloid deposits on cardiac biopsy or scintigraphy with heart contralateral greater than 1.5 (Grade III) -AND- Cardiac involvement supported by cardiac magnetic resonance, echocardiography or serum cardiac biomarker (e.g. B-type natriuretic peptide, cardiac troponin) - AND- Primary (light chain) amyloidosis has been ruled out by immunohistochemistry, mass spectrometry or scintigraphy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement or delayed disease progression from baseline demonstrated by 6-minute walk test, cardiac function (e.g. LVEF, NYHA class), Kansas City Cardiomyopathy Questionnaire-Overall Summary, number of cardiovascular-related hospitalizations or serum cardiac biomarkers (e.g. B-type natriuretic peptide, cardiac troponin)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Atypical Antipsychotics

Products Affected

- **ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET WITH SENSOR AND STRIP 15 MG, 2 MG, 20 MG, 30 MG, 5 MG**
 - **ABILIFY MYCITE STARTER KIT**
- **ORAL TABLET WITH SENSOR, STRIP, POD 10 MG**
 - **ABILIFY ORAL TABLET**
 - *aripiprazole*
 - **REXULTI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI). For Rexulti used for schizophrenia, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Aubagio

Products Affected

- **AUBAGIO**
- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as fingolimod, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	For brand Aubagio, therapeutic failure or intolerance to generic teriflunomide is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Augtyro

Products Affected

- AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive. Documentation of solid tumors - AND- disease harbors a NTRK gene fusion -AND- one of the following (1-2): 1) disease is locally advanced or metastatic, or 2) surgical resection is likely to result in severe morbidity -AND- one of the following (3-4): 3) disease has progressed following treatment, or 4) the member has no satisfactory alternative therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Auryxia

Products Affected

- AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of iron deficiency anemia
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Austedo

Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 6 MG (14)-12 MG (14)-24 MG (14)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3) 1) Chorea associated with Huntington's disease 2) In patients with comorbid depression, attestation of adequate treatment for depression is required. 3) Attestation that patient is not actively suicidal. -OR- 4) Tardive Dyskinesia
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Auvelity

Products Affected

- AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ayvakit

Products Affected

- **AYVAKIT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) PDGFRA exon 18 mutation status 2) platelet count greater than or equal to 50 x 10 ⁹ /L
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Baclofen Solution

Products Affected

- *baclofen oral solution 10 mg/5 ml (2 mg/ml)*
- **OZOBAX DS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member has an inability to swallow generic baclofen tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bafiertam

Products Affected

- **BAFIERTAM**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Balversa

Products Affected

- **BALVERSA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) FGFR3 mutation status as detected by an FDA approved test 2) disease progression on or after at least on prior line of systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Banzel

Products Affected

- **BANZEL**
- *rufinamide*

PA Criteria	Criteria Details
Exclusion Criteria	Patients with familial short QT syndrome
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Benlysta

Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of active systemic lupus erythematosus (SLE) -AND- documentation of positive anti-nuclear antibody (ANA) titer (greater than or equal to 1:80) or anti-double-stranded DNA antibody (anti-dsDNA) greater than or equal to 30IU/mL -AND- member will continue to receive concomitant standard of care treatment with use of at least one of the following (alone or in combination): 1.) corticosteroids (e.g. prednisone) 2.) antimalarials (e.g. hydroxychloroquine) 3.) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate) -OR- Documentation of active lupus nephritis -AND- documentation of positive ANA titer (greater than or equal to 1:80) or anti-dsDNA greater than or equal to 30 IU/mL -AND- member will continue to receive concomitant standard of care treatment which includes corticosteroids (e.g. prednisone) with at least one of the following: 1.) mycophenolate for induction followed by mycophenolate for maintenance 2.) cyclophosphamide for induction followed by azathioprine for maintenance</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	<p>For SLE reauthorization, attestation of disease stability or improvement -AND- attestation the member will continue to receive standard of care therapy with corticosteroids, antimalarials, or immunosuppressives. For active LN reauthorization, attestation of disease stability or improvement -AND- attestation the member will continue to receive standard of care therapy with mycophenolate or azathioprine. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.</p>

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Benznidazole

Products Affected

- *benznidazole*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 2 years of age.
Prescriber Restrictions	
Coverage Duration	2 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Berinert

Products Affected

- **BERINERT INTRAVENOUS KIT**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	<p>For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-4): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. 4) Documentation of member's weight. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (5-9): 5) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 6) Documentation of family history of HAE or FXII mutation 7) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 8) Medications known to cause angioedema have been evaluated and discontinued. 9) Documentation of member's weight.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For 18 years of age or older, therapeutic failure, intolerance or contraindication to icatibant.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Besremi

Products Affected

- **BESREMI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bimzelx

Products Affected

- **BIMZELX**
- **BIMZELX AUTOINJECTOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Taltz, Otezla, Stelara SC, Enbrel and Skyrizi SC. Must follow recommended dosing guidelines based upon weight. Induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bonjesta

Products Affected

- **BONJESTA**
- **DICLEGIS**
- *doxylamine-pyridoxine (vit b6)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	9 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bosulif

Products Affected

- **BOSULIF ORAL CAPSULE 100 MG, 50 MG**
- **BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy. For pediatric patients 1 year of age and older, one of the following (3, 4): 3) newly diagnosed PH-positive CML in the chronic phase 4) diagnosis of Ph-positive CML in the chronic phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Bosutinib capsules and 18 years of age or older, inability to swallow tablets is required. For Bosutinib 100mg capsules and pediatric 1 year of age or older, inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Braftovi

Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	Use in wild-type BRAF melanoma or wild-type BRAF CRC
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) BRAF V600E or V600K mutation status 2) alternatives tried/failed 3) concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bronchitol

Products Affected

- BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis -AND- Passed a Bronchitol Tolerance Test -AND- Used in conjunction with standard therapies for the management of cystic fibrosis to improve pulmonary function (e.g. bronchodilators, antibiotics, anti-inflammatory therapy).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increase in FEV1
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Brukinsa

Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma (MCL), previous treatment with at least 1 prior therapy. For marginal zone lymphoma (MZL), previous treatment with at least 1 anti-CD20-based regimen. For follicular lymphoma (FL), using in combination with obinutuzumab and previous treatment with at least 2 prior lines of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Buphenyl

Products Affected

- **BUPHENYL**
- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Buphenyl, trial and failure of generic sodium phenylbutyrate is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bylvay

Products Affected

- **BYLVAY ORAL CAPSULE 1,200 MCG, 400 MCG**
- **BYLVAY ORAL PELLETT 200 MCG, 600 MCG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pruritis with progressive familial intrahepatic cholestasis or cholestatic pruritus with Alagille syndrome -AND- The member does not have cirrhosis, portal hypertension, or history of hepatic decompensation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial, 12 months reauthorization
Other Criteria	For reauthorization, attestation of improvement in pruritis -AND- attestation that the member has not progressed to any of the following (1-3): 1) portal hypertension, 2) cirrhosis, or 3) experienced a hepatic decompensation event.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cablivi

Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of acquired thrombocytopenic purpura (aTTP) -AND- Used in conjunction with plasma exchange and immunosuppressive therapy (i.e. systemic corticosteroids or rituximab)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	75 days initial authorization, 28 days reauthorization
Other Criteria	For reauthorization, attestation of remaining signs and symptoms of persistent disease (e.g. suppressed ADAMTS 13 activity level remain present)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cabometyx

Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) alternatives tried/failed 2) attestation of first line use 3) concomitant therapy 4) radioactive iodine refractory status
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Calquence

Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if applicable to diagnosis, alternatives tried/failed
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Camzyos

Products Affected

- CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy -AND- Left ventricular ejection fraction of greater than or equal to 55% -AND- Valsalva left ventricular outflow tract peak gradient of greater than or equal to 50 mmHg at rest or after provocation -AND- Therapeutic failure or intolerance to one of the following (1 or 2) or contraindication to all: 1) Non-vasodilating beta blocker (e.g. metoprolol) 2) non-dihydropyridine calcium channel blocker (e.g. diltiazem) -AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, prescriber attestation of no NYHA class worsening - AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Caplyta

Products Affected

- CAPLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Caprelsa

Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic or progressive medullary thyroid cancer in patients with unresectable locally advanced or metastatic disease
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Carac

Products Affected

- **CARAC**
- *fluorouracil topical cream 0.5 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Trial and failure of generic topical fluorouracil solution or fluorouracil 5% cream.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If requesting brand Carac, trial and failure of fluorouracil 0.5% cream.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Carbaglu

Products Affected

- **CARBAGLU**
- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use as an adjunct therapy for acute hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA) -OR- maintenance therapy for chronic hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Carbinoxamine

Products Affected

- *carbinoxamine maleate oral liquid*
- *carbinoxamine maleate oral tablet 4 mg*
- **RYVENT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Failure, contraindication or intolerance to 2 antihistamines indicated for diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cayston

Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cerdelga

Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of type 1 Gaucher disease confirmed by the following A. or B. A) With one of the following symptoms (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm ³ . 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B) Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles -AND- Documentation of CYP2D6 metabolizer status (e.g. intermediate metabolizer).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CF drugs

Products Affected

- **BETHKIS**
 - **KITABIS PAK**
 - **TOBI**
 - **TOBI PODHALER**
- *tobramycin in 0.225 % nacl*
 - *tobramycin inhalation*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis. For Bethkis: failure on, intolerance to, or contraindication to generic tobramycin inhalation solution
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization of tobramycin products, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Chenodal

Products Affected

- CHENODAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of radiolucent gallstones AND an inadequate response or intolerance to ursodiol therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months for initial approval with an additional 12 months upon renewal
Other Criteria	Safety of use beyond 24 months is not established. For reauthorization, attestation of partial dissolution of gallstones
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cholbam

Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of bile acid synthesis disorders due to single enzyme defects (e.g. AKR1D1, CTX, CYP7A1) -OR- documentation of use as adjunctive therapy for peroxisomal disorders (PDs), including Zellweger spectrum disorders, in patients who exhibit manifestations of liver disease, steatorrhea, or complications from decreased fat soluble vitamin absorption.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cialis

Products Affected

- **CIALIS ORAL TABLET 2.5 MG, 5 MG**
- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of benign prostatic hyperplasia (BPH) and trial/failure of at least two alternative medications in the following classes (alpha-1 adrenergic blockers and/or 5-alpha reductase inhibitors)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Cialis, trial and failure of generic tadalafil is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cibinqo

Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: Dupixent and Rinvoq.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cimzia

Products Affected

- CIMZIA
- CIMZIA POWDER FOR RECONST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For non-radiographic axial spondyloarthritis, inadequate response or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs). For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	<p>For Crohn's disease, patients must have therapeutic failure or intolerance to 2 of the following preferred biologic products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Stelara, Rinvoq, and Skyrizi SC. For Rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Actemra SC, Xeljanz/Xeljanz XR and Rinvoq. For plaque psoriasis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Taltz, Otezla, Stelara SC, Skyrizi SC, and Enbrel. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Enbrel, Taltz, Rinvoq, and Xeljanz/Xeljanz XR. For Psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cosentyx, Taltz, Enbrel, a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For initial and induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended initial and induction therapy dosing regimens per indication.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cinryze

Products Affected

- CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Citalopram Capsule

Products Affected

- *citalopram oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- citalpram 20 mg has been received for greater than or equal to 7 days -AND- Therapeutic failure or intolerance to generic citalopram tablets -AND- Therapeutic failure, intolerance or contraindication to at least one other antidepressant (e.g. SNRI, SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Clemastine Syrup

Products Affected

- *clemastine oral syrup*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of allergic rhinitis or mild, uncomplicated allergic skin manifestations of urticaria and angioedema -AND- Therapeutic failure, contraindication or intolerance to one generic antihistamine product - AND- Therapeutic failure or intolerance to generic clemastine fumarate tablets or an inability to swallow tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cometriq

Products Affected

- **COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressive, metastatic medullary thyroid cancer
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Copiktra

Products Affected

- **COPIKTRA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if applicable to diagnosis, alternatives tried/failed
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Corlanor

Products Affected

- **CORLANOR ORAL SOLUTION**
- **CORLANOR ORAL TABLET 5 MG, 7.5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis AND all of the following: 1) Normal sinus rhythm. 2) Resting heart rate greater than or equal to 70 beats per minute. 3) Left ventricular ejection fraction less than or equal to 35 percent, when applicable. 4) In adult patients (greater than or equal to 18 years), trial/failure of maximum tolerated dose of one beta-blocker used for treatment of heart failure (e.g., bisoprolol, carvedilol, metoprolol succinate) OR contraindication to beta-blocker use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For oral solution, attestation of inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cosentyx

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs. For enthesitis-related arthritis, inadequate response or intolerance to at least 1 non-biologic disease modifying anti-rheumatic drugs (DMARD), or all are contraindicated.
Age Restrictions	Deny if less than 6 years of age for moderate to severe plaque psoriasis -OR- less than 2 years of age for psoriatic arthritis -OR- less than 4 years of age for enthesitis-related arthritis -OR- less than 18 years of age for all other indications
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication. For hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cotellic

Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) BRAF V600E or V600K mutation status 2) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cresemba

Products Affected

- CRESEMBA ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of invasive aspergillosis infection -AND- therapeutic failure, contraindication, or intolerance to voriconazole -OR- Documentation of invasive mucormycosis infection.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of presence of continued indicators of active disease (e.g. histopathology, fungal culture).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Crinone

Products Affected

- CRINONE

PA Criteria	Criteria Details
Exclusion Criteria	Use to promote fertility
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cuvrior

Products Affected

- CUVRIOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of stable Wilson's disease -AND- Previously tolerated one penicillamine product used for de-coppering (e.g., generic penicillamine tablet or capsule, Depen) -AND- Therapeutic failure or intolerance to generic trientine hydrochloride
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cysteamine Ophthalmic Drops

Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystinosis -AND- Attestation of accumulation of corneal cystine crystals
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daraprim

Products Affected

- **DARAPRIM**
- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For primary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 100 cells/mm ³ -AND- Toxoplasma IgG positive -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For secondary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 200 cells/mm ³ . For secondary prophylaxis of cystoisosporiasis with CD4 count less than 200 cells/mm ³ or acute cystoisosporiasis infection: failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For primary prophylaxis of Pneumocystis jirovecii pneumonia: diagnosis of HIV - AND- CD4 count less than 200 cells/mm ³ -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Daraprim, trial and failure of generic pyrimethamine is required.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Daurismo

Products Affected

- **DAURISMO ORAL TABLET 100 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of newly diagnosed Acute Myeloid Leukemia -AND- Used in combination with cytarabine -AND- At least one comorbidity that preclude use of intensive induction chemotherapy defined as one of the following: 1) Age greater than or equal to 75 2) Severe cardiac or pulmonary comorbidity 3) Reduced renal function 4) Hepatic impairment 5.) Physician attests patient is not a candidate for intensive induction therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daybue

Products Affected

- DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Rett syndrome confirmed by all of the following (1 thru 4): 1) Partial or complete loss of acquired purposeful hand skills, 2) Partial or complete loss of acquired spoken language, 3) Gait abnormalities: impaired or absence of ability, 4) Stereotypic hand movements such as hand wringing/squeezing, clapping/tapping, mouthing, and washing/rubbing automatisms
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daytrana

Products Affected

- **DAYTRANA**
- *methylphenidate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial/failure or intolerance to 2 of the following generic medications: methylphenidate, atomoxetine, or dextroamphetamine/amphetamine.
Age Restrictions	Deny if less than 6 years of age or greater than 17 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Daytrana, trial/failure of generic methylphenidate patch is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Deferasirox

Products Affected

- *deferasirox*
- **EXJADE**
- **JADENU**
- **JADENU SPRINKLE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic iron overload due to blood transfusions, transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -And- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw). For Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia (NTDT) Syndrome, LIC of at least 5 mg Fe/g dw -AND- serum ferritin greater than 300 mcg/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Jadenu and brand Exjade, initial authorization requires trial and failure of deferasirox (generic Exjade) and reauthorization requires failure of generic deferasirox (generic Exjade) if not previously trialed. For reauthorization of chronic iron overload due to blood transfusion, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw. For reauthorization of chronic iron overload in NTDT syndrome, LIC greater than or equal to 3 mg Fe/g dw.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Diacomit

Products Affected

- **DIACOMIT ORAL CAPSULE 250 MG, 500 MG**
- **DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravets syndrome - AND- Used in combination with clobazam
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dihydroergotamine

Products Affected

- *dihydroergotamine nasal*
- **MIGRANAL**
- **TRUDHESA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- requires non-oral route of administration -AND- therapeutic failure or intolerance to generic sumatriptan nasal spray.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dojolvi

Products Affected

- DOJOLVI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitantly with another medium chain triglyceride product
Required Medical Information	Documentation of molecularly confirmed long-chain fatty acid oxidation disorders
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Doptelet

Products Affected

- **DOPTELET (10 TAB PACK)**
- **DOPTELET (15 TAB PACK)**
- **DOPTELET (30 TAB PACK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -OR- Documentation of chronic immune thrombocytopenia -AND- Trial, intolerance, or inadequate response to corticosteroid therapy, immunoglobulin therapy or splenectomy -AND- One of the following (1 or 2): 1) Platelet count less than or equal to $50 \times 10^9/L$ and has significant mucous member bleeding or at least one risk factor for bleeding (e.g. hypertension, peptic ulcer disease). 2) Platelets count of less than or equal to $30 \times 10^9/L$
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For thrombocytopenia with chronic liver disease- 1 mo. For chronic immune thrombocytopenia- 12 mo.
Other Criteria	Platelet count is provided for applicable dosing.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Doxepin Cream

Products Affected

- *doxepin topical*
- **PRUDOXIN**
- **ZONALON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- course of therapy will not exceed 8 days -AND- experienced therapeutic failure, intolerance, or contraindication to one of the following (1 or 2): 1) at least 2 generic formulary topical corticosteroids. 2) generic topical tacrolimus or pimecrolimus for topic dermatitis with the facial or anogenital involvement.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Duexis2

Products Affected

- *ibuprofen-famotidine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) Trial/failure of ibuprofen used in combination with famotidine. -AND- 2) Trial/failure of one additional generic formulary NSAID (other than ibuprofen) used in combination with another generic formulary H2-receptor blocker (other than famotidine).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Duobrii

Products Affected

- DUOBRII

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of plaque psoriasis -AND- therapeutic failure or intolerance to generic tazarotene cream -AND- therapeutic failure or intolerance to 1 high-potency topical corticosteroid (e.g. betamethasone dipropionate 0.05%, halobetasol propionate 0.05%)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dupixent

Products Affected

- **DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML**
- **DUPIXENT SYRINGE**
- **SUBCUTANEOUS SYRINGE 100 MG/0.67 ML, 200 MG/1.14 ML, 300 MG/2 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-2): 1) moderate to severe atopic dermatitis 2) trial & failure, or intolerance to at least one topical corticosteroid -OR- If 2 years of age or older, topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin -OR- Documentation of the following (3-6): 3) moderate-to-severe asthma 4) documented pretreatment FEV1 less than 80 percent predicted in adults or FEV1 less than 90 percent predicted in adolescents or FEV1 reversibility of at least 12% after albuterol administration 5) Blood eosinophils greater than or equal to 150 cells/uL -OR- patient is currently taking daily or alternate-day oral corticosteroids 6) using a medium- or high-dose inhaled corticosteroid and a long acting beta agonist -OR- Documentation of the following (7-8): 7) chronic rhinosinusitis with nasal polyposis (CRSwNP) 8) trial & failure, intolerance or contraindication to intra-nasal corticosteroid and 14 day course of oral corticosteroids -OR- Documentation of the following (9-11): 9) eosinophilic esophagitis 10) esophageal eosinophil count greater than or equal to 15 eos/hpf on esophageal biopsy 11) clinical symptoms of esophageal dysfunction -OR- Documentation of the following (12): 12) prurigo nodularis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization, attestation of positive clinical response to therapy, for atopic dermatitis -OR- attestation of decreased rescue medication or oral corticosteroid use, decreased frequency of severe asthma exacerbation, increased pulmonary function from baseline (e.g. FEV1) or reduction in reported asthma related symptoms, for asthma -OR- attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score, for CRSwNP -OR- attestation of histological remission (less than 15 eos/hpf) on esophageal biopsy or reduced severity or frequency of clinical symptoms of esophageal dysfunction, for EoE -OR- attestation of reduction in itch or number of nodules or lesions from baseline, for prurigo nodularis
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EGFR Tyrosine Kinase Inhibitors

Products Affected

- *erlotinib*
- **GILOTRIF**

PA Criteria	Criteria Details
Exclusion Criteria	Afatinib products: tumors with resistant EGFR mutations. Erlotinib products: use in NSCLC tumors with mutations other than those in FDA-approved indications. Use in combination with platinum based chemotherapy.
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis 1) Epidermal growth factor receptor (EGFR) mutations, including exon 19 deletions, exon 21 (L858R) substitution mutations or exon 20 insertion mutations 2) Alternatives tried/failed 3) Concomitant therapy 4) Line of therapy in which medication will be used
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Egrifta

Products Affected

- EGRIFTA SV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of HIV and lipodystrophy, member must actively be receiving antiretroviral therapy including protease inhibitors, nucleoside reverse transcriptase inhibitors, or non-nucleoside reverse transcriptase inhibitors
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Elyxyb

Products Affected

- **ELYXYB**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of migraine diagnosis -AND- trial/failure or intolerance to 2 generic NSAIDs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Emflaza

Products Affected

- *deflazacort*
- **EMFLAZA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy (DMD) with mutation of dystrophin gene -AND- onset of weakness or history of DMD starting before age 5 -AND- One of the following (1, 2, or 3). 1) Documented trial/failure, intolerance or contraindication to prednisone. 2) Documented trial of prednisone has resulted in intolerable adverse events (e.g. diabetes, hypertension that is difficult to manage, Cushingoid features, truncal obesity, greater than or equal to 10 percent increase in body weight over a 6 month period). 3) Documented severe behavioral adverse event while on prednisone that warrants prednisone dose reduction impacting efficacy for management of DMD (i.e. abnormal behavior, aggression, irritability, disturbance in mood)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Emflaza tablets and 5 years of age or older, therapeutic failure or intolerance to generic deflazacort tablets is required. For brand Emflaza suspension and 5 years of age or older, therapeutic failure or intolerance to generic deflazacort tablets or inability to swallow tablets is required, and therapeutic failure or intolerance to generic deflazacort suspension. For generic deflazacort suspension and 5 years of age or older, therapeutic failure or intolerance to generic deflazacort tablets or inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Emgality

Products Affected

- **EMGALITY PEN**
- **EMGALITY SYRINGE
SUBCUTANEOUS SYRINGE 120
MG/ML, 300 MG/3 ML (100 MG/ML X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For episodic cluster headaches, characterized by severe or very severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes when left untreated -AND- Attack frequency of at least one attack every other day during the cluster period.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency is required. For reauthorization of cluster headache, attestation of reduction in the number of mean weekly cluster headaches from baseline is required.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Enbrel

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE
25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Ankylosing Spondylitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis or Less than 4 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Endari

Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Sickle Cell Disease with 2 or more sickle cell acute complications (e.g. vaso-occlusive crisis, acute anemia, acute chest syndrome, etc.) -AND-documentation of previous trial of hydroxyurea or plans of continued therapy while taking Endari
Age Restrictions	Deny if less than 5 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of stability in sickle cell acute complications or decrease in number of sickle cell acute complications is required (e.g. vaso-occlusive crisis, acute anemia, acute chest syndrome, etc.)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Enspryng

Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of neuromyelitis optica spectrum disorder (NMOSD) - AND- Attestation of anti-aquaporin-4 (AQP4) antibody positive -AND- Not used in combination with another monoclonal antibody used for the treatment of NMOSD.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in number of NMOSD relapses. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Entadfi

Products Affected

- **ENTADFI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use for the initial treatment of benign prostatic hyperplasia (BPH) -AND- Trial/failure of finasteride used in combination with tadalafil.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	For reauthorization, attestation that the member is reinitiating treatment for BPH.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Entyvio

Products Affected

- ENTYVIO PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation of clinical response or remission following at least 2 doses of IV Entyvio at least 6 weeks before initiating therapy with Entyvio SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For moderate to severe ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Rinvoq, Xeljanz/Xeljanz XR, and Stelara SC. For moderate to severe active Crohn's disease, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Rinvoq, Skyrizi SC, and Stelara SC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epclusa

Products Affected

- **EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG**
- **EPCLUSA ORAL TABLET**
- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epidiolex

Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Lennox-Gastaut, Dravet syndromes or Tuberous Sclerosis Complex. For Lennox-Gastaut, trial and failure or intolerance of at least two standard of care treatments (e.g. lamotrigine, clobazam). For Lennox-Gastaut and Dravet syndromes, treatment is in combination with other conventional agents.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Eprontia

Products Affected

- EPRONTIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow pills or food. For migraine, therapeutic failure, intolerance, or contraindication to two generic preventive migraine therapies. For partial onset seizures, primary generalized tonic-clonic seizures, or adjunctive treatment of Lennox-Gastaut Syndrome, therapeutic failure, contraindication, or intolerance to two generic anti-epileptic drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ergotamine

Products Affected

- *ergotamine-caffeine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use to abort a vascular headache -AND- therapeutic failure or intolerance to a generic triptan -OR- documentation of use to prevent a vascular headache -AND- therapeutic failure or intolerance to generic prophylactic migraine medication
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Erivedge

Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if disease is locally advanced all of the following: 1) disease has recurred following surgery, or is not a candidate for surgery, 2) is not a candidate for radiation.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Erleada

Products Affected

- **ERLEADA ORAL TABLET 240 MG, 60 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member meets one of the following (1 or 2) 1. Documentation of use in combination with a GnRH analog -OR- 2. The member has had a bilateral orchiectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Eucrisa

Products Affected

- EUCRISA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1 and 2): 1) mild to moderate atopic dermatitis 2) trial & failure, intolerance, or contraindication to at least one topical corticosteroid -OR- documentation of facial or anogenital involvement
Age Restrictions	Deny if less than 3 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If 2 years of age or older, therapeutic failure of one of the following is required: topical tacrolimus -OR- topical pimecrolimus. Reauthorization or continuation of therapy will be approved when documentation of improvement or response to therapy is provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Evekeo

Products Affected

- *amphetamine sulfate*
- EVEKEO

PA Criteria	Criteria Details
Exclusion Criteria	Obesity
Required Medical Information	Documentation of diagnosis. For narcolepsy the following are required: Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B)Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP.
Age Restrictions	Deny if less than 6 years of age for narcolepsy or 3 years of age for ADHD
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For narcolepsy, trial and failure, intolerance to 2 of the following generic alternatives is required: immediate release amphetamine/dextroamphetamine, dextroamphetamine, and methylphenidate. For ADHD, trial/failure or intolerance to 2 unique generic stimulants (e.g. methylphenidate) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Evenity

Products Affected

- **EVENTITY SUBCUTANEOUS SYRINGE
210MG/2.34ML (105MG/1.17MLX2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Postmenopausal woman at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug. A cumulative lifetime approval of romosozumab will be limited to a coverage duration of 12 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Evrysdi

Products Affected

- EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of spinal muscular atrophy -AND- Baseline motor function test results (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT) -AND- Not using concomitantly with Spinraza -AND- Molecular genetic testing of 5q SMA showing Homozygous gene deletion, Homozygous conversion mutation or Compound heterozygote
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	For reauthorization, attestation of stable or clinically significant improvement in Spinal Muscular Atrophy associated symptoms (e.g. stabilization or decreased decline in motor function compared to the predicted natural history trajectory of disease) -OR- Stable or improved motor function results compared to baseline (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fabhalta

Products Affected

- FABHALTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of paroxysmal nocturnal hemoglobinuria (PNH) -AND- meets one of the following (1 or 2): 1) PNH mutant clones confirmed by flow cytometry, or 2) glycosylphosphatidylinositol-anchored proteins (GPI-AP)-deficient polymorphonuclear cells (PMNs) confirmed by flow cytometry -AND- meets one of the following (3-6): 3) anemia secondary to PNH (e.g. hemoglobin less than 10.5 g/dL with symptoms of anemia), 4) elevated lactate dehydrogenase (LDH) greater than or equal to 1.5 times the upper limit of normal, 5) history of a thromboembolic event, or 6) clinical findings of systemic complications (e.g. fatigue, hemoglobinuria, abdominal pain, dyspnea, dysphagia, erectile dysfunction, history of blood cell transfusion due to PNH) -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)) unless initially cross-titrating.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response defined as one of the following (1-3): 1) hemoglobin stabilization or increase from baseline, 2) decrease in transfusions from baseline, or 3) decrease in LDH levels from baseline or reduction of hemolysis -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Fasenra

Products Affected

- **FASENRA PEN**
- **FASENRA SUBCUTANEOUS SYRINGE**
10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe asthma -AND- history of at least one asthma exacerbations requiring oral or systemic corticosteroid treatment in the past 12 months -AND- documented pretreatment FEV1 less than 80 percent predicted in adults or FEV1 less than 90 percent predicted in children or FEV1 reversibility of at least 12% after albuterol administration despite regular treatment with (a. or b.): a) high dose inhaled corticosteroid and additional asthma controller medication or b.) a medium or high dose inhaled corticosteroid plus a long-acting beta agonist with or without oral corticosteroids and additional asthma controller medication
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Blood eosinophil count (in the absence of other potential causes of eosinophilia, including hypereosinophilic syndromes, neoplastic disease, and known or suspected parasitic infection) greater than or equal to 150 cells/microliter within 6 weeks of initiation of therapy or greater than or equal to 300 cells/uL within 12 months of screening. For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Fecal Microbiota Products

Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a recent diagnosis of recurrent <i>Clostridioides difficile</i> infection (CDI) -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	For reauthorization, attestation of recurrent CDI episodes after administration of the initial fecal microbiota product -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ferriprox

Products Affected

- *deferiprone*
- **FERRIPROX**
- **FERRIPROX (2 TIMES A DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of transfusional iron overload due to thalassemia syndromes, sickle cell disease and other anemias -AND- transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -AND- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Initial authorization requires trial and failure of deferasirox (generic Exjade) and reauthorization requires failure of generic deferasirox (generic Exjade) if not previously trialed. For reauthorization, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Fetzima

Products Affected

- **FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)** **CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG**
- **FETZIMA ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder and trial and failure of two other antidepressants.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Filspari

Products Affected

- FILSPARI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with renin-angiotensin system inhibitors (i.e., ACE inhibitors, ARBs, aliskiren) or endothelin receptor antagonists (e.g., Letairis, Opsumit, Tracleer)
Required Medical Information	Documentation of primary immunoglobulin A nephropathy confirmed by biopsy -AND- Risk for rapid disease progression, evidenced by urine protein-to-creatinine ratio greater than or equal to 1.5 g/g or proteinuria greater than or equal to 1 g/day -AND- Therapeutic failure, contraindication, or intolerance to an angiotensin converting enzyme inhibitor or an angiotensin receptor blocker
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, reduction in urine protein-to-creatinine ratio or proteinuria from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Filsuvez

Products Affected

- **FILSUVEZ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB) -AND- at least one open wound.
Age Restrictions	Deny if less than 6 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement in targeted wound(s) - AND- member requires additional courses of treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fintepla

Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravet syndrome or Lennox-Gastaut syndrome. For Lennox-Gastaut syndrome, therapeutic failure, contraindication, or intolerance to at least 2 standard of care treatments (e.g. lamotrigine, clobazam).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Firazyr

Products Affected

- **FIRAZYR**
- *icatibant*
- **SAJAZIR**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Firazyr, therapeutic failure, intolerance or contraindication to icatibant.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Firdapse

Products Affected

- **FIRDAPSE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Flector

Products Affected

- *diclofenac epolamine*
- **FLECTOR**
- **LICART**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis AND one of the following (1,2 or 3): 1) trial/failure, intolerance, or contraindication to 2 oral generic NSAIDs one of which must be diclofenac 2) hypersensitivity to oral NSAIDs 3) history or high risk for adverse gastrointestinal effects associated with oral NSAID use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fleqsuvy

Products Affected

- *baclofen oral suspension*
- **FLEQSUVY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member has an inability to swallow generic baclofen tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Forteo

Products Affected

- **FORTEO**
- *teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. For brand Forteo, trial and failure of teriparatide is required. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Forteo/teriparatide will be limited to a coverage duration of 24 months in the absence of provider attestation that the member remains at or has returned to having a high risk for fracture
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fotivda

Products Affected

- FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if applicable to diagnosis, previous therapies tried/failed
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fruzaqla

Products Affected

- **FRUZAQLA ORAL CAPSULE 1 MG, 5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Furoscix

Products Affected

- **FUROSCIX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of heart failure classified as NYHA Class II/III - AND- chronic heart failure medication regimen includes an oral loop diuretic (e.g. furosemide, bumetanide, torsemide) -AND- treatment with oral diuretics will replace the use of Furoscix as soon as practical.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gabapentin

Products Affected

- *gabapentin oral capsule 100 mg, 300 mg, 400 mg*
- *gabapentin oral solution 250 mg/5 ml*
- *gabapentin oral tablet 600 mg, 800 mg*
- **NEURONTIN ORAL CAPSULE 100 MG, 300 MG, 400 MG**
- **NEURONTIN ORAL SOLUTION**
- **NEURONTIN ORAL TABLET 600 MG, 800 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Galafold

Products Affected

- GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Fabry disease confirmed by biochemical or genetic test -AND- Presence of an amenable GLA variant causing Fabry disease in the clinical context of the patient -AND- Will not be used concomitantly with enzyme replacement therapy (ERT) e.g. Fabrazyme.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy - AND- Will not be used concomitantly with enzyme replacement therapy (ERT) e.g. Fabrazyme.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gattex

Products Affected

- **GATTEX 30-VIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of short bowel syndrome (SBS) having less than 200 cm of functional small bowel AND if age 18 and older, dependence on parenteral nutrition or intravenous nutritional support for at least 12 months AND requiring parenteral nutrition at least 3 times per week -OR- Documentation of SBS having less than 200 cm of functional small bowel AND age 1 to 17 years of age AND Dependence on parenteral nutrition AND weight of at least 10 kg.
Age Restrictions	Deny if less than 1 year of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gavreto

Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis 1) RET mutant or fusion status 2) Radioactive iodine-refractory (if radioactive iodine is appropriate)
Age Restrictions	Deny if less than 18 years of age for NSCLC or less than 12 years of age for thyroid cancer
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gilenya

Products Affected

- *fingolimod*
- GILENYA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	Brand Gilenya requires therapeutic failure or intolerance to generic fingolimod. For Brand Gilenya 0.25mg, failure of generic is not required if age is between 10-17 years old and weight is less than or equal to 40kg.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gimoti

Products Affected

- GIMOTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute or recurrent diabetic gastroparesis -AND- Attestation of no signs or symptoms of tardive dyskinesia -AND- Therapeutic failure or intolerance to generic metoclopramide (tablets, orally disintegrating tablets, or solution) or not a candidate for oral dosage forms -AND- If over 65 years of age, member was titrated to a stable dose of metoclopramide tablets or solution at 10mg four times a day before switching to Gimoti therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	For reauthorization, treatment is for a new episode of diabetic gastroparesis -AND- There has been a 2 week drug holiday without Gimoti since its last administration -AND- Attestation of no signs or symptoms of tardive dyskinesia -AND- Attestation that extended therapy with Gimoti outweighs risk of developing tardive dyskinesia.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Glatiramer

Products Affected

- **COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- **GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gleevec

Products Affected

- **GLEEVEC ORAL TABLET 100 MG, 400 MG**
- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and all of the following, if applicable to diagnosis 1) Alternatives tried 2) Concomitant therapy 3) mutation status, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Gleevec, documentation of trial and failure of generic imatinib is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gleostine

Products Affected

- GLEOSTINE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) Prior surgical procedures and/or radiotherapeutic procedures 2) Concomitant therapy 3) Alternatives tried/failed
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLP1

Products Affected

- **BYDUREON BCISE** INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- **BYETTA SUBCUTANEOUS PEN INJECTOR 10 MCG/DOSE(250 MCG/ML) 2.4 ML, 5 MCG/DOSE (250 MCG/ML) 1.2 ML**
- **MOUNJARO**
- **OZEMPIC SUBCUTANEOUS PEN**
- **RYBELSUS**
- **TRULICITY**
- **VICTOZA 3-PAK**

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Bydureon Bcise or Byetta, therapeutic failure or intolerance to two, or contraindication to all of the following: Mounjaro, Ozempic, Rybelsus, Trulicity, or Victoza.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Glycate

Products Affected

- **GLYCATE**
- *glycopyrrolate oral tablet 1.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using as an adjunctive treatment for peptic ulcer disease -AND- Therapeutic failure or intolerance to both generic glycopyrrolate 1mg tablet AND 2 mg tablet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy - AND- Additional courses of therapy are required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gocovri

Products Affected

- **GOCOVRI ORAL
CAPSULE,EXTENDED RELEASE 24HR
137 MG, 68.5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For dyskinesia associated with Parkinson's disease, documentation of concurrent levodopa-based therapy -AND- trial and failure, contraindication, or intolerance to immediate-release amantadine. For off-episodes of Parkinson's disease, documentation of concurrent carbidopa/levodopa therapy -AND- trial and failure, contraindication, or intolerance to immediate-release amantadine -AND- trial and failure, contraindication, or intolerance to one (1) of the following agents: entacapone, pramipexole, rasagiline, ropinirole, or selegiline.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gralise

Products Affected

- *gabapentin oral tablet extended release 24 hr 300 mg, 600 mg* **MG, 750 MG, 900 MG**
- **GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 300 MG, 450 MG, 600**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) -AND- Trial and failure or intolerance to generic immediate-release gabapentin. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Grastek

Products Affected

- **GRASTEK**

PA Criteria	Criteria Details
Exclusion Criteria	Asthma (severe, unstable or uncontrolled), concomitant sublingual or subcutaneous immunotherapy, therapy initiation during active allergy season
Required Medical Information	Documentation of allergic rhinitis and use for Timothy grass pollen or cross reactive grass pollens (Sweet Vernal, Orchard, Perennial Rye, Timothy, Kentucky Blue Grass pollen, Redtop, or meadow fescue) -AND- allergic rhinitis with or without conjunctivitis has been confirmed by a pollen specific positive skin test or in vitro testing for pollen-specific IgE antibodies -AND- trial and failure or intolerance to an intranasal steroid and an oral non-sedating antihistamine, intranasal antihistamine or intranasal anticholinergic agent
Age Restrictions	Deny if less than 5 years of age or greater than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Member must also be prescribed an epinephrine auto injector. For reauthorization, attestation of improved allergy symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Growth Hormone

Products Affected

- GENOTROPIN
- GENOTROPIN MINIQUICK
- HUMATROPE INJECTION CARTRIDGE
- NGENLA
- NORDITROPIN FLEXPRO
- NUTROPIN AQ NUSPIN
- OMNITROPE
- SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG
- ZOMACTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, bone age, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria. Additionally for growth failure due to chronic kidney disease, glomerular filtration rate is less than 89ml/min per 1.73m ² . For HIV wasting and cachexia, Concurrent use of antiretroviral therapy -AND- weight loss of at least 10 percent from baseline. For short bowel syndrome, receiving management for short bowel syndrome, including specialized nutritional support -AND- less than 200 cm of functional small bowel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines. For reauthorization of HIV wasting and cachexia, attestation of increase in weight from start of therapy. For reauthorization of short bowel syndrome, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Haegarda

Products Affected

- HAEGARDA

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	<p>For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-4): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. 4) Documentation of member's weight. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (5-9): 5) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 6) Documentation of family history of HAE or FXII mutation 7) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 8) Medications known to cause angioedema have been evaluated and discontinued. 9) Documentation of member's weight.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Harvoni

Products Affected

- **HARVONI ORAL PELLETS IN PACKET**
- **HARVONI ORAL TABLET 90-400 MG**
- *ledipasvir-sofosbuvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hemady

Products Affected

- **HEMADY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- used in combination with other anti-myeloma agents -AND- therapeutic failure or intolerance to generic dexamethasone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hetlioz

Products Affected

- HETLIOZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use -OR- Documented diagnosis of Smith-Magenis Syndrome as confirmed by chromosome analysis -AND- patient is experiencing nighttime sleep disturbances (e.g. difficulty falling asleep, shortened sleep cycles, inability to enter REM sleep, or frequent awaking during the night and early in the morning)
Age Restrictions	Deny if less than 16 years of age for nighttime sleep disturbances in Smith-Magenis Syndrome or deny if less than 18 years of age for Non-24 Hour Sleep-Wake Disorder
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For brand Hetlioz, documentation of trial and failure of generic tasimelteon is required. For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder -OR- attestation of positive clinical response to therapy with minimal side effects for Smith-Magenis Syndrome
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Hetlioz LQ

Products Affected

- HETLIOZ LQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Smith-Magenis Syndrome as confirmed by chromosome analysis -AND- patient is experiencing nighttime sleep disturbances (e.g. difficulty falling asleep, shortened sleep cycles, inability to enter REM sleep, or frequent awaking during the night and early in the morning)
Age Restrictions	Deny if less than 3 or greater than 15 years of age.
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of positive clinical response to therapy with minimal side effects for Smith-Magenis Syndrome -AND- member is between 3 and 15 years of age
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High-risk meds

Products Affected

- *amitriptyline*
- *amitriptyline-chlordiazepoxide*
- **ANAFRANIL**
- *benztropine oral*
- *carisoprodol*
- *chlorzoxazone*
- *clomipramine*
- *cycloheptadine*
- *doxepin oral capsule*
- *doxepin oral concentrate*
- *doxepin oral tablet*
- *ergoloid*
- *hydroxyzine hcl oral solution 10 mg/5 ml*
- *hydroxyzine hcl oral tablet*
- *hydroxyzine pamoate*
- *imipramine hcl*
- *imipramine pamoate*
- **LORZONE**
- *metaxalone*
- **NORGESIC**
- **NORGESIC FORTE**
- *orphenadrine citrate oral*
- *orphenadrine-asa-caffeine oral tablet 25-385-30 mg*
- *perphenazine-amitriptyline*
- *promethazine oral*
- **SILENOR**
- **SOMA**
- *trimipramine*
- **VISTARIL ORAL CAPSULE 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For all medications subject to this PA group, the following information (1 through 3) is required: 1. Documentation of diagnosis 2. Explanation of risk-benefit profile favoring use of the high-risk medication 3. Attestation of an intent to monitor and address treatment-related adverse events. For target tricyclic antidepressants (TCAs), in addition to criteria 1 through 3 above, trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs for the same indication, if available, is required (e.g. SSRIs and SNRIs). If using a TCA for a medically-accepted indication not shared by the safer alternatives listed, then no trial of alternatives is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Doxepin doses less than or equal to 6 mg per day will receive automatic approval.

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

High-risk meds phenobarbital

Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia
Required Medical Information	For use in seizures the following are required: 1. Explanation of risk-benefit profile favoring use of the high-risk medication 2. Attestation of an intent to monitor and address treatment-related adverse events. 3. For new starts, the trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs used for seizures (e.g. carbamazepine, lamotrigine) is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Seizure disorders
Part B Prerequisite	No

Homozygous FH

Products Affected

- JUXTAPID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of homozygous familial hypercholesterolemia (HoFH) supported by genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9, or LDLRAP1 gene -OR- untreated LDL-C greater than 400mg/dL or TC greater than 500mg/dl with cutaneous or tendon xanthoma before age 10 yrs or HeFH in both parents -AND- The member has a current LDL-C greater than 100mg/dL despite use of maximally tolerated statin OR statin intolerance -AND- The member will continue to receive concurrent lipid-lowering therapies for the treatment of HoFH
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	Patients must have a trial/failure or contraindication to the preferred product Repatha. For reauthorization, documentation showing an LDL-C reduction on Juxtapid therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Horizant

Products Affected

- **HORIZANT ORAL TABLET
EXTENDED RELEASE 300 MG, 600 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of moderate to severe active primary restless leg syndrome -AND- trial and failure of pramipexole or ropinirole -AND- trial and failure of an additional dopaminergic agent, clonidine or pregabalin -OR- Documentation of post herpetic neuralgia and trial and failure of generic gabapentin. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Humira

Products Affected

- *adalimumab-adaz*
- *adalimumab-adbm subcutaneous pen injector kit 40 mg/0.4 ml, 40 mg/0.8 ml*
- *adalimumab-adbm subcutaneous syringe kit 10 mg/0.2 ml, 20 mg/0.4 ml, 40 mg/0.4 ml, 40 mg/0.8 ml*
- **ADALIMUMAB-ADBM(CF) PEN CROHNS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 40 MG/0.8 ML**
- **ADALIMUMAB-ADBM(CF) PEN PS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 40 MG/0.8 ML**
- **CYLTEZO(CF)**
- **CYLTEZO(CF) PEN**
- **CYLTEZO(CF) PEN CROHN'S-UC-HS**
- **CYLTEZO(CF) PEN PSORIASIS-UV**
- **HUMIRA PEN**
- **HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML**
- **HUMIRA(CF)**
- **HUMIRA(CF) PEN**
- **HUMIRA(CF) PEN CROHNS-UC-HS**
- **HUMIRA(CF) PEN PEDIATRIC UC**
- **HUMIRA(CF) PEN PSOR-UV-ADOL HS**
- **HYRIMOZ PEN CROHN'S-UC STARTER**
- **HYRIMOZ PEN PSORIASIS STARTER**
- **HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML, 80 MG/0.8 ML- 40 MG/0.4 ML**
- **HYRIMOZ(CF) PEN SUBCUTANEOUS PEN INJECTOR 40 MG/0.4 ML, 80 MG/0.8 ML**
- **HYRIMOZ(CF) SUBCUTANEOUS SYRINGE 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 1 immunosuppressant or corticosteroid, or all are contraindicated.

PA Criteria	Criteria Details
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. Please Note for New Starts Only: preferred adalimumab products include Humira with NDC starting 00074, Cyltezo with NDC starting 00597, Hyrimoz with NDC starting 61314, adalimumab-adaz with NDC starting 61314, and adalimumab-adbm with NDC starting 00597. Non-preferred adalimumab products require therapeutic failure or intolerance to 2 preferred adalimumab products.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Hyftor

Products Affected

- **HYFTOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of tuberous sclerosis complex with facial angiofibromas.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of positive clinical response to therapy (e.g. decrease in size of facial angiofibromas, decrease in redness of facial angiofibromas)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ibrance

Products Affected

- **IBRANCE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of HR-positive, HER2-negative advanced or metastatic breast cancer -AND- meets one of the following (1 or 2): 1) documentation of use with an aromatase inhibitor as initial endocrine-based therapy -OR- 2) documentation of use with fulvestrant in patients with disease progression following endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ibsrela

Products Affected

- **IBSRELA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of irritable bowel syndrome with constipation -AND- failure or intolerance to Linzess -AND- if member is female, failure or intolerance to lubiprostone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Iclusig

Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of newly-diagnosed chronic phase CML
Required Medical Information	Documentation of T3151+ chronic myeloid leukemia (CML) -OR- documentation of chronic phase CML and member has experienced resistance or intolerance to at least two prior kinase inhibitors -OR- documentation of accelerated phase or blast phase CML and no other kinase inhibitor is indicated -OR- member is using Iclusig as monotherapy and meets one of the following (1-2): 1) documentation of T3151+ acute lymphoblastic leukemia (ALL) 2) documentation of Ph+ ALL and no other tyrosine kinase inhibitor therapy is indicated -OR- member has newly diagnosed Ph+ ALL and is using Iclusig in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Idhifa

Products Affected

- **IDHIFA ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IG

Products Affected

- **BIVIGAM**
- **GAMMAGARD LIQUID**
- **GAMMAGARD S-D (IGA**
- **GAMMAKED INJECTION SOLUTION**
- **GAMMAPLEX**
- **GAMMAPLEX (WITH SORBITOL)**
- **GAMUNEX-C INJECTION SOLUTION**
- **1 GRAM/10 ML (10 %)**
- **OCTAGAM**
- **PANZYGA**
- **PRIVIGEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis. For select diagnoses the following apply 1) For Myasthenia Gravis Syndrome, documentation that the patient is refractory to other standard therapies (e.g., cholinesterase inhibitors, corticosteroids, azathioprine) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 2) For Multiple Sclerosis, patient is refractory to other standard therapies (e.g., interferons) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 3) For Inflammatory Myopathies, the patient is refractory to corticosteroids given in therapeutic doses over at least 4 months OR is intolerant/contraindication to corticosteroids or immunosuppressants. 4) For B-cell CLL, associated with recurrent bacterial infections OR with Associated Hypogammaglobulinemia defined as IgG level less than 600mg/dL or evidence of a specific antibody deficiency. 5) For Bone Marrow Transplantation, the member is 20 years of age or older and within the first 100 days after transplantation. 6) For Dermatomyositis/Polymyositis, trial and failure, intolerance, or contraindication to standard fist line therapy (i.e. corticosteroids or immunosuppressants). 7) For Pediatric HIV, the patient is less than 13 y.o. who have entry CD4 lymphocyte count greater than or equal to 200/mcl and IgG less than 400mg/dL OR recurrent bacterial infections. 8) For Guillain-Barre syndrome, impaired function by objective assessment and/or objective findings on physical exam at the time of initial therapy and IVIG therapy must be initiated within 2 weeks of symptom onset. 9) For Autoimmune Mucocutaneous Blistering Diseases (e.g. Stevens-Johnson Syndrome), trial and failure, intolerance, or contraindication to conventional therapy (e.g. corticosteroids) or the patient has rapidly progressive disease in which a clinical response could not be affected quickly enough using conventional agents.</p>

PA Criteria	Criteria Details
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when administered in the home to a member with a diagnosis of primary immunodeficiency disease
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Myasthenia Gravis syndrome, Multiple Sclerosis, Inflammatory Myopathies, Polymyositis, Bone Marrow Transplant, Pediatric HIV, Guillain-Barre syndrome, Autoimmune Mucocutaneous Blistering Diseases
Part B Prerequisite	No

Ilumya

Products Affected

- ILUMYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate, cyclosporine) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For psoriasis, patients must have therapeutic failure or intolerance to 2 of the preferred products: Taltz, a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Otezla, Stelara, Enbrel and Skyrizi. For psoriasis induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Imbruvica

Products Affected

- **IMBRUVICA ORAL CAPSULE 140 MG, 280 MG, 420 MG, 70 MG**
- **IMBRUVICA ORAL SUSPENSION**
- **IMBRUVICA ORAL TABLET 140 MG,**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) 17p deletion status 2) Alternatives tried/failed 3) concomitant therapy. For suspension, members must also have an inability to swallow oral tablets or oral capsules.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inbrija

Products Affected

- **INBRIJA INHALATION CAPSULE,
W/INHALATION DEVICE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use for the treatment of intermittent off episodes of Parkinson's disease while on carbidopa/levodopa
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Increlex

Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe primary IGF-1 deficiency and all of the following: 1) Normal or elevated response (greater than 10 ng/ml) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 2) Serum IGF-1 concentration that is less than or equal to three (3) standard deviations below the normal value based on laboratory reference range. 3) Height less than or equal to three (3) standard deviations below normal (at or below the third percentile for gender and age). 4) If female, bone age less than or equal to 14 years. If male, bone age less than or equal to 16. - OR- Documentation of diagnosis of growth hormone deficiency caused by gene deletion and all of the following: 1) Growth velocity at least 2 standard deviations below the age-appropriate mean or height at least 2.25 standard deviations below the age-appropriate mean. 2) Subnormal response (less than 10 ng/mL) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 3) Development of neutralizing antibodies to growth hormone product(s). 4) If female, bone age of less than or equal to 14 years. If male, bone age less than or equal to 16.
Age Restrictions	Deny if greater than 17 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Ingrezza

Products Affected

- **INGREZZA INITIATION PK(TARDIV)**
- **INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG**
- **INGREZZA SPRINKLE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of tardive dyskinesia -OR- documentation of chorea associated with Huntington's disease.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inlyta

Products Affected

- INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) Alternatives tried/failed 2) Concomitant therapy 3) Line of therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inqovi

Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation chronic myelomonocytic leukemia. Documentation of de novo or secondary myelodysplastic syndrome -AND- One of the following (1 or 2): 1) French American-British MDS subtypes of refractory anemia, refractory anemia with ringed sideroblasts or refractory anemia with excess blasts. 2) International Prognostic Scoring System group of intermediate-1, intermediate-2 or high-risk.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inrebic

Products Affected

- **INREBIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate-2 or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) -AND- If a new start, baseline platelet count of greater than $50 \times 10^9/L$
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Interferon Alfa

Products Affected

- PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Interferon Beta

Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT
- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML
- REBIF (WITH ALBUMIN)
- REBIF REBIDOSE SUBCUTANEOUS PEN INJECTOR 22 MCG/0.5 ML, 44 MCG/0.5 ML, 8.8MCG/0.2ML-22 MCG/0.5ML (6)
- REBIF TITRATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Interleukin-1b Blockers

Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), documentation of need for maintenance of remission. For Recurrent Pericarditis (RP), documentation of trial/failure or intolerance to one, or contraindication to all of the following: oral nonsteroidal anti-inflammatory drug (NSAID), systemic corticosteroid, or colchicine.
Age Restrictions	Deny if less than 12 years of age for Recurrent Pericarditis and Cryopyrin-Associated Periodic Syndromes
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For DIRA: patient must weigh 10kg or more
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Intrarosa

Products Affected

- INTRAROSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IPF Agents

Products Affected

- **ESBRIET ORAL CAPSULE**
- **ESBRIET ORAL TABLET**
- **OFEV**
- *pirfenidone oral capsule*
- *pirfenidone oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 50% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30%. For Ofev only, documentation of systemic sclerosis-associated interstitial lung disease -AND- baseline forced vital capacity (FVC) of at least 40% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- documentation of a high-resolution chest computed tomography (CT) scan demonstrating greater than or equal to 10% pulmonary fibrosis. For Ofev only, documentation of chronic fibrosing interstitial lung disease with progressive phenotype -AND- high resolution chest computing tomography (HRCT) scan demonstrating greater than 10% fibrosing disease -AND- baseline forced vital capacity (FVC) of at least 45% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- disease progression in previous 24 months shown by one of the following : 1. Relative decline in FVC greater than or equal to 10% predicted 2. Relative decline in FVC greater than or equal to 5% but less than 10% predicted and either worsening of respiratory symptoms or increased extent of fibrotic changes on HRCT 3. Worsening of respiratory symptoms and increasing extent of fibrotic changes on HRCT
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Esbriet, documentation of failure on generic pirfenidone tablets is required.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Iressa

Products Affected

- *gefitinib*
- **IRESSA**

PA Criteria	Criteria Details
Exclusion Criteria	Use in tumors with EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.
Required Medical Information	Documentation of diagnosis -AND- the following: 1) EGFR exon 19 deletion mutations or exon 21 (L858R) mutations as detected by an FDA-approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Isturisa

Products Affected

- **ISTURISA ORAL TABLET 1 MG, 5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Cushing's disease AND patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of mean urine free cortisol (mUFC) decrease from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Itraconazole

Products Affected

- *itraconazole*
- SPORANOX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If using for diagnosis of onychomycosis, confirmation through positive laboratory testing (e.g. KOH preparation, fungal culture, or nail biopsy) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Onychomycosis: 3 months. All other indications: 3 months initial, 12 months reauth
Other Criteria	Documentation of trial/failure or intolerance of amphotericin b must be provided for approval in patients with aspergillosis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ivermectin Oral

Products Affected

- *ivermectin oral*
- **STROMEKTOL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of strongyloidiasis of the intestinal tract (non-disseminated disease) or onchocerciasis -AND- Member weighs greater than or equal to 15kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Iwilfin

Products Affected

- IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of high-risk neuroblastoma (HRNB) -AND- partial response to anti-glycolipid disialoganglioside (GD2) immunotherapy (e.g., dinutuximab, naxitamab).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jakafi

Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) and if a new start, baseline platelet count of greater than $50 \times 10^9/L$ -OR- documentation of polycythemia vera and inadequate response or intolerance to hydroxyurea -OR- Documentation of steroid refractory acute graft-versus-host disease and prior therapy with at least one systemic corticosteroid -OR- Documentation of chronic graft-versus-host disease with prior failure of at least one systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jaypirca

Products Affected

- **JAYPIRCA ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if applicable to diagnosis, alternatives tried/failed
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Joenja

Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of activated phosphoinositide 3-kinase delta syndrome (APDS) with genetic confirmation of variant in PIK3CD or PIK3R1 gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jynarque

Products Affected

- JYNARQUE ORAL TABLET
- JYNARQUE ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of rapidly progressing autosomal dominant polycystic kidney disease defined by one of the following: 1.) Historical decline in eGFR greater than or equal to 5mL/min/1.73 m*2 within a 12 month period. 2.) Decline in eGFR of greater than or equal to 2.5mL/min/1.73m*2 over a period of 5 years. 3.) 5% increase in total kidney volume per year by 3 repeat CT or MRI. 4.) Average kidney length greater than 16.5cm. 5.) Family history of end-stage renal disease before age 58. 6.) Mayo imaging classification of 1C, 1D, or 1E. 7.) Kidney bleeds.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, prescriber attestation of slowed decline of kidney function
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kalydeco

Products Affected

- **KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG**
- **KALYDECO ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to ivacaftor based on clinical and or in vitro assay (e.g. G551D, G1244E, G1349D)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kerendia

Products Affected

- **KERENDIA**

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than or equal to 25 mL/min/1.73 m*2, serum potassium greater than 5.5 mEq/L
Required Medical Information	Documentation of type 2 diabetes mellitus with chronic kidney disease - AND- one of the following (1. 2. or 3.): 1) concomitant use of a sodium-glucose Cotransporter-2 (SGLT2) inhibitor 2) therapeutic failure to at least one SGLT2 inhibitor or 3) contraindication or intolerance to at least one SGLT2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that signs or symptoms of hyperkalemia are not present.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kesimpta

Products Affected

- **KESIMPTA PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Keveyis

Products Affected

- KEVEYIS
- ORMALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of one of the following: 1. Primary hyperkalemic periodic paralysis 2. Primary hypokalemic periodic paralysis 3. Related variants of primary periodic paralysis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	2 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation the number of muscle weakness attacks per week has decreased from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kevzara

Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For polymyalgia rheumatica (PMR), one of the following (1-3): 1) inadequate response to corticosteroids, 2) intolerance to corticosteroid taper, or 3) used in combination with corticosteroid tapering course. For juvenile idiopathic arthritis, weight greater than or equal to 63 kg -AND- one of the following (4-5): 4) inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide), or 5) requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis and polymyalgia rheumatica.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to two of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Actemra SC, Xeljanz/Xeljanz XR and Rinvoq. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Rinvoz/Rinvoq LQ, Xeljanz/Xeljanz solution and Actemra SC.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Kineret

Products Affected

- KINERET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), therapeutic failure or intolerance to at least one (1) corticosteroid, or all corticosteroids are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products Humira, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kisqali

Products Affected

- **KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG**
- **KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following. 1) HR mutation status and HER2 mutation status 2) Alternatives tried/failed 3) Concomitant therapy, if applicable to diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Klisyri

Products Affected

- **KLISYRI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of actinic keratosis of the face or scalp -AND- Therapeutic failure or intolerance to 1 of the following 1) generic topical imiquimod 5% cream 2) generic fluorouracil 5% topical cream 3) generic fluorouracil topical solution
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation that the member has previously experienced complete or partial clearance of actinic keratosis lesions with Klisyri -AND- additional course of therapy is required for recurrence of actinic keratosis -AND- member is restarting therapy at least 60 days after cessation of an initial Klisyri 5-day course.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Korlym

Products Affected

- **KORLYM**
- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperglycemia secondary to hypercortisolism in patients with endogenous Cushing's syndrome -AND- Patient is not a candidate for surgery or where surgery has failed -AND- one of the following (1 or 2): 1) Diagnosis of diabetes with trial and failure, intolerance, or contraindication to one previous therapy for Type 2 Diabetes (e.g. metformin, sulfonylureas, insulin) or using in addition to a therapy for Type 2 diabetes. 2) Glucose intolerance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Korlym, a trial and failure of generic mifepristone is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Koselugo

Products Affected

- **KOSELUGO ORAL CAPSULE 10 MG,
25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- for neurofibromatosis type 1 (NF1), documentation of symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Krazati

Products Affected

- KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- member is using Krazati as a single agent. Documentation of locally advanced or metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA-approved test -AND- member using using Krazati in combination with cetuximab -AND- member has received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kuvan

Products Affected

- **JAVYGTOR**
- **KUVAN**
- *sapropterin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of PKU -AND- documented baseline Phe level greater than 6 mg/dL -AND- clinical documentation of current weight -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, initial therapy has resulted in decrease in phenylalanine levels from baseline or current phenylalanine levels within the range of 120-360 micromol/L -AND- clinical documentation of current weight is required -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day. For brand Kuvan, a trial and failure of generic sapropterin dihydrochloride is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lampit

Products Affected

- LAMPIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Weight of at least 2.5 kg
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	2 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Latuda

Products Affected

- **LATUDA ORAL TABLET 120 MG, 20 MG, 40 MG, 60 MG, 80 MG**
- *lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for bipolar 1 disorder, documentation of trial and failure or intolerance to one other formulary medication indicated in bipolar 1 disorder (e.g. quetiapine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Latuda, documentation of trial and failure of generic lurasidone is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lenvima

Products Affected

- LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following if applicable to diagnosis: 1) Radioactive iodine refractory status 2) Microsatellite instability-high status or mismatch repair deficient status 3) Alternatives tried/failed or attestation of first line use 4) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Leukine

Products Affected

- LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis: following induction chemotherapy in patients who are 55 years or older with acute myelogenous leukemia (AML) -OR- mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation -OR- acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation -OR- acceleration of myeloid reconstitution following allogeneic BMT -OR- treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic BMT -OR- following acute exposure to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Leukotriene Modifiers

Products Affected

- *zileuton*
- **ZYFLO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of asthma -AND- trial/failure of generic montelukast and generic zafirlukast
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Libervant

Products Affected

- **LIBERVANT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 2 years of age or greater than 5 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lidoderm

Products Affected

- *lidocaine topical adhesive patch, medicated 5 %*
- **LIDOCAN III**
- **LIDODERM**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) -AND- trial and failure of 1 other agent used to treat PHN (e.g. gabapentin), inability to swallow oral medication or unable to take an oral medication due to potential adverse events (e.g. sedation) -OR- documentation of diabetic peripheral neuropathy (DPN) -AND- trial and failure of one other agent used to treat DPN (e.g. duloxetine), inability to swallow oral medication or unable to take an oral medication due to potential adverse events (e.g. sedation)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	diabetic peripheral neuropathy
Part B Prerequisite	No

Litfulo

Products Affected

- LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For severe alopecia areata, therapeutic failure/intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Livmarli

Products Affected

- LIVMARLI ORAL SOLUTION 9.5 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Cholestatic Pruritus -AND- Documentation of one of the following diagnoses (1-2): 1) Alagille syndrome, 2) progressive familial intrahepatic cholestasis confirmed by genetic testing that is not PFIC type 2 with a ABCB11 variant resulting in nonfunctional or absent bile salt export pump (BSEP) protein -AND- Documentation that the member does not have any of the following diagnoses (3-5): 3) decompensated cirrhosis, 4) portal hypertension, 5) history of a hepatic decompensation event.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial, 12 months reauthorization
Other Criteria	For reauthorization, attestation of improvement in pruritus -AND- attestation that the member has not progressed to any of the following (1-3): 1) portal hypertension, 2) cirrhosis, or 3) experienced a hepatic decompensation event.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Livtency

Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of CMV prophylaxis.
Required Medical Information	Documentation of refractory cytomegalovirus infection or disease as evidenced by antigenemia or polymerase chain reaction (PCR) test -AND- all of the following (1-3): 1) member weighs at least 35 kg, 2) member is a recipient of hematopoietic stem cell transplant -OR- solid organ transplant. 3) member has experienced therapeutic failure to one of the following: ganciclovir, valganciclovir, cidofovir, or foscarnet.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of a previous reduction in CMV DNA level -AND- documentation of one of the following (1-3): 1) new onset symptomatic CMV infection, 2) virologic relapse with treatment-emergent maribavir resistance or 3) continued antiviral treatment is required to achieve virologic clearance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lodoco

Products Affected

- LODOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of established atherosclerotic disease (e.g., ASCVD, experienced CV event) or multiple risk factors for cardiovascular disease (e.g., high cholesterol, high blood pressure, obesity, diabetes, smoking) - AND- being used to reduce the risk of myocardial infarction, stroke, coronary revascularization, or cardiovascular death -AND- trial and failure of maximally tolerated statin or documentation of statin intolerance.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lokelma

Products Affected

- **LOKELMA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 7.4 mmol/L on at least two (2) screenings -AND- Modification of medications to reduce serum potassium levels were not successful, when applicable
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in serum potassium levels following Lokelma administration and continued treatment for hyperkalemia is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lonsurf

Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer as a single agent or in combination with bevacizumab in patients who have previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy -OR- documentation of metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/neu-targeted therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lorbrena

Products Affected

- **LORBRENA ORAL TABLET 100 MG,
25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ALK-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lotronex

Products Affected

- *alosetron oral tablet 0.5 mg, 1 mg*
- **LOTROXEX ORAL TABLET 0.5 MG, 1 MG**

PA Criteria	Criteria Details
Exclusion Criteria	For irritable bowel syndrome (IBS): Exclude if male gender
Required Medical Information	Documentation of chronic severe diarrhea-predominant IBS -AND- trial and failure or intolerance to one anti-diarrheal (e.g. loperamide), anti-spasmodic, or tricyclic antidepressant, or contraindication to all
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that symptoms of IBS continue to persist AND positive clinical response.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lovaza

Products Affected

- LOVAZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For hypertriglyceridemia, triglycerides greater than or equal to 500mg/dL indicating sever hypertriglyceridemia -AND- Therapeutic failure or intolerance to a prescription, generic omega 3 acid ethyl ester capsules
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, reduction in triglyceride levels from baseline for hypertriglyceridemia
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lumakras

Products Affected

- **LUMAKRAS ORAL TABLET 120 MG, 320 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and all of the following, if applicable to diagnosis 1) KRAS G12C mutation status, as detected by an FDA-approved test 2) Alternatives tried
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lumryz

Products Affected

- LUMRYZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	<p>If no diagnosis of cataplexy- trial and failure, intolerance, or contraindication to generic modafinil AND a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lupkynis

Products Affected

- LUPKYNIS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of active lupus nephritis -AND- Documented positive ANA titer (greater than or equal to 1:80) or anti-dsDNA greater than or equal to 30 IU/mL -AND- Trial, contraindication, intolerance, or inadequate response to at least 2 of the following standard of care drug classes: 1.) corticosteroids (e.g. prednisone) 2.) antimalarials (e.g. hydroxychloroquine) 3.) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate) -AND- Member will continue to receive concomitant standard of care treatment with corticosteroids (e.g. prednisone) and mycophenolate mofetil.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	24 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of disease stability or disease improvement -AND- member will continue to receive concomitant standard of care treatment with corticosteroids (e.g. prednisone) and mycophenolate mofetil.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lupron Depot Ped

Products Affected

- **LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG**
- **LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)**
- **LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of central precocious puberty -AND- advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3 mIU/L or leuprolide-stimulating LH level greater than 3.3-5 mIU/L
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy defined as pre-pubertal slowing/decline, normalization of FSH, normalization LH, normalization of bone age, normalization of estradiol level or normalization of testosterone level
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lybalvi

Products Affected

- LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of schizophrenia or bipolar I disorder -AND- One of the following (1 or 2): 1) Therapeutic failure, intolerance or contraindication to generic risperidone and generic quetiapine 2) Currently stable and responding to olanzapine but experiencing weight gain from the medication
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lynparza

Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) BRCA mutations 2) Genomic instability status 3) Homologous recombinant repair gene mutations 4) HER2 status 5) Alternatives tried/failed 6) Concomitant therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lyrica

Products Affected

- LYRICA CR
 - LYRICA ORAL CAPSULE 100 MG, 150 MG, 200 MG, 225 MG, 25 MG, 300 MG, 50 MG, 75 MG
 - LYRICA ORAL SOLUTION
- *pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg*
 - *pregabalin oral solution*
 - *pregabalin oral tablet extended release 24 hr*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For immediate release and controlled release tablets, documentation of DPN and trial/failure or intolerance to duloxetine -OR- PHN and trial/failure or intolerance to gabapentin. For immediate release capsules, documentation of partial-onset seizures and trial/failure or intolerance to two AEDS -OR- neuropathic pain associated with spinal cord injury -OR- documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine. When using pregabalin products concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lytgobi

Products Affected

- **LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) FGFR2 fusion or other rearrangement 2) Previous therapies tried/failed
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lyvispah

Products Affected

- LYVISPAH

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member has an inability to swallow generic baclofen tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mavenclad

Products Affected

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Mavenclad and other disease modifying agents such as interferons, Copaxone, Tysabri. Treatment duration greater than 24 months.
Required Medical Information	Documentation of diagnosis of relapse-remitting multiples sclerosis or active secondary progressive disease -AND- therapeutic failure or intolerance to one other disease modifying therapy (e.g. Avonex, Gilenya, Copaxone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Coverage beyond 24 months will not be approved.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mavyret

Products Affected

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mayzent

Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Mayzent and other disease modifying agents such as interferons, Copaxone, Tysabri.
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Testing for CYP2C9 variants has confirmed member does not have CYP2C9*3/*3 genotype
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Megace

Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mekinist

Products Affected

- **MEKINIST ORAL TABLET 0.5 MG, 2 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following. 1) BRAF mutations, if applicable to diagnosis. 2) Alternatives tried/failed. 3) Concomitant therapy, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mekinist Suspension

Products Affected

- MEKINIST ORAL RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following. 1) BRAF mutations, if applicable to diagnosis. 2) Alternatives tried/failed. 3) Concomitant therapy, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation of inability to swallow Mekinist tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mektovi

Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) BRAF mutation status 2) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Methamphetamine

Products Affected

- *methamphetamine*

PA Criteria	Criteria Details
Exclusion Criteria	Obesity
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For ADHD, trial/failure or intolerance to 2 of the following generic medications: methylphenidate, atomoxetine, or dextroamphetamine/amphetamine is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Metyrosine

Products Affected

- **DEMSER**
- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pheochromocytoma defined by 1 of the following (1 or 2): 1) Elevated metanephrines in plasma or urine, or 2) Tumor evidence from CT scan or MRI -AND- Documentation of 1 of the following (3., 4., or 5.): 3) Planned resection surgery, 4) Resection surgery is contraindicated, or 5) malignant pheochromocytoma. -AND- therapeutic failure, contraindication, or intolerance to a selective alpha-blocker (e.g., doxazosin, prazosin, terazosin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Demser, documentation of failure on generic metyrosine is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Motegrity

Products Affected

- **MOTTEGRITY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic idiopathic constipation -AND- Therapeutic failure, contraindication or intolerance to Linzess and lubiprostone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Motpoly XR

Products Affected

- **MOTPOLY XR ORAL
CAPSULE,EXTENDED RELEASE 24HR
100 MG, 150 MG, 200 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial onset seizures or adjunctive therapy for the treatment of primary generalized tonic-clonic seizures -AND- weight greater than or equal to 50kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Therapeutic failure or intolerance to generic lacosamide tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MTX Injection

Products Affected

- OTREXUP (PF)
- RASUVO (PF)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis AND trial/failure, intolerance, or contraindication to oral generic methotrexate tablets. For rheumatoid arthritis and juvenile idiopathic arthritis, disease is classified as severe and active -AND- therapeutic failure or intolerance to first-line therapy (e.g., nonsteroidal anti-inflammatory drug [NSAID]). For psoriasis, disease is classified as severe, recalcitrant, and disabling -AND- therapeutic failure or intolerance to at least one (1) other standard psoriasis therapy (e.g., cyclosporine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mulpleta

Products Affected

- MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -AND- trial and failure or intolerance to Doptelet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Myalept

Products Affected

- MYALEPT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of congenital or acquired generalized lipodystrophy - AND- Leptin levels less than 8 ng/mL for males or less than 12 ng/mL for females -AND- the member has a diagnosis of diabetes or fasting insulin levels greater than 30uU/mL or fasting hypertriglyceridemia greater than 200mg/dL -AND- Therapeutic failure to one previous diabetes therapy (e.g. metformin, insulin) or hypertriglyceridemia therapy (e.g. statin, fibrate)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, decrease in HbA1c from baseline, decrease in fasting plasma glucose from baseline or decrease in fasting triglycerides from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Myasthenia Gravis

Products Affected

- **ZILBRYSQ SUBCUTANEOUS
SYRINGE 16.6 MG/0.416 ML, 23
MG/0.574 ML, 32.4 MG/0.81 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized myasthenia gravis (gMG) -AND- Anti-acetylcholine receptor (AChR) antibody-positive -AND- Therapeutic failure, contraindication, or intolerance to generic pyridostigmine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of gMG signs/symptoms improvement (e.g., speech, swallowing, mobility, and/or respiratory function) -OR- decreased gMG exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mycapssa

Products Affected

- MYCAPSSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acromegaly -AND- High pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- Previous response to and tolerated treatment with octreotide or lanreotide.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Myfembree

Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of severe hepatic impairment or osteoporosis.
Required Medical Information	Documentation of premenopausal woman with uterine leiomyomas - AND- Experiencing heavy menstrual bleeding -AND- For women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months -OR- Documentation of premenopausal woman with endometriosis with moderate to severe pain -AND- For women of childbearing age, attestation of not pregnant - AND- Therapeutic failure or contraindication to 2 standard of care treatments (i.e. generic NSAID, combined hormonal contraceptive, progestin, GnRH agonist) -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist in combination with estrogen and progestin does not exceed 24 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 18 months reauthorization
Other Criteria	For reauthorization of uterine leiomyomas, attestation of continued experience of heavy menstrual bleeding -AND- Attestation of decrease in menstrual blood loss -AND- for women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months. For reauthorization of endometriosis, continued pain associated with endometriosis -AND- Attestation of reduction in pain - AND - For women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Namenda

Products Affected

- **NAMENDA TITRATION PAK**
- **NAMENDA XR ORAL CAPSULE,SPRINKLE,ER 24HR 14 MG, 21 MG, 28 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and trial/failure of generic memantine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Namzarin

Products Affected

- NAMZARIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and trial/failure of generic memantine and generic donepezil
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nayzilam

Products Affected

- NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nerlynx

Products Affected

- **NERLYNX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following. 1) HER2 mutations, if applicable to diagnosis. 2) Alternatives tried/failed. 3) Concomitant therapy, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nexavar

Products Affected

- **NEXAVAR**
- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally recurrent or metastatic, progressive, differentiated thyroid carcinoma, refractory to radioactive iodine treatment
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Nexavar, documentation of trial and failure of generic sorafenib is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nexletol

Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>For HeFH, diagnosis supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite Simon Broome register criteria, or definite on the Make Early Diagnosis to Prevent Early Deaths tool -AND- LDL-C greater than 100 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe. For Hyperlipidemia with ASCVD or Hyperlipidemia with attestation of high risk for CVD, LDL-C greater than 70 mg/dL -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe. For Primary Hyperlipidemia not associated with ASCVD or HeFH, LDL-C greater than 70 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For reauthorization, documentation showing an LDL-C reduction from baseline -AND- for HeFH and Primary Hyperlipidemia (not associated with ASCVD or HeFH), attestation of continued use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statin which resolved upon discontinuation of statin or documentation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ninlaro

Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- previous treatment with at least 1 prior therapy -AND- used in combination with lenalidomide and dexamethasone
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nitisinone

Products Affected

- *nitisinone*
- **NITYR**
- **ORFADIN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary tyrosinemia type 1 (HT-1) confirmed by biochemical or genetic testing -AND- dietary restriction of tyrosine and phenylalanine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Nityr and Brand Orfadin capsule, documentation of failure on generic nitisinone capsules is required. For Orfadin suspension, documentation of failure on generic nitisinone capsules or an inability to swallow capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Northera

Products Affected

- *droxidopa oral capsule 100 mg, 200 mg, 300 mg*
- **NORTHERA ORAL CAPSULE 100 MG, 200 MG, 300 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of neurogenic orthostatic hypotension caused by primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency or non-diabetic autonomic neuropathy -AND- documentation of inadequate response, intolerance or contraindication to preferred generic alternative midodrine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For brand Northera, initial authorization requires documentation of trial and failure of generic droxidopa and reauthorization requires failure of generic droxidopa if not previously trialed. For reauthorization, attestation of increase from baseline of systolic or diastolic blood pressure upon standing -OR- attestation of decrease from baseline of neurogenic orthostatic hypotension symptoms upon standing (e.g., dizziness, feeling faint, etc.).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nourianz

Products Affected

- NOURIANZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease experiencing off episodes -AND- Used as adjunct to levodopa/carbidopa -AND- trial/failure, contraindication or intolerance to one of the following generic products: ropinirole, pramipexole, entacapone, selegiline, or rasagiline
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nubeqa

Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following if applicable to diagnosis: 1. Concomitant therapy 2. History of a bilateral orchiectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nucala

Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of diagnosis of severe asthma evidenced by pretreatment FEV1 less than 80 percent predicted in adults or FEV1 less than 90 percent predicted in adolescents or FEV1 reversibility of at least 12% after albuterol administration -AND- 1.) A history of 2 or more exacerbations in the previous year or inadequate symptom control with inhaled corticosteroid in combination with 3 months of controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), unless intolerant of or contraindication to all agents -AND- 2 or 3. 2) Greater than or equal to 150 cells/uL screening within 6 weeks of dosing. 3) Greater than or equal to 300 cells/uL within 12 months of screening. -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) in patients who have a history of relapsing or refractory disease and will be receiving concomitant glucocorticoid treatment with or without immunosuppressive therapy -OR- Documentation of hypereosinophilic syndrome (HES) without an identifiable non-hematologic secondary cause for greater than or equal to 6 months -AND- At least 2 HES flares (HES-related worsening of clinical symptoms or blood eosinophil counts requiring an escalation in therapy) within the past 12 months -AND- Stable on HES therapy for at least 4 weeks (chronic or episodic oral corticosteroids, immunosuppressive or cytotoxic therapy) -OR- Documentation of chronic rhinosinusitis with nasal polyps (CRSwNP) and trial/failure, contraindication, or intolerance to an intranasal corticosteroid.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For Severe Asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EGPA reauthorization, attestation of one of the following is required (5-8): 5) reduction in frequency and/or severity of relapses, 6) reduction or discontinuation of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms. For HES reauthorization, attestation of one of the following is required (9-10): 9) reduction in frequency of HES flares, or 10) maintenance or reduction in background HES therapy requirements. For CRSwNP reauthorization, attestation of one of the following is required (11-12): 11) decrease in nasal polyp score, or 12) reduction in nasal congestion/obstruction severity score.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuedexta

Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting improvement in symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuplazid

Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hallucinations and delusions associated with Parkinson's disease psychosis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nurtec

Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine Prevention, defined as 4-14 migraine days per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For acute treatment of migraine with or without aura, trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency. For reauthorization of acute treatment of migraine, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuvigil

Products Affected

- *armodafinil*
- NUVIGIL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required. For brand Nuvigil, documentation of failure on modafinil (generic Provigil).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ocaliva

Products Affected

- OCALIVA

PA Criteria	Criteria Details
Exclusion Criteria	Compensated cirrhosis with evidence of portal hypertension
Required Medical Information	Documentation of primary biliary cholangitis -AND- trial and failure, contraindication, or intolerance to ursodiol monotherapy -AND- will use concomitantly with ursodiol unless contraindicated or intolerant.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Octreotide

Products Affected

- *octreotide acetate injection solution*
- **SANDOSTATIN INJECTION SOLUTION 100 MCG/ML, 50 MCG/ML, 500 MCG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- therapeutic failure or cannot be treated with surgical resection, pituitary irradiation or bromocriptine mesylate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Odactra

Products Affected

- ODACTRA

PA Criteria	Criteria Details
Exclusion Criteria	Asthma (severe, unstable or uncontrolled), concomitant sublingual or subcutaneous immunotherapy
Required Medical Information	Documentation of allergic rhinitis due to house dust mites -AND- allergic rhinitis with or without conjunctivitis has been confirmed by skin testing for licensed house dust mite allergen extracts or in vitro testing for IgE antibodies to D. pteronyssinus or D. Farina -AND- trial and failure or intolerance to an intranasal steroid and an oral non-sedating antihistamine, intranasal antihistamine or intranasal anticholinergic agent
Age Restrictions	Deny if less than 12 years of age or greater than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Member must also be prescribed an epinephrine auto injector. For reauthorization, attestation of improved allergy symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Odomzo

Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced basal cell carcinoma (laBCC) that has recurred following surgery or radiation therapy or for use in patients who are not candidates for surgery or radiation therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ogsiveo

Products Affected

- **OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressing desmoid tumor(s) requiring systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ojemda

Products Affected

- **OJEMDA ORAL SUSPENSION FOR RECONSTITUTION**
- **OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory pediatric low-grade glioma - AND- one of the following (1-2): 1) BRAF fusion or rearrangement, or 2) BRAF V600K mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ojjaara

Products Affected

- OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of anemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Olpruva

Products Affected

- **OLPRUVA ORAL PELLETS IN PACKET 2 GRAM, 3 GRAM, 4 GRAM, 5 GRAM, 6 GRAM, 6.67 GRAM**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase -AND- use as an adjunctive therapy to dietary management.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Trial and failure of generic sodium phenylbutyrate is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Olumiant

Products Affected

- OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis an inadequate response or intolerance to at least one non-biologic DMARD (e.g., methotrexate, leflunomide). For severe alopecia areata, inadequate response or intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Alopecia Areata
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products Humira, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR, with at least 1 being a tumor necrosis factor blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OmvoH

Products Affected

- OMVOH PEN
- OMVOH SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation of clinical response or remission following 3 doses of IV Omvoh within 3 months of initiating therapy with Omvoh SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Rinvoq, Xeljanz/Xeljanz XR, and Stelara SC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Onfi

Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*
- **ONFI ORAL SUSPENSION**
- **ONFI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ongentys

Products Affected

- **ONGENTYS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease experiencing off episodes -AND- Used as adjunct to levodopa/carbidopa -AND- trial/failure, contraindication or intolerance to the following (1. and 2.): 1) entacapone 2) one of the following generic products: rasagiline, pramipexole, ropinirole, selegiline
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Onureg

Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute myeloid leukemia that has achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy -AND- Inability to complete intensive curative therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Opzelura

Products Affected

- OPZELURA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-4): 1) mild to moderate atopic dermatitis (AD), 2) attestation of up to 20 percent of body surface area (excluding scalp) with AD involvement, 3) trial & failure, intolerance, or contraindication to at least one topical corticosteroid -OR- documentation of facial or anogenital involvement, 4) trial & failure, intolerance, or contraindication to topical generic tacrolimus or topical generic pimecrolimus -OR- Documentation of all of the following (5-7): 5) Nonsegmental vitiligo, 6) body surface area with vitiligo involvement does not exceed 10%, 7) trial & failure or intolerance to at least 1 generic, formulary high or ultrahigh potency topical corticosteroid -OR- vitiligo with facial or anogenital involvement
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	Initial authorization: 8 weeks for AD, 24 weeks for vitiligo. Reauthorization: 12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy for atopic dermatitis -OR- meaningful repigmentation of affected areas for vitiligo
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Oralair

Products Affected

- **ORALAIR SUBLINGUAL TABLET 300
INDX REACTIVITY**

PA Criteria	Criteria Details
Exclusion Criteria	Asthma (severe, unstable or uncontrolled), concomitant sublingual or subcutaneous immunotherapy, therapy initiation during active allergy season
Required Medical Information	Documentation of allergic rhinitis and use for Sweet Vernal, Orchard, Perennial Rye, Timothy or Kentucky Blue Grass pollens -AND- allergic rhinitis with or without conjunctivitis has been confirmed by a pollen specific positive skin test or in vitro testing for pollen-specific IgE antibodies -AND- trial and failure or intolerance to an intranasal steroid and an oral non-sedating antihistamine, intranasal antihistamine or intranasal anticholinergic agent
Age Restrictions	Deny if less than 5 years of age or greater than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Member must also be prescribed an epinephrine auto injector. For reauthorization, attestation of improved allergy symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orencia

Products Affected

- **ORENCIA CLICKJECT**
- **ORENCIA SUBCUTANEOUS SYRINGE**
125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients over 18 years of age must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Enbrel, Taltz, Xeljanz/Xeljanz XR, Otezla and Stelara SC, Rinvoq, and Skyrizi SC. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Xeljanz/Xeljanz solution and Actemra SC.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Orgovyx

Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Oriahnn

Products Affected

- **ORIAHNN**

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of severe hepatic impairment or osteoporosis.
Required Medical Information	Documentation of premenopausal woman with uterine leiomyomas -AND- Experiencing heavy menstrual bleeding -AND- For women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 18 months reauthorization
Other Criteria	For reauthorization, attestation of continued experience of heavy menstrual bleeding -AND- Attestation of decrease in menstrual blood loss -AND- for women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orilissa

Products Affected

- **ORILISSA ORAL TABLET 150 MG, 200 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of severe hepatic impairment or osteoporosis.
Required Medical Information	Documentation of premenopausal woman with diagnosis of endometriosis with moderate to severe pain -AND- For women of childbearing age, attestation of not pregnant -AND- Inadequate response, failure or contraindication to 2 standard of care treatments (e.g. NSAIDS, combined hormonal contraceptives, progestin, GnRH agonist, Danazol) -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or with out estrogen and progestin does not exceed 24 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 18 months reauthorization
Other Criteria	For reauthorization, Orilissa is continued to be used for pain associated with endometriosis -AND- attestation of reduction in pain -AND- For women of childbearing age, attestation of not pregnant -AND- Treatment with a gonadotropin-releasing hormone receptor antagonist with or without estrogen and progestin does not exceed 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orkambi

Products Affected

- **ORKAMBI ORAL GRANULES IN PACKET**
- **ORKAMBI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis and homozygous F508del mutation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orladeyo

Products Affected

- ORLADEYO

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Orserdu

Products Affected

- **ORSERDU ORAL TABLET 345 MG, 86 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) ER mutation status, HER2 mutation status, and ESR1 mutation status 2) Alternatives tried/failed
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Osmolex

Products Affected

- **OSMOLEX ER ORAL TABLET, IR - ER, BIPHASIC 24HR 129 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease or drug-induced extrapyramidal symptoms -AND- trial and failure, contraindication, or intolerance to immediate-release amantadine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ospheña

Products Affected

- OSPHENA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Otezla

Products Affected

- OTEZLA ORAL TABLET 30 MG
- OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For plaque psoriasis, all of the following (1-2): 1) if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg and member has moderate-to-severe disease -AND- 2) inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For oral ulcers associated with Behcet's Disease, inadequate response or intolerance to one systemic therapy colchicine for prevention of recurrent oral ulcers
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Oxbryta

Products Affected

- **OXBRYTA ORAL TABLET 300 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of sickle cell disease -AND- Failure, intolerance or contraindication to hydroxyurea
Age Restrictions	Deny if less than 4 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement of sickle cell disease signs, symptoms or complications
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Oxbryta for Suspension

Products Affected

- **OXBRYTA ORAL TABLET FOR SUSPENSION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of sickle cell disease -AND- Failure, intolerance or contraindication to hydroxyurea -AND- one of the following (1 or 2): 1) is eleven years of age or younger -OR- 2) has an inability to swallow tablets
Age Restrictions	Deny if less than 4 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement of sickle cell disease signs, symptoms or complications
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Oxervate

Products Affected

- **OXERVATE**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 8 weeks per eye
Required Medical Information	Documentation of diagnosis -AND- affected eye (e.g. right eye, both eyes).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	8 weeks
Other Criteria	Coverage beyond 8 weeks per eye will not be approved
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Palynziq

Products Affected

- **PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of phenylketonuria. Member meets the following criteria 1.) Baseline Phe level greater than 600 micromoles/L -AND- 2.) Failure or intolerance to existing management (i.e. Kuvan therapy) -AND- 3.) Has a prescription for epinephrine agent unless contraindicated.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in baseline pretreatment Phe levels -OR- blood Phe levels are less than or equal to 600 micromoles/L -OR- attestation that additional therapy with Palynziq is needed to allow adequate trial of maximum dose of 60mg per day for 16 weeks or member is not currently on 60mg per day dose.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Panretin

Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions in patients with AIDS-related Kaposi Sarcoma (KS) who are not receiving systemic therapy for KS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pemazyre

Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) FGFR2 fusion or other rearrangement as detected by an FDA-approved test 2) Attestation of previous treatment in alignment with FDA indication
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pheburane

Products Affected

- PHEBURANE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure or intolerance to generic sodium phenylbutyrate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Phenoxybenzamine

Products Affected

- **DIBENZYLINE**
- *phenoxybenzamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of excessive sweating and hypertension associated with pheochromocytoma supported by one of the following (1. or 2.): 1. Elevated metanephrines in plasma or urine. 2. Tumor evidence from CT scan or MRI
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Phospholine Iodide

Products Affected

- **PHOSPHOLINE IODIDE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For diagnosis of elevated intraocular pressure (IOP), the following criteria apply (1 and 2): therapeutic failure or intolerance to generic latanoprost, 2) therapeutic failure, contraindication, or intolerance to one generic ophthalmic alternative that lowers IOP (a. through f.): a) prostaglandin analog, b) ophthalmic beta-blocker, c) alpha-adrenergic agonist, d) carbonic anhydrase inhibitor, e) ophthalmic cholinergic agonist, f) combination products of these classes
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Piqray

Products Affected

- **PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR-positive, HER2-negative tumor status, 2) PIK3CA mutation positive as detected by an FDA-approved test, 3) disease progression on or after an endocrine-based regimen, 4) Concomitant therapy with fulvestrant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pomalyst

Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma, and combination use with dexamethasone, and previous trial of at least 2 therapies including lenalidomide and a proteasome inhibitor, and disease progression on or within 60 days of completion of the last therapy -OR- Documentation of AIDS-related Kaposi sarcoma (KS) after failure of highly active antiretroviral therapy (HAART) or in patients with KS who are HIV negative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ponvory

Products Affected

- **PONVORY**
- **PONVORY 14-DAY STARTER PACK**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Posaconazole Powder Packet

Products Affected

- NOXAFIL ORAL SUSP, DELAYED
RELEASE FOR RECON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 2 years of age or greater than 17 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Posaconazole Suspension

Products Affected

- **NOXAFIL ORAL SUSPENSION**
- *posaconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight less than or equal to 40 kg. For oropharyngeal candidiasis treatment, trial/failure or intolerance to generic fluconazole or generic itraconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of failure on generic posaconazole delayed-release tablets or an inability to swallow tablets is required when being utilized for the same FDA approved indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Posaconazole Tablet

Products Affected

- **NOXAFIL ORAL TABLET,DELAYED RELEASE (DR/EC)**
- *posaconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight greater than 40 kg. For invasive aspergillosis infection, trial/failure or contraindication to voriconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Noxafil delayed-release tablets, documentation of failure on generic posaconazole delayed-release tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Praluent

Products Affected

- PRALUENT PEN

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
Required Medical Information	<p>1.HoFH supported by genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9, or LDLRAP1 gene OR untreated LDL-C greater than 400mg/dL or TC greater than 500mg/dl with cutaneous or tendon xanthoma before age 10 yrs or HeFH in both parents AND The member has a current LDL-C of greater than 135 mg/dL (if 17 years of age or younger) or greater than 100mg/dL (18 years of age or older) despite use of maximally tolerated statin or statin intolerance AND The member will continue to receive concurrent lipid-lowering therapies for the treatment of HoFH. 2.HeFH supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool AND LDL-C greater than 100 mg/dL if 18 and older or LDL-C greater than 130 if 17 and younger despite use of maximally tolerated statin or statin intolerance. If 17 and younger will continue to receive concurrent lipid-lowering therapies. 3.Hypercholesterolemia ASCVD or Primary Hyperlipidemia AND LDL-C greater than 70 mg/dL despite use of maximally tolerated statin or statin intolerance</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For HoFH and Hypercholesterolemia with ASCVD, patients must have a trial/failure or contraindication to the preferred product Repatha . For HeFH and 10 years and older, patients must have a trial/failure or contraindication to the preferred product Repatha. For reauthorization, documentation showing an LDL-C reduction on Praluent therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prenatal Vitamins

Products Affected

- **PRENATAL VITAMIN PLUS LOW IRON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of nutritional supplementation required in a female of child-bearing potential during pre-conception, pregnancy, or lactation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prescription Drug Combo

Products Affected

- *acetaminophen-caff-dihydrocod*
- *acetaminophen-codeine oral solution 120-12 mg/5 ml*
- *acetaminophen-codeine oral tablet*
- **ALPRAZOLAM INTENSOL**
- *alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg, 2 mg, 3 mg*
- *alprazolam oral tablet,disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- **AMBIEN**
- **AMBIEN CR**
- **ASCOMP WITH CODEINE**
- **ATIVAN ORAL TABLET 0.5 MG, 1 MG, 2 MG**
- **BELBUCA**
- *buprenorphine*
- *butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg, 50-325-40-30 mg*
- **BUTRANS**
- *chlordiazepoxide hcl*
- *clonazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- *clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg*
- *codeine sulfate*
- *codeine-butalbital-asa-caff*
- **CONZIP**
- **DEMEROL (PF) INJECTION SYRINGE 25 MG/ML**
- **DEMEROL INJECTION**
- **DIAZEPAM INTENSOL**
- *diazepam oral solution 5 mg/5 ml (1 mg/ml)*
- *diazepam oral tablet*
- **DILAUDID ORAL LIQUID**
- **DILAUDID ORAL TABLET**
- **EDLUAR**
- **ENDOCET**
- *estazolam*
- *eszopiclone*
- *fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 37.5 mcg/hour, 50 mcg/hr, 62.5 mcg/hour, 75 mcg/hr, 87.5 mcg/hour*
- **FIORICET WITH CODEINE**
- *flurazepam*
- **HALCION ORAL TABLET 0.25 MG**
- *hydrocodone bitartrate oral capsule, oral only, er 12hr*
- *hydrocodone bitartrate oral tablet,oral only,ext.rel.24 hr*
- *hydrocodone-acetaminophen oral solution 7.5-325 mg/15 ml*
- *hydrocodone-acetaminophen oral tablet 10-300 mg, 10-325 mg, 5-300 mg, 5-325 mg, 7.5-300 mg, 7.5-325 mg*
- *hydrocodone-ibuprofen*
- *hydromorphone (pf) injection solution 10 (mg/ml) (5 ml), 10 mg/ml*
- *hydromorphone oral liquid*
- *hydromorphone oral tablet*
- *hydromorphone oral tablet extended release 24 hr 12 mg, 16 mg, 32 mg, 8 mg*
- **HYSINGLA ER**
- **KLONOPIN ORAL TABLET 0.5 MG, 1 MG, 2 MG**
- *levorphanol tartrate*
- **LORAZEPAM INTENSOL**
- *lorazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- **LOREEV XR ORAL CAPSULE,EXTENDED RELEASE 24HR 1 MG, 1.5 MG, 2 MG, 3 MG**
- **LUNESTA**
- *meperidine (pf) injection solution 100 mg/ml, 25 mg/ml, 50 mg/ml*
- *meperidine oral solution*
- *meperidine oral tablet 50 mg*
- *methadone oral solution 10 mg/5 ml, 5 mg/5 ml*
- *methadone oral tablet 10 mg, 5 mg*
- *morphine concentrate oral solution*
- *morphine oral capsule, er multiphase 24 hr 120 mg, 30 mg, 45 mg, 60 mg, 75 mg, 90 mg*
- *morphine oral capsule,extend.release pellets*

- 10 mg, 100 mg, 20 mg, 30 mg, 50 mg, 60 mg, 80 mg
- *morphine oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)*
- *morphine oral tablet*
- *morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg*
- **MS CONTIN ORAL TABLET EXTENDED RELEASE 100 MG, 15 MG, 200 MG, 30 MG, 60 MG**
- **NALOCET**
- **NUCYNTA**
- **NUCYNTA ER**
- *oxazepam*
- *oxycodone oral capsule*
- *oxycodone oral concentrate*
- *oxycodone oral solution*
- *oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg*
- *oxycodone oral tablet, oral only, ext. rel. 12 hr 10 mg, 20 mg*
- *oxycodone-acetaminophen oral solution 5-325 mg/5 ml*
- *oxycodone-acetaminophen oral tablet 10-300 mg, 10-325 mg, 2.5-325 mg, 5-300 mg, 5-325 mg, 7.5-300 mg, 7.5-325 mg*
- **OXYCONTIN ORAL TABLET, ORAL ONLY, EXT. REL. 12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG**
- *oxymorphone oral tablet*
- *oxymorphone oral tablet extended release 12 hr 10 mg, 15 mg, 20 mg, 30 mg, 40 mg, 5 mg, 7.5 mg*
- **PERCOCET**
- **PROLATE ORAL SOLUTION**
- **PROLATE ORAL TABLET**
- **QDOLO**
- **RESTORIL**
- **ROXICODONE ORAL TABLET 15 MG, 30 MG**
- **ROXYBOND**
- **SEGLENTIS**
- *temazepam*
- *tramadol oral capsule, er biphasic 24 hr 17-83*
- *tramadol oral capsule, er biphasic 24 hr 25-75 100 mg, 200 mg*
- *tramadol oral solution*
- *tramadol oral tablet 100 mg, 25 mg, 50 mg*
- *tramadol oral tablet extended release 24 hr*
- *tramadol oral tablet, er multiphasic 24 hr*
- *tramadol-acetaminophen*
- **TREZIX**
- *triazolam*
- **VALIUM**
- **XANAX ORAL TABLET 0.25 MG, 0.5 MG, 1 MG, 2 MG**
- **XANAX XR ORAL TABLET EXTENDED RELEASE 24 HR 0.5 MG, 1 MG, 2 MG, 3 MG**
- **XTAMPZA ER**
- *zaleplon oral capsule 10 mg, 5 mg*
- *zolpidem*

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	<p>For concomitant use of an opiate agonist and substance abuse therapy, documentation that the member has an acute pain condition (e.g. acute traumatic injury) in which treatment with other agents would cause insufficient pain control or if the member requires treatment for pain related to a terminal illness. For concomitant use of an opiate agonist, benzodiazepine, and a centrally acting skeletal muscle relaxant, documentation that the member has tried/failed at least 2 other skeletal muscle relaxants (e.g. methocarbamol, metaxalone), understanding these skeletal muscle relaxants are high-risk medications in geriatric patients AND attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For concomitant use of an opiate agonist and other opiate potentiators (e.g. gabapentinoids, sedative-hypnotics) attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For long acting (e.g. extended release) opioid medications, the following apply (1-5). 1)Pain is severe enough to require daily, around-the-clock, long-term opioid treatment. 2)Patient is not opioid naive. 3)Attestation that non-opiate alternative therapies have been explored (e.g. NSAIDs). 4)Attestation that controlled substance Rx history has been reviewed in the state Prescription Drug Monitoring Program. 5)Attestation of counseling on the potential adverse effects of opioid analgesics, including the risk of misuse, abuse, and addiction.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Opiate tx for pain+subs. abuse, app. opiate x 1mo. All other combos and dx approve x 12mo.
Other Criteria	<p>Opiate agonists will receive automatic approval if no recent claims for a substance abuse therapy (e.g. buprenorphine-naloxone) OR a benzodiazepine (e.g. triazolam, alprazolam) OR a benzodiazepine with a centrally acting skeletal muscle relaxant (e.g., carisoprodol) OR a gabapentinoid OR a sedative-hypnotic. Benzodiazepines (e.g. triazolam, alprazolam) will receive automatic approval if no recent claims for an opiate agonist (e.g. oxycodone, hydrocodone, oxymorphone) or an opiate agonist with a centrally acting skeletal muscle relaxant (e.g. carisoprodol). Sedative-hypnotics (e.g. zolpidem) will receive automatic approval if no recent claims for an opiate agonist. Infusible opiate agonists will be covered under Part B when administered via infusion pump.</p>
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Pretomanid

Products Affected

- *pretomanid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of extensively drug resistant, treatment intolerant or nonresponsive multidrug resistant tuberculosis -AND- Used as part of a combination regimen with bedaquiline and linezolid -AND- Therapeutic failure, contraindication or intolerance to both of the following (1 and 2): 1) A fluoroquinolone antibiotic or amikacin, kancamycin, or capreomycin 2) isoniazid, rifampin, rifabutin, or rifapentine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	26 weeks
Other Criteria	For reauthorization, additional therapy required due to doses of the regimen being missed for safety reasons
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Procysbi

Products Affected

- **PROCYSBI ORAL GRANULES DEL
RELEASE IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- previous trial and failure, intolerance, or contraindication to Cystagon (cysteamine bitartrate immediate-release)
Age Restrictions	Deny if less than 1 year of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prolia

Products Affected

- PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -And- For osteoporosis at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 (i.e. osteopenia) -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For osteoporosis and osteopenia, documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Provigil

Products Affected

- *modafinil*
- **PROVIGIL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications. Diagnosis of fatigue associated with Multiple Sclerosis (MS)</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required. For brand Provigil, documentation of failure on generic modafinil.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Fatigue associated with Multiple Sclerosis (MS)
Part B Prerequisite	No

Pulmonary Arterial Hypertension

Products Affected

- **ADCIRCA**
- **ADEMPAS**
- **ALYQ**
- *ambrisentan*
- *bosentan*
- **LETAIRIS**
- **LIQREV**
- **OPSUMIT**
- **OPSYNVI**
- **ORENITRAM MONTH 1 TITRATION KT**
- **ORENITRAM MONTH 2 TITRATION KT**
- **ORENITRAM MONTH 3 TITRATION KT**
- **ORENITRAM ORAL TABLET EXTENDED RELEASE 0.125 MG, 0.25 MG, 1 MG, 2.5 MG, 5 MG**
- **REVATIO ORAL SUSPENSION FOR RECONSTITUTION**
- **REVATIO ORAL TABLET**
- *sildenafil (pulm.hypertension) oral suspension for reconstitution*
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*
- **TADLIQ**
- **TRACLEER ORAL TABLET**
- **TRACLEER ORAL TABLET FOR SUSPENSION**
- **TYVASO DPI INHALATION CARTRIDGE WITH INHALER 16 MCG, 16(112)-32(112) -48(28) MCG, 32 MCG, 32-48 MCG, 48 MCG, 64 MCG**
- **UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG**
- **UPTRAVI ORAL TABLETS,DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group. For bosentan and sildenafil (i.e. Revatio) in pediatric individuals, an exception to RHC will be allowed when the risk of RHC outweighs the benefit -AND- prescriber attests alternative studies have been completed (i.e. CT, MRI or specified test ruling out other causes of pulmonary hypertension). For Adempas, additional diagnosis of CTEPH as documented by right heart catheterization and V/Q scan substantiating mPAP greater than 20 mmHg at rest and PWP less than or equal to 15 mmHg and documented presence of occlusive thrombi within the pulmonary arteries will be approved.</p>
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Adcirca, trial and failure of generic tadalafil or Alyq is required. For brand Letairis, trial and failure of generic ambrisentan is required. For brand Revatio, trial and failure of generic sildenafil is required. For brand Tracleer 62.5mg and 125mg, trial and failure of generic bosentan is required. For brand Tadliq, trial and failure of generic tadalafil or Alyq - OR- Inability to swallow tablets is required. For Liquev, trial and failure of generic sildenafil for oral suspension is required. For Opsynvi, therapeutic failure, contraindication, or intolerance to a generic endothelin-1 receptor antagonist, phosphodiesterase type 5 inhibitor and soluble guanylate cyclase stimulator.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pulmozyme

Products Affected

- PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis -AND- Used in conjunction with standard therapies for management of cystic fibrosis to improve pulmonary function.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization, attestation of increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Purified Cortrophin Gel

Products Affected

- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Covered for the following indications: 1. Acute exacerbations of multiple sclerosis (MS) for patients receiving concurrent immunomodulator therapy 2. Rheumatic disorders for patients receiving maintenance therapy 3. Collagen diseases for members receiving maintenance therapy 4. Dermatologic diseases, if using for severe psoriasis, then the member is concurrently receiving maintenance therapy 5. Allergic states (i.e. serum sickness and transfusion reaction due to serum protein reaction), if using for atopic dermatitis, then the member is concurrently receiving maintenance therapy 6. Ophthalmic diseases 7. Respiratory diseases 8. Gout and unable to take first-line therapies. 9. Pediatric acquired epileptic aphasia. 10. Proteinuria in nephrotic syndrome and trial/failure or contraindication to two therapies from any of the following different classes: corticosteroids (e.g., cortisone or dexamethasone), calcineurin inhibitors (e.g., cyclosporine or tacrolimus, per DRUGDEX). For covered indications 1 through 9, limited/unsatisfactory response or intolerance (i.e. severe anaphylaxis) to two corticosteroids (i.e. IV methylprednisolone, IV dexamethasone, or high dose oral steroids) must be documented.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month

PA Criteria	Criteria Details
Other Criteria	For reauthorization, the following (1. 2. and 3.) must be met. 1) Prescriber attestation that the member cannot use corticosteroids (e.g. IV methylprednisolone, high dose oral corticosteroids) due to unsatisfactory response, intolerance (e.g. severe anaphylaxis) or experienced a severe adverse event to corticosteroids (e.g. psychosis). 2) If the reauthorization is for the treatment of multiple sclerosis, a rheumatic disorder, dermatologic disease, or nephrotic syndrome, the prescriber attests that Cortrophin is being used for a new acute exacerbation and not on a routine basis to prevent an exacerbation as supported by Compendia. 3) If the reauthorization is for treatment of multiple sclerosis, a rheumatic disorder, or collagen disease, the member continues to receive maintenance therapy.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Epileptic Aphasia
Part B Prerequisite	No

Pyrukynd

Products Affected

- **PYRUKYND ORAL TABLET 20 MG, 5 MG, 5 MG (4-WEEK PACK), 50 MG**
- **PYRUKYND ORAL TABLETS,DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hemolytic anemia with pyruvate kinase deficiency, as supported by one of the following (1-2): 1) mutations in the PKLR gene, 2) reduced activity of the pyruvate kinase enzyme.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	24 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of increase in hemoglobin level from baseline -OR- decrease in transfusion burden from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Qelbree

Products Affected

- **QELBREE ORAL
CAPSULE,EXTENDED RELEASE 24HR
100 MG, 150 MG, 200 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant and generic atomoxetine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Qinlock

Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced gastrointestinal stromal tumor -AND- Prior treatment with imatinib and 2 other kinase inhibitors.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Quinine

Products Affected

- **QUALAQUIN**
- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment or prevention of leg cramps
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	Doses for duration greater than 10 days will not be approved
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Qulipta

Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, of which 8 or more are migraine days. The following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Radicava ORS

Products Affected

- **RADICAVA ORS STARTER KIT SUSP**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of amyotrophic lateral sclerosis (ALS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ravicti

Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	Urea cycle disorders due to N-acetylglutamate synthetase deficiency, Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of a urea cycle disorders (UCDs)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Recorlev

Products Affected

- **RECORLEV**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endogenous Cushing's syndrome AND patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of mean urine free cortisol (mUFC) less than starting baseline value.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Regranex

Products Affected

- **REGRANEX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of lower-extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond and have an adequate blood supply -AND- being used as an adjunct to standard ulcer care practices (e.g. sharp debridement, non-weight bearing regimen, infection control) -AND- attestation of a wound care plan.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	20 weeks
Other Criteria	For reauthorization, one of the following (1-2): 1) documentation of decrease in ulcer(s) size without complete ulcer(s) closure -OR- 2) documentation of new lower extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond with an adequate blood supply, attestation of being used as an adjunct to standard ulcer care practices, and attestation of a wound care plan.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Relistor

Products Affected

- **RELISTOR ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-2): 1) chronic non-cancer pain, or 2) chronic pain related to prior cancer or its treatment who do not require frequent (e.g. weekly) opioid dosage escalation -AND- documentation of opioid medication use for at least one month -AND- trial and failure, contraindication, or intolerance to 2 of the following (3-5): 3) Laxatives, 4) lubiprostone, 5) Movantik
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Relistor SC

Products Affected

- **RELISTOR SUBCUTANEOUS SOLUTION**
- **RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-3): 1) chronic non-cancer pain, 2) advanced illness or active cancer in palliative care or 3) chronic pain related to prior cancer or its treatments who do not require frequent (e.g., weekly) opioid dosage escalation -AND- documentation of opioid medication use for at least one month -AND- trial and failure, contraindication, or intolerance to 2 of the following (4 to 6): 4) Laxatives 5) lubiprostone 6) Movantik.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Repatha

Products Affected

- **REPATHA PUSHTRONEX**
- **REPATHA SURECLICK**
- **REPATHA SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
Required Medical Information	<p>1.HoFH supported by genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9, or LDLRAP1 gene OR untreated LDL-C greater than 400mg/dL or TC greater than 500mg/dl with cutaneous or tendon xanthoma before age 10 yrs or HeFH in both parents AND The member has a current LDL-C of greater than 135 mg/dL (if 17 years of age or younger) or greater than 100mg/dL (18 years of age or older) despite use of maximally tolerated statin or statin intolerance AND The member will continue to receive concurrent lipid-lowering therapies for the treatment of HoFH. 2.HeFH supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool, AND LDL-C greater than 100 mg/dL if 18 and older or LDL-C greater than 130 if 17 and younger despite use of maximally tolerated statin or statin intolerance. If 17 and younger will continue to receive concurrent lipid-lowering therapies. 3.Hypercholesterolemia ASCVD or Primary Hyperlipidemia AND LDL-C greater than 70 mg/dL despite use of maximally tolerated statin or statin intolerance</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For reauthorization, documentation showing an LDL-C reduction on Repatha therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Retevmo

Products Affected

- **RETEVMO ORAL CAPSULE 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restrictions	Pending CMS Review
Prescriber Restrictions	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Indications	Pending CMS Review
Off Label Uses	Pending CMS Review
Part B Prerequisite	No

Revlimid

Products Affected

- *lenalidomide*
- **REVLIMID**

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of chronic lymphocytic leukemia outside of a controlled clinical trial
Required Medical Information	Diagnosis of multiple myeloma in combination with dexamethasone -OR- diagnosis of multiple myeloma, as maintenance following autologous hematopoietic stem cell transplant (auto-HSCT) -OR- diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities -OR- diagnosis of mantle cell lymphoma (MCL) in which disease has relapsed or progressed after two prior therapies, one of which included bortezomib -OR- diagnosis of follicular lymphoma in combination with a rituximab product -OR- diagnosis of marginal zone lymphoma in combination with a rituximab product after previous treatment.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Rezdiffra

Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in conjunction with diet and exercise. Diagnosis of NASH confirmed by a liver biopsy or non-invasive tests (NITs) performed within the previous 6 months indicating F2 or F3 fibrosis - AND- member does not have any evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of continued use in conjunction with diet and exercise -AND- member has experienced stabilization of fibrosis as demonstrated by NIT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rezlidhia

Products Affected

- **REZLIDHIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rezurock

Products Affected

- **REZUROCK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic graft-versus-host disease (cGVHD) -AND- therapeutic failure or intolerance to 2 lines of systemic therapy
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Riluzole

Products Affected

- **EXSERVAN**
- **TEGLUTIK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of amyotrophic lateral sclerosis (ALS) -AND- Inability to swallow tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rinvoq

Products Affected

- **RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate) or all non-biologic DMARDs are contraindicated. For moderate to severe refractory atopic dermatitis whose disease is not adequately controlled with other systemic drug products, documentation of one of the following (1 or 2): 1) trial & failure, or intolerance to at least one topical corticosteroid -OR- topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) 2) The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For Non-radiographic Axial Spondyloarthritis, trial & failure or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or contraindication to all. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For diagnoses in which tumor necrosis factor (TNF) blockers are also indicated (e.g., Rheumatoid Arthritis, Psoriatic Arthritis), the member has experienced therapeutic failure or intolerance to at least 1 TNF blocker.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rivfloza

Products Affected

- **RIVFLOZA SUBCUTANEOUS SOLUTION**
- **RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3): 1) genetically confirmed diagnosis of primary hyperoxaluria type 1 (PH1), 2) relatively preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m ²), and 3) at least two elevated urinary oxalate levels greater than 1.5 times the upper reference limit.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, continued preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m ²) -AND- a reduction in urinary oxalate levels from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rozlytrek

Products Affected

- **ROZLYTREK ORAL CAPSULE 100 MG, 200 MG**
- **ROZLYTREK ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, the tumor status is ROS1-positive. For solid tumors with NTRK gene fusion without a known acquired resistance mutation, the tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or the tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rubraca

Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following. 1) BRCA mutations, if applicable to diagnosis. 2) Alternatives tried/failed. 3) Concomitant therapy, if applicable to diagnosis.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ruconest

Products Affected

- RUCONEST

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	For the treatment of attacks of hereditary angioedema (HAE) type I & II with the following (1-4): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. 4) Documentation of member's weight. For the treatment of attacks of hereditary angioedema (HAE) type III with the following (5-9): 5) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 6) Documentation of family history of HAE or FXII mutation 7) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 8) Medications known to cause angioedema have been evaluated and discontinued. 9) Documentation of member's weight.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For 18 years of age or older, therapeutic failure, intolerance or contraindication to icatibant.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Rydapt

Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Use as single agent induction therapy for AML
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) FLT3 mutations 2) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sabril

Products Affected

- **SABRIL**
- *vigabatrin*
- **VIGADRONE**
- **VIGPODER**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of refractory complex partial seizures -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance to at least two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, tiagabine) -OR- documentation of use as monotherapy in treatment of infantile spasms
Age Restrictions	Deny if less than 2 years of age in treatment of refractory complex partial seizures -OR- if less than 1 month old and greater than 2 years of age in treatment of infantile spasms
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Sabril, trial and failure of generic vigabatrin is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Samsca

Products Affected

- SAMSCA
- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic hypervolemic or euvoletic hyponatremia evidenced by (1. or 2.): 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	Doses must be initiated in the hospital setting to closely monitor serum sodium. Treatment should be limited to 30 days to minimize risk of liver injury. For brand Samsca, initial authorization requires a trial and failure of generic tolvaptan and reauthorization requires failure of generic tolvaptan if not previously trialed. For reauthorization, treatment is for a new episode of a clinically significant euvoletic or hypervolemic hyponatremia -AND- on of the following (1. or 2.) 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Saphris

Products Affected

- *asenapine maleate*
- **SAPHRIS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Brand Saphris, trial and failure of generic asenapine sublingual tablets
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Savella

Products Affected

- SAVELLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Scemblix

Products Affected

- **SCSEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restrictions	Pending CMS Review
Prescriber Restrictions	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Indications	Pending CMS Review
Off Label Uses	Pending CMS Review
Part B Prerequisite	No

Secuado

Products Affected

- **SECUADO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure of generic asenapine sublingual tablets -AND- trial/failure or intolerance to 1 of the following or all are contraindicated (1-3): 1) olanzapine, 2) quetiapine, 3) risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sertraline Capsule

Products Affected

- *sertraline oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder or obsessive-compulsive disorder -AND- Sertraline 100mg or sertraline 125mg has been received for greater than or equal to 7 days -AND- Therapeutic failure or intolerance to generic sertraline immediate release tablets -AND- Therapeutic failure, intolerance or contraindication to at least one other antidepressant (e.g. SNRI, SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Seysara

Products Affected

- SEYSARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of inflammatory lesions of non-nodular moderate to severe acne -AND- trial and failure or intolerance of at least 1 generic topical acne medication (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) -AND- trial and failure or intolerance of at least 1 generic oral acne antibiotic (e.g. minocycline, azithromycin, trimethoprim)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Signifor

Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Cushing's disease AND patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in urinary free cortisol levels from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Siliq

Products Affected

- SILIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For psoriasis, patients must have therapeutic failure or intolerance to 2 preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Taltz, Otezla, Stelara SC, Enbrel, and Skyrizi SC. For psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Simponi

Products Affected

- **SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML, 50 MG/0.5 ML**
- **SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML, 50 MG/0.5 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Simponi 50mg: For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide) and Simponi will be used in combination with methotrexate. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). Diagnosis of psoriatic arthritis Simponi 100mg: Diagnosis of moderate to severe ulcerative colitis.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	<p>For Rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Enbrel, Actemra SC, Xeljanz/Xeljanz XR and Rinvoq. For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Enbrel, Taltz, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Enbrel, Taltz, Rinvoq, and Xeljanz/Xeljanz XR. For ulcerative colitis, patients must have therapeutic failure or intolerance to the preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Stelara SC, Rinvoq, and Xeljanz/Xeljanz XR. For ulcerative colitis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sirturo

Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of extensively drug resistant tuberculosis, treatment-intolerant tuberculosis, or nonresponsive multidrug-resistant tuberculosis - AND- Therapeutic failure, contraindication, or intolerance to 1 of the following: isoniazid, rifampin, rifabutin, or rifapentine -AND- Used in combination with at least 3 other drugs for tuberculosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 weeks
Other Criteria	For reauthorization, additional therapy required due to doses of the regimen being missed for safety reasons
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skyclarys

Products Affected

- SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Friedreichs ataxia confirmed by genetic testing (i.e., FXN gene mutation).
Age Restrictions	Deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skyrizi

Products Affected

- **SKYRIZI SUBCUTANEOUS PEN INJECTOR**
- **SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML**
- **SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohns disease, attestation of clinical response or remission following IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC. For moderate to severe ulcerative colitis, attestation of clinical response or remission following IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skytrofa

Products Affected

- SKYTROFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, bone age, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sogroya

Products Affected

- SOGROYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, bone age, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sohonos

Products Affected

- SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 2.5 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	Deny if female and less than 8 years of age -OR- if male and less than 10 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in volume of new heterotopic ossification from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Solaraze

Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure, intolerance, or contraindication to one of the following (1 or 2): 1) topical fluorouracil solution or fluorouracil 5% cream 2) topical imiquimod 5% cream
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of 30 day washout period since optimal therapeutic effect may not be evident until 30 days following cessation of therapy AND attestation of previous response to diclofenac sodium 3% topical gel therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Somavert

Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- inadequate or partial response to surgery or radiotherapy or not a candidate for surgery or radiotherapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sotyktu

Products Affected

- SOTYKTU

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Cosentyx, Taltz, Otezla, Stelara, Enbrel and Skyrizi.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sovaldi

Products Affected

- SOVALDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for all regimens recommended by the AASLD/IDSA guidelines containing the following agents: ledipasvir/sofosbuvir, sofosbuvir/velpatasvir, glecaprevir/pibrentasvir.
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Spevigo SC

Products Affected

- **SPEVIGO SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized pustular psoriasis (GPP) -AND- Previous GPP flares have been experienced -AND- A flare is not currently being experienced -AND- Prevention for future GPP flares is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sprycel

Products Affected

- **SPRYCEL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) Philadelphia Chromosome status (e.g. positive) 2) Alternatives tried/failed 3) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Statin Liquid

Products Affected

- ATORVALIQ
- FLOLIPID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Inability to swallow tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Stelara

Products Affected

- **STELARA SUBCUTANEOUS SOLUTION**
- **STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease, attestation of clinical response or remission following single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC. For Ulcerative colitis, attestation of clinical response or remission following single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC
Age Restrictions	Deny if less than 18 years of age for Crohn's Disease and Ulcerative Colitis or less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Must follow recommended dosing guidelines based upon weight. Induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Stivarga

Products Affected

- **STIVARGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer and trial of a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy, AND an anti-VEGF therapy AND if RAS wild-type, an anti-EGFR therapy -OR- documentation of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with both imatinib and sunitinib -OR- documentation of hepatocellular cancer AND previous treatment with sorafenib
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sucraid

Products Affected

- SUCRAID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of congenital-sucrase-isomaltase deficiency (CSID) supported by one of the following (1 or 2): 1) small bowel biopsy with disaccharidase assay showing absent or reduced sucrase activity, reduced or normal isomaltase activity, reduced maltase activity, and reduced or normal lactase activity. 2) sucrase deficiency evidenced by sucrose breath hydrogen test or carbon-sucrose breath test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sunosi

Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	<p>For narcolepsy and OSAHS, documentation of trial and failure, contraindication or intolerance to modafinil and armodafinil. For reauthorization, provider attestation of improvement in daytime sleepiness is required.</p>
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Sutent

Products Affected

- *sunitinib malate*
- SUTENT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) disease progression on or intolerance to imatinib mesylate 2) high risk of recurrent RCC following nephrectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Sutent, documentation of trial and failure of generic sunitinib is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Symdeko

Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have either the homozygous F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to tezacaftor/ivacaftor based on clinical and/or in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sympazan

Products Affected

- SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy -AND- unable to tolerate generic clobazam
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Symproic

Products Affected

- SYMPROIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-2): 1) chronic non-cancer pain, or 2) chronic pain related to prior cancer or its treatments who do not require frequent (e.g., weekly) opioid dosage escalation -AND- documentation of opioid medication use -AND- trial and failure, contraindication, or intolerance to at least 2 of the following (3-5): 3) Laxatives, 4) lubiprostone, 5) Movantik
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Synarel

Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For central precocious puberty (CPP), advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L. For female with endometriosis, attestation of not pregnant if of childbearing age -AND- Therapeutic failure, contraindication or intolerance to 2 of the following standard of care treatments: NSAIDs, combination hormonal contraceptive, progestin (i.e. medroxyprogesterone injection), GnRH agonist (i.e. Leuprolide) or danazol
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty. Deny if less than 18 years of age for endometriosis.
Prescriber Restrictions	
Coverage Duration	Endometriosis: 6 months, CPP: 6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization for CPP, attestation of pubertal development slowing from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Syndros

Products Affected

- SYNDROS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of one of the following (1 or 2): 1) anorexia associated with weight loss in patients with AIDS -and- trial and failure, contraindication, or intolerance to generic dronabinol capsules -OR- 2) nausea and vomiting associated with cancer chemotherapy in adults who have trial and failure, contraindication, or intolerance to a conventional antiemetic treatment (e.g., metoclopramide, promethazine, ondansetron, perphenazine, etc.) -and- trial and failure, contraindication, or intolerance to generic dronabinol capsules
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when the following are met: 1) used for chemotherapy-induced nausea and vomiting. 2) used as full replacement for IV anti-emetic therapy. 3) using within 48 hours of receiving chemotherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tabrecta

Products Affected

- **TABRECTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with MET exon 14 skipping mutation as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tafinlar for Suspension

Products Affected

- **TAFINLAR ORAL TABLET FOR SUSPENSION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- both of the following. 1) BRAF mutations, if applicable to diagnosis. 2) Alternatives tried/failed and concomitant therapy, if applicable to diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation of inability to swallow Tafinlar capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tagrisso

Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) and one of the following (1-4): 1) Adjuvant therapy after tumor resection -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, 2) Locally advanced disease -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations -AND- used in combination with pemetrexed and platinum-based chemotherapy, 3) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations -AND- used with or without combination therapy of pemetrexed and platinum-based chemotherapy, 4) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR T790M mutations -AND- has progressed on or after EGFR TKI therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Takhzyro

Products Affected

- **TAKHZYRO SUBCUTANEOUS SOLUTION (150 MG/ML)**
- **TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Taltz

Products Affected

- **TALTZ AUTOINJECTOR**
- **TALTZ SYRINGE SUBCUTANEOUS SYRINGE 80 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs.
Age Restrictions	Deny if less than 18 years of age for Psoriatic Arthritis, Ankylosing Spondylitis and non-radiographic axial spondyloarthritis or less than 6 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Talzenna

Products Affected

- **TALZENNA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of deleterious or suspected deleterious gBRCAm, HER2-negative locally advanced or metastatic breast cancer as a single agent - OR- Documentation of HRR gene-mutated metastatic castration-resistant prostate cancer in combination with enzalutamide
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Targretin

Products Affected

- *bexarotene oral*
- *bexarotene topical*
- **TARGRETIN ORAL**
- **TARGRETIN TOPICAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations of cutaneous T-cell lymphoma in patients who are refractory to at least one prior systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Targretin, documentation of trial and failure of generic bexarotene is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tarpeyo

Products Affected

- TARPEYO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one (1) of the following (A or B): A) Used in combination with one (1) of the following (1 or 2): 1) angiotensin-converting enzyme inhibitor (ACE-I), or 2) angiotensin II receptor blocker (ARB), -OR- B) Intolerance or contraindication to both of the following (3 and 4): 3) ACE-I, and 4) ARB.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 months
Other Criteria	For reauthorizations, documentation of diagnosis -AND- member requires re-continuation of therapy with Tarpeyo -AND- one of the following (A or B): A) Used in combination with one of the following (1 or 2): 1) angiotensin-converting enzyme inhibitor (ACE-I), or 2) angiotensin II receptor blocker (ARB), -OR- B) Intolerance or contraindication to both of the following (3 and 4): 3) ACE-I, and 4) ARB
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tascenso ODT

Products Affected

- TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Inability to swallow capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tasigna

Products Affected

- TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) Philadelphia Chromosome status (e.g. positive) 2) Alternatives tried/failed
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tasimelteon

Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tavalisse

Products Affected

- TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For diagnosis of ITP, the following criteria apply (1 and 2): 1) trial, intolerance, or inadequate response to a corticosteroid, immunoglobulin, or splenectomy. 2) One of the following (A or B): A) Platelet count less than or equal to $50 \times 10^9/L$ and has significant mucous member bleeding or at least one risk factor for bleeding (e.g. hypertension, peptic ulcer disease). B) Platelet count of less than or equal to $30 \times 10^9/L$.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tavneos

Products Affected

- TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe active granulomatosis with polyangiitis or severe active microscopic polyangiitis -AND- Prescriber attests to positive test for anti-PR3 or positive test for anti-MPO -AND- Member will continue to receive concomitant standard of care treatment with systemic glucocorticoids (e.g. prednisone) and immunosuppressives (e.g. mycophenolate mofetil, azathioprine).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of therapeutic response defined by disease stability or disease improvement -AND- Member will continue to receive concomitant standard of care treatment with glucocorticoids (e.g. prednisone) and immunosuppressives (e.g. mycophenolate mofetil, azathioprine).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tazarotene

Products Affected

- **ARAZLO**
- **FABIOR**
- *tazarotene topical foam*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tazorac

Products Affected

- *tazarotene topical cream*
- *tazarotene topical gel*
- **TAZORAC TOPICAL CREAM**
- **TAZORAC TOPICAL GEL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 of the following (A or B). A) Documentation of plaque psoriasis -AND- trial and failure or intolerance to at least one topical corticosteroid (e.g. fluocinonide, mometasone, triamcinolone, betamethasone). B) Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tazverik

Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic epithelioid sarcoma - AND- Disease is not eligible for complete resection. Documentation of relapsed or refractory follicular lymphoma -AND-Tumors are EZH2 mutation positive, as detected by FDA approved test, in a member that has received at least 2 prior systemic therapies -OR- Prescriber attests there are no satisfactory alternative treatment options.
Age Restrictions	For epithelioid sarcoma, deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tecfidera

Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg* **CAPSULE, DELAYED RELEASE(DR/EC) 120 MG, 120 MG (14)- 240 MG (46), 240 MG**
- **TECFIDERA ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	For brand Tecfidera, documentation of failure on generic dimethyl fumarate
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tegsedi

Products Affected

- TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of polyneuropathy associated with hereditary TTR (hATTR) amyloidosis with mutation in TTR gene confirmed by genetic testing -AND- Neurologic examination shows clinical signs and symptoms of the disease (e.g. peripheral/autonomic neuropathy, motor disability, carpal tunnel, etc.) -AND- Attestation of peripheral neuropathy impairment score (NIS) of 10 or greater -AND- Not simultaneously utilizing other gene targeted therapy for polyneuropathy of hATTR
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement of polyneuropathy from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tepmetko

Products Affected

- **TEPMETKO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with a MET exon 14 skipping alteration
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Testosterone (androgens)

Products Affected

- **ANDROGEL TRANSDERMAL GEL IN METERED-DOSE PUMP**
- **AVEED**
- **DEPO-TESTOSTERONE**
- **JATENZO ORAL CAPSULE 158 MG, 198 MG, 237 MG**
- **NATESTO**
- **TESTIM**
- *testosterone cypionate*
- *testosterone enanthate*
- *testosterone transdermal gel in metered-dose pump*
- *testosterone transdermal gel in packet*
- *testosterone transdermal solution in metered pump w/app*
- **TLANDO**
- **VOGELXO TRANSDERMAL GEL**
- **VOGELXO TRANSDERMAL GEL IN METERED-DOSE PUMP**
- **XYOSTED**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of primary or secondary hypogonadism in males with testicular failure due to cryptorchidism, bilateral torsions, orchitis, vanishing testis syndrome, orchidectomy, Klinefelter's syndrome, chemotherapy, radiation or toxic damage -OR- documentation of primary or secondary hypogonadism in males with multiple symptoms of hypogonadism including at least one of the following specific symptoms: height loss due to vertebral fractures, low trauma fractures, low bone density, incomplete or delayed sexual development, breast discomfort, loss of axillar and/or pubic body hair, hot flushes -OR- documentation of HIV infection in men with weight loss -OR- documentation of chronic steroid treatment in men. In all previously noted indications, members must also have documented low total testosterone level below the normal range for the laboratory -OR- a total testosterone level near the lower limit of the normal range with a low free testosterone level which is less than normal based upon the laboratory reference range -OR- the member is not producing any testosterone. Additional approvable indications include female patients with metastatic breast cancer (testosterone enanthate only), primary or secondary hypogonadism in males with testicular failure due to double orchidectomy, and delayed puberty in males (testosterone enanthate only).</p>
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	HIV Wasting
Part B Prerequisite	No

Thalomid

Products Affected

- **THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Use as monotherapy for ENL treatment in the presence of moderate to severe neuritis
Required Medical Information	Documentation of multiple myeloma in combination with dexamethasone -OR- documentation for use in the treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) -OR- documentation of therapy for prevention and suppression of the cutaneous manifestations of ENL recurrence
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Thiola

Products Affected

- **THIOLA**
- **THIOLA EC**
- *tiopronin oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Confirmation of cystinuria by at least one 24-hour urine collection with measurement of urinary cysteine levels greater than 400 mg/day -AND- Attestation of failure of urine alkalization with potassium citrate (to achieve pH of 7.0).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For brand Thiola or brand Thiola EC, trial and failure of generic tiopronin. For reauthorization, attestation of urine cystine concentration decreased from baseline -OR- decrease in production of cystine stones is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Thrombopoiesis Stimulating Agents

Products Affected

- ALVAIZ ORAL TABLET 18 MG, 36 MG, 54 MG, 9 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG
- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of chronic immune idiopathic thrombocytopenia purpura and trial and failure of corticosteroid or immunoglobulin therapy or splenectomy -OR- documentation of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy -OR- severe aplastic anemia who have had an insufficient response to immunosuppressive therapy -OR- For eltrombopag olamine only, documentation of first line treatment for severe aplastic anemia and used in combination with at least two immunosuppressive therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Platelet count to be provided. For Alvaiz (eltrombopag choline), therapeutic failure, contraindication, or intolerance to Promacta (eltrombopag olamine) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tibsovo

Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) IDH1 mutation status, 2) alternatives tried/failed, 3) comorbidity that precludes use of intensive induction chemotherapy (i.e. age greater than or equal to 75 years, severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) 4) Concomitant therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tigan

Products Affected

- *trimethobenzamide oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tolsura

Products Affected

- TOLSURA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Unable to tolerate generic itraconazole capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation the member is still unable to tolerate generic itraconazole capsules
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Topical Lidocaine

Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical ointment*
- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Transmucosal Fentanyl Citrate

Products Affected

- *fentanyl citrate buccal lozenge on a handle*
1,200 mcg, 1,600 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg
- *fentanyl citrate buccal tablet, effervescent*
100 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg
- *FENTORA BUCCAL TABLET, EFFERVESCENT 100 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of therapeutic use and the member is currently receiving and tolerant to long acting opioid therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tremfya

Products Affected

- TREMFYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For psoriasis, patients must have therapeutic failure or intolerance to 2 of the preferred products: Cosentyx, Taltz, a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Otezla, Stelara SC, Enbrel and Skyrizi SC. For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the preferred products: Cosentyx, Taltz, a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Otezla, Stelara SC, Enbrel, Rinvoq, Skyrizi SC, and Xeljanz/Xeljanz XR. For psoriasis and psoriatic arthritis induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tretinoin

Products Affected

- *adapalene topical cream*
- *adapalene topical gel 0.3 %*
- *adapalene topical swab*
- *adapalene-benzoyl peroxide*
- **AKLIEF**
- **ALTRENO**
- **ATRALIN**
- **CABTREO**
- *clindamycin-tretinoin*
- **DIFFERIN TOPICAL CREAM**
- **DIFFERIN TOPICAL GEL WITH PUMP**
- **DIFFERIN TOPICAL LOTION**
- **EPIDUO FORTE**
- **EPIDUO TOPICAL GEL WITH PUMP**
- **RETIN-A**
- **RETIN-A MICRO**
- **RETIN-A MICRO PUMP TOPICAL GEL WITH PUMP 0.06 %, 0.08 %**
- *tretinoin*
- *tretinoin microspheres topical gel*
- *tretinoin microspheres topical gel with pump 0.08 %*
- **TWYNEO**
- **VELTIN**
- **WINLEVI**
- **ZIANA**

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two generic topical non-retinoid acne medications (e.g. clindamycin, sulfacetamide, erythromycin)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Trientine

Products Affected

- *trientine oral capsule 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Wilson's Disease -AND- intolerant of penicillamine (e.g., generic penicillamine capsule or tablet).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Therapeutic failure or intolerance to generic trientine hydrochloride 250mg capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Trikafta

Products Affected

- **TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL**
- **TRIKAFTA ORAL TABLETS, SEQUENTIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have at least one F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to elexacaftor/tezacaftor/ivacaftor based on in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Truqap

Products Affected

- TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) HR mutation status, HER2 mutation status, and PIK3CA/AKT1/PTEN status 2) concomitant therapy with fulvestrant 3) disease progression on at least one endocrine-based regimen in the metastatic setting -OR- recurrence on or within 12 months of completing adjuvant therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tukysa

Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) HER2 mutations 2) RAS genetic status 3) Alternatives tried/failed 4) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Turalio

Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor associated with severe morbidity and functional limitations -AND- patient is not amenable to improvement with surgery or not a candidate for surgery
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tykerb

Products Affected

- *lapatinib*
- **TYKERB**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following, if applicable to diagnosis: 1) HER2 mutations 2) Alternatives tried/failed 3) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Tykerb, trial and failure of generic lapatinib is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tymlos

Products Affected

- TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 24 months.
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Additional documentation of trial/failure, intolerance or contraindication to preferred parathyroid hormone analog teriparatide. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Tymlos will be limited to a coverage duration of 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ubrelvy

Products Affected

- **UBRELVY ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of acute treatment of migraine with or without aura -AND- trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	Patients must have therapeutic failure, intolerance, or contraindication to Nurtec ODT. For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Uloric

Products Affected

- *febuxostat*
- **ULORIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic management of hyperuricemia due to gout -And- trial/failure, intolerance or contraindication to allopurinol.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Valchlor

Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received at least one prior skin-directed therapy (e.g. topical corticosteroids, topical chemotherapy, local radiation and topical retinoids).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Valtoco

Products Affected

- VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vancomycin

Products Affected

- **VANCOGIN ORAL CAPSULE 125 MG, 250 MG**
- *vancomycin oral capsule 125 mg, 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For brand Vancocin, trial and failure of generic vancomycin capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vanflyta

Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) Induction, consolidation, or maintenance therapy 2) FLT3 ITD mutation status, as detected by an FDA-approved test 3) Concomitant therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Velsipity

Products Affected

- VELSIPITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inadequate response or intolerance to one systemic therapy (e.g. corticosteroids, azathioprine, cyclophosphamide).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Patients must have therapeutic failure or intolerance to 2 of the following preferred products Humira, Rinvoq, Xeljanz/Xeljanz XR and Stelara SC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Veltassa

Products Affected

- VELTASSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 6.4 mmol/L on at least two (2) screenings -AND- Modification of medications to reduce serum potassium levels were not successful, when applicable.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in serum potassium levels following Veltassa administration and continued treatment for hyperkalemia is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Venbysi XR

Products Affected

- *venlafaxine besylate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- venlafaxine extended-release product at a total daily dose greater than or equal to 75 mg has been received for greater than or equal to 4 days.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Venclexta

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and concomitant therapy, if applicable to diagnosis. For newly-diagnosed AML, age greater than or equal to 75 years or presence of at least one comorbidity that precludes use of intensive induction chemotherapy (i.e. severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) is required.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Veozah

Products Affected

- VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of moderate to severe vasomotor symptoms due to menopause -AND- one of the following (1 or 2): 1) therapeutic failure, contraindication, or intolerance to a generic hormone therapy product, or 2) attestation that hormone therapy is not clinically appropriate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verkazia

Products Affected

- VERKAZIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure or intolerance to 2 of the following medication classes or all are contraindicated: 1) generic ophthalmic antihistamines (e.g., olopatadine), 2) generic ophthalmic mast cell stabilizers (e.g., cromolyn sodium), 3) Generic ophthalmic corticosteroids (e.g., dexamethasone, prednisolone, fluorometholone).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verquvo

Products Affected

- VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of heart failure (NYHA Class II to IV) -AND- Left ventricular ejection fraction less than 45% -AND- Hospitalization for heart failure or received outpatient IV diuretics for heart failure -AND- Used in combination with a angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker or Entresto -AND- Used in combination with bisoprolol, carvedilol IR/ER or metoprolol succinate ER.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verzenio

Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR mutation status and HER2 mutation status 2) Alternatives tried/failed 3) Concomitant therapy, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Viberzi

Products Affected

- **VIBERZI**

PA Criteria	Criteria Details
Exclusion Criteria	Severe (Child-Pugh C) hepatic impairment
Required Medical Information	Documentation of diarrhea predominant, irritable bowel syndrome (IBS-D) -AND- trial/failure or intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vioice

Products Affected

- **VIJOICE ORAL GRANULES IN PACKET** **MG**
- **VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe manifestations of PIK3CA Related Overgrowth Spectrum (PROS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vimovo

Products Affected

- VIMOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) Trial/failure of ibuprofen/famotidine. -AND- 2) Trial/failure of naproxen/esomeprazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vimovo2

Products Affected

- *naproxen-esomeprazole*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) Trial/failure of naproxen used in combination with omeprazole. -AND- 2) Trial/failure of one additional generic formulary NSAID (other than naproxen) used in combination with another generic formulary PPI (other than omeprazole).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vittrakvi

Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation -AND- Tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vivjoa

Products Affected

- VIVJOA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of recurrent vulvovaginal candidiasis defined as at least 3 episodes of vulvovaginal candidiasis in less than one year -AND Documentation the member is NOT of reproductive potential defined as postmenopausal or another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vivlodex

Products Affected

- *meloxicam submicronized*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure or intolerance to generic meloxicam tablets and one additional generic NSAID
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vizimpro

Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer -AND- one of the following (1 or 2): 1. Epidermal growth factor (EGFR) exon 19 deletions - OR- 2. Epidermal growth factor receptor (EGFR) exon 21 L858R substitution mutations.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vonjo

Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of a platelet count of less than $50 \times 10^9/L$.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voquezna

Products Affected

- **VOQUEZNA ORAL TABLET 10 MG, 20 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For healing of erosive esophagitis and relief of heartburn associated with erosive esophagitis, therapeutic failure, contraindication, or intolerance to omeprazole and pantoprazole. For maintenance of erosive esophagitis healing and relief of heartburn associated with erosive esophagitis, therapeutic failure, contraindication, or intolerance to omeprazole and pantoprazole. For treatment of H. pylori infection, used in combination with amoxicillin -OR- amoxicillin and clarithromycin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	H.pylori tx: 14 days. Erosive esophagitis healing: 8 wks. Erosive esophagitis Maintenance: 6 mo.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voquezna Combo Pak

Products Affected

- **VOQUEZNA DUAL PAK**
- **VOQUEZNA TRIPLE PAK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voriconazole

Products Affected

- **VFEND IV**
- *voriconazole intravenous*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary cannot take oral voriconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of continued indicators of active disease (e.g. histopathology, positive cultures) is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vosevi

Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for all regimens recommended by the AASLD/IDSA guidelines containing the following agents: ledipasvir/sofosbuvir, sofosbuvir/velpatasvir, glecaprevir/pibrentasvir.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Votrient

Products Affected

- *pazopanib*
- **VOTRIENT**

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of adipocytic soft tissue sarcoma or gastrointestinal stromal tumor
Required Medical Information	Documentation of diagnosis. For advanced soft-tissue sarcoma, trial/failure of at least one prior chemotherapy regimen.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voxzogo

Products Affected

- VOXZOGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of achondroplasia as evidenced by an FGFR3 gene mutation detected by genetic testing -AND- all of the following (1-2): 1) baseline annualized growth velocity (AGV), 2) attestation that the member's epiphyses have not closed.
Age Restrictions	Deny if greater than 17 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increase in annualized growth velocity (AGV) from baseline and attestation that the member's epiphyses have not closed are required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vraylar

Products Affected

- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For a diagnosis of schizophrenia and bipolar I disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vtama

Products Affected

- VTAMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3). 1) Documentation of plaque psoriasis -AND- 2) trial and failure or intolerance to at least one generic, formulary medium, high, or ultrahigh potency topical corticosteroid (e.g. fluocinonide, triamcinolone, betamethasone) -OR- psoriasis on facial or intertriginous areas -AND- 3) trial and failure, contraindication, or intolerance to one generic, formulary vitamin D analog (e.g., calcipotriene, calcitriol)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vuity

Products Affected

- VUITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 18 years of age.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vumerity

Products Affected

- VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Wainua

Products Affected

- WAINUA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of polyneuropathy associated with hereditary TTR (hATTR) amyloidosis with mutation in TTR gene confirmed by genetic testing -AND- Neurologic examination shows clinical signs and symptoms of the disease (e.g. peripheral/autonomic neuropathy, motor disability, carpal tunnel, etc.) -AND- Attestation of peripheral neuropathy impairment score (NIS) of 10 or greater -AND- Not simultaneously utilizing other gene targeted therapy for polyneuropathy of hATTR
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement of polyneuropathy from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Wakix

Products Affected

- **WAKIX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If no diagnosis of cataplexy, trial and failure, intolerance, or contraindication to generic modafinil and a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, provider attestation of improvement in symptoms of narcolepsy or improvement in symptoms of cataplexy (if applicable).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Wegovy

Products Affected

- **WEGOVY SUBCUTANEOUS PEN INJECTOR 0.25 MG/0.5 ML, 0.5 MG/0.5 ML, 1 MG/0.5 ML, 1.7 MG/0.75 ML, 2.4 MG/0.75 ML**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment to reduce body weight and maintain weight reduction
Required Medical Information	Documentation of use in combination with reduced calorie diet and increased physical activity to reduce the risk of major adverse cardiovascular (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease and either obesity or overweight -AND- Baseline BMI is greater than or equal to 27 kg/m ² -AND- Will not be used in combination with GLP1 RA or GLP1 RA combinations.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of continued use in combination with reduced calorie diet and increased physical activity for reduction of major adverse cardiovascular event risk -AND- Requested dose is 1.7mg or 2.4mg weekly.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Welireg

Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For von Hippel Lindau (VHL) syndrome, one of the following diagnoses not requiring immediate surgery (1, 2, or 3): 1) Renal cell carcinoma (RCC) 2) CNS hemangioblastoma 3) Pancreatic neuroendocrine tumor. For advanced RCC, prior treatment with a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) -AND- a programmed death receptor-1 (PD-1) or a programmed death-ligand (PD-L1) inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Winrevair

Products Affected

- WINREVAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group -AND- Concomitant use of at least one of the following (1-4): 1) generic endothelin-1 receptor antagonist, 2) phosphodiesterase type 5 inhibitor, 3) soluble guanylate cyclase stimulator, 4) generic prostacyclin agent -AND- Background therapy with PAH-specific therapies will be continued while being treated with Winrevair.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xalkori

Products Affected

- **XALKORI ORAL CAPSULE**
- **XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer (NSCLC), disease is ALK-positive or ROS1-positive. For relapsed or refractory anaplastic large cell lymphoma (ALCL), disease is ALK-positive. For unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT), disease is ALK-positive.
Age Restrictions	Deny if less than 18 years of age for NSCLC. Deny if less than 1 year of age or greater than 21 years of age for ALCL.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For crizotinib oral pellets and NSCLC, inability to swallow capsules is required. For crizotinib oral pellets and ALCL / IMT, inability to swallow oral capsules -OR- body surface area less than 1.34 m ² is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xcopri

Products Affected

- **XCOPRI**
- **XCOPRI MAINTENANCE PACK**
- **XCOPRI TITRATION PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Therapeutic failure, intolerance or contraindication to 1 other anti-epileptic drug (e.g. carbamazepine, levetiracetam)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xdemvy

Products Affected

- XDEMVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis confirmed by identification of Demodex infection via microscopic examination of pulled eyelashes -OR- identification of collarettes via slit-lamp evaluation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xeljanz

Products Affected

- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis and an inadequate response or intolerance to methotrexate. Xeljanz immediate release for juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID).
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis, psoriatic arthritis, ulcerative colitis, ankylosing spondylitis. For Xeljanz regular release tablet, deny if less than 2 years of age for juvenile idiopathic arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced inadequate response or intolerance to at least 1 tumor necrosis factor (TNF) blocker. Doses greater than 10 mg per day for Xeljanz and 11 mg per day for Xeljanz XR will not be approved for rheumatoid arthritis and psoriatic arthritis. Doses greater than 20mg per day for Xeljanz and 22 mg per day for Xeljanz XR will not be approved for ulcerative colitis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xeljanz Solution

Products Affected

- XELJANZ ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of juvenile idiopathic arthritis -AND- Inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xenazine

Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*
- **XENAZINE ORAL TABLET 12.5 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if requesting brand Xenazine, trial and failure or intolerance to generic tetrabenazine has been documented - AND- attestation that the beneficiary is not actively suicidal
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	In patients with comorbid depression, attestation of adequate treatment for depression is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xermelo

Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of carcinoid syndrome diarrhea AND used in combination with a somatostatin analog AND trial and failure of somatostatin analog monotherapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in average number of daily bowel movements -AND- the member will continue to use in combination with a somatostatin analog.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xgeva

Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For hypercalcemia of malignancy, refractory to bisphosphonates. For giant cell tumor of bone, unresectable or surgical resection is likely to result in severe morbidity -AND- one of the following (1. or 2.)- 1.) the member is 18 years old or older -OR- 2.) the member is a skeletally mature adolescent (e.g. has at least one mature long bone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xifaxan

Products Affected

- XIFAXAN ORAL TABLET 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 or 2. 1) Diagnosis of hepatic encephalopathy AND trial/failure, intolerance, or contraindication to lactulose. 2) Diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D) AND trial/failure, intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Hepatic encephalopathy: 1 year. IBS-D: 14 days.
Other Criteria	No more than three courses of rifaximin for the treatment of IBS-D will be approved per lifetime.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xolair

Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of chronic idiopathic urticaria with trial/failure or intolerance of a second-generation non-sedating H1 antihistamine at the maximum recommended doses (e.g., cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine) -OR- Documentation of moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen -AND- Baseline IgE titer greater than or equal to 30 IU/mL -AND- symptoms that are inadequately controlled despite a 3 month trial of both 1. and 2. 1) medium-dose inhaled corticosteroid or systemic steroid 2) a long-acting beta-agonist or leukotriene antagonist - AND- patient is currently on a long-acting beta2-agonist, leukotriene modifier, or theophylline -OR- Documentation of chronic rhinosinusitis with nasal polyps -AND- will use concomitantly with nasal corticosteroid maintenance treatment -OR- Documentation of IgE mediated food allergy confirmed by skin prick test or food-specific antibodies -AND- previous allergic reaction to food -AND- using for the reduction of allergic reactions (type 1), including anaphylaxis -AND- used in conjunction with food allergen avoidance -AND- member has a documented prescription for epinephrine.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, attestation of decreased rescue medication or oral corticosteroid use, decreased frequency of asthma exacerbations, increased pulmonary function from baseline (e.g. FEV1), or reduction in reported asthma related symptoms in treatment of asthma -OR- improved symptoms in treatment of CIU -OR- reduction in nasal polyp score or nasal congestion/obstruction severity score in treatment of nasal polyps -OR- member requires continuation of therapy and will continue food allergen avoidance in IgE-mediated food allergy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xolremdi

Products Affected

- **XOLREMDI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of WHIM syndrome (warts, hypogammaglobulinemia, infections, myelokathexis).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in incidence of infections is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xospata

Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- FLT3 mutations, if applicable to diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xpovio

Products Affected

- **XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use in combination with dexamethasone for relapse or refractory multiple myeloma with failure, intolerance or contraindication to 5 therapies (e.g. bortezomib, carfilzomib, lenalidomide, pomalidomide and daratumumab) -OR- Documentation of use in combination with both bortezomib and dexamethasone for relapse or refractory multiple myeloma after receiving 1 prior multiple myeloma therapy -OR- Documentation of relapsing or refractory diffuse large B-cell lymphoma with failure, intolerance or contraindication to at least 2 lines of systemic therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xtandi

Products Affected

- **XTANDI ORAL CAPSULE**
- **XTANDI ORAL TABLET 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1 or 2): 1) Concomitant GnRH analog, 2) The member has had a bilateral orchiectomy, if applicable to diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xuriden

Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary orotic aciduria
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xyrem

Products Affected

- *sodium oxybate*
- **XYREM**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	<p>If no diagnosis of cataplexy- trial and failure, intolerance, or contraindication to generic modafinil AND a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xywav

Products Affected

- XYWAV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP -AND- Sensitivity to sodium intake due to heart failure, hypertension or impaired renal function. If the member has a diagnosis of cataplexy, provision of baseline number of cataplexy episodes is required. -OR- (II) Diagnosis of idiopathic hypersomnia - AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- documentation the member does not have cataplexy -AND- documentation of less than 2 SOREMPs -AND- documentation of the following (1, 2, or 3): 1) MSLT documenting MSL less than or equal to 8 minutes -OR- 2) polysomnography demonstrating total sleep time greater than or equal to 660 minutes per 24 hours -OR- 3) wrist actigraphy demonstrating total sleep time greater than or equal to 660 minutes per 24 hours.</p>
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	If diagnosis of narcolepsy without cataplexy- trial and failure, intolerance, or contraindication to generic modafinil AND a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy, idiopathic hypersomnia and cataplexy (if applicable) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Yonsa

Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) Alternatives tried/failed 2) Concomitant therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zavesca

Products Affected

- *miglustat*
- YARGESA
- ZAVESCA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of mild to moderate type 1 Gaucher disease confirmed by the following (A. or B.) A. (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm ³ -OR- 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B. Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation enzyme replacement therapy (e.g. Cerezyme, Elelyso, or VPRIV) is not a therapeutic option. For brand Zavesca, documentation of failure on generic miglustat.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zavzpret

Products Affected

- ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- Therapeutic failure, contraindication or intolerance to one generic triptan -AND- Inability to swallow capsules/tablets
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zejula

Products Affected

- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restrictions	Pending CMS Review
Prescriber Restrictions	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Indications	Pending CMS Review
Off Label Uses	Pending CMS Review
Part B Prerequisite	No

Zelboraf

Products Affected

- TAFINLAR ORAL CAPSULE
- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Documentation of diagnosis -AND- both of the following. 1) BRAF mutations, if applicable to diagnosis. 2) Alternatives tried/failed and concomitant therapy, if applicable to diagnosis (e.g. diagnosis of V600K metastatic melanoma and drug regimen of Zelboraf and Cotellic)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zepatier

Products Affected

- ZEPATIER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for all regimens recommended by the AASLD/IDSA guidelines containing the following agents: ledipasvir/sofosbuvir, sofosbuvir/velpatasvir, glecaprevir/pibrentasvir.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zeposia

Products Affected

- ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Zeposia and other disease modifying agents such as interferons, Copaxone, Tysabri.
Required Medical Information	Documentation of diagnosis. For moderate to severe active ulcerative colitis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate, cyclosporine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Ulcerative Colitis: 12 months, Multiple Sclerosis: 24 months
Other Criteria	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products Humira, Xeljanz/Xeljanz XR and Stelara SC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zokinvy

Products Affected

- ZOKINVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Hutchinson-Gilford Progeria Syndrome with mutation of the LMNA gene and BSA of 0.39 square meters or above -OR- Documentation of processing-deficient Progeroid Laminopathies with BSA of 0.39 square meters or above -AND- one of the following (1 or 2): 1) Heterozygous LMNA mutation and progerin-like protein accumulation 2.) Homozygous ZMPSTE24 mutations or compound heterozygous ZMPSTE23 mutations
Age Restrictions	Deny if less than 12 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zolinza

Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations in patients with cutaneous T-cell lymphoma (CTCL) who have progressive, persistent, or recurrent disease on or following 2 systemic therapies. Systemic therapies include bexarotene, interferon alpha, extracorporeal photochemotherapy, PUVA, single agent or combination chemotherapies (e.g. cyclophosphamide, vinblastine, romidepsin)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zolpidem Capsule

Products Affected

- *zolpidem*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of insomnia characterized by difficulties with sleep onset - AND- therapeutic failure or intolerance to 2 of the following (1-3): 1) eszopiclone, 2) zaleplon, 3) generic zolpidem tablets/zolpidem ER tablets. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	Deny if greater than 64 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zonisade

Products Affected

- ZONISADE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Documentation of adjunctive therapy -AND- Inability to swallow capsules -AND- Therapeutic failure/intolerance to 2 or contraindication to all of the following (1-3): 1) generic carbamazepine suspension/chewable tablet/extended-release capsule, 2) generic lacosamide solution, 3) generic oxcarbazepine suspension.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zoryve

Products Affected

- ZORYVE TOPICAL CREAM 0.3 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3). 1) Documentation of plaque psoriasis -AND- 2) trial and failure or intolerance to at least one generic, formulary medium, high, or ultrahigh potency topical corticosteroid (e.g. fluocinonide, triamcinolone, betamethasone) -OR- psoriasis on facial or intertriginous areas -AND- 3) trial and failure, contraindication, or intolerance to one generic, formulary vitamin D analog (e.g., calcipotriene, calcitriol)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zoryve Foam

Products Affected

- ZORYVE TOPICAL FOAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3). 1) Documentation of seborrheic dermatitis -AND- 2) trial and failure or intolerance to at least one generic, formulary topical corticosteroid -OR- seborrheic dermatitis on facial or intertriginous areas -AND- 3) if the member is 12 years of age or older, trial and failure, contraindication, or intolerance to one generic, topical antifungal agent for seborrheic dermatitis (e.g., ketoconazole, ciclopirox).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ztalmy

Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures associated with CDKL5 deficiency confirmed by genetic testing -AND- therapeutic failure or intolerance to 2 previous antiepileptic therapies
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ZTLido

Products Affected

- ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) -AND- One of the following (1,2 or 3): 1) trial and failure of 1 other agent used to treat PHN (e.g. gabapentin) 2) Inability to swallow oral medication 3) Unable to take an oral medication due to potential adverse events (e.g. sedation)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Patients must have therapeutic failure or contraindication to Lidoderm or lidocaine patch 5%.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zurzuvaе

Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of moderate to severe postpartum depression -AND- less than or equal to 12 months postpartum.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zydelig

Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	First line treatment. Combination use with benadmustine and/or rituximab for the treatment of FL.
Required Medical Information	Documentation of diagnosis -AND- all of the following, if applicable to diagnosis: 1) Alternatives tried/failed 2) Concomitant therapy 3) Candidacy for use of rituximab alone
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zykadia

Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- ALK mutations, as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zymfentra

Products Affected

- ZYMFENTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation of clinical response or remission following at least 3 doses of IV infliximab at least 10 weeks before initiating therapy with Zymfentra.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Rinvoq, Xeljanz/Xeljanz XR, and Stelara SC. For moderate to severe active Crohn's disease, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product (i.e. Humira, Cyltezo, Hyrimoz, adalimumab-adaz, or adalimumab-adbm), Rinvoq, Skyrizi SC, and Stelara SC
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zytiga

Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*
- **ZYTIGA ORAL TABLET 250 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- concomitant therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For brand Zytiga, trial and failure of generic abiraterone is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Index of Drugs

ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET WITH SENSOR AND STRIP 15 MG, 2 MG, 20 MG, 30 MG, 5 MG	27	<i>adapalene-benzoyl peroxide</i>	426
ABILIFY MYCITE STARTER KIT ORAL TABLET WITH SENSOR, STRIP, POD 10 MG	27	ADBRY SUBCUTANEOUS SYRINGE	8
ABILIFY ORAL TABLET	27	ADCIRCA	327
<i>abiraterone oral tablet 250 mg, 500 mg</i>	511	ADEMPAS	327
ABRILADA(CF)	6	ADLARITY	10
ABRILADA(CF) PEN	6	AFINITOR	11
<i>acetaminophen-caff-dihydrocod</i>	318	AFINITOR DISPERZ ORAL TABLET FOR SUSPENSION 2 MG, 3 MG, 5 MG .	11
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i>	318	AGAMREE	12
<i>acetaminophen-codeine oral tablet</i>	318	AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML, 70 MG/ML	13
<i>acitretin</i>	1	AJOVY AUTOINJECTOR	14
ACTEMRA ACTPEN	2	AJOVY SYRINGE	14
ACTEMRA SUBCUTANEOUS	2	AKEEGA	15
ACTHAR	3	AKLIEF	426
ACTIMMUNE	5	ALECENSA	16
<i>adalimumab-aacf subcutaneous pen injector kit</i>	6	ALKINDI SPRINKLE	17
<i>adalimumab-aaty</i>	6	<i>alose tron oral tablet 0.5 mg, 1 mg</i>	225
<i>adalimumab-adaz</i>	165	ALPRAZOLAM INTENSOL	318
<i>adalimumab-adbm subcutaneous pen injector kit 40 mg/0.4 ml, 40 mg/0.8 ml</i>	165	<i>alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	318
<i>adalimumab-adbm subcutaneous syringe kit 10 mg/0.2 ml, 20 mg/0.4 ml, 40 mg/0.4 ml, 40 mg/0.8 ml</i>	165	<i>alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg, 2 mg, 3 mg</i>	318
ADALIMUMAB-ADBM(CF) PEN CROHNS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 40 MG/0.8 ML	165	<i>alprazolam oral tablet, disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	318
ADALIMUMAB-ADBM(CF) PEN PS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.4 ML, 40 MG/0.8 ML	165	ALTRENO	426
<i>adalimumab-fkjp subcutaneous pen injector kit</i>	6	ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG	19
<i>adalimumab-fkjp subcutaneous syringe kit</i>	6	ALUNBRIG ORAL TABLETS, DOSE PACK	19
<i>adalimumab-ryvk subcutaneous auto-injector, kit</i>	6	ALVAIZ ORAL TABLET 18 MG, 36 MG, 54 MG, 9 MG	419
<i>adapalene topical cream</i>	426	ALYQ	327
<i>adapalene topical gel 0.3 %</i>	426	AMBIEN	318
<i>adapalene topical swab</i>	426	AMBIEN CR	318
		<i>ambrisentan</i>	327
		<i>amitriptyline</i>	160
		<i>amitriptyline-chlordiazepoxide</i>	160
		AMJEVITA(CF) AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 40 MG/0.4 ML, 40 MG/0.8 ML, 80 MG/0.8 ML	6

AMJEVITA(CF) SUBCUTANEOUS SYRINGE 10 MG/0.2 ML, 20 MG/0.2 ML, 20 MG/0.4 ML, 40 MG/0.4 ML, 40 MG/0.8 ML6	<i>baclofen oral solution 10 mg/5 ml (2 mg/ml)</i>34
<i>amphetamine sulfate</i>116	<i>baclofen oral suspension</i>133
AMPYRA21	BAFIERTAM35
ANAFRANIL160	BALVERSA36
ANDROGEL TRANSDERMAL GEL IN METERED-DOSE PUMP415	BANZEL37
APOKYN23	BELBUCA318
<i>apomorphine</i>23	BENLYSTA SUBCUTANEOUS38
ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG18	<i>benznidazole</i>40
ARAZLO409	<i>benztropine oral</i>160
ARCALYST185	BERINERT INTRAVENOUS KIT41
ARIKAYCE24	BESREMI43
<i>aripiprazole</i>27	BETASERON SUBCUTANEOUS KIT184
<i>armodafinil</i>271	BETHKIS63
ASCOMP WITH CODEINE318	<i>bexarotene oral</i>402
<i>asenapine maleate</i>361	<i>bexarotene topical</i>402
ASPRUZYO SPRINKLE25	BIMZELX44
ATIVAN ORAL TABLET 0.5 MG, 1 MG, 2 MG318	BIMZELX AUTOINJECTOR44
ATORVALIQ383	BIVIGAM172
ATRALIN426	BONJESTA45
AUBAGIO28	<i>bosentan</i>327
AUGTYRO29	BOSULIF ORAL CAPSULE 100 MG, 50 MG46
AURYXIA30	BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG46
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG31	BRAFTOVI47
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG31	BRONCHITOL48
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 6 MG (14)-12 MG (14)-24 MG (14)31	BRUKINSA49
AUVELITY32	BUPHENYL50
AVEED415	<i>buprenorphine</i>318
AVONEX INTRAMUSCULAR PEN INJECTOR KIT184	<i>butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg, 50-325-40-30 mg</i>318
AVONEX INTRAMUSCULAR SYRINGE KIT184	BUTRANS318
AYVAKIT33	BYDUREON BCISE147
	BYETTA SUBCUTANEOUS PEN INJECTOR 10 MCG/DOSE(250 MCG/ML) 2.4 ML, 5 MCG/DOSE (250 MCG/ML) 1.2 ML147
	BYLVAY ORAL CAPSULE 1,200 MCG, 400 MCG51
	BYLVAY ORAL PELLETT 200 MCG, 600 MCG51
	CABLIVI INJECTION KIT52
	CABOMETYX53
	CABTREO426
	CALQUENCE54

CALQUENCE (ACALABRUTINIB MAL)	54	CORLANOR ORAL SOLUTION	76
CAMZYOS	55	CORLANOR ORAL TABLET 5 MG, 7.5 MG	76
CAPLYTA	56	CORTROPHIN GEL	330
CAPRELSA ORAL TABLET 100 MG, 300 MG	57	COSENTYX (2 SYRINGES)	77
CARAC	58	COSENTYX PEN (2 PENS)	77
CARBAGLU	59	COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML	77
<i>carbinoxamine maleate oral liquid</i>	60	COSENTYX UNOREADY PEN	77
<i>carbinoxamine maleate oral tablet 4 mg</i>	60	COTELLIC	78
<i>carglumic acid</i>	59	CRESEMBA ORAL	79
<i>carisoprodol</i>	160	CRINONE	80
CAYSTON	61	CUVRIOR	81
CERDELGA	62	CYLTEZO(CF)	165
CHENODAL	64	CYLTEZO(CF) PEN	165
<i>chlordiazepoxide hcl</i>	318	CYLTEZO(CF) PEN CROHN'S-UC-HS	165
<i>chlorzoxazone</i>	160	CYLTEZO(CF) PEN PSORIASIS-UV ..	165
CHOLBAM	65	<i>cyproheptadine</i>	160
CIALIS ORAL TABLET 2.5 MG, 5 MG	66	CYSTDROPS	82
CIBINQO	67	CYSTARAN	82
CIMZIA	68	<i>dalfampridine</i>	21
CIMZIA POWDER FOR RECONST	68	DARAPRIM	83
CINRYZE	70	DAURISMO ORAL TABLET 100 MG, 25 MG	84
<i>citalopram oral capsule</i>	72	DAYBUE	85
<i>clemastine oral syrup</i>	73	DAYTRANA	86
<i>clindamycin-tretinoin</i>	426	<i>deferasirox</i>	87
<i>clobazam oral suspension</i>	283	<i>deferiprone</i>	124
<i>clobazam oral tablet</i>	283	<i>deflazacort</i>	100
<i>clomipramine</i>	160	DEMEROL (PF) INJECTION SYRINGE 25 MG/ML	318
<i>clonazepam oral tablet 0.5 mg, 1 mg, 2 mg</i>	318	DEMEROL INJECTION	318
<i>clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	318	DEM SER	244
<i>clonidine hcl oral tablet extended release 12 hr</i>	9	DEPO-TESTOSTERONE	415
<i>clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg</i>	318	DIACOMIT ORAL CAPSULE 250 MG, 500 MG	88
<i>codeine sulfate</i>	318	DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG	88
<i>codeine-butalbital-asa-caff</i>	318	DIAZEPAM INTENSOL	318
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)	74	<i>diazepam oral solution 5 mg/5 ml (1 mg/ml)</i>	318
CONZIP	318	<i>diazepam oral tablet</i>	318
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML	144	DIBENZYLINE	307
COPIKTRA	75	DICLEGIS	45
		<i>diclofenac epolamine</i>	132

<i>diclofenac sodium topical gel 3 %</i>	377	ENDOCET	318
DIFFERIN TOPICAL CREAM	426	ENSPRYNG	106
DIFFERIN TOPICAL GEL WITH PUMP	426	ENTADFI	107
DIFFERIN TOPICAL LOTION	426	ENTYVIO PEN	108
<i>dihydroergotamine nasal</i>	89	EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG	109
DILAUDID ORAL LIQUID	318	EPCLUSA ORAL TABLET	109
DILAUDID ORAL TABLET	318	EPIDIOLEX	110
<i>dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg</i>	412	EPIDUO FORTE	426
DOJOLVI	90	EPIDUO TOPICAL GEL WITH PUMP	426
DOPTELET (10 TAB PACK)	91	EPRONTIA	111
DOPTELET (15 TAB PACK)	91	<i>ergoloid</i>	160
DOPTELET (30 TAB PACK)	91	<i>ergotamine-caffeine</i>	112
<i>doxepin oral capsule</i>	160	ERIVEDGE	113
<i>doxepin oral concentrate</i>	160	ERLEADA ORAL TABLET 240 MG, 60 MG	114
<i>doxepin oral tablet</i>	160	<i>erlotinib</i>	97
<i>doxepin topical</i>	92	ESBRIET ORAL CAPSULE	187
<i>doxylamine-pyridoxine (vit b6)</i>	45	ESBRIET ORAL TABLET	187
<i>droxidopa oral capsule 100 mg, 200 mg, 300 mg</i>	263	<i>estazolam</i>	318
DUOBRII	94	<i>eszopiclone</i>	318
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML	95	EUCRISA	115
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 100 MG/0.67 ML, 200 MG/1.14 ML, 300 MG/2 ML	95	EVEKEO	116
EDLUAR	318	EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML (105MG/1.17MLX2)	117
EGRIFTA SV	98	<i>everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg</i>	11
ELYXYB	99	<i>everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg</i>	11
EMFLAZA	100	EVRYSDI	118
EMGALITY PEN	102	EXJADE	87
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3)	102	EXSERVAN	350
ENBREL MINI	104	FABHALTA	119
ENBREL SUBCUTANEOUS SOLUTION	104	FABIOR	409
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)	104	FASENRA PEN	121
ENBREL SURECLICK	104	FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML	121
ENDARI	105	<i>febuxostat</i>	435
		<i>fentanyl citrate buccal lozenge on a handle 1,200 mcg, 1,600 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg</i>	424
		<i>fentanyl citrate buccal tablet, effervescent 100 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg</i>	424

<i>fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 37.5 mcg/hour, 50 mcg/hr, 62.5 mcg/hour, 75 mcg/hr, 87.5 mcg/hour</i>	318	GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)	172
FENTORA BUCCAL TABLET, EFFERVESCENT 100 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG	424	GATTEX 30-VIAL	140
FERRIPROX	124	GAVRETO	141
FERRIPROX (2 TIMES A DAY)	124	<i>gefitinib</i>	189
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)	125	GENOTROPIN	152
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG	125	GENOTROPIN MINIQUICK	152
FILSPARI	126	GILENYA	142
FILSUVEZ	127	GILOTRIF	97
<i> fingolimod</i>	142	GIMOTI	143
FINTEPLA	128	GLASSIA	18
FIORICET WITH CODEINE	318	<i>glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml</i>	144
FIRAZYR	129	GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML	144
FIRDAPSE	131	GLEEVEC ORAL TABLET 100 MG, 400 MG	145
FLECTOR	132	GLEOSTINE	146
FLEQSUVY	133	GLYCATE	148
FLOLIPID	383	<i>glycopyrrolate oral tablet 1.5 mg</i>	148
<i>fluorouracil topical cream 0.5 %</i>	58	GOCOVRI ORAL CAPSULE,EXTENDED RELEASE 24HR 137 MG, 68.5 MG	149
<i>flurazepam</i>	318	GRALISE ORAL TABLET EXTENDED RELEASE 24 HR 300 MG, 450 MG, 600 MG, 750 MG, 900 MG	150
FORTEO	134	GRASTEK	151
FOTIVDA	135	<i>guanfacine oral tablet extended release 24 hr</i>	9
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG	136	HADLIMA	6
FUROSCIX	137	HADLIMA PUSHTOUCH	6
<i>gabapentin oral capsule 100 mg, 300 mg, 400 mg</i>	138	HADLIMA(CF)	6
<i>gabapentin oral solution 250 mg/5 ml</i>	138	HADLIMA(CF) PUSHTOUCH	6
<i>gabapentin oral tablet 600 mg, 800 mg</i>	138	HAEGARDA	153
<i>gabapentin oral tablet extended release 24 hr 300 mg, 600 mg</i>	150	HALCION ORAL TABLET 0.25 MG...	318
GALAFOLD	139	HARVONI ORAL PELLETS IN PACKET	155
GAMMAGARD LIQUID	172	HARVONI ORAL TABLET 90-400 MG	155
GAMMAGARD S-D (IGA < 1 MCG/ML)	172	HEMADY	156
GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)	172	HETLIOZ	157
GAMMAPLEX	172	HETLIOZ LQ	159
GAMMAPLEX (WITH SORBITOL)	172	HORIZANT ORAL TABLET EXTENDED RELEASE 300 MG, 600 MG	164

HULIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT	6	HYRIMOZ(CF) SUBCUTANEOUS SYRINGE 10 MG/0.1 ML, 20 MG/0.2 ML, 40 MG/0.4 ML	165
HULIO(CF) SUBCUTANEOUS SYRINGE KIT	6	HYSINGLA ER	318
HUMATROPE INJECTION CARTRIDGE	152	IBRANCE	168
HUMIRA PEN	165	IBSRELA	169
HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML	165	<i>ibuprofen-famotidine</i>	93
HUMIRA(CF)	165	<i>icatibant</i>	129
HUMIRA(CF) PEN	165	ICLUSIG	170
HUMIRA(CF) PEN CROHNS-UC-HS ..	165	IDACIO(CF)	6
HUMIRA(CF) PEN PEDIATRIC UC ..	165	IDACIO(CF) PEN CROHN-UC STARTR	6
HUMIRA(CF) PEN PSOR-UV-ADOL HS	165	IDACIO(CF) PEN PSORIASIS START ...	6
<i>hydrocodone bitartrate oral capsule, oral only, er 12hr</i>	318	IDACIO(CF) PEN SUBCUTANEOUS PEN INJECTOR KIT	6
<i>hydrocodone bitartrate oral tablet, oral only, ext. rel. 24 hr</i>	318	IDHIFA ORAL TABLET 100 MG, 50 MG	171
<i>hydrocodone-acetaminophen oral solution 7.5-325 mg/15 ml</i>	318	ILUMYA	174
<i>hydrocodone-acetaminophen oral tablet 10-300 mg, 10-325 mg, 5-300 mg, 5-325 mg, 7.5-300 mg, 7.5-325 mg</i>	318	<i>imatinib oral tablet 100 mg, 400 mg</i>	145
<i>hydrocodone-ibuprofen</i>	318	IMBRUVICA ORAL CAPSULE 140 MG, 70 MG	175
<i>hydromorphone (pf) injection solution 10 (mg/ml) (5 ml), 10 mg/ml</i>	318	IMBRUVICA ORAL SUSPENSION	175
<i>hydromorphone oral liquid</i>	318	IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG	175
<i>hydromorphone oral tablet</i>	318	<i>imipramine hcl</i>	160
<i>hydromorphone oral tablet extended release 24 hr 12 mg, 16 mg, 32 mg, 8 mg</i> ..	318	<i>imipramine pamoate</i>	160
<i>hydroxyzine hcl oral solution 10 mg/5 ml</i> ..	160	INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE	176
<i>hydroxyzine hcl oral tablet</i>	160	INCRELEX	177
<i>hydroxyzine pamoate</i>	160	INGREZZA INITIATION PK(TARDIV)	179
HYFTOR	167	INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG	179
HYRIMOZ PEN CROHN'S-UC STARTER	165	INGREZZA SPRINKLE	179
HYRIMOZ PEN PSORIASIS STARTER	165	INLYTA	180
HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML, 80 MG/0.8 ML- 40 MG/0.4 ML	165	INQOVI	181
HYRIMOZ(CF) PEN SUBCUTANEOUS PEN INJECTOR 40 MG/0.4 ML, 80 MG/0.8 ML	165	INREBIC	182
		INTRAROSA	186
		INTUNIV ER	9
		IRESSA	189
		ISTURISA ORAL TABLET 1 MG, 5 MG	190
		<i>itraconazole</i>	191
		<i>ivermectin oral</i>	192
		IWILFIN	193
		JADENU	87

JADENU SPRINKLE	87	LENVIMA	213
JAKAFI	194	LETAIRIS	327
JATENZO ORAL CAPSULE 158 MG, 198 MG, 237 MG	415	LEUKINE INJECTION RECON SOLN	214
JAVYGTOR	210	<i>levorphanol tartrate</i>	318
JAYPIRCA ORAL TABLET 100 MG, 50 MG	195	LIBERVANT	216
JOENJA	196	LICART	132
JUXTAPID	163	<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i>	423
JYNARQUE ORAL TABLET	197	<i>lidocaine topical adhesive patch,medicated 5 %</i>	217
JYNARQUE ORAL TABLETS, SEQUENTIAL	197	<i>lidocaine topical ointment</i>	423
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG	198	<i>lidocaine-prilocaine topical cream</i>	423
KALYDECO ORAL TABLET	198	LIDOCAN III	217
KATERZIA	20	LIDODERM	217
KERENDIA	199	LIQREV	327
KESIMPTA PEN	200	LITFULO	218
KEVEYIS	201	LIVMARLI ORAL SOLUTION 9.5 MG/ML	219
KEVZARA	202	LIVTENCITY	220
KINERET	204	LODOCO	221
KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG	205	LOKELMA	222
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)	205	LONSURF	223
KITABIS PAK	63	LORAZEPAM INTENSOL	318
KLISYRI	206	<i>lorazepam oral tablet 0.5 mg, 1 mg, 2 mg</i> ..	318
KLONOPIN ORAL TABLET 0.5 MG, 1 MG, 2 MG	318	LORBRENA ORAL TABLET 100 MG, 25 MG	224
KORLYM	207	LOREEV XR ORAL CAPSULE,EXTENDED RELEASE 24HR 1 MG, 1.5 MG, 2 MG, 3 MG	318
KOSELUGO ORAL CAPSULE 10 MG, 25 MG	208	LORZONE	160
KRAZATI	209	LOTRONEX ORAL TABLET 0.5 MG, 1 MG	225
KUVAN	210	LOVAZA	226
LAMPIT	211	LUMAKRAS ORAL TABLET 120 MG, 320 MG	227
<i>lapatinib</i>	432	LUMRYZ	228
LATUDA ORAL TABLET 120 MG, 20 MG, 40 MG, 60 MG, 80 MG	212	LUNESTA	318
<i>ledipasvir-sofosbuvir</i>	155	LUPKYNIS	229
<i>lenalidomide</i>	346	LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	230

LUPRON DEPOT-PED	
INTRAMUSCULAR KIT 7.5 MG (PED)	
.....	230
LUPRON DEPOT-PED	
INTRAMUSCULAR SYRINGE KIT	230
<i>lurasidone oral tablet 120 mg, 20 mg, 40</i>	
<i>mg, 60 mg, 80 mg</i>	212
LYBALVI	231
LYNPARZA	232
LYRICA CR	233
LYRICA ORAL CAPSULE 100 MG,	
150 MG, 200 MG, 225 MG, 25 MG, 300	
MG, 50 MG, 75 MG	233
LYRICA ORAL SOLUTION	233
LYTGOBI ORAL TABLET 12	
MG/DAY (4 MG X 3), 16 MG/DAY (4	
MG X 4), 20 MG/DAY (4 MG X 5)	234
LYVISPAH	235
MAVENCLAD (10 TABLET PACK)	236
MAVENCLAD (4 TABLET PACK)	236
MAVENCLAD (5 TABLET PACK)	236
MAVENCLAD (6 TABLET PACK)	236
MAVENCLAD (7 TABLET PACK)	236
MAVENCLAD (8 TABLET PACK)	236
MAVENCLAD (9 TABLET PACK)	236
MAVYRET ORAL PELLETS IN	
PACKET	237
MAVYRET ORAL TABLET	237
MAYZENT ORAL TABLET 0.25 MG,	
1 MG, 2 MG	238
MAYZENT STARTER(FOR 1MG	
MAINT)	238
MAYZENT STARTER(FOR 2MG	
MAINT)	238
<i>megestrol oral suspension 400 mg/10 ml</i>	
<i>(40 mg/ml), 625 mg/5 ml (125 mg/ml)</i>	239
<i>megestrol oral tablet</i>	239
MEKINIST ORAL RECON SOLN	241
MEKINIST ORAL TABLET 0.5 MG, 2	
MG	240
MEKTOVI	242
<i>meloxicam submicronized</i>	454
<i>mepredine (pf) injection solution 100</i>	
<i>mg/ml, 25 mg/ml, 50 mg/ml</i>	318
<i>mepredine oral solution</i>	318
<i>mepredine oral tablet 50 mg</i>	318
<i>metaxalone</i>	160
<i>methadone oral solution 10 mg/5 ml, 5</i>	
<i>mg/5 ml</i>	318
<i>methadone oral tablet 10 mg, 5 mg</i>	318
<i>methamphetamine</i>	243
METHITEST	22
<i>methylphenidate</i>	86
<i>methyltestosterone oral capsule</i>	22
<i>metyrosine</i>	244
<i>mifepristone oral tablet 300 mg</i>	207
<i>miglustat</i>	493
MIGRANAL	89
<i>modafinil</i>	325
<i>morphine concentrate oral solution</i>	318
<i>morphine oral capsule, er multiphase 24 hr</i>	
<i>120 mg, 30 mg, 45 mg, 60 mg, 75 mg, 90</i>	
<i>mg</i>	318
<i>morphine oral capsule, extend. release</i>	
<i>pellets 10 mg, 100 mg, 20 mg, 30 mg, 50</i>	
<i>mg, 60 mg, 80 mg</i>	318
<i>morphine oral solution 10 mg/5 ml, 20</i>	
<i>mg/5 ml (4 mg/ml)</i>	318
<i>morphine oral tablet</i>	318
<i>morphine oral tablet extended release 100</i>	
<i>mg, 15 mg, 200 mg, 30 mg, 60 mg</i>	318
MOTEGRITY	245
MOTPOLY XR ORAL	
CAPSULE, EXTENDED RELEASE	
24HR 100 MG, 150 MG, 200 MG	246
MOUNJARO	147
MS CONTIN ORAL TABLET	
EXTENDED RELEASE 100 MG, 15	
MG, 200 MG, 30 MG, 60 MG	318
MULPLETA	248
MYALEPT	249
MYCAPSSA	251
MYFEMBREE	252
NALOCET	318
NAMENDA TITRATION PAK	254
NAMENDA XR ORAL	
CAPSULE, SPRINKLE, ER 24HR 14	
MG, 21 MG, 28 MG	254
NAMZARIC	255
<i>naproxen-esomeprazole</i>	451
NATESTO	415
NAYZILAM	256

NERLYNX.....	257	OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG.....	277
NEURONTIN ORAL CAPSULE 100 MG, 300 MG, 400 MG.....	138	OJEMDA ORAL SUSPENSION FOR RECONSTITUTION.....	278
NEURONTIN ORAL SOLUTION.....	138	OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5).....	278
NEURONTIN ORAL TABLET 600 MG, 800 MG.....	138	OJJAARA.....	279
NEXAVAR.....	258	OLPRUVA ORAL PELLETS IN PACKET 2 GRAM, 3 GRAM, 4 GRAM, 5 GRAM, 6 GRAM, 6.67 GRAM.....	280
NEXLETOL.....	259	OLUMIANT.....	281
NEXLIZET.....	259	OMNITROPE.....	152
NGENLA.....	152	OMVOH PEN.....	282
NINLARO.....	261	OMVOH SUBCUTANEOUS.....	282
<i>nitisinone</i>	262	ONFI ORAL SUSPENSION.....	283
NITYR.....	262	ONFI ORAL TABLET.....	283
NORDITROPIN FLEXPRO.....	152	ONGENTYS.....	284
NORGESIC.....	160	ONUREG.....	285
NORGESIC FORTE.....	160	OPSUMIT.....	327
NORLIQVA.....	20	OPSYNVI.....	327
NORTHERA ORAL CAPSULE 100 MG, 200 MG, 300 MG.....	263	OPZELURA.....	286
NOURIANZ.....	264	ORALAIR SUBLINGUAL TABLET 300 INDX REACTIVITY.....	287
NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON.....	312	ORENCIA CLICKJECT.....	288
NOXAFIL ORAL SUSPENSION.....	313	ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML.....	288
NOXAFIL ORAL TABLET,DELAYED RELEASE (DR/EC).....	314	ORENITRAM MONTH 1 TITRATION KT.....	327
NUBEQA.....	265	ORENITRAM MONTH 2 TITRATION KT.....	327
NUCALA SUBCUTANEOUS AUTO- INJECTOR.....	266	ORENITRAM MONTH 3 TITRATION KT.....	327
NUCALA SUBCUTANEOUS RECON SOLN.....	266	ORENITRAM ORAL TABLET EXTENDED RELEASE 0.125 MG, 0.25 MG, 1 MG, 2.5 MG, 5 MG.....	327
NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML.....	266	ORFADIN.....	262
NUCYNTA.....	318	ORGOVYX.....	290
NUCYNTA ER.....	318	ORIAHNN.....	291
NUEDEXTA.....	268	ORILISSA ORAL TABLET 150 MG, 200 MG.....	292
NUPLAZID.....	269	ORKAMBI ORAL GRANULES IN PACKET.....	293
NURTEC ODT.....	270	ORKAMBI ORAL TABLET.....	293
NUTROPIN AQ NUSPIN.....	152	ORLADEYO.....	294
NUVIGIL.....	271	ORMALVI.....	201
OALIVA.....	273		
OCTAGAM.....	172		
<i>octreotide acetate injection solution</i>	274		
ODACTRA.....	275		
ODOMZO.....	276		
OFEV.....	187		

<i>orphenadrine citrate oral</i>	160	PANRETIN	304
<i>orphenadrine-asa-caffeine oral tablet 25-385-30 mg</i>	160	PANZYGA	172
ORSERDU ORAL TABLET 345 MG, 86 MG	296	<i>pazopanib</i>	461
OSMOLEX ER ORAL TABLET, IR - ER, BIPHASIC 24HR 129 MG	297	PEGASYS	183
OSPHERA	298	PEMAZYRE	305
OTEZLA ORAL TABLET 30 MG	299	PERCOCET	318
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)	299	<i>perphenazine-amitriptyline</i>	160
OTREXUP (PF)	247	PHEBURANE	306
<i>oxazepam</i>	318	<i>phenobarbital</i>	162
OXBRYTA ORAL TABLET 300 MG, 500 MG	300	<i>phenoxybenzamine</i>	307
OXBRYTA ORAL TABLET FOR SUSPENSION	301	PHOSPHOLINE IODIDE	308
OXERVATE	302	PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	309
<i>oxycodone oral capsule</i>	318	<i>pirfenidone oral capsule</i>	187
<i>oxycodone oral concentrate</i>	318	<i>pirfenidone oral tablet</i>	187
<i>oxycodone oral solution</i>	318	PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	184
<i>oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg</i>	318	PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	184
<i>oxycodone oral tablet,oral only,ext.rel.12 hr 10 mg, 20 mg</i>	318	PLIAGLIS	423
<i>oxycodone-acetaminophen oral solution 5-325 mg/5 ml</i>	318	POMALYST	310
<i>oxycodone-acetaminophen oral tablet 10-300 mg, 10-325 mg, 2.5-325 mg, 5-300 mg, 5-325 mg, 7.5-300 mg, 7.5-325 mg</i>	318	PONVORY	311
OXYCONTIN ORAL TABLET,ORAL ONLY,EXT.REL.12 HR 10 MG, 15 MG, 20 MG, 30 MG, 40 MG, 60 MG, 80 MG	318	PONVORY 14-DAY STARTER PACK	311
<i>oxymorphone oral tablet</i>	318	<i>posaconazole oral</i>	313, 314
<i>oxymorphone oral tablet extended release 12 hr 10 mg, 15 mg, 20 mg, 30 mg, 40 mg, 5 mg, 7.5 mg</i>	318	PRALUENT PEN	315
OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)	147	<i>pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg</i>	233
OZOBAX DS	34	<i>pregabalin oral solution</i>	233
PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML	303	<i>pregabalin oral tablet extended release 24 hr</i>	233
		PRENATAL VITAMIN PLUS LOW IRON	317
		<i>pretomanid</i>	322
		PRIVIGEN	172
		PROCYSBI ORAL GRANULES DEL RELEASE IN PACKET	323
		PROLASTIN-C INTRAVENOUS SOLUTION	18
		PROLATE ORAL SOLUTION	318
		PROLATE ORAL TABLET	318
		PROLIA	324
		PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG	419

PROMACTA ORAL TABLET 12.5		RETIN-A MICRO PUMP TOPICAL	
MG, 25 MG, 50 MG, 75 MG	419	GEL WITH PUMP 0.06 %, 0.08 %	426
<i>promethazine oral</i>	160	REVATIO ORAL SUSPENSION FOR	
PROVIGIL	325	RECONSTITUTION	327
PRUDOXIN	92	REVATIO ORAL TABLET	327
PULMOZYME	329	REVLIMID	346
<i>pyrimethamine</i>	83	REXULTI ORAL TABLET	27
PYRUKYND ORAL TABLET 20 MG, 5		REZDIFFRA	347
MG, 5 MG (4-WEEK PACK), 50 MG ...	332	REZLIDHIA	348
PYRUKYND ORAL TABLETS,DOSE		REZUROCK	349
PACK	332	RINVOQ ORAL TABLET EXTENDED	
QDOLO	318	RELEASE 24 HR 15 MG, 30 MG, 45	
QELBREE ORAL		MG	351
CAPSULE,EXTENDED RELEASE		RIVFLOZA SUBCUTANEOUS	
24HR 100 MG, 150 MG, 200 MG	333	SOLUTION	353
QINLOCK	334	RIVFLOZA SUBCUTANEOUS	
QUALAQUIN	335	SYRINGE 128 MG/0.8 ML, 160	
<i>quinine sulfate</i>	335	MG/ML	353
QULIPTA	336	ROXICODONE ORAL TABLET 15	
RADICAVA ORS STARTER KIT		MG, 30 MG	318
SUSP	337	ROXYBOND	318
RASUVO (PF)	247	ROZLYTREK ORAL CAPSULE 100	
RAVICTI	338	MG, 200 MG	354
REBIF (WITH ALBUMIN)	184	ROZLYTREK ORAL PELLETS IN	
REBIF REBIDOSE SUBCUTANEOUS		PACKET	354
PEN INJECTOR 22 MCG/0.5 ML, 44		RUBRACA	355
MCG/0.5 ML, 8.8MCG/0.2ML-22		RUCONEST	356
MCG/0.5ML (6)	184	<i>rufinamide</i>	37
REBIF TITRATION PACK	184	RYBELSUS	147
RECORLEV	339	RYDAPT	358
REGRANEX	340	RYVENT	60
RELISTOR ORAL	341	SABRIL	359
RELISTOR SUBCUTANEOUS		SAJAZIR	129
SOLUTION	342	SAMSCA	360
RELISTOR SUBCUTANEOUS		SANDOSTATIN INJECTION	
SYRINGE 12 MG/0.6 ML, 8 MG/0.4		SOLUTION 100 MCG/ML, 50	
ML	342	MCG/ML, 500 MCG/ML	274
REPATHA PUSHTRONEX	343	SAPHRIS	361
REPATHA SURECLICK	343	<i>sapropterin</i>	210
REPATHA SYRINGE	343	SAVELLA	362
RESTORIL	318	SCSEMBLIX ORAL TABLET 100 MG,	
RETEVMO ORAL CAPSULE 40 MG,		20 MG, 40 MG	363
80 MG	345	SECUADO	364
RETIN-A	426	SEGLENTIS	318
RETIN-A MICRO	426	SEROSTIM SUBCUTANEOUS	
		RECON SOLN 4 MG, 5 MG, 6 MG	152

<i>sertraline oral capsule</i>	365	<i>sunitinib malate</i>	389
SEYSARA	366	SUNOSI	387
SIGNIFOR	367	SUTENT	389
<i>sildenafil (pulm.hypertension) oral</i>		SYMDEKO	390
<i>suspension for reconstitution</i>	327	SYMPAZAN	391
<i>sildenafil (pulm.hypertension) oral tablet</i> ..	327	SYMPROIC	392
SILENOR	160	SYNAREL	393
SILIQ	368	SYNDROS	394
SIMLANDI(CF) AUTOINJECTOR	6	TABRECTA	395
SIMPONI SUBCUTANEOUS PEN		<i>tadalafil (pulm. hypertension)</i>	327
INJECTOR 100 MG/ML, 50 MG/0.5		<i>tadalafil oral tablet 2.5 mg, 5 mg</i>	66
ML	369	TADLIQ	327
SIMPONI SUBCUTANEOUS		TAFINLAR ORAL CAPSULE	496
SYRINGE 100 MG/ML, 50 MG/0.5 ML	369	TAFINLAR ORAL TABLET FOR	
SIRTURO	371	SUSPENSION	396
SKYCLARYS	372	TAGRISO	397
SKYRIZI SUBCUTANEOUS PEN		TAKHZYRO SUBCUTANEOUS	
INJECTOR	373	SOLUTION	398
SKYRIZI SUBCUTANEOUS		TAKHZYRO SUBCUTANEOUS	
SYRINGE 150 MG/ML	373	SYRINGE 150 MG/ML, 300 MG/2 ML	
SKYRIZI SUBCUTANEOUS		(150 MG/ML)	398
WEARABLE INJECTOR 180 MG/1.2		TALTZ AUTOINJECTOR	400
ML (150 MG/ML), 360 MG/2.4 ML (150		TALTZ SYRINGE SUBCUTANEOUS	
MG/ML)	373	SYRINGE 80 MG/ML	400
SKYTROFA	374	TALZENNA	401
<i>sodium oxybate</i>	489	TARGRETIN ORAL	402
<i>sodium phenylbutyrate</i>	50	TARGRETIN TOPICAL	402
<i>sofosbuvir-velpatasvir</i>	109	TARPEYO	403
SOGROYA	375	TASCENSO ODT	404
SOHONOS ORAL CAPSULE 1 MG, 1.5		TASIGNA	405
MG, 10 MG, 2.5 MG, 5 MG	376	<i>tasimelteon</i>	406
SOMA	160	TAVALISSE	407
SOMAVERT	378	TAVNEOS	408
<i>sorafenib</i>	258	<i>tazarotene topical cream</i>	410
SOTYKTU	379	<i>tazarotene topical foam</i>	409
SOVALDI	380	<i>tazarotene topical gel</i>	410
SPEVIGO SUBCUTANEOUS	381	TAZORAC TOPICAL CREAM	410
SPORANOX	191	TAZORAC TOPICAL GEL	410
SPRYCEL	382	TAZVERIK	411
STELARA SUBCUTANEOUS		TECFIDERA ORAL	
SOLUTION	384	CAPSULE,DELAYED	
STELARA SUBCUTANEOUS		RELEASE(DR/EC) 120 MG, 120 MG	
SYRINGE 45 MG/0.5 ML, 90 MG/ML ..	384	(14)- 240 MG (46), 240 MG	412
STIVARGA	385	TEGLUTIK	350
STROMEKTOL	192	TEGSEDI	413
SUCRAID	386	<i>temazepam</i>	318

TEPMETKO	414	<i>triazolam</i>	318
<i>teriflunomide</i>	28	<i>trientine oral capsule 500 mg</i>	427
<i>teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)</i>	134	TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	428
TESTIM	415	TRIKAFTA ORAL TABLETS, SEQUENTIAL	428
<i>testosterone cypionate</i>	415	<i>trimethobenzamide oral</i>	421
<i>testosterone enanthate</i>	415	<i>trimipramine</i>	160
<i>testosterone transdermal gel in metered-dose pump</i>	415	TRUDHESA	89
<i>testosterone transdermal gel in packet</i>	415	TRULICITY	147
<i>testosterone transdermal solution in metered pump w/app</i>	415	TRUQAP	429
<i>tetrabenazine oral tablet 12.5 mg, 25 mg</i> ...478		TUKYSA ORAL TABLET 150 MG, 50 MG	430
THALOMID ORAL CAPSULE 100 MG, 150 MG, 200 MG, 50 MG	417	TURALIO ORAL CAPSULE 125 MG ..	431
THIOLA	418	TWYNEO	426
THIOLA EC	418	TYKERB	432
TIBSOVO	420	TYMLOS	433
<i>tiopronin oral tablet</i>	418	TYVASO DPI INHALATION CARTRIDGE WITH INHALER 16 MCG, 16(112)-32(112) -48(28) MCG, 32 MCG, 32-48 MCG, 48 MCG, 64 MCG ...327	
TLANDO	415	UBRELVY ORAL TABLET 100 MG, 50 MG	434
TOBI	63	ULORIC	435
TOBI PODHALER	63	UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG	327
<i>tobramycin in 0.225 % nacl</i>	63	UPTRAVI ORAL TABLETS,DOSE PACK	327
<i>tobramycin inhalation</i>	63	VALCHLOR	436
TOLSURA	422	VALIUM	318
<i>tolvaptan</i>	360	VALTOCO	437
TRACLEER ORAL TABLET	327	VANCOCIN ORAL CAPSULE 125 MG, 250 MG	438
TRACLEER ORAL TABLET FOR SUSPENSION	327	<i>vancomycin oral capsule 125 mg, 250 mg</i> ..	438
<i>tramadol oral capsule,er biphas 24 hr 17-83</i>	318	VANFLYTA	439
<i>tramadol oral capsule,er biphas 24 hr 25-75 100 mg, 200 mg</i>	318	VELSIPITY	440
<i>tramadol oral solution</i>	318	VELTASSA	441
<i>tramadol oral tablet 100 mg, 25 mg, 50 mg</i>	318	VELTIN	426
<i>tramadol oral tablet extended release 24 hr</i>	318	VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG	443
<i>tramadol oral tablet, er multiphas 24 hr</i> ..	318	VENCLEXTA STARTING PACK	443
<i>tramadol-acetaminophen</i>	318	<i>venlafaxine besylate</i>	442
TREMFYA	425	VEOZAH	444
<i>tretinoin</i>	426	VERKAZIA	445
<i>tretinoin microspheres topical gel</i>	426		
<i>tretinoin microspheres topical gel with pump 0.08 %</i>	426		
TREZIX	318		

VERQUVO.....	446	WINLEVI.....	426
VERZENIO.....	447	WINREVAIR.....	472
VFEND IV.....	459	XALKORI ORAL CAPSULE.....	473
VIBERZI.....	448	XALKORI ORAL PELLETT 150 MG, 20	
VICTOZA 3-PAK.....	147	MG, 50 MG.....	473
<i>vigabatrin</i>	359	XANAX ORAL TABLET 0.25 MG, 0.5	
VIGADRONE.....	359	MG, 1 MG, 2 MG.....	318
VIGPODER.....	359	XANAX XR ORAL TABLET	
VIJOICE ORAL GRANULES IN		EXTENDED RELEASE 24 HR 0.5 MG,	
PACKET.....	449	1 MG, 2 MG, 3 MG.....	318
VIJOICE ORAL TABLET 125 MG, 250		XCOPRI.....	474
MG/DAY (200 MG X1-50 MG X1), 50		XCOPRI MAINTENANCE PACK.....	474
MG.....	449	XCOPRI TITRATION PACK.....	474
VIMOVO.....	450	XDEMVI.....	475
VISTARIL ORAL CAPSULE 25 MG....	160	XELJANZ ORAL SOLUTION.....	477
VITRAKVI ORAL CAPSULE 100 MG,		XELJANZ ORAL TABLET.....	476
25 MG.....	452	XELJANZ XR.....	476
VITRAKVI ORAL SOLUTION.....	452	XENAZINE ORAL TABLET 12.5 MG,	
VIVJOA.....	453	25 MG.....	478
VIZIMPRO.....	455	XERMELO.....	479
VOGELXO TRANSDERMAL GEL.....	415	XGEVA.....	480
VOGELXO TRANSDERMAL GEL IN		XIFAXAN ORAL TABLET 550 MG....	481
METERED-DOSE PUMP.....	415	XOLAIR.....	482
VONJO.....	456	XOLREMDI.....	484
VOQUEZNA DUAL PAK.....	458	XOSPATA.....	485
VOQUEZNA ORAL TABLET 10 MG,		XPOVIO ORAL TABLET 100	
20 MG.....	457	MG/WEEK (50 MG X 2), 40	
VOQUEZNA TRIPLE PAK.....	458	MG/WEEK (40 MG X 1), 40MG	
<i>voriconazole intravenous</i>	459	TWICE WEEK (40 MG X 2), 60	
VOSEVI.....	460	MG/WEEK (60 MG X 1), 60MG	
VOTRIENT.....	461	TWICE WEEK (120 MG/WEEK), 80	
VOWST.....	123	MG/WEEK (40 MG X 2), 80MG	
VOXZOGO.....	462	TWICE WEEK (160 MG/WEEK).....	486
VRAYLAR ORAL CAPSULE.....	463	XTAMPZA ER.....	318
VTAMA.....	464	XTANDI ORAL CAPSULE.....	487
VUITY.....	465	XTANDI ORAL TABLET 40 MG, 80	
VUMERITY.....	466	MG.....	487
VYNDAMAX.....	26	XURIDEN.....	488
VYNDAQEL.....	26	XYOSTED.....	415
WAINUA.....	467	XYREM.....	489
WAKIX.....	468	XYWAV.....	490
WEGOVY SUBCUTANEOUS PEN		YARGESA.....	493
INJECTOR 0.25 MG/0.5 ML, 0.5		YONSA.....	492
MG/0.5 ML, 1 MG/0.5 ML, 1.7 MG/0.75		YUFLYMA(CF).....	6
ML, 2.4 MG/0.75 ML.....	470	YUFLYMA(CF) AI CROHN'S-UC-HS.....	6
WELIREG.....	471	YUFLYMA(CF) AUTOINJECTOR.....	6

YUSIMRY(CF) PEN	6
<i>zaleplon oral capsule 10 mg, 5 mg</i>	318
ZAVESCA	493
ZAVZPRET	494
ZEJULA ORAL TABLET	495
ZELBORAF	496
ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG	18
ZEPATIER	497
ZEPOSIA	498
ZEPOSIA STARTER KIT (28-DAY)	498
ZEPOSIA STARTER PACK (7-DAY)	498
ZIANA	426
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML, 23 MG/0.574 ML, 32.4 MG/0.81 ML	250
<i>zileuton</i>	215
ZOKINVY	499
ZOLINZA	500
<i>zolpidem</i>	318, 501
ZOMACTON	152
ZONALON	92
ZONISADE	502
ZORYVE TOPICAL CREAM 0.3 %	503
ZORYVE TOPICAL FOAM	504
ZTALMY	505
ZTLIDO	506
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG	507
ZYDELIG	508
ZYFLO	215
ZYKADIA	509
ZYMFENTRA	510
ZYTIGA ORAL TABLET 250 MG, 500 MG	511

Brand ADHD

Products Affected

- **ADDERALL 20 MG TABLET**
- **ADDERALL 5 MG TABLET**
- **ADDERALL 7.5 MG TABLET**
- **ADDERALL XR 10 MG CAPSULE,EXTENDED RELEASE**
- **ADDERALL XR 15 MG CAPSULE,EXTENDED RELEASE**
- **ADDERALL XR 20 MG CAPSULE,EXTENDED RELEASE**
- **ADDERALL XR 25 MG CAPSULE,EXTENDED RELEASE**
- **ADDERALL XR 30 MG CAPSULE,EXTENDED RELEASE**
- **ADDERALL XR 5 MG CAPSULE,EXTENDED RELEASE**
- **ADZENYS XR-ODT 12.5 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **ADZENYS XR-ODT 15.7 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **ADZENYS XR-ODT 18.8 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **ADZENYS XR-ODT 3.1 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **ADZENYS XR-ODT 6.3 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **ADZENYS XR-ODT 9.4 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **APTENSIO XR 10 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 15 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 20 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 30 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 40 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 50 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **APTENSIO XR 60 MG CAPSULE,EXTENDED RELEASE SPRINKLE**
- **AZSTARYS 26.1 MG-5.2 MG CAPSULE**
- **AZSTARYS 39.2 MG-7.8 MG CAPSULE**
- **AZSTARYS 52.3 MG-10.4 MG CAPSULE**
- **CONCERTA 18 MG TABLET,EXTENDED RELEASE**
- **CONCERTA 27 MG TABLET,EXTENDED RELEASE**
- **CONCERTA 36 MG TABLET,EXTENDED RELEASE**
- **CONCERTA 54 MG TABLET,EXTENDED RELEASE**
- **COTEMPLA XR-ODT 17.3 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **COTEMPLA XR-ODT 25.9 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **COTEMPLA XR-ODT 8.6 MG EXTENDED RELEASE DISINTEGRATING TABLET**
- **DEXEDRINE SPANSULE 10 MG CAPSULE,EXTENDED RELEASE**
- *dextroamphetamine-amphetamine er 12.5 mg capsule, 3 bead, ext rel 24hr*
- *dextroamphetamine-amphetamine er 25 mg capsule,3 bead,ext release 24hr*
- *dextroamphetamine-amphetamine er 37.5 mg capsule, 3 bead, ext rel 24hr*
- *dextroamphetamine-amphetamine er 50 mg capsule,3 bead,ext release 24hr*
- **DYANAVEL XR 10 MG TABLET, EXTENDED RELEASE**

- **DYANAVEL XR 15 MG TABLET, EXTENDED RELEASE**
- **DYANAVEL XR 2.5 MG/ML ORAL 24 HR EXTENDED RELEASE SUSPENSION**
- **DYANAVEL XR 20 MG TABLET, EXTENDED RELEASE**
- **DYANAVEL XR 5 MG TABLET, EXTENDED RELEASE**
- **FOCALIN 10 MG TABLET**
- **FOCALIN 2.5 MG TABLET**
- **FOCALIN 5 MG TABLET**
- **FOCALIN XR 10 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 15 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 20 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 25 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 30 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 35 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 40 MG CAPSULE,EXTENDED RELEASE**
- **FOCALIN XR 5 MG CAPSULE,EXTENDED RELEASE**
- **JORNAY PM 100 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE**
- **JORNAY PM 20 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE**
- **JORNAY PM 40 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE**
- **JORNAY PM 60 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE**
- **JORNAY PM 80 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE**
- **SPRINKLE**
- *lisdexamfetamine 10 mg capsule*
- *lisdexamfetamine 10 mg chewable tablet*
- *lisdexamfetamine 20 mg capsule*
- *lisdexamfetamine 20 mg chewable tablet*
- *lisdexamfetamine 30 mg capsule*
- *lisdexamfetamine 30 mg chewable tablet*
- *lisdexamfetamine 40 mg capsule*
- *lisdexamfetamine 40 mg chewable tablet*
- *lisdexamfetamine 50 mg capsule*
- *lisdexamfetamine 50 mg chewable tablet*
- *lisdexamfetamine 60 mg capsule*
- *lisdexamfetamine 60 mg chewable tablet*
- *lisdexamfetamine 70 mg capsule*
- **METADATE CD 10 MG CAPSULE,EXTENDED RELEASE**
- **METADATE CD 20 MG CAPSULE,EXTENDED RELEASE**
- **METADATE CD 30 MG CAPSULE,EXTENDED RELEASE**
- **METADATE CD 40 MG CAPSULE,EXTENDED RELEASE**
- **METADATE CD 50 MG CAPSULE,EXTENDED RELEASE**
- **METADATE CD 60 MG CAPSULE,EXTENDED RELEASE**
- **METHYLIN 10 MG/5 ML ORAL SOLUTION**
- **METHYLIN 5 MG/5 ML ORAL SOLUTION**
- *methylphenidate er 45 mg tablet,extended release 24 hr*
- *methylphenidate er 63 mg tablet,extended release 24 hr*
- *methylphenidate er 72 mg tablet,extended release 24 hr*
- **MYDAYIS 12.5 MG CAPSULE EXTENDED RELEASE 24 HR**
- **MYDAYIS 25 MG CAPSULE EXTENDED RELEASE 24 HR**
- **MYDAYIS 37.5 MG CAPSULE EXTENDED RELEASE 24 HR**
- **MYDAYIS 50 MG CAPSULE EXTENDED RELEASE 24 HR**
- **QUILLICHEW ER 20 MG CHEWABLE TABLET, EXTENDED RELEASE**
- **QUILLICHEW ER 30 MG CHEWABLE**

- **TABLET, EXTENDED RELEASE**
- **QUILLICHEW ER 40 MG CHEWABLE, EXTENDED RELEASE TABLET**
- **QUILLIVANT XR 5 MG/ML (25 MG/5 ML) ORAL SUSPENSION, EXTENDED RELEASE 24HR**
- **RELEXXII 18 MG TABLET, EXTENDED RELEASE**
- **RELEXXII 27 MG TABLET, EXTENDED RELEASE**
- **RELEXXII 36 MG TABLET, EXTENDED RELEASE**
- **RELEXXII 45 MG TABLET, EXTENDED RELEASE**
- **RELEXXII 63 MG TABLET, EXTENDED RELEASE**
- **RITALIN 10 MG TABLET**
- **RITALIN 20 MG TABLET**
- **RITALIN 5 MG TABLET**
- **RITALIN LA 10 MG CAPSULE, EXTENDED RELEASE**
- **RITALIN LA 20 MG CAPSULE, EXTENDED RELEASE**
- **RITALIN LA 30 MG CAPSULE, EXTENDED RELEASE**
- **RITALIN LA 40 MG CAPSULE, EXTENDED RELEASE**
- **STRATTERA 10 MG CAPSULE**
- **STRATTERA 100 MG CAPSULE**
- **STRATTERA 18 MG CAPSULE**
- **STRATTERA 25 MG CAPSULE**
- **STRATTERA 40 MG CAPSULE**
- **STRATTERA 60 MG CAPSULE**
- **STRATTERA 80 MG CAPSULE**
- **VYVANSE 10 MG CAPSULE**
- **VYVANSE 10 MG CHEWABLE TABLET**
- **VYVANSE 20 MG CAPSULE**
- **VYVANSE 20 MG CHEWABLE TABLET**
- **VYVANSE 30 MG CAPSULE**
- **VYVANSE 30 MG CHEWABLE TABLET**
- **VYVANSE 40 MG CAPSULE**
- **VYVANSE 40 MG CHEWABLE TABLET**
- **VYVANSE 50 MG CAPSULE**
- **VYVANSE 50 MG CHEWABLE TABLET**
- **VYVANSE 60 MG CAPSULE**
- **VYVANSE 60 MG CHEWABLE TABLET**
- **VYVANSE 70 MG CAPSULE**
- **XELSTRYM 13.5 MG/9 HOUR TRANSDERMAL 24 HOUR PATCH**
- **XELSTRYM 18 MG/9 HOUR TRANSDERMAL 24 HOUR PATCH**
- **XELSTRYM 4.5 MG/9 HOUR TRANSDERMAL 24 HOUR PATCH**
- **XELSTRYM 9 MG/9 HOUR TRANSDERMAL 24 HOUR PATCH**

Details

Criteria	Require a 1 month trial of 2 of the following generic ADHD medications (Step 1 drug) when being utilized for the same medically accepted indication: methylphenidate, atomoxetine, dextroamphetamine/amphetamine (generic Adderall/Adderal XR), or dexamethylphenidate in the last 180 days
-----------------	---

Brand Albuterol

Products Affected

- **AIRSUPRA 90 MCG-80 MCG/ACTUATION HFA AEROSOL INHALER**
• *albuterol sulfate hfa 90 mcg/actuation*
- *aerosol inhaler (nda020983)*
- **PROAIR RESPICLICK 90 MCG/ACTUATION BREATH ACTIVATED**

Details

Criteria	Require a 1 month trial of albuterol (generic Proair HFA) in the last 90 days
-----------------	---

Brand Epinephrine

Products Affected

- **AUVI-Q 0.1 MG/0.1 ML INJECTION,AUTO-INJECTOR**
- **AUVI-Q 0.15 MG/0.15 ML AUTO-INJECTOR (FOR 33 LB TO 66 LB PATIENTS)**
- **AUVI-Q 0.3 MG/0.3 ML INJECTION, AUTO-INJECTOR**

Details

Criteria	Require a trial of 2 of the following (Step 1 drug): generic epinephrine injection or EpiPen in the last 180 days
-----------------	---

Brand Glaucoma

Products Affected

- **IYUZEH (PF) 0.005 % EYE DROPS IN A DROPPERETTE**
- **RHOPRESSA 0.02 % EYE DROPS**
- **ROCKLATAN 0.02 %-0.005 % EYE DROPS**
- **VYZULTA 0.024 % EYE DROPS**

Details

Criteria	Require a 1 month trial of generic latanoprost (Step 1 drug) plus one other preferred generic formulary glaucoma drug (Step 1 drug) in the last 180 days
-----------------	--

Brand Levalbuterol

Products Affected

- **XOPENEX HFA 45 MCG/ACTUATION
AEROSOL INHALER**

Details

Criteria	Require a 1 month trial of albuterol (generic Proair HFA) and generic levalbuterol in the last 180 days
-----------------	---

Breo Ellipta AG

Products Affected

- *fluticasone furoate 100 mcg-vilanterol 25 mcg/dose inhalation powder*
- *fluticasone furoate 200 mcg-vilanterol 25 mcg/dose inhalation powder*

Details

Criteria	Require a 1 month trial of Breo Ellipta (Step 1 drug) in the last 90 days
----------	---

Celecoxib

Products Affected

- **CELEBREX 100 MG CAPSULE**
 - **CELEBREX 200 MG CAPSULE**
 - **CELEBREX 400 MG CAPSULE**
 - **CELEBREX 50 MG CAPSULE**
- *celecoxib 100 mg capsule*
 - *celecoxib 200 mg capsule*
 - *celecoxib 400 mg capsule*
 - *celecoxib 50 mg capsule*

Details

Criteria	Require a 1 month trial of 2 formulary generic NSAIDs (Step 1 drug) in the last 180 days
-----------------	--

Clonidine ER

Products Affected

- *clonidine hcl er 0.17 mg tablet, extended release 24 hr*
- **NEXICLON XR 0.17 MG
TABLET, EXTENDED RELEASE**

Details

Criteria	Require a 1 month trial of generic clonidine IR tablet (Step 1 drug) in the last 90 days
-----------------	--

Conjupri

Products Affected

- *levamlodipine 2.5 mg tablet*
- *levamlodipine 5 mg tablet*

Details

Criteria	Require a trial of 2 of the following generic medications (Step 1 drug): amlodipine tablet, felodipine extended-release tablet, or nifedipine 24hr extended-release tablet in the last 180 days
-----------------	---

DPP4 Agents

Products Affected

- *alogliptin 12.5 mg tablet*
- *alogliptin 12.5 mg-metformin 1,000 mg tablet*
- *alogliptin 12.5 mg-metformin 500 mg tablet*
- *alogliptin 12.5 mg-pioglitazone 30 mg tablet*
- *alogliptin 25 mg tablet*
- *alogliptin 25 mg-pioglitazone 15 mg tablet*
- *alogliptin 25 mg-pioglitazone 30 mg tablet*
- *alogliptin 25 mg-pioglitazone 45 mg tablet*
- *alogliptin 6.25 mg tablet*
- **KAZANO 12.5 MG-1,000 MG TABLET**
- **KAZANO 12.5 MG-500 MG TABLET**
- **NESINA 12.5 MG TABLET**
- **NESINA 25 MG TABLET**
- **NESINA 6.25 MG TABLET**
- **OSENI 12.5 MG-30 MG TABLET**
- **OSENI 25 MG-15 MG TABLET**
- **OSENI 25 MG-30 MG TABLET**
- **OSENI 25 MG-45 MG TABLET**
- *saxagliptin 2.5 mg tablet*
- *saxagliptin 2.5 mg-metformin er 1,000 mg tablet,extend release 24hr mp*
- *saxagliptin 5 mg tablet*
- *saxagliptin 5 mg-metformin er 1,000 mg tablet,extend release 24hr mp*
- *saxagliptin 5 mg-metformin er 500 mg tablet,extend release 24hr mp*
- *sitagliptin 100 mg tablet*
- *sitagliptin 25 mg tablet*
- *sitagliptin 50 mg tablet*
- *sitagliptin 50 mg-metformin 1,000 mg tablet*
- *sitagliptin 50 mg-metformin 500 mg tablet*
- **ZITUVIO 100 MG TABLET**
- **ZITUVIO 25 MG TABLET**
- **ZITUVIO 50 MG TABLET**

Details

Criteria	Require a 1 month trial of a Januvia/Janumet/Janumet XR product and Tradjenta/Jentadueto/Jentadueto XR product in the last 180 days
-----------------	---

DPP4-SGLT2 Combo

Products Affected

- STEGLUJAN 15 MG-100 MG TABLET
- STEGLUJAN 5 MG-100 MG TABLET

Details

Criteria	Require a 1 month trial of 2 of the following (Step 1 drug): Glyxambi, Trijardy or Qtern in the last 180 days
-----------------	---

Dry Eye

Products Affected

- **CEQUA 0.09 % EYE DROPS IN A DROPPERETTE**
- **MIEBO (PF) 100 % EYE DROPS**
- **TYRVAYA 0.03 MG/SPRAY NASAL SPRAY**
- **VEVYE 0.1 % EYE DROPS**

Details

Criteria	Require a 1 month trial of Xiidra and one of the following: Restasis or cyclosporine eye drop (i.e. generic Restasis), in the last 180 days
-----------------	---

Dymista

Products Affected

- **DYMISTA 137 MCG-50 MCG/SPRAY NASAL SPRAY**
- **RYALTRIS 665 MCG-25 MCG/SPRAY NASAL SPRAY**

Details

Criteria	Require a 1 month trial of generic azelastine/fluticasone spray in the last 90 days
-----------------	---

Erythroid Stimulants

Products Affected

- ARANESP 10 MCG/0.4 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 100 MCG/0.5 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 100 MCG/ML (IN POLYSORBATE) INJECTION
- ARANESP 150 MCG/0.3 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 200 MCG/0.4 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 200 MCG/ML (IN POLYSORBATE) INJECTION
- ARANESP 25 MCG/0.42 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 25 MCG/ML (IN POLYSORBATE) INJECTION
- ARANESP 300 MCG/0.6 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 40 MCG/0.4 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 40 MCG/ML (IN POLYSORBATE) INJECTION
- ARANESP 500 MCG/ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 60 MCG/0.3 ML (IN POLYSORBATE) INJECTION SYRINGE
- ARANESP 60 MCG/ML (IN POLYSORBATE) INJECTION
- EPOGEN 2,000 UNIT/ML INJECTION SOLUTION
- EPOGEN 20,000 UNIT/2 ML INJECTION SOLUTION
- EPOGEN 20,000 UNIT/ML INJECTION SOLUTION
- EPOGEN 3,000 UNIT/ML INJECTION SOLUTION
- EPOGEN 4,000 UNIT/ML INJECTION SOLUTION

Details

Criteria	Require a 1 month trial of Procrit and Retacrit (Step 1 drugs) in the last 180 days
----------	---

Filgrastim

Products Affected

- **GRANIX 300 MCG/0.5 ML SUBCUTANEOUS SYRINGE**
- **GRANIX 300 MCG/ML SUBCUTANEOUS SOLUTION**
- **GRANIX 480 MCG/0.8 ML SUBCUTANEOUS SYRINGE**
- **GRANIX 480 MCG/1.6 ML SUBCUTANEOUS SOLUTION**
- **NEUPOGEN 300 MCG/0.5 ML INJECTION SYRINGE**
- **NEUPOGEN 300 MCG/ML INJECTION SOLUTION**
- **NEUPOGEN 480 MCG/0.8 ML INJECTION SYRINGE**
- **NEUPOGEN 480 MCG/1.6 ML INJECTION SOLUTION**
- **RELEUKO 300 MCG/0.5 ML SUBCUTANEOUS SYRINGE**
- **RELEUKO 480 MCG/0.8 ML SUBCUTANEOUS SYRINGE**

Details

Criteria	Require a 1 month trial of Zarxio and Nivestym (step 1 drugs) in the last 180 days
-----------------	--

Fortamet

Products Affected

- *metformin er 1,000 mg tablet, extended release 24hr (osmotic)*
- *metformin er 500 mg tablet, extended release 24hr (osmotic)*

Details

Criteria	Require a 1 month trial of generic metformin IR tablets and generic metformin ER (i.e. generic Glucophage XR) tablets in the last 180 days
-----------------	--

Glumetza

Products Affected

- **GLUMETZA 1,000 MG TABLET,EXTENDED RELEASE**
 - **GLUMETZA 500 MG TABLET,EXTENDED RELEASE**
- *metformin er 1,000 mg 24 hr tablet,extended release (gastric reten.)*
 - *metformin er 500 mg 24 hr tablet,extended release (gastric retention)*

Details

Criteria	Require a 1 month trial of generic metformin IR tablets and generic metformin ER (i.e. generic Glucophage XR) tablets in the last 180 days
-----------------	--

GnRH Agonists

Products Affected

- *leuprolide 22.5 mg (3 month) intramuscular suspension*
- **LUPRON DEPOT 11.25 MG (3 MONTH) INTRAMUSCULAR SYRINGE KIT**
- **LUPRON DEPOT 22.5 MG (3 MONTH) INTRAMUSCULAR SYRINGE KIT**
- **LUPRON DEPOT 3.75 MG INTRAMUSCULAR SYRINGE KIT**
- **LUPRON DEPOT 30 MG (4 MONTH) INTRAMUSCULAR SYRINGE KIT**
- **LUPRON DEPOT 45 MG (6 MONTH) INTRAMUSCULAR SYRINGE KIT**
- **LUPRON DEPOT 7.5 MG INTRAMUSCULAR SYRINGE KIT**
- **TRELSTAR 11.25 MG IM SUSPENSION**
- **TRELSTAR 22.5 MG IM SUSPENSION**
- **TRELSTAR 3.75 MG IM SUSPENSION**

Details

Criteria	Require a trial of Eligard (Step 1 drug) in the last 180 days when being utilized for the same medically accepted indication
----------	--

Herpetic Keratitis

Products Affected

- ZIRGAN 0.15 % EYE GEL

Details

Criteria	Require a 1 month trial of generic trifluridine eye drops (Step 1 drug) in the last 90 days
-----------------	---

Inhaled LAMA

Products Affected

- **INCRUSE ELLIPTA 62.5 MCG/ACTUATION POWDER FOR INHALATION**
- **TUDORZA PRESSAIR 400 MCG/ACTUATION BREATH**
- **ACTIVATED TUDORZA PRESSAIR 400 MCG/ACTUATION BREATH ACTIVATED (30 ACTUAT)**

Details

Criteria	Require a 1 month trial of Spiriva (Step 1 drug) in the last 90 days
-----------------	--

Lactulose

Products Affected

- **KRISTALOSE 10 GRAM ORAL PACKET**
 - **KRISTALOSE 20 GRAM ORAL PACKET**
- *lactulose 10 gram oral packet*

Details

Criteria	Require a 1 month trial of lactulose solution (Step 1 drug) in the last 90 days
-----------------	---

Mirabegron ER

Products Affected

- *mirabegron er 25 mg tablet, extended release 24 hr*
- *mirabegron er 50 mg tablet, extended release 24 hr*

Details

Criteria	Require a 1 month trial of Myrbetriq (Step 1 drug) in the last 90 days
-----------------	--

Mupirocin

Products Affected

- *mupirocin calcium 2 % topical cream*

Details

Criteria	Require a 1 month trial of generic mupirocin ointment (Step 1 drug) in the last 90 days
-----------------	---

Pegfilgrastim

Products Affected

- **FYLNETRA 6 MG/0.6 ML
SUBCUTANEOUS SYRINGE**
- **NYVEPRIA 6 MG/0.6 ML
SUBCUTANEOUS SYRINGE**
- **STIMUFEND 6 MG/0.6 ML
SUBCUTANEOUS SYRINGE**
- **UDENYCA 6 MG/0.6 ML
SUBCUTANEOUS SYRINGE**
- **UDENYCA AUTOINJECTOR 6 MG/0.6
ML SUBCUTANEOUS AUTO-
INJECTOR**

Details

Criteria	Require a trial of 2 of the following (Step 1 drug): Neulasta, Fulphila, or Ziextenzo in the last 180 days
-----------------	--

Pennsaid

Products Affected

- *diclofenac 20 mg/gram/actuation (2 %) topical soln metered-dose pump* **(2 %) TOPICAL SOLN IN METERED-DOSE PUMP**
- **PENNSAID 20 MG/GRAM/ACTUATION**

Details

Criteria	Require a 1 month trial of Diclofenac 1.5% (Step 1 drug) in the last 90 days
-----------------	--

Rapid Acting Insulin

Products Affected

- **ADMELOG SOLOSTAR U-100 INSULIN LISPRO 100 UNIT/ML SUBCUTANEOUS PEN**
- **ADMELOG U-100 INSULIN LISPRO 100 UNIT/ML SUBCUTANEOUS SOLUTION**
- **APIDRA SOLOSTAR U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS PEN**
- **APIDRA U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION**
- **FIASP FLEXTOUCH U-100 INSULIN 100 UNIT/ML (3 ML) SUBCUTANEOUS PEN**
- **FIASP PENFILL U-100 INSULIN 100 UNIT/ML (3 ML) SUBCUTANEOUS CARTRIDGE**
- **FIASP U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION**
- *insulin aspar prot-insulin aspart 100 unit/ml (70-30) subcutaneous pen*
- *insulin aspar prt-insulin aspart 100 unit/ml (70-30) subcutaneous soln*
- *insulin aspart (u-100) 100 unit/ml (3 ml) subcutaneous pen*
- *insulin aspart u-100 100 unit/ml subcutaneous cartridge*
- *insulin aspart u-100 100 unit/ml subcutaneous solution*
- **LYUMJEV KWIKPEN U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS**
- **LYUMJEV KWIKPEN U-200 INSULIN 200 UNIT/ML (3 ML) SUBCUTANEOUS**
- **LYUMJEV TEMPO PEN (U-100) INSULIN 100 UNIT/ML SUBCUTANEOUS PEN, SENSOR**
- **LYUMJEV U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION**
- **NOVOLOG FLEXPEN U-100 INSULIN ASPART 100 UNIT/ML (3 ML) SUBCUTANEOUS**
- **NOVOLOG MIX 70-30 FLEXPEN U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS PEN**
- **NOVOLOG MIX 70-30 U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION**
- **NOVOLOG PENFILL U-100 INSULIN ASPART 100 UNIT/ML SUBCUTANEOUS CARTRIDG**
- **NOVOLOG U-100 INSULIN ASPART 100 UNIT/ML SUBCUTANEOUS SOLUTION**

Details

Criteria	Require a 1 month trial of Humalog or insulin lispro product (Step 1 drug) in the last 90 days when utilized for the same medically accepted indication
-----------------	---

Riomet

Products Affected

- *metformin 500 mg/5 ml oral solution*

Details

Criteria	Require a 1 month trial of generic metformin IR tablets in the last 90 days -OR- documentation supporting the inability to swallow or difficulty swallowing tablets containing metformin
-----------------	---

Rytary

Products Affected

- **DHIVY 25 MG-100 MG TABLET**
- **RYTARY 23.75 MG-95 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 36.25 MG-145 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 48.75 MG-195 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 61.25 MG-245 MG CAPSULE,EXTENDED RELEASE**

Details

Criteria	Require a trial of generic carbidopa/levodopa product (Step 1 drug) in the last 90 days
-----------------	---

Short Acting Insulin

Products Affected

- **NOVOLIN 70/30 U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SUSPENSION**
- **NOVOLIN 70-30 FLEXPEN U-100 INSULIN 100 UNIT/ML (70-30) SUBCUTANEOUS**
- **NOVOLIN N FLEXPEN 100 UNIT/ML (3 ML) SUBCUTANEOUS INSULIN PEN**
- **NOVOLIN N NPH U-100 INSULIN ISOPHANE 100 UNIT/ML SUBCUTANEOUS SUSP**
- **NOVOLIN R FLEXPEN 100 UNIT/ML (3 ML) SUBCUTANEOUS INSULIN PEN**
- **NOVOLIN R REGULAR U-100 INSULIN 100 UNIT/ML INJECTION SOLUTION**

Details

Criteria	Require a 1 month trial of Humulin product (Step 1 drug) in the last 90 days
-----------------	--

Sleeper Meds

Products Affected

- **BELSOMRA 10 MG TABLET**
- **BELSOMRA 15 MG TABLET**
- **BELSOMRA 20 MG TABLET**
- **BELSOMRA 5 MG TABLET**
- **DAYVIGO 10 MG TABLET**
- **DAYVIGO 5 MG TABLET**
- **QUVIVIQ 25 MG TABLET**
- **QUVIVIQ 50 MG TABLET**

Details

Criteria	Require a trial of 2 of the following (Step 1 drug): generic zolpidem tablet/zolpidem ER tablet, zaleplon or eszopiclone in the last 180 days
-----------------	---

Soanz

Products Affected

- SOAANZ 20 MG TABLET
- SOAANZ 40 MG TABLET
- SOAANZ 60 MG TABLET

Details

Criteria	Require a 1 month trial of generic furosemide tablets and generic torsemide tablets in the last 180 days
-----------------	--

Suboxone

Products Affected

- **SUBOXONE 12 MG-3 MG SUBLINGUAL FILM**
- **SUBOXONE 2 MG-0.5 MG SUBLINGUAL FILM**
- **SUBOXONE 4 MG-1 MG SUBLINGUAL FILM**
- **SUBOXONE 8 MG-2 MG SUBLINGUAL FILM**

Details

Criteria	Require a 1 month trial of Zubsolv (Step 1 drug) in the last 90 days
-----------------	--

Symbicort

Products Affected

- **SYMBICORT 160 MCG-4.5 MCG/ACTUATION HFA AEROSOL INHALER**
- **SYMBICORT 80 MCG-4.5 MCG/ACTUATION HFA AEROSOL INHALER**

Details

Criteria	Require a 1 month trial of budesonide-formoterol or Breyna (Step 1 drug) in the last 90 days
-----------------	--

Topical Antifungals

Products Affected

- **ERTACZO 2 % TOPICAL CREAM**
- **EXELDERM 1 % TOPICAL CREAM**
- **EXELDERM 1 % TOPICAL SOLUTION**
- **KETODAN 2 % TOPICAL FOAM**
- *naftifine 1 % topical cream*
- *naftifine 2 % topical cream*
- *naftifine 2 % topical gel*
- **NAFTIN 1 % TOPICAL GEL**
- **NAFTIN 2 % TOPICAL GEL**
- *oxiconazole 1 % topical cream*
- **OXISTAT 1 % LOTION**
- **OXISTAT 1 % TOPICAL CREAM**

Details

Criteria	Require a 1 month trial of generic econazole cream and one of the following: generic ketoconazole cream or ketoconazole shampoo (Step 1 drugs), when being utilized for the same medically accepted indication, in the last 180 days
-----------------	--

Topical Psoriasis

Products Affected

- *calcipotriene 0.005 % topical foam*
- *calcipotriene-betamethasone 0.005 %-0.064 % topical ointment*
- *calcipotriene-betamethasone 0.005 %-0.064 % topical suspension*
- *calcitriol 3 mcg/gram topical ointment*
- **ENSTILAR 0.005 %-0.064 % TOPICAL FOAM**
- **SORILUX 0.005 % TOPICAL FOAM**
- **TACLONEX 0.005 %-0.064 % TOPICAL OINTMENT**
- **TACLONEX 0.005 %-0.064 % TOPICAL SUSPENSION**
- **VECTICAL 3 MCG/GRAM TOPICAL OINTMENT**

Details

Criteria	Require a 1 month trial of generic calcipotriene cream, ointment or solution (Step 1 drug) in the last 90 days when utilized for the same medically accepted indication
-----------------	---

Ursodiol

Products Affected

- **RELTONE 200 MG CAPSULE**
- **RELTONE 400 MG CAPSULE**
- **URSO 250 250 MG TABLET**
- **URSO FORTE 500 MG TABLET**
- *ursodiol 200 mg capsule*
- *ursodiol 400 mg capsule*

Details

Criteria	Require a 1 month trial of generic ursodiol 250mg tablet, 300mg capsule or 500mg tablet (step 1 drug) in the last 90 days
-----------------	---

Index of Drugs

ADDERALL 20 MG TABLET1	<i>alogliptin 12.5 mg-metformin 1,000 mg</i>
ADDERALL 5 MG TABLET1	<i>tablet</i> 12
ADDERALL 7.5 MG TABLET1	<i>alogliptin 12.5 mg-metformin 500 mg</i>
ADDERALL XR 10 MG	<i>tablet</i> 12
CAPSULE,EXTENDED RELEASE1	<i>alogliptin 12.5 mg-pioglitazone 30 mg</i>
ADDERALL XR 15 MG	<i>tablet</i> 12
CAPSULE,EXTENDED RELEASE1	<i>alogliptin 25 mg tablet</i>12
ADDERALL XR 20 MG	<i>alogliptin 25 mg-pioglitazone 15 mg tablet</i> ..12
CAPSULE,EXTENDED RELEASE1	<i>alogliptin 25 mg-pioglitazone 30 mg tablet</i> ..12
ADDERALL XR 25 MG	<i>alogliptin 25 mg-pioglitazone 45 mg tablet</i> ..12
CAPSULE,EXTENDED RELEASE1	<i>alogliptin 6.25 mg tablet</i>12
ADDERALL XR 30 MG	APIDRA SOLOSTAR U-100 INSULIN
CAPSULE,EXTENDED RELEASE1	100 UNIT/ML SUBCUTANEOUS PEN ...28
ADDERALL XR 5 MG	APIDRA U-100 INSULIN 100 UNIT/ML
CAPSULE,EXTENDED RELEASE1	SUBCUTANEOUS SOLUTION 28
ADMELOG SOLOSTAR U-100	APTENSIO XR 10 MG
INSULIN LISPRO 100 UNIT/ML	CAPSULE,EXTENDED RELEASE
SUBCUTANEOUS PEN 28	SPRINKLE 1
ADMELOG U-100 INSULIN LISPRO	APTENSIO XR 15 MG
100 UNIT/ML SUBCUTANEOUS	CAPSULE,EXTENDED RELEASE
SOLUTION 28	SPRINKLE 1
ADZENYS XR-ODT 12.5 MG	APTENSIO XR 20 MG
EXTENDED RELEASE	CAPSULE,EXTENDED RELEASE
DISINTEGRATING TABLET 1	SPRINKLE 1
ADZENYS XR-ODT 15.7 MG	APTENSIO XR 30 MG
EXTENDED RELEASE	CAPSULE,EXTENDED RELEASE
DISINTEGRATING TABLET 1	SPRINKLE 1
ADZENYS XR-ODT 18.8 MG	APTENSIO XR 40 MG
EXTENDED RELEASE	CAPSULE,EXTENDED RELEASE
DISINTEGRATING TABLET 1	SPRINKLE 1
ADZENYS XR-ODT 3.1 MG	APTENSIO XR 50 MG
EXTENDED RELEASE	CAPSULE,EXTENDED RELEASE
DISINTEGRATING TABLET 1	SPRINKLE 1
ADZENYS XR-ODT 6.3 MG	APTENSIO XR 60 MG
EXTENDED RELEASE	CAPSULE,EXTENDED RELEASE
DISINTEGRATING TABLET 1	SPRINKLE 1
ADZENYS XR-ODT 9.4 MG	ARANESP 10 MCG/0.4 ML (IN
EXTENDED RELEASE	POLYSORBATE) INJECTION
DISINTEGRATING TABLET 1	SYRINGE 16
AIRSUPRA 90 MCG-80	ARANESP 100 MCG/0.5 ML (IN
MCG/ACTUATION HFA AEROSOL	POLYSORBATE) INJECTION
INHALER 4	SYRINGE 16
<i>albuterol sulfate hfa 90 mcg/actuation</i>	ARANESP 100 MCG/ML (IN
<i>aerosol inhaler (nda020983)</i>4	POLYSORBATE) INJECTION 16
<i>alogliptin 12.5 mg tablet</i>12	

ARANESP 150 MCG/0.3 ML (IN POLYSORBATE) INJECTION SYRINGE	16	<i>calcipotriene 0.005 % topical foam</i>	37
ARANESP 200 MCG/0.4 ML (IN POLYSORBATE) INJECTION SYRINGE	16	<i>calcipotriene-betamethasone 0.005 %-0.064 % topical ointment</i>	37
ARANESP 200 MCG/ML (IN POLYSORBATE) INJECTION	16	<i>calcipotriene-betamethasone 0.005 %-0.064 % topical suspension</i>	37
ARANESP 25 MCG/0.42 ML (IN POLYSORBATE) INJECTION SYRINGE	16	<i>calcitriol 3 mcg/gram topical ointment</i>	37
ARANESP 25 MCG/ML (IN POLYSORBATE) INJECTION	16	CELEBREX 100 MG CAPSULE	9
ARANESP 300 MCG/0.6 ML (IN POLYSORBATE) INJECTION SYRINGE	16	CELEBREX 200 MG CAPSULE	9
ARANESP 40 MCG/0.4 ML (IN POLYSORBATE) INJECTION SYRINGE	16	CELEBREX 400 MG CAPSULE	9
ARANESP 40 MCG/ML (IN POLYSORBATE) INJECTION	16	CELEBREX 50 MG CAPSULE	9
ARANESP 500 MCG/ML (IN POLYSORBATE) INJECTION SYRINGE	16	<i>celecoxib 100 mg capsule</i>	9
ARANESP 60 MCG/0.3 ML (IN POLYSORBATE) INJECTION SYRINGE	16	<i>celecoxib 200 mg capsule</i>	9
ARANESP 60 MCG/ML (IN POLYSORBATE) INJECTION	16	<i>celecoxib 400 mg capsule</i>	9
AUVI-Q 0.1 MG/0.1 ML INJECTION,AUTO-INJECTOR	5	<i>celecoxib 50 mg capsule</i>	9
AUVI-Q 0.15 MG/0.15 ML AUTO-INJECTOR (FOR 33 LB TO 66 LB PATIENTS)	5	CEQUA 0.09 % EYE DROPS IN A DROPPERETTE	14
AUVI-Q 0.3 MG/0.3 ML INJECTION, AUTO-INJECTOR	5	<i>clonidine hcl er 0.17 mg tablet,extended release 24 hr</i>	10
AZSTARYS 26.1 MG-5.2 MG CAPSULE	1	CONCERTA 18 MG TABLET,EXTENDED RELEASE	1
AZSTARYS 39.2 MG-7.8 MG CAPSULE	1	CONCERTA 27 MG TABLET,EXTENDED RELEASE	1
AZSTARYS 52.3 MG-10.4 MG CAPSULE	1	CONCERTA 36 MG TABLET,EXTENDED RELEASE	1
BELSOMRA 10 MG TABLET	32	CONCERTA 54 MG TABLET,EXTENDED RELEASE	1
BELSOMRA 15 MG TABLET	32	COTEMPLA XR-ODT 17.3 MG EXTENDED RELEASE	
BELSOMRA 20 MG TABLET	32	DISINTEGRATING TABLET	1
BELSOMRA 5 MG TABLET	32	COTEMPLA XR-ODT 25.9 MG EXTENDED RELEASE	
		DISINTEGRATING TABLET	1
		COTEMPLA XR-ODT 8.6 MG EXTENDED RELEASE	
		DISINTEGRATING TABLET	1
		DAYVIGO 10 MG TABLET	32
		DAYVIGO 5 MG TABLET	32
		DEXEDRINE SPANSULE 10 MG CAPSULE,EXTENDED RELEASE	1
		<i>dextroamphetamine-amphetamine er 12.5 mg capsule, 3 bead, ext rel 24hr</i>	1
		<i>dextroamphetamine-amphetamine er 25 mg capsule,3 bead,ext release 24hr</i>	1
		<i>dextroamphetamine-amphetamine er 37.5 mg capsule, 3 bead, ext rel 24hr</i>	1

<i>dextroamphetamine-amphetamine er 50 mg capsule,3 bead,ext release 24hr</i>	1	FOCALIN 10 MG TABLET	1
DHIVY 25 MG-100 MG TABLET	30	FOCALIN 2.5 MG TABLET	1
<i>diclofenac 20 mg/gram/actuation (2 %)</i>		FOCALIN 5 MG TABLET	1
<i>topical soln metered-dose pump</i>	27	FOCALIN XR 10 MG CAPSULE,EXTENDED RELEASE	1
DYANAVEL XR 10 MG TABLET, EXTENDED RELEASE	1	FOCALIN XR 15 MG CAPSULE,EXTENDED RELEASE	1
DYANAVEL XR 15 MG TABLET, EXTENDED RELEASE	1	FOCALIN XR 20 MG CAPSULE,EXTENDED RELEASE	1
DYANAVEL XR 2.5 MG/ML ORAL 24 HR EXTENDED RELEASE SUSPENSION	1	FOCALIN XR 25 MG CAPSULE,EXTENDED RELEASE	1
DYANAVEL XR 20 MG TABLET, EXTENDED RELEASE	1	FOCALIN XR 30 MG CAPSULE,EXTENDED RELEASE	1
DYANAVEL XR 5 MG TABLET, EXTENDED RELEASE	1	FOCALIN XR 35 MG CAPSULE,EXTENDED RELEASE	1
DYMISTA 137 MCG-50 MCG/SPRAY NASAL SPRAY	15	FOCALIN XR 40 MG CAPSULE,EXTENDED RELEASE	1
ENSTILAR 0.005 %-0.064 % TOPICAL FOAM	37	FOCALIN XR 5 MG CAPSULE,EXTENDED RELEASE	1
EPOGEN 2,000 UNIT/ML INJECTION SOLUTION	16	FYLNETRA 6 MG/0.6 ML SUBCUTANEOUS SYRINGE	26
EPOGEN 20,000 UNIT/2 ML INJECTION SOLUTION	16	GLUMETZA 1,000 MG TABLET,EXTENDED RELEASE	19
EPOGEN 20,000 UNIT/ML INJECTION SOLUTION	16	GLUMETZA 500 MG TABLET,EXTENDED RELEASE	19
EPOGEN 3,000 UNIT/ML INJECTION SOLUTION	16	GRANIX 300 MCG/0.5 ML SUBCUTANEOUS SYRINGE	17
EPOGEN 4,000 UNIT/ML INJECTION SOLUTION	16	GRANIX 300 MCG/ML SUBCUTANEOUS SOLUTION	17
ERTACZO 2 % TOPICAL CREAM	36	GRANIX 480 MCG/0.8 ML SUBCUTANEOUS SYRINGE	17
EXELDERM 1 % TOPICAL CREAM	36	GRANIX 480 MCG/1.6 ML SUBCUTANEOUS SOLUTION	17
EXELDERM 1 % TOPICAL SOLUTION	36	INCRUSE ELLIPTA 62.5 MCG/ACTUATION POWDER FOR INHALATION	22
FIASP FLEXTOUCH U-100 INSULIN 100 UNIT/ML (3 ML) SUBCUTANEOUS PEN	28	<i>insulin aspar prot-insulin aspart 100 unit/ml (70-30) subcutaneous pen</i>	28
FIASP PENFILL U-100 INSULIN 100 UNIT/ML (3 ML) SUBCUTANEOUS CARTRIDGE	28	<i>insulin aspar prt-insulin aspart 100 unit/ml (70-30) subcutaneous soln</i>	28
FIASP U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION	28	<i>insulin aspart (u-100) 100 unit/ml (3 ml) subcutaneous pen</i>	28
<i>fluticasone furoate 100 mcg-vilanterol 25 mcg/dose inhalation powder</i>	8	<i>insulin aspart u-100 100 unit/ml subcutaneous cartridge</i>	28
<i>fluticasone furoate 200 mcg-vilanterol 25 mcg/dose inhalation powder</i>	8	<i>insulin aspart u-100 100 unit/ml subcutaneous solution</i>	28

IYUZEH (PF) 0.005 % EYE DROPS IN A DROPPERETTE	6	<i>lisdexamfetamine 70 mg capsule</i>	1
JORNAY PM 100 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE	1	LUPRON DEPOT 11.25 MG (3 MONTH) INTRAMUSCULAR SYRINGE KIT	20
JORNAY PM 20 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE	1	LUPRON DEPOT 22.5 MG (3 MONTH) INTRAMUSCULAR SYRINGE KIT	20
JORNAY PM 40 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE	1	LUPRON DEPOT 3.75 MG INTRAMUSCULAR SYRINGE KIT	20
JORNAY PM 60 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE	1	LUPRON DEPOT 30 MG (4 MONTH) INTRAMUSCULAR SYRINGE KIT	20
JORNAY PM 80 MG CAPSULE,DELAYED RELEASE,EXTENDED RELEASE SPRINKLE	1	LUPRON DEPOT 45 MG (6 MONTH) INTRAMUSCULAR SYRINGE KIT	20
KAZANO 12.5 MG-1,000 MG TABLET	12	LUPRON DEPOT 7.5 MG INTRAMUSCULAR SYRINGE KIT	20
KAZANO 12.5 MG-500 MG TABLET	12	LYUMJEV KWIKPEN U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS	28
KETODAN 2 % TOPICAL FOAM	36	LYUMJEV KWIKPEN U-200 INSULIN 200 UNIT/ML (3 ML) SUBCUTANEOUS	28
KRISTALOSE 10 GRAM ORAL PACKET	23	LYUMJEV TEMPO PEN (U-100) INSULIN 100 UNIT/ML SUBCUTANEOUS PEN, SENSOR	28
KRISTALOSE 20 GRAM ORAL PACKET	23	LYUMJEV U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION	28
<i>lactulose 10 gram oral packet</i>	23	METADATE CD 10 MG CAPSULE,EXTENDED RELEASE	1
<i>leuprolide 22.5 mg (3 month) intramuscular suspension</i>	20	METADATE CD 20 MG CAPSULE,EXTENDED RELEASE	1
<i>levamlodipine 2.5 mg tablet</i>	11	METADATE CD 30 MG CAPSULE,EXTENDED RELEASE	1
<i>levamlodipine 5 mg tablet</i>	11	METADATE CD 40 MG CAPSULE,EXTENDED RELEASE	1
<i>lisdexamfetamine 10 mg capsule</i>	1	METADATE CD 50 MG CAPSULE,EXTENDED RELEASE	1
<i>lisdexamfetamine 10 mg chewable tablet</i>	1	METADATE CD 60 MG CAPSULE,EXTENDED RELEASE	1
<i>lisdexamfetamine 20 mg capsule</i>	1	<i>metformin 500 mg/5 ml oral solution</i>	29
<i>lisdexamfetamine 20 mg chewable tablet</i>	1	<i>metformin er 1,000 mg 24 hr tablet,extended release (gastric reten.)</i>	19
<i>lisdexamfetamine 30 mg capsule</i>	1	<i>metformin er 1,000 mg tablet,extended release 24hr (osmotic)</i>	18
<i>lisdexamfetamine 30 mg chewable tablet</i>	1	<i>metformin er 500 mg 24 hr tablet,extended release (gastric retention)</i>	19
<i>lisdexamfetamine 40 mg capsule</i>	1	<i>metformin er 500 mg tablet,extended release 24hr (osmotic)</i>	18
<i>lisdexamfetamine 40 mg chewable tablet</i>	1		
<i>lisdexamfetamine 50 mg capsule</i>	1		
<i>lisdexamfetamine 50 mg chewable tablet</i>	1		
<i>lisdexamfetamine 60 mg capsule</i>	1		
<i>lisdexamfetamine 60 mg chewable tablet</i>	1		

METHYLIN 10 MG/5 ML ORAL SOLUTION	1	NOVOLIN 70-30 FLEXPEN U-100 INSULIN 100 UNIT/ML (70-30) SUBCUTANEOUS	31
METHYLIN 5 MG/5 ML ORAL SOLUTION	1	NOVOLIN N FLEXPEN 100 UNIT/ML (3 ML) SUBCUTANEOUS INSULIN PEN	31
<i>methylphenidate er 45 mg tablet,extended release 24 hr</i>	1	NOVOLIN N NPH U-100 INSULIN ISOPHANE 100 UNIT/ML SUBCUTANEOUS SUSP	31
<i>methylphenidate er 63 mg tablet,extended release 24 hr</i>	1	NOVOLIN R FLEXPEN 100 UNIT/ML (3 ML) SUBCUTANEOUS INSULIN PEN	31
<i>methylphenidate er 72 mg tablet,extended release 24 hr</i>	1	NOVOLIN R REGULAR U-100 INSULIN 100 UNIT/ML INJECTION SOLUTION	31
MIEBO (PF) 100 % EYE DROPS	14	NOVOLOG FLEXPEN U-100 INSULIN ASPART 100 UNIT/ML (3 ML) SUBCUTANEOUS	28
<i>mirabegron er 25 mg tablet,extended release 24 hr</i>	24	NOVOLOG MIX 70-30 FLEXPEN U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS PEN	28
<i>mirabegron er 50 mg tablet,extended release 24 hr</i>	24	NOVOLOG MIX 70-30 U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SOLUTION	28
<i>mupirocin calcium 2 % topical cream</i>	25	NOVOLOG PENFILL U-100 INSULIN ASPART 100 UNIT/ML SUBCUTANEOUS CARTRIDG	28
MYDAYIS 12.5 MG CAPSULE EXTENDED RELEASE 24 HR	1	NOVOLOG U-100 INSULIN ASPART 100 UNIT/ML SUBCUTANEOUS SOLUTION	28
MYDAYIS 25 MG CAPSULE EXTENDED RELEASE 24 HR	1	NYVEPRIA 6 MG/0.6 ML SUBCUTANEOUS SYRINGE	26
MYDAYIS 37.5 MG CAPSULE EXTENDED RELEASE 24 HR	1	OSENI 12.5 MG-30 MG TABLET	12
MYDAYIS 50 MG CAPSULE EXTENDED RELEASE 24 HR	1	OSENI 25 MG-15 MG TABLET	12
<i>naftifine 1 % topical cream</i>	36	OSENI 25 MG-30 MG TABLET	12
<i>naftifine 2 % topical cream</i>	36	OSENI 25 MG-45 MG TABLET	12
<i>naftifine 2 % topical gel</i>	36	<i>oxiconazole 1 % topical cream</i>	36
NAFTIN 1 % TOPICAL GEL	36	OXISTAT 1 % LOTION	36
NAFTIN 2 % TOPICAL GEL	36	OXISTAT 1 % TOPICAL CREAM	36
NESINA 12.5 MG TABLET	12	PENNSAID 20 MG/GRAM/ACTUATION (2 %) TOPICAL SOLN IN METERED-DOSE PUMP	27
NESINA 25 MG TABLET	12	PROAIR RESPICLICK 90 MCG/ACTUATION BREATH ACTIVATED	4
NESINA 6.25 MG TABLET	12		
NEUPOGEN 300 MCG/0.5 ML INJECTION SYRINGE	17		
NEUPOGEN 300 MCG/ML INJECTION SOLUTION	17		
NEUPOGEN 480 MCG/0.8 ML INJECTION SYRINGE	17		
NEUPOGEN 480 MCG/1.6 ML INJECTION SOLUTION	17		
NEXICLON XR 0.17 MG TABLET,EXTENDED RELEASE	10		
NOVOLIN 70/30 U-100 INSULIN 100 UNIT/ML SUBCUTANEOUS SUSPENSION	31		

QUILLICHEW ER 20 MG CHEWABLE TABLET, EXTENDED RELEASE.....	1	RYTARY 23.75 MG-95 MG CAPSULE,EXTENDED RELEASE.....	30
QUILLICHEW ER 30 MG CHEWABLE TABLET, EXTENDED RELEASE.....	1	RYTARY 36.25 MG-145 MG CAPSULE,EXTENDED RELEASE.....	30
QUILLICHEW ER 40 MG CHEWABLE, EXTENDED RELEASE TABLET.....	1	RYTARY 48.75 MG-195 MG CAPSULE,EXTENDED RELEASE.....	30
QUILLIVANT XR 5 MG/ML (25 MG/5 ML) ORAL SUSPENSION,EXTEND RELEASE 24HR.....	1	RYTARY 61.25 MG-245 MG CAPSULE,EXTENDED RELEASE.....	30
QUVIVIQ 25 MG TABLET.....	32	<i>saxagliptin 2.5 mg tablet.....</i>	<i>12</i>
QUVIVIQ 50 MG TABLET.....	32	<i>saxagliptin 2.5 mg-metformin er 1,000 mg tablet,extend release 24hr mp.....</i>	<i>12</i>
RELEUKO 300 MCG/0.5 ML SUBCUTANEOUS SYRINGE.....	17	<i>saxagliptin 5 mg tablet.....</i>	<i>12</i>
RELEUKO 480 MCG/0.8 ML SUBCUTANEOUS SYRINGE.....	17	<i>saxagliptin 5 mg-metformin er 1,000 mg tablet,extend release 24hr mp.....</i>	<i>12</i>
RELEXXII 18 MG TABLET,EXTENDED RELEASE.....	1	<i>saxagliptin 5 mg-metformin er 500 mg tablet,extend release 24hr mp.....</i>	<i>12</i>
RELEXXII 27 MG TABLET,EXTENDED RELEASE.....	1	<i>sitagliptin 100 mg tablet.....</i>	<i>12</i>
RELEXXII 36 MG TABLET,EXTENDED RELEASE.....	1	<i>sitagliptin 25 mg tablet.....</i>	<i>12</i>
RELEXXII 45 MG TABLET,EXTENDED RELEASE.....	1	<i>sitagliptin 50 mg tablet.....</i>	<i>12</i>
RELEXXII 63 MG TABLET,EXTENDED RELEASE.....	1	<i>sitagliptin 50 mg-metformin 1,000 mg tablet.....</i>	<i>12</i>
RELTONE 200 MG CAPSULE.....	38	<i>sitagliptin 50 mg-metformin 500 mg tablet..</i>	<i>12</i>
RELTONE 400 MG CAPSULE.....	38	SOAANZ 20 MG TABLET.....	33
RHOPRESSA 0.02 % EYE DROPS.....	6	SOAANZ 40 MG TABLET.....	33
RITALIN 10 MG TABLET.....	1	SOAANZ 60 MG TABLET.....	33
RITALIN 20 MG TABLET.....	1	SORILUX 0.005 % TOPICAL FOAM....	37
RITALIN 5 MG TABLET.....	1	STEGLUJAN 15 MG-100 MG TABLET..	13
RITALIN LA 10 MG CAPSULE,EXTENDED RELEASE.....	1	STEGLUJAN 5 MG-100 MG TABLET..	13
RITALIN LA 20 MG CAPSULE,EXTENDED RELEASE.....	1	STIMUFEND 6 MG/0.6 ML SUBCUTANEOUS SYRINGE.....	26
RITALIN LA 30 MG CAPSULE,EXTENDED RELEASE.....	1	STRATTERA 10 MG CAPSULE.....	1
RITALIN LA 40 MG CAPSULE,EXTENDED RELEASE.....	1	STRATTERA 100 MG CAPSULE.....	1
ROCKLATAN 0.02 %-0.005 % EYE DROPS.....	6	STRATTERA 18 MG CAPSULE.....	1
RYALTRIS 665 MCG-25 MCG/SPRAY NASAL SPRAY.....	15	STRATTERA 25 MG CAPSULE.....	1
		STRATTERA 40 MG CAPSULE.....	1
		STRATTERA 60 MG CAPSULE.....	1
		STRATTERA 80 MG CAPSULE.....	1
		SUBOXONE 12 MG-3 MG SUBLINGUAL FILM.....	34
		SUBOXONE 2 MG-0.5 MG SUBLINGUAL FILM.....	34
		SUBOXONE 4 MG-1 MG SUBLINGUAL FILM.....	34
		SUBOXONE 8 MG-2 MG SUBLINGUAL FILM.....	34

SYMBICORT 160 MCG-4.5		VYVANSE 50 MG CAPSULE	1
MCG/ACTUATION HFA AEROSOL		VYVANSE 50 MG CHEWABLE	
INHALER	35	TABLET	1
SYMBICORT 80 MCG-4.5		VYVANSE 60 MG CAPSULE	1
MCG/ACTUATION HFA AEROSOL		VYVANSE 60 MG CHEWABLE	
INHALER	35	TABLET	1
TACLONEX 0.005 %-0.064 %		VYVANSE 70 MG CAPSULE	1
TOPICAL OINTMENT	37	VYZULTA 0.024 % EYE DROPS	6
TACLONEX 0.005 %-0.064 %		XELSTRYM 13.5 MG/9 HOUR	
TOPICAL SUSPENSION	37	TRANSDERMAL 24 HOUR PATCH	1
TRELSTAR 11.25 MG IM		XELSTRYM 18 MG/9 HOUR	
SUSPENSION	20	TRANSDERMAL 24 HOUR PATCH	1
TRELSTAR 22.5 MG IM SUSPENSION	20	XELSTRYM 4.5 MG/9 HOUR	
TRELSTAR 3.75 MG IM SUSPENSION	20	TRANSDERMAL 24 HOUR PATCH	1
TUDORZA PRESSAIR 400		XELSTRYM 9 MG/9 HOUR	
MCG/ACTUATION BREATH		TRANSDERMAL 24 HOUR PATCH	1
ACTIVATED	22	XOPENEX HFA 45	
TUDORZA PRESSAIR 400		MCG/ACTUATION AEROSOL	
MCG/ACTUATION BREATH		INHALER	7
ACTIVATED (30 ACTUAT)	22	ZIRGAN 0.15 % EYE GEL	21
TYRVAYA 0.03 MG/SPRAY NASAL		ZITUVIO 100 MG TABLET	12
SPRAY	14	ZITUVIO 25 MG TABLET	12
UDENYCA 6 MG/0.6 ML		ZITUVIO 50 MG TABLET	12
SUBCUTANEOUS SYRINGE	26		
UDENYCA AUTOINJECTOR 6			
MG/0.6 ML SUBCUTANEOUS AUTO-			
INJECTOR	26		
URSO 250 250 MG TABLET	38		
URSO FORTE 500 MG TABLET	38		
<i>ursodiol 200 mg capsule</i>	38		
<i>ursodiol 400 mg capsule</i>	38		
VECTICAL 3 MCG/GRAM TOPICAL			
OINTMENT	37		
VEVYE 0.1 % EYE DROPS	14		
VYVANSE 10 MG CAPSULE	1		
VYVANSE 10 MG CHEWABLE			
TABLET	1		
VYVANSE 20 MG CAPSULE	1		
VYVANSE 20 MG CHEWABLE			
TABLET	1		
VYVANSE 30 MG CAPSULE	1		
VYVANSE 30 MG CHEWABLE			
TABLET	1		
VYVANSE 40 MG CAPSULE	1		
VYVANSE 40 MG CHEWABLE			
TABLET	1		