

## PHARMACY COVERAGE GUIDELINE

**NURTEC™ ODT (rimegepant)**  
**QULIPTA™ (atogepant)**  
**REYVOW™ (lasmiditan)**  
**UBRELVY™ (ubrogepant)**  
**ZAVZPRET™ (zavegepant)**  
**Generic Equivalent (if available)**

---

### **This Pharmacy Coverage Guideline (PCG):**

- Provides information about the reasons, basis, and information sources we use for coverage decisions
- Is not an opinion that a drug (collectively “Service”) is clinically appropriate or inappropriate for a patient
- Is not a substitute for a provider’s judgment (Provider and patient are responsible for all decisions about appropriateness of care)
- Is subject to all provisions e.g. (benefit coverage, limits, and exclusions) in the member’s benefit plan; and
- Is subject to change as new information becomes available.

### **Scope**

- This PCG applies to Commercial and/or Marketplace plans
- This PCG does not apply to the Federal Employee Program, Medicare Advantage, Medicaid or members of out-of-state Blue Cross and/or Blue Shield Plans

### **Instructions & Guidance**

- To determine whether a member is eligible for the Service, read the entire PCG.
- This PCG is used for FDA approved indications including, but not limited to, a diagnosis and/or treatment with dosing, frequency, and duration.
- Use of a drug outside the FDA approved guidelines, refer to the appropriate Off-Label Use policy.
- The “Criteria” section outlines the factors and information we use to decide if the Service is medically necessary as defined in the Member’s benefit plan.
- The “Description” section describes the Service.
- The “Definition” section defines certain words, terms or items within the policy and may include tables and charts.
- The “Resources” section lists the information and materials we considered in developing this PCG
- **We do not accept patient use of samples as evidence of an initial course of treatment, justification for continuation of therapy, or evidence of adequate trial and failure.**
- Information about medications that require prior authorization is available at [www.azblue.com/pharmacy](http://www.azblue.com/pharmacy). You must fully complete the [request form](#) and provide chart notes, lab workup and any other supporting documentation. The prescribing provider must sign the form. Fax the form to BCBSAZ Pharmacy Management at (602) 864-3126 or email it to [Pharmacyprecert@azblue.com](mailto:Pharmacyprecert@azblue.com).

---

## Medical Necessity Requirements for **NURTEC ODT** (rimegepant)

### **Criteria for Initial Therapy:**

#### **Prescriber Qualifications**

- **ONE** of the following:
  - A Neurologist
  - A licensed professional **and ONE** of the following:

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

1. Is prescribing in consultation with a Neurologist or Pain Specialist
2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Indication

- Acute migraine of moderate to severe headache pain intensity
- Episodic migraine (moderate to severe headache pain intensity) defined as 4 to 14 headache days per month, with at least 4 migraine days

#### Age Requirement

- 18 years or older

#### Baseline Clinical Evaluation

- Confirmed diagnosis of acute or episodic migraine
- Does not have chronic migraine (15 or more headache days per month for more than 3 months, with migraine features on at least 8 days/month)

#### Alternative Therapies

- **Acute (abortive) treatment of migraine:**
  - Failure (used for at least two months), contraindication, or intolerance to at least **TWO** triptan drugs (naratriptan, rizatriptan, sumatriptan, and zolmitriptan)
  - If has 4 or more headache days per month, currently treated with **OR** documented failure (used for at least 2-months), contraindication, or intolerance to **ONE** of the following agents for migraine prevention:
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)
- **Episodic migraine prevention:**
  - Failure (used for at least 2-months), contraindication, or intolerance to **TWO** of the following agents for migraine prevention (must not be within the same class):
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)
- **Acute (abortive) treatment AND prevention of episodic migraine:**
  - Failure (used for at least two months), contraindication, or intolerance to at least **ONE** triptan drugs (naratriptan, rizatriptan, sumatriptan, and zolmitriptan)
  - Failure (used for at least 2-months), contraindication, or intolerance to **TWO** of the following agents for migraine prevention (must not be within the same class):
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

#### Safety

- **For acute migraine treatment:** No concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
- **For episodic migraine prevention:** No concomitant use with another CGRP antagonist for prevention (Aimovig, Ajovy, Emgality, Nurtec ODT, Qulipta, Vyepti)
- Not used in end-stage renal disease (creatinine clearance less than 15 mL/min)
- Not used in severe hepatic impairment (Child-Pugh Class C)
- No significant interacting drugs such as:
  - Strong CYP3A4 inhibitors (itraconazole, ketoconazole, posaconazole, etc.)
  - Strong/moderate CYP3A4 inducers (rifampin, phenytoin, St. John's Wort, phenobarbital, primidone, etc.)
  - P-glycoprotein or BCRP inhibitors (amiodarone, clarithromycin, cyclosporine, verapamil, etc.)

#### Documentation Requirements

- A completed request form must be submitted, including:
  - Chart notes
  - Lab results
  - Supporting clinical documentation

#### Initial Therapy Criteria Approval Duration

- 12 months OR end of plan year
- Acute treatment only: no more than 8 tablets per month
- Episodic prevention only: no more than 15 tablets per month
- Safety limit: no more than 18 tablets per 30 days

---

### Criteria for Continuation of Therapy (renewal therapy)

**Note: Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy**

#### Prescriber Qualification

- Continues to be seen by **ONE** of the following
  - Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Clinical Response

- **ALL** the following:
  - Achieved and maintains reduction of headache pain to no pain and absence of most bothersome symptom (photophobia, phonophobia, nausea)
  - Significant reduction in emergency room or urgent care visits
  - No evidence of disease progression

#### Adherence

- Adherence to prescribed therapy regimen has been documented

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

#### Safety

- **For acute migraine treatment:** No concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
- **For episodic migraine prevention:** No concomitant use with another CGRP antagonist for prevention (Aimovig, Ajovy, Emgality, Nurtec ODT, Qulipta, Vyepti)
- Not used in end-stage renal disease (creatinine clearance less than 15 mL/min)
- Not used in severe hepatic impairment (Child-Pugh Class C)
- No significant interacting drugs such as:
  - Strong CYP3A4 inhibitors (itraconazole, ketoconazole, posaconazole, etc.)
  - Strong/moderate CYP3A4 inducers (rifampin, phenytoin, St. John's Wort, phenobarbital, primidone, etc.)
  - P-glycoprotein or BCRP inhibitors (amiodarone, clarithromycin, cyclosporine, verapamil, etc.)

#### Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement
- Lab values confirming safe use

#### Continuation Therapy Criteria Approval Duration

- 12 months OR end of plan year
- When used for acute treatment only: no more than 8 ODT per month
- When used for Episodic Migraine prevention only: no more than 15 ODT per month
- Per labeling safety concerns: no more than 18 ODT per 30 days

---

## Medical Necessity Requirements for QULIPTA (atogepant)

---

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Criteria for Initial Therapy:

##### Prescriber Qualifications

- **ONE** of the following:
  - A Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

##### Indication

- Episodic migraine (moderate to severe headache pain intensity) defined as 4 to 14 headache days per month, with at least 4 migraine days
- Chronic migraine (moderate to severe headache pain intensity) defined as 15 or more headache days a month for more than 3 months, with migraine features on at least 8 days per month

##### Age Requirement

- 18 years or older

##### Baseline Clinical Evaluation

- **NONE** of the following:
  - Persistent daily headache
  - Trigeminal autonomic cephalgia (cluster headache)
  - Painful cranial neuropathy

##### Alternative Therapies

- Failure (trial for at least two months duration), contraindication, intolerance to at least **TWO** of the following (must not be within the same class)::
  - Beta-blocker (atenolol, metoprolol, nadolol, propranolol, timolol)
  - Anticonvulsant (topiramate, divalproex sodium, sodium valproate)
  - Antidepressant (amitriptyline, venlafaxine)

##### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the Food and Drug Administration (see Definitions section)

##### Safety

- No concomitant use with another CGRP antagonist for prevention (Aimovig, Ajovy, Emgality, Nurtec ODT, Qulipta, Vyepti)
- Not used in severe hepatic impairment (Child-Pugh Class C)
- **For Qulipta 60mg dose for chronic migraine, NONE** of the following:

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

- Strong cytochrome CYP3A4 inhibitors (itraconazole, etc.)
- Strong, moderate, or weak CYP3A4 inducers (rifampin, dexamethasone, topiramate, etc.)
- Severe renal impairment and end-stage renal disease (CrCl less than 30 mL/min)

#### Documentation Requirements

- A completed request form must be submitted, including:
  - Chart notes
  - Lab results
  - Supporting clinical documentation

#### Initial Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

#### Criteria for Continuation of Therapy (renewal therapy)

**Note: Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy**

#### Prescriber Qualification

- Continues to be seen by **ONE** of the following
  - Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Clinical Response

- **ALL** the following:
  - Achieved and maintains at least a 50 percent reduction in number of monthly migraine days from baseline
  - Achieved and maintain a reduction in number of monthly headache days
  - Achieved and maintains a reduction in the number of days of use of acute migraine-specific medications from baseline
  - Significant reduction in emergency room or urgent care visits for acute migraine treatment

#### Adherence

- Adherence to the prescribed therapy regimen has been documented

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Safety

- No concomitant use with another CGRP antagonist for prevention (Aimovig, Ajovy, Emgality, Nurtec ODT, Qulipta, Vyepti)
- Not used in severe hepatic impairment (Child-Pugh Class C)
- **For Qulipta 60mg dose for chronic migraine, NONE** of the following:
  - Strong cytochrome CYP3A4 inhibitors (itraconazole, etc.)
  - Strong, moderate, or weak CYP3A4 inducers (rifampin, dexamethasone, topiramate, etc.)
  - Severe renal impairment and end-stage renal disease (CrCl less than 30 mL/min)
- No persistent daily headache, trigeminal autonomic cephalgia (cluster headache) or painful cranial neuropathy

#### Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement in given indication
- Lab values that confirm safe use

#### Continuation Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

## Medical Necessity Requirements for REYVOW (lasmiditan)

---

### Criteria for Initial Therapy:

#### Prescriber Qualifications

- **ONE** of the following:
  - A Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Indication

- Acute migraine of moderate to severe headache pain intensity

#### Age Requirement

- 18 years or older

#### Baseline Clinical Evaluation

No more than 4 migraine attacks in a 30-day period

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Alternative Therapies

- **ALL** the following:
  - Failure (used for at least two months), contraindication, or intolerance to at least **TWO** triptan drugs (naratriptan, rizatriptan, sumatriptan, and zolmitriptan)
  - Failure (used for at least 2-months), contraindication, or intolerance to **BOTH** of the following:
    1. Nurtec
    2. Ubrelvy
  - If has 4 or more headache days per month, currently treated with **OR** documented failure (used for at least 2-months), contraindication, or intolerance to **ONE** of the following agents for migraine prevention:
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the Food and Drug Administration (see Definitions section)

#### Safety

- **NONE** of the following:
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
  - Reyvow for treatment of migraine prevention
  - Severe hepatic impairment (Child-Pugh Class C)
  - End-stage renal disease (creatinine clearance less than 15 milliliters per minute)
  - Significant interacting drugs including P-glycoprotein or BCRP substrates (e.g., dabigatran, digoxin, fexofenadine, rosuvastatin, sulfasalazine, etc.)

#### Documentation Requirements

- A completed request form must be submitted, including:
  - Chart notes
  - Lab results
  - Supporting clinical documentation

#### Initial Therapy Criteria Approval Duration

- 12 months OR end of plan year
- No more than 1 tab per 24 hours and no more than 4 tabs per 30-day period

---

#### Criteria for Continuation of Therapy (renewal therapy)

**Note: Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy**

#### Prescriber Qualification

- Continues to be seen by **ONE** of the following

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

- Neurologist
- A licensed professional **and ONE** of the following:
  1. Is prescribing in consultation with a Neurologist or Pain Specialist
  2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
  3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Clinical Response

- **ALL** the following:
  - Achieved and maintains reduction of headache pain to no pain and absence of most bothersome symptom (photophobia, phonophobia, nausea)
  - Significant reduction in emergency room or urgent care visits
  - No evidence of disease progression

#### Adherence

- Adherence to the prescribed therapy regimen has been documented

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

#### Safety

- **NONE** of the following:
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
  - Reyvow for treatment of migraine prevention
  - Severe hepatic impairment (Child-Pugh Class C)
  - End-stage renal disease (creatinine clearance less than 15 milliliters per minute)
  - Significant interacting drugs including P-glycoprotein or BCRP substrates (e.g., dabigatran, digoxin, fexofenadine, rosuvastatin, sulfasalazine, etc.)
  - Significant adverse effects such as:
    1. Central nervous system depression, including dizziness and sedation
    2. Serotonin syndrome

#### Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement in given indication
- Lab values that confirm safe use

#### Continuation Therapy Criteria Approval Duration

- 12 months OR end of plan year
- No more than 1 tab per 24 hours and no more than 4 tabs per 30-day period

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

## Medical Necessity Requirements for **UBRELVY** (ubrogepant)

### Criteria for Initial Therapy:

#### Prescriber Qualifications

- **ONE** of the following:
  - A Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Indication

- Acute migraine of moderate to severe headache pain intensity

#### Age Requirement

- 18 years or older

#### Alternative Therapies

- **BOTH** of the following:
  - Failure (used for at least two months), contraindication, or intolerance to at least **TWO** triptan drugs (naratriptan, rizatriptan, sumatriptan, and zolmitriptan)
  - If has 4 or more headache days per month, currently treated with **OR** documented failure (used for at least 2-months), contraindication, or intolerance to **ONE** of the following agents for migraine prevention:
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the Food and Drug Administration (see Definitions section)

#### Safety

- **NONE** of the following:
  - FDA-label contraindications, such as concurrent use with a strong CYP3A4 inhibitor (e.g., itraconazole, ketoconazole, clarithromycin, etc.)
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelyvy, Zavzpret)
  - End-stage renal disease (creatinine clearance less than 15 milliliters per minute)
  - Ubrelyvy for treatment of migraine prevention
  - No strong CYP3A4 inducers (e.g., rifampin, phenytoin, St. John's Wort, etc.)

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Documentation Requirements

- A completed request form must be submitted, including:
  - Chart notes
  - Lab results
  - Supporting clinical documentation

#### Initial Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

#### Criteria for Continuation of Therapy (renewal therapy)

**Note: Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy**

#### Prescriber Qualification

- Continues to be seen by **ONE** of the following
  - Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Clinical Response

- **ALL** the following:
  - Achieved and maintains reduction of headache pain to no pain and absence of most bothersome symptom (photophobia, phonophobia, nausea)
  - Significant reduction in emergency room or urgent care visits
  - No evidence of disease progression

#### Adherence

- Adherence to the prescribed therapy regimen has been documented

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

#### Safety

- **NONE** of the following:
  - FDA-label contraindications, such as concurrent use with a strong CYP3A4 inhibitor (e.g., itraconazole, ketoconazole, clarithromycin, etc.)
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelyvy, Zavzpret)
  - End-stage renal disease (creatinine clearance less than 15 milliliters per minute)
  - Ubrelyvy for treatment of migraine prevention

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

- No strong CYP3A4 inducers (e.g., rifampin, phenytoin, St. John's Wort, etc.)

#### Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement in given indication
- Lab values that confirm safe use

#### Continuation Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

### Medical Necessity Requirements for ZAVZPRET (zavegepant)

---

#### Criteria for Initial Therapy:

##### Prescriber Qualifications

- **ONE** of the following:
  - A Neurologist
  - A licensed professional **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

##### Indication

- Acute migraine of moderate to severe headache pain intensity

##### Age Requirement

- Request is for at least 18 years or older

##### Alternative Therapies

- **ALL** the following:
  - Failure (used for at least two months), contraindication, or intolerance to at least **TWO** triptan drugs (naratriptan, rizatriptan, sumatriptan, and zolmitriptan)
  - Failure (used for at least 2-months), contraindication, or intolerance to **BOTH** of the following:
    1. Nurtec
    2. Ubrelvy
  - If has 4 or more headache days per month, currently treated with **OR** documented failure (used for at least 2-months), contraindication, or intolerance to **ONE** of the following agents for migraine prevention:
    1. Beta-blocker (atenolol, metoprolol, nadolol, propranolol, or timolol)
    2. Anticonvulsant (topiramate, divalproex sodium, or sodium valproate)
    3. Antidepressant (amitriptyline or venlafaxine)

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the Food and Drug Administration (see Definitions section)

#### Safety

- **NONE** of the following:
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
  - Renal impairment (creatinine clearance less than 30 milliliters per minute)
  - Zavzpret for treatment of migraine prevention
  - Combination with drugs that inhibit organic anion transporting polypeptide 1B3 (OATP1B3) or sodium taurocholate co-transporting polypeptide (NTCP) transporters (e.g., rifampin)
  - Combination with drugs that induce OATP1B3 or NTCP transporters

#### Documentation Requirements

- A completed request form must be submitted, including:
  - Chart notes
  - Lab results
  - Supporting clinical documentation

#### Initial Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

#### Criteria for Continuation of Therapy (renewal therapy)

**Note: Manufacturer assistance (e.g., coupons, samples, etc.) are not considered for continuation of therapy**

#### Prescriber Qualification

- Continues to be seen by **ONE** of the following
  - Neurologist
  - A licensed **and ONE** of the following:
    1. Is prescribing in consultation with a Neurologist or Pain Specialist
    2. Is certified as a headache specialist by the United Council for Neurologic Subspecialties (UCNS)
    3. Has earned a Certificate in Added Qualifications (CAQ) in Headache Medicine from the National Headache Foundation

#### Clinical Response

- **ALL** the following:
  - Achieved and maintains reduction of headache pain to no pain and absence of most bothersome symptom (photophobia, phonophobia, nausea)
  - Significant reduction in emergency room or urgent care visits
  - No evidence of disease progression

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

#### Adherence

- Adherence to the prescribed therapy regimen has been documented

#### Brand Specific Criteria

- Have failure, contraindication or intolerance with **THREE** generic equivalents (when available) for at least three months each. **Note:** Any failure, contraindication, or intolerance to the generic drugs should be reported to the FDA (see Definitions section)

#### Safety

- **NONE** of the following:
  - Concomitant use with another CGRP antagonist for acute treatment (Nurtec ODT, Ubrelvy, Zavzpret)
  - Renal impairment (creatinine clearance less than 30 milliliters per minute)
  - Zavzpret for treatment of migraine prevention
  - Combination with drugs that inhibit organic anion transporting polypeptide 1B3 (OATP1B3) or sodium taurocholate co-transporting polypeptide (NTCP) transporters (e.g., rifampin)
  - Combination with drugs that induce OATP1B3 or NTCP transporters

#### Documentation Requirements

- Chart notes
- Supporting clinical documentation with evidence of improvement in given indication
- Lab values that confirm safe use

#### Continuation Therapy Criteria Approval Duration

- 12 months OR end of plan year
- 

### Criteria for Off-Label Use Requests:

Criteria for a request for non-FDA use or indication, treatment with dosing, frequency, or duration outside the FDA-approved dosing, frequency, and duration, refer to one of the following Pharmacy Coverage Guideline:

1. Off-Label Use of Non-Cancer Medications
  2. Off-Label Use of Cancer Medications
- 

#### Description:

Nurtec ODT (rimegepant) is a calcitonin gene-related peptide (CGRP) receptor antagonist indicated for the acute treatment of migraine with or without aura in adults and for preventive treatment of episodic migraine in adults. Reyvow (lasmiditan) is a serotonin (5-HT) 1F receptor agonist indicated for the acute treatment of migraine with or without aura in adults.

Ubrelvy (ubrogepant) is a CGRP receptor antagonist indicated for the acute treatment of migraine with or without aura. It is not indicated for the preventive treatment of migraine.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

Qulipta (atogepant) is a CGRP receptor antagonist indicated for the preventive treatment of episodic migraine in adults.

Zavzpret (zavegepant) is a CGRP receptor antagonist indicated for the acute treatment of migraine with or without aura in adults. It is not indicated for the preventive treatment of migraine. The safety of treating more than 8 migraines in a 30-day period has not been established.

The CGRP pathway is important in pain modulation, and CGRP has been observed to increase during a migraine. CGRP is a 37-amino acid peptide and functions as a neurotransmitter in the central and peripheral nervous system and as a vasodilator. The involvement of CGRP in migraine was suggested in the 1980s. Since then, new agents affecting the CGRP pathway have been developed and studied. Some approaches focused on small molecule CGRP receptor antagonists to be used to treat migraine attacks, or monoclonal antibodies to be used for migraine prevention.

Lasmiditan binds with high affinity to the 5-HT<sub>1F</sub> receptor. Lasmiditan presumably exerts its therapeutic effects in the treatment of migraine through agonist effects at the 5-HT<sub>1F</sub> receptor; however, the precise mechanism is unknown.

Migraine is a common episodic disorder, the hallmark of which is a disabling headache generally associated with nausea, and/or light and sound sensitivity. Migraine with aura refers to the occurrence of transient visual, sensory, language, or motor disturbance that is followed by a migraine headache. The exact mechanisms of migraine are unknown, but currently it is believed to initiate from a primary neuronal dysfunction that leads to a sequence of intracranial and extracranial changes accounting for migraine, including the four phases of premonitory symptoms, aura, headache and post-drome. Specifically, activation of the trigeminovascular system, cortical spreading depression, and neuronal sensitization all seem to play a role.

Pharmacologic therapies for migraine can be categorized broadly into agents used for treatment once symptoms have started (“acute” or “abortive” medications) and agents used to decrease the frequency or severity of migraines (“preventive” or “prophylactic” therapies).

Selection of medication for acute treatment is directed by the severity of the attacks, presence of associated nausea and vomiting, treatment setting, and patient-specific factors. Abortive treatments are usually more effective if they are given early in the course of the headache (within in the first hour if possible). A 2015 guideline assessment published by the American Headache Society lists the following medications as Level A (established as effective) for acute migraine treatment: all triptan drugs, NSAIDs (naproxen, ibuprofen, aspirin, diclofenac), combination of sumatriptan and naproxen, acetaminophen/aspirin/caffeine, acetaminophen (for acute treatment of non-incapacitating migraine), and dihydroergotamine nasal spray.

Prophylactic headache treatment is indicated if the headaches are frequent, long lasting, or account for a significant amount of total disability. A number of drug classes are used for the prevention of migraine. Medications that are effective in controlled trials include: beta blockers (metoprolol, propranolol, and timolol); anticonvulsants (valproate, divalproex, and topiramate); and antidepressants (amitriptyline and venlafaxine).

Outcomes of clinical trials of acute treatment of migraine commonly include relief of symptoms including pain, nausea/vomiting, photophobia and phonophobia; pain freedom; freedom from the most bothersome symptom (MBS); pain relief; and sustained symptom response.

The Migraine Disability Assessment (MIDAS) is a brief, 7-item, self-administered questionnaire designed to quantify headache-related disability. Respondents answer five questions about activity limitations in the past 3

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

months due to migraine including (1) missed work or school days, (2) missed household chores days, (3) missed non-work activity days, and days at work or school (4) plus days of household chores (5) where productivity was reduced by half or more. Two additional questions about the number of headaches and average pain level associated with headaches over the past 3 months are not used in deriving the MIDAS score, but they are for use by the respondent's clinician.

The MIDAS score is the sum of the number of days reported for each of the five questions. Respondents with a MIDAS score of 0-5 are rated as having little or no disability, 6-10 as having mild disability, 11-20 as having moderate disability, and 21 or greater as having severe disability.

---

#### **Definitions:**

U.S. Food and Drug Administration (FDA) MedWatch Forms for FDA Safety Reporting  
[MedWatch Forms for FDA Safety Reporting | FDA](#)

#### **Migraine day:**

- Any calendar day in which the patient experiences a qualified migraine headache (onset, continuation, or recurrence of the migraine headache)
- A qualified migraine is defined as migraine with or without aura, lasting  $\geq 30$  minutes that meets at least one of the following:
  - $\geq 2$  of the following pain features: unilateral, throbbing, moderate to severe, or exacerbated with exercise/physical activity
  - $> 1$  of the following associated non-pain features: nausea and or vomiting, or both photophobia, and phonophobia
- Any calendar day on which acute migraine-specific medication was used is counted as a migraine day

#### **Treatment considerations:**

- There are no strict definitions for the precise frequency or duration of migraine headaches that would prompt preventive therapy
- Migraine prevention therapy may be indicated for those with migraine headaches that are frequent (ex. as  $\geq 4$  headaches/month) or long-lasting (ex.  $\geq 12$  hours) and those that cause significant disability or diminished quality of life
- The goals of preventive therapy are to reduce the frequency, severity, and duration of headaches, to improve treatment responsiveness of therapies for acute attacks, prevent progression or transformation of episodic migraine to chronic migraine and to improve overall function or reduce the risk of neurologic impairment

#### **Episodic migraine:**

- Individual with migraine who has between 1 to 14 headache days per month, of which at least 4 were migraine days

#### **Chronic migraine:**

- Chronic migraine is defined as headache occurring 15 or more days a month for more than 3 months, which 8 days per month meet the features of migraine with or without aura.
- Some patients with an episodic migraine pattern ( $< 15$  headache days a month) transition to a chronic migraine pattern ( $\geq 15$  headache days a month), a transition that has been called "transformation" and "chronification"

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

- Features of migraine headache include:
  - Lasts 4-72 hours **AND** has at least 2 of the following 4 characteristics:
    - Unilateral, pulsating, moderate or severe pain intensity, aggravates or causes avoidance of routine physical activity
  - **AND** associated with at least one of the following during the headache:
    - Nausea and/or vomiting or photophobia and phonophobia.
- The management of chronic migraine should focus on prophylactic therapy and avoidance of acute headache medication overuse

#### **Pain freedom:**

- A **reduction in severity** of headache from mild, moderate or severe pain **at baseline to none** at a given follow-up time point

#### **Freedom from most bothersome symptoms (MBS):**

- **Total absence** of nausea/vomiting, photophobia or phonophobia at a given follow-up time point

#### **Pain relief:**

- **Having mild to no pain** at a given follow-up time point

#### **Sustained symptom response after 2-hours:**

- Those with an initial response that is sustained at subsequent follow-up time points **without** the use of repeat dosing or rescue medications

#### **Migraine Disability Assessment (MIDAS):**

Please answer the following questions about **ALL** of the headaches you have had over the last 3 months. Select zero if you did not have the activity in the last 3 months.

1. \_\_\_\_\_ On how many days in the last 3 months did you miss work or school because of your headaches?
2. \_\_\_\_\_ How many days in the last 3 months was your productivity at work or school reduced by half or more because of your headaches? (Do not include days you counted in question 1 where you missed work or school.)
3. \_\_\_\_\_ On how many days in the last 3 months did you not do household work (such as housework, home repairs and maintenance, shopping, caring for children and relatives) because of your headaches?
4. \_\_\_\_\_ How many days in the last 3 months was your productivity in household work reduced by half or more because of your headaches? (Do not include days you counted in question 3 where you did not do household work.)
5. \_\_\_\_\_ On how many days in the last 3 months did you miss family, social or leisure activities because of your headaches?
6. Total number of days (from questions 1 through 5): \_\_\_\_\_

#### **Answer the following for your provider:**

1. \_\_\_\_\_ On how many days in the last 3 months did you have a headache? (If a headache lasted more than 1 day, count each day.)

**PHARMACY COVERAGE GUIDELINE**

**NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™**

2. \_\_\_\_\_ On a scale of 0 - 10, on average how painful were these headaches? (where 0 = no pain at all, and 10 = pain as bad as it can be.)

MIDAS Grade	Definition	MIDAS Score
I	Little or No disability	0-5
II	Mild disability	6-10
III	Moderate disability	11-20
IV	Severe disability	21+

<b>AHS Consensus 2021</b>
Patients with migraine should be considered for preventive treatment in <b>any</b> of the following situations:
• Attacks significantly interfere with patients' daily routines despite acute treatment
• Frequent attacks as described in following table
• Contraindication to, failure, or overuse of acute treatments, with overuse defined as follows: <ul style="list-style-type: none"> <li>a. Ten or more days per month for ergot derivatives, triptans, opioids, combination analgesics, and a combination of drugs from different classes that are not individually overused</li> <li>b. Fifteen or more days per month for nonopioid analgesics, acetaminophen, and NSAIDs</li> </ul>
• Adverse events with acute treatments
• Patient preference

<b>AHS Consensus 2021</b>		
Identifying Patients for Preventive Treatment		
Prevention should be:	Headache days/month	Degree of MIDAS required
Offered	6 or more	None
	4 or more	Some
	3 or more	Severe
Considered	4 or 5	None
	3	Some
	2	Moderate

**Migraine Severity:**

- **Mild Pain Level / Pain Score**
  - Does not interfere with most activities and is easy to manage both physically and psychologically. Individual able to adapt to these levels of pain with low doses of medication (e.g., acetaminophen), dietary changes, or bed rest.
- **Moderate Uncomfortable Pain Level / Pain Score**
  - Interferes with many activities of daily living and requires changes to daily lifestyle to manage pain symptoms. Migraine pain is more noticeable but is not incapacitating.
- **Severe Pain Level / Pain Score**
  - Individual is no longer able to engage in normal activities and seeks stronger medications to help improve ability to function independently.

<b>Identification of headache type: migraine, tension, or cluster</b>			
	<b>Migraine</b>	<b>Tension</b>	<b>Cluster</b>

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

**PHARMACY COVERAGE GUIDELINE**

**NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™**

Location	Unilateral	Bilateral	Supraorbital/temporal
Pain intensity <sup>1</sup>	Moderate to severe	Mild to moderate	Severe
Duration	4–72 hours	30 minutes to 7 days	15–180 minutes
Characterization of pain	Pulsing	Pressure/squeezing	Boring/stabbing
Sensitivity to light/sound	One or both may be present	Both are absent or only one is present	No
Nausea/vomiting	One or both may be present	No	One or both may be present
Aggravated by routine activity	Yes	No	No
Aura	May be present	No	No
Associated symptoms	None	None	Miosis, ptosis, rhinorrhea
<sup>1</sup> Pain intensity <ul style="list-style-type: none"> <li>• Mild—Patient is aware of a headache but is able to continue daily routine with minimum alterations.</li> <li>• Moderate—The headache inhibits daily activities; migraine pain is more noticeable but is not incapacitating.</li> <li>• Severe—The headache is incapacitating such that patient is no longer able to engage in normal activities.</li> </ul>			

**2013 Canadian Headache Society (CHS) – medications for acute migraine:**

2013 Canadian Headache Society (CHS) Summary of Recommendations*		
Recommended for Use in Episodic Migraine** (Use)		
Drug	Recommendation Strength	Quality of Evidence
Almotriptan	Strong	High
Eletriptan	Strong	High
Frovatriptan	Strong	High
Naratriptan	Strong	High
Rizatriptan	Strong	High
Sumatriptan	Strong	High
Zolmitriptan	Strong	High
Aspirin	Strong	High
Diclofenac	Strong	High
Ibuprofen	Strong	High
Naproxen	Strong	High
Acetaminophen	Strong	High
Domperidone	Strong	Low
Metoclopramide	Strong	Moderate
Dihydroergotamine	Weak	Moderate
Ergotamine	Weak, not recommended for routine use	Moderate
Opioid containing compounds	Weak, not recommended for routine use	Low
Tramadol containing compounds	Weak, not recommended for routine use	Moderate
Not Recommended for Use in Episodic Migraine** (Do not use***)		
Butalbital containing compounds	Strong	Low
Butorphanol	Strong	Low
*Utilizing Grading of Recommendations Assessment, Development and Evaluation (GRADE) Criteria **Migraine with headache on less than 15 days a month *** Except under exceptional circumstances		

Metoclopramide strongly recommended for use when necessary

**Abortive (symptomatic) treatment of acute migraine:**

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

---

- Simple analgesics such as non-steroidal anti-inflammatory drugs (NSAIDs) or acetaminophen, triptans, antiemetics, calcitonin gene-related peptide (CGRP) antagonists (remigepant, ubrogepant, zavegepant), lasmiditan, and dihydroergotamine

#### **Non-Calcitonin gene-related peptide (Non-CGRP) preventative (episodic or chronic) migraine agent(s):**

- Beta-blocker: atenolol, metoprolol, nadolol, propranolol, or timolol
- Antidepressant: amitriptyline or venlafaxine
- Anticonvulsant: topiramate, divalproex sodium, or sodium valproate

#### **Botulinum toxin injection:**

- Treatment of chronic migraine:
  - Botox (onabotulinumtoxinA)

#### **CGRP related agents:**

- Preventive treatment of episodic or chronic migraine:
  - Vyepti (eptinezumab-jjmr)
  - Aimovig (erenumab)
  - Ajovy (fremanezumab-vfrm)
  - Emgality (galcanezumab) – also used in cluster headache
- Preventive treatment of episodic migraine:
  - Qulipta (atogepant)
  - Nurtec ODT (rimegepant)
- Acute (abortive) treatment of migraine:
  - Oral:
    - Nurtec ODT (rimegepant)
    - Ubrelvy (ubrogepant)
  - Nasal:
    - Zavzpret (zavegepant)

#### **Serotonin (5-HT) 1F receptor agonist:**

- Acute (abortive) treatment of migraine
    - Reyvow (lasmiditan)
- 

#### **Resources:**

Nurtec ODT (rimegepant) product information, revised by Pfizer Laboratories Div Pfizer Inc. 04-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 30, 2024.

Qulipta (atogepant) product information, revised by AbbVie Inc. 06-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 30, 2024.

Reyvow (lasmiditan) product information, revised by Eli Lilly and Company 09-2022. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 30, 2024.

Ubrelvy (ubrogepant) product information, revised by Allergan, Inc. 06-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 30, 2024.

Zavzpret (zavegepant) product information, revised by Pfizer Labs Division of Pfizer Inc. 03-2023. Available at DailyMed <http://dailymed.nlm.nih.gov>. Accessed August 30, 2024.

ORIGINAL EFFECTIVE DATE: 02/13/2020 | ARCHIVE DATE: | LAST REVIEW DATE: 05/15/2025 | LAST CRITERIA REVISION DATE: 05/15/2025

BLUE CROSS®, BLUE SHIELD® and the Cross and Shield Symbols are registered service marks of the Blue Cross and Blue Shield Association, an association of independent Blue Cross and Blue Shield Plans. All other trademarks and service marks contained in this guideline are the property of their respective owners, which are not affiliated with BCBSAZ.

## PHARMACY COVERAGE GUIDELINE

### NURTEC™ ODT QULIPTA™ REYVOW™ UBRELVY™ ZAVZPRET™

Schwedt TJ, Garza I. Acute treatment of migraine in adults. In: UpToDate, Swanson JW, Goddeau RP (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated March 25, 2025. Accessed March 28, 2025.

Schwedt TJ, Garza I. Preventive treatment of episodic migraine in adults. In: UpToDate, Swanson JW, Goddeau RP (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated March 25, 2025. Accessed March 28, 2025.

Garza I, Schwedt TJ. Chronic migraine. In: UpToDate, Robertson CE, Goddeau RP (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated October 03, 2024. Accessed March 28, 2025.

Cutrer FM. Pathophysiology, clinical manifestations, and diagnosis of migraine in adults. In: UpToDate, Swanson JW, Goddeau RP (Eds), UpToDate, Waltham MA.: UpToDate Inc. Available at <http://uptodate.com>. Literature current through February 2025. Topic last updated November 06, 2024. Accessed March 28, 2025.

Ailani J, Burch RC, Robbins MS. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. *Headache* 2021;61:1021–1039. Accessed April 25, 2023. Re-evaluated March 27, 2025.

Simpson DM, Hallett M, Ashman EJ, et al. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache. Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology* 2016;86:1818–1826. Accessed on April 25, 2023. Re-evaluated March 27, 2025.

Institute for Clinical and Economic Review (ICER): Calcitonin Gene-Related Peptide (CGRP) Inhibitors as Preventive Treatments for Patients with Episodic or Chronic Migraine: Effectiveness and Value. Final Evidence Report; July 03, 2018; 1-275. Available at <https://icer.org/>. Accessed April 27, 2023. Re-evaluated March 27, 2025.

Puledda F, Sacco S, Hans-Christop D, et al.: International Headache Society global practice recommendations for the acute pharmacological treatment of migraine. *Cephalalgia* 2024; 44(8): 1–45. Accessed March 27, 2025.

Qassem A, Tice JA, Exteandia-Ikobaltzeta I, et al.: Pharmacologic Treatments of Acute Episodic Migraine Headache in Outpatient Settings: A Clinical Guideline from the American College of Physicians. *Ann Intern Med*. doi:10.7326/ANNALS-24-03095. Accessed March 27, 2025.

Qassem A, Cooney T, Exteandia-Ikobaltzeta I, et al.: Prevention of Episodic Migraine Headache using Pharmacologic Treatments in Outpatient Settings: A Clinical Guideline from the American College of Physicians. 2025; 178:426-433. doi:10.7326/ANNALS-24-01052. Accessed March 27, 2025.