

lt's Wholecare.

Prior Authorization Criteria Vyndaqel (tafamidis meglumine) and Vyndamax (tafamidis)

All requests for Vyndaqel (tafamidis meglumine) and Vyndamax (tafamidis) require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Coverage may be provided with a <u>diagnosis</u> of cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) and the following criteria is met:

- Must be 18 years of age or older
- The diagnosis is confirmed by presence of amyloid deposits on biopsy analysis from cardiac or non-cardiac sites (e.g., fat aspirate, gastrointestinal sites, salivary glands, bone marrow) or by technetium-labeled bone scintigraphy tracing
- Cardiac involvement was confirmed by echocardiography or cardiac magnetic resonance imaging (MRI) (e.g., end-diastolic interventricular septal wall thickness exceeding 12 mm)
- For members with hereditary ATTR-CM, presence of a mutation of the TTR gene was confirmed
- For members with wild type ATTR-CM, presence of transthyretin precursor proteins was confirmed by immunohistochemical analysis, scintigraphy, or mass spectrometry
- The member exhibits clinical symptoms of cardiomyopathy and heart failure (e.g., dyspnea, fatigue, orthostatic hypotension, syncope, peripheral edema)
- Member has a New York Heart Association Class I, II or III heart failure
- The member will not receive Vyndaqel or Vyndamax in combination with either of the following
 - Tetramer stabilizers (e.g. diflunisal)
- Must be prescribed by or in consultation with a cardiologist or a physician who specializes in the treatment of amyloidosis
- Member does not have a history of liver or heart transplantation
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
- Initial Duration of Approval: 12 months
- Reauthorization criteria
 - Documentation confirming the member demonstrates a beneficial response to treatment (e.g., improvement on the 6-minute walk test, the Kansas City Cardiomyopathy Questionnaire–Overall Summary (KCCQ-OS) score, cardiovascular-related hospitalizations, NYHA classification of heart failure, left ventricular stroke volume, NT-proBNP level)
- Reauthorization Duration of Approval: 12 months

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or



Updated: 08/2021 PARP Approved: 10/2021

peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

lt's

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.



VYN	VYNDAQEL (tafamidis meglumine) and VYNDAMAX (tafamidis)					
PRIOR AUTHORIZATION FORM						
Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Gateway Health SM Pharmacy Services. FAX: (888) 245-2049						
If needed, you may call to speak to a Pharmacy Services Representative.						
PHONE : (800) 392-1147 Monday through Friday 8:30am to 5:00pm						
PROVIDER INFORMATION						
Requesting Provider:						
Provider Specialty:			Office Contact:			
State license #:						
Office Address:			Office Phone:			
			Office Fax:			
MEMBER INFORMATION						
Member Name:		DOB:				
Gateway ID:		Member weight:	Height:			
	REQUESTED DR	UG INFORMATION				
Medication:		Strength:				
Directions:		Quantity:	Refills:			
Is the member currently receiving receiving	quested medication? 🗌 Yes	No Date N	Aedication Initiated:			
	Billing I	Information				
This medication will be billed:	t a pharmacy OR med	ically, JCODE:				
Place of Service: Hospital	Provider's office 🗌 Meml	per's home 🗌 Other				
	Place of Serv	vice Information				
Name:		NPI:				
Address:			Phone:			
	MEDICAL HISTORY (
Diagnosis: Cardiomyopathy of wild-type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) Other: ICD-10 Code:						
	 ov presence of amyloid der					
Has the diagnosis been confirmed by presence of amyloid deposits on biopsy analysis from cardiac or non-cardiac sites (e.g., fat aspirate, gastrointestinal sites, salivary glands, bone marrow) or by technetium-labeled bone scintigraphy tracing?						
	alivary glands, bone marro		beled bone scintigraphy tracing?			
Yes No		w) or by technetium-la				
Yes No Has cardiac involvement been cont	firmed by echocardiograph	w) or by technetium-la y or cardiac magnetic i	resonance imaging? Yes No			
Yes No Has cardiac involvement been cont For members with hereditary ATT	firmed by echocardiograph R-CM, has the presence of	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l	resonance imaging? Yes No			
Yes No Has cardiac involvement been conf For members with hereditary ATT For members with wild type ATTR	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by			
Yes No Has cardiac involvement been conf For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci	firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectro	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l ransthyretin precursor metry? Yes No	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l ransthyretin precursor metry? Yes No	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yor	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l ransthyretin precursor metry? Yes No ny and heart failure (e.g	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yor Is the member receiving Vyndaqel	Firmed by echocardiograph R-CM, has the presence of the trigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yorl Is the member receiving Vyndaqel Does the member have a history of	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g I, II or III heart failure? ion with tetramer stabil on? Yes No	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yor Is the member have a New Yor Is the member have a history of Is the medication being prescribed	Firmed by echocardiograph R-CM, has the presence of R-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g I, II or III heart failure? ion with tetramer stabil on? Yes No	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yorl Is the member receiving Vyndaqel Does the member have a history of	Firmed by echocardiograph R-CM, has the presence of t c-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination liver or heart transplantation by or in consultation with a	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l ransthyretin precursor metry? Yes No ny and heart failure (e.g ion with tetramer stabil on? Yes No a cardiologist or a phys	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			
Yes No Has cardiac involvement been conf For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yorl Is the member receiving Vyndaqel Does the member have a history of Is the medication being prescribed amyloidosis? Yes No	Firmed by echocardiograph R-CM, has the presence of the trigraphy, or mass spectrom ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination i liver or heart transplantation by or in consultation with a	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g I, II or III heart failure? ion with tetramer stabil on? Yes No a cardiologist or a phys	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			
Yes No Has cardiac involvement been cont For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yor Is the member have a New Yor Is the member have a history of Is the medication being prescribed	Firmed by echocardiograph R-CM, has the presence of t c-CM, has the presence of t ntigraphy, or mass spectron ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination liver or heart transplantation by or in consultation with a	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l ransthyretin precursor metry? Yes No ny and heart failure (e.g ion with tetramer stabil on? Yes No a cardiologist or a phys	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			
Yes No Has cardiac involvement been conf For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yorl Is the member receiving Vyndaqel Does the member have a history of Is the medication being prescribed amyloidosis? Yes No	Firmed by echocardiograph R-CM, has the presence of the trigraphy, or mass spectrom ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination i liver or heart transplantation by or in consultation with a	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g I, II or III heart failure? ion with tetramer stabil on? Yes No a cardiologist or a phys	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			
Yes No Has cardiac involvement been conf For members with hereditary ATT For members with wild type ATTR immunohistochemical analysis, sci Does the member exhibit clinical s hypotension, syncope, peripheral e Does the member have a New Yorl Is the member receiving Vyndaqel Does the member have a history of Is the medication being prescribed amyloidosis? Yes No	Firmed by echocardiograph R-CM, has the presence of the trigraphy, or mass spectrom ymptoms of cardiomyopath dema)? Yes No k Heart Association Class I or Vyndamax in combination i liver or heart transplantation by or in consultation with a	w) or by technetium-la y or cardiac magnetic r a TTR gene mutation l rransthyretin precursor metry? Yes No ny and heart failure (e.g I, II or III heart failure? ion with tetramer stabil on? Yes No a cardiologist or a phys	resonance imaging? Yes No peen confirmed? Yes No proteins been confirmed by g., dyspnea, fatigue, orthostatic Yes No lizers (e.g. diflunisal)? Yes No			



lt's Wholecare.

VYNDAQEL (tafamidis meglumine) and VYNDAMAX (tafamidis)						
PRIOR AUTHORIZATION FORM (CONTINUED)-PAGE 2 of 2						
Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation						
as applicable to Gateway Health SM Pharmacy Services. FAX: (888) 245-2049						
If needed, you may call to speak to a Pharmacy Services Representative.						
PHONE : (800) 392-1147 Monday through Friday 8:30am to 5:00pm						
MEMBER INFORMATION						
Member Name:	DOB:					
Gateway ID:	Member weight:	Height:				
REAUTHORIZATION						
Is there documentation confirming the member has demonstrated a beneficial response to treatment (e.g., improvement on the 6-						
minute walk test, Kansas City Cardiomyopathy Questionnaire-Overall Summary (KCCQ-OS) score, cardiovascular-related						
hospitalizations, NYHA classification of heart failure, left ventricular stroke volume, NT-proBNP level)? Yes No						
SUPPORTING INFORMATION or CLINICAL RATIONALE						
Prescribing Provider Signature		Date				