



Prior Authorization Criteria
Non-TNF Biologic Therapies

All requests for Non-TNF Biologic Therapies require a prior authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Drugs addressed in this policy: Orencia (abatacept), Actemra (tocilizumab), Arcalyst (rilonacept), Cosentyx (secukinumab), Entyvio (vedolizumab), Ilaris (canakinumab), Kineret (anakinra), Otezla (apremilast), Stelara (ustekinumab), Taltz (ixekizumab), Tremfya (Guselkumab), Siliq (Brodalumab), Ilumya (tildrakizumab-asmn), Olumiant (baricitinib)

Indication	Product
Rheumatoid Arthritis (RA)	Orencia, Actemra, Olumiant, or Kineret
Polyarticular Juvenile Idiopathic Arthritis (JIA)	Orencia, Actemra
Systemic Juvenile Idiopathic Arthritis (JIA)	Actemra, Ilaris
Plaque Psoriasis	Cosentyx, Stelara, Taltz, Otezla, Siliq, Ilumya, or Tremfya
Psoriatic Arthritis (PsA)	Orencia, Otezla, Cosentyx, Stelara, or Taltz
Cytokine Release Syndrome (CRS)	Actemra
Giant Cell Arteritis	Actemra
Cryoprin-Associated Periodic Syndrome (CAPS)	Arcalyst, Ilaris, or Kineret
Ankylosing Spondylitis	Cosentyx
Ulcerative Colitis	Entyvio
Crohn's Disease	Entyvio, Stelara
Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS)	Ilaris
Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD)	Ilaris
Familial Mediterranean Fever (FMF)	Ilaris

For all requests for Non-TNF Biologic Therapies all of the following criteria must be met:

- 1) The prescribing physician must be a Rheumatologist, Dermatologist, Gastroenterologist, Oncologist, or Ophthalmologist.
- 2) The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines.
- 3) Members who are currently established on therapy will not be required to change to a formulary/preferred product.

Coverage may be provided with a diagnosis of **Rheumatoid Arthritis (RA)** and the following criteria is met:

- Prescribed medication is Orencia, Actemra, Olumiant, or Kineret.
- Member is 18 years of age or older
- Member must have a history of trial and failure, contraindication, or intolerance to ALL of the following
 - Three-month trial with methotrexate, or another DMARD.
 - Three-month trial with Xeljanz*
 - Three-month trial with either Humira* OR Enbrel* with or without MTX unless there is clinical rationale for using a non-TNF over a TNF inhibitorFor Actemra:
 - Member must have ALL of the following lab values:
 - $ANC \geq 2000/mm^3$ at baseline.
 - Platelet count $\geq 100,000/mm^3$ at baseline.
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- For Olumiant
 - Member must avoid use of live vaccines
 - Member must have ALL of the following lab values
 - Hemoglobin greater than 8g/dL at baseline
 - Absolute Lymphocytic Count greater than 500 cells/mm³ at baseline
 - Absolute Neutrophil Count greater than 1000 cells/mm³ at baseline
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- **Initial Duration of Approval:**
 - For Olumiant: 3 months
 - For Actemra, Orencia and Kineret: 6 months
- **Reauthorization Criteria**
 - For Olumiant
 - Member must continue to avoid use of live vaccines
 - Member must have ALL of the following lab values following 3 months of therapy:
 - Hemoglobin greater than 8g/dL
 - Absolute Lymphocytic Count greater than 500 cells/mm³
 - Absolute Neutrophil Count greater than 1000 cells/mm³
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months
 - For Actemra, Orencia and Kineret
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment

- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Polyarticular Juvenile Idiopathic Arthritis (PJIA)** and the following criteria is met:

- Prescribed medication is Orenzia, or Actemra.
- Member is 2 years of age or older.
- Member must have a history of trial and failure, contraindication, or intolerance to a three-month trial with either Humira*, OR Enbrel*.
- Member must meet ONE of the following:
 - The member has an AJC (active joint count)>0 and continued disease activity after 3 months of MTX or leflunomide.
 - The member has an AJC (active joint count)>0 and continued disease activity after 1 month of anakinra*.
- Specific for Actemra:
 - The member has ALL of the following lab values:
 - $ANC \geq 2000/mm^3$ at baseline.
 - Platelet count $\geq 100,000/mm^3$ at baseline.
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria**
 - Reauthorization benefit will be approved if there is documented, significant improvement in AJC (active joint count) with prior courses of treatment.
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Systemic Juvenile Idiopathic Arthritis (SJIA)** and the following criteria is met:

- Prescribed medication is Actemra or Ilaris.
- Member is age 2 years of age or older.
- The member must meet ONE of the following:
 - The member has an MD global ≥ 5 and continued disease activity after 1 month of NSAIDs.
 - The member has continued disease activity after 2 weeks of steroid monotherapy.
 - The member has continued disease activity after 1 month of anakinra* therapy.
- Specific for Actemra:
 - The member has ALL of the following lab values:
 - $ANC \geq 2000/mm^3$ at baseline.
 - Platelet count $\geq 100,000/mm^3$ at baseline.
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- **Initial Duration of Approval:**

- For Ilaris: 6 months
- **Reauthorization Criteria**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Plaque Psoriasis** and the following criteria is met:

- Prescribed medication is Cosentyx, Stelara, Taltz, Otezla, Siliq, Ilumya, or Tremfya.
- Member is age 18 years of age or older. (12 years or older for Stelara only).
- Member must have clinical documentation of a diagnosis of moderate to severe plaque psoriasis characterized by greater than or equal to 5% body surface involved or disease affecting crucial body areas such as hands, feet, face, or genitals.
- Members must have therapeutic failure of a three or more month trial or a contraindication to ALL of the following:
 - Psoralens with UVA light (PUVA) or UVB light
 - Systemic treatments including ONE of the following:
 - Immunomodulators (i.e. Methotrexate, Cyclosporine)
 - Retinoids (i.e. Soriatane)
- Member must have a history of trial and failure, contraindication, or intolerance to a three-month trial with either Humira*, OR Enbrel*.
- Specific for Otezla:
 - Will not be used in combination with a strong cytochrome P450 enzyme inducer (e.g. rifampin, phenobarbital, carbamazepine, phenytoin)
- Specific for Siliq
 - Member should not have a diagnosis of Crohn's disease.
- Specific for Siliq, Ilumya, and Tremfya
 - Will avoid use of live vaccines while on medication.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Clinical documentation that supports a decrease in percent of body surface area involvement when compared to baseline must be submitted
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Psoriatic Arthritis (PsA)** and the following criteria is met:

- Prescribed medication is Orencia, Otezla, Cosentyx, Stelara, or Taltz
- Member is age 18 years of age or older.
- Member has moderately to severely active psoriatic arthritis which must include documentation of either active psoriatic lesions or documented history of psoriasis.
- Member must have a history of trial and failure, contraindication, or intolerance to ALL of the following:

- Three-month trial with either Humira* OR Enbrel*
- Three-month trial with Xeljanz*
- Member must have a history of trial and failure, contraindication, or intolerance to ALL of the following:
 - Member without axial disease:
 - Four- week trial each of at least 2 NSAIDs.
 - Eight week trial of methotrexate or other DMARD
 - Member with axial disease
 - Four- week trial each of at least 2 NSAIDs.
 - Member with psoriatic arthritis with enthesitis
 - Four- week trial each of at least 2 NSAIDs.
- Specific for Otezla:
 - Will not be used in combination with a strong cytochrome P450 enzyme inducer (e.g. rifampin, phenobarbital, carbamazepine, phenytoin)
- **Initial Duration of Approval:** 6 months
- **Reauthorization:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Cytokine Release Syndrome (CRS)** and the following criteria is met:

- Prescribed medication is Actemra.
- Member is age 2 years of age or older.
- Member meets one of the following criteria for Cytokine Release Syndrome (CRS):
 - Grade 1 CRS with persistent or refractory fever
 - Grade 2 with any ONE of the following:
 - Hypotension refractory to fluid boluses
 - Presence of hypoxia.
 - Organ toxicity.
 - Documented diagnosis of chimeric antigen receptor (CAR) T cell-induced severe (grade 3) CRS with any ONE of the following:
 - Defined as: oxygen requirement $\geq 40\%$.
 - Hypotension requiring high dose or multiple vasopressors.
 - Grade 3 organ toxicity or grade 4 transaminitis.
 - Life threatening (grade 4) CRS with any ONE of the following:
 - Defined as: requirement for ventilator support.
 - Grade 4 organ toxicity (excluding transaminitis).
- The member has ALL of the following lab values:
 - ANC $\geq 2000/\text{mm}^3$ at baseline.
 - Platelet count $\geq 100,000/\text{mm}^3$ at baseline.
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- **Initial Duration of Approval:** 3 months

- **Reauthorization Criteria**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Giant Cell Arteritis** and the following criteria is met:

- Prescribed medication is Actemra
- The member is 18 years of age or older
- Diagnosis is confirmed by ONE of the following:
 - Temporal artery biopsy or cross-sectional imaging
 - Acute-phase reactant elevation (i.e. high erythrocyte sedimentation rate and/ or high serum C-reactive protein)
- Member must have a history of trial and failure, contraindication, or intolerance to a one-week trial of glucocorticoids (e.g., prednisone, methylprednisone).
- The member has the following lab values:
 - ANC \geq 2000/mm³ at baseline.
 - Platelet count \geq 100,000/mm³ at baseline.
 - ALT/AST does not exceed 1.5 times the upper limit of normal at baseline.
- **Initial Duration of Approval:** 3 months
- **Reauthorization Criteria**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Cyropyrin-Associated Periodic Syndrome (CAPS)** and the following criteria is met:

- Prescribed medication is Arcalyst, Ilaris, or Kineret
- Specific for Arcalyst:
 - The member is 12 years of age or older
 - The member has genetic evidence of an CIAS1 (NLRP3) mutation based on DNA sequencing.
 - The member has documented signs and symptoms associated with ONE of the following conditions:
 - Familial Cold Autoinflammatory Syndrome (FCAS).
 - Recurrent, intermittent fever
 - Rash that is often exacerbated by exposure to generalized cool ambient temperature
 - Muckle-Wells Syndrome (MWS).
 - Rash of waxing and waning intensity sometimes exacerbated by exposure to generalized cool ambient temperature.

- Specific for Ilaris
 - The member is 4 years of age or older.
 - The member has genetic evidence of a CIAS1 (NLRP3) mutation based on DNA sequencing.
 - The member has documented signs and symptoms associated with ONE of the following conditions:
 - Familial Cold Autoinflammatory Syndrome (FCAS).
 - Recurrent, intermittent fever
 - Rash that is often exacerbated by exposure to generalized cool ambient temperature
 - Muckle-Wells Syndrome (MWS).
 - Rash of waxing and waning intensity sometimes exacerbated by exposure to generalized cool ambient temperature.
- Specific for Kineret:
 - The member has a documented diagnosis of Neonatal-Onset Multisystem Inflammatory Disease (NOMID)
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Ankylosing Spondylitis** and the following criteria is met:

- Prescribed medication is Cosentyx.
- Member is age 18 years of age or older.
- Member must have a history of trial and failure, contraindication, or intolerance to ALL of the following:
 - Four- week trial each of at least 2 NSAIDs
 - Three-month trial with either Humira*, OR Enbrel*.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
 - **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Ulcerative Colitis** and the following criteria is met:

- Prescribed medication is Entyvio
- Member is 18 years of age or older.
- Member must have a history of trial and failure, contraindication, or intolerance to conventional treatments including all of the following for at least 3 months:

- Aminosalicylates, 5-ASAs (*i.e.*, Sulfasalazine, Pentasa[®], Asacol[®], Colazal[®])
- Steroids (*i.e.*, prednisone, Entocort[®])
- Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Member must have a history of trial and failure, contraindication, or intolerance to Humira* for at least 3 months.
- Member must have a history of trial and failure, contraindication, or intolerance to Xeljanz* for at least 3 months.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Crohn's Disease** and the following criteria is met:

- Prescribed medication is Entyvio or Stelara.
- Member is 18 years of age or older.
- Member must have a history of trial and failure, contraindication, or intolerance to conventional treatments including two or more of the following for at least 3 months of each medication:
 - Aminosalicylates, 5-ASAs (*i.e.*, Sulfasalazine, Pentasa[®], Asacol[®], Colazal[®])
 - Antibiotics (*i.e.*, Metronidazole, Ciprofloxacin)
 - Steroids (*i.e.*, prednisone, Entocort[®])
 - Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Member must have a history of trial and failure, contraindication, or intolerance to Humira* for at least 3 months.
- **Initial Duration of Approval:** 6 months
- **Reauthorization criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Fistulizing Crohn's Disease** and the following criteria is met:

- Prescribed medication is Entyvio or Stelara.
- Member is 18 years of age or older.
- Member must have clinical documentation of Crohn's disease with actively draining fistulas.
- Member must have a history of trial and failure, contraindication, or intolerance to conventional treatments including all of the following for at least 3 months of each medication:
 - Antibiotics (*i.e.*, Metronidazole, Ciprofloxacin)

- Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Member must have a history of trial and failure, contraindication, or intolerance to Humira* for at least 3 months.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) in adults/pediatrics** and the following criteria is met:

- The prescribed medication is Ilaris
- Member is 2 years of age or older.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD in adults/pediatrics)** and the following criteria is met:

- The prescribed medication is Ilaris.
- Member is 2 years of age or older
- Member must meet ONE of the following:
 - Member has diagnosis confirmed by either genetic mevalonate kinase gene (MVK) or enzymatic (MKD) findings.
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**
 - Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.
- **Reauthorization Duration of Approval:** 12 months

Coverage may be provided with a diagnosis of **Familial Mediterranean Fever (FMF)** and the following criteria is met:

- The prescribed medication is Ilaris
- Member is 2 years of age or older
- Member must have a history of trial and failure, contraindication, or intolerance to colchicine for at least 3 months
- **Initial Duration of Approval:** 6 months
- **Reauthorization Criteria:**



Updated: 07/2018
PARP Approved: 09/2018

- Reauthorization benefit will be approved if there is documented, significant improvement with prior courses of treatment.

Reauthorization Duration of Approval: 12 months

*Humira, Enbrel, Xeljanz, and Kineret may require a prior authoriz

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

When criteria are not met, the request will be forwarded to a Medical Director for review. The physician reviewer must override criteria when, in their professional judgment, the requested medication is medically necessary.

**NON-TNF MEDICATIONS
PRIOR AUTHORIZATION FORM**

Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation as applicable to Gateway HealthSM Pharmacy Services. **FAX:** (888) 245-2049
If needed, you may call to speak to a Pharmacy Services Representative.
PHONE: (800) 392-1147 Monday through Friday 8:30am to 5:00pm

PROVIDER INFORMATION

Requesting Provider:	NPI:
Provider Specialty:	Office Contact:
Office Address:	Office Phone:
	Office Fax:

MEMBER INFORMATION

Member Name:	DOB:
Gateway ID:	Member weight: _____ pounds or _____ kg

REQUESTED DRUG INFORMATION

Medication:	Strength:
Frequency:	Duration:
Is the member currently receiving requested medication? <input type="checkbox"/> Yes <input type="checkbox"/> No	
Date Medication Initiated:	

Billing Information

This medication will be billed: at a pharmacy **OR**
 medically (if medically please provide a JCODE: _____)

Place of Service: Hospital Provider's office Member's home Other

Place of Service Information

Name:	NPI:
Address:	Phone:

MEDICAL HISTORY (Complete for ALL requests)

- 1) Will the medication be prescribed by a rheumatologist, dermatologist, gastroenterologist, oncologist, or ophthalmologist?
 Yes No

- 2) Which of the following diagnoses will the medication be used for
 - a. Rheumatoid Arthritis, if yes please answer the following questions:
 Yes No
 - i. Is member 18 years of age or older?
 Yes No
 - ii. Does the member have a history of failure, contraindication, or intolerance to a three-month trial with methotrexate, or another DMARD?
 Yes No
 - iii. Does the member have a history of failure, contraindication, or intolerance to a three-month trial with Xeljanz?
 Yes No

- iv. Does the member have a history of failure, contraindication, or intolerance to a 3-month trial with either Humira OR Enbrel with or without MTX unless there is clinical rationale for using a non-TNF over a TNF inhibitor?

Yes No

- v. If the request is for Actemra, please provide ALL of the following lab values:
ANC:

Platelet Count:

AST/ALT:

- vi. If request is for Olumiant, will prescriber attest that member will avoid use of live vaccines?

Yes No

- vii. If request is for Olumiant, please provide ALL of the following lab values:
Hemoglobin:

Absolute Lymphocytic Count

Absolute Neutrophil Count

AST/ALT:

- b. Polyarticular Juvenile Idiopathic Arthritis, if yes please answer the following questions:

Yes No

- i. Is the member 2 years of age or older?

Yes No

- ii. Does the member have a history of trial and failure, contraindication, or intolerance to a three-month trial with either Humira, OR Enbrel?

Yes No

- iii. Does member meet any of the following:
- The member has an AJC (active joint count) >0 and continued disease activity after 3 months of MTX or leflunomide.
 Yes No
 - The member has an AJC (active joint count) >0 and continued disease activity after 1 month of anakinra*.
 Yes No

- iv. If requested medication is Actemra, please provide ALL of the following lab values”
ANC:

Platelet Count:

ALT/AST:

- c. Systemic Juvenile Idiopathic Arthritis, if yes please answer the following questions:
 Yes No

- i. Is member 2 years of age or older?
 Yes No

- ii. Does the member have an MD global score of ≥ 5 and continued disease activity after 1 month of NSAIDs?
 Yes No

- iii. Does the member have continued disease activity after 2 weeks of steroid monotherapy?
 Yes No

- iv. Does the member have continued disease activity after 1 month of anakinra therapy?
 Yes No

- v. If the medication being requested is Actemra, please provide ALL of the following lab values:
ANC:

Platelet Count:

ALT/AST:

d. Plaque Psoriasis, if yes if yes please answer the following questions:

Yes No

i. If requested medication is Cosentyx, Taltz, Otelza, Siliq, Ilumya, or Tremfya; is member 18 years of age or older?

Yes No

ii. If requested medication is Stelara, is member 12 years of age or older?

Yes No

iii. Is there clinical documentation that member has a diagnosis of moderate to severe plaque psoriasis characterized by greater than or equal to 5% body surface involved or disease affecting crucial body areas such as hands, feet, face, or genitals?

Yes No

iv. Does the member have a therapeutic failure of a three or more month trial or a contraindication to ANY of the following:

o Psoralens with UVA light (PUVA) or UVB light

Yes No

o Systemic treatments including ONE of the following:

a. Immunomodulators (i.e. Methotrexate, Cyclosporine)

Yes No

b. Retinoids (i.e. Soriatane)

Yes No

v. Does the member have a history of trial and failure, contraindication, or intolerance to a three-month trial with either Humira*, OR Enbrel*?

Yes No

- vi. Specific for Otezla: Will medication be used in combination with a strong cytochrome P450 enzyme inducer (e.g. rifampin, phenobarbital, carbamazepine, phenytoin)?
 Yes No

- vii. Specific for Siliq: Does the member should have a diagnosis of Crohn's disease?
 Yes No

- viii. Specific for Siliq, Ilumya, and Tremfya: Will prescriber avoid use of live vaccines while on medication?
 Yes No

- e. Psoriatic Arthritis (PsA)
 Yes No
 - Is member age 18 years of age or older?
 Yes No

 - Does the member have moderately to severely active psoriatic arthritis which must include documentation of either active psoriatic lesions or documented history of psoriasis?
 Yes No

 - Does the member have a history of trial and failure, contraindication, or intolerance to ALL of the following?
 Yes No
 - Three-month trial with either Humira* OR Enbrel*
 - Three-month trial with Xeljanz*

 - Does the member have a history of trial and failure, contraindication, or intolerance to ALL of the following:
 Yes No
 - Member without axial disease:
 - Four- week trial each of at least 2 NSAIDs.
 - Eight week trial of methotrexate or other DMARD
 - Member with axial disease
 - Four- week trial each of at least 2 NSAIDs.
 - Member with psoriatic arthritis with enthesitis

- Four- week trial each of at least 2 NSAIDs.
- Specific for Otezla: The prescriber attests that medication will not be used in combination with a strong cytochrome P450 enzyme inducer (e.g. rifampin, phenobarbital, carbamazepine, phenytoin)?
 Yes No

f. Cytokine Release Syndrome (CRS)

Yes No

i. Is the member 2 years of age or older?

Yes No

ii. Does the member meet one of the following criteria for Cytokine Release Syndrome (CRS):

○ Grade 1 CRS with persistent or refractory fever

Yes No

○ Grade 2 with any ONE of the following:

a. Hypotension refractory to fluid boluses

Yes No

b. Presence of hypoxia.

Yes No

c. Organ toxicity.

Yes No

○ Is there documented diagnosis of chimeric antigen receptor (CAR) T cell-induced severe (grade 3) CRS with any ONE of the following:

a. Defined as: oxygen requirement $\geq 40\%$.

Yes No

b. Hypotension requiring high dose or multiple vasopressors.

Yes No

c. Grade 3 organ toxicity or grade 4 transaminitis.

Yes No

- o Life threatening (grade 4) CRS with any ONE of the following:
 - a. Defined as: requirement for ventilator support.
 Yes No
 - b. Grade 4 organ toxicity (excluding transaminitis).
 Yes No

iii. Please provide ALL of the following lab values:
ANC:

Platelet count:

ALT/AST:

g. Giant Cell Arteritis

Yes No

i. Is the member 18 years of age or older?

Yes No

ii. Is the diagnosis confirmed by ONE of the following:

o Temporal artery biopsy or cross-sectional imaging
 Yes No

o Acute-phase reactant elevation (i.e. high erythrocyte sedimentation rate and/ or high serum C-reactive protein)
 Yes No

o Does the member have a history of trial and failure, contraindication, or intolerance to a one-week trial of glucocorticoids (e.g., prednisone, methylprednisone)?
 Yes No

o Please provide the following lab values:
ANC:

Platelet Count:

ALT/AST

h. Cyropyrin-Associated Periodic Syndrome (CAPS)

Yes No

i. Specific for Arcalyst

o Is the member 12 years of age or older?

Yes No

o Does the member have genetic evidence of a CIAS1 (NLRP3) mutation based on DNA sequencing?

Yes No

o Does the member have documented signs and symptoms associated with ONE of the following conditions:

a. Familial Cold Autoinflammatory Syndrome (FCAS).

i. Recurrent, intermittent fever

Yes No

ii. Rash that is often exacerbated by exposure to generalized cool ambient temperature

Yes No

b. Muckle-Wells Syndrome (MWS)

i. Rash of waxing and waning intensity sometimes exacerbated by exposure to generalized cool ambient temperature.

Yes No

ii. Specific for Ilaris

o Is the member 12 years of age or older?

Yes No

- Does the member have genetic evidence of a CIAS1 (NLRP3) mutation based on DNA sequencing?
 Yes No

- Does the member have documented signs and symptoms associated with ONE of the following conditions:
 - a. Familial Cold Autoinflammatory Syndrome (FCAS).
 - i. Recurrent, intermittent fever
 Yes No

 - ii. Rash that is often exacerbated by exposure to generalized cool ambient temperature
 Yes No

 - b. Muckle-Wells Syndrome (MWS)
 - i. Rash of waxing and waning intensity sometimes exacerbated by exposure to generalized cool ambient temperature.
 Yes No

 - iii. Specific for Kineret
 - Does the member have a documented diagnosis of Neonatal-Onset Multisystem Inflammatory Disease (NOMID)?
 Yes No

- i. Ankylosing Spondylitis
 Yes No
 - i. Is the member 18 years of age or older?
 Yes No

- Does the member have a history of trial and failure, contraindication, or intolerance to ALL of the following?
 Yes No
 - Three-month trial with either Humira* OR Enbrel*
 - Four-week trial each of at least 2 NSAIDs

j. Ulcerative Colitis

Yes No

i. Is member 18 years of age or older?

Yes No

ii. Does the member have a history of trial and failure, contraindication, or intolerance to conventional treatments including all of the following for at least 3 months:

Yes No

- Aminosalicylates, 5-ASAs (*i.e.*, Sulfasalazine, Pentasa[®], Asacol[®], Colazal[®])
- Steroids (*i.e.*, prednisone, Entocort[®])
- Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Humira

k. Crohn's Disease

Yes No

i. Is member 18 years of age or older?

Yes No

ii. Does the member have a history of trial and failure, contraindication, or intolerance to conventional treatments including two or more of the following for at least 3 months of each medication:

Yes No

- Aminosalicylates, 5-ASAs (*i.e.*, Sulfasalazine, Pentasa[®], Asacol[®], Colazal[®])
- Antibiotics (*i.e.*, Metronidazole, Ciprofloxacin)
- Steroids (*i.e.*, prednisone, Entocort[®])
- Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Humira

l. Fistulizing Crohn's Disease

Yes No

i. Is member 18 years of age or older?

Yes No

ii. Is there clinical documentation of Crohn's disease with actively draining fistulas?

Yes No

iii. Does the member have a history of trial and failure, contraindication, or intolerance to conventional treatments including ALL of the following for at least 3 months of each medication?

Yes No

- Antibiotics (*i.e.*, Metronidazole, Ciprofloxacin)
- Immunomodulators (*i.e.*, Azathioprine, 6-Mercaptopurine, Methotrexate)
- Humira

m. Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) in adults/pediatrics

Yes No

i. Is the member 2 years of age or older?

Yes No

n. Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD in adults/pediatrics)

Yes No

i. Is the member 2 years of age or older?

Yes No

ii. Has the member's diagnosis been confirmed by either genetic mevalonate kinase gene (MVK) or enzymatic (MKD) findings?

o. Familial Mediterranean Fever (FMF)

Yes No

i. Is the member 2 years of age or older?

Yes No

ii. Does the member have a history of trial and failure, contraindication, or intolerance to colchicine for at least 3 months?

Yes No

p. Other: _____

CURRENT or PREVIOUS THERAPY

Medication Name	Strength/ Frequency	Dates of Therapy	Status (Discontinued & Why/Current)

REAUTHORIZATION

1) For reauthorization of Rheumatoid Arthritis?

a. Has the member experienced a significant improvement with treatment?

Yes No

Please describe:

b. If request is for Olumiant, will prescriber attest that member will avoid use of live vaccines?
 Yes No

c. If request is for Olumiant, please provide ALL of the following lab values:
Hemoglobin:

Absolute Lymphocytic Count:

Absolute Neutrophil Count:

AST/ALT:

2) For reauthorization of Polyarticular Juvenile Idiopathic Arthritis?

a. Is there documented, significant improvement in AJC (active joint count) with prior courses of treatment?
 Yes No
Please describe:

3) For reauthorization of Plaque Psoriasis?

a. Is there clinical documentation that supports a decrease in percent of body surface area involvement when compared to baseline?
 Yes No

b. Please provide the following:
i. Baseline Body Surface Area Involvement:

ii. Current Body Surface Area Involvement:

4) For all other diagnoses:

a. Has the member experienced a significant improvement with treatment?

Yes No

Please describe:

SUPPORTING INFORMATION or CLINICAL RATIONALE

Prescribing Provider Signature

Date