



Samaritan  
Health Plans

# Prior Authorization Criteria

Samaritan Choice

**PLEASE READ: This document contains information about the criteria for coverage for this plan.**

Updated on 07/01/2024. For more recent information or other questions, please contact Pharmacy Services at **541-768-4550** or toll free **800-832-4580** (TTY 800-735-2900) or visit **[samhealthplans.org](https://www.samhealthplans.org)**. Pharmacy Services is available Monday through Friday, from 8 a.m. to 5 p.m.

# Abatacept (ORENCIA)

## Products Affected

- ORENCIA
- ORENCIA CLICKJECT

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Adult Rheumatoid Arthritis (RA)** Diagnosis of moderately to severely active rheumatoid arthritis **AND** trial and failure, contraindication, or intolerance to ONE nonbiologic disease-modifying antirheumatic drug (DMARD) (e.g., methotrexate [Rheumatrex/Trexall], Arava [leflunomide], Azulfidine [sulfasalazine]) **AND** trial and failure, contraindication, or intolerance to TWO of the following, or attestation demonstrating a trial may be inappropriate Cimzia (certolizumab pegol) adalimumab Rinvoq (upadacitinib) Simponi (golimumab) Xeljanz/XR (tofacitinib/ER).

**Polyarticular Juvenile Idiopathic Arthritis (PJIA)** Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, **AND** trial and failure, contraindication, or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate : Enbrel (etanercept), adalimumab, Xeljanz (tofacitinib).

**Psoriatic Arthritis (PsA)** Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to **TWO** of the following: Cimzia (certolizumab pegol), adalimumab, Simponi (golimumab) Stelara (ustekinumab), Tremfya (guselkumab), Skyrizi (risankizumab-rzaa), Rinvoq (upadacitinib), Xeljanz/XR (tofacitinib/ER).

**Prophylaxis for Acute Graft versus Host Disease (aGVHD)** Used for prophylaxis of acute graft versus host disease (aGVHD) **AND** patient will receive hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor **AND** recommended antiviral prophylactic treatment for Epstein-Barr Virus (EBV) reactivation (e.g., acyclovir) will be administered prior to Orencia **AND** continued for six

## Criteria Details

	months after HSCT <b>AND</b> Used in combination with both of the following: calcineurin inhibitor (e.g., cyclosporine, tacrolimus) methotrexate.
<b>Age Restrictions</b>	<b>aGVHD:</b> 2 years of age or older
<b>Prescriber Restrictions</b>	<b>RA, PJIA:</b> Prescribed by or in consultation with a rheumatologist <b>PsA:</b> Prescribed by or in consultation with a dermatologist or rheumatologist
<b>Coverage Duration</b>	<b>RA, PJIA, PsA: Initial:</b> 6 months. <b>Renewal:</b> 12 months <b>aGVHD: Initial:</b> 2 months. <b>Renewal:</b> N/A
<b>Renewal Criteria</b>	Documented positive clinical response to therapy  <b>RA, PJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.  <b>PsA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Abemaciclib (VERZENIO)

## Products Affected

- VERZENIO TAB 50MG
- VERZENIO TAB 100MG
- VERZENIO TAB 150MG
- VERZENIO TAB 200MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Breast Cancer- Early:</b> Hormone receptor (HR) positive <b>AND</b> HER2 negative breast cancer <b>AND</b> Node-positive disease at high risk or recurrence <b>AND</b> Patient is using in combination with anastrozole, exemestane, or letrozole OR Patient is using in combination with tamoxifen.</p> <p><b>Breast Cancer – Recurrent or Metastatic:</b> HR positive <b>AND</b> HER2 negative breast cancer <b>AND</b> Recurrent or metastatic breast cancer diagnosis <b>AND</b> Patient has tried chemotherapy for metastatic breast cancer <b>AND</b> Medication will be used in combination with anastrozole, exemestane, or letrozole OR medication will be used in combination with fulvestrant OR medication will be used as monotherapy.</p>
<b>Age Restrictions</b>	Must be at least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in collaboration with an oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months; <b>Renewal:</b> 3 months
<b>Renewal Criteria</b>	Clinical documentation of provider follow-up indicating safety <b>AND</b> efficacy with medication adherence over previous approval duration
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Adalimumab & Biosimilars

## Products Affected

- Adalimumab-adaz- **40 MG/0.4ML (auto-injector and prefilled syringe)**
- Adalimumab-fkjp - **40MG/0.8ML, 20 MG/0.4ML, (auto-injector and prefilled syringe)**
- Hadlima **40/0.4M, 40/0.8ML (auto-injector and prefilled syringe)**
- Yusimry **40/0.8ML**

## Prior Authorization Criteria

### Criteria Details

<p><b>Required Medical Information</b></p>	<p><b>Rheumatoid arthritis (RA):</b> Diagnosis of moderately to severely active RA <b>AND</b> trial and failure, contraindication, or intolerance to one non-biologic disease-modifying antirheumatic drug (DMARD) [e.g., methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)].</p> <p><b>Polyarticular Juvenile idiopathic arthritis (PJIA):</b> Diagnosis of moderate to severely active polyarticular JIA <b>AND</b> trial and failure, contraindication, or intolerance to one of the following non-biologic disease-modifying antirheumatic drugs (DMARDs): Arava (leflunomide), methotrexate (Rheumatrex/Trexall).</p> <p><b>Psoriatic arthritis (PsA):</b> Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.</p> <p><b>Ankylosing spondylitis (AS):</b> Diagnosis of active ankylosing spondylitis <b>AND</b> trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen)</p> <p><b>Crohn’s disease (CD):</b> Diagnosis of moderately to severely active Crohn’s disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220. <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional</p>
--	--

## Criteria Details

	<p>therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate</p> <p><b>Ulcerative Colitis (UC):</b> Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone).</p> <p><b>Plaque Psoriasis (PP):</b> Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement. <b>AND</b> a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.</p> <p><b>Hidradenitis Suppurativa (HS):</b> Diagnosis of moderate to severe hidradenitis suppurativa (i.e., Hurley Stage II or III)</p> <p><b>Uveitis (UV):</b> Diagnosis of non-infectious uveitis <b>AND</b> uveitis is classified as one of the following: intermediate, posterior or panuveitis.</p>
<p><b>Age Restrictions</b></p>	
<p><b>Prescriber Restrictions</b></p>	<p><b>RA, PJIA, AS:</b> Prescribed by or in consultation with a rheumatologist  <b>PsA:</b> Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist.  <b>CD, UC:</b> Prescribed by or in consultation with a gastroenterologist  <b>PsO, HS:</b> Prescribed by or in consultation with a dermatologist  <b>UV:</b> Prescribed by or in consultation with one of the following: ophthalmologist or rheumatologist</p>
<p><b>Coverage Duration</b></p>	<p><b>RA, PJIA, PsA, PsO, AS, CD, HS, UV Initial:</b> 6 months; <b>Renewal:</b> 12 months  <b>UC: Initial:</b> 12 weeks; <b>Renewal:</b> 12 months</p>
<p><b>Renewal Criteria</b></p>	<p><b>RA, PJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.</p>

## Criteria Details

**PsA:** Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.

**AS:** Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.

**CD:** Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.

**UC:** One of the following: For patients who initiated adalimumab therapy within the past 12 weeks: Documentation of clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on adalimumab therapy for longer than 12 weeks: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.

**PsO:** Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.

**HS, UV:** Documentation of positive clinical response to therapy.

<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Alpelisib (VIJOICE)

## Products Affected

- Vioice TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmed diagnosis of PROS <b>AND</b> has at least one severe clinical manifestation of PROS <b>AND</b> has a PIK3CA mutation that is confirmed by genetic testing
<b>Age Restrictions</b>	At least 2 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a provider who specializes in treatment of genetic disorders
<b>Coverage Duration</b>	<b>Initial:</b> 24 weeks. <b>Renewal:</b> 6 months.
<b>Renewal Criteria</b>	Documentation of a reduction in volume from baseline in at least one lesion <b>AND</b> an improvement in at least one symptom of PROS from baseline
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Ambrisentan (LETAIRIS)

## Products Affected

- AMBRISENTAN

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension (PAH) World Health Organization (WHO Group 1) confirmed by right heart catheterization <b>OR</b> patient is currently on any therapy for the diagnosis of PAH <b>AND</b> documented failure or incomplete response to or being co-prescribed with tadalafil
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.  <i>Note: Letairis (ambrisentan) has a black box warning for embryo-fetal toxicity. Because of the risks of birth defects, Letairis is available for females only through a special restricted distribution program under a Risk Evaluation and Mitigation Strategy (REMS).</i>
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Apremilast (OTEZLA)

## Products Affected

- OTEZLA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Plaque Psoriasis (PsO):</b> Diagnosis of plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement <b>AND</b> a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.</p> <p><b>Psoriatic Arthritis (PsA):</b> Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.</p> <p><b>Oral Ulcers Associated with Behçet’s Disease:</b> Diagnosis of Behçet’s Disease <b>AND</b> Patient has active oral ulcers.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p><b>Plaque psoriasis:</b> Prescribed by or in consultation with a dermatologist.</p> <p><b>Psoriatic arthritis:</b> Prescribed by or in consultation with a dermatologist or rheumatologist.</p>
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	<p><b>PsO:</b> Documentation of positive clinical response to therapy as evidenced by <b>ONE</b> of the following: Reduction the body surface area (BSA) involvement from baseline <b>OR</b> improvement in symptoms (e.g., pruritus, inflammation) from baseline.</p> <p><b>PsA:</b> Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms</p> <p><b>Behçet’s Disease:</b> Documentation of positive clinical response to therapy (e.g., reduction in pain from oral ulcers or reduction in number of oral ulcers).</p>

## Criteria Details

**Effective Date**

**P&T Approval Date**

**P&T Revision Date**

# Aripiprazole (ABILIFY)

## Products Affected

- ARIPIPRAZOLE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Diagnosed with Major Depressive Disorder (MDD):</b></p> <ul style="list-style-type: none"> <li>• Failure or intolerance to at least 2 first line antidepressant agents (sertraline, citalopram, venlafaxine, bupropion)</li> <li>• Prescribed for concurrent use with an antidepressant             <ul style="list-style-type: none"> <li>○ <b>OR</b> patient is new to the plan (request within 3 months of starting on plan) <b>AND</b> has been receiving aripiprazole therapy for greater than 4 weeks.</li> </ul> </li> <li>• One of the following if requesting oral solution:             <ul style="list-style-type: none"> <li>○ Pediatric member age 10 or under</li> <li>○ Documentation inability of the member to use the preferred tablet formulation</li> </ul> </li> </ul> <p><b>Diagnosed with schizophrenia, Bipolar I Disorder with acute manic or mixed episodes, irritability due to autism spectrum disorder, Tourette's disorder, Attention-deficit/hyperactivity disorder (ADHD), or Conduct disorder:</b></p> <ul style="list-style-type: none"> <li>• One of the following:             <ul style="list-style-type: none"> <li>○ Failure or intolerance to at least one of the following generic atypical antipsychotics</li> <li>○ Both of the following                 <ul style="list-style-type: none"> <li>▪ Patient is new to plan</li> <li>▪ Patient has been receiving aripiprazole therapy with success</li> </ul> </li> </ul> </li> <li>• One of the following if requesting oral solution:             <ul style="list-style-type: none"> <li>○ Pediatric member age 10 or under</li> <li>○ Documentation inability of the member to use the preferred tablet formulation</li> </ul> </li> </ul>
<b>Age Restrictions</b>	<p><b>For diagnoses of Irritability in autism, conduct disorder, or Tourette's disorder:</b> Patient is 6 years of age or older</p> <p><b>For diagnosis of ADHD:</b> Patient is 8 years of age or older</p> <p><b>For diagnosis of Bipolar I disorder:</b> Patient is 10 years of age or older</p>

## Criteria Details

	<b>For diagnosis of Schizophrenia:</b> Patient is 13 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Asciminib (SCEMBLIX)

## Products Affected

- SCEMBLIX

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Philadelphia positive CML that has been treated with at least two other TKIs OR Philadelphia positive CML with the T3151 mutation <b>AND</b> ECOG performance status of 0 or 1
<b>Age Restrictions</b>	Patient must be 18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by a hematologist or oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Belzutifan (WELIREG)

## Products Affected

- WELIREG

### Prior Authorization Criteria

Criteria Details	
<b>Required Medical Information</b>	Confirmed diagnosis of Von Hippel-Lindau disease with VHL alteration confirmation <b>AND</b> require therapy for either associated renal cell carcinoma, associated pancreatic neuroendocrine tumors, or associated CNS hemangioblastoma <b>AND</b> confirmation that patient is not eligible currently for surgery <b>AND</b> Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Beumosudil (REXUROCK)

## Products Affected

- REZUROCK

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosed with chronic graft-versus-host disease (cGVHD) <b>AND</b> who have tried and failed of at least two prior lines of systemic therapy for cGVHD <b>AND</b> not currently taking Imbruvica (ibrutinib)
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by oncologist or transplant specialist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 6 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Bimekizumab-bkzx (BIMEZELX)

## Products Affected

- Bimzelx

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Plaque Psoriasis</b></p> <ul style="list-style-type: none"> <li>• Severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool)</li> <li>• One or more of the following:             <ul style="list-style-type: none"> <li>○ Both of the following:                 <ul style="list-style-type: none"> <li>▪ At least 10% of body surface area involved</li> <li>▪ Hand, foot, face, or mucous membrane involvement</li> </ul> </li> <li>○ The patient on a current biologic product and experiencing intolerable side effects</li> </ul> </li> <li>• The patient tried and failed or have contraindications to ALL of the following?             <ul style="list-style-type: none"> <li>○ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.)</li> <li>○ At least one other topical agent: calcipotriene, tazarotene, anthralin, tar, etc.</li> <li>○ PUVA or UVB Phototherapy</li> <li>○ Methotrexate</li> <li>○ At least 1 other second line systemic agent such as cyclosporine or acitretin</li> </ul> </li> <li>• The patient tried and failed BOTH first line agents (infliximab or biosimilar AND Humira or biosimilar)</li> </ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.

## Criteria Details

<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy as evidenced by ONE of the following: <ul style="list-style-type: none"><li>• Reduction of body surface area (BSA) involvement from baseline</li><li>• Improvement in symptoms (e.g. pruritus, inflammation) from baseline</li></ul> Evidence of functional improvement
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024
<b>P&amp;T Revision Date</b>	

# Binimetinib (MEKTOVI)

## Products Affected

- Mektovi

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Diagnosis of unresectable or metastatic melanoma or Non-Small Cell Lung Cancer (NSCLC) :</b> <ul style="list-style-type: none"><li>• Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li><li>• Used in combination with encorafenib</li><li>• Trial and failure, contraindication or intolerance to one of the following:<ul style="list-style-type: none"><li>○ Cotellic</li><li>○ Mekinist</li></ul></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Bosentan (TRACLEER)

## Products Affected

- BOSENTAN

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Pulmonary arterial hypertension (PAH):</b> Diagnosed with PAH WHO Group 1 confirmed by right heart catheterization. Documentation of NYHA Functional Classification II, III, or IV symptoms <b>AND</b> documented normal liver function tests prior to initiation.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Bosutinib (BOSULIF)

## Products Affected

- Bosutinib capsules
- Bosutinib tablets

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Chronic Myelogenous/Myeloid Leukemia:</b> <ul style="list-style-type: none"><li>• Diagnosis of Philadelphia chromosome-positive chronic myelogenous/myeloid leukemia (Ph+ CML) AND</li><li>• One of the following:<ul style="list-style-type: none"><li>○ Disease is in the accelerated or blast phase OR</li><li>○ Disease is in the chronic phase and patient is 1 year of age or older</li></ul></li><li>• One of the following:<ul style="list-style-type: none"><li>○ Trial and failure or intolerance to generic imatinib</li><li>○ Continuation of prior therapy</li></ul></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Brexipiprazole (REXULTI)

## Products Affected

- REXULTI

## Prior Authorization Criteria

Criteria Details	
<b>Required Medical Information</b>	<b>Major Depressive Disorder (MDD):</b> A diagnosis of MDD <b>AND</b> prior treatment failure (at least 3 weeks) of or contraindication to 3 prior antidepressants <b>AND</b> one antipsychotic FDA approved as adjunct treatment for MDD <b>AND</b> to be used concurrently with an antidepressant. <b>Schizophrenia:</b> A diagnosis of schizophrenia <b>AND</b> prior treatment failure with a minimum of 2 antipsychotics <b>AND</b> Vraylar.
<b>Age Restrictions</b>	<b>Schizophrenia:</b> Aged 13 or older <b>Major Depressive Disorder (MDD):</b> Aged 18 or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of treatment success <b>AND</b> continued need for Rexulti.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Brigantinib (ALUNBRIG)

## Products Affected

- ALUNBRIG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment supported for the diagnosis in NCCN guidelines. Treatment being used according to FDA indication
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by oncologist or hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months unless otherwise specified in drug specific criteria <b>Renewal:</b> Up to 12 months
<b>Renewal Criteria</b>	Clinical documentation showing continued adherence <b>AND</b> toleration with lack of disease progression..
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# BUTRANS, BUPRENORPHINE PATCH, BELBUCA

## Products Affected

- BELBUCA
- BUPRENORPHINE PATCH

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Cancer or End-of-Life Care: Patient is being treated for cancer related pain or pain associated with end-of-life:</b> Documented trial and failure of, scheduled short-acting opioid therapy <b>AND</b> documented trial and failure or contraindication to long-acting morphine sulfate therapy. Documented trial/failure of, or reason why fentanyl is not appropriate.</p> <p><b>Other Chronic Pain:</b> Documented above the line diagnosis, FDA indicated, or guideline supported condition. Documented severe chronic pain (greater than 3mo) that is severe enough to require around the clock analgesic therapy <b>AND</b> documented trial and failure or contraindication to short-acting opioid therapy <b>AND</b> documented trial and failure of, or contraindication to long-acting morphine sulfate therapy <b>AND</b> documented trial and failure of, or reason why fentanyl is not appropriate.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial cancer/end of life:</b> 12 months. <b>Initial non-cancer/end of life:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# C1 Inhibitor (CINRYZE)

## Products Affected

- CINRYZE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Hereditary angioedema (HAE):</b> Diagnosed with HAE <b>AND</b> prescribed for routine prophylaxis against angioedema attacks.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by an immunologist, allergist, or rheumatologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Cabozantinib (CABOMETYX)

## Products Affected

- CABOMETYX TAB 20MG
- CABOMETYX TAB 40MG
- CABOMETYX TAB 60MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment is supported for the diagnosis in NCCN guidelines. <b>For HCC:</b> trial and failure or a contraindication to either Stivarga or Cyramza. Cabometyx is first line for RCC and DTC
<b>Age Restrictions</b>	Aged 12 years and older
<b>Prescriber Restrictions</b>	Treatment being prescribed or supervised by a hematologist, or oncologist as appropriate for the type of cancer.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Clinical documentation showing continued adherence and toleration of Cabometyx with lack of disease progression.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ceritinib (ZYKADIA)

## Products Affected

- Zykadia

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment supported for the diagnosis in NCCN guidelines. Treatment being used according to FDA indication
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by oncologist or hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months unless otherwise specified in drug specific criteria <b>Renewal:</b> Up to 12 months
<b>Renewal Criteria</b>	Clinical documentation showing continued adherence <b>AND</b> toleration with lack of disease progression.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Certolizumab Pegol (CMIZIA)

## Products Affected

- Cimzia

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Crohn's Disease (CD):** Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220. **AND** trial and failure, contraindication, or intolerance to ONE of the following conventional therapies: 6-mercaptopurine, Azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone), Methotrexate.

**Rheumatoid Arthritis (RA):** Diagnosis of moderately to severely active RA **AND** trial and failure, contraindication or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine.

**Psoriatic Arthritis (PsA):** Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.

**Ankylosing Spondylitis (AS):** Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses.

**Plaque Psoriasis (PsO):** Diagnosis of moderate to severe plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement **AND** a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.

## Criteria Details

	<p><b>Non-radiographic Axial Spondyloarthritis (nr-axSpA):</b> Diagnosis of active non-radiographic axial spondyloarthritis <b>AND</b> patient has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) <b>AND</b> minimum duration of one month trial and failure, contraindication, or intolerance to two different NSAIDs (e.g., ibuprofen, naproxen) at maximally tolerated doses</p>
<p><b>Age Restrictions</b></p>	
<p><b>Prescriber Restrictions</b></p>	<p><b>CD:</b> Prescribed by or in consultation with a gastroenterologist  <b>RA, AS, nr-axSpA:</b> Prescribed by or in consultation with a rheumatologist  <b>PsA:</b> Prescribed by or in consultation with one of the following:  Dermatologist or Rheumatologist  <b>PsO:</b> Prescribed by or in consultation with a dermatologist</p>
<p><b>Coverage Duration</b></p>	<p><b>CD: Initial:</b> 16 weeks; <b>Renewal:</b> 12months  <b>RA, PsA, AS, PsO, nr-axSpA: Initial:</b> 6 months; <b>Renewal:</b> 12 months</p>
<p><b>Renewal Criteria</b></p>	<p><b>CD:</b> Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.</p> <p><b>RA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.</p> <p><b>PsA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.</p> <p><b>AS, nr-axSpA:</b> Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial</p>

### Criteria Details

status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.

**PsO:** Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.

**Effective Date**

**P&T Approval Date**

**P&T Revision Date**

# Clobazam (ONFI)

## Products Affected

- Clobazam 10mg Tablets
- Clobazam 20mg Tablets
- Clobazam 2.5mg/mL suspension

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Lennox-Gastaut Syndrome</b> <ul style="list-style-type: none"><li>• Confirmed diagnosis.</li></ul> <b>Refractory Seizures</b> <p>Documentation showing appropriate trial of 2 or more tolerated anticonvulsant therapies.</p>
<b>Age Restrictions</b>	<b>Solution only</b> <p>One of the following:</p> <ul style="list-style-type: none"><li>• Pediatric member age 10 or under</li><li>• Documentation inability of the member to use the preferred tablet formulation</li></ul>
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Compounds (standard criteria for all compounded medications)

## Products Affected

- All compounded medications

### Prior Authorization Criteria

## Criteria Details

<b>Required Medical Information</b>	<ul style="list-style-type: none"> <li>• Each active ingredient in the compounded drug is FDA-approved or national compendia* supported for the condition being treated.</li> <li>• The requested amounts are supported by national compendia* or two peer-reviewed literature for the condition being treated in the requested route of delivery.</li> <li>• If any prescription ingredients require prior authorization and/or step therapy, all drug-specific criteria must be also met.</li> <li>• The patient has tried and failed therapy or had an intolerance to two FDA-approved commercially-available prescription therapeutic alternatives, one of which is the same route of administration as the requested compound, unless one of the following criteria are met:             <ul style="list-style-type: none"> <li>○ Patient has a contraindication to commercially available products</li> <li>○ Only one or no other therapeutic alternatives are commercially available</li> <li>○ Prepared strength(s) is/are not commercially available or currently in short supply</li> <li>○ Prepared in a different dosage form for a patient who is unable to take the commercially available formulation (mixing or reconstituting commercially available products based on the manufacturer's instructions or the product's approved labeling does NOT meet this criteria).</li> <li>○ Patient has an allergy or sensitivity to inactive ingredients (e.g. dyes, preservatives, sugars, etc.) that are found in commercially available products.</li> </ul> </li> </ul>
<b>Age Restrictions</b>	



## Criteria Details

<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Continuous Glucose Monitor (CGM)

## Products Affected

- DEXCOM G6 & G7 SYSTEMS
- FREESTYLE LIBRE SYSTEMS

## Prior Authorization Criteria

### Criteria Details

<p><b>Required Medical Information</b></p>	<p>Patient has documented diagnosis of type 1 or type 2 diabetes mellitus. Patient must have ALL of the following:</p> <ul style="list-style-type: none"> <li>• Intensive insulin regimen (3 or more insulin injections per day, or use of continuous subcutaneous insulin infusion pump).</li> <li>• Patient consistently monitors blood glucose 3 or more times per day.</li> <li>• Patient is motivated and knowledgeable about use of continuous glucose monitoring, is adherent to diabetic treatment plan, and participates in ongoing education and support.</li> <li>• Patient must have 1 OR more of the following:             <ul style="list-style-type: none"> <li>○ Dawn phenomenon, known or suspected, Hypoglycemic unawareness (i.e., patient does not have symptoms with hypoglycemia).</li> <li>○ Nocturnal hypoglycemia, known or suspected.</li> <li>○ Postprandial hyperglycemia, known or suspected.</li> <li>○ Significant change to diabetes treatment regimen (e.g., initiation of insulin, change from multiple-dose insulin to insulin pump therapy).</li> </ul> </li> </ul> <p>Unexplained hyperglycemia.</p>
<p><b>Age Restrictions</b></p>	
<p><b>Prescriber Restrictions</b></p>	
<p><b>Coverage Duration</b></p>	<p><b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.</p>
<p><b>Renewal Criteria</b></p>	<p>Documentation of positive clinical response to therapy.</p>
<p><b>Effective Date</b></p>	<p>01/01/2024</p>
<p><b>P&amp;T Approval Date</b></p>	<p>11/14/2023</p>

## Criteria Details

P&T Revision Date	
-------------------	--

# Crizotinib (XALKORI)

## Products Affected

- XALKORI

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>NSCLC:</b> Confirmed diagnosis of ALK-positive NSCLC or NSCLC with ROS1 rearrangement. <b>ALCL:</b> Confirmed diagnosis of ALK positive ALCL <b>AND</b> Trial <b>AND</b> failure of at least one prior systemic therapy
<b>Age Restrictions</b>	<b>NSCLC:</b> 18 years of age or older. <b>ALCL:</b> 1 year and older.
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Cyclosporine ophthalmic emulsion (RESTASIS)

## Products Affected

- RESTASIS
- RESTASIS MULTIDOSE

### Prior Authorization Criteria

#### Criteria Details

<b>Required Medical Information</b>	The patient has a diagnosis of lack of tear production due to ocular inflammation associated with keratoconjunctivitis sicca <b>AND ONE</b> of the following: The patient is not currently using a topical ophthalmic anti-inflammatory drug or punctal plug <b>OR</b> the patients current use of topical ophthalmic anti-inflammatory drug or punctal plug will be discontinued before starting the requested agent <b>AND</b> the patient has previously tried or is currently using aqueous enhancements (e.g. artificial tears, gels, ointments) <b>OR</b> the patient has a documented intolerance, contraindication, or hypersensitivity to aqueous enhancements <b>AND</b> the patient is not currently using Xiidra <b>OR</b> the patients current use of Xiidra will be discontinued before starting Restasis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Deutetrabenazine (AUSTEDO)

## Products Affected

- Austedo 6mg TAB
- Austedo 9mg TAB
- Austedo 12mg TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Chorea associated with Huntington’s Disease:</b> Documentation of the degree of chorea present and the impact on functional ability and/or quality of life as a baseline <b>AND</b> documentation of mental status, specifically depression and suicidality.</p> <p><b>Tardive Dyskinesia:</b> Clinical documentation of tardive dyskinesia including 1) At least one month of past or current exposure to a dopamine receptor blocker, 2) Dyskinetic or dystonic involuntary movements, 3) Exclusion of other causes of abnormal movements <b>AND</b> clear documentation that tardive dyskinesia causes functional impairment <b>AND</b> documentation of the degree of tardive dyskinesia with the AIMS scale as a baseline <b>AND</b> one of the following: tried and failed an 8-week trial of at least 2 other agents within the same therapeutic category at a clinically effective and maximally tolerated dose for the patient’s primary neuropsychiatric diagnosis <b>OR</b> evidence the medications precipitating tardive dyskinesia are medically necessary <b>AND</b> trial and failure or contraindication to clonazepam and amantadine.</p>
<b>Age Restrictions</b>	Age 18 and older
<b>Prescriber Restrictions</b>	<b>Huntington’s Disease:</b> neurologist <b>Tardive Dyskinesia:</b> neurologist or psychiatrist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	<b>Huntington’s Chorea:</b> clinical response such as improvement in chorea, ability to perform ADLs, reduction in falls, or increase in quality of life. <b>AND</b> documentation of continued monitoring of mental status specifically for depression and suicidality.

### Criteria Details

	<b>Tardive Dyskinesia:</b> Follow-up AIMS assessment showing improvement from Baseline <b>AND</b> documented improvement in functioning such as ability to perform ADLs, reduction in falls and increase in quality of life.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Dimethyl Fumarate (TECFIDERA)

## Products Affected

- DIMETHYL FUMARATE
- DIMETHYL FUMARATE STARTER PACK

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Multiple sclerosis:</b> Patient is diagnosed with relapsing forms of multiple sclerosis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by a neurologist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Direct-Acting Antivirals (use in Hepatitis C)

## Products Affected

- LEDIPASVIR-SOFOSBUVIR
- SOFOSBUVIR-VELPATASVIR
- MAVYRET

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Treatment of Hepatitis C:</b></p> <ul style="list-style-type: none"><li>• Expected survival from non-HCV-associated morbidities more than 1 year.</li><li>• Must have all pretreatment testing completed: including genotype, HBV, HIV, and cirrhosis status.</li><li>• Care must be provided by or in consultation with a specialist (hepatologist, gastroenterologist, or infectious disease specialist).</li><li>• Attestation that the patient and provider will comply with case management to promote the best possible outcome for the patient and adhere to monitoring requirements required by the Oregon Health Authority, including measuring and reporting of a posttreatment viral load OR attestation from the patient and provider that they have opted out of OHA case management. Case management includes assessment of treatment barriers and offer of patient support to mitigate potential barriers to regimen adherence as well as facilitation of SVR12 evaluation to assess treatment success.</li><li>• Documentation if the patient has a GT 1a infection or GT 3 infection and the patient had a baseline NS5a resistance test that documents a resistant variant to Elbasvir/grazoprevir or Daclatasvir + sofosbuvir. Note: Baseline NS5A resistance testing is required.</li><li>• Documentation of the prescribed regimen includes a NS3/4a protease inhibitor (glecaprevir, simeprevir, paritaprevir, voxilaprevir).</li><li>• Documentation if the patient has moderate-severe hepatic impairment (Child-Pugh B or Child-Pugh C).</li></ul> <p>Documentation if the prescribed regimen for the retreatment after failure of a DAA due to noncompliance or loss of follow-up and the prescribed</p>
-------------------------------------	--

### Criteria Details

	drug regimen is a recommended regimen based on the patient's genotype, age, treatment status (retreatment or treatment naive) and cirrhosis status.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a hepatologist, gastroenterologist, or infectious disease specialist
<b>Coverage Duration</b>	<b>Initial:</b> 2-4 months.
<b>Renewal Criteria</b>	
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Disposable Insulin Pump (OMNIPOD)

## Products Affected

- OMNIPOD 5
- OMNIPOD DASH

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Insulin dependent diabetes mellitus – pediatric (under age 18)</b></p> <ul style="list-style-type: none"> <li>• Documentation of Type 1 Diabetes Mellitus or Diabetes with C-reactive protein levels indicating insulin dependence.</li> <li>• On intensive insulin therapy (&gt;3 daily insulin injections) requiring frequent self-adjustments for at least 6 months prior to initiation of the insulin pump.</li> <li>• Documentation self-testing of blood glucose at least 4 times per day during the previous 2 months</li> <li>• Evidence of completion of a comprehensive diabetes education program in the last 12 months (member or caregiver/parent).</li> </ul> <p><b>Insulin dependent diabetes mellitus – adult</b></p> <ul style="list-style-type: none"> <li>• All of the above pediatric requirements AND</li> <li>• Documentation of 1 of the following:               <ul style="list-style-type: none"> <li>○ HbA1c &gt;7%</li> <li>○ History of recurring hypoglycemia</li> <li>○ Wide fluctuations in blood glucose before mealtime</li> <li>○ Dawn phenomenon with fasting blood sugars frequently exceeding 200mg/dL</li> <li>○ History of severe glycemic excursions</li> </ul> </li> <li>• Inability to use a traditional (non-disposable) insulin pump.</li> </ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 6 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy and in-person visit with provider within the last 6 months.

## Criteria Details

<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	01/09/024
<b>P&amp;T Revision Date</b>	

# Dupilumab (DUPIXENT)

## Products Affected

- DUPIXENT

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Moderate to severe Asthma with inadequate control of asthma symptoms with one of the following:** inhaled corticosteroids **AND** long acting beta2 agonist **OR** inhaled corticosteroids **AND** long-acting muscarinic antagonist.

**Atopic Dermatitis:** Diagnosed with severe atopic dermatitis **AND** patient is experiencing functional impairment (i.e., inability to use hands or feet for activities of daily living or significant facial involvement preventing normal social interaction) **AND** one or more of the following: At least 10% of body surface area involvement **OR** hand foot or mucous membrane involvement **AND** documented contraindication or failed trial to **ALL** of the following: failed, contraindicated or intolerance to a 12-week trial of a topical calcineurin inhibitor (e.g. tacrolimus) **AND** failed, contraindicated or intolerance to a 12-week trial of at least 2 prescription strength topical corticosteroids (e.g., betamethasone, fluticasone, mometasone) **AND** failed, contraindicated or intolerance to a 12-week trial of oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil, oral corticosteroids).

**Eosinophilic Esophagitis:** Confirmed diagnosis of EoE **AND** Weight  $\geq$  40 kg (88lbs) **AND** two or more episodes of dysphagia per week **AND** Inadequate response (8-week trial) or intolerance/contraindication to high-dose PPI therapy and inadequate response (8- to 12- week trial), intolerance, or contraindication to swallowed inhaled respiratory corticosteroid therapy.

**CRSwNP:** Diagnosis of CRSwNP, which includes objective evidence of the presence of bilateral nasal polyps **AND** not used in combination with other biologics for eosinophilic indications **AND** failure to adequately reduce symptoms after at least 2 months of saline nasal irrigations **AND** INCS use at doses appropriate for nasal polyp treatment **AND** systemic corticosteroid treatment for NPs at least once within the last 2 years or

## Criteria Details

	<p>prior NP removal surgery <b>AND</b> concomitant use of an INCS (unless not tolerated or contraindicated) while on biologic therapy</p> <p><b>Prurigo Nodularis:</b> Diagnosis of PN verified by a dermatologist <b>AND</b> has had the diagnosis of PN for at least 3 months <b>AND</b> severe or very severe itch (WI-NRS score <math>\geq 7</math>) reported within the past week <b>AND</b> at least 20 PN lesions in total on both leg and/or both arms and/or trunk.</p>
<b>Age Restrictions</b>	<p><b>Moderate to Severe Asthma:</b> 6 years and older</p> <p><b>Atopic Dermatitis:</b> 6 months and older</p> <p><b>Eosinophilic Esophagitis:</b> 12 years and older</p> <p><b>CRSwNP &amp; Prurigo Nodularis:</b> 18 years and older</p>
<b>Prescriber Restrictions</b>	<p><b>Atopic dermatitis:</b> Prescribed by a Dermatologist</p> <p><b>Eosinophilic Esophagitis:</b> Prescribed by Gastroenterologist or Immunologist</p> <p><b>CRSwNP:</b> Prescribed by ENT or Immunologist</p> <p><b>Prurigo Nodularis:</b> Prescribed by Dermatologist</p>
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Elagolix (ORLISSA)

## Products Affected

- ORLISSA TAB 150MG

- ORLISSA TAB 200MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of moderate to severe pain associated with endometriosis <b>AND</b> trial and failure, contraindication, or intolerance to a 3-month trial of prescription strength NSAIDs <b>AND</b> trial and failure, contraindication, or intolerance to two 3-month trials of hormonal therapies (eg combined oral contraceptives, progestins, or levonorgestrel IUD, etc.).  <b>Additional info required for 200 mg tablet twice daily:</b> documentation of coexisting dyspareunia
<b>Age Restrictions</b>	At least 18 years old but not yet through menopause
<b>Prescriber Restrictions</b>	Prescribed by obstetrician or gynecologist
<b>Coverage Duration</b>	<b>200MG dose: Initial:</b> 6 months; <b>Renewal:</b> No Renewals allowed <b>150MG dose: Initial:</b> 6 months; <b>Renewal:</b> 18months
<b>Renewal Criteria</b>	<b>150MG ONLY</b> Documentation of positive clinical response to therapy <b>AND</b> total therapy durations is less than 24 months.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Elexacaftor-tezacaftor-ivacaft (TRIKAFTA)

## Products Affected

- TRIKAFTA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical documentation of cystic fibrosis diagnosis with at least one F508del mutation (heterozygous or homozygous).
<b>Age Restrictions</b>	12 years of age <b>AND</b> older
<b>Prescriber Restrictions</b>	Prescribed by pulmonologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Encorafenib (BRAFTOVI)

## Products Affected

- Braftovi

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>BRAF V600E or V600K unresectable or metastatic melanoma:</b></p> <ul style="list-style-type: none"> <li>• Cancer is BRAF V600E or V600K mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li> <li>• Used in combination with Mektovi (binimetinib)</li> <li>• Trial and failure, contraindication or intolerance to one of the following:             <ul style="list-style-type: none"> <li>○ Zelboraf</li> <li>○ Tafinlar</li> </ul> </li> </ul> <p><b>Colorectal Cancer (CRC):</b></p> <ul style="list-style-type: none"> <li>• One of the following             <ul style="list-style-type: none"> <li>○ Unresectable or advanced disease</li> <li>○ Metastatic disease</li> </ul> </li> <li>• Patient has received prior therapy</li> <li>• Cancer is BRAFV600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li> <li>• Used in combination with Erbitux (cetuximab)</li> </ul> <p><b>Non-Small Cell Lung Cancer (NSCLC):</b></p> <ul style="list-style-type: none"> <li>• Cancer is BRAF V600E mutant type as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li> <li>• Used in combination with Mektovi (binimetinib)</li> </ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	

### Criteria Details

<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Entrectinib (ROZLYTREK)

## Products Affected

ROZLYTREK

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p>Non-Small Cell Lung Cancer (NSCLC):</p> <ul style="list-style-type: none"><li>• Patient has ROS1 rearrangement positive tumor(s)</li></ul> <p>Solid Tumors:</p> <ul style="list-style-type: none"><li>• Disease has neurotrophic tyrosine receptor kinase (NTRK) gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1, TPR-NTRK1, etc.) as detected by an FDA-approved test (THxID-BRAF Kit) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li><li>• Disease is without a known acquired resistance mutation (e.g., TRKA G595R, TRKA G667C or TRKC G623R substitutions)</li><li>• Disease is one of the following:<ul style="list-style-type: none"><li>○ Metastatic</li><li>○ Unresectable (including cases where surgical resection is likely to result in severe morbidity)</li></ul></li><li>• One of the following:<ul style="list-style-type: none"><li>○ Disease has progressed following previous treatment (e.g., surgery, radiation therapy, or systemic therapy) [3]</li><li>○ Disease has no satisfactory alternative treatments</li></ul></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	03/01/2024

<b>Criteria Details</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Epoetin Alpha (PROCRIT) (EPOGEN)

## Products Affected

- PROCRIT

### Prior Authorization Criteria

#### Criteria Details

<b>Required Medical Information</b>	<p><b>Anemia due to Chronic Kidney Disease (CKD):</b> Anemia with hematocrit less than 30% or hemoglobin less than 10g/dL within 30 days of request <b>AND</b> patient is on dialysis <b>OR</b> patient is not on dialysis but the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion <b>AND</b> reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal.</p> <p><b>Anemia in HIV Patients:</b> Anemia with hematocrit less than 36% or hemoglobin is less than 12 g/dL collected within 30 days of request, Serum erythropoietin less than or equal to 500mU/mL. Patient is receiving zidovudine therapy or diagnosed with HIV.</p> <p><b>Anemia due to Chemotherapy:</b> Anemia with hematocrit less than 30% &amp; hemoglobin less than 10 g/dL collected within the prior 2 weeks of request. All other causes of anemia have been ruled out, cancer is a non-myeloid malignancy <b>AND</b> patient is concurrently on chemo <b>OR</b> will receive concomitant chemo for a minimum of 2 months <b>OR</b> anemia is caused by cancer chemo (will not be approved if patient is not receiving cancer chemotherapy).</p> <p><b>Preoperative for reduction of allogeneic blood transfusion:</b> Patient scheduled for an elective, non-cardiac, non-vascular surgery. Perioperative hemoglobin is greater than 10 to less than or equal to 13 g/dL <b>AND</b> patient is at high risk of blood loss <b>AND</b> patient is unwilling or unable to donate autologous blood pre-operatively.</p> <p><b>Anemia in Myelodysplastic Syndrome (MDS):</b> Diagnosis of MDS <b>AND</b> serum erythropoietin less than or equal to 500mU/mL <b>OR</b> diagnosis of transfusion dependent MDS.</p>
<b>Age Restrictions</b>	

## Criteria Details

<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Preop Initial:</b> 1 month.
<b>Renewal Criteria</b>	Patient has a documented continued need for therapy demonstrated by an improvement in the hematocrit <b>AND</b> hemoglobin levels or by a significant decrease in transfusion requirements.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Erenumbab (AIMOVIG)

## Products Affected

- **AIMOVIG 70MG/ML**
- **AIMOVIG 140MG/ML**

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Diagnosis of episodic migraines:</b> Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Diagnosis of chronic migraines:</b> Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months <b>AND</b> medication overuse headache has been considered and potentially offending medication(s) have been discontinued</p> <p style="text-align: center;"><b>AND</b></p> <p><b>Two of the following:</b> History of failure or contraindication (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine) <b>OR</b> history of failure or contraindications (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) <b>OR</b> history of failure or contraindication (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol <b>OR</b> history of failure or contraindication (after at least a two month trial) or intolerance to Atacand (candesartan) <b>AND</b> medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines</p>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity <b>AND</b> use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan,

### Criteria Details

sumatriptan)] has decreased since the start of CGRP therapy **AND** medication will not be used in combination with another CGRP inhibitor for preventive treatment of migraines.

\* **AND For Chronic Migraine only:** Patient continues to be monitored for medication overuse headache

**Effective Date**

**P&T Approval Date**

**P&T Revision Date**



# Etrasimod arginine (VELSIPITY)

## Products Affected

- VELSIPITY TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Ulcerative Colitis (UC):</b> <ul style="list-style-type: none"><li>• Documentation of moderate-to-severe ulcerative colitis</li><li>• The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following:<ul style="list-style-type: none"><li>○ Mesalamine, sulfasalazine OR</li><li>○ Mercaptopurine, azathioprine, OR</li><li>○ Corticosteroids (prednisone, methylprednisolone)</li></ul></li><li>• Trial and failure of both infliximab and adalimumab</li></ul>
<b>Age Restrictions</b>	Must be at least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in collaboration with a Gastroenterologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission.
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	01/09/2024
<b>P&amp;T Revision Date</b>	

# Etanercept (ENBREL)

## Products Affected

- ENBREL
- ENBREL SURECLICK

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>ALL:</b> must have a negative tuberculin test (TB)</p> <p><b>AS:</b> Patient has a documented diagnosis of ankylosing spondylitis. Clinical documentation showing an inadequate response, intolerance, or contraindication to one or more non-steroidal anti-inflammatory drugs NSAIDs (trial at maximum dose for at least 2-3 weeks before considering them as failures) or analgesic agents if NSAIDs do not completely control the pain, or sulfasalazine (if peripheral joint involvement is present).</p> <p><b>CD:</b> Clinical documentation showing an inadequate response, intolerance, or contraindication to budesonide, mesalamine, or corticosteroids, or non-biologic DMARDs (i.e., azathioprine, methotrexate, mercaptopurine. JIA: Clinical documentation showing inadequate response, intolerance, or contraindication to one or more NSAID <b>AND</b> one or more non-biologic DMARD (i.e., methotrexate, sulfasalazine).</p> <p><b>PP:</b> Patient has documented diagnosis of moderate to severe plaque psoriasis for at least 6 months with at least one of the following: Incapacitation due to plaque location (e.g., head and neck, palms, soles, or genitalia) <b>OR</b> Involvement of at least 10 percent of body surface area (BSA) <b>OR</b> Psoriasis Area and Severity Index (PASI) score of 12 or greater, <b>AND</b> patient is free of any clinically important active infections <b>AND</b> clinical documentation of inadequate or non-candidate to a 3-month minimum trial of at least 1 systemic agent (e.g., immunosuppressive, retinoic acid derivatives, and/or methotrexate, <b>AND</b> did not respond or non-candidate to a 3-month minimum trial of phototherapy.</p>
<b>Age Restrictions</b>	

## Criteria Details

<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Fezolinetant (VEOZAH)

## Products Affected

- VEOZAH TAB 45MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Vasomotor Symptoms (VMS):</b> <ul style="list-style-type: none"><li>• Diagnosis of moderate-to-severe VMS due to menopause</li><li>• Documented contraindication, intolerance, or inadequate response to at least 2 hormonal therapies AND</li></ul> Documented contraindication, intolerance, or inadequate response to two nonhormonal therapies (e.g., one SNRI and one SSRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Gynecologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of at least 50% reduction in VMS from baseline.
<b>Effective Date</b>	09/01/2023
<b>P&amp;T Approval Date</b>	7/11/2023
<b>P&amp;T Revision Date</b>	

# Fingolimod (GILENYA)

## Products Affected

- GILENYA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Fremanezumab-vfrm (AJOVY)

## Products Affected

- AJOVY

### Prior Authorization Criteria

#### Criteria Details

<b>Required Medical Information</b>	<p><b>Diagnosis of episodic migraines:</b> Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month</p> <p style="text-align: center;"><b>OR</b></p> <p><b>Diagnosis of chronic migraines:</b> Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months <b>AND</b> medication overuse headache has been considered and potentially offending medication(s) have been discontinued</p> <p style="text-align: center;"><b>AND</b></p> <p><b>Two of the following:</b> History of failure or contraindication (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine) <b>OR</b> history of failure or contraindications (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) <b>OR</b> history of failure or contraindication (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol <b>OR</b> history of failure or contraindication (after at least a two month trial) or intolerance to Atacand (candesartan) <b>AND</b> medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines</p>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity <b>AND</b> use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy <b>AND</b>

### Criteria Details

	medication will not be used in combination with another CGRP inhibitor for preventive treatment of migraines.  * <b>AND For Chronic Migraine only:</b> Patient continues to be monitored for medication overuse headache
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Fruquintinib (FRUZAQLA)

## Products Affected

- Fruzaqla

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Metastatic colorectal cancer (mCRC): <ul style="list-style-type: none"><li>• Patient has been previously treated with both of the following:<ul style="list-style-type: none"><li>○ Fluoropyrimidine-, oxaliplatin-, irinotecan-based chemotherapy (e.g., FOLFOX, FOLFIRI, FOLFOXIRI)</li><li>○ Anti-VEGF biological therapy (e.g., Avastin [bevacizumab], Zaltrap [ziv-aflibercept])</li></ul></li><li>• One of the following:<ul style="list-style-type: none"><li>○ Patient has RAS mutant tumors</li><li>○ Patient has RAS wild-type tumors<ul style="list-style-type: none"><li>▪ Patient has been previously treated with both of the following: An anti-EGFR biological therapy (e.g., Vectibix [panitumumab], Erbitux [cetuximab])</li><li>▪ One of the following:<ul style="list-style-type: none"><li>• Lonsurf [trifluridine/tipiracil]</li><li>• Stivarga [regorafenib]</li></ul></li></ul></li></ul></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Galcanezumab-gnlm (EMGALITY)

## Products Affected

- Emgality 100mg/mL

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of episodic cluster headache <b>AND</b> patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least 3 months <b>AND</b> medication will not be used in combination with another CGRP inhibitor.
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity <b>AND</b> Medication will not be used in combination with another CGRP inhibitor.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# General Oncology

## Products Affected

- AKEEGA TAB
- AUGTYRO TAB
- CALQUENCE TAB
- COTELLIC TAB
- EMCYT CAP
- EVEROLIMUS
- FRUZAQLA CAP
- GLEOSTINE CAP
- HYCAMTIN CAP
- IBRANCE CAP
- IBRANCE TAB
- KRAZATI TAB
- LENALIDOMIDE
- LEUKERAN TAB
- LONSURF TAB
- LYNPARZA TAB
- LYTGOBI TAB
- MEKINIST TAB
- NINLARO CAP
- ORSERDU
- OSGIVEO
- PEMAZYRE TAB
- RETEVMO TAB
- REZLIDHIA CAP
- SORAFENIB TAB
- SPRYCEL TAB
- TALZENNA CAP
- TAFINLAR CAP
- THALOMID CAP
- TRUQAP TAB
- TUKYSA
- VANFLYTA TAB
- VONJO CAP
- VOTRIENT TAB
- ZOLINZA TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Medication is being used for an FDA approved age <b>AND</b> medication is being used for FDA approved indication <b>OR</b> Medication is being used according to National Comprehensive Cancer Network (NCCN) guidelines
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> up to 6 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	9/12/2023

# Glatiramer (GLATOPA)

## Products Affected

- GLATIRAMER INJ 20MG/ML
- GLATIRAMER INJ 40MG/ML

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> up to 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Glucagon-Like Peptide-1 (GLP1s) Receptor Agonist

## Products Affected

- BYDUREON BCISE
- BYETTA 10 MCG PEN
- BYETTA 5MCG PEN
- TRULICITY
- VICTOZA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Patient must have clinically diagnosed Type 2 Diabetes <b>AND</b> patient must have adequate trial of, or contraindication to an SGLT-2 if patient has HF or high risk/established ASCVD or a DPP-4 if no high risk/established ASCVD <b>AND</b> a maximal tolerated doses of metformin.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months, <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Golimumab (SIMPONI)

## Products Affected

- SIMPONI INJ 50/0.5ML
- SIMPONI INJ 100MG/ML

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Rheumatoid Arthritis (RA):</b> Diagnosis of moderately to severely active RA AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine <b>AND</b> patient is receiving concurrent therapy with methotrexate <b>OR</b> has a contraindication or intolerance to methotrexate.</p> <p><b>Psoriatic Arthritis (PsA):</b> Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.</p> <p><b>Ankylosing Spondylitis (AS):</b> Diagnosis of active ankylosing spondylitis <b>AND</b> minimum duration of one month trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen).</p> <p><b>Polyarticular Juvenile Idiopathic Arthritis (PJIA):</b> Diagnosis of moderate to severely active PJIA AND trial and failure, contraindication, or intolerance to one of the following non-biologic disease-modifying antirheumatic drugs (DMARDs): leflunomide, methotrexate.</p> <p><b>Ulcerative Colitis (UC):</b> Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids. AND trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone).</p>
<b>Age Restrictions</b>	

### Criteria Details

<b>Prescriber Restrictions</b>	<p><b>RA, AS, PJIA:</b> Prescribed by or in consultation with a rheumatologist</p> <p><b>PsA:</b> Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist</p> <p><b>UC:</b> Prescribed by or in consultation with a gastroenterologist</p>
<b>Coverage Duration</b>	<p><b>RA, PsA, AS, PJIA: Initial:</b> 6 months; <b>Renewal:</b> 12 months</p> <p><b>UC: Initial:</b> 10 weeks; <b>Renewal:</b> 12 months</p>
<b>Renewal Criteria</b>	<p><b>RA, PJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.</p> <p><b>PsA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.</p> <p><b>AS:</b> Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.</p> <p><b>UC:</b> Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.</p>
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Grass Pollen Allergen Extract – Timothy Grass (GRASTEK)

## Products Affected

- GRASTEK

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Grass pollen-induced allergic rhinitis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by Allergy and Immunology specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Guselkumab (TREMFYA)

## Products Affected

- TREMFYA INJ 100MG/ML

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Plaque Psoriasis (PsO):</b> Diagnosis of moderate-to-severe plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement <b>AND</b> a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.</p> <p><b>Psoriatic Arthritis (PsA):</b> Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p><b>PsO:</b> Prescribed by or in consultation with a dermatologist</p> <p><b>PsA:</b> Prescribed by or in consultation with one of the following: dermatologist or rheumatologist</p>
<b>Coverage Duration</b>	<b>Initial:</b> 6 months; <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	<p><b>PsO:</b> Documentation of positive clinical response to therapy as evidenced by <b>ONE</b> of the following: reduction the body surface area (BSA) involvement from baseline <b>OR</b> improvement in symptoms (e.g., pruritus, inflammation) from baseline</p> <p><b>PsA:</b> Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms (e.g., improvement in number of swollen/tender joints, pain, or stiffness).</p>
<b>Effective Date</b>	



## Criteria Details

**P&T Approval Date**

**P&T Revision Date**

# Ibrutinib (IMBRUVICA)

## Products Affected

- IMBRUVICA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Documentation of one of the following: <ul style="list-style-type: none"><li>• Diagnosis of Mantle Cell Lymphoma (MCL) <b>AND</b> patient has relapsed or is refractory to at least one prior therapy for the treatment of MCL.</li><li>• Diagnosis of Chronic Lymphocytic Leukemia (CLL) OR Small Lymphocytic Lymphoma (SLL).</li><li>• Diagnosis of Marginal Zone Lymphoma (MZL) <b>AND</b> patient has received at least one prior anti-CD20- based therapy.</li></ul> Diagnosis of Waldenstrms macroglobulinemia (WM) of Waldenstrms macroglobulinemia/lymphoplasmacytic lymphoma.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Infigratinib (TRUSELTIQ)

## Products Affected

- TRUSELTIQ (100MG DAILY DOSE)
- TRUSELTIQ (125MG DAILY DOSE)
- TRUSELTIQ (50MG DAILY DOSE)
- TRUSELTIQ (75MG DAILY DOSE)

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmation of trial and failure of guideline directed therapy.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Insulin Degludec (TRESIBA)

## Products Affected

- TRESIBA FLEXTOUCH U-100 & U-200

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>U-100 &amp; *U-200:</b> Must have tried and failed formulary long-acting insulin analogues <b>OR</b> have documented intolerance or contraindication to formulary long-acting insulin analogues <b>AND</b> have significant barriers to sardized administration requiring flexibility in dose timing.  <b>*(U-200)</b> Patient must require greater than 160 units of insulin per dose <b>AND</b> have difficulty with multiple daily injections.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Interferon Alfa-2b (INTRON A)

## Products Affected

- INTRON A

### Prior Authorization Criteria

#### Criteria Details

<b>Required Medical Information</b>	<p><b>Chronic hepatitis B:</b> Diagnosed with chronic hepatitis B infection <b>AND</b> patient is without decompensated liver disease.</p> <p><b>Chronic hepatitis C:</b> Diagnosed with chronic hepatitis C infection <b>AND</b> patient is without decompensated liver disease <b>AND</b> patient has not previously been treated with interferon <b>AND</b> is prescribed for use with ribavirin <b>OR</b> patient has intolerance or contraindication to ribavirin.</p> <p><b>Metastatic renal cell carcinoma (RCC):</b> Diagnosed with metastatic RCC <b>AND</b> prescribed in combination with Avastin (bevacizumab).</p> <p><b>AIDS-related Kaposi sarcoma (KS):</b> Diagnosed with AIDS-related KS.</p> <p><b>Condylomata acuminata (CA):</b> Diagnosed with CA involving external surfaces of the genital &amp; perianal areas.</p> <p><b>Follicular lymphoma (FL):</b> Diagnosed with clinically aggressive follicular non-Hodgkin lymphoma. Prescribed in conjunction with anthracycline-containing combination chemotherapy.</p> <p><b>Hairy cell leukemia (HCL):</b> Diagnosed with HCL. Melanoma: Diagnosed with malignant melanoma. Prescribed as adjuvant to surgical treatment who are free of disease but at high risk for systemic recurrence <b>AND</b> must be administered within 56 days of surgery.</p>
<b>Age Restrictions</b>	Patient must be 18 years or older.
<b>Prescriber Restrictions</b>	Prescribed by a specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.

## Criteria Details

<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Interferon beta-1a (AVONEX)

## Products Affected

- AVONEX PEN
- AVONEX PREFILLED

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of a relapsing form of Multiple Sclerosis <b>AND</b> trial and failure, contraindication, or intolerance to all of the following: dimethyl fumarate, fingolimod, glatiramer acetate/glatopa.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Interferon beta-1a (REBIF)

## Products Affected

- REBIF INJ 22/0.5ML
- REBIF INJ 44/0.5 ML
- REBIF REBIDO INJ 22/0.5ML
- REBIF REBIDO INJ 44/0.5ML

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis <b>AND</b> trial and failure, contraindication, or intolerance to all of the following: dimethyl fumarate, fingolimod, glatiramer acetate/glatopa, avonex.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Interferon beta-1b (EXTAVIA)

## Products Affected

- EXTAVIA INJ 0.3MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis <b>AND</b> trial and failure, contraindication, or intolerance to all of the following: dimethyl fumarate, fingolimod, glatiramer acetate/glatopa.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ivacaftor (KALYDECO)

## Products Affected

- KALYDECO

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of Cystic Fibrosis with documentation showing at least one CFTR gene mutation that has shown to be responsive to Kalydeco
<b>Age Restrictions</b>	6 months of age and older
<b>Prescriber Restrictions</b>	Prescribed by pulmonologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ivermectin (STROMEKTOL)

## Products Affected

- IVERMECTIN

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment of FDA approved diagnosis including Strongyloidiasis, Onchocerciasis, Infestation by Phthirus pubis, Scabies, Enterobiasis
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 6 months, <b>Renewals:</b> reinfection 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ivosidenib (TIBSOVO)

## Products Affected

- TIBSOVO

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Acute Myeloid Leukemia (AML):</b> Test confirmed IDH1 mutation <b>Cholangiocarcinoma:</b> Test confirmed IDH1 mutation <b>AND</b> previous treatment with at least one chemotherapy regimen (e.g. FOLFOX)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by an oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 6 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ixekizumab (TALTZ)

## Products Affected

- TALTZ

## Prior Authorization Criteria

### Criteria Details

**Required Medical Information**

**Plaque Psoriasis (PsO):** Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement **AND** a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar **AND** trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, adalimumab, Skyrizi, Stelara, or Tremya.

**Psoriatic Arthritis (PsA):** Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, Enbrel, adalimumab, Simponi, Stelara, Tremfya, Skyrizi, Rinvoq, or Xeljanz

**Ankylosing Spondylitis (AS):** Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to **TWO** non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen) **AND** trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, Enbrel, adalimumab, Simponi, Rinvoq, Xeljanz.

**Non-radiographic Axial Spondyloarthritis (nr-axSpA):** Diagnosis of active non-radiographic axial spondyloarthritis **AND** patient has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) **AND** minimum duration of one month trial and failure, contraindication, or intolerance to **TWO** non-steroidal anti-inflammatory drugs (NSAIDs) (e.g.,

Criteria Details	
	diclofenac, ibuprofen, meloxicam, naproxen) <b>AND</b> trial and failure, contraindication, or intolerance to Cimzia.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p><b>Plaque Psoriasis (PP):</b> Prescribed by or in consultation with a dermatologist</p> <p><b>Psoriatic Arthritis (PsA):</b> Prescribed by or in consultation with a dermatologist or rheumatologist</p> <p><b>Ankylosing Spondylitis (AS):</b> Prescribed by or in consultation with a rheumatologist</p> <p><b>Non-radiographic Axial Spondyloarthritis:</b> Prescribed by or in consultation with a rheumatologist</p>
<b>Coverage Duration</b>	<b>Initial:</b> 12 months; <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	<p><b>PsO:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.</p> <p><b>PsA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.</p> <p><b>AS, nr-axSpA:</b> Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.</p>
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Lacosamide (VIMPAT)

## Products Affected

- Lacosamide **TAB 50MG, 100MG, 150MG, 200MG**
- Lacosamide **Solution 10MG/ML**

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Focal seizures OR Primary generalized tonic-clonic seizures:</b> Documented epilepsy or seizure disorder
<b>Age Restrictions</b>	<b>Solution only</b> One of the following: <ul style="list-style-type: none"><li>• Pediatric member age 10 or under</li><li>• Documentation inability of the member to use the preferred tablet formulation</li></ul>
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	7/11/2023

# Lanthanum Carbonate (FOSRENOL)

## Products Affected

- Lanthanum carbonate 500MG
- Lanthanum carbonate 750MG
- Lanthanum carbonate 1000MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of hyperphosphatemia in chronic kidney disease <b>AND</b> trial and failure, contraindication, or intolerance (at least 6 weeks) to both maximally tolerated calcium acetate and sevelamer carbonate
<b>Age Restrictions</b>	6 years or older
<b>Prescriber Restrictions</b>	Nephrologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Lasmiditan (REYVOW)

## Products Affected

- REYVOW 100MG TAB
- REYVOW 50MG TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Medication intended for use for acute use for the treatment of migraine headaches <b>AND</b> documentation patient is on preventative therapy <b>AND</b> trial and failure (defined as trial period of 6 weeks per agent) or contraindication to at least 3 generic oral formulary triptans use at up to the maximally indicated dosing and in combination with NSAID therapy (naproxen) <b>OR</b> trial and failure to intolerance to NSAID treatment alone if triptans contraindicated <b>OR</b> contraindication to all triptans and NSAIDs
<b>Age Restrictions</b>	Patient is 18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by neurologist or headache specialist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documented positive clinical response to therapy
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Lenvatinib (LENVIMA)

## Products Affected

- LENVIMA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Endometrial carcinoma (EC):</b> Has advanced EC that is not microsatellite instability-high (MIS-H) or mismatch repair deficient (dMMR) <b>AND</b> has tried at least one systemic therapy <b>AND</b> is not a candidate for curative therapy.</p> <p><b>Hepatocellular Cancer (HCC):</b> Has unresectable or metastatic disease</p> <p><b>Renal Cell Carcinoma (RCC):</b> Has advanced disease <b>AND</b> is either being used in combination with Keytruda <b>OR</b> an everolimus.</p> <p><b>Thyroid Carcinoma, differentiated (DTC):</b> Diagnosed with differentiated thyroid carcinoma AMD disease is refractory to radioactive iodine therapy.</p>
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Lidocaine Topical Anesthetic (LIDODERM)

## Products Affected

- LIDOCAINE EXTERNAL PATCH 5%

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of post-herpetic neuralgia.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Lifitegrast (XIIDRA)

## Products Affected

- XIIDRA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p>The patient has a diagnosis of lack of tear production due to ocular inflammation associated with keratoconjunctivitis sicca <b>AND ONE</b> of the following:</p> <ul style="list-style-type: none"><li>• The patient is not currently using a topical ophthalmic anti-inflammatory drug or punctal plug <b>OR</b></li><li>• The patients current use of topical ophthalmic anti-inflammatory drug or punctal plug will be discontinued before starting the requested agent <b>AND</b> The patient has previously tried or is currently using aqueous enhancements (e.g. artificial tears, gels, ointments) <b>OR</b></li><li>• The patient has a documented intolerance, contraindication, or hypersensitivity to aqueous enhancements.</li></ul> <p>The patient is not currently using Restasis <b>OR</b> the patients current use of Restasis will be discontinued before starting Xiidra.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Linezolid (ZYVOX)

## Products Affected

- LINEZOLID
- LINEZOLID IN SODIUM CHLORIDE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinically documented infection that is susceptible to linezolid if the patient has a severe allergy to beta lactamase inhibitors or any antibiotic that the organism is susceptible <b>OR</b> clinically documented infection that is susceptible to linezolid if the patient has failed treatment with antibiotics that the organism is susceptible <b>OR</b> clinically documented Vancomycin-Resistant Enterococcus faecium infection <b>OR</b> clinically documented MRSA <b>AND</b> has failed or is intolerant to Vancomycin if the organism is susceptible to Vancomycin.
<b>Age Restrictions</b>	<p><b>Solution only</b> One of the following:</p> <ul style="list-style-type: none"> <li>• Pediatric member age 10 or under</li> <li>• Documentation inability of the member to use the preferred tablet formulation</li> </ul>
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an Infectious Disease specialist.
<b>Coverage Duration</b>	<b>Initial:</b> length of treatment. <b>Renewal:</b> length of treatment.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Lisdexamfetamine (VYVANSE)

## Products Affected

- VYVANSE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>ADHD:</b> Clinical documentation of trial and failure (defined as at least 6 weeks of treatment) of generic Adderall XR, generic Concerta, and generic Focalin XR in the treatment of ADHD. In cases of concern of stimulant abuse, must provide clinical documentation of trial and failure of one long-acting formulary stimulant <b>OR</b> clinical justification as to why formulary long-acting stimulants are contraindicated for the patient. <b>BED:</b> Clinical documentation confirming binge eating disorder diagnosis per DSM-5 criteria <b>AND</b> trial and failure of at least two therapeutic alternatives including SSRIs, topiramate, and/or methylphenidate.
<b>Age Restrictions</b>	<b>BED:</b> 18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Long Acting Opiates AND Dolophine

## Products Affected

- FENTANYL PATCH 72 HOUR 100 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 12 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 25 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 37.5 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 50 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 62.5 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 75 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 87.5 MCG/HR TRANSDERMAL
- HYDROCODONE BITARTRATE ER
- HYDROMORPHONE HCL ER
- METHADONE HCL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 120 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 30 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 45 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 60 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 75 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 90 MG ORAL
- MORPHINE SULFATE ER ORAL CAPSULE EXTENDED RELEASE 24 HOUR
- MORPHINE SULFATE ER ORAL TABLET EXTENDED RELEASE
- **NUCYNTA ER**
- OXYCODONE HCL ER
- **OXYCONTIN**
- OXYMORPHONE HCL ER
- **XTAMPZA ER**

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Cancer, end of life, or palliative care:</b> No coverage restrictions.</p> <p><b>Non-cancer/end of life care:</b> Documented use of current and/or recent usage of short-acting opioids for at least 15 days prior to long-acting opioids.</p> <ul style="list-style-type: none"> <li>• <b>For opioid naive (14 or fewer days filled in previous 120 days):</b> 7-day maximum quantity limit, equal to or less than 50 MED [morphine equivalents per day].</li> </ul>
-------------------------------------	---

### Criteria Details

	<ul style="list-style-type: none"><li>• <b>For opioid experienced (greater than or equal to 15 days filled in previous 120 days):</b> equal to or less than 90 MED [morphine equivalents per day]. Restricted to 2 fills in a 60-day period for both naive <b>AND</b> experienced.</li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Lorlatinib (LORBRENA)

## Products Affected

- LORBRENA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment being prescribed or supervised by a hematologist, or oncologist as appropriate for the type of cancer <b>AND</b> treatment supported for the diagnosis in NCCN guidelines <b>AND</b> treatment being used according to FDA indication <b>AND</b> trial and failure of one of the following agents: <ul style="list-style-type: none"><li>• For diagnosis of ALK-positive arrangement-positive NSCLC, no prior treatment:<ul style="list-style-type: none"><li>○ Alecensa (alectinib) OR Alunbrig (brigatinib)</li></ul></li><li>• For diagnosis of ALK-positive arrangement-positive NSCLC when the ALK-rearrangement is discovered during first-line systemic therapy: Alunbrig (brigatinib), OR Zykadia (ceritinib)</li></ul>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Clinical documentation showing continued adherence and toleration of Lorbrena with lack of disease progression
<b>Effective Date</b>	02/01/2022
<b>P&amp;T Approval Date</b>	01/11/2022
<b>P&amp;T Revision Date</b>	

# Lotilaner (XDEMVY)

## Products Affected

- Xdemvy 0.25% Ophthalmic solution

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Diagnosis:</b> Demodex Blepharitis <ul style="list-style-type: none"><li>• Documentation of at least mild erythema of the upper eyelid margin</li><li>• Presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination</li></ul>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Optometrist or Ophthalmologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 weeks. <b>Renewal:</b> No renewals allowed
<b>Renewal Criteria</b>	
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Lumacaftor/ivacaftor (ORKAMBI)

## Products Affected

- ORKAMBI ORAL PACKET
- ORKAMBI ORAL TABLET

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical documentation of cystic fibrosis diagnosis with homozygous F508del mutation.
<b>Age Restrictions</b>	2 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by pulmonologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Methylphenidate solution/chewable

## Products Affected

- METHYLPHENIDATE HCL ORAL SOLUTION
- METHYLPHENIDATE HCL ORAL TABLET CHEWABLE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Documentation that the patient has difficulty swallowing pills and/or has tried and failed methylphenidate tablets.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Midostaurin (RYDAPT)

## Products Affected

- Rydapt

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Acute Myeloid Leukemia: <ul style="list-style-type: none"><li>FMS-like tyrosine kinase 3 (FLT3) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test (e.g., LeukoStrat CDx FLT3 Mutation Assay) or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)</li><li>Used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation</li></ul> Diagnosis of Aggressive Systemic Mastocytosis, Systemic Mastocytosis with Associated Hematological Neoplasm, or Mast Cell Leukemia
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Patient does not show evidence of progressive disease while on therapy
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Mitapivat (PYRUKYND)

## Products Affected

- PYRUKYND

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of PKD with at least two mutations within the PKLR gene, including a missense mutation <b>AND</b> confirmation of current hemoglobin is $\leq 10\text{mg/dL}$ <b>AND</b> patient is not homozygous for the R479H mutation <b>AND</b> does not have two non-missense variants in the PKLR gene, without the presence of another missense variant <b>AND</b> patient has had at least 6 RBC transfusions within the previous year for hemolytic anemia due to PKD <b>AND</b> prescriber confirmed concomitant use of daily folic acid <b>AND</b> confirmation that the patient does not have moderate or severe hepatic dysfunction.
<b>Age Restrictions</b>	At least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in consultations with a hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 6 months.
<b>Renewal Criteria</b>	Clinical documentation showing an increase in Hb at least 1.5 mg/dL over baseline and/or a reduction in frequency of transfusions.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Mirikizumab-mrkz (OMVOH)

## Products Affected

- Omvoh

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>All diagnoses:</b></p> <ul style="list-style-type: none"><li>• Initial testing for latent TB and treatment, if necessary, before starting treatment.</li><li>• No current active infection at initiation of therapy.</li><li>• Risks and benefits documented in cases of chronic or recurrent infection.</li><li>• Will NOT be used in combination with another biologic or Otezla</li></ul> <p><b>Ulcerative Colitis (UC):</b></p> <ul style="list-style-type: none"><li>• Documentation of moderate-to-severe ulcerative colitis</li><li>• The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following:<ul style="list-style-type: none"><li>• Mesalamine, sulfasalazine OR</li><li>• Mercaptopurine, azathioprine, OR</li><li>• Corticosteroids (prednisone, methylprednisolone)</li><li>• Trial and failure of both infliximab and adalimumab</li></ul></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultations with a Gastroenterologist.
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024

## Criteria Details

P&T Revision Date	
-------------------	--



# Mobocertinib (EXKIVITY)

## Products Affected

- Exkivity

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Treatment supported for the diagnosis in NCCN guidelines <b>AND</b> treatment being used according to FDA indication <b>AND</b> prior trial and failure of contraindication to Rybrevant (amivantamab).
<b>Age Restrictions</b>	18 and older
<b>Prescriber Restrictions</b>	Oncologist or Hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Clinical documentation showing continued adherence and toleration with lack of disease progression
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Naltrexone (VIVITROL)

## Products Affected

- VIVITROL INJ.

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	The medication will be sent directly to the administering provider and will not be dispensed directly to the member
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Nilotinib (TASIGNA)

## Products Affected

- TASIGNA CAP 50MG
- TASIGNA CAP 150MG
- TASIGNA CAP 200MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Ph+ CML:</b> Newly diagnosed Ph+ CML in chronic phase <b>OR</b> Chronic phase <b>AND</b> accelerated phase Ph+ CML that is resistant or intolerant to prior therapy, including imatinib or the patient has a Sokal risk score >1.2 (High risk)
<b>Age Restrictions</b>	At least one year of age
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 3 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Omalizumab (XOLAIR)

## Products Affected

- Xolair

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Severe asthma:</b> Confirmed diagnosis of moderate to severe persistent asthma <b>AND</b> documentation of smoking status <b>AND</b> positive skin test or RAST to a perennial aeroallergen <b>AND</b> baseline IgE serum level within FDA label <b>AND</b> documentation of steps taken to avoid within reason environmental allergens and other triggers <b>AND</b> documented trial <b>AND</b> failure of the following: High dose inhaled corticosteroid with a long-acting beta agonist (e.g. Advair), Long acting anti-muscarinic (e.g., Spiriva), Leukotriene Inhibitor (e.g., Singulair) <b>AND</b> documented trial and failure of, or contraindication to allergen immunotherapy <b>AND</b> medical or claims history of compliance/adherence with prescribed asthma medications</p> <p><b>Nasal Polyps:</b> Documentation of recurrent nasal polyps after prior sinus surgery <b>AND</b> trial and failure of at least 2 intranasal corticosteroids <b>AND</b> Sinuva nasal implant <b>AND</b> documented adherence to a nasal corticosteroid with Xolair intended as adjunct therapy <b>AND</b> documented risk of another sinus surgery, or a statement why sinus surgery is not medically appropriate.</p> <p><b>Idiopathic chronic urticaria- refractory:</b> Documentation of chronic spontaneous or idiopathic urticaria <b>AND</b> documented trial and failure (including dose escalation of both first and second-generation antihistamines) for at least 6 weeks: (1<sup>st</sup> generation – doxepin, hydroxyzine), (2<sup>nd</sup> generation – Cetirizine, Levocetirizine, Fexofenadine, Loratadine, Desloratadine) <b>AND</b> documented trial and failure of an H2 antihistamine (e.g., Famotidine, Cimetidine) <b>AND</b> documented trial and failure (at least 4 weeks) of, or contraindication to a leukotriene inhibitor (e.g., Montelukast, Zafirlukast).</p>
<b>Age Restrictions</b>	<p><b>Asthma:</b> 6 years of age and older  <b>CIU:</b> 12 years of age and older  <b>Nasal Polyps:</b> 18 years of age and older</p>

### Criteria Details

<b>Prescriber Restrictions</b>	<b>Asthma:</b> Prescribed by or in consultation with a pulmonologist or immunologist <b>CIU:</b> Prescribed by or in consultation with an immunologist <b>Nasal Polyps:</b> Prescribed by or in consultation with an allergist or ENT
<b>Coverage Duration</b>	<b>SA/NP Initial:</b> 6 months. <b>Renewal:</b> 6 months. <b>CIU Initial:</b> 4 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Paliperidone (INVEGA HAFYERA)

## Products Affected

- INVEGA HAFYERA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical documentation of a diagnosis of schizophrenia <b>AND</b> trial and failure (defined by at least 6 months of treatment) of Invega Trinza <b>OR</b> Invega Sustenna <b>AND</b> clinical need or concern for adherence which could be improved upon with twice yearly dosing.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Pancrelipase (CREON) (PANCREAZE)

## Products Affected

- PANCREAZE
- CREON

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p>Confirmed diagnosis of cystic fibrosis <b>OR</b> history of pancreatectomy <b>OR</b> diagnosis of exocrine pancreatic cancer <b>OR</b> diagnosis of chronic pancreatitis confirmed by imaging <b>OR</b> confirmed diagnosis of pancreatic insufficiency confirmed with one of the following methods:</p> <ul style="list-style-type: none"> <li>• Steatorrhea with fecal fat determination <b>OR</b></li> <li>• Measurement of fecal elastase <b>OR</b> Secretin or CCK pancreatic function testing <b>OR</b></li> </ul> <p>Two of the following CFTR mutations (G542X, W1282X, R553X, 621+1G&gt;T, 1717-1G&gt;A, 3120+1G&gt;A, R1162X, 3659delC, 1898+1G&gt;A, 2184delA, 711+1G&gt;T, F508del, I507del, G551D, N1303K, R560T).</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# PCSK9 inhibitors

## Products Affected

- PRALUENT
- REPATHA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Clinical ASCVD:</b> Documented history of clinical ASCVD or has experienced a cardiovascular event <b>AND</b> documentation of a current LDL greater than or equal to 70 mg/dl <b>AND</b> documentation of at least one of the following: Patient is receiving maximally tolerated statin therapy (or is statin intolerant) <b>AND</b> is receiving ezetimibe or documented intolerance to ezetimibe</p> <p><b>Primary or familial hyperlipidemia:</b> Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL <b>AND</b> documentation of current LDL greater than 100 mg/dL <b>AND</b> documentation of at least one of the following: Patient is receiving maximally tolerated statin therapy (or is statin intolerant) <b>AND</b> is receiving ezetimibe or documented intolerance to ezetimibe.</p> <p><b>Homozygous Familial Hyperlipidemia:</b> Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL <b>AND</b> documentation of current LDL greater than 100 mg/dL <b>AND</b> Documentation of at least one of the following: Patient is receiving maximally tolerated statin therapy (or is statin intolerant) <b>AND</b> is receiving ezetimibe or documented intolerance to ezetimibe.</p>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	



## Criteria Details

**P&T Approval Date**

**P&T Revision Date**

# Peanut Powder (PALFORZIA)

## Products Affected

- PALFORZIA (12 MG DAILY DOSE)
  - PALFORZIA (120 MG DAILY DOSE)
  - PALFORZIA (160 MG DAILY DOSE)
  - PALFORZIA (20 MG DAILY DOSE)
  - PALFORZIA (200 MG DAILY DOSE)
  - PALFORZIA (240 MG DAILY DOSE)
  - PALFORZIA (3 MG DAILY DOSE)
  - PALFORZIA (300 MG MAINTENANCE)
  - PALFORZIA (300 MG TITRATION)
  - PALFORZIA (40 MG DAILY DOSE)
  - PALFORZIA (6 MG DAILY DOSE)
  - PALFORZIA (80 MG DAILY DOSE)
- PALFORZIA INITIAL ESCALATION

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmed positive skin test or peanut-specific serum IgE greater than 0.35 kUA/L Concurrent prescription with injectable epinephrine medical justification supports necessity for oral immunotherapy despite peanut avoidance.
<b>Age Restrictions</b>	Patient must be between 4 and 17 at therapy initiation
<b>Prescriber Restrictions</b>	Prescribed by allergist or immunologist enrolled in Palforzia REMS program
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Currently receiving medication byway of previously approved SHP authorization or documents showing <b>Initial</b> approval criteria was/has been met. For patients who required use of injectable epinephrine while on Palforzia, must have medical justification that supports continued need for Palforzia. If greater than 18 years old, must have medical justification that supports continued need for oral immunotherapy despite peanut avoidance and documentation that Initial dose escalation happened between age 4 and 17.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Pirtobrutinib (JAYPIRCA)

## Products Affected

- JAYPIRCA 50MG
- JAYPIRCA 100MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Mantle Cell Lymphoma (MCL):</b> <ul style="list-style-type: none"><li>• Diagnosis of MCL confirmed by histology</li><li>• Tried and failed at least 2 prior therapies of which one was with a BTKi</li><li>• ECOG performance status score of 0 to 2</li></ul> <b>Other Diagnosis</b> <ul style="list-style-type: none"><li>• Diagnosis as supported in National Comprehensive Cancer Network (NCCN) guidelines.</li></ul>
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 2 months. <b>Renewal:</b> 4 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	09/01/2023
<b>P&amp;T Approval Date</b>	07/11/2023
<b>P&amp;T Revision Date</b>	

# Pramlintide Acetate (SYMLIN)

## Products Affected

- SYMLINPEN 60
- SYMLINPEN 120

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p>Coverage is provided for the use of pramlintide as an adjunct treatment in type 1 and type 2 diabetic patients 18 or older who use mealtime insulin therapy <b>AND</b> who meet all of the following criteria:</p> <ul style="list-style-type: none"><li>• Are currently on mealtime insulin.</li><li>• Have an HbA1c less than or equal to 9%.</li><li>• Are monitoring blood glucose levels regularly <b>AND</b> reliably (3 or more times per day).</li><li>• Are capable of monitoring blood glucose levels pre- <b>AND</b> post-meals <b>AND</b> at bedtime.</li><li>• Have failed to achieve adequate control of blood glucose levels despite individualized management of their insulin therapy.</li></ul> <p>Are receiving ongoing care under the guidance of a health care provider skilled in use of insulin <b>AND</b> supported by the services of a diabetes educator.</p>
<b>Age Restrictions</b>	Patient must be 18 years or older.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Priftin (rifapentine)

## Products Affected

- Priftin

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Latent tuberculosis:</b> <ul style="list-style-type: none"><li>• Used in combination with isoniazid</li></ul> <b>Active tuberculosis:</b> <ul style="list-style-type: none"><li>• One of the following:<ul style="list-style-type: none"><li>○ The member does NOT have multidrug resistant disease</li><li>○ The member has multidrug resistant disease and the drug is prescribed by or in consultation with infectious disease specialist.</li><li>○ Priftin will be used as part of multi-drug regimen</li></ul></li><li>• The drug will NOT be prescribed by a county clinic with a state funded TB program(for these programs the drug is funded directly through the state).</li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Latent TB Initial:</b> 3 months. <b>Renewal:</b> N/A <b>Active TB Initial:</b> 6 months. <b>Renewal:</b> N/A
<b>Renewal Criteria</b>	
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024

# Ranolazine (RANEXA)

## Products Affected

- RANOLAZINE ER

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of chronic angina not controlled with other antianginal therapy. May be used with beta-blockers, nitrates, calcium channel blockers, anti-platelet therapy, lipid-lowering therapy, ACE inhibitors, and angiotensin receptor blockers.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Resmetirom (REZDIFFRA)

## Products Affected

- Rezdiffra 60MG/80MG/100MG TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<ul style="list-style-type: none"><li>• Diagnosis of metabolic dysfunction-associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH)</li><li>• Patient does not have cirrhosis (e.g. decompensated cirrhosis)</li><li>• Submission of medical records (e.g. chart notes) showing diagnosis has been confirmed by one of the following:<ul style="list-style-type: none"><li>○ FibroScan-aspartate aminotransferase (FAST)</li><li>○ MRI-aspartate aminotransferase (MAST)</li><li>○ Liver biopsy</li></ul></li><li>• Submission of medical records (e.g. chart notes) showing disease is fibrosis stage F2 or F3 as confirmed by one of the following:<ul style="list-style-type: none"><li>○ FibroScan</li><li>○ Fibrosis-4 index (FIB-4)</li><li>○ Magnetic resonance Elastography (MRE)</li></ul></li><li>• Presence of greater than or equal to 3 metabolic risk factors (e.g., Type 2 diabetes, hypertension, obesity)</li><li>• Submission of medical records (e.g. chart notes) confirming drug is used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)</li></ul>
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by Gastroenterologist; Hepatologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Patient demonstrates positive response to therapy (e.g., MASH resolution, fibrosis stage improvement, etc.) AND Submission of medical

### Criteria Details

	records (e.g., chart notes) confirming drug will continue to be used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program)
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024
<b>P&amp;T Revision Date</b>	



# Ribavirin (VIRAZOLE)

## Products Affected

- RIBAVIRIN INHALATION

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Respiratory Syncytial Virus (RSV) Infection:</b> Chart notes / written medical summary documenting diagnosis of RSV.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Request is initiated by an infectious disease specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 3 months.
<b>Renewal Criteria</b>	
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ribociclib Succinate (KISQALI)

## Products Affected

- KISQALI

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Breast Cancer in Women:</b> Has recurrent or metastatic disease <b>AND</b> has hormone receptor positive (HR+) <b>AND</b> has human epidermal growth factor receptor 2 (HER2)-negative breast cancer <b>AND</b> is either post-menopausal <b>OR</b> is pre/perimenopausal <b>AND</b> is receiving gonadotropin-releasing hormone agonist <b>OR</b> has had bilateral oophorectomy or ovarian irradiation <b>AND</b> medication will be used in combination with anastrozole, exemestane, or letrozole <b>OR</b> will be used in combination with fulvestrant.</p> <p><b>Breast Cancer in Men:</b> Has recurrent or metastatic disease <b>AND</b> has hormone receptor positive (HR+) <b>AND</b> has human epidermal growth factor receptor 2 (HER2)-negative breast cancer <b>AND</b> is receiving gonadotropin-releasing hormone analog <b>AND</b> medication will be used in combination with anastrozole, exemestane, or letrozole <b>OR</b> will be used in combination with fulvestrant.</p>
<b>Age Restrictions</b>	At least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in consultations with an oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Rifaximin (XIFAXAN)

## Products Affected

- XIFAXAN

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>IBS-D:</b> Diagnosis of IBS with diarrhea <b>Hepatic Encephalopathy:</b> must have one of the following: used as add-on therapy to lactulose <b>AND</b> unable to achieve an optimal clinical response with lactulose monotherapy <b>OR</b> a history of contraindication or intolerance to lactulose.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>IBS-D Initial:</b> 14 Days. <b>Renewal:</b> 30 Days. <b>HE Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Rimegepant (NURTEC)

## Products Affected

- NURTEC ODT

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Acute treatment:</b> Diagnosis of migraine with or without aura <b>AND</b> will be used for the acute treatment of migraine <b>AND</b> patient has fewer than 15 headache days per month <b>AND</b> trial and failure or intolerance to 3 generic triptans or contraindications to all triptans and NSAID combined treatment <b>OR</b> trial and failure or intolerance to NSAID treatment alone if triptans contraindicated <b>OR</b> contraindication to all triptans and NSAIDs <b>AND</b> medication will not be used in combination with another CGRP inhibitor <b>AND</b> if patient has 4 or more headache days per month, then patient must currently be treated with amitriptyline, venlafaxine, divalproex, topiramate, candesartan or a beta-blocker or have a contraindication or intolerance to all of these medications.</p> <p><b>Prophylaxis treatment:</b> Diagnosis of episodic migraines <b>AND</b> patient has 4 to 18 migraine days per month but no more than 18 headache days per month <b>AND</b> trial and failure of at least 2 months intolerance or contraindication of two of the following sets: 1) amitriptyline or venlafaxine; 2) divalproex or topiramate; 3) one of the following beta blocker: atenolol, propranolol, nadolol, timolol, or metoprolol; 4) candesartan <b>AND</b> not used in combination with another CGRP inhibitor.</p>
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist
<b>Coverage Duration</b>	<b>Acute Initial:</b> 3 months; <b>Renewal:</b> 12 months <b>Prophylaxis Initial:</b> 6 months; <b>Renewal</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy
<b>Effective Date</b>	

<b>Criteria Details</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Risankizumab (SKYRIZI)

## Products Affected

- SKYRIZI

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Plaque Psoriasis (PsO):</b> Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement <b>AND</b> a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.</p> <p><b>Psoriatic Arthritis (PsA):</b> Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.</p> <p><b>Crohn’s disease (CD):</b> Diagnosis of moderately to severely active Crohn’s disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220 <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate.*induction dose is IV, maintenance is subcutaneous.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p><b>PsA:</b> Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist.</p> <p><b>CD:</b> Prescribed by or in consultation with a gastroenterologist</p> <p><b>PsO:</b> Prescribed by or in consultation with a dermatologist</p>
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	<b>PsO:</b> Documentation of positive clinical response to therapy as evidenced by ONE of the following: Reduction in the body surface area

## Criteria Details

(BSA) involvement from baseline **OR** improvement in symptoms (e.g., pruritus, inflammation) from baseline.

**PsA:** Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.

**UC:** Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.

**Effective Date**

**P&T Approval Date**

**P&T Revision Date**

# Risdiplam (EVRYSDI)

## Products Affected

- EVRYSDI SOL 0.75MG/ML

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Spinal Muscular Atrophy (SMA):</b> <ul style="list-style-type: none"><li>• Confirmed (via genetic testing) diagnosis of 5q-autosomal recessive SMA (type 1, 2 or 3)</li><li>• Patient is not dependent on invasive ventilation or tracheostomy OR use of non-invasive ventilation beyond uses for sleeping</li><li>• Is not receiving concomitant chronic SMN modifying therapy such as Spinraza</li></ul> <p>Patient has not previously received gene replacement therapy for the treatment of SMA (e.g. Zolgensma)</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a neurologist with expertise in the treatment of SMA
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> up to 12 months.
<b>Renewal Criteria</b>	Documentation of clinical improvement from baseline in motor functionality confirmed by standard exams (e.g. BSID-III, CHOP INTEND, HINE-2, RULM test)
<b>Effective Date</b>	09/01/2023
<b>P&amp;T Approval Date</b>	7/11/2023
<b>P&amp;T Revision Date</b>	



# Sacubitril/Valsartan (ENTRESTO)

## Products Affected

- ENTRESO

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	The patient has a diagnosis of New York Heart Association class II to IV heart failure <b>AND</b> patient is receiving concomitant therapy with one of the following beta blockers: carvedilol, bisoprolol, sustained-released metoprolol, unless unable to tolerate or contraindicated <b>AND</b> patient will discontinue use of any concomitant *ACE inhibitor or ARB before initiating therapy.  <b>*ACE inhibitors must be discontinued at least 36 hours prior to ENTRESTO.</b>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Cardiologist or in consultation with a cardiologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Sargramostin (LEUKINE)

## Products Affected

- LEUKINE

### Prior Authorization Criteria

## Criteria Details

<b>Required Medical Information</b>	<p><b>Acute myelogenous leukemia (AML):</b> To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in older adults (greater than or equal to 55 years of age).</p> <p><b>Bone marrow transplant (allogeneic or autologous):</b> For graft failure or engraftment delay.</p> <p><b>Myeloid reconstitution after allogeneic bone marrow transplantation:</b> To accelerate myeloid recovery following transplantation in non-Hodgkin lymphoma (NHL), acute lymphoblastic leukemia (ALL), Hodgkin lymphoma. Febrile neutropenia Primary prophylaxis of neutropenia in patients receiving chemotherapy (outside transplant and AML) or who are at high risk for neutropenic fever.</p> <p><b>Peripheral stem cell transplantation:</b> Mobilization of hematopoietic progenitor cells for leukapheresis <b>AND</b> myeloid reconstitution following autologous peripheral stem cell transplantation.</p> <p><b>Acute Radiation Syndrome:</b> Treatment of radiation-induced myelosuppression of the bone marrow.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Requested by specialist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	

## Criteria Details

**P&T Approval Date**

**P&T Revision Date**

# Secukinumab (COSENTYX)

## Products Affected

- COSENTYX (200 MG DOSE)
- COSENTYX SENSOREADY (300 MG)

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**All:** patient has a negative tuberculin test (TB) prior to initiating therapy.

**PP:** Patient has documented diagnosis of moderate to severe plaque psoriasis for at least 6 months with at least one of the following: incapacitation due to plaque locations (e.g., head and neck, palms, soles, or genitalia), **OR** involvement of at least 10% of body surface area (BSA) **OR** psoriasis area **AND** severity index (PASI) score of 12 or greater, **AND** patient is free of any clinically important active infections, **AND** patient did not respond adequately (or is not a candidate) to a 3-month trial of at least 1 systemic agent **AND** patient did not respond adequately (or is not a candidate) to a 3 month trial of phototherapy **AND** patient has had a trial **AND** failure of adalimumab **AND** Enbrel with clinical documentation.

**PA:** Patient has active psoriatic arthritis for at least 6 months defined as: greater than 3 swollen joints **AND** greater than 3 tender joints **AND** patient has had an inadequate response, intolerance or contraindication (clinical documentation required) with the following, one or more non-steroidal anti-inflammatory drugs (NSAIDs) trialed at a max dose for at least 2 weeks **AND** one or more non-biologic disease modifying anti-rheumatic drugs **AND** patient has had a trial **AND** failure of adalimumab OR Xeljanz OR Orencia.

**AS:** Patient has had an inadequate response, intolerance or contraindication with the following, one or more non-steroidal anti-inflammatory drugs (NSAIDs) trialed at a max dose for at least 2 weeks **OR** analgesic agents if NSAIDs do not control pain OR sulfasalazine (if peripheral joint involvement is present).

**Hidradenitis Suppurativa (HS):**

- Documentation of one of the following:
  - Moderate to severe hidradenitis suppurative (Hurley Stage II or

## Criteria Details

	<p>Hurley Stage III)</p> <ul style="list-style-type: none"> <li>• Patient is on a current biologic product and experiencing intolerable side effects.</li> <li>• The patient is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of the following: <ul style="list-style-type: none"> <li>○ 90-day trial of conventional therapy (e.g. oral antibiotics)</li> <li>○ Trial and failure of both infliximab and adalimumab</li> </ul> </li> </ul>
<b>Age Restrictions</b>	Patient must be 18 years or older.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024
<b>P&amp;T Revision Date</b>	

# Semaglutide (WEYGOVY)

## Products Affected

- Wegovy

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Reduction of Major Adverse Cardiovascular Events</b> <ul style="list-style-type: none"><li>• Wegovy is being used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease and either obesity or overweight</li><li>• Wegovy is being used as adjunct to lifestyle modification (e.g. dietary or caloric restriction, exercise, behavioral support, community-based program)</li><li>• Patient has established cardiovascular disease as evidenced by one of the following: prior MI, prior stroke, peripheral arterial disease (e.g. intermittent claudication with ankle-brachial index &lt;0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease)</li><li>• BMI greater than or equal to 27 kg/m<sup>2</sup></li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	<ul style="list-style-type: none"><li>• Documentation of treatment success (BMI reduction of 5% or more)</li><li>• Documentation of continuation of lifestyle modification program with reduced calorie diet and regular physical activity alongside continuous Wegovy use (80% adherence)</li></ul>
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024

## Criteria Details

P&T Revision Date	
-------------------	--

# Sildenafil Citrate (REVATIO)

## Products Affected

- SILDENAFIL CITRATE INTRAVENOUS                      SILDENAFIL CITRATE ORAL TABLET
- SILDENAFIL CITRATE ORAL SUSPENSION  
RECONSTITUTED

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical diagnosis of pulmonary arterial hypertension (PAH).
<b>Age Restrictions</b>	<b>Solution only</b> One of the following: <ul style="list-style-type: none"><li>• Pediatric member age 10 or under</li><li>• Documentation inability of the member to use the preferred tablet formulation</li></ul>
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	5/1/2024
<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	3/12/2024



# Sodium-Glucose CO-Transporter 2 (Sglt2) Inhibitors

## Products Affected

- FARXIGA TABLET 10MG ORAL
- JARDIANCE TABLET 20MG ORAL
- JARDIANCE 25MG ORAL
- FARXIGA TABLET 5MG ORAL

## Prior Authorization Criteria

Criteria Details	
<b>Required Medical Information</b>	Trial and failure or contraindication to any of the following: metformin, glipizide-metformin, glyburide-metformin, pioglitazone-metformin OR one of the following generics or preferred brands: formulary ACEi or ARBs, spironolactone, eplerenone, or Entresto
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Somatropins

## Products Affected

- GENOTROPIN
- GENOTROPIN MINIQUICK
- HUMATROPE
- NORDITROPIN FLEXPRO
- NUTROPIN AQ NUSPIN 10
- NUTROPIN AQ NUSPIN 20
- NUTROPIN AQ NUSPIN 5
- OMNITROPE
- SAIZEN
- SAIZENPREP
- SEROSTIM
- ZORBTIVE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>For children:</b> Growth hormone deficiency in individuals less than 16 years of age or radiographic evidence of non-closure of epiphyseal plates. Appropriate medical work up: Assessment <b>AND</b> evaluation must indicate absolute growth less than 4.5 cm per year without growth hormone. Subnormal growth, greater than or equal to 2 standard deviations below the mean for age.</p> <p><b>For adults:</b> Biochemical diagnosis of adult growth hormone deficiency by means of a subnormal response to a standard growth hormone stimulation test (peak growth hormone Less than or equal to 5 mcg/L). Confirmatory testing may not be required in patients with congenital/genetic growth hormone deficiency or multiple pituitary hormone deficiencies due to organic diseases.</p> <p><b>Adult-onset:</b> Patients who have adult growth hormone deficiency whether alone or with multiple hormone deficiencies (hypopituitarism) as a result of pituitary disease, hypothalamic disease, surgery, radiation therapy, or trauma. Turners Syndrome in females is an approved indication.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.

## Criteria Details

<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Sotorasib (LUMAKRAS)

## Products Affected

- LUMAKRAS

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Trial and failure of at least one prior systemic therapy.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an oncologist.
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Sparsentan (FILSPARI)

## Products Affected

- FILSPARI TAB 200MG
- FILSPARI TAB 400MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Primary immunoglobulin A nephropathy:</b> <ul style="list-style-type: none"><li>Urine protein-to-creatinine ratio (UPCR) <math>\geq 1.5</math> and eGFR <math>\geq 30</math> mL/min/1.73 m<sup>2</sup></li><li>Biopsy-verified primary IgA nephropathy</li><li>No history of kidney transplant and not currently receiving dialysis</li></ul> Member has failed to achieve a reduction in proteinuria to under 1 gram/day while receiving maximally tolerated doses of an ACE inhibitor or ARB for at least 12 weeks
<b>Age Restrictions</b>	18 or older
<b>Prescriber Restrictions</b>	Nephrologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Improved or stable kidney function compared to baseline <b>OR</b> reduction in proteinuria
<b>Effective Date</b>	09/01/2023
<b>P&amp;T Approval Date</b>	7/11/2023
<b>P&amp;T Revision Date</b>	

# Tacrolimus (PROTOPIC)

## Products Affected

- TACROLIMUS EXTERNAL

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Atopic Dermatitis:</b> Clinically diagnosed moderate-to-severe atopic dermatitis (10 percent BSA, hand, foot or mucous membrane involvement, or functional impairment) <b>AND</b> trial and failure of topical steroids, UVB phototherapy, or reason why they would not be medically appropriate. <b>Psoriasis:</b> diagnosis of moderate to severe Psoriasis (10 percent BSA, hand, foot or mucous membrane involvement, or functional impairment) <b>AND</b> trial and failure or contraindication to a high potency topical corticosteroid and/or UVB phototherapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<b>Psoriasis:</b> prescribed by or in consultation with a dermatologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Tadalafil (ADCIRCA)

## Products Affected

- TADALAFIL (PAH)

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical diagnosis of pulmonary arterial hypertension (PAH).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Tazarotene (AVAGE) (TAZORAC)

## Products Affected

- TAZAROTENE 0.1% CREAM

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Psoriasis:</b> diagnosis of moderate to severe Psoriasis (10% BSA, hand, foot or mucous membrane involvement <b>AND</b> functional impairment) <b>AND</b> trial and failure of high potency topical corticosteroids or medical reason why they would be inappropriate  <b>Other FDA approved indications (i.e., severe acne):</b> trial and failure/contraindication to two formulary alternatives used to treat the approved indication.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Tenapanor (XPHOZA)

## Products Affected

- XPHOZA 20MG
- XPHOZA 30MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Add on therapy for hyperphosphatemia in chronic kidney disease, on chronic hemodialysis:</b> <ul style="list-style-type: none"><li>• On chronic maintenance hemodialysis 3x per week for at least 3 months or chronic peritoneal dialysis for a minimum of 6 months</li><li>• Other contributing factors for hyperphosphatemia have been addressed:<ul style="list-style-type: none"><li>○ On a stable dose of calcimimetic or active vitamin D for at least 4 weeks</li><li>○ Serum parathyroid hormone is &lt;1200 pg/mL</li><li>○ Dietary restrictions are in place to limit phosphate intake</li></ul></li><li>• Serum phosphate levels remain elevated despite use of at least 3 different phosphate binder agents (calcium acetate, sevelamer, lanthanum) used at therapeutic doses with adherence to therapy (at least 80% adherence based on fill history)</li><li>• Medication will be used as add-on therapy with phosphate binder treatment</li></ul>
<b>Age Restrictions</b>	Must be at least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in collaboration with a Nephrologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Clinical documentation of provider follow-up indicating safety and efficacy with medication adherence over previous approval duration
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	01/09/2024
<b>P&amp;T Revision Date</b>	

# Tepotinib (TEPMETKO)

## Products Affected

- TEPMETKO

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of NSCLC containing MET exon 14-skipping alterations <b>AND</b> no EGFR-activating mutations predictive of sensitivity to anti-EGFR therapy <b>AND</b> no ALK rearrangements predictive of sensitivity to anti-ALK therapy <b>AND</b> Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1.  <b>Maximum daily dose of 2 tablets</b>
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> up to 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Terbinafine Hydrochloride (LAMISIL)

## Products Affected

- TERBINAFINE HCL

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	For the treatment of onychomycosis of the toenail or fingernail due to dermatophytes (tinea unguium) <b>AND</b> patient is experiencing pain which limits normal activity (i.e., unable to wear shoes, difficulty walking, etc.) <b>OR</b> patient has diabetes <b>OR</b> patient has peripheral vascular diseases, <b>OR</b> patient is immunocompromised.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 3 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Tezacaftor/ivacaftor (SYMDEKO)

## Products Affected

- SYMDEKO

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Clinical documentation of cystic fibrosis diagnosis with homozygous F508del mutation.
<b>Age Restrictions</b>	6 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by a pulmonologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 6 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ticagrelor (BRILINTA)

## Products Affected

- BRILINTA 60MG
- BRILINTA 90MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Acute Coronary Syndrome</b> <ul style="list-style-type: none"><li>• Patient has acute coronary syndrome (ACS)<ul style="list-style-type: none"><li>○ Either NSTEMI-ACS or STEMI and has had a coronary stent implanted OR</li><li>○ Patient has NSTEMI-ACS and is treated with medical therapy alone (i.e., has not had revascularization)?</li></ul></li></ul>
<b>Age Restrictions</b>	18 or older
<b>Prescriber Restrictions</b>	Gynecologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	09/01/2023
<b>P&amp;T Approval Date</b>	7/11/2023
<b>P&amp;T Revision Date</b>	

# Tranexamic Acid

## Products Affected

• Tranexamic Acid 650MG TAB

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Hemophilia Diagnosis</b> <ul style="list-style-type: none"><li>• Intending to use for hemorrhage prophylaxis for tooth extraction(s)</li></ul> <b>Abnormal Uterine Bleeding</b> <ul style="list-style-type: none"><li>• Currently using or documented trial and failure or contradiction to ALL the following treatments:<ul style="list-style-type: none"><li>○ Cobombined Oral Contraceptive therapy</li><li>○ Profestin therapy (oral or LM) or Levonogrestrel IUD</li></ul></li><li>• NSAID therapy</li></ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by Hematologist, Hemophilia specialist, Dentist, Gynecologist
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024
<b>P&amp;T Revision Date</b>	

# Tivozanib (FOTIVDA)

## Products Affected

- FOTIVDA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of metastatic renal cell carcinoma <b>AND</b> tried and failed at least two systemic therapies with at least one including a VEGF-TKI <b>AND</b> Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1 <b>AND</b> maximum monthly dose of 21 per 28 days
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> up to 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Tocilizumab SC (ACTEMRA)

## Products Affected

- ACTEMRA INJ 80MG/4ML
- ACTEMRA INJ 200/10ML
- ACTEMRA INJ 400/20ML
- ACTEMRA INJ 162/0.9
- ACTEMRA INJ ACTPEN

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Rheumatoid Arthritis (RA):** Diagnosis of moderately to severely active rheumatoid arthritis **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine **AND** trial and failure, contraindication, or intolerance to **TWO** of the following, or attestation demonstrating a trial may be inappropriate: Cimzia (certolizumab pegol), Enbrel (etanercept), adalimumab, Rinvoq (upadacitinib), Simponi (golimumab), Xeljanz/XR (tofacitinib/ER).

**Systemic Juvenile Idiopathic Arthritis (SJIA):** Diagnosis of active systemic juvenile idiopathic arthritis **AND** trial and failure, contraindication, or intolerance to **ONE** of the following: Non-steroidal anti-inflammatory drug (NSAID) (e.g., ibuprofen, naproxen), Systemic glucocorticoid (e.g., prednisone), methotrexate.

**Polyarticular Juvenile Idiopathic Arthritis (PJIA):** Diagnosis of active polyarticular juvenile idiopathic arthritis **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, **AND** trial and failure, contraindication, or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), adalimumab, Xeljanz (tofacitinib).

**Giant Cell Arteritis (GCA):** Diagnosis of giant cell arteritis **AND** trial and failure, contraindication, or intolerance to a glucocorticoid.

**Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD):** Diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) as documented by the following: exclusion of other known causes of



## Criteria Details

	<p>interstitial lung disease (ILD) (e.g., domestic and occupational environmental exposures, connective tissue disease, drug toxicity) <b>AND</b> one of the following: In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (e.g., fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] <b>AND</b> centrilobular fibrosis) pattern on high-resolution computed tomography (HRCT) revealing SSc-ILD or probable SSc-ILD OR In patients subjected to a lung biopsy, both HRCT <b>AND</b> surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD.</p> <p><b>Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy:</b> Patient will receive or is receiving chimeric antigen receptor (CAR) T-cell immunotherapy (i.e., Kymriah [tisagenlecleucel], Yescarta [axicabtagene ciloleucel])</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p><b>RA, SJIA, PJIA, GCA:</b> Prescribed by or in consultation with a rheumatologist.</p> <p><b>SSc-ILD:</b> Prescribed by or in consultation with a pulmonologist or rheumatologist.</p> <p><b>CRS:</b> Prescribed by or in consultation with an oncologist or hematologist</p>
<b>Coverage Duration</b>	<p><b>RA, SJIA, PJIA, GCA, SSc-ILD: Initial:</b> 6 months; <b>Renewal:</b> 12 months</p> <p><b>Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy: Initial:</b> 2 months.</p>
<b>Renewal Criteria</b>	<p><b>RA, PJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen <b>AND</b> tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.</p> <p><b>SJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen <b>AND</b> tender) joint count from baseline, or improvement in clinical features or symptoms (e.g., pain, fever, inflammation, rash, lymphadenopathy, serositis) from baseline.</p> <p><b>GCA, SSc-ILD:</b> Documentation of positive clinical response to therapy.</p>
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Tofacitinib (XELJANZ)

## Products Affected

- XELJANZ
- XELJANZ XR

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Rheumatoid Arthritis (RA):** Diagnosis of moderately to severely active rheumatoid arthritis **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine **AND** patient has had an inadequate response or intolerance to one or more TNF inhibitors (e.g., Cimzia, adalimumab, Simponi) **AND** patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

**Psoriatic Arthritis (PsA):** Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** patient is will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

**Ankylosing Spondylitis (AS):** Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to **TWO** non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen) **AND** Trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

**Ulcerative Colitis (UC):** Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids. **AND** trial and failure, contraindication, or intolerance to one of the following conventional

## Criteria Details

	<p>therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone) <b>AND</b> trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. adalimumab, Simponi) <b>AND</b> patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g., azathioprine or cyclosporine).</p> <p><b>Polyarticular Juvenile Idiopathic Arthritis (PJIA):</b> Diagnosis of active polyarticular course juvenile idiopathic arthritis <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide <b>AND</b> trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Enbrel, adalimumab) <b>AND</b> patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).</p>
<p><b>Age Restrictions</b></p>	<p><b>Solution only</b> One of the following:</p> <ul style="list-style-type: none"> <li>• Pediatric member age 10 or under</li> <li>• Documentation inability of the member to use the preferred tablet formulation</li> </ul>
<p><b>Prescriber Restrictions</b></p>	<p><b>RA, AS, PJIA:</b> Prescribed by or in consultation with a rheumatologist <b>PsA:</b> Prescribed by or in consultation with a dermatologist or rheumatologist <b>UC:</b> Prescribed by or in consultation with a gastroenterologist</p>
<p><b>Coverage Duration</b></p>	<p><b>RA, PsA, AS, PJIA: Initial:</b> 6 months. <b>Renewal</b> 12 months <b>Ulcerative Colitis (UC): Initial:</b> 4 months. <b>Renewal</b> 12 months</p>
<p><b>Renewal Criteria</b></p>	<p><b>RA, PJIA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline <b>AND</b> will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).</p> <p><b>PsA:</b> Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline <b>AND</b> will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).</p>

## Criteria Details

**AS:** Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count. **AND** will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**UC:** Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state **AND** will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**Effective Date**

5/1/2024

**P&T Approval Date**

3/12/2024

**P&T Revision Date**

3/12/2024

# Treprostinil (ORENITRAM)

## Products Affected

- ORENITRAM

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmed diagnosis of pulmonary arterial hypertension with right heart catheterization <b>AND</b> WHO functional class II through IV <b>AND</b> evidence of either an unfavorable acute response to vasodilators or evidence of being refractory to or unable to tolerate calcium channel blockers (e.g., extended release nifedipine or extended-release diltiazem) <b>AND</b> documented failure or incomplete response to sildenafil or tadalafil/ambrisentan combination <b>AND</b> trial and failure of Remodulin or clinical justification for the need of an alternative route of administration.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Treprostinil (REMODULIN)

## Products Affected

- TREPROSTINIL

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmed diagnosis of pulmonary arterial hypertension with right heart catheterization <b>AND</b> WHO functional class II through IV <b>AND</b> evidence of either an unfavorable acute response to vasodilators or evidence of being refractory to or unable to tolerate calcium channel blockers (e.g., extended release nifedipine or extended-release diltiazem) <b>AND</b> documented failure or incomplete response to sildenafil or tadalafil/ambriksentan combination.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a cardiologist or pulmonologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Treprostinil (TYVASO)

## Products Affected

- TYVASO
- TYVASO REFILL
- TYVASO STARTER

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Confirmed diagnosis of pulmonary hypertension associated with interstitial lung disease (WHO Group 3).  *Any other indication would be required to try and fail formulary alternatives.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by pulmonologist or cardiologist.
<b>Coverage Duration</b>	<b>Initial:</b> 12 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Ubrogепant (UBRELVY)

## Products Affected

- UBRELVY

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	Diagnosis of migraine with or without aura <b>AND</b> will be used for the acute treatment of migraine <b>AND</b> patient has fewer than 15 headache days per month <b>AND</b> trial and failure or intolerance to 3 generic triptans and NSAID (ibuprofen, naproxen, diclofenac) combined treatment <b>OR</b> trial and failure or intolerance to NSAID treatment alone if triptans contraindicated <b>OR</b> contraindication to all triptans and NSAIDs <b>AND</b> medication will not be used in combination with another CGRP inhibitor <b>AND</b> if patient has 4 or more headache days per month, then patient must currently be treated with amitriptyline, venlafaxine, divalproex, topiramate, candesartan or a beta-blocker or have a contraindication or intolerance to all of these medications.
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with one of the following specialists: Neurologist, pain specialist, headache specialist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy and will not be used for preventative treatment of migraine.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	



# Umbralisib (UKONIQ)

## Products Affected

- Ukoniq

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Marginal zone lymphoma (MZL):</b> Diagnosis of relapsed or refractory marginal zone lymphoma (MZL) <b>AND</b> must have received a prior therapy that included an anti-CD20 antibody agent  <b>Follicular lymphoma (FL):</b> Must have received at least three prior therapies, including both an anti-CD20 antibody <b>AND</b> an alkylating agent  *Maximum daily dose of 4 tablets
<b>Age Restrictions</b>	18 years of age <b>AND</b> older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> up to 12 months.
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Upadacitinib (RINVOQ)

## Products Affected

- RINVOQ

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

**Rheumatoid Arthritis (RA):** Diagnosis of moderately to severely active rheumatoid arthritis **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**Psoriatic Arthritis (PsA):** Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**Ankylosing spondylitis (AS):** Diagnosis of active ankylosing spondylitis minimum duration of one month trial and failure, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen). **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**Atopic Dermatitis (AD):** Diagnosis of moderate to severe atopic dermatitis **AND** one of the following: Involvement of at least 10% body surface area (BSA), SCORing Atopic Dermatitis (SCORAD) index value of at least 25 **AND** a minimum duration of a 30-day trial and failure, contraindication, or intolerance to at least one of the following: Medium

## Criteria Details

	<p>or higher potency topical corticosteroid, Pimecrolimus cream, Tacrolimus ointment, or Eucrisa (crisaborole) ointment <b>AND</b> a minimum duration of 12-week trial and failure, contraindication, or intolerance of at least one systemic drug for the treatment of AD (e.g. Dupixent) <b>AND</b> not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).</p> <p><b>Crohn's disease (CD):</b> Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220 <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate <b>AND</b> trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. adalimumab, Simponi) <b>AND</b> not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).</p>
<b>Age Restrictions</b>	<b>AD:</b> Age 12 or older
<b>Prescriber Restrictions</b>	<p><b>RA, AS:</b> Prescribed by or in consultation with a rheumatologist</p> <p><b>PsA:</b> Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist.</p> <p><b>AD:</b> Prescribed by or in consultation with one of the following: Dermatologist or Allergist/Immunologist.</p> <p><b>UC:</b> Prescribed by or in consultation with a gastroenterologist</p>
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	<p><b>RA:</b> Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in total active joint count, improvement in symptoms (e.g., improvement in number of swollen/tender joints, pain, or stiffness). <b>AND</b> Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).</p> <p><b>PsA:</b> Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms(e.g., improvement in number of swollen/tender joints, pain, or stiffness) <b>AND</b> Rinvoq will not be used in combination with other JAK inhibitors, biologic</p>

## Criteria Details

DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**AS:** Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count **AND** Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**AD:** Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline **AND** Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**UC:** Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state **AND** Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

**Effective Date**

**P&T Approval Date**

**P&T Revision Date**

# Ustekinumab (STELARA)

## Products Affected

- STELARA

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<p><b>Plaque Psoriasis (PsO):</b> Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement <b>AND</b> a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar. For Stelara SC 90mg/1 mL: patient weight is greater than 100 kg (220 lbs).</p> <p><b>Psoriatic Arthritis (PsA):</b> Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement. For Stelara SC 90mg/1 mL: patient weight is greater than 100 kg (220 lbs) and has diagnosis of co-existent moderate to severe psoriasis.</p> <p><b>Crohn's Disease (CD):</b> Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220 <b>AND</b> Trial and failure, contraindication, or intolerance to <b>ONE</b> of the following conventional therapies: 6-mercaptopurine, Azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone), Methotrexate.</p> <p><b>Ulcerative Colitis (UC):</b> Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids <b>AND</b> trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine</p>
-------------------------------------	---

### Criteria Details

	sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone).
<b>Age Restrictions</b>	<b>PsO/PsA:</b> 6 and older
<b>Prescriber Restrictions</b>	<b>PsO:</b> Prescribed by or in consultation with a dermatologist <b>PsA:</b> Prescribed by or in consultation with a dermatologist or rheumatologist <b>CD/UC:</b> Prescribed by or in consultation with a gastroenterologist
<b>Coverage Duration</b>	<b>Initial:</b> 6 months. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	<p><b>PsO:</b> Documentation of positive clinical response to therapy as evidenced by <b>ONE</b> of the following: Reduction the body surface area (BSA) involvement from baseline <b>OR</b> improvement in symptoms (e.g., pruritus, inflammation) from baseline.</p> <p><b>PsA:</b> Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms (e.g., improvement in number of swollen/tender joints, pain, or stiffness).</p> <p><b>CD:</b> Documentation of positive clinical response to therapy as evidenced by improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline or reversal of high fecal output state.</p> <p><b>UC:</b> Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.</p>
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Vedolizumab (ENTYVIO)

## Products Affected

- ENTYVIO

## Prior Authorization Criteria

### Criteria Details

#### Required Medical Information

#### Crohn's Disease

- Documentation of moderately to severely active Crohn's disease
- One of the following:
  - Frequently diarrhea and abdominal pain
  - At least 10% weight loss
  - Complications such as obstruction, fever abdominal mass
  - Abnormal lab values (e.g., C-reactive protein [CRP])
- Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:
  - 6-mercaptopurine
  - azathioprine
  - corticosteroids (e.g., prednisone)
  - methotrexate
- One of the following:
  - Trial and failure, contraindication, or intolerance to TWO of the following:
    - Cimzia (certolizumab pegol)
    - Humira (adalimumab), Amjevita, Cyltezo, Hyrimoz, or Brand Adalimumab-adaz
    - Stelara (ustekinumab)
    - Skyrizi (risankizumab-rzaa)
  - For continuation of prior Entyvio therapy, defined as no more than a 45-day gap in therapy

#### Ulcerative Colitis

- Diagnosis of moderately to severely active ulcerative colitis
- One of the following:
  - Greater than 6 stools per day
  - Frequent blood in the stools
  - Frequent urgency
  - Presence of ulcers

## Criteria Details

	<ul style="list-style-type: none"> <li>○ Abnormal lab values (e.g., hemoglobin, ESR, CRP)</li> <li>○ Dependent on, or refractory to, corticosteroids</li> <li>● Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies :             <ul style="list-style-type: none"> <li>○ 6-mercaptopurine</li> <li>○ Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)</li> <li>○ Azathioprine</li> <li>○ Corticosteroids (e.g., prednisone)</li> </ul> </li> <li>● One of the following:             <ul style="list-style-type: none"> <li>○ Trial and failure, contraindication, or intolerance to TWO of the following, or attestation demonstrating a trial may be inappropriate*:                 <ul style="list-style-type: none"> <li>▪ Humira (adalimumab), Amjevita, Cyltezo, Hyrimoz, or Brand Adalimumab-adaz</li> <li>▪ Simponi (golimumab)</li> <li>▪ Stelara (ustekinumab)</li> <li>▪ Rinvoq (upadacitinib)</li> <li>▪ Xeljanz/XR (tofacitinib/ER)</li> </ul> </li> <li>○ For continuation of prior Entyvio therapy, defined as no more than a 45-day gap in therapy</li> </ul> </li> </ul>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Prescribed by or in collaboration with a Gastroenterologist
<b>Coverage Duration</b>	<b>Initial:</b> 14 weeks. <b>Renewal:</b> 12 months.
<b>Renewal Criteria</b>	<p><b>Crohn's Disease:</b> Documentation of positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> <li>• Reversal of high fecal output state</li> </ul> <p><b>Ulcerative Colitis:</b> Documentation of positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> <li>• Reversal of high fecal output state</li> </ul>
<b>Effective Date</b>	5/1/2024



## Criteria Details

<b>P&amp;T Approval Date</b>	3/12/2024
<b>P&amp;T Revision Date</b>	

# Vonoprazan (VOQUEZNA)

## Products Affected

- VOQUEZNA 10MG
- VOQUEZNA 20MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Erosive esophagitis-</b> <ul style="list-style-type: none"><li>• Imaging confirmed LA Classification Grade C/D erosive esophagitis AND</li><li>• Documented contraindication, intolerance, or inadequate response to 2 or more PPIs (i.e., lansoprazole, omeprazole, esomeprazole, etc.) at maximum tolerated twice-daily dosing for at least 8 weeks each.</li></ul> <b>H.pylori eradication –</b> <ul style="list-style-type: none"><li>• Confirmed H. pylori positive infection AND</li><li>• Documented contraindication, intolerance, or inadequate response to standard first-line therapies for H.pylori infection (e.g. PPI (standard or double dose), clarithromycin, amoxicillin (or metronidazole)) AND</li><li>• Documented contraindication, intolerance, or inadequate response to a quadruple bismuth regimen (e.g. standard twice daily dose PPI, bismuth subsalicylate, tetracycline, metronidazole) AND</li><li>• Co-prescribed in combination with antibiotics.</li></ul>
<b>Age Restrictions</b>	Must be at least 18 years of age
<b>Prescriber Restrictions</b>	Prescribed by or in collaboration with a Gastroenterologist or Infectious Disease specialist
<b>Coverage Duration</b>	<b>Initial healing of erosive esophagitis:</b> 2 months <b>Maintenance of healing of erosive esophagitis:</b> 6 months <b>H. Pylori eradication:</b> 14 days
<b>Renewal Criteria</b>	Renewals past the above timelines are not allowed
<b>Effective Date</b>	03/01/2024
<b>P&amp;T Approval Date</b>	01/09/2024

## Criteria Details

P&T Revision Date	
-------------------	--

# Zanubrutinib (BRUKINSA)

## Products Affected

- BRUKINSA CAP 80MG

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Follow general oncology criteria with the following parameters:</b> One prior treatment for mantle cell lymphoma( MCL) <b>OR</b> one prior treatment (anti-CD20 based) for refractory marginal zone lymphoma (MZL) <b>OR</b> diagnosis of Waldenstrom's macroglobulinemia (WM).
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by oncologist
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> 12 months
<b>Renewal Criteria</b>	Documentation of positive clinical response to therapy.
<b>Effective Date</b>	
<b>P&amp;T Approval Date</b>	
<b>P&amp;T Revision Date</b>	

# Zuranolone (ZURZUVAE)

## Products Affected

- ZURZUVAE

## Prior Authorization Criteria

### Criteria Details

<b>Required Medical Information</b>	<b>Postpartum Depression</b> <ul style="list-style-type: none"><li>• Physician attestation of moderate to severe postpartum depression (PPD) diagnosis and submission of validated screening tool result(s) (e.g. EPDS, PHQ-9) that will be used to monitor a patient's response to Zurzuvae therapy</li><li>• Physician attestation that patient has not had a major depressive episode prior to third trimester of pregnancy and no later than the first 4 weeks following delivery</li><li>• Patient has tried/failed generic SSRI or SNIR for PPD</li></ul>
<b>Age Restrictions</b>	18 years of age and older
<b>Prescriber Restrictions</b>	Prescribed by psychiatrist or OB/GYN
<b>Coverage Duration</b>	<b>Initial:</b> 3 months. <b>Renewal:</b> N/A
<b>Renewal Criteria</b>	
<b>Effective Date</b>	7/1/2024
<b>P&amp;T Approval Date</b>	5/14/2024
<b>P&amp;T Revision Date</b>	

