

Prior Authorization Criteria

Samaritan Choice

PLEASE READ: This document contains information about the criteria for coverage for this plan.

Abatacept (ORENCIA)

Products Affected

ORENCIA

ORENCIA CLICKJECT

Prior Authorization Criteria

Criteria Details

Adult Rheumatoid Arthritis (RA) Diagnosis of moderately to severely active rheumatoid arthritis AND trial and failure, contraindication, or intolerance to ONE nonbiologic disease-modifying antirheumatic drug (DMARD) (e.g., methotrexate [Rheumatrex/Trexall], Arava [leflunomide], Azulfidine [sulfasalazine]) AND trial and failure, contraindication, or intolerance to TWO of the following, or attestation demonstrating a trial may be inappropriate Cimzia (certolizumab pegol) adalimumab Rinvoq (upadacitinib) Simponi (golimumab) Xeljanz/XR (tofacitinib/ER).

Required Medical Information

Polyarticular Juvenile Idiopathic Arthritis (PJIA) Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, AND trial and failure, contraindication, or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), adalimumab, Xeljanz (tofacitinib).

Psoriatic Arthritis (PsA) Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement AND trial and failure, contraindication, or intolerance to TWO of the following: Cimzia (certolizumab pegol), adalimumab, Simponi (golimumab), ustekinumab, Tremfya (guselkumab), Skyrizi (risankizumab-rzaa), Rinvoq (upadacitinib), Xeljanz/XR (tofacitinib/ER).

Prophylaxis for Acute Graft versus Host Disease (aGVHD) Used for prophylaxis of acute graft versus host disease (aGVHD) AND patient will receive hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor AND recommended antiviral prophylactic treatment for Epstein-Barr Virus (EBV) reactivation (e.g.,

	Criteria Details
	acyclovir) will be administered prior to Orencia AND continued for six months after HSCT AND Used in combination with both of the following: calcineurin inhibitor (e.g., cyclosporine, tacrolimus) methotrexate.
Age Restrictions	aGVHD: 2 years of age or older
Prescriber Restrictions	RA, PJIA: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with a dermatologist or rheumatologist
Coverage Duration	RA, PJIA, PsA: Initial: 6 months. Renewal: 12 months aGVHD: Initial: 2 months. Renewal: N/A
Renewal Criteria	RA, PJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline. PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Adalimumab Biosimilars

Products Affected

- Adalimumab-adaz- 40 MG/0.4ML (autoinjector and prefilled syringe)
- Adalimuimab-fkjp 40MG/0.8ML, 20 MG/0.4ML, (auto-injector and prefilled syringe)
- Hadlima 40MG/0.4ML, 40/0.8ML (autoinjector and prefilled syringe)
- Yusimry 40MG/0.8ML
- Simlandi 40MG/0.4ML

Prior Authorization Criteria

Criteria Details

Rheumatoid arthritis (RA): Diagnosis of moderately to severely active RA **AND** trial and failure, contraindication, or intolerance to one non-biologic disease-modifying antirheumatic drug (DMARD) [e.g., methotrexate (Rheumatrex/Trexall), Arava (leflunomide), Azulfidine (sulfasalazine)].

Polyarticular Juvenile idiopathic arthritis (PJIA): Diagnosis of moderate to severely active polyarticular JIA **AND** trial and failure, contraindication, or intolerance to one of the following non-biologic disease-modifying antirheumatic drugs (DMARDs): Arava (leflunomide), methotrexate (Rheumatrex/Trexall).

Required Medical Information

Psoriatic arthritis (PsA): Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.

Ankylosing spondylitis (AS): Diagnosis of active ankylosing spondylitis **AND** trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen)

Crohn's disease (CD): Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220. **AND** trial and failure,

	Criteria Details
	contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate
	Ulcerative Colitis (UC): Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids AND trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone).
	Plaque Psoriasis (PP): Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement. AND a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.
	Hidradenitis Suppurativa (HS): Diagnosis of moderate to severe hidradenitis suppurativa (i.e., Hurley Stage II or III)
	Uveitis (UV): Diagnosis of non-infectious uveitis AND uveitis is classified as one of the following: intermediate, posterior or panuveitis.
Age Restrictions	
Prescriber Restrictions	RA, PJIA, AS: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist. CD, UC: Prescribed by or in consultation with a gastroenterologist PsO, HS: Prescribed by or in consultation with a dermatologist UV: Prescribed by or in consultation with one of the following: ophthalmologist or rheumatologist
Coverage Duration	RA, PJIA, PsA, PsO, AS, CD, HS, UV Initial: 6 months; Renewal: 12 months UC: Initial: 12 weeks; Renewal: 12 months
Renewal Criteria	RA, PJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen
	, , , , , , , , , , , , , , , , , , , ,

Criteria Details

and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.

PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.

AS: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.

CD: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.

UC: One of the following: For patients who initiated adalimumab therapy within the past 12 weeks: Documentation of clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on adalimumab therapy for longer than 12 weeks: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.

PsO: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.

HS, UV: Documentation of positive clinical response to therapy.

Effective Date	
P&T Approval Date	

	Criteria Details
P&T Revision Date	

Alpelisib (VIJOICE)

Products Affected

Vijoice TAB

	Criteria Details
Required Medical Information	Confirmed diagnosis of PROS AND has at least one severe clinical manifestation of PROS AND has a PIK3CA mutation that is confirmed by genetic testing
Age Restrictions	At least 2 years of age
Prescriber Restrictions	Prescribed by or in consultation with a provider who specializes in treatment of genetic disorders
Coverage Duration	Initial: 24 weeks. Renewal: 6 months.
Renewal Criteria	Documentation of a reduction in volume from baseline in at least one lesion AND an improvement in at least one symptom of PROS from baseline
Effective Date	
P&T Approval Date	
P&T Revision Date	

Apremilast (OTEZLA)

Products Affected

OTEZLA

	Criteria Details
Required Medical Information	Plaque Psoriasis (PsO): Diagnosis of plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement AND a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar. Psoriatic Arthritis (PsA): Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement. Oral Ulcers Associated with Behçet's Disease: Diagnosis of Behçet's Disease AND Patient has active oral ulcers.
Age Restrictions	
Prescriber Restrictions	Plaque psoriasis: Prescribed by or in consultation with a dermatologist. Psoriatic arthritis: Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	PsO: Documentation of positive clinical response to therapy as evidenced by ONE of the following: Reduction the body surface area (BSA) involvement from baseline OR improvement in symptoms (e.g., pruritus, inflammation) from baseline. PsA: Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms Behçet's Disease: Documentation of positive clinical response to therapy (e.g., reduction in pain from oral ulcers or reduction in number of oral ulcers).

	Criteria Details
Effective Date	
P&T Approval Date	
P&T Revision Date	

Bedaquiline (SIRTURO)

Products Affected

Sirturo tablets

	Criteria Details
Required Medical Information	 Pulmonary tuberculosis Evidence of active pulmonary tuberculosis caused by mycobacterium tuberculosis that is resistant to at least rifampin and isoniazid. The member weighs at least 15kg. Sirturo is prescribed as part of a guideline recommended multi-drug treatment regimen.
Exclusion Criteria	Medication is being received through a county clinic with a state funded TB program.
Age Restrictions	5 years of age or older
Prescriber Restrictions	Prescribed by Infectious Disease
Coverage Duration	Pulmonary tuberculosis: 24 weeks
Renewal Criteria	N/A
Effective Date	05/01/2025
P&T Approval Date	03/11/2025
P&T Revision Date	03/11/2025

Bempedoic acid (NEXLETOL)

Products Affected

NEXLETOL TABLETS

NEXLIZET TABLETS

	Criteria Details
Required Medical Information	 Established clinical ASCVD: Documentation of very high risk ASCVD as evidenced by either: History of multiple major ASCVD events OR One major ASCVD event AND multiple high-risk conditions. Documentation of a current LDL greater than or equal to 55 mg/dl. Documentation that: Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins AND Is receiving ezetimibe or has a documented intolerance to ezetimibe. Documentation of failure of PCSK9 inhibitor (Repatha or Praluent).
	 Primary or familial hyperlipidemia: Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL. Documentation of current LDL greater than 100 mg/dL. Documentation that: Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins AND Is receiving ezetimibe or has a documented intolerance to ezetimibe. Documentation of failure of PCSK9 inhibitor (Repatha or Praluent).
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.

	Criteria Details
Coverage Duration	Initial: 6 months. Renewal: 12 months
Renewal Criteria	Documented positive clinical response to therapy (significant decrease in lipid levels).
Effective Date	11/1/2024
P&T Approval Date	9/10/2024
P&T Revision Date	9/10/2024

Belumosudil (REZUROCK)

Products Affected

REZUROCK

	Criteria Details
Required Medical Information	Diagnosed with chronic graft-versus-host disease (cGVHD) AND who have tried and failed of at least two prior lines of systemic therapy for cGVHD AND not currently taking Imbruvica (ibrutinib)
Age Restrictions	12 years of age or older
Prescriber Restrictions	Prescribed by oncologist or transplant specialist
Coverage Duration	Initial: 3 months. Renewal: 6 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Bimekizumab-bkzx (BIMZELX)

Products Affected

- Bimzelx 160mg/mL PFS
- Bimzelx 160mg/mL auto-injector
- Bimzelx 320mg/2mL PFS
- Bimzelx 320mg/2mL auto-injector

Prior Authorization Criteria

Criteria Details

Plaque Psoriasis

- Diagnosis of severe plaque psoriasis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool)
- One or more of the following:
 - At least 10% of body surface area involved
 - Hand, foot, face, or mucous membrane involvement
 - The patient is on a current biologic product and experiencing intolerable side effects
- The patient tried and failed or have contraindications to ALL of the following
 - High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.)
 - At least one other topical agent: calcipotriene, tazarotene, anthralin, tar, etc.
 - PUVA or UVB Phototherapy
 - Methotrexate
 - At least 1 other second line systemic agent such as cyclosporine or acitretin
- Trial and failure of both infliximab and adalimumab

Psoriatic Arthritis

- Documentation of psoriatic arthritis based on at least 3 out of 5 of the following:
 - Psoriasis (1 point for personal or family history, 2 points for current)
 - Psoriatic nail dystrophy
 - Negative test result for RF

Required Medical Information

Criteria Details

- Dactylitis (current of history)
- Radiological evidence of juxta-articular new bone formation
- The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following:
 - NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND
 - Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine
- Trial and failure of both infliximab and adalimumab
- Trial and failure of Cosentyx, ustekinumab, and Taltz

Ankylosing Spondylitis/Axial Spondyloarthritis

- Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by:
 - o Back pain and stiffness for more than 3 months AND
 - Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND
 - BASDAI score of >=4
- The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR Documented failure of conventional therapy with both of the following:
 - At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND
 - Physical therapy/exercise program
- Trial and failure of both infliximab and adalimumab
- Trial and failure of Cosentyx, and Taltz

Hidradenitis Suppurativa

- Documentation of one of the following:
 - Moderate to severe hidradenitis suppurative (Hurley Stage II or Hurley Stage III)
 - Patient is on a current biologic product and experiencing intolerable side effects.
- The patient is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of the following:
 - o 90-day trial of conventional therapy (e.g. oral antibiotics)
- Trial and failure of both infliximab and adalimumab
- Trial and failure of both Cosentyx and ustekinumab

	Criteria Details
Age Restrictions	Refer to FDA label
Prescriber Restrictions	 Plaque Psoriasis, Hidradenitis Suppurativa: Dermatologist. Psoriatic Arthritis: Dermatologist or Rheumatologist. Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	 Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. Hidradenitis Suppurativa: Evidence of a reduction of 25% or more of the total abscess and inflammatory nodule count and no increase in abscesses and draining fistulas.
Effective Date	9/1/2025
P&T Approval Date	5/13/2024
P&T Revision Date	7/8/2025, 5/13/2024

Brexipiprazole (REXULTI)

Products Affected

REXULTI

	Criteria Details
Required Medical Information	Major Depressive Disorder (MDD): A diagnosis of MDD AND prior treatment failure (at least 3 weeks) of or contraindication to 3 prior antidepressants AND one antipsychotic FDA approved as adjunct treatment for MDD AND to be used concurrently with an antidepressant.
	Schizophrenia: A diagnosis of schizophrenia AND prior treatment failure with a minimum of 2 antipsychotics AND Vraylar.
Age Restrictions	Schizophrenia: Aged 13 or older Major Depressive Disorder (MDD): Aged 18 or older
Prescriber Restrictions	
Coverage Duration	Initial: 3 months. Renewal: 12 months.
Renewal Criteria	Documentation of treatment success AND continued need for Rexulti.
Effective Date	
P&T Approval Date	
P&T Revision Date	

BUTRANS, BUPRENORPHINE PATCH, BELBUCA

Products Affected

BELBUCA

BUPRENORPHINE PATCH

	Criteria Details
	Cancer or End-of-Life Care: Patient is being treated for cancer related pain or pain associated with end-of-life: Documented trial and failure of, scheduled short-acting opioid therapy AND documented trial and failure or contraindication to long-acting morphine sulfate therapy. Documented trial/failure of, or reason why fentanyl is not appropriate.
Required Medical Information	Other Chronic Pain: Documented above the line diagnosis, FDA indicated, or guideline supported condition. Documented severe chronic pain (greater than 3mo) that is severe enough to require around the clock analgesic therapy AND documented trial and failure or contraindication to short-acting opioid therapy AND documented trial and failure of, or contraindication to long-acting morphine sulfate therapy AND documented trial and failure of, or reason why fentanyl is not appropriate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial cancer/end of life: 12 months. Initial non-cancer/end of life: 6 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Cenegermin (Oxervate)

Products Affected

Oxervate Ophthalmic

Criteria Details	
Required Medical Information	 Confirmed diagnosis of stage 2 or 3 neurotrophic keratitis Trial and failure of at least one ocular lubricant used for at least 2 weeks
Prescriber Restrictions	Prescribed by or in consultation with an opthalmologist
Coverage Duration	Initial: 8 weeks. Renewal: N/A, no renewal allowed
Effective Date	11/01/2025
P&T Approval Date	09/09/2025
P&T Revision Date	09/09/2025

Certolizumab Pegol (CIMZIA)

Products Affected

Cimzia

Prior Authorization Criteria

Criteria Details

Crohn's Disease (CD): Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. Creactive protein), CD Activity Index greater than 220. AND trial and failure, contraindication, or intolerance to ONE of the following conventional therapies: 6-mercaptopurine, Azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone), Methotrexate.

Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active RA **AND** trial and failure, contraindication or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine.

Required Medical Information

Psoriatic Arthritis (PsA): Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.

Ankylosing Spondylitis (AS): Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses.

Plaque Psoriasis (PsO): Diagnosis of moderate to severe plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement **AND** a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.

Criteria Details	
	Non-radiographic Axial Spondyloarthritis (nr-axSpA): Diagnosis of active non-radiographic axial spondyloarthritis AND patient has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) AND minimum duration of one month trial and failure, contraindication, or intolerance to two different NSAIDs (e.g., ibuprofen, naproxen) at maximally tolerated doses
Age Restrictions	
Prescriber Restrictions	CD: Prescribed by or in consultation with a gastroenterologist RA, AS, nr-axSpA: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist PsO: Prescribed by or in consultation with a dermatologist
Coverage Duration	CD: Initial: 16 weeks; Renewal: 12months RA, PsA, AS, PsO, nr-axSpA: Initial: 6 months; Renewal: 12 months
Renewal Criteria	CD: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.
	RA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.
	PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.
	AS, nr-axSpA: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial

Criteria Details		
	status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.	
	Ps0: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Clobazam (ONFI)

Products Affected

- Clobazam 10mg Tablets
- Clobazam 20mg Tablets

• Clobazam 2.5mg/mL suspension

	Criteria Details
Required Medical	Lennox-Gastaut Syndrome • Confirmed diagnosis. Refractory Seizures
	Documentation showing appropriate trial of 2 or more tolerated anticonvulsant therapies.
Age Restrictions	Solution only One of the following: Pediatric member age 10 or under Documentation inability of the member to use the preferred tablet formulation
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 12 months. Renewal: Lifetime.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	5/1/2025
P&T Approval Date	3/12/2024
P&T Revision Date	3/11/2025, 3/12/2024

Compounds (standard criteria for all compounded medications)

Products Affected

• All compounded medications

	Criteria Details
Required Medical Information	 Each active ingredient in the compounded drug is FDA-approved or national compendia* supported for the condition being treated. The requested amounts are supported by national compendia* or two peer-reviewed literature for the condition being treated in the requested route of delivery. If any prescription ingredients require prior authorization and/or step therapy, all drug-specific criteria must be also met. The patient has tried and failed therapy or had an intolerance to two FDA-approved commercially-available prescription therapeutic alternatives, one of which is the same route of administration as the requested compound, unless one of the following criteria are met: Patient has a contraindication to commercially available products Only one or no other therapeutic alternatives are commercially available Prepared strength(s) is/are not commercially available or currently in short supply Prepared in a different dosage form for a patient who is unable to take the commercially available products based on the manufacturer's instructions or the product's approved labeling does NOT meet this criteria). Patient has an allergy or sensitivity to inactive ingredients (e.g. dyes, preservatives, sugars, etc.) that are found in commercially available products.
Age Restrictions	

Prescriber Restrictions	
Restrictions	
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Continuous Glucose Monitor (CGM)

Products Affected

DEXCOM G6 & G7 SYSTEMS

• FREESTYLE LIBRE SYSTEMS

	Criteria Details	
Required Medical Information	Patient has documented diagnosis of type 1 or type 2 diabetes mellitus. Patient must have ALL of the following: Intensive insulin regimen (3 or more insulin injections per day, or use of continuous subcutaneous insulin infusion pump). Patient consistently monitors blood glucose 3 or more times per day. Patient is motivated and knowledgeable about use of continuous glucose monitoring, is adherent to diabetic treatment plan, and participates in ongoing education and support. Patient must have 1 OR more of the following: Dawn phenomenon, known or suspected, Hypoglycemic unawareness (i.e., patient does not have symptoms with hypoglycemia). Nocturnal hypoglycemia, known or suspected. Postprandial hyperglycemia, known or suspected. Significant change to diabetes treatment regimen (e.g., initiation of insulin, change from multiple-dose insulin to insulin pump therapy). Unexplained hyperglycemia.	
Age Restrictions		
Prescriber Restrictions		
Coverage Duration	Initial: 12 months. Renewal: 12 months.	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date	01/01/2024	
P&T Approval Date	11/14/2023	

Criteria Details	
P&T Revision Date	

Colony-Stimulating Factors

Products Affected

- NIVESTYM
- ZARXIO

- NUELASTA/NEULASTA ONPRO
- UDENYCA/UDENYCA ONPRO

Prior Authorization Criteria

Criteria Details

Bone Marrow/Stem Cell Transplant:

- One of the following:
 - Patient has non-myeloid malignancies undergoing myeloablative chemotherapy followed by autologous or allogeneic bone marrow transplant (BMT) **OR**
 - Used for mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis OR
 - Patient has had a peripheral stem cell transplant (PSCT) and has received myeloablative chemotherapy.

Acute Myeloid Leukemia (AML) Induction or Consolidation Therapy:

- Diagnosis of acute myeloid leukemia (AML).
- Patient has completed induction or consolidation chemotherapy.

Required Medical Information

Febrile Neutropenia Prophylaxis:

- Patient will be receiving prophylaxis for febrile neutropenia (FN) due to one of the following:
 - Patient is receiving National Cancer Institute's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer.
 - Patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown.
 - Patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN.
 - Patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN AND has one or more risk factors associated with chemotherapy induced infection, FN, or neutropenia.
 - Patient is receiving myelosuppressive anticancer drugs associated with neutropenia AND has a history of FN or dose-

	Critorio Dotoilo
	Criteria Details limiting event during a previous course of chemotherapy (secondary prophylaxis).
	 Treatment of High-Risk Febrile Neutropenia: Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. Diagnosis of febrile neutropenia (FN). Patient is at high risk for infection-associated complications.
	Severe Chronic Neutropenia (SCN): For patients with severe chronic neutropenia (SCN) (i.e., congenital, cyclic, and idiopathic neutropenias with chronic absolute neutrophil count [ANC] less than or equal to 500 cells/mm^3)
	Acute Radiation Syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation cells/mm^3).
	 Human Immunodeficiency Virus (HIV) Related Neutropenia: Patient is infected with HIV virus. ANC less than or equal to 1,000 (cells/mm3).
Age Restrictions	
Prescriber Restrictions	HIV Related Neutropenia – Hematologist, Oncologist, or Infectious Disease Specialist All Other Diagnosis - Hematologist or Oncologist
Coverage Duration	Bone Marrow/Stem Cell Transplant: 3 months or duration of therapy. AML Induction/Consolidation Therapy: 3 months or duration of therapy. Febrile Neutropenia (FN) Prophylaxis: 3 months or duration of therapy. Treatment of High-Risk FN: 3 months or duration of therapy. Severe Chronic Neutropenia (SCN): 12 months. Acute Radiation Syndrome (ARS): 1 month
Renewal Criteria	
Effective Date	09/01/2024
P&T Approval Date	07/09/2024

Cyclosporine ophthalmic emulsion (RESTASIS)

Products Affected

RESTASIS

RESTASIS MULTIDOSE

	Criteria Details
Required Medical Information	The patient has a diagnosis of lack of tear production due to ocular inflammation associated with keratoconjunctivitis sicca AND ONE of the following: The patient is not currently using a topical ophthalmic anti-inflammatory drug or punctal plug OR the patients current use of topical ophthalmic anti-inflammatory drug or punctal plug will be discontinued before starting the requested agent AND the patient has previously tried or is currently using aqueous enhancements (e.g. artificial tears, gels, ointments) OR the patient has a documented intolerance, contraindication, or hypersensitivity to aqueous enhancements AND the patient is not currently using Xiidra OR the patients current use of Xiidra will be discontinued before starting Restasis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Deutetrabenazine (AUSTEDO)

Products Affected

- Austedo 6mg TAB
- Austedo 9mg TAB

• Austedo 12mg TAB

	Criteria Details
Required Medical Information	Chorea associated with Huntington's Disease: Documentation of the degree of chorea present and the impact on functional ability and/or quality of life as a baseline AND documentation of mental status, specifically depression and suicidality. Tardive Dyskinesia: Clinical documentation of tardive dyskinesia including 1) At least one month of past or current exposure to a dopamine receptor blocker, 2) Dyskinetic or dystonic involuntary movements, 3) Exclusion of other causes of abnormal movements AND clear documentation that tardive dyskinesia causes functional impairment AND documentation of the degree of tardive dyskinesia with the AIMS scale as a baseline AND one of the following: tried and failed an 8-week trial of at least 2 other agents within the same therapeutic category at a clinically effective and maximally tolerated dose for the patient's primary neuropsychiatric diagnosis OR evidence the medications precipitating tardive dyskinesia are medically necessary AND trial and failure or contraindication to clonazepam and amantadine.
Age Restrictions	Age 18 and older
Prescriber Restrictions	Huntington's Disease: neurologist Tardive Dyskinesia: neurologist or psychiatrist
Coverage Duration	Initial: 3 months. Renewal: 12 months.
Renewal Criteria	Huntington's Chorea: clinical response such as improvement in chorea, ability to perform ADLs, reduction in falls, or increase in quality of life. AND documentation of continued monitoring of mental status specifically for depression and suicidality.

Criteria Details		
	Tardive Dyskinesia: Follow-up AIMS assessment showing improvement from Baseline AND documented improvement in functioning such as ability to perform ADLs, reduction in falls and increase in quality of life.	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Dimethyl Fumarate (TECFIDERA)

Products Affected

DIMETHYL FUMARATE

• DIMETHYL FUMARATE STARTER PACK

	Criteria Details
Required Medical Information	Multiple sclerosis: Patient is diagnosed with relapsing forms of multiple sclerosis.
Age Restrictions	
Prescriber Restrictions	Prescribed by a neurologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Direct-Acting Antivirals (use in Hepatitis C)

Products Affected

LEDIPASVIR-SOFOSBUVIR

SOFOSBUVIR-VELPATASVIR

MAVYRET

Prior Authorization Criteria

Criteria Details

Treatment of Hepatitis C:

- Expected survival from non-HCV-associated morbidities more than 1 year.
- Must have all pretreatment testing completed: including genotype, HBV, HIV, and cirrhosis status.
- Care must be provided by or in consultation with a specialist (hepatologist, gastroenterologist, or infectious disease specialist).
- Attestation that the patient and provider will comply with case management to promote the best possible outcome for the patient and adhere to monitoring requirements required by the Oregon Health Authority, including measuring and reporting of a posttreatment viral load OR attestation from the patient and provider that they have opted out of OHA case management. Case management includes assessment of treatment barriers and offer of patient support to mitigate potential barriers to regimen adherence as well as facilitation of SVR12 evaluation to assess treatment success.
- Documentation if the patient has a GT 1a infection or GT 3 infection and the patient had a baseline NS5a resistance test that documents a resistant variant to Elbasvir/grazoprevir or Daclatasvir + sofosbuvir. Note: Baseline NS5A resistance testing is required.
- Documentation of the prescribed regimen includes a NS3/4a protease inhibitor (glecaprevir, simeprevir, paritaprevir, voxilaprevir).
- Documentation if the patient has moderate-severe hepatic impairment (Child-Pugh B or Child-Pugh C).

Documentation if the prescribed regimen for the retreatment after failure of a DAA due to noncompliance or loss of follow-up and the prescribed

Required Medical Information

Criteria Details		
Age Restrictions		
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist, gastroenterologist, or infectious disease specialist	
Coverage Duration	Initial: 2-4 months.	
Renewal Criteria		
Effective Date		
P&T Approval Date		
P&T Revision Date		

Disposable Insulin Pump (OMNIPOD)

Products Affected

OMNIPOD 5

OMNIPOD DASH

Criteria Details		
Required Medical Information	Insulin dependent diabetes mellitus – pediatric (under age 18) • Documentation of Type 1 Diabetes Mellitus or Diabetes with Creactive protein levels indicating insulin dependence. • On intensive insulin therapy (>3 daily insulin injections) requiring frequent self-adjustments for at least 6 months prior to initiation of the insulin pump. • Documentation self-testing of blood glucose at least 4 times per day during the previous 2 months • Evidence of completion of a comprehensive diabetes education program in the last 12 months (member or caregiver/parent). Insulin dependent diabetes mellitus – adult • All of the above pediatric requirements AND • Documentation of 1 of the following: • HbA1c >7% • History of recurring hypoglycemia • Wide fluctuations in blood glucose before mealtime • Dawn phenomenon with fasting blood sugars frequently exceeding 200mg/dL • History of severe glycemic excursions • Inability to use a traditional (non-disposable) insulin pump.	
Age Restrictions		
Prescriber Restrictions		
Coverage Duration	Initial: 6 months Renewal: 12 months	
Renewal Criteria	Documentation of positive clinical response to therapy and in-person visit with provider within the last 6 months.	

Criteria Details	
Effective Date	03/01/2024
P&T Approval Date	01/09/024
P&T Revision Date	

Dupilumab (DUPIXENT)

Products Affected

DUPIXENT

Prior Authorization Criteria

Criteria Details

Moderate to Severe Asthma:

- Documentation of inadequate control of asthma symptoms with one of the following:
 - o inhaled corticosteroids and long acting beta2 agonist **OR**
 - o inhaled corticosteroids and long-acting muscarinic antagonist.

Atopic Dermatitis:

- Diagnosed with severe atopic dermatitis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) ≥ 11 or Children's Dermatology Life Quality Index (CDLQI) ≥ 13 (or severe score on another validated tool)
- One or more of the following:
 - At least 10% of body surface area involvement
 - Hand, foot, or mucous membrane involvement
- Documented contraindication or failed trial to ALL of the following:
 - Moderate-high potency corticosteroid (e.g., clobetasol, fluocinonide, fluticasone)
 - Topical calcineurin inhibitor (e.g. tacrolimus)
 - Oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil) **OR** the member is oral corticosteroid dependent.

Eosinophilic Esophagitis:

- Confirmed diagnosis of EoE
- Weight ≥ 15 kg
- Two or more episodes of dysphagia per week
- Inadequate response to an 8-week trial, intolerance, or contraindication to high-dose PPI therapy
- Inadequate response to and 8-to-12-week trial, intolerance, or contraindication to swallowed inhaled respiratory corticosteroid therapy.

Required Medical Information

Criteria Details

Chronic Rhinosinusitis with Nasal Polyps (CRSwNP):

- Diagnosis of CRSwNP, including objective evidence of the presence of bilateral nasal polyps
- Will not be used in combination with other biologics for eosinophilic indications.
- Trial and failure to adequately reduce symptoms with:
 - At least 2 months of saline nasal irrigations and inhaled nasal corticosteroids used at doses appropriate for nasal polyp treatment.
 - Systemic corticosteroid treatment for nasal polyps at least once within the last 2 years or prior nasal polyp removal surgery.
- Inhaled nasal corticosteroids will be used concomitantly with dupilumab (unless not tolerated or contraindicated).

Prurigo Nodularis (PN):

- Diagnosis of PN verified by a dermatologist and the patient has had the diagnosis for at least 3 months.
- Severe or very severe itch (WI-NRS score ≥7) reported within the past week.
- At least 20 PN lesions in total on both legs and/or both arms and/or trunk.
- Trial and failure (inadequate efficacy after 4 week trial, intolerable side effects) or contraindication to recommended first line agents for the treatment of PN including:
 - High potency topical steroids
 - Phototherapy
 - At least one systemic agent (immunosuppressant, gabapentinoid, or antidepressant).

Chronic Obstructive Pulmonary Disease (COPD):

- Diagnosis of COPD confirmed by post-bronchodilator FEV1/FVC < 0.7 on spirometry.
- Blood eosinophil count (BEC) ≥300 cells/µL within the past 3 months.
- Chronic bronchitis, defined as a chronic productive cough for ≥ 3 months in the past year, in the absence of other known causes of chronic cough.
- ≥ 2 moderate COPD exacerbation (defined as requiring treatment with either systemic corticosteroids and/or antibiotics) or ≥ 1 severe COPD exacerbation (defined as requiring hospitalization or observation for over 24 hours in emergency department of urgent

	Criteria Details care) within the past year despite the adherent use of inhaled LABA + LAMA+ ICS triple therapy [or LABA + LAMA dual therapy if ICS are contraindicated].	
Age Restrictions	Moderate to Severe Asthma: 6 years and older Atopic Dermatitis: 6 months and older Eosinophilic Esophagitis: 1 year and older CRSwNP: 12 years and older Prurigo Nodularis: 18 years and older COPD: 18 years and older	
Prescriber Restrictions	Atopic dermatitis: Prescribed by a Dermatologist Eosinophilic Esophagitis: Prescribed by Gastroenterologist or Immunologist CRSwNP: Prescribed by ENT or Immunologist Prurigo Nodularis: Prescribed by Dermatologist Asthma/COPD: Pulmonologist	
Coverage Duration	Initial: 6 months. Renewal: 12 months.	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date	03/01/2025	
P&T Approval Date	11/08/2022	
P&T Revision Date	01/14/2025	

Elagolix (ORLISSA)

Products Affected

• ORILISSA TAB 150MG

ORILISSA TAB 200MG

	Criteria Details
Required Medical Information	Diagnosis of moderate to severe pain associated with endometriosis AND trial and failure, contraindication, or intolerance to a 3-month trial of prescription strength NSAIDs AND trial and failure, contraindication, or intolerance to two 3-month trials of hormonal therapies (eg combined oral contraceptives, progestins, or levonorgestrel IUD, etc.).
	Additional info required for 200 mg tablet twice daily: documentation of coexisting dyspareunia
Age Restrictions	At least 18 years old but not yet through menopause
Prescriber Restrictions	Prescribed by obstetrician or gynecologist
Coverage Duration	200MG dose: Initial: 6 months; Renewal: No Renewals allowed 150MG dose: Initial: 6 months; Renewal: 18months
Renewal Criteria	150MG ONLY Documentation of positive clinical response to therapy AND total therapy durations is less than 24 months.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Elefibranor (IQIRVO)

Products Affected

• IQIRVO TABLETS

Criteria Details		
Required Medical Information	 Primary biliary cholangitis: Diagnosis of primary biliary cholangitis (PBC) confirmed by two of the following: Biochemical evidence of cholestasis based on ALP elevation Presence of AMA or other PBC-specific autoantibodies Histology confirmation after biopsy Trial and failure of 12 months of ursodiol. No current decompensated cirrhosis. 	
Age Restrictions		
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist or hepatologist.	
Coverage Duration	Initial: 6 months; Renewal: 12 months	
Renewal Criteria	Documented positive clinical response to therapy.	
Effective Date	11/1/2024	
P&T Approval Date	9/10/2024	
P&T Revision Date	9/10/2024	

Elexacaftor-tezacaftor-ivacaftor (TRIKAFTA)

Products Affected

TRIKAFTA

Criteria Details		
Required Medical Information	 Cystic Fibrosis Presence of at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA): F508del mutation A mutation in the CFTR gene that is responsive based on clinical, in vitro, or extrapolated data. Not to be used in combination with other CFTR modulator treatments 	
Age Restrictions	6 years of age and older	
Prescriber Restrictions	Pulmonologist or Specialist affiliated with a CF care center.	
Coverage Duration	Initial: 6 months Renewal: 12 months	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date	05/01/2025	
P&T Approval Date	03/11/2025	
P&T Revision Date	05/01/2021, 09/01/2021, 03/11/2025	
References	Alyftrek [package insert]. Boston, MA: Vertx Pharmaceuticals Inc.; 2025.	

Endothelin Receptor Antagonists

Products Affected

Ambrisentan tablets

• Bosentan Tablets

Criteria Details		
Required Medical Information	 Pulmonary Arterial Hypertension Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension) 	
Age Restrictions	Ambrisentan: 18 years of age and overBosentan: 3 years and up	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist.	
Coverage Duration	Initial: 6 months. Renewal: 12 months.	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date	9/1/2025	
P&T Approval Date	7/13/2021	
P&T Revision Date	7/8/2025, 7/13/2021	

Erythropoietic Agents

Products Affected

- PROCRIT
- RETACRIT

ARANESP

Prior Authorization Criteria

Criteria Details

Anemia due to Chronic Kidney Disease (CKD):

- Diagnosis of chronic kidney disease.
- Verification of adequate iron stores.
- Verification of anemia with hematocrit less than 30% or hemoglobin less than 10g/dL within 30 days of request AND
 - o Patient is on dialysis. OR
 - Patient is not on dialysis but the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion AND reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal.

Anemia in HIV Patients:

- Verification of adequate iron stores.
- Verification of anemia with hematocrit less than 36% or hemoglobin less than 12g/dL within 30 days of request.
- Serum erythropoietin less than or equal to 500mU/mL.
- Patient is receiving zidovudine therapy or diagnosed with HIV.

Anemia due to Chemotherapy:

- Verification that other causes of anemia have been ruled out.
- Verification of adequate iron stores.
- Verification of anemia with hematocrit less than 30% or hemoglobin less than 10g/dL within 2 weeks of request.
- Verification that the cancer is a non-myeloid malignancy.
- Patient is receiving chemotherapy.

Preoperative for reduction of allogeneic blood transfusion:

- Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery.
- Hemoglobin is greater than 10 to less than or equal to 13.

Required Medical Information

	Criteria Details
	 Patient is at high risk for perioperative transfusions. Patient is unwilling or unable to donate autologous blood preoperatively. Verification of adequate iron stores.
	 Anemia in Myelodysplastic Syndrome (MDS): Diagnosis of MDS. Serum erythropoietin less than or equal to 500mU/mL OR diagnosis of transfusion dependent MDS. Verification of adequate iron stores.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months. Renewal: 12 months. Anemia due to chemotherapy Initial: 3 months. Renewal: 3 months. Anemia in MDS Initial: 3 months. Renewal: 12 months. Preop Initial: 1 month.
Renewal Criteria	 Anemia due to CKD: One of the following: Patient is on dialysis and most recent or average Hct over 3 months is 33% or less (Hgb 11g/dL or less) Patient is not on dialysis and most recent or average Hct over 3 months is 30% or less (Hgb 10g/dL or less) Request is for a pediatric member and most recent or average Hct over 3 months is 36% or less (Hgb 12g/dL or less) Decrease in the need for blood transfusion OR Hemoglobin increased greater than or equal to 1d/dL from pre-treatment level Verification of adequate iron stores Anemia in HIV Patients: Most recent or average Hct over 3 months is 36% or less (Hgb 12g/dL or less). Decrease in the need for blood transfusion OR Hemoglobin increased greater than or equal to 1d/dL from pre-treatment level.

	Criteria Details
	 Anemia due to chemotherapy: Most recent or average Hct over 3 months is 30% or less (Hgb 10g/dL or less). Decrease in the need for blood transfusion OR Hemoglobin increased greater than or equal to 1d/dL from pre-treatment level. Patient is receiving chemotherapy.
	 Anemia in MDS Patients: Most recent or average Hct over 3 months is 36% or less (Hgb 12g/dL or less). Decrease in the need for blood transfusion OR Hemoglobin increased greater than or equal to 1.5d/dL from pre-treatment level.
Effective Date	9/1/2024
P&T Approval Date	7/9/2024
P&T Revision Date	7/9/2024

Erenumbab (AIMOVIG)

Products Affected

AIMOVIG 70MG/ML

AIMOVIG 140MG/ML

Criteria Details		
	Diagnosis of episodic migraines: Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month OR Diagnosis of chronic migraines: Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months AND medication overuse headache has been considered and potentially offending medication(s) have been discontinued AND	
Required Medical Information	Two of the following: History of failure or contraindication (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine) OR history of failure or contraindications (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) OR history of failure or contraindication (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol OR history of failure or contraindication (after at least a two month trial) or intolerance to Atacand (candesartan) AND medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines	
Age Restrictions	18 years of age or older	
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.	
Coverage Duration	Initial: 6 months. Renewal: 12 months.	
Renewal Criteria	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity AND use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan,	

Criteria Details		
	sumatriptan)] has decreased since the start of CGRP therapy AND medication will not be used in combination with another CGRP inhibitor for preventive treatment of migraines.	
	* AND For Chronic Migraine only: Patient continues to be monitored for medication overuse headache	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Etrasimod arginine (VELSIPITY)

Products Affected

VELSIPITY TAB

Criteria Details	
Required Medical Information	 Ulcerative Colitis (UC): Documentation of moderate-to-severe ulcerative colitis The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following: Mesalamine, sulfasalazine OR Mercaptopurine, azathioprine, OR Corticosteroids (prednisone, methylprednisolone) Trial and failure of both infliximab and adalimumab
Age Restrictions	Must be at least 18 years of age
Prescriber Restrictions	Prescribed by or in collaboration with a Gastroenterologist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission.
Effective Date	03/01/2024
P&T Approval Date	01/09/2024
P&T Revision Date	

Etanercept (ENBREL)

Products Affected

ENBREL

ENBREL SURECLICK

Prior Authorization Criteria

Criteria Details

ALL: must have a negative tuberculin test (TB)

AS: Patient has a documented diagnosis of ankylosing spondylitis. Clinical documentation showing an inadequate response, intolerance, or contraindication to one or more non-steroidal anti-inflammatory drugs NSAIDs (trial at maximum dose for at least 2-3 weeks before considering them as failures) or analgesic agents if NSAIDs do not completely control the pain, or sulfasalazine (if peripheral joint involvement is present).

Required Medical Information

CD: Clinical documentation showing an inadequate response, intolerance, or contraindication to budesonide, mesalamine, or corticosteroids, or non-biologic DMARDs (i.e., azathioprine, methotrexate, mercaptopurine. JIA: Clinical documentation showing inadequate response, intolerance, or contraindication to one or more NSAID **AND** one or more non-biologic DMARD (i.e., methotrexate, sulfasalazine).

PP: Patient has documented diagnosis of moderate to severe plaque psoriasis for at least 6 months with at least one of the following: Incapacitation due to plaque location (e.g., head and neck, palms, soles, or genitalia) OR Involvement of at least 10 percent of body surface area (BSA) OR Psoriasis Area and Severity Index (PASI) score of 12 or greater, AND patient is free of any clinically important active infections AND clinical documentation of inadequate or non-candidate to a 3-month minimum trial of at least 1 systemic agent (e.g., immunosuppressive, retinoic acid derivatives, and/or methotrexate, AND did not respond or non-candidate to a 3-month minimum trial of phototherapy.

Age Restrictions

Criteria Details	
Initial: 12 months. Renewal: 12 months.	
Documentation of positive clinical response to therapy.	

Fezolinetant (VEOZAH)

Products Affected

VEOZAH TAB 45MG

	Criteria Details
Required Medical Information	 Vasomotor Symptoms (VMS): Diagnosis of moderate-to-severe VMS due to menopause Documented contraindication, intolerance, or inadequate response to at least 2 hormonal therapies AND Documented contraindication, intolerance, or inadequate response to two nonhormonal therapies (e.g., one SNRI and one SSRI).
Age Restrictions	
Prescriber Restrictions	Gynecologist
Coverage Duration	Initial: 3 months. Renewal: 12 months.
Renewal Criteria	Documentation of at least 50% reduction in VMS from baseline.
Effective Date	09/01/2023
P&T Approval Date	7/11/2023
P&T Revision Date	

Fingolimod (GILENYA)

Products Affected

GILENYA

Criteria Details	
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Fremanezumab-vfrm (AJOVY)

Products Affected

AJOVY

	Criteria Details
	Diagnosis of episodic migraines: Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month OR
	Diagnosis of chronic migraines : Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months AND medication overuse headache has been considered and potentially offending medication(s) have been discontinued
	AND
Required Medical Information	Two of the following: History of failure or contraindication (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine) OR history of failure or contraindications (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) OR history of failure or contraindication (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol OR history of failure or contraindication (after at least a two month trial) or intolerance to Atacand (candesartan) AND medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity AND use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy AND

	Criteria Details
	medication will not be used in combination with another CGRP inhibitor for preventive treatment of migraines.
	* AND For Chronic Migraine only: Patient continues to be monitored for medication overuse headache
Effective Date	
P&T Approval Date	
P&T Revision Date	

Galcanezumab-gnlm (EMGALITY)

Products Affected

• Emgality 100mg/mL

	Criteria Details
Required Medical Information	Diagnosis of episodic cluster headache AND patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least 3 months AND medication will not be used in combination with another CGRP inhibitor.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist.
Coverage Duration	Initial: 3 months Renewal: 12 months
Renewal Criteria	Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity AND Medication will not be used in combination with another CGRP inhibitor.
Effective Date	
P&T Approval Date	
P&T Revision Date	

General Oncology

Products Affected

- Abemaciclib (Verzenio)
- Abiraterone
- Acalabrutinib (Calquence)
- Adagrasib (Krazati)
- Alectinib (Alecensa)
- Alpelisib (Piqray)
- Ascinimib (Scemblix)
- Asparaginase Erwinia (Rylaze)
- Avutometinib/Defactinib (Avmapki/Fazynja Pak)
- Belzutifan (Welireg)
- Binimetinib (Mektovi)
- Bosutinib (Bosulif)
- Brigatinib (Alunbrig)
- Busulfan (Myleran)
- Cabozantinib (Cabometyx)
- Capivasertib (Trugap)
- Capmatinib (Tabrecta)
- Ceritinib (Zykadia)
- Chlorambucil (Leukeran)
- Crizotinib (Xalkori)
- Dabrafenib (Tafinlar)
- Dasatinib
- Darolutamide (Nubega)
- Elacestrant (Orserdu)
- Encorafenib (Braftovi)
- Entrectinib (Rozlytrek)
- Erlotinib
- Estramustine (Emcyt)
- Everolimus
- Fruquintinib (Fruzagla)
- Futibatinib (Lytgobi)
- Gefitinib
- Ibrutinib (Imbruvica)
- Inavolisib (Itovebi)
- Infigratinib (Truseltig)
- Ivosidenib (Tibsovo)

- Lazertinib (Lazcluze)
- Lenvatinib (Lenvima)
- Lomustine (Gleostine)
- Lorlatinib (Lobrena)
- Midostaurin (Rydapt)
- Mobocertinib (Exkivity)
- Nilotinib (Tasigna)
- Niraparib/Abiraterone (Akeega)
- Nirogacestat (Ogsiveo)
- Olaparib (Lynparza)
- Olutasidenib (Rezlidhia)
- Osimertinib (Tagrisso)
- Pacritinib (Vonjo)
- Palbociclib (Ibrance)
- Pazopanib
- Pemigatinib (Pemazyre)
- Pexidartinib (Turalio)
- Pirtobrutinib (Jaypirca)
- Ponatinib (Iclusig)
- Quizartinib (Vanflyta)
- Repotrectinib (Augtyro)
- Ribociclib (Kisgali)
- Selpercatinib (Retevmo)
- Sorafenib
- Sotorasib (Lumakras)
- Sunitinib
- Talazoparib (Talzenna)
- Taletrectinib (Ibtrozi)
- Temozolomide
- Tepotinib (Tepmetko)
- Tivozanib (Fotivda)
- Topotecan (Hycamtin)
- Tovorafenib (Ojemda)
- Trametinib (Mekinist)
- Trifluridine/Tipiracil (Lonsurf)
- Tucatinib (Tukysa)
- Vimseltinib (Romvimza)

- Vorasidinib (Voranigo)Vorinostat (Zolinza)

	Criteria Details
Required Medical Information	Medication is being used for an FDA approved age AND medication is being used for FDA approved indication OR Medication is being used according to National Comprehensive Cancer Network (NCCN) guidelines
Age Restrictions	
Prescriber Restrictions	Prescribed by oncologist
Coverage Duration	Initial: 3 months. Renewal: up to 6 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	9/1/2025
P&T Approval Date	10/01/2022
P&T Revision Date	9/9/2025, 7/8/2025, 5/13/2025, 3/11/2025, 5/14/2024, 03/01/2024, 01/09/2024, 11/1/2023, 09/01/2023, 7/11/2023, 05/09/2023, 03/14/2023, 01/10/2023, 10/01/2022

Glatiramer (GLATOPA)

Products Affected

• GLATIRAMER INJ 20MG/ML

• GLATIRAMER INJ 40MG/ML

Criteria Details	
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis
Age Restrictions	
Prescriber Restrictions	Prescribed by Neurologist
Coverage Duration	Initial: 12 months. Renewal: up to 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy
Effective Date	
P&T Approval Date	
P&T Revision Date	

Glucagon-Like Peptide-1 (GLP1s) Receptor Agonist

Products Affected

- BYDUREON BCISE
- BYETTA 10 MCG PEN
- BYETTA 5MCG PEN

- TRULICITY
- VICTOZA

	Criteria Details
Required Medical Information	Patient must have clinically diagnosed Type 2 Diabetes AND patient must have adequate trial of, or contraindication to an SGLT-2 if patient has HF or high risk/established ASCVD or a DPP-4 if no high risk/established ASCVD AND a maximal tolerated doses of metformin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months, Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	
	•

Golimumab (SIMPONI)

Products Affected

SIMPONI INJ 50/0.5ML

SIMPONI INJ 100MG/ML

Prior Authorization Criteria

Criteria Details

Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active RA AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine AND patient is receiving concurrent therapy with methotrexate OR has a contraindication or intolerance to methotrexate.

Psoriatic Arthritis (PsA): Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.

Required Medical Information

Ankylosing Spondylitis (AS): Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to two NSAIDs (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen).

Polyarticular Juvenile Idiopathic Arthritis (PJIA): Diagnosis of moderate to severely active PJIA AND trial and failure, contraindication, or intolerance to one of the following non-biologic disease-modifying antirheumatic drugs (DMARDs): leflunomide, methotrexate.

Ulcerative Colitis (UC): Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids. AND trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone).

Age Restrictions

	Criteria Details
Prescriber Restrictions	RA, AS, PJIA: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist UC: Prescribed by or in consultation with a gastroenterologist
Coverage Duration	RA, PsA, AS, PJIA: Initial: 6 months; Renewal: 12 months UC: Initial: 10 weeks; Renewal: 12 months
	RA, PJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline.
	PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.
Renewal Criteria	AS: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.
	UC: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Gonadotropin-Releasing Hormone Agonists

Products Affected

- Lupron Depot
- Eligard

- Lupron
- Leuprorelin

Prior Authorization Criteria

Criteria Details

Endometriosis

- Diagnosis of endometriosis
- One of the following:
 - History of inadequate pain control response following a trial of at least 6 months, or history of intolerance or contraindication to one of the following:
 - Danazol
 - Combination (estrogen/progestin) oral contraceptive
 - **Progestins**
 - Patient has had surgical ablation to prevent recurrence

Uterine Leiomyomata (Fibroids) - For the reduction of the size of fibroids [off-label]

For use prior to surgery to reduce the size of fibroids to facilitate a surgical procedure (e.g., myomectomy, hysterectomy)

Information

Uterine Leiomyomata (Fibroids) - Anemia

- For the treatment of anemia
- Anemia is caused by uterine leiomyomata (fibroids)
- Patient has tried and had an inadequate response to at least 1 month of monotherapy with iron
- Used in combination with iron therapy
- For use prior to surgery

Central Precocious Puberty (CPP)

- Diagnosis of central precocious puberty (idiopathic or neurogenic)
- Early onset of secondary sexual characteristics in one of the following:
 - Females less than 8 years of age
 - Males less than 9 years of age

Required Medical

	Criteria Details
	Criteria Details Advanced bone age of at least one year compared with chronological age One of the following: Both of the following: Patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing Peak luteinizing hormone (LH) level above prepubertal range Patient has a random LH level in the pubertal range Patient had one of the following diagnostic evaluations to rule out tumors, when suspected: Diagnostic imaging of the brain (MRI or CT scan) (in patients with symptoms suggestive of a brain tumor or in those 6 years of age or younger) Pelvic/testicular/adrenal ultrasound (if steroid levels suggest suspicion) Adrenal steroids to rule out congenital adrenal hyperplasia (when pubarche precedes thelarche or gonadarche) Patient has no suspected tumors Prostate Cancer Diagnosis of advanced or metastatic prostate cancer Trial and failure, contraindication, or intolerance to any brand Lupron formulation Gender Dysphoria/Gender Incongruence (off-label) Using gonadotropin for suppression of puberty
	Diagnosis of gender dysphoria/gender incongruence
Age Restrictions	
Prescriber Restrictions	Central Precocious Puberty (CPP): Pediatric endocrinologist
Coverage Duration	Endometriosis: Initial: 6 months; Renewal: 6 months Uterine Leiomyomata (Fibroids): Initial: 4 months; Renewal: 3 months Uterine Leiomyomata (Fibroids) – Anemia Initial: 3 months Central Precocious Puberty (CPP): Initial: 12 months; Renewal: 12 months Prostate Cancer: Initial: 12 months; Renewal: 12 months Gender Dysphoria/Gender Incongruence: Initial: 12 months; Renewal:
	12 months

	Criteria Details
	 Recurrence of symptoms following a trial of at least 6 months with leuprolide acetate Used in combination with one of the following: Norethindrone 5 mg daily Other "add-back" sex-hormones (e.g., estrogen, medroxyprogesterone) Other bone-sparing agents (e.g., bisphosphonates) Central Precocious Puberty (CPP) LH levels have been suppressed to pre-pubertal levels Prescribed by or in consultation with a pediatric endocrinologist Prostate Cancer Diagnosis of advanced or metastatic prostate cancer
Effective Date	8/1/2024
P&T Approval Date	5/13/2024
P&T Revision Date	

Grass Pollen Allergen Extract – Timothy Grass (GRASTEK)

Products Affected

• GRASTEK

Criteria Details		
Required Medical Information	Grass pollen-induced allergic rhinitis.	
Age Restrictions		
Prescriber Restrictions	Prescribed by Allergy and Immunology specialist.	
Coverage Duration	Initial: 3 months. Renewal: 3 months.	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Guselkumab (TREMFYA)

Products Affected

- Tremfya Auto-Injector
- Tremfya IV Solution

Tremfya Prefilled Syringe

Prior Authorization Criteria

Criteria Details

Plaque Psoriasis

- Diagnosis of moderate to severe plaque psoriasis
- One of the following:
 - Greater than or equal to 3% body surface area involvement
 - Severe scalp psoriasis
 - o Palmoplantar (i.e., palms, soles), facial, or genital involvement
- Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies:
 - o Corticosteroids (e.g., betamethasone, clobetasol)
 - Vitamin d analogs (e.g., calcitriol, calcipotriene)
 - Tazarotene
 - Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
 - Anthralin
 - o Coal tar

Required Medical Information

Psoriatic Arthritis

- Diagnosis of active psoriatic arthritis
- One of the following:
 - Actively inflamed joints
 - o Dactylitis
 - o Enthesitis
 - Axial disease
 - Active skin and/or nail involvement

Crohn's Disease

- Diagnosis of moderately to severely active Crohn's Disease
- One of the following (for SC or IV induction):
 - Documentation of one of the following:
 - Frequent Diarrhea and abdominal pain
 - At least 10% weight loss

	Criteria Details
	 Complications such as obstruction, fever, abdominal mass Abnormal lab values (e.g. C-reactive protein) CD Activity Index (CDAI) greater than 220 For SC dosing: will be used for maintenance dosing following IV induction dose
	Ulcerative Colitis ■ Diagnosis of moderately to severely active ulcerative colitis ■ One of the following (for IV induction): □ Documentation of one of the following: ■ Greater than 6 stools per day ■ Frequent blood in the stools ■ Frequent urgency ■ Presence of ulcers ■ Abnormal lab values (e.g., hemoglobin, erythrocyte sedimentation rate, C-reactive protein) ■ Dependent on, or refractory to, corticosteroids ○ For SC dosing: will be used for maintenance dosing following IV induction dose
Age Restrictions	
Prescriber Restrictions	 Plaque Psoriasis: Dermatologist Psoriatic Arthritis: Dermatologist Rheumatologist Crohn's Disease and Ulcerative Colitis: Gastroenterologist
Coverage Duration	 Plaque Psoriasis and Psoriatic Arthritis: Initial: 6 months Renewal: 12 months Crohn's Disease and Ulcerative Colitis: Initial IV: 3 months Initial SC: 6 months Renewal SC: 12 months
Renewal Criteria	 Plaque Psoriasis – Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction the body surface area (BSA) involvement from baseline Improvement in symptoms (e.g., pruritus, inflammation) from baseline

	Criteria Details
	 Psoriatic Arthritis - Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in the total active (swollen and tender) joint count from baseline Improvement in symptoms (e.g., pain, stiffness, pruritus, inflammation) from baseline Reduction in the body surface area (BSA) involvement from baseline Crohn's Disease, Ulcerative Colitis - Documentation of positive clinical response to therapy as evidenced by at least one of the following: Improvement in intestinal inflammation (e.g. mucosal healing, improvement in lab values) from baseline Reversal of high fecal output state
Effective Date	9/1/2025
P&T Approval Date	10/28/2022
P&T Revision Date	7/8/2025, 5/8/2023, 10/28/2022

Inclisiran (LEQVIO)

Products Affected

• LEQVIO SOLUTION

Criteria Details		
Required Medical Information	 Established clinical ASCVD: Documentation of very high risk ASCVD as evidenced by either: History of multiple major ASCVD events OR One major ASCVD event AND multiple high-risk conditions. Documentation of a current LDL greater than or equal to 55 mg/dl. Documentation that: Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins AND Is receiving ezetimibe or has a documented intolerance to ezetimibe. Documentation of failure of PCSK9 inhibitor (Repatha or Praluent). 	
	 Primary or familial hyperlipidemia: Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL. Documentation of current LDL greater than 100 mg/dL. Documentation that:	
Age Restrictions		
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.	

Criteria Details	
Coverage Duration	Initial: 6 months; Renewal: 12 months
Renewal Criteria	Documented positive clinical response to therapy (significant decrease in lipid levels).
Effective Date	11/1/2024
P&T Approval Date	9/10/2024
P&T Revision Date	9/10/2024

Infigratinib (TRUSELTIQ)

Products Affected

- TRUSELTIQ (100MG DAILY DOSE)
- TRUSELTIQ (125MG DAILY DOSE)
- TRUSELTIQ (50MG DAILY DOSE) TRUSELTIQ (75MG DAILY DOSE)

Criteria Details	
Required Medical Information	Confirmation of trial and failure of guideline directed therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	Initial: 3 months. Renewal: 3 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Insulin Degludec (TRESIBA)

Products Affected

• TRESIBA FLEXTOUCH U-100 & U-200

	Criteria Details
Required Medical Information	U-100 & *U-200: Must have tried and failed formulary long-acting insulin analogues OR have documented intolerance or contraindication to formulary long-acting insulin analogues AND have significant barriers to sardized administration requiring flexibility in dose timing.
	*(U-200) Patient must require greater than 160 units of insulin per dose AND have difficulty with multiple daily injections.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	5/1/2024
P&T Approval Date	3/12/2024
P&T Revision Date	3/12/2024

Interferon Alfa-2b (INTRON A)

Products Affected

INTRON A

	Criteria Details
	Chronic hepatitis B: Diagnosed with chronic hepatitis B infection AND patient is without decompensated liver disease.
	Chronic hepatitis C: Diagnosed with chronic hepatitis C infection AND patient is without decompensated liver disease AND patient has not previously been treated with interferon AND is prescribed for use with ribavirin OR patient has intolerance or contraindication to ribavirin.
	Metastatic renal cell carcinoma (RCC): Diagnosed with metastatic RCC AND prescribed in combination with Avastin (bevacizumab).
Required Medical Information	AIDS-related Kaposi sarcoma (KS): Diagnosed with AIDS-related KS.
iniormation	Condylomata acuminata (CA) : Diagnosed with CA involving external surfaces of the genital & perianal areas.
	Follicular lymphoma (FL): Diagnosed with clinically aggressive follicular non-Hodgkin lymphoma. Prescribed in conjunction with anthracycline-containing combination chemotherapy.
	Hairy cell leukemia (HCL): Diagnosed with HCL. Melanoma: Diagnosed with malignant melanoma. Prescribed as adjuvant to surgical treatment who are free of disease but at high risk for systemic recurrence AND must be administered within 56 days of surgery.
Age Restrictions	Patient must be 18 years or older.
Prescriber Restrictions	Prescribed by a specialist.
Coverage Duration	Initial: 12 months. Renewal: 12 months.

	Criteria Details
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Interferon beta-1a (AVONEX)

Products Affected

AVONEX PEN

AVONEX PREFILLED

	Outhoute Dotatle
	Criteria Details
Required Medical Information	Diagnosis of a relapsing form of Multiple Sclerosis AND trial and failure, contraindication, or intolerance to all of the following: dimethyl furmate, fingolimod, glatiramer acetate/glatopa.
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Interferon beta-1a (REBIF)

Products Affected

- REBIF INJ 22/0.5ML
- REBIF INJ 44/0.5 ML

- REBIF REBIDO INJ 22/0.5ML
- REBIF REBIDO INJ 44/0.5ML

	Criteria Details
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis AND trial and failure, contraindication, or intolerance to all of the following: dimethyl furmate, fingolimod, glatiramer acetate/glatopa, avonex.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Interferon beta-1b (EXTAVIA)

Products Affected

EXTAVIA INJ 0.3MG

	Criteria Details
Required Medical Information	Diagnosis of relapsing forms of multiple sclerosis AND trial and failure, contraindication, or intolerance to all of the following: dimethyl furmate, fingolimod, glatiramer acetate/glatopa.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ivacaftor (KALYDECO)

Products Affected

KALYDECO

	Criteria Details
Required Medical Information	Diagnosis of Cystic Fibrosis with documentation showing at least one CFTR gene mutation that has shown to be responsive to Kalydeco
Age Restrictions	6 months of age and older
Prescriber Restrictions	Prescribed by pulmonologist
Coverage Duration	Initial: 3 months Renewal: 6 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ivermectin (STROMECTOL)

Products Affected

IVERMECTIN

	Criteria Details
Required Medical Information	Treatment of FDA approved diagnosis including Strongyloidiasis, Onchocerciasis, Infestation by Phthirus pubis, Scabies, Enterobiasis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 6 months, Renewals: reinfection 6 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ivosidenib (TIBSOVO)

Products Affected

TIBSOVO

Criteria Details	
Required Medical Information	Acute Myeloid Leukemia (AML): Test confirmed IDH1 mutation Cholangiocarcinoma: Test confirmed IHD1 mutation AND previous treatment with at least one chemotherapy regimen (e.g. FOLFOX)
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by an oncologist
Coverage Duration	Initial: 3 months. Renewal: 6 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ixekizumab (TALTZ)

Products Affected

TALTZ

Prior Authorization Criteria

Criteria Details

Plaque Psoriasis (PsO): Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement AND a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar AND trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, adalimumab, Skyrizi, Ustekinumab, or Tremya.

Required Medical Information

Psoriatic Arthritis (PsA): Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, Enbrel, adalimumab, Simponi, Ustekinumab, Tremfya, Skyrizi, Rinvoq, or Xeljanz

Ankylosing Spondylitis (AS): Diagnosis of active ankylosing spondylitis **AND** minimum duration of one month trial and failure, contraindication, or intolerance to **TWO** non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen) **AND** trial and failure, contraindication, or intolerance to ONE of the following: Cimzia, Enbrel, adalimumab, Simponi, Rinvog, Xeljanz.

Non-radiographic Axial Spondyloarthritis (nr-axSpA): Diagnosis of active non-radiographic axial spondyloarthritis AND patient has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints.) AND minimum duration of one month trial and failure, contraindication, or intolerance to **TWO** non-steroidal anti-inflammatory drugs (NSAIDs) (e.g.,

	Criteria Details
	diclofenac, ibuprofen, meloxicam, naproxen) AND trial and failure,
	contraindication, or intolerance to Cimzia.
Age Restrictions	
Prescriber Restrictions	Plaque Psoriasis (PP): Prescribed by or in consultation with a dermatologist Psoriatic Arthritis (PsA): Prescribed by or in consultation with a dermatologist or rheumatologist Ankylosing Spondylitis (AS): Prescribed by or in consultation with a rheumatologist Non-radiographic Axial Spondyloarthritis: Prescribed by or in consultation with a rheumatologist
Coverage Duration	Initial: 12 months; Renewal: 12 months
Renewal Criteria	Ps0: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or improvement in symptoms (e.g., pruritus, inflammation) from baseline.
	PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline.
	AS, nr-axSpA: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Lacosamide (VIMPAT)

Products Affected

- Lacosamide TAB 50MG, 100MG, 150MG, 200MG
- Lacosamide Solution 10MG/ML

	Criteria Details
Required Medical Information	 Focal Seizures Documentation confirming epilepsy or seizure disorder Solution only: Member under age 10 or unable to use tablets Primary Generalized Tonic-Clonic Seizures Documentation confirming epilepsy or seizure disorder Solution only: Member under age 10 or unable to use tablets
Age Restrictions	 Solution only One of the following: Pediatric member age 10 or under Documentation inability of the member to use the preferred tablet formulation
Prescriber Restrictions	Neurologist
Coverage Duration	Initial: 12 months. Renewal: Lifetime.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	5/1/2025
P&T Approval Date	7/11/2023
P&T Revision Date	3/11/2025, 3/12/2024, 7/11/2023

Lanthanum Carbonate (FOSRENOL)

Products Affected

- Lanthanum carbonate 500MG
- Lanthanum carbonate 750MG
- Lanthanum carbonate 1000MG

	Outhority Doubling
	Criteria Details
Required Medical Information	Diagnosis of hyperphosphatemia in chronic kidney disease AND trial and failure, contraindication, or intolerance (at least 6 weeks) to both maximally tolerated calcium acetate and sevelamer carbonate
Age Restrictions	6 years or older
Prescriber Restrictions	Nephrologist
Coverage Duration	Initial: 6 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Lasmiditan (REYVOW)

Products Affected

REYVOW 100MG TAB

• REYVOW 50MG TAB

	Criteria Details
Required Medical Information	Medication intended for use for acute use for the treatment of migraine headaches AND documentation patient is on preventative therapy AND trial and failure (defined as trial period of 6 weeks per agent) or contraindication to at least 3 generic oral formulary triptans use at up to the maximally indicated dosing and in combination with NSAID therapy (naproxen) OR trial and failure to intolerance to NSAID treatment alone if triptans contraindicated OR contraindication to all triptans and NSAIDs
Age Restrictions	Patient is 18 years of age or older
Prescriber Restrictions	Prescribed by neurologist or headache specialist
Coverage Duration	Initial: 3 months Renewal: 6 months
Renewal Criteria	Documented positive clinical response to therapy
Effective Date	
P&T Approval Date	
P&T Revision Date	

Lenacapavir (Sunlenca)

Products Affected

Sunlenca Therapy Pack

• Sunlenca Subcutaneous

	Criteria Details
Required Medical Information	 Multi-Drug-Resistant HIV Diagnosis of MDR HIV-1 infection with resistance to at least two drugs in each of at least three of the following classes: NRTIs, NNRTIs, PIs, and INSTIs Will be used in combination with an optimized baseline regimen (OBR) Current ARV regimen has been stable for at least 2 months HIV-1 RNA is ≥400 copies/mL
Prescriber Restrictions	HIV Specialist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	 Continues to be used in combination with an optimized background regimen (OBR) Provider states that patient continues to receive clinical benefit from the treatment
Effective Date	4/1/2023
P&T Approval Date	3/14/2023
P&T Revision Date	3/14/2023

Lenacapavir (Yeztugo)

Products Affected

Yeztugo inj.

• Yeztugo Tablets

	Criteria Details
Required Medical Information	 Pre-Exposure Prophylaxis (PrEP) Confirmation the drug is being used for PrEP (a different product is used for treatment) Submission of medical records documenting both of the following U.S. Food and Drug (FDA)-approved tests prior to use of Yeztugo: Negative HIV-1 antigen/antibody test Negative HIV-1 RNA assay Trial and failure, contraindication, or intolerance to generic emtricitabine-tenofovir disoproxil fumarate 200/300mg.
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	 Provider attests that patient is adherent to the testing appointments and scheduled injections of Yeztugo Submission of medical records documenting both of the following U.S. Food and Drug (FDA)-approved tests prior to use of Yeztugo: Negative HIV-1 antigen/antibody test Negative HIV-1 RNA assay
Effective Date	11/1/2025
P&T Approval Date	9/9/2025
P&T Revision Date	9/9/2025

Lebrikizumab (EBGLYSS)

Products Affected

Ebglyss autoinjector

• Ebglyss prefilled syringe

	Criteria Details
Required Medical Information	Atopic dermatitis: Diagnosed with severe atopic dermatitis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) ≥ 11 or Children's Dermatology Life Quality Index (CDLQI) ≥ 13 (or severe score on another validated tool) One or more of the following: At least 10% of body surface area involvement Hand, foot, or mucous membrane involvement Documented contraindication or failed trial to ALL of the following: Moderate-high potency corticosteroid (e.g., clobetasol, fluocinonide, fluticasone) Topical calcineurin inhibitor (e.g. tacrolimus) Oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil) OR the member is oral corticosteroid dependent.
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist.
Coverage Duration	Initial: 6 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy
Effective Date	1/1/2025
P&T Approval Date	11/12/2024
P&T Revision Date	11/12/2024

Lidocaine Topical Anesthetic (LIDODERM)

Products Affected

• LIDOCAINE EXTERNAL PATCH 5%

	Criteria Details
Required Medical Information	Diagnosis of post-herpetic neuralgia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 3 months. Renewal: 3 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Lifitegrast (XIIDRA)

Products Affected

XIIDRA

	Criteria Details
Required Medical Information	The patient has a diagnosis of lack of tear production due to ocular inflammation associated with keratoconjunctivitis sicca AND ONE of the following: • The patient is not currently using a topical ophthalmic anti-inflammatory drug or punctal plug OR • The patients current use of topical ophthalmic anti-inflammatory drug or punctal plug will be discontinued before starting the requested agent AND The patient has previously tried or is currently using aqueous enhancements (e.g. artificial tears, gels, ointments) OR • The patient has a documented intolerance, contraindication, or hypersensitivity to aqueous enhancements. The patient is not currently using Restasis OR the patients current use of Restasis will be discontinued before starting Xiidra.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Linezolid (ZYVOX)

Products Affected

LINEZOLID

LINEZOLID IN SODIUM CHLORIDE

	Criteria Details
Required Medical Information	Clinically documented infection that is susceptible to linezolid if the patient has a severe allergy to beta lactamase inhibitors or any antibiotic that the organism is susceptible OR clinically documented infection that is susceptible to linezolid if the patient has failed treatment with antibiotics that the organism is susceptible OR clinically documented Vancomycin-Resistant Enterococcus faecium infection OR clinically documented MRSA AND has failed or is intolerant to Vancomycin if the organism is susceptible to Vancomycin.
Age Restrictions	Solution only One of the following: Pediatric member age 10 or under Documentation inability of the member to use the preferred tablet formulation
Prescriber Restrictions	Prescribed by or in consultation with an Infectious Disease specialist.
Coverage Duration	Initial: length of treatment. Renewal: length of treatment.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	5/1/2024
P&T Approval Date	3/12/2024
P&T Revision Date	3/12/2024

Lisdexamfetamine (VYVANSE)

Products Affected

LISDEXAMFETAMINE CAPS

	Criteria Details
Required Medical Information	ADHD: Prior trial (30-day trial) of an extended-release amphetamine product (amphetamine salts ER, dextroamphetamine ER, etc.) and an extended-release methylphenidate product (dexmethylphenidate ER, methylphenidate ER). BED: Clinical documentation confirming binge eating disorder diagnosis
	per DSM-5 criteria AND trial and failure of at least two therapeutic alternatives including SSRIs, topiramate, and/or methylphenidate.
Age Restrictions	BED: 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	03/01/2025
P&T Approval Date	01/14/2025
P&T Revision Date	01/14/2025

Long Acting Opiates AND Dolophine

Products Affected

- FENTANYL PATCH 72 HOUR 100 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 12 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 25 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 37.5 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 50 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 62.5 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 75 MCG/HR TRANSDERMAL
- FENTANYL PATCH 72 HOUR 87.5 MCG/HR TRANSDERMAL
- HYDROCODONE BITARTRATE ER
- HYDROMORPHONE HCL ER
- METHADONE HCL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 120 MG ORAL

- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 30 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 45 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 60 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 75 MG ORAL
- MORPHINE SULFATE ER BEADS CAPSULE EXTENDED RELEASE 24 HOUR 90 MG ORAL
- MORPHINE SULFATE ER ORAL CAPSULE EXTENDED RELEASE 24 HOUR
- MORPHINE SULFATE ER ORAL TABLET EXTENDED RELEASE
- NUCYNTA ER
- OXYCODONE HCL ER
- OXYCONTIN
- OXYMORPHONE HCL ER

XTAMPZA ER

	Criteria Details
	Cancer, end of life, or palliative care: No coverage restrictions.
Required Medical Information	Non-cancer/end of life care: Documented use of current and/or recent usage of short-acting opioids for at least 15 days prior to long-acting opioids. • For opioid naive (14 or fewer days filled in previous 120 days): 7-day maximum quantity limit, equal to or less than 50 MED [morphine equivalents per day].

Criteria Details	
	For opioid experienced (greater than or equal to 15 days filled in previous 120 days): equal to or less than 90 MED [morphine equivalents per day]. Restricted to 2 fills in a 60-day period for both naive AND experienced.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Lotilaner (XDEMVY)

Products Affected

Xdemvy 0.25% Ophthalmic solution

	Criteria Details
Required Medical Information	Diagnosis: Demodex Blepharitis Documentation of at least mild erythema of the upper eyelid margin Presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination
Age Restrictions	18 years of age or older
Prescriber Restrictions	Optometrist or Ophthalmologist
Coverage Duration	Initial: 6 weeks. Renewal: No renewals allowed
Renewal Criteria	
Effective Date	5/1/2024
P&T Approval Date	3/12/2024
P&T Revision Date	3/12/2024

Lumacaftor/ivacaftor (ORKAMBI)

Products Affected

ORKAMBI ORAL PACKET

ORKAMBI ORAL TABLET

	Criteria Details
Required Medical Information	Clinical documentation of cystic fibrosis diagnosis with homozygous F508del mutation.
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by pulmonologist
Coverage Duration	Initial: 3 months Renewal: 6 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Methylphenidate solution/chewable

Products Affected

 METHYLPHENIDATE HCL ORAL SOLUTION METHYLPHENIDATE HCL ORAL TABLET CHEWABLE

	Criteria Details
Required Medical Information	Documentation that the patient has difficulty swallowing pills and/or has tried and failed methylphenidate tablets.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Mirabegron (Myrbetriq)

Products Affected

• Mirabegron ER tablets

Covered Uses	 All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design Overactive Bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency Neurogenic detrusor overactivity in pediatric members
Required Medical Information and Criteria	Overactive Bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency Documented trial and failure, intolerance, or contraindication to at least 3 of the following: Oxybutynin IR or ER Fesoterodine Solifenacin Tolterodine IR or ER Trospium IR or ER (requires step therapy through oxybutynin) Neurogenic Detrusor Overactivity in pediatric members Is there documented trial and failure, intolerance, or contraindication to both of the following: Oxybutynin IR or ER Solifenacin
Renewal Criteria	Documentation of positive clinical response to therapy
Age Restriction	3 years of age and older
Prescriber Restriction	• N/A
Coverage Duration	All Diagnoses: Initial: 12 months Renewal: 12 months

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

References

- Cameron AP, Chung DE, Dielubanza EJ, et al. The AUA/SUFU guideline on the diagnosis and treatment of idiopathic overactive bladder. J Urol. Published online April 23, 2024.
- Myrbetriq [package insert]. Northbrook, IL: Astellas.; 2021.

Mitapivat (PYRUKYND)

Products Affected

PYRUKYND

	Criteria Details
Required Medical Information	Diagnosis of PKD with at least two mutations within the PKLR gene, including a missense mutation AND confirmation of current hemoglobin is ≤ 10mg/dL AND patient is not homozygous for the R479H mutation AND does not have two non-missense variants in the PKLR gene, without the presence of another missense variant AND patient has had at least 6 RBC transfusions within the previous year for hemolytic anemia due to PKD AND prescriber confirmed concomitant use of daily folic acid AND confirmation that the patient does not have moderate or severe hepatic dysfunction.
Age Restrictions	At least 18 years of age
Prescriber Restrictions	Prescribed by or in consultations with a hematologist
Coverage Duration	Initial: 3 months. Renewal: 6 months.
Renewal Criteria	Clinical documentation showing an increase in Hb at least 1.5 mg/dL over baseline and/or a reduction in frequency of transfusions.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Mirikizumab-mrkz (OMVOH)

Products Affected

Omvoh

	Criteria Details
Required Medical Information	All diagnoses: Initial testing for latent TB and treatment, if necessary, before starting treatment. No current active infection at initiation of therapy. Risks and benefits documented in cases of chronic or recurrent infection. Will NOT be used in combination with another biologic or Otezla Ulcerative Colitis (UC): Documentation of moderate-to-severe ulcerative colitis The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following: Mesalamine, sulfasalazine OR Mercaptopurine, azathioprine, OR Corticosteroids (prednisone, methylprednisolone) Trial and failure of both infliximab and adalimumab
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultations with a Gastroenterologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Effective Date	7/1/2024
P&T Approval Date	5/14/2024

	Criteria Details
P&T Revision Date	

Naltrexone (VIVITROL)

Products Affected

• VIVITROL INJ.

	Criteria Details
Required Medical Information	The medication will be sent directly to the administering provider and will not be dispensed directly to the member
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	5/1/2024
P&T Approval Date	3/12/2024
P&T Revision Date	3/12/2024

Omalizumab (XOLAIR)

Products Affected

XOLAIR AUTO-INJECTOR

XOLAIR PREFILLED SYRINGE

Prior Authorization Criteria

Criteria Details

Severe Asthma:

- Confirmed diagnosis of moderate to severe persistent asthma.
- Positive skin test or RAST to a perennial aeroallergen.
- Baseline IgE serum level within FDA label.
- Documentation of steps taken to avoid, within reason, environmental allergens and other triggers environmental allergens and other triggers.
- Documented trial and failure, with claims history of adherence to:
 - High dose inhaled corticosteroid with a long-acting beta agonist (e.g., Advair),
 - o Long acting anti-muscarinic (e.g., Spiriva).
 - Leukotriene Inhibitor (e.g., Singulair).
- Documented trial and failure of, or contraindication to allergen immunotherapy.

Required Medical Information

Chronic Rhinosinusitis with Nasal Polyps (CRSwNP):

- Confirmed diagnosis of chronic rhinosinusitis with nasal polyps (CRSwNP).
- Documentation of recurrent nasal polyps after prior sinus surgery.
- Documented risk of another sinus surgery, or a statement why sinus surgery is not medically appropriate.
- Documented trial and failure, with claims history of adherence to:
 - At least 2 intranasal corticosteroids (e.g., fluticasone, mometasone),
 - o Sinuva.
- Documentation that Xolair is intended as adjunct therapy with nasal corticosteroids.

Chronic Idiopathic Urticaria- refractory (CIU):

Documentation of chronic spontaneous or idiopathic urticaria.

	Criteria Details
	 Documented trial and failure of at least 6 weeks of maximally tolerated doses of all the following: 1st generation antihistamine – (e.g., doxepin, hydroxyzine) 2nd generation antihistamine – (e.g., cetirizine, levocetirizine, fexofenadine, loratadine, desloratadine) Histamine Type-2 Receptor Antagonists (e.g., famotidine, cimetidine) Leukotriene inhibitor (e.g., montelukast, zafirlukast) IgE-Mediated Food Allergy: Diagnosis of IgE Mediated Food Allergy as evidenced by one of the following:
Age Restrictions	epinephrine at home. Asthma: 6 years of age and older CIU: 12 years of age and older CRSwNP: 18 years of age and older IgE Mediated Food Allergy: 1 year of age and older
Prescriber Restrictions	Asthma: Prescribed by or in consultation with a pulmonologist or immunologist. CIU: Prescribed by or in consultation with an immunologist. CRSwNP: Prescribed by or in consultation with an allergist or ENT. IgE Mediated Food Allergy: Prescribed by or in consultation with an allergist or immunologist.
Coverage Duration	Asthma - Initial: 6 months. Renewal: 12 months. CRSwNP - Initial: 6 months. Renewal: 12 months. CIU - Initial: 4 months. Renewal: 6 months.

Criteria Details	
Renewal Criteria	 IgE Mediated Food Allergy: Patient demonstrates positive clinical response to therapy (e.g., reduction of type 1 allergic reactions, including anaphylaxis, following accidental exposure to one or more foods). Used in conjunction with food allergen avoidance. Dosing will continue to be based on body weight and pretreatment IgE serum levels. All Other Diagnoses: Documentation of clinically significant improvement in symptoms.
Effective Date	9/1/2024
P&T Approval Date	7/9/2024
P&T Revision Date	7/9/2024

Paliperidone (INVEGA HAFYERA)

Products Affected

• INVEGA HAFYERA

	Criteria Details
Required Medical Information	Clinical documentation of a diagnosis of schizophrenia AND trial and failure (defined by at least 6 months of treatment) of Invega Trinza OR Invega Sustenna AND clinical need or concern for adherence which could be improved upon with twice yearly dosing.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Pancrelipase (CREON) (PANCREAZE)

Products Affected

PANCREAZECREON

	Criteria Details
Required Medical Information	Confirmed diagnosis of cystic fibrosis OR history of pancreatectomy OR diagnosis of exocrine pancreatic cancer OR diagnosis of chronic pancreatitis confirmed by imaging OR confirmed diagnosis of pancreatic insufficiency confirmed with one of the following methods: • Steatorrhea with fecal fat determination OR • Measurement of fecal elastase OR Secretin or CCK pancreatic function testing OR Two of the following CFTR mutations (G542X, W1282X, R553X, 621+1G>T, 1717-1G>A, 3120+1G>A, R1162X, 3659delC, 1898+1G>A, 2184delA, 711+1G>T, F508del, I507del, G551D, N1303K, R560T).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy
Effective Date	
P&T Approval Date	
P&T Revision Date	

PCSK9 inhibitors

Products Affected

• PRALUENT • REPATHA

Criteria Details	
Required Medical Information	 Established clinical atherosclerotic cardiovascular disease (ASCVD): Confirmed diagnosis of atherosclerotic cardiovascular disease (ASCVD). Documentation of a current LDL greater than or equal to 55 mg/dl. Documentation that: Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins AND Is receiving ezetimibe or has a documented intolerance to ezetimibe. Primary or familial hyperlipidemia: Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL. Documentation of current LDL greater than 100 mg/dL. Documentation that:
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Documented positive clinical response to therapy (significant decrease in lipid levels).

Criteria Details	
Effective Date	11/1/2024
P&T Approval Date	1/11/2022
P&T Revision Date	01/11/2022, 09/10/2024

Peanut Powder (PALFORZIA)

Products Affected

- PALFORZIA (12 MG DAILY DOSE)
- PALFORZIA (120 MG DAILY DOSE)
- PALFORZIA (160 MG DAILY DOSE)
- PALFORZIA (20 MG DAILY DOSE)
- PALFORZIA (200 MG DAILY DOSE)
- PALFORZIA (240 MG DAILY DOSE)
- PALFORZIA (3 MG DAILY DOSE)

- PALFORZIA (300 MG MAINTENANCE)
- PALFORZIA (300 MG TITRATION)
- PALFORZIA (40 MG DAILY DOSE)
- PALFORZIA (6 MG DAILY DOSE)
- PALFORZIA (80 MG DAILY DOSE)

PALFORZIA INITIAL ESCALATION

	Criteria Details
Required Medical Information	Confirmed positive skin test or peanut-specific serum IgE greater than 0.35 kUA/L Concurrent prescription with injectable epinephrine medical justification supports necessity for oral immunotherapy despite peanut avoidance.
Age Restrictions	Patient must be between 4 and 17 at therapy initiation
Prescriber Restrictions	Prescribed by allergist or immunologist enrolled in Palforzia REMS program
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Currently receiving medication byway of previously approved SHP authorization or documents showing Initial approval criteria was/has been met. For patients who required use of injectable epinephrine while on Palforzia, must have medical justification that supports continued need for Palforzia. If greater than 18 years old, must have medical justification that supports continued need for oral immunotherapy despite peanut avoidance and documentation that Initial dose escalation happened between age 4 and 17.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Phosphodiesterase Type 5 (PDE5) inhibitors

Products Affected

- Sildenafil 20mg tablet
- Sildenafil oral solution

Tadalafil 20mg tablet

Criteria Details	
Required Medical Information	 Pulmonary Arterial Hypertension Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension)
Age Restrictions	Solution only One of the following: • Pediatric member age 10 or under • Documentation inability of the member to use the preferred tablet formulation
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist.
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	9/1/2025
P&T Approval Date	7/13/2021
P&T Revision Date	7/8/2025, 3/12/2024, 7/13/2021

Pitolisant (WAKIX)

Products Affected

• WAKIX 4.45MG TAB

WAKIX 17.8NG TAB

Cuitavia Dataila	
Required Medical Information	 Criteria Details Confirmation of diagnosis of narcolepsy based on polysomnography AND a multiple sleep latency test. Documentation that CYP2D6 testing has been done, and the dosing will be adjusted if the patient is a poor metabolizer. For Excessive Daytime Sleepiness (EDS) the following is required: Documentation of fatigue severity using a validated measure (e.g. Epworth score, Brief Fatigue Inventory, or other validated measure Trial and failure or contraindication to ALL the following: Modafinil (at least 200mg dose) AND armodafinil Mixed amphetamine salts, methylphenidate or dexmethylphenidate, AND dextroamphetamine Sunosi (solriamfetol) A sodium oxybate product For Cataplexy the following is required:
Age Restrictions	18 years of age and older
Exclusions	 Severe renal or hepatic impairment Pregnant or actively trying to conceive
Prescriber Restrictions	Prescribed by Sleep Specialist or Neurologist
Coverage Duration	Initial: 3 months. Renewal: 6 months
Renewal Criteria	Documentation of positive clinical response to therapy.

Criteria Details	
Effective Date	03/01/2025
P&T Approval Date	01/14/2025
P&T Revision Date	01/14/2025

Pramlintide Acetate (SYMLIN)

Products Affected

SYMLINPEN 60

SYMLINPEN 120

	Criteria Details		
Required Medical Information	Coverage is provided for the use of pramlintide as an adjunct treatment in type 1 and type 2 diabetic patients 18 or older who use mealtime insulin therapy AND who meet all of the following criteria: • Are currently on mealtime insulin. • Have an HbA1c less than or equal to 9%. • Are monitoring blood glucose levels regularly AND reliably (3 or more times per day). • Are capable of monitoring blood glucose levels pre- AND postmeals AND at bedtime. • Have failed to achieve adequate control of blood glucose levels despite individualized management of their insulin therapy. Are receiving ongoing care under the guidance of a health care provider skilled in use of insulin AND supported by the services of a diabetes educator.		
Age Restrictions	Patient must be 18 years or older.		
Prescriber Restrictions			
Coverage Duration	Initial: 12 months. Renewal: 12 months.		
Renewal Criteria	Documentation of positive clinical response to therapy.		
Effective Date			
P&T Approval Date			
P&T Revision Date			

Pretomanid

Products Affected

PRETOMANID TAB

Criteria Details	
Required Medical Information	 Pulmonary tuberculosis Evidence of extensively drug-resistant active pulmonary tuberculosis (XDR-TB) caused by mycobacterium tuberculosis. XDR-TB is defined as TB that is resistant to rifampicin and isoniazid, at least one fluoroquinolone (levofloxacin or moxifloxacin) and a second-line injectable (amikacin, capreomycin, and kanamycin) OR Isoniazid, rifampin a fluoroquinolone AND bedaquiline or linezolid. Pretomanid is prescribed as part of a guideline recommended multidrug treatment regimen.
Age Restrictions	Patient must be 14 years or older.
Exclusion Criteria	 Use outside of recognized treatment guidelines. Medication is being received through a county clinic with a state funded TB program.
Prescriber Restrictions	Infectious Disease
Coverage Duration	Pulmonary tuberculosis: 24 weeks
Renewal Criteria	
Effective Date	05/01/2025
P&T Approval Date	03/11/2025
P&T Revision Date	03/11/2025

Priftin (rifapentine)

Products Affected

• Priftin

	Criteria Details
Required Medical Information	Latent tuberculosis: • Used in combination with isoniazid Active tuberculosis: • Priftin will be used as part of multi-drug regimen
Age Restrictions	 Age ≥ 2 years old with latent TB Age ≥ 12 years old with active TB
Exclusion Criteria	 Medication is being received through a county clinic with a state funded TB program.
Prescriber Restrictions	 Latent TB: Not limited by specialty Active TB: Infectious disease specialist required for multidrug resistant cases only
Coverage Duration	Latent TB: 3 months. Active TB: 6 months.
Renewal Criteria	N/A
Effective Date	5/1/2025
P&T Approval Date	3/12/2024
P&T Revision Date	3/11/2025, 3/12/2024

Prostacyclin Agonists

Products Affected

- Orenitram tablets
- Remodulin

- Treprostinil
- Tyvaso

	Criteria Details
Required Medical Information	 Pulmonary Arterial Hypertension Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension) Clinical documentation of WHO functional class III or IV Evidence of an unfavorable response or intolerance to all of the following: Phosphodiesterase Type 5 inhibitor (sildenafil, tadalafil) Endothelin Receptor Antagonist (ambrisentan, bosentan) Combination therapy with a Phosphodiesterase Type 5 inhibitor + Endothelin Receptor Antagonist (ambrisentan + tadalafil) The requested medication will be an add on to an already established first line agent or agents (sildenafil, tadalafil, ambrisentan, bosentan)
Age Restrictions	Age 16 or older
Prescriber Restrictions	Pulmonary Arterial Hypertension: Cardiologist or Pulmonologist
Coverage Duration	 Pulmonary Arterial Hypertension: Initial: 6 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	9/1/2025
P&T Approval Date	7/13/2021
P&T Revision Date	7/8/2025, 7/13/2021

Ranolazine (RANEXA)

Products Affected

• RANOLAZINE ER

	Criteria Details
Required Medical Information	Diagnosis of chronic angina not controlled with other antianginal therapy. May be used with beta-blockers, nitrates, calcium channel blockers, anti-platelet therapy, lipid-lowering therapy, ACE inhibitors, and angiotensin receptor blockers.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Resmetirom (REZDIFFRA)

Products Affected

Rezdiffra 60MG/80MG/100MG TAB

	Criteria Details
	Diagnosis of metabolic dysfunction-associated steatohepatitis (MASH), formulary known as nonalcoholic steatohepatitis (NASH)
	Patient does not have cirrhosis (e.g. decompensated cirrhosis)
	 Submission of medical records (e.g. chart notes) showing diagnosis has been confirmed by one of the following: FibroScan-aspartate aminotransferase (FAST) MRI-aspartate aminotransferase (MAST) Liver biopsy
Required Medical Information	 Submission of medical records (e.g. chart notes) showing disease is fibrosis stage F2 or F3 as confirmed by one of the following: FibroScan Fibrosis-4 index (FIB-4)
	 Magnetic resonance Elastography (MRE) Presence of greater than or equal to 3 metabolic risk factors (e.g., Type 2 diabetes, hypertension, obesity)
	 Submission of medical records (e.g. chart notes) confirming drug is used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by Gastroenterologist; Hepatologist
Coverage Duration	Initial: 6 months. Renewal: 12 months
Renewal Criteria	Patient demonstrates positive response to therapy (e.g., MASH resolution, fibrosis stage improvement, etc.) AND Submission of medical

Criteria Details	
	records (e.g., chart notes) confirming drug will continue to be used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program)
Effective Date	7/1/2024
P&T Approval Date	5/14/2024
P&T Revision Date	

Ribavirin (VIRAZOLE)

Products Affected

RIBAVIRIN INHALATION

	Critorio Dotoilo
Required Medical Information	Criteria Details Respiratory Syncytial Virus (RSV) Infection: Chart notes / written medical summary documenting diagnosis of RSV.
Age Restrictions	
Prescriber Restrictions	Request is initiated by an infectious disease specialist.
Coverage Duration	Initial: 3 months.
Renewal Criteria	
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ribociclib Succinate (KISQALI)

Products Affected

KISQALI

	Criteria Details
Required Medical Information	Breast Cancer in Women: Has recurrent or metastatic disease AND has hormone receptor positive (HR+) AND has human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND is either postmenopausal OR is pre/perimenopausal AND is receiving gonadotropin-releasing hormone agonist OR has had bilateral oophorectomy or ovarian irradiation AND medication will be used in combination with anastrozole, exemestane, or letrozole OR will be used in combination with fulvestrant. Breast Cancer in Men: Has recurrent or metastatic disease AND has hormone receptor positive (HR+) AND has human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND is receiving
	gonadotropin-releasing hormone analog AND medication will be used in combination with anastrozole, exemestane, or letrozole OR will be used in combination with fulvestrant.
Age Restrictions	At least 18 years of age
Prescriber Restrictions	Prescribed by or in consultations with an oncologist
Coverage Duration	Initial: 3 months. Renewal: 3 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Rifaximin (XIFAXAN)

Products Affected

XIFAXAN

	Criteria Details	
	IBS-D: Diagnosis of IBS with diarrhea	
Required Medical Information	Hepatic Encephalopathy: must have one of the following: used as add-on therapy to lactulose AND unable to achieve an optimal clinical response with lactulose monotherapy OR a history of contraindication or intolerance to lactulose.	
Age Restrictions		
Prescriber Restrictions		
Coverage Duration	IBS-D Initial: 14 Days. Renewal: 30 Days. HE Initial: 12 months. Renewal: 12 months.	
Renewal Criteria	Documentation of positive clinical response to therapy.	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Rimegepant (NURTEC)

Products Affected

NURTEC ODT

	Criteria Details
Required Medical Information	Acute treatment: Diagnosis of migraine with or without aura AND will be used for the acute treatment of migraine AND patient has fewer than 15 headache days per month AND trial and failure or intolerance to 3 generic triptans or contraindications to all triptans and NSAID combined treatment OR trial and failure or intolerance to NSAID treatment alone if triptans contraindicated OR contraindication to all triptans and NSAIDs AND medication will not be used in combination with another CGRP inhibitor AND if patient has 4 or more headache days per month, then patient must currently be treated with amitriptyline, venlafaxine, divalproex, topiramate, candesartan or a beta-blocker or have a contraindication or intolerance to all of these medications. Prophylaxis treatment: Diagnosis of episodic migraines AND patient has 4 to 18 migraine days per month but no more than 18 headache days per month AND trial and failure of at least 2 months intolerance or contraindication of two of the following sets: 1) amitriptyline or venlafaxine; 2) divalproex or topiramate; 3) one of the following beta blocker: atenolol, propranolol, nadolol, timolol, or metoprolol; 4) candesartan AND not used in combination with another CGRP inhibitor.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, pain specialist, or headache specialist
Coverage Duration	Acute Initial: 3 months; Renewal: 12 months Prophylaxis Initial: 6 months; Renewal 12 months
Renewal Criteria	Documentation of positive clinical response to therapy
Effective Date	

	Criteria Details
P&T Approval Date	
P&T Revision Date	

Risankizumab (SKYRIZI)

Products Affected

SKYRIZI

Prior Authorization Criteria

Criteria Details

Plaque Psoriasis (PsO): Diagnosis of moderate to severe chronic plaque psoriasis with one of the following: 1) greater than or equal to 3% body surface area involvement, 2) severe scalp psoriasis, 3) palmoplantar, facial, or genital involvement **AND** a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids, vitamin D analogs, tazarotene, calcineurin inhibitors, anthralin, coal tar.

Psoriatic Arthritis (PsA): Diagnosis of active PsA with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement.

Required Medical Information

Crohn's disease (CD): Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220 AND trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate.*induction dose is IV, maintenance is subcutaneous.

Ulcerative Colitis (UC): Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g., hemoglobin, erythrocyte sedimentation rate, C-reactive protein), 6) dependent on, or refractory to, corticosteroids. **AND** trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine). *induction dose is IV, maintenance is subcutaneous.

	Criteria Details
Age Restrictions	
Prescriber Restrictions	PsA: Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist. CD/UC: Prescribed by or in consultation with a gastroenterologist PsO: Prescribed by or in consultation with a dermatologist
Coverage Duration	Initial: 6 months. Renewal: 12 months
Renewal Criteria	PsO: Documentation of positive clinical response to therapy as evidenced by ONE of the following: Reduction in the body surface area (BSA) involvement from baseline OR improvement in symptoms (e.g., pruritus, inflammation) from baseline. PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline. CD/UC: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state.
Effective Date	10/1/2024
P&T Approval Date	
P&T Revision Date	7/9/2024

Risdiplam (EVRYSDI)

Products Affected

EVRYSDI SOL 0.75MG/ML

	Criteria Details
Required Medical Information	 Spinal Muscular Atrophy (SMA): Confirmed (via genetic testing) diagnosis of 5q-autosomal recessive SMA (type 1, 2 or 3) Patient is not dependent on invasive ventilation or tracheostomy OR use of non-invasive ventilation beyond uses for sleeping Is not receiving concomitant chronic SMN modifying therapy such as Spinraza Patient has not previously received gene replacement therapy for the treatment of SMA (e.g. Zolgensma)
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a neurologist with expertise in the treatment of SMA
Coverage Duration	Initial: 12 months. Renewal: up to 12 months.
Renewal Criteria	Documentation of clinical improvement from baseline in motor functionality confirmed by standard exams (e.g. BSID-III, CHOP INTEND, HINE-2, RULM test)
Effective Date	09/01/2023
P&T Approval Date	7/11/2023
P&T Revision Date	

Sacubitril/Valsartan (ENTRESTO)

Products Affected

ENTRESO

	Criteria Details
Required Medical Information	The patient has a diagnosis of New York Heart Association class II to IV heart failure AND patient is receiving concomitant therapy with one of the following beta blockers: carvedilol, bisoprolol, sustained-released metoprolol, unless unable to tolerate or contraindicated AND patient will discontinue use of any concomitant *ACE inhibitor or ARB before initiating therapy. *ACE inhibitors must be discontinued at least 36 hours prior to ENTRESTO.
Age Restrictions	
Prescriber Restrictions	Cardiologist or in consultation with a cardiologist
Coverage Duration	Initial: 12 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Sargramostin (LEUKINE)

Products Affected

LEUKINE

Criteria Details	
Required Medical Information	Acute myelogenous leukemia (AML): To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in older adults (greater than or equal to 55 years of age). Bone marrow transplant (allogeneic or autologous): For graft failure or engraftment delay. Myeloid reconstitution after allogeneic bone marrow transplantation: To accelerate myeloid recovery following transplantation in non-Hodgkin lymphoma (NHL), acute lymphoblastic leukemia (ALL), Hodgkin lymphoma. Febrile neutropenia Primary prophylaxis of neutropenia in patients receiving chemotherapy (outside transplant and AML) or who are at high risk for neutropenic fever. Peripheral stem cell transplantation: Mobilization of hematopoietic progenitor cells for leukapheresis AND myeloid reconstitution following autologous peripheral stem cell transplantation. Acute Radiation Syndrome: Treatment of radiation-induced myelosuppression of the bone marrow.
Age Restrictions	
Prescriber Restrictions	Requested by specialist.
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	

	Criteria Details
P&T Approval Date	
P&T Revision Date	

Seladelpar (LIVDELZI)

Products Affected

• Livdelzi Capsules

	Criteria Details
Required Medical Information	 Primary biliary cholangitis: Diagnosis of primary biliary cholangitis (PBC) confirmed by two of the following: Biochemical evidence of cholestasis based on ALP elevation Presence of AMA or other PBC-specific autoantibodies Histology confirmation after biopsy Documentation of at least 12 months of inadequate response to ursodiol No current decompensated cirrhosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hepatologist or gastroenterologist
Coverage Duration	Initial: 6 months. Renewal: up to 12 months.
Renewal Criteria	Documented adherence to medication regimen and clinical benefit
Effective Date	01/01/2025
P&T Approval Date	11/12/2024
P&T Revision Date	11/12/2024

Semaglutide (WEGOVY)

Products Affected

Wegovy

Criteria Details	
	2 22 2 22 2
Required Medical Information	 Wegovy is being used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease and either obesity or overweight Wegovy is being used as adjunct to lifestyle modification (e.g. dietary or caloric restriction, exercise, behavioral support, community-based program) Patient has established cardiovascular disease as evidenced by one of the following: prior MI, prior stroke, peripheral aterial disease (e.g. intermittent claudication with ankle-brachial index <0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease) BMI greater than or equal to 27 kg/m2
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	 Documentation of treatment success (BMI reduction of 5% or more) Documentation of continuation of lifestyle modification program with reduced calorie diet and regular physical activity alongside continuous Wegovy use (80% adherence)
Effective Date	7/1/2024
P&T Approval Date	5/14/2024

	Criteria Details
P&T Revision Date	5/14/2024

Sodium Oxybate

Products Affected

Sodium Oxybate 500mg/mL

	Criteria Details
	 Cataplexy and Narcolepsy Confirmation of diagnosis of narcolepsy and cataplexy based on BOTH: Polysomnography A multiple sleep latency test with cataplexy Trial and failure or contraindication to ALL the following SSRI antidepressant (e.g. fluoxetine) SNRI antidepressant (e.g. venlafaxine or duloxetine) Tricyclic antidepressant (e.g. clomipramine)
Required Medical Information	 Excessive Somnolence due to Narcolepsy Confirmation of diagnosis of narcolepsy based on BOTH: Polysomnography A multiple sleep latency test Baseline documentation of fatigue severity using a validated measure (e.g. Epworth score, Brief Fatigue Inventory, or other validated tool). Trial and failure or contraindication to ALL the following: Modafinil (maximum recommended/tolerated dose) At least 2 stimulant medications (amphetamine, methylphenidate, dextroamphetamine, etc.) Sunosi (maximum recommended/tolerated dose)
Age Restrictions	Age ≥7 years old and weight ≥20kg
Prescriber Restrictions	Sleep specialist, Neurologist, Pulmonologist
Coverage Duration	Initial: 3 months. Renewal: 12 months.

Criteria Details	
Renewal Criteria	 Request for a continued maintenance dose within FDA approved limits based on indication. Documented clinical efficacy and tolerability to therapy compared to baseline (for Epworth Sleepiness scale—improvement of at least 3 points is considered clinically significant).
Effective Date	11/1/2025
P&T Approval Date	9/9/2025
P&T Revision Date	9/9/2025

Solriamfetol (Sunosi)

Products Affected

Sunosi Tablets

Prior Authorization Criteria

Criteria Details

Excessive somnolence secondary to narcolepsy

- Confirmed diagnosis of narcolepsy based on BOTH
 - Polysomnography
 - A multiple sleep latency test
- Baseline documentation of fatigue severity using a validated measure (e.g. Epworth score, Brief Fatigue Inventory, or other validated tool).
- Documentation of recent cardiovascular risk assessment (including blood pressure) with a physician attestation that benefits of therapy outweigh the risk.
- Trial and failure, adverse reaction, or contraindication to the following:
 - Modafinil titrated to maximum recommended/tolerated dose
 - Methylphenidate titrated to maximum recommended/tolerated dose
 - Dextroamphetamine titrated to maximum recommended/tolerated dose

Required Medical Information

Excessive somnolence secondary to Obstructive Sleep Apnea

- The medication is intended for the treatment of residual excessive daytime sleepiness in Obstructive Sleep Apnea in a patient treated with CPAP (AHI <5/hour cannot be achieved)
 - CPAP use has been optimized
 - Compliance using device
- Baseline documentation of fatigue severity using a validated measure (e.g. Epworth score, Brief Fatigue Inventory, or other validated tool).
- Documentation of recent cardiovascular risk assessment (including blood pressure) with a physician attestation that benefits of therapy outweigh the risk.

Ouitouis Dataila	
	 Criteria Details Trial and failure, adverse reaction, or contraindication to preferred formulary options Modafinil titrated to maximum recommended/tolerated dose Methylphenidate titrated to maximum recommended/tolerated dose
Age Restrictions	Age 18 and older
Prescriber Restrictions	Sleep specialist, Neurologist, Pulmonologist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	 Excessive somnolence secondary to narcolepsy Documentation of blood pressure evaluation in the last 3 months. Documentation of clinical benefit and tolerability to therapy compared to baseline (must use same clinical measure used to diagnose EDS or fatigue at baseline [Epworth Sleepiness Scale—improvement of at least 3 points is considered clinically significant]). Excessive somnolence secondary to Obstructive Sleep Apnea Documentation of blood pressure evaluation in the last 3 months. Documentation of adherence to primary OSA treatment (e.g. CPAP). Documentation of clinical benefit and tolerability to therapy compared to baseline (must use same clinical measure used to diagnose EDS or fatigue at baseline [Epworth Sleepiness Scale—improvement of at least 3 points is considered clinically significant]).
Effective Date	11/1/2025
P&T Approval Date	9/9/2025
P&T Revision Date	9/9/2025

Somatropins

Products Affected

- GENOTROPIN
- GENOTROPIN MINIQUICK
- HUMATROPE
- NORDITROPIN FLEXPRO
- NUTROPIN AQ NUSPIN 10
- NUTROPIN AQ NUSPIN 20

- NUTROPIN AQ NUSPIN 5
- OMNITROPE
- SAIZEN
- SAIZENPREP
- SEROSTIM

ZORBTIVE

Criteria Details	
	For children: Growth hormone deficiency in individuals less than 16 years of age or radiographic evidence of non-closure of epiphyseal plates. Appropriate medical work up: Assessment AND evaluation must indicate absolute growth less than 4.5 cm per year without growth hormone. Subnormal growth, greater than or equal to 2 standard deviations below the mean for age.
Required Medical Information	For adults: Biochemical diagnosis of adult growth hormone deficiency by means of a subnormal response to a standard growth hormone stimulation test (peak growth hormone Less than or equal to 5 mcg/L). Confirmatory testing may not be required in patients with congenital/genetic growth hormone deficiency or multiple pituitary hormone deficiencies due to organic diseases.
	Adult-onset: Patients who have adult growth hormone deficiency whether alone or with multiple hormone deficiencies (hypopituitarism) as a result of pituitary disease, hypothalamic disease, surgery, radiation therapy, or trauma. Turners Syndrome in females is an approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.

Renewal Criteria Documentation of positive clinical response to therapy. Effective Date P&T Approval Date P&T Revision Date	Criteria Details	
P&T Approval Date	Renewal Criteria	Documentation of positive clinical response to therapy.
	Effective Date	
P&T Revision Date	P&T Approval Date	
	P&T Revision Date	

Sotatercept (WINREVAIR)

Products Affected

WINREVAIR INJECTION

Criteria Details	
Required Medical Information	 Pulmonary Arterial Hypertension (PAH): Diagnosis of symptomatic PAH (WHO Group 1 PH) confirmed by right heart catheterization. WHO functional class II or III symptoms. On a stable dose of both
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	11/1/2024
P&T Approval Date	9/10/2024
P&T Revision Date	9/10/2024

Sparsentan (FILSPARI)

Products Affected

FILSPARI TAB 200MG

• FILSPARI TAB 400MG

	Criteria Details
Required Medical Information	 Primary immunoglobulin A nephropathy: Urine protein-to-creatine ratio (UPCR) ≥ 1.5 and eGFR ≥ 30 mL/min?1.73 m2 Biopsy-verified primary IgA nephropathy No history of kidney transplant and not currently receiving dialysis Member has failed to achieve a reduction in proteinuria to under 1 gram/day while receiving maximally tolerated doses of an ACE inhibitor or ARB for at least 12 weeks
Age Restrictions	18 or older
Prescriber Restrictions	Nephrologist
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	Improved or stable kidney function compared to baseline OR reduction in proteinuria
Effective Date	09/01/2023
P&T Approval Date	7/11/2023
P&T Revision Date	

Suzetrigine (Journavx)

Products Affected

• Journavx tablets

Covered Uses	 All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design Acute, moderate to severe pain
Required Medical Information and Criteria	 Acute moderate to severe pain Clinical documentation of post-operative use following one of the following: Abdominoplasty Bunionectomy Documentation of one of the following: Diagnosis of opioid use disorder Prescriber has a specific concern for opioid abuse.
Renewal Criteria	Not eligible for renewal, patients will need to meet initial criteria with new surgery to be eligible for a new prescription.
Exclusion Criteria	 Use for more than 14 days. Any use outside of acute post-procedural pain.
Age Restriction	Age 18 or older.
Coverage Duration	 Acute moderate to severe pain: Initial: 14 days Renewal: N/A

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

References

- Chou, Roger et al. Management of Postoperative Pain: A Clinical Practice Guideline From the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. The Journal of Pain, Volume 17, Issue 2, 131 - 157
- Dowel D, Ragan KR, Jones CM, Baldwin GT, Chou R. Center for Disease Control and Prevention. CDC Clinical Practice Guideline for Prescribing Opioids for Pain, United States, 2022. CDC Morbidity and Mortality Weekly Report. Recommendations and Reports; Nov 4, 2022: 71 (3); 1-95. Found at: CDC Clinical Practice Guideline for Prescribing Opioids for Pain — United States, 2022 | MMWR
- Hegmann, Kurt T. MD, MPH; Weiss, Michael S. MD, MPH; Bowden, Kirk PhD; Branco, Fernando MD; DuBrueler, Kimberly PharmD, RPh; Els, Charl MBChB, FCPsych, MMed Psych; Mandel, Steven MD; McKinney, David W. MD, MPH; Miguel, Rafael MD; Mueller, Kathryn L. MD, MPH; Nadig, Robert J. MD, MPH; Schaffer, Michael I. PhD, MS, DABFT, NRCC-TC; Studt, Larry MD; Talmage, James B. MD; Travis, Russell L. MD; Winters, Thomas MD; Thiese, Matthew S. PhD, MSPH; Harris, Jeffrey S. MD, MPH. ACOEM Practice Guidelines: Opioids for Treatment of Acute, Subacute, Chronic, and Postoperative Pain. Journal of Occupational and Environmental Medicine 56(12):p e143-e159, December 2014. | DOI: 10.1097/JOM.000000000000352.
- Herzig SJA, Mosher HJR, Calcaterra SLN, Jena ABA, Nuckols TKL. Improving the Safety of Opioid Use for Acute Noncancer Pain in Hospitalized Adults: A Consensus Statement From the Society of Hospital Medicine. J Hosp Med. 2018 Apr;13(4):263-71. https://pubmed.ncbi.nlm.nih.gov/29624189. doi:10.12788/jhm.2980.
- Hsu, Joseph R. MD*; Mir, Hassan MD†; Wally, Meghan K. MSPH*; Seymour, Rachel B. PhD*; the Orthopaedic Trauma Association Musculoskeletal Pain Task Force. Clinical Practice Guidelines for Pain Management in Acute Musculoskeletal Injury. Journal of Orthopaedic Trauma 33(5):p e158-e182, May 2019. | DOI: 10.1097/BOT.000000000001430
- Journavx (suzetrigine) [Package Insert]. Vertex Pharmaceuticals, Inc; Boston, MA: 2025.
 Accessed at: https://pi.vrtx.com/files/uspi_suzetrigine.pdf.
- Vertex Pharmaceuticals. Evaluation of Efficacy and Safety of VX-548 for Acute Pain After a Bunionectomy. NCT05553366. Updated March 26, 2025. Accessed March 26, 2025. https://www.clinicaltrials.gov/study/NCT05553366.
- Vertex Pharmaceuticals. Evaluation of Efficacy and Safety of VX-548 for Acute Pain After an Abdominoplasty. NCT05558410. Updated August 27, 2024, Accessed March 26, 2025. https://www.clinicaltrials.gov/study/NCT05558410.

Tacrolimus (PROTOPIC)

Products Affected

TACROLIMUS EXTERNAL

	Criteria Details
Required Medical Information	Atopic Dermatitis: Clinically diagnosed moderate-to-severe atopic dermatitis (10 percent BSA, hand, foot or mucous membrane involvement, or functional impairment) AND trial and failure of topical steroids, UVB phototherapy, or reason why they would not be medically appropriate.
	Psoriasis: diagnosis of moderate to severe Psoriasis (10 percent BSA, hand, foot or mucous membrane involvement, or functional impairment) AND trial and failure or contraindication to a high potency topical corticosteroid and/or UVB phototherapy.
Age Restrictions	
Prescriber Restrictions	Psoriasis: prescribed by or in consultation with a dermatologist.
Coverage Duration	Initial: 12 months. Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Tazarotene (AVAGE) (TAZORAC)

Products Affected

TAZAROTENE 0.1% CREAM

	Criteria Details
Required Medical Information	Psoriasis: diagnosis of moderate to severe Psoriasis (10% BSA, hand, foot or mucous membrane involvement AND functional impairment) AND trial and failure of high potency topical corticosteroids or medical reason why they would be inappropriate Other FDA approved indications (i.e., severe acne): trial and failure/contraindication to two formulary alternatives used to treat the approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Terbinafine Hydrochloride (LAMISIL)

Products Affected

TERBINAFINE HCL

	Criteria Details
Required Medical Information	For the treatment of onychomycosis of the toenail or fingernail due to dermatophytes (tinea unguium) AND patient is experiencing pain which limits normal activity (i.e., unable to wear shoes, difficulty walking, etc.) OR patient has diabetes OR patient has peripheral vascular diseas, OR patient is immunocompromised.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initial: 3 months. Renewal: 3 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Tezacaftor/ivacaftor (SYMDEKO)

Products Affected

• SYMDEKO

	Criteria Details
Required Medical Information	Clinical documentation of cystic fibrosis diagnosis with homozygous F508del mutation.
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by a pulmonologist
Coverage Duration	Initial: 3 months Renewal: 6 months
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Ticagrelor

Products Affected

Ticagrelor tablets

Criteria Details	
Required Medical Information	Acute Coronary Syndrome • Member has: ○ Either non-ST-elevation acute coronary syndrome (NSTE-ACS) or ST-elevation myocardial infarction (STEMI) AND ○ Has had percutaneous coronary intervention (PCI) AND ○ Has a contraindication to prasugrel OR • Member has NSTE-ACS and is treated with medical therapy alone (has not had PCI) Minor Ischemic Stroke • Member has had a minor non-cardioembolic ischemic stroke (NIHSS score ≤5) in the immediate past • Did not receive IV alteplase • Has a reason that clopidogrel can't be used
Age Restrictions	18 years of age and older
Prescriber Restrictions	Acute Coronary Syndrome: Cardiologist Minor Look area Condictor and Advanced area
Kestrictions	Minor Ischemic Stroke: Cardiologist, Neurologist Acute Corpora Syndrome:
Coverage Duration	 Acute Coronary Syndrome: Initial: 12 months Renewal: 12 months Minor Ischemic Stroke: Initial: 1 month Renewal: N/A
Renewal Criteria	Acute Coronary Syndrome: Documentation of positive clinical response to therapy and continued need for treatment

Criteria Details	
	Minor ischemic stroke: Renewal not appropriate
Effective Date	9/1/2025
P&T Approval Date	7/11/2023
P&T Revision Date	7/8/2025, 7/11/2023

Tranexamic Acid

Products Affected

Tranexamic Acid 650MG TAB

	Criteria Details
	Hemophilia Diagnosis
	 Intending to use for hemorrhage prophylaxis for tooth extraction(s)
Required Medical	Abnormal Uterine Bleeding
Information	 Currently using or documented trial and failure or contradiction to ALL the following treatments: Cobombined Oral Contraceptive therapy
	 Profestin therapy (oral or LM) or Levonogrestrel IUD NSAID therapy
Age Restrictions	
Prescriber Restrictions	Prescribed by Hematologist, Hemophilia specialist, Dentist, Gynecologist
Coverage Duration	Initial: 12 months. Renewal: 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	7/1/2024
P&T Approval Date	5/14/2024
P&T Revision Date	

Tobramycin Solution

Products Affected

Tobramycin nebulization solution

	Criteria Details	
Required Medical Information	Use must be for cystic fibrosis or any FDA-approved or compendia supported indication.	
Age Restrictions		
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist, pulmonologist, or cystic fibrosis specialist	
Coverage Duration	Cystic Fibrosis: Lifetime Other diagnoses: Initial: 3 months; Renewal: 12 months	
Renewal Criteria	Confirmed diagnosis with clinical evidence supporting chronic use	
Effective Date	01/01/2025	
P&T Approval Date	11/12/2024	
P&T Revision Date	11/12/2024	

Tocilizumab SC (ACTEMRA)

Products Affected

ACTEMRA INJ PF syringe

TYENNE INJ PF syringe

• TYENNE INJ AUTO-Inj

TYENNE IV

ACTEMRA INJ ACTPEN

Prior Authorization Criteria

Criteria Details

Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active rheumatoid arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine AND trial and failure, contraindication, or intolerance to TWO of the following, or attestation demonstrating a trial may be inappropriate: Cimzia (certolizumab pegol), Enbrel (etanercept), adalimumab, Rinvoq (upadacitinib), Simponi (golimumab), Xeljanz/XR (tofacitinib/ER).

Systemic Juvenile Idiopathic Arthritis (SJIA): Diagnosis of active systemic juvenile idiopathic arthritis AND trial and failure, contraindication, or intolerance to ONE of the following: Non-steroidal anti-inflammatory drug (NSAID) (e.g., ibuprofen, naproxen), Systemic glucocorticoid (e.g., prednisone), methotrexate.

Required Medical Information

Polyarticular Juvenile Idiopathic Arthritis (PJIA): Diagnosis of active polyarticular juvenile idiopathic arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, AND trial and failure, contraindication, or intolerance to two of the following, or attestation demonstrating a trial may be inappropriate: Enbrel (etanercept), adalimumab, Xeljanz (tofacitinib).

Giant Cell Arteritis (GCA): Diagnosis of giant cell arteritis **AND** trial and failure, contraindication, or intolerance to a glucocorticoid.

Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD): Diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) as documented by the following: exclusion of other known causes of

	Criteria Details	
	interstitial lung disease (ILD) (e.g., domestic and occupational environmental exposures, connective tissue disease, drug toxicity) AND one of the following: In patients not subjected to surgical lung biopsy, the presence of idiopathic interstitial pneumonia (e.g., fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] AND centrilobular fibrosis) pattern on high-resolution computed tomography (HRCT) revealing SSc-ILD or probable SSc-ILD OR In patients subjected to a lung biopsy, both HRCT AND surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD.	
	Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy: Patient will receive or is receiving chimeric antigen receptor (CAR) T-cell immunotherapy (i.e., Kymriah [tisagenlecleucel], Yescarta [axicabtagene ciloleucel])	
Age Restrictions		
Prescriber Restrictions	RA, SJIA, PJIA, GCA: Prescribed by or in consultation with a rheumatologist. SSc-ILD: Prescribed by or in consultation with a pulmonologist or rheumatologist. CRS: Prescribed by or in consultation with an oncologist or hematologist	
Coverage Duration	RA, SJIA, PJIA, GCA, SSc-ILD: Initial: 6 months; Renewal: 12 months Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy: Initial: 2 months.	
Renewal Criteria	RA, PJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen AND tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline. SJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen AND tender) joint count from baseline, or improvement in clinical features or symptoms (e.g., pain, fever, inflammation, rash, lymphadenopathy, serositis) from baseline.	
	GCA, SSc-ILD: Documentation of positive clinical response to therapy.	
Effective Date	9/1/2024	
P&T Approval Date		
P&T Revision Date	7/9/2024	

Tofacitinib (XELJANZ)

Products Affected

XELJANZ

XELJANZ XR

Prior Authorization Criteria

Criteria Details

Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active rheumatoid arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine AND patient has had an inadequate response or intolerance to one or more TNF inhibitors (e.g., Cimzia, adalimumab, Simponi) AND patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

Psoriatic Arthritis (PsA): Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** patient is will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

Required Medical Information

Ankylosing Spondylitis (AS): Diagnosis of active ankylosing spondylitis AND minimum duration of one month trial and failure, contraindication, or intolerance to TWO non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen) AND Trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) AND patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).

Ulcerative Colitis (UC): Diagnosis of moderately to severely active ulcerative colitis with one of the following: 1) Greater than 6 stools per day, 2) frequent blood in the stools, 3) frequent urgency, 4) presence of ulcers, 5) abnormal lab values (e.g. hemoglobin, ESR, CRP), 6) dependent on, or refractory to, corticosteroids. AND trial and failure, contraindication, or intolerance to one of the following conventional

Criteria Details		
	therapies: 6-mercaptopurine, aminosalicylate [e.g., mesalamine sulfasalazine, azathioprine, Corticosteroids (e.g., prednisone, methylprednisolone) AND trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. adalimumab, Simponi) AND patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g., azathioprine or cyclosporine).	
	Polyarticular Juvenile Idiopathic Arthritis (PJIA): Diagnosis of active polyarticular course juvenile idiopathic arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide AND trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Enbrel, adalimumab) AND patient will not be taking Xeljanz in combination with a potent immunosuppressant (e.g azathioprine or cyclosporine).	
Age Restrictions	Solution only One of the following: Pediatric member age 10 or under Documentation inability of the member to use the preferred tablet formulation	
Prescriber Restrictions	RA, AS, PJIA: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with a dermatologist or rheumatologist UC: Prescribed by or in consultation with a gastroenterologist	
overage Duration	RA, PsA, AS, PJIA: Initial: 6 months. Renewal 12 months Ulcerative Colitis (UC): Initial: 4 months. Renewal 12 months	
Renewal Criteria	RA, PJIA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, or improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline AND will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine). PsA: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the total active (swollen and tender) joint count from baseline, improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline, or reduction in the body surface area (BSA) involvement from baseline AND will not be used in	

Criteria Details	
	AS: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count. AND will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).
	UC: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state AND will not be used in combination with biologic DMARDs or potent immunosuppressants (e.g., azathioprine or cyclosporine).
Effective Date	5/1/2024
P&T Approval Date	3/12/2024
P&T Revision Date	3/12/2024

Ubrogepant (UBRELVY)

Products Affected

UBRELVY

	Criteria Details
Required Medical Information	Diagnosis of migraine with or without aura AND will be used for the acute treatment of migraine AND patient has fewer than 15 headache days per month AND trial and failure or intolerance to 3 generic triptans and NSAID (ibuprofen, naproxen, diclofenac) combined treatment OR trial and failure or intolerance to NSAID treatment alone if triptans contraindicated OR contraindication to all triptans and NSAIDs AND medication will not be used in combination with another CGRP inhibitor AND if patient has 4 or more headache days per month, then patient must currently be treated with amitriptyline, venlafaxine, divalproex, topiramate, candesartan or a beta-blocker or have a contraindication or intolerance to all of these medications.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with one of the following specialists: Neurologist, pain specialist, headache specialist
Coverage Duration	Initial: 3 months Renewal: 12 months
Renewal Criteria	Documentation of positive clinical response to therapy and will not be used for preventative treatment of migraine.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Umbralisib (UKONIQ)

Products Affected

Ukoniq

	Criteria Details
Required Medical Information	Marginal zone lymphoma (MZL): Diagnosis of relapsed or refractory marginal zone lymphoma (MZL) AND must have received a prior therapy that included an anti-CD20 antibody agent Follicular lymphoma (FL): Must have received at least three prior therapies, including both an anti-CD20 antibody AND an alkylating agent *Maximum daily dose of 4 tablets
Age Restrictions	18 years of age AND older
Prescriber Restrictions	Prescribed by oncologist
Coverage Duration	Initial: 3 months. Renewal: up to 12 months.
Renewal Criteria	Documentation of positive clinical response to therapy.
Effective Date	
P&T Approval Date	
P&T Revision Date	

Upadacitinib (RINVOQ)

Products Affected

RINVOQ TABLETS

RINVOQ LQ SOLUTION

Prior Authorization Criteria

Criteria Details

Rheumatoid Arthritis (RA): Diagnosis of moderately to severely active rheumatoid arthritis AND trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine AND trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) AND not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

Psoriatic Arthritis (PsA): Diagnosis of active psoriatic arthritis with one of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, active skin and/or nail involvement **AND** trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) **AND** not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

Required Medical Information

Ankylosing spondylitis (AS): Diagnosis of active ankylosing spondylitis minimum duration of one month trial and failure, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., diclofenac, ibuprofen, indomethacin, meloxicam, naproxen). AND trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. Cimzia, Enbrel, adalimumab, Simponi) AND not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).

Atopic Dermatitis (AD): Diagnosis of moderate to severe atopic dermatitis **AND** one of the following: Involvement of at least 10% body surface area (BSA), SCORing Atopic Dermatitis (SCORAD) index value of at least 25 **AND** a minimum duration of a 30-day trial and failure, contraindication, or intolerance to at least one of the following: Medium

Criteria Details		
or higher potency topical corticosteroid, Pimecrolimus cream, Tacrolimus ointment, or Eucrisa (crisaborole) ointment AND a minimum duration of 12-week trial and failure, contraindication, or intolerance of at least one systemic drug for the treatment of AD (e.g. Dupixent) AND not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).		
Age Restrictions	Crohn's disease (CD): Diagnosis of moderately to severely active Crohn's disease with one of the following: 1) frequent diarrhea and abdominal pain, 2) at least 10% weight loss, 3) complications such as obstruction, fever, abdominal mass, 4) abnormal lab values (e.g. C-reactive protein), CD Activity Index greater than 220 AND trial and failure, contraindication, or intolerance to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroids (e.g., prednisone, methylprednisolone), methotrexate AND trial and failure, contraindication, or intolerance to one or more TNF inhibitors (e.g. adalimumab, Simponi) AND not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine). AD: Age 12 or older	
Prescriber Restrictions	RA, AS: Prescribed by or in consultation with a rheumatologist PsA: Prescribed by or in consultation with one of the following: Dermatologist or Rheumatologist. AD: Prescribed by or in consultation with one of the following: Dermatologist or Allergist/Immunologist. UC: Prescribed by or in consultation with a gastroenterologist	
Coverage Duration	Initial: 6 months. Renewal: 12 months	
Renewal Criteria	RA: Documentation of positive clinical response to therapy as evidenced by one of the following: reduction in total active joint count, improvement in symptoms (e.g., improvement in number of swollen/tender joints, pain, or stiffness). AND Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).	
	PsA: Documentation of positive clinical response to therapy as evidenced by one of the following: Reduction in BSA from baseline, reduction in total active joint count, improvement in symptoms(e.g., improvement in number of swollen/tender joints, pain, or stiffness) AND Rinvoq will not be used in combination with other JAK inhibitors, biologic	

Criteria Details		
	DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).	
	AS: Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following: disease activity (e.g., pain, fatigue, inflammation, stiffness), lab values (erythrocyte sedimentation rate, C-reactive protein level), function, axial status (e.g., lumbar spine motion, chest expansion), or total active (swollen and tender) joint count AND Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).	
	AD: Documentation of a positive clinical response to therapy as evidenced by one of the following: reduction in the body surface area (BSA) involvement from baseline, or reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline AND Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).	
	UC: Documentation of positive clinical response to therapy as evidenced by at least one of the following: improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline, or reversal of high fecal output state AND Rinvoq will not be used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine).	
Effective Date		
P&T Approval Date		
P&T Revision Date		

Ustekinumab

Products Affected

- Selarsdi
- Stegeyma

Yesintek

Prior Authorization Criteria

Criteria Details

All diagnoses

- Initial testing for latent TB and treatment, if necessary, before starting treatment.
- No current active infection at initiation of therapy.
- Risks and benefits documented in cases of chronic or recurrent infection.
- Will NOT be used in combination with another biologic or Otezla

Crohn's Disease (CD)

- Documentation of moderate-to-severe Crohn's Disease
- One of the following:
 - The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR
 - Documented trial and failure of at least 1 of the following: 6mercaptopurine, azathioprine, corticosteroid, methotrexate
- Trial and failure of both infliximab and adalimumab

Plaque Psoriasis (PP)

- Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following:
 - At least 10% of body surface area involved
 - Hand, foot, face, or mucous membrane involvement
- The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of all the following:
 - High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.)

Required Medical Information

Criteria Details

- At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.)
- PUVA or UVB Phototherapy
- Methotrexate
- At least 1 other second line systemic agent such as cyclosporine or acitretin
- Trial and failure of both infliximab and adalimumab

Psoriatic Arthritis (PsA)

- Documentation of psoriatic arthritis based on at least 3 out of 5 of the following:
 - Psoriasis (1 point for personal or family history, 2 points for current)
 - Psoriatic nail dystrophy
 - Negative test result for RF
 - Dactylitis (current of history)
 - Radiological evidence of juxta-articular new bone formation
- The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following:
 - NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND
 - Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine
- Trial and failure of both infliximab and adalimumab

<u>Ulcerative Colitis (UC)</u>

- Documentation of moderate-to-severe ulcerative colitis
- The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following:
 - Mesalamine, sulfasalazine OR
 - Mercaptopurine, azathioprine, OR
 - Corticosteroids (prednisone, methylprednisolone)
- Trial and failure of both infliximab and adalimumab

Age Restrictions	PsO/PsA: 6 and older	
Exclusion Criteria	 Not to be used in combination with other biologics for the same indication. 	

Criteria Details	
Prescriber Restrictions	 Crohn's Disease and Ulcerative Colitis: Gastroenterologist. Plaque Psoriasis: Dermatologist. Psoriatic Arthritis: Dermatologist or Rheumatologist.
Coverage Duration	Initial: 6 months. Renewal: 12 months.
Renewal Criteria	 CD: Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission. PP: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. PsA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. UC: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Effective Date	05/01/2025
P&T Approval Date	03/11/2025
P&T Revision Date	3/11/2025, 7/11/2023, 1/11/2022
References	 Selarsdi [package insert]. Parsippany, NJ: Teva Pharmaceuticals; 2025. Yesintek [package insert]. Cambridge, MA: Biocon Biologics; 2024. Steqeyma [package insert]. Jersey City, NJ: Celltrion USA Inc.; 2024.

Vedolizumab (ENTYVIO)

Products Affected

ENTYVIO

Prior Authorization Criteria

Criteria Details

Crohn's Disease

- Documentation of moderately to severely active Crohn's disease
- One of the following:
 - Frequently diarrhea and abdominal pain
 - At least 10% weight loss
 - Complications such as obstruction, fever abdominal mass
 - Abnormal lab values (e.g., C-reactive protein [CRP])
- Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:
 - o 6-mercaptopurine
 - azathioprine
 - o corticosteroids (e.g., prednisone)
 - methotrexate
 - One of the following:
 - Trial and failure, contraindication, or intolerance to TWO of the following:
 - Cimzia (certolizumab pegol)
 - Humira (adalimumab), Amjevita, Cyltezo, Hyrimoz, or Brand Adalimumab-adaz
 - Ustekinumab
 - Skyrizi (risankizumab-rzaa)
 - For continuation of prior Entyvio therapy, defined as no more than a 45-day gap in therapy

Ulcerative Colitis

- Diagnosis of moderately to severely active ulcerative colitis
- One of the following:
 - Greater than 6 stools per day
 - Frequent blood in the stools
 - Frequent urgency
 - Presence of ulcers

Required Medical Information

	Criteria Details	
	 Abnormal lab values (e.g., hemoglobin, ESR, CRP) Dependent on, or refractory to, corticosteroids Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies: 6-mercaptopurine Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine) Azathioprine Corticosteroids (e.g., prednisone) One of the following: Trial and failure, contraindication, or intolerance to TWO of the following, or attestation demonstrating a trial may be inappropriate*:	
Age Restrictions	more than a 45-day gap in therapy	
Prescriber Restrictions	Prescribed by or in collaboration with a Gastroenterologist	
Coverage Duration	Initial: 14 weeks. Renewal: 12 months.	
Renewal Criteria	Crohn's Disease: Documentation of positive clinical response to therapy as evidenced by at least one of the following: • Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline • Reversal of high fecal output state Ulcerative Colitis: Documentation of positive clinical response to therapy as evidenced by at least one of the following: • Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline • Reversal of high fecal output state	
Effective Date	5/1/2024	

Criteria Details		
P&T Approval Date	3/12/2024	
P&T Revision Date		

Vonoprazan (VOQUEZNA)

Products Affected

VOQUEZNA 10MG

• VOQUEZNA 20MG

	Criteria Details	
	Erosive esophagitis-	
	 Imaging confirmed LA Classification Grade C/D erosive esophagitis AND Documented contraindication, intolerance, or inadequate response to 2 or more PPIs (i.e., lansoprazole, omeprazole, esomeprazole, etc.) at maximum tolerated twice-daily dosing for at least 8 weeks each. 	
Required Medical Information	 H.pylori eradication – Confirmed H. pylori positive infection AND Documented contraindication, intolerance, or inadequate response to standard first-line therapies for H.pylori infection (e.g. PPI (standard or double dose), clarithromycin, amoxicillin (or metronidazole)) AND Documented contraindication, intolerance, or inadequate response to a quadruple bismuth regimen (e.g. standard twice daily dose PPI, bismuth subsalicylate, tetracycline, metronidazole) AND Co-prescribed in combination with antibiotics. 	
Age Restrictions	Must be at least 18 years of age	
Prescriber Restrictions	Prescribed by or in collaboration with a Gastroenterologist or Infectious Disease specialist	
Coverage Duration	Initial healing of erosive esophagitis: 2 months Maintenance of healing of erosive esophagitis: 6 months H. Pylori eradication: 14 days	
Renewal Criteria	Renewals past the above timelines are not allowed	
Effective Date	03/01/2024	
P&T Approval Date	01/09/2024	

	Criteria Details
P&T Revision Date	

Xanomeline and trospium (COBENFY)

Products Affected

COBENFY CAPS

Criteria Details					
Required Medical Information	 Schizophrenia: Confirmed diagnosis of schizophrenia. Used as monotherapy (not used in combination with 1st or 2nd generation antipsychotic therapy). Documentation of prior therapy, intolerance, or contraindication to 2 generic antipsychotics indicated for the treatment of schizophrenia (risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, paliperidone, asenapine, or lurasidone). 				
Age Restrictions	18 years of age and older				
Prescriber Restrictions	Psychiatry				
Coverage Duration	Initial: 12 months. Renewal: 12 months				
Renewal Criteria	Documentation of positive clinical response to therapy.				
Effective Date	03/01/2025				
P&T Approval Date	01/14/2025				
P&T Revision Date	01/14/2025				

Zuranolone (ZURZUVAE)

Products Affected

ZURZUVAE

Criteria Details					
Required Medical Information	Postpartum Depression Physician attestation of moderate to severe postpartum depression (PPD) diagnosis and submission of validated screening tool result(s) (e.g. EPDS, PHQ-9) that will be used to monitor a patient's response to Zurzuvae therapy Physician attestation that patient has not had a major depressive episode prior to third trimester of pregnancy and no later than the first 4 weeks following delivery Patient has tried/failed generic SSRI or SNRI for PPD				
Age Restrictions	18 years of age and older				
Prescriber Restrictions	Prescribed by psychiatrist or OB/GYN				
Coverage Duration	Initial: 3 months. Renewal: N/A				
Renewal Criteria					
Effective Date	7/1/2024				
P&T Approval Date	5/14/2024				
P&T Revision Date					