

Prior Authorization Criteria

InterCommunity Health Network CCO

You can get this letter in other languages, large print, Braille or a format you prefer. You can also ask for an interpreter. This help is free. Call 866-203-3435 or (TTY 800-735-2900 or 711). We accept relay calls.

Puede obtener este documento en otros idiomas, en letra grande, braille o en un formato que usted prefiera.

También puede recibir los servicios de un intérprete. Esta ayuda es gratuita. Llame al servicio de atención al cliente 866-203-3435 o (TTY 800-735-2900 o 711). Aceptamos todas las llamadas de retransmisión.

PLEASE READ: This document contains information about the criteria for coverage of this plan

Updated on 09/01/2025. For more recent information or other questions, please contact Pharmacy Services at **541-768-7863** or toll free at **866-203-3435 (TTY 800-735-2900 or 711)** or visit **samhealthplans.org**. Pharmacy Services is available Monday through Friday, from 8 a.m. to 5 p.m.

Abatacept (Orencia)

Products Affected

- Orenzia PFS
- Orenzia Clickjet

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Juvenile Idiopathic Arthritis○ Psoriatic Arthritis○ Rheumatoid Arthritis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none">• Initial testing for latent TB and treatment, if necessary, before starting treatment.• No current active infection at initiation of therapy.• Risks and benefits documented in cases of chronic or recurrent infection.• Will NOT be used in combination with another biologic or Otezla <p><u>Juvenile Idiopathic Arthritis (JIA)</u></p> <ul style="list-style-type: none">• Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features• The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of the following:<ul style="list-style-type: none">○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated○ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids• Documented intolerance or contraindication to DMARDs OR DMARD will be continued• Trial and failure of both infliximab and adalimumab <p><u>Psoriatic Arthritis (PsA)</u></p> <ul style="list-style-type: none">• Documentation of psoriatic arthritis based on at least 3 out of 5 of the following:<ul style="list-style-type: none">○ Psoriasis (1 point for personal or family history, 2 points for current)○ Psoriatic nail dystrophy○ Negative test result for RF

	<ul style="list-style-type: none"> ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following <ul style="list-style-type: none"> ○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ○ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine • Trial and failure of both infliximab and adalimumab <p><u>Rheumatoid Arthritis (RA)</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • JIA: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • PsA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • RA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	<ul style="list-style-type: none"> • Psoriatic Arthritis: Dermatologist or Rheumatologist • Rheumatoid Arthritis, Juvenile Idiopathic Arthritis: Rheumatologist
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	09/01/2023
P&T Approval Date:	07/11/2023
P&T Revision Date:	07/11/2023

Acne Combo Products

Products Affected

- Clindamycin/Benzoyl Peroxide 1-5% Gel
- Erythromycin/Benzoyl Peroxide 3-5% Gel

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Acne vulgaris
Required Medical Information and Criteria	<p><u>Acne Vulgaris</u></p> <ul style="list-style-type: none"> • Documentation of trial and failure, intolerance, or contraindication to clindamycin/benzoyl peroxide 1.2-5% gel
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	
Coverage Duration	<ul style="list-style-type: none"> • Initial: 12 months • Renewal: 12 months

Effective Date:	02/01/2023
P&T Approval Date:	01/10/2023
P&T Revision Date:	01/10/2023

Adalimumab

Products Affected

- Adalimumab-adaz 40mg/0.4mL, 80mg/0.8mL (auto-injector and prefilled syringe)
- Adalimuimab-fkjp 40mg/0.8mL, 20 mg/0.4mL, (auto-injector and prefilled syringe)
- Hadlima 40mg/0.4mL, 40mg/0.8mL (auto-injector and prefilled syringe)
- Yusimry 40mg/0.8mL
- Simlandi 40mg/0.4mL

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Crohn's Disease ○ Hidradenitis Suppurativa ○ Juvenile Idiopathic Arthritis ○ Plaque Psoriasis ○ Psoriatic Arthritis ○ Rheumatoid Arthritis ○ Ulcerative Colitis ○ Uveitis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla. <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR

	<ul style="list-style-type: none"> ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program <p><u>Crohn's Disease (CD)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe Crohn's Disease • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: 6-mercaptopurine, azathioprine, corticosteroid, methotrexate. <p><u>Hidradenitis Suppurativa (HS)</u></p> <ul style="list-style-type: none"> • Documentation of a diagnosis of moderate to severe hidradenitis suppurativa (Hurley Stage II or Hurley Stage III) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy (e.g. oral antibiotics). <p><u>Juvenile Idiopathic Arthritis (JIA)</u></p> <ul style="list-style-type: none"> • Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids • Documented intolerance or contraindication to DMARDs OR DMARD will be continued with adalimumab.
--	--

Plaque Psoriasis (PP):

- Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following:
 - At least 10% of body surface area involved
 - Hand, foot, face, or mucous membrane involvement
- One of the following:
 - The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR
 - Documented trial and failure of all the following:
 - High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.)
 - At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.)
 - PUVA or UVB Phototherapy
 - Methotrexate
 - At least 1 other second line systemic agent such as cyclosporine or acitretin.

Psoriatic Arthritis (PsA):

- Documentation of psoriatic arthritis based on at least 3 out of 5 of the following:
 - Psoriasis (1 point for personal or family history, 2 points for current)
 - Psoriatic nail dystrophy
 - Negative test result for RF
 - Dactylitis (current or history)
 - Radiological evidence of juxta-articular new bone formation
- One of the following:
 - The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR
 - Documented trial and failure of conventional therapy with both of the following:
 - NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND
 - Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine.

	<p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine <p><u>Ulcerative Colitis (UC):</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: mesalamine, sulfasalazine, mercaptopurine, azathioprine, or corticosteroids (prednisone, methylprednisolone). <p><u>Uveitis:</u></p> <ul style="list-style-type: none"> • Documentation of non-infectious, intermediate-, posterior- or pan-uveitis • Documented trial and failure of all the following: <ul style="list-style-type: none"> ○ Topical glucocorticoids for at least 1 month OR periocular steroid injection, and ○ Oral corticosteroids, and ○ At least one of the following: mycophenolate, tacrolimus, cyclosporine, azathioprine, or methotrexate
<p>Renewal Criteria</p>	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Crohn's Disease: Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission. • Hidradenitis Suppurativa: Evidence of a reduction of 25% or more of the total abscess and inflammatory nodule count and no increase in abscesses and draining fistulas.

	<ul style="list-style-type: none"> • Juvenile Idiopathic Arthritis: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Ulcerative Colitis: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity. • Uveitis: Evidence that that disease activity has been controlled, such as a lack of inflammation, no new inflammatory vascular lesions, no vitreous haze or decreases in visual acuity.
Prescriber Restriction	<ul style="list-style-type: none"> • Crohn's Disease and Ulcerative Colitis: Gastroenterologist. • Hidradenitis Suppurativa and Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Rheumatoid Arthritis, Juvenile Idiopathic Arthritis: Rheumatologist. • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Uveitis: Ophthalmologist or Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/01/2023
P&T Approval Date:	07/11/2023
P&T Revision Date:	07/11/2023

Alpelisib (Vijoice)

Products Affected

- Vijoice tab

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ PIK3CA-related overgrowth spectrum (PROS)
Required Medical Information and Criteria	<u>PIK3CA-related overgrowth spectrum (PROS)</u> <ul style="list-style-type: none">• Confirmed diagnosis of PROS• At least one severe clinical manifestation of PROS• A PIK3CA mutation that is confirmed by genetic testing.
Renewal Criteria	<ul style="list-style-type: none">• Documentation of a reduction in volume from baseline in at least one lesion AND• Improvement in at least one symptom of PROS from baseline
Exclusion Criteria	
Age Restriction	<ul style="list-style-type: none">• Two years of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Prescribed by or in consultation with a provider who specializes in treatment of genetic disorders.
Coverage Duration	<ul style="list-style-type: none">• Initial: 24 weeks• Renewal: 6 months

Effective Date:	10/01/2022
P&T Approval Date:	09/13/2022
P&T Revision Date:	09/13/2022

Anakinra (Kineret)

Products Affected

- Kineret 100MG/0.67ML

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Rheumatoid Arthritis
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla <p><u>Rheumatoid Arthritis (RA)</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	<ul style="list-style-type: none"> • Rheumatoid Arthritis: Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • Rheumatoid Arthritis <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	09/01/2023
P&T Approval Date:	07/11/2023

P&T Revision Date:	07/11/2023
-----------------------	------------

Apremilast (Otezla)

Products Affected

- Otezla tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Plaque Psoriasis ○ Psoriatic Arthritis
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic <p><u>Plaque Psoriasis</u></p> <ul style="list-style-type: none"> • Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face or mucous membrane involvement • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of all the following: <ul style="list-style-type: none"> ▪ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ▪ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ▪ PUVA or UVB Phototherapy ▪ Methotrexate ▪ At least 1 other second line systemic agent such as cyclosporine or acitretin. • Trial and failure of both infliximab and adalimumab. <p><u>Psoriatic Arthritis</u></p>

	<ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of all the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine • Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	<ul style="list-style-type: none"> • Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	09/01/2023
P&T Approval Date:	07/11/2023
P&T Revision Date:	07/11/2023, 01/11/2022

Aprepitant (Emend)

Products Affected

- Aprepitant Capsule

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Prevention of nausea and vomiting associated with initial and repeat courses of highly emetogenic chemotherapy ○ Prevention of nausea and vomiting associated with initial and repeat courses of moderately emetogenic chemotherapy
Required Medical Information and Criteria	<p><u>Prevention of chemotherapy induced nausea and vomiting</u></p> <ul style="list-style-type: none"> • Documentation of patient receiving treatment with a moderate to highly emetogenic chemotherapy agent. • Documentation patient is receiving concurrent treatment with all the following: <ul style="list-style-type: none"> ○ IV or oral ondansetron, granisetron or palonosetron ○ Dexamethasone
Renewal Criteria	<ul style="list-style-type: none"> • Positive clinical response to therapy
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	
Coverage Duration	<ul style="list-style-type: none"> • Initial: 12 months • Renewal: 12 months

Effective Date:	10/01/2021
P&T Approval Date:	09/14/2021
P&T Revision Date:	09/14/2021

Asthma Triple Combination Inhaler Step Therapy

Products Affected

- Trelegy
- Breztri

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of at least 4 weeks of 2 of the following<ul style="list-style-type: none">○ A Long-Acting Beta Agonist (LABA)○ An Inhaled Corticosteroid (ICS)○ A Long-Acting Muscarinic Antagonist (LAMA)
------------------------------	---

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	5/14/2024

Atovaquone-proguanil (MALARONE)

Products Affected

- Atovaquone-proguanil 62.5mg-25mg tablet
- Atovaquone-proguanil 250mg-100mg tablet

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Prevention of malaria infection○ Treatment of malaria infection
Required Medical Information and Criteria	<p>Prevention of malaria infection:</p> <ul style="list-style-type: none">• Travel to a location where CDC recommends the use of atovaquone-proguanil• Clinical contraindication to doxycycline OR doxycycline is not recommended by CDC for the travel location. <p>Treatment of malaria infection:</p> <ul style="list-style-type: none">• Recommended by the CDC
Renewal Criteria	N/A
Age Restriction	Refer to FDA label
Coverage Duration	<ul style="list-style-type: none">• Prevention of malaria infection:<ul style="list-style-type: none">○ Initial: 3○ Renewal: N/A• Treatment of malaria infection:<ul style="list-style-type: none">○ Initial: 1○ Renewal: N/A

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024

Baricitinib (OLUMIANT)

Products Affected

- Olumiant tablet

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Rheumatoid Arthritis (RA)
Required Medical Information and Criteria	<p>All diagnoses:</p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla <p>Rheumatoid Arthritis (RA):</p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab • Trial and failure of Actemra, Cimzia, Kineret, Orencia, and rituximab
Renewal Criteria	<ul style="list-style-type: none"> • Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Exclusion Criteria	
Prescriber Restriction	<ul style="list-style-type: none"> • Rheumatoid Arthritis: Rheumatologist
Coverage Duration	<ul style="list-style-type: none"> • Rheumatoid Arthritis <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2023
-----------------	----------

P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023; 1/11/2022

Bedaquiline (Sirturo)

Products Affected

- Sirturo Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Pulmonary Tuberculosis
Required Medical Information and Criteria	<p><u>Pulmonary Tuberculosis</u></p> <ul style="list-style-type: none">• Evidence of active pulmonary tuberculosis caused by mycobacterium tuberculosis that is resistant to at least rifampin and isoniazid• The member weighs at least 15kg• Sirturo is prescribed as part of a guideline recommended multi-drug treatment regimen.
Renewal Criteria	<ul style="list-style-type: none">• Renewal not appropriate
Exclusions	<ul style="list-style-type: none">• Medication is being received through a county clinic with a state funded TB program.
Age Restriction	<ul style="list-style-type: none">• 5 years of age and older.
Prescriber Restriction	<ul style="list-style-type: none">• Pulmonary Tuberculosis: Infectious Disease
Coverage Duration	<ul style="list-style-type: none">• Pulmonary Tuberculosis:<ul style="list-style-type: none">○ Initial: 24 weeks○ Renewal: N/A

Effective Date:	5/1/2025
P&T Approval Date:	3/11/2025
P&T Revision Date:	3/11/2025

References

- Sirturo [package insert]. Horsham, PA: Janssen Products, LP; 2024.

Belumosudil (Rezurock)

Products Affected

- Rezurock tablet

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Chronic graft-versus-host disease
Required Medical Information and Criteria	<ul style="list-style-type: none">• Chronic graft-versus-host disease• Trial and failure of at least two prior lines of systemic therapy for cGVHD• Not currently taking Imbruvica (ibrutinib)
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Exclusion Criteria	
Age Restriction	<ul style="list-style-type: none">• 12 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• cGVHD: oncologist or transplant specialist
Coverage Duration	<ul style="list-style-type: none">• cGVHD:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 6 months

Effective Date:	4/1/2022
P&T Approval Date:	3/8/2022
P&T Revision Date:	3/8/2022

Bempedoic acid (Nexletol/Nexlizet)

Products Affected

- Nexletol tablet
- Nexlizet tablet

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Clinical ASCVD ○ Primary or familial hyperlipidemia
Required Medical Information and Criteria	<p>Clinical ASCVD</p> <ul style="list-style-type: none"> • Documentation of very high risk ASCVD as evidenced by either: <ul style="list-style-type: none"> ○ History of multiple major ASCVD events ○ One major ASCVD event AND multiple high-risk conditions • Documentation of a current LDL greater than or equal to 55 mg/dl • Documentation of all the following <ul style="list-style-type: none"> ○ Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins ○ Is receiving ezetimibe or has a documented intolerance to ezetimibe. • Documentation of failure of PCSK9 inhibitor (Repatha or Prluent) <p>Primary or familial hyperlipidemia</p> <ul style="list-style-type: none"> • Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL • Documentation of current LDL greater than 100 mg/dL • Documentation of all the following <ul style="list-style-type: none"> ○ Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins ○ Is receiving ezetimibe or has a documented intolerance to ezetimibe. • Documentation of failure of PCSK9 inhibitor (Repatha or Prluent)
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy (significant decrease in lipid levels).

Exclusion Criteria	
Age Restriction	
Prescriber Restriction	<ul style="list-style-type: none"> • ASCVD & Primary or familial hyperlipidemia: Cardiologist, Endocrinologist, or lipid specialist
Coverage Duration	ASCVD & Primary or familial hyperlipidemia: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024

Bimekizumab (Bimzelx)

Products Affected

- Bimzelx 160mg/mL PFS
- Bimzelx 320mg/2mL PFS
- Bimzelx 160mg/mL auto-injector
- Bimzelx 320mg/2mL auto-injector

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Plaque Psoriasis ○ Psoriatic Arthritis ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Hidradenitis Suppurativa
Required Medical Information and Criteria	<p><u>Plaque Psoriasis</u></p> <ul style="list-style-type: none"> • Defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) • One or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face, or mucous membrane involvement ○ The patient on a current biologic product and experiencing intolerable side effects • The patient tried and failed or have contraindications to ALL of the following <ul style="list-style-type: none"> ○ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ○ At least one other topical agent: calcipotriene, tazarotene, anthralin, tar, etc. ○ PUVA or UVB Phototherapy ○ Methotrexate ○ At least 1 other second line systemic agent such as cyclosporine or acitretin • Trial and failure of both infliximab and adalimumab • Trial and failure of Cosentyx and ustekinumab <p><u>Psoriatic Arthritis</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF

	<ul style="list-style-type: none"> ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ○ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine • Trial and failure of both infliximab and adalimumab • Trial and failure of Cosentyx, ustekinumab, and Taltz <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ○ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ○ Physical therapy/exercise program • Trial and failure of both infliximab and adalimumab • Trial and failure of Cosentyx, and Taltz <p><u>Hidradenitis Suppurativa</u></p> <ul style="list-style-type: none"> • Documentation of one of the following: <ul style="list-style-type: none"> ○ Moderate to severe hidradenitis suppurative (Hurley Stage II or Hurley Stage III) ○ Patient is on a current biologic product and experiencing intolerable side effects. • The patient is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of the following: <ul style="list-style-type: none"> ○ 90-day trial of conventional therapy (e.g. oral antibiotics) • Trial and failure of both infliximab and adalimumab • Trial and failure of both Cosentyx and ustekinumab
--	---

Renewal Criteria	<ul style="list-style-type: none"> • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Hidradenitis Suppurativa: Evidence of a reduction of 25% or more of the total abscess and inflammatory nodule count and no increase in abscesses and draining fistulas.
Exclusion Criteria	<ul style="list-style-type: none"> • Not to be used in combination with other biologics for the same indication
Age Restriction	Refer to FDA label
Prescriber Restriction	<ul style="list-style-type: none"> • Plaque Psoriasis, Hidradenitis Suppurativa: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	5/13/2024
P&T Revision Date:	7/8/2025, 5/13/2024

References
<ul style="list-style-type: none"> • Bimzelx (bimekizumab-bkzx). Prescribing Information. UCB, Inc. Smyrna, GA 2024.

- Gordon KB, et al. Bimekizumab efficacy and safety in moderate to severe plaque psoriasis (BE READY): a multicentre, double-blind, placebo-controlled, randomised withdrawal phase 3 trial. *Lancet*. 2021; 37:475-86.
- Gordon KB, et al. Bimekizumab safety in patients with moderate to severe plaque psoriasis: Pooled results from Phase 2 and Phase 3 randomized clinical trials. *JAMA Dermatol*. 2022;158(7):735-744.
- Gordon KB, et al. Bimekizumab safety in patients with moderate-to-severe plaque psoriasis: pooled data from up to 3 years of treatment in randomized phase III trials. *Br J Dermatol* 2024; 190: 477-485.

Buprenorphine Patch (Butrans)

Products Affected

- Buprenorphine Patch

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Cancer or End-of-life Care ○ Other Chronic Pain
Required Medical Information and Criteria	<p>Cancer or End-of-life Care</p> <ul style="list-style-type: none"> • Patient is being treated for cancer related pain or pain associated with end-of-life • Documented trial and failure of scheduled short-acting opioid therapy • Documented trial and failure of, or contraindication to long-acting morphine sulfate therapy • Documented trial/failure of, or reason why fentanyl is not appropriate. <p>Other Chronic Pain</p> <ul style="list-style-type: none"> • Documented above the line diagnosis, FDA indicated, or guideline supported condition • Documented chronic pain (greater than 3mo) that is severe enough to require around the clock analgesic therapy • Documented trial and failure or contraindication to short-acting opioid therapy • Documented trial and failure of, or contraindication to long-acting morphine sulfate therapy • Documented trial and failure of, or reason why fentanyl is not appropriate.
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> • Cancer or End-of-Life Care: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months • Other Chronic Pain: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	10/1/2021
P&T Approval Date:	9/14/2021
P&T Revision Date:	9/14/2021

Calcitonin Gene Related Peptide (CGRP) Antagonists

Products Affected

- Aimovig (erenumab)
- Ajoovy (fremanezumab)
- Emgality (galcanezumab)

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Migraine prophylaxis ○ Cluster headache (Emgality only)
Required Medical Information and Criteria	<p><u>Migraine Prophylaxis</u></p> <ul style="list-style-type: none"> • Documentation showing member experiences at least 4 migraines per month. • For members with chronic migraine (≥ 15 headache days & 8 migraine episodes per month) Trial and failure of at least 8 weeks of: <ul style="list-style-type: none"> ○ 3 or more prophylactic medications from at least 2 of the following classes: <ul style="list-style-type: none"> ▪ beta blockers ▪ antidepressants ▪ anticonvulsants ○ Botox. • For members without a diagnosis of chronic migraine: • Trial and failure of: <ul style="list-style-type: none"> ○ at least 3 prophylactic medications from 2 or more of the following classes: <ul style="list-style-type: none"> ▪ beta blockers ▪ antidepressants ▪ anticonvulsants • The requested treatment will not be used in combination with another CGRP inhibitor. <p><u>Cluster Headache</u></p> <ul style="list-style-type: none"> • The request is for Emgality 300mg dose • The member does not have any of the following exclusions: <ul style="list-style-type: none"> ○ ECG abnormalities compatible with an acute CV event, ○ history of unstable angina, percutaneous coronary intervention, coronary artery bypass grafting, deep vein

	<p>thrombosis, or pulmonary embolism within the past 6 months,</p> <ul style="list-style-type: none"> ○ history of stroke, intracranial or carotid aneurysm, intracranial hemorrhage, vasospastic angina, or peripheral vascular disease • Trial and failure of at least a 3-month trial of both of the following: <ul style="list-style-type: none"> ○ Verapamil ○ Topiramate • The requested treatment will not be used in combination with another CGRP inhibitor.
Renewal Criteria	<ul style="list-style-type: none"> • Migraine Prophylaxis: <ul style="list-style-type: none"> ○ Reduction in monthly headache days by at least 2 days from pre-CGRP treatment baseline ○ Clinical documented improvement in migraine-related disability • Cluster Headache <ul style="list-style-type: none"> ○ Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> • 18 years of age and older
Prescriber Restriction	<ul style="list-style-type: none"> • Migraine Prophylaxis and Cluster Headache: Prescribed by or in consultation with a neurologist
Coverage Duration	<ul style="list-style-type: none"> • Migraine Prophylaxis: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months • Cluster Headache: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 6 months

Effective Date:	9/1/2025
P&T Approval Date:	11/08/2022
P&T Revision Date:	7/8/2025, 12/1/2022, 1/11/2022

References

- Qaseem A, et al. Prevention of Episodic Migraine Headache Using Pharmacologic Treatment in Outpatient Settings: A Clinical Guideline From the American College of Physicians. *Ann Intern Med.* 2025;178:426-433.
- Ailani J., et al. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. *Headache.* 2021;61:1021-1039.
- Timotheussen Lund NL, et al. Current treatment options for cluster headache: limitations and the unmet need for better and specific treatments—a consensus article. *The Journal of Headache and Pain.* 2023;24:121.

Certolizumab (Cimzia)

Products Affected

- Cimzia PFS
- Cimzia Auto-injector

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA) ○ Crohn's Disease (CD) ○ Juvenile Idiopathic Arthritis (JIA) ○ Plaque Psoriasis (PsO) ○ Psoriatic Arthritis (PsA) ○ Rheumatoid Arthritis (RA)
Required Medical Information and Criteria	<p>All diagnoses</p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment • No current active infection at initiation of therapy • Risks and benefits documented in cases of chronic or recurrent infection • Will NOT be used in combination with another biologic or Otezla <p>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months ○ Signs of active inflammation on MRI ○ Radiological evidence of sacroiliitis OR HLA-B27 positive ○ BASDAI score of ≥ 4 • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR Documented failure of conventional therapy with both of the following <ul style="list-style-type: none"> ○ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated ○ Physical therapy/exercise program • Trial and failure of both infliximab and adalimumab <p>Crohn's Disease (CD)</p>

	<ul style="list-style-type: none"> • Documentation of moderate-to-severe Crohn's Disease • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR Documented trial and failure of at least 1 of the following: 6-mercaptopurine, azathioprine, corticosteroid, methotrexate • Trial and failure of both infliximab and adalimumab <p>Juvenile Idiopathic Arthritis (JIA)</p> <ul style="list-style-type: none"> • Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of the following: <ul style="list-style-type: none"> ○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ○ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids • Documented intolerance or contraindication to DMARDs OR DMARD will be continued • Trial and failure of both infliximab and adalimumab <p>Plaque Psoriasis (PsO)</p> <ul style="list-style-type: none"> • Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) • One or more of the following <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face, or mucous membrane involvement • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of all the following <ul style="list-style-type: none"> ○ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ○ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ○ PUVA or UVB Phototherapy ○ Methotrexate
--	--

	<ul style="list-style-type: none"> ○ At least 1 other second line systemic agent such as cyclosporine or acitretin • Trial and failure of both infliximab and adalimumab <p>Psoriatic Arthritis (PsA)</p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following <ul style="list-style-type: none"> Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following <ul style="list-style-type: none"> ○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated ○ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine • Trial and failure of both infliximab and adalimumab <p>Rheumatoid Arthritis (RA)</p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • AS/SpA: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • CD: Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission.

	<ul style="list-style-type: none"> • JIA: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • PsO: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • PsA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • RA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Exclusion Criteria	
Age Restriction	
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Rheumatologist. • Crohn's Disease: Gastroenterologist • Juvenile Idiopathic Arthritis: Rheumatologist • Plaque Psoriasis: Dermatologist • Psoriatic Arthritis: Dermatologist or Rheumatologist • Rheumatoid Arthritis: Rheumatologist
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

Chloroquine Phosphate

Products Affected

- Chloroquine tablet

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of malaria○ Extraintestinal amebiasis○ Prevention of malarial infection
Required Medical Information and Criteria	<p>Prevention of malaria infection:</p> <ul style="list-style-type: none">• Travel to a location where CDC recommends the use of chloroquine <p>Treatment of malaria</p> <ul style="list-style-type: none">• Confirmed diagnosis of Malaria <p>Extraintestinal amebiasis</p> <ul style="list-style-type: none">• Confirmed diagnosis of amebiasis
Renewal Criteria	N/A
Coverage Duration	<ul style="list-style-type: none">• Treatment of malaria & extraintestinal amebiasis:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: N/A

Effective Date:	11/01/2024
P&T Approval Date:	7/15/2013
P&T Revision Date:	7/15/2013, 9/10/2024

Cinacalcet Hydrochloride

Products Affected

- Cinacalcet tablet

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Treatment of secondary hyperparathyroidism in patients with CKD on dialysis ○ Treatment of hypercalcemia in patients with parathyroid carcinoma. ○ Treatment of severe hypercalcemia in patients with primary hyperparathyroidism
Required Medical Information and Criteria	<p>Secondary hyperparathyroidism in patients with CKD</p> <ul style="list-style-type: none"> • Confirmed diagnosis of secondary hyperparathyroidism in patients with CKD on dialysis. <p>Hypercalcemia in patients with parathyroid carcinoma</p> <ul style="list-style-type: none"> • Confirmed diagnosis of hypercalcemia in patients with parathyroid carcinoma <p>Severe hypercalcemia in patients with primary hyperparathyroidism</p> <ul style="list-style-type: none"> • Confirmed diagnosis of severe hypercalcemia in patients with primary hyperparathyroidism who are unable to undergo parathyroidectomy
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> • 18 years of age and older
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Clobazam (Onfi)

Products Affected

- Clobazam tablets
- Clobazam suspension

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Lennox-Gastaut Syndrome○ Refractory Seizures
Required Medical Information and Criteria	<p><u>Lennox-Gastaut Syndrome</u></p> <ul style="list-style-type: none">• Confirmation of diagnosis• Suspension only: Member under age 10 or unable to use tablets <p><u>Refractory Seizures</u></p> <ul style="list-style-type: none">• Documentation showing appropriate trial of 2 or more tolerated anticonvulsant therapies• Suspension only: Member under age 10 or unable to use tablets
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Age 2 or older• Suspension only: Member under age 10 or unable to use tablets
Prescriber Restriction	<ul style="list-style-type: none">• All Diagnoses: Neurologist
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: Lifetime

Effective Date:	05/01/2025
P&T Approval Date:	07/11/2023
P&T Revision Date:	03/12/2024, 03/11/2025

References
<ul style="list-style-type: none">• Onfi [package insert]. Deerfield, IL: Lundbeck.; 2024.

Clotrimazole Troche Step Therapy

Products Affected

- Clotrimazole Troche

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Formulary nystatin
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Colony-Stimulating Factors (Filgrastim, Pegfilgrastim)

Products Affected

- Nivestym Syringe and Inj.
- Zarxio Syringe
- Udenyca Syringe, Pen and Onbody
- Neulasta Syringe, Onpro
- Ziextenzo Syringe

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Bone Marrow/Stem Cell Transplant Acute Myeloid Leukemia (AML) Induction or Consolidation Therapy Febrile Neutropenia Prophylaxis Treatment of High-Risk Febrile Neutropenia Severe Chronic Neutropenia Acute Radiation Syndrome Human Immunodeficiency Virus (HIV) Related Neutropenia
Required Medical Information and Criteria	<p><u>Bone Marrow/Stem Cell Transplant</u></p> <ul style="list-style-type: none"> One of the following <ul style="list-style-type: none"> Patient has non-myeloid malignancies undergoing myeloablative chemotherapy followed by autologous or allogeneic bone marrow transplant (BMT) Used for mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis Patient has had a peripheral stem cell transplant (PSCT) and has received myeloablative chemotherapy <p><u>Acute Myeloid Leukemia (AML) Induction or Consolidation Therapy</u></p> <ul style="list-style-type: none"> Diagnosis of acute myeloid leukemia (AML) Patient has completed induction or consolidation chemotherapy <p><u>Febrile Neutropenia Prophylaxis</u></p> <ul style="list-style-type: none"> Patient will be receiving prophylaxis for febrile neutropenia (FN) due to one of the following:

	<ul style="list-style-type: none"> ○ Patient is receiving National Cancer Institute's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer. ○ Patient is receiving a dose-dense chemotherapy regimen for which the incidence of FN is unknown. ○ Patient is receiving chemotherapy regimen(s) associated with greater than 20% incidence of FN. ○ Patient is receiving chemotherapy regimen(s) associated with 10-20% incidence of FN and has one or more risk factors associated with chemotherapy induced infection, FN, or neutropenia. ○ Patient is receiving myelosuppressive anticancer drugs associated with neutropenia and has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). <p><u>Treatment of High-Risk Febrile Neutropenia</u></p> <ul style="list-style-type: none"> • Patient has received or is receiving myelosuppressive anticancer drugs associated with neutropenia. • Diagnosis of febrile neutropenia (FN). • Patient is at high risk for infection-associated complications. <p><u>Severe Chronic Neutropenia (SCN)</u></p> <ul style="list-style-type: none"> • For patients with severe chronic neutropenia (SCN) (i.e., congenital, cyclic, and idiopathic neutropenias with chronic absolute neutrophil count [ANC] less than or equal to 500 cells/mm³) <p><u>Acute Radiation Syndrome (ARS)</u></p> <ul style="list-style-type: none"> • Patient was/will be acutely exposed to myelosuppressive doses of radiation cells/mm³) <p><u>Human Immunodeficiency Virus (HIV) Related Neutropenia</u></p> <ul style="list-style-type: none"> • Patient is infected with HIV virus. • ANC less than or equal to 1,000 (cells/mm³).
Renewal Criteria	<ul style="list-style-type: none"> • Renewal criteria
Prescriber Restriction	<ul style="list-style-type: none"> • HIV Related Neutropenia: Hematologist, Oncologist, or Infectious Disease Specialist • All Other Diagnoses: Hematologist or Oncologist
Coverage Duration	<ul style="list-style-type: none"> • Severe Chronic Neutropenia:

	<ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months • Acute Radiation Syndrome: <ul style="list-style-type: none"> ○ Initial: 1 month ○ Renewal: N/A • All other Diagnoses: <ul style="list-style-type: none"> ○ Initial: 3 months or duration of therapy ○ Renewal: 3 months or duration of therapy
--	---

Effective Date:	9/1/2024
P&T Approval Date:	7/9/2024
P&T Revision Date:	7/9/2024

References	
<ul style="list-style-type: none"> • Neulasta Prescribing Information. Amgen Inc. Thousand Oaks, CA. February 2021. • Neupogen Prescribing Information. Amgen Inc. Thousand Oaks, CA. February 2021. 	

Compounds (standard criteria for all compounded medications)

Products Affected

- All compounded medications

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p>All diagnoses</p> <ul style="list-style-type: none"> • The requested medication is being used to treat an above the line diagnosis on the OHA Prioritized list OR the member has an above the line comorbid condition that will be treated indirectly by the requested medication OR the member is under the age of 21. • Each active ingredient in the compounded drug is FDA-approved or national compendia supported for the condition being treated. • The requested amounts are supported by national compendia or at least two peer-reviewed studies for the condition being treated in the requested route of delivery. • If any prescription ingredients require prior authorization and/or step therapy, all drug-specific criteria must be also met. • The patient has tried and failed therapy, or had an intolerance to all FDA-approved commercially available prescription therapeutic alternatives, unless one of the following criteria are met: <ul style="list-style-type: none"> ○ Patient has a contraindication to all commercially available products. ○ No other therapeutic alternatives are commercially available. ○ Prepared strength(s) is/are not commercially available or currently in short supply and the strength is medically necessary. ○ Prepared in a different dosage form for a patient who is unable to take the commercially available formulation (mixing or reconstituting commercially available products based on the manufacturer's instructions or the product's approved labeling does NOT meet this criteria). ○ Patient has an allergy or sensitivity to inactive ingredients (e.g. dyes, preservatives, sugars, etc.) that are found in commercially available products.

Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Conjugated Estrogens (Premarin Tablets)

Products Affected

- Premarin TAB 0.3MG, 0.45MG, 0.625MG, 0.9MG, 1.25MG

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	All FDA indicated or guideline supported diagnoses <ul style="list-style-type: none">• Trial and failure of generic estradiol tablets and patches
Coverage Duration	<ul style="list-style-type: none">• All FDA indications<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	3/1/2024
P&T Approval Date:	1/9/2024
P&T Revision Date:	

Conjugated estrogens and medroxyprogesterone acetate (Prempro/Premphase)

Products Affected

- Prempro
- Premphase

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	All FDA indicated or guideline supported diagnoses <ul style="list-style-type: none">• One of the following:<ul style="list-style-type: none">○ Trial and failure of generic combination products○ Estradiol tablets/patches used in combination with medroxyprogesterone capsules.
Coverage Duration	<ul style="list-style-type: none">• All FDA indications:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	3/1/2024
P&T Approval Date:	1/9/2024
P&T Revision Date:	1/9/2024

Continuous Glucose Monitors (CGM)

Products Affected

- Dexcom G6/G7
- Freestyle Libre 2/3

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Type 1 Diabetes Mellitus ○ Type 2 Diabetes ○ Gestational Diabetes
Required Medical Information and Criteria	<p>Type 1 Diabetes Mellitus</p> <ul style="list-style-type: none"> • One of the following: <ul style="list-style-type: none"> ○ Use of continuous insulin infusion via pump ○ Child or adolescent under the age of 21 ○ Member is pregnant or plans to become pregnant within 6 months ○ Using short or intermediate acting insulin • The criteria listed under “applies to all requests” has been met. <p>Type 2 Diabetes Mellitus OR Gestational Diabetes</p> <ul style="list-style-type: none"> • Using short or intermediate acting insulin • The criteria listed under “applies to all requests” has been met. <p>Applies to all requests</p> <ul style="list-style-type: none"> • Must meet at least one of the following: <ul style="list-style-type: none"> ○ Baseline HbA1c levels greater than or equal to 8% ○ Frequent or severe hypoglycemia ○ Impaired awareness of hypoglycemia (including presence of these conditions prior to initiation of CGM) ○ Diabetes related complications (e.g. peripheral neuropathy, end-organ damage, etc.) • Member has received or will receive diabetes education specific to the use of the CGM device. <p>** If the request is for a Dexcom one of the following criteria must apply:</p> <ul style="list-style-type: none"> • Use of an insulin pump compatible with the requested Dexcom and not Freestyle • Pediatric member under the age of 16 years old • Inability to use preferred Freestyle CGM device
Renewal Criteria	<ul style="list-style-type: none"> • Use of CGM device for at least 50% of the time for a 90-day period by the time of first follow-up visit (within 3-6 months) and

	provider visits within the last 6 months. **Note: two trials per year of CGM are allowed to meet adherence for continuation of coverage.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	6/1/2022
P&T Approval Date:	5/8/2022
P&T Revision Date:	5/8/2022

Cyclosporine Oral

Products Affected

- Cyclosporine capsules
- Cyclosporine Modified capsules
- Cyclosporine solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Solid Organ Transplant○ Rheumatoid Arthritis○ Psoriasis
Required Medical Information and Criteria	<u>All diagnoses</u> <ul style="list-style-type: none">• Documentation of an FDA approved indication
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Liquid only: Member is under age 10 or unable to use tablets/capsules.
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References
<ul style="list-style-type: none">• Gengraf [package insert]. North Chicago, IL: Abbvie.; 2024.• Sandimmune/Neoral [package insert]. East Hanover, NJ: Novartis.; 2020.

Cyclosporine ophthalmic (Restasis)

Products Affected

- Cyclosporine 0.05% emulsion

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Dry eye syndrome or keratoconjunctivitis sicca for EPSDT members
Required Medical Information and Criteria	<p><u>Dry Eye Syndrome or Keratoconjunctivitis Sicca for members under age 21</u></p> <ul style="list-style-type: none"> • Clinically documented trial and failure of both: <ul style="list-style-type: none"> ○ Ocular lubricants in solution form ○ Ocular lubricants in ointment form. • Clinical documentation that topical and systemic contributors to dry eye disease have been discussed and either: <ul style="list-style-type: none"> ○ Eliminated or ○ Contributing agents are medically necessary. <p><u>Dry Eye Syndrome or Keratoconjunctivitis Sicca for members age 21 or older</u></p> <ul style="list-style-type: none"> • Documented clinical evidence of a funded (above the line) comorbid condition for which the following applies: <ul style="list-style-type: none"> ○ Clinical evidence shows that the funded treatments are not working or are contraindicated ○ Treating dry eye syndrome or keratoconjunctivitis sicca would significantly improve the outcome of treating the funded condition. • Clinically documented trial and failure of both: <ul style="list-style-type: none"> ○ Ocular lubricants in solution form. ○ Ocular lubricants in ointment form. • Clinical documentation that topical and systemic contributors to dry eye disease have been discussed and either: <ul style="list-style-type: none"> ○ Eliminated or ○ Contributing agents are medically necessary.
Renewal Criteria	<ul style="list-style-type: none"> • Clinical documentation of efficacy.
Age Restriction	<ul style="list-style-type: none"> • 16 years of age or older.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months

	○ Renewal: 12 months
--	----------------------

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

References	
<ul style="list-style-type: none"> • Dermatologic and Eyes, Ears, Nose, and Throat, and Immunologic Disorders: Presented by Jamie L. McConaha. • https://www.aao.org/education/preferred-practice-pattern/dry-eye-syndrome-ppp-2023 • Restasis [package insert]. North Chicago, IL: Abbvie.; 2024. 	

Deferasirox

Products Affected

- Deferasirox 125MG, 250MG, 500MG tablets

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Chronic iron overload due to blood transfusion Non-transfusion-dependent thalassemia syndromes
Required Medical Information and Criteria	<p>Chronic iron overload due to blood transfusion</p> <ul style="list-style-type: none"> One of the following: <ul style="list-style-type: none"> A creatinine clearance of greater than or equal to 40 mL/minute Serum creatinine less than or equal to 2 times the age-appropriate level Patient has a serum ferritin levels consistently greater than 1,000 mcg/L (as demonstrated with at least two lab values within the previous two months) Patient has had a failure or contraindication to deferoxamine injection <p>Non-transfusion-dependent thalassemia syndromes</p> <ul style="list-style-type: none"> One of the following: <ul style="list-style-type: none"> A creatinine clearance of greater than or equal to 40 mL/minute Serum creatinine less than or equal to 2 times the age-appropriate level Patient has a serum ferritin levels consistently greater than 1,000 mcg/L (as demonstrated with at least two lab values within the previous two months) Patient has had a failure or contraindication to deferoxamine injection
Renewal Criteria	<ul style="list-style-type: none"> Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> Patient is 2 years of age or older
Coverage Duration	<ul style="list-style-type: none"> Both Diagnoses: <ul style="list-style-type: none"> Initial: 12 months Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Desmopressin

Products Affected

- Desmopressin Acetate Nasal Spray & Injections

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Diabetes Insipidus○ Maintenance of hemostasis and control of bleeding in hemophilia A with factor VIII coagulant activity levels greater than 5%○ Mild-to-moderate classic von Willebrand's disease (type 1) with factor VIII coagulant activity greater than 5%.
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none">• Documentation of one of the conditions listed under “covered uses”
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to therapy.
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
<ul style="list-style-type: none">• Stimate [package insert]. King of Prussia, PA: CSL Behring LLC.; 2015.

Deutetrabenazine (Austedo)

Products Affected

- Austedo tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Chorea associated with Huntington's Disease ○ Tardive Dyskinesia
Required Medical Information and Criteria	<p><u>Chorea associated with Huntington's Disease</u></p> <ul style="list-style-type: none"> • Documentation of the degree of chorea present and the impact on functional ability and/or quality of life. • Documentation of mental status, specifically depression and suicidality. <p><u>Tardive Dyskinesia</u></p> <ul style="list-style-type: none"> • Clinical documentation of tardive dyskinesia including: <ul style="list-style-type: none"> ○ At least one month of past or current exposure to a dopamine receptor blocker ○ Dyskinetic or dystonic involuntary movements ○ Exclusion of other causes of abnormal movements • Clear documentation that tardive dyskinesia causes functional impairment • Documentation of the degree of tardive dyskinesia with the AIMS scale as a baseline • One of the following: <ul style="list-style-type: none"> ○ Discontinuation of the medication precipitating TD ○ Documentation that the patient has tried and failed an 8-week trial of at least 2 other agents within the same therapeutic category at a clinically effective and maximally tolerated dose for the patient's primary neuropsychiatric diagnosis ○ Evidence the medications precipitating tardive dyskinesia are medically necessary • Trial and failure of both of the following: <ul style="list-style-type: none"> ○ Clonazepam ○ Amantadine
Renewal Criteria	Huntington's Chorea

	<ul style="list-style-type: none"> • Clinical response such as improvement in chorea, ability to perform ADLs, reduction in falls, or increase in quality of life. AND • Documentation of continued monitoring of mental status specifically for depression and suicidality. <p>Tardive Dyskinesia</p> <ul style="list-style-type: none"> • Follow-up AIMS assessment showing improvement from Baseline AND • Documented improvement in functioning such as ability to perform ADLs, reduction in falls and increase in quality of life.
Age Restriction	<ul style="list-style-type: none"> • Age 18 and older
Prescriber Restriction	<ul style="list-style-type: none"> • Huntington's Chorea: Neurologist • Tardive Dyskinesia: Neurologist or Psychiatrist
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 12 months

Effective Date:	7/1/2023
P&T Approval Date:	5/9/2023
P&T Revision Date:	5/9/2023

References
<ul style="list-style-type: none"> • Austedo [package insert]. Parsippany, NJ: Teva Neuroscience, Inc.; 2025.

Dihydroergotamine Injection

Products Affected

- Dihydroergotamine Inj

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Acute treatment of migraine○ Acute treatment of cluster headaches
Required Medical Information and Criteria	<p><u>Acute treatment of migraine</u></p> <ul style="list-style-type: none">• Diagnosis of migraine with or without aura• Trial and failure of, or contraindication to 5-HT_{1B/1D} agonist (triptans) <p><u>Acute treatment of cluster headaches</u></p> <ul style="list-style-type: none">• Diagnosis of cluster headache• Trial and failure of, or contraindication to 5-HT_{1B/1D} agonist (triptans)
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	7/1/2023
P&T Approval Date:	
P&T Revision Date:	5/9/2023

References
<ul style="list-style-type: none">• Migranal [package insert]. Bridgewater, NJ: Bausch Health US, LLC; 2022.

Dimethyl Fumarate

Products Affected

- Dimethyl Fumarate DR Capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Multiple sclerosis
Required Medical Information and Criteria	<u>Multiple Sclerosis</u> <ul style="list-style-type: none">• Documented diagnosis of a relapsing form of multiple sclerosis.
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple sclerosis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	2/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References

- Tecfidera [package insert]. Cambridge, MA: Biogen INC.; 2024.

Disposable Insulin Pump (Omnipod)

Products Affected

- Omnipod 5
- Omnipod Dash

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Insulin dependent diabetes mellitus
Required Medical Information and Criteria	<p><u>Insulin Dependent Diabetes Mellitus – Pediatric</u></p> <ul style="list-style-type: none"> • Documentation of Type 1 Diabetes Mellitus or Diabetes with C-reactive protein levels indicating insulin dependence • On intensive insulin therapy (>3 daily insulin injections) requiring frequent self-adjustments for at least 6 months prior to initiation of the insulin pump • Documentation self-testing of blood glucose at least 4 times per day during the previous 2 months • Evidence of completion of a comprehensive diabetes education program in the last 12 months (member or caregiver/parent) <p><u>Insulin Dependent Diabetes Mellitus – Adult</u></p> <ul style="list-style-type: none"> • All of the above pediatric requirements AND • Documentation of one of the following: <ul style="list-style-type: none"> ○ HbA1c >7% ○ History of recurring hypoglycemia ○ Wide fluctuations in blood glucose before mealtime ○ Dawn phenomenon with fasting blood sugars frequently exceeding 200mg/dL ○ History of severe glycemic excursions • Inability to use a traditional (non-disposable) insulin pump
Renewal Criteria	<ul style="list-style-type: none"> • Clinical documentation of positive clinical response to therapy, and an in-person visit with the prescribing provider within the last 6 months
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	3/1/2024
-----------------	----------

P&T Approval Date:	1/9/2024
P&T Revision Date:	1/9/2024

References
•

Donepezil

Products Affected

- Donepezil 5mg tablets
- Donepezil 10mg tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Alzheimer's disease – dementia○ Alzheimer's disease, prophylaxis – Impaired cognition○ Multi-Infarct dementia
Required Medical Information and Criteria	<u>All diagnoses</u> <ul style="list-style-type: none">• Confirmed diagnosis of one of the above listed conditions
Renewal Criteria	<ul style="list-style-type: none">• Positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	8/1/2013
P&T Approval Date:	4/5/2011
P&T Revision Date:	7/15/2013, 4/5/2011

References
<ul style="list-style-type: none">• Aricept [package insert]. Woodcliff Lake, NJ: Eisai Inc.; 2013.

Dipeptidyl Peptidase 4 (DPP-4) Inhibitor Step Therapy

Products Affected

- Janumet
- Janumet XR
- Januvia
- Kombiglyze XR
- Onglyza
- Tradjenta

Step Therapy Criteria	<ul style="list-style-type: none">• Clinical diagnosis of Type 2 Diabetes Mellitus (T2DM)• Trial and failure of the following:<ul style="list-style-type: none">○ Metformin○ Sulfonylurea or insulin
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Dronabinol

Products Affected

- Dronabinol capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Nausea and vomiting associated with cancer chemotherapy (CINV)○ AIDS anorexia
Required Medical Information and Criteria	<p><u>Nausea and Vomiting Associated with Cancer Chemotherapy</u></p> <ul style="list-style-type: none">• Confirmation of treatment with chemotherapy• Trial and failure or contraindication to all of the following:<ul style="list-style-type: none">○ At least one 5HT-3 receptor antagonist (e.g. ondansetron, granisetron, etc.)○ At least one of:<ul style="list-style-type: none">▪ Prochlorperazine▪ Dexamethasone▪ Haloperidol▪ Olanzapine <p><u>AIDS Anorexia</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of anorexia with weight loss in patients with AIDS• Patient is on antiretroviral therapy
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">• Marinol [package insert]. North Chicago, IL: Abbvie; 2017.

Dupilumab (Dupixent)

Products Affected

- Dupixent Prefilled Syringe
- Dupixent Auto-Injector

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Moderate to Severe Asthma ○ Atopic Dermatitis ○ Eosinophilic Esophagitis ○ Chronic Rhinosinusitis with Nasal Polyps ○ Prurigo Nodularis ○ Chronic Obstructive Pulmonary Disease
Required Medical Information and Criteria	<p><u>Moderate to Severe Asthma</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of moderate to severe asthma • Inadequate control of asthma symptoms despite fully maximized treatment with one of the following: <ul style="list-style-type: none"> ○ Inhaled corticosteroids combined with long-acting beta-2 agonist ○ Inhaled corticosteroids combined with long-acting muscarinic antagonist <p><u>Atopic Dermatitis</u></p> <ul style="list-style-type: none"> • Diagnosed with severe atopic dermatitis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) ≥ 11 or Children's Dermatology Life Quality Index (CDLQI) ≥ 13 (or severe score on another validated tool) • One or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involvement ○ Hand, foot, or mucous membrane involvement • Documented contraindication or failed trial to ALL of the following: <ul style="list-style-type: none"> ○ Moderate-high potency corticosteroid (e.g., clobetasol, fluocinonide, fluticasone) ○ Topical calcineurin inhibitor (e.g. tacrolimus) ○ Oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil) OR the member is oral corticosteroid dependent. <p><u>Eosinophilic Esophagitis</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of Eosinophilic Esophagitis

	<ul style="list-style-type: none"> • Weight \geq 15 kg • Two or more episodes of dysphagia per week • Inadequate response to an 8-week trial, intolerance, or contraindication to high-dose PPI therapy • Inadequate response to and 8-to-12-week trial, intolerance, or contraindication to swallowed inhaled respiratory corticosteroid therapy. <p><u>Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)</u></p> <ul style="list-style-type: none"> • Diagnosis of CRSwNP, including objective evidence of the presence of bilateral nasal polyps • Will not be used in combination with other biologics for eosinophilic indications. • Trial and failure to adequately reduce symptoms with: <ul style="list-style-type: none"> ○ At least 2 months of saline nasal irrigations and inhaled nasal corticosteroids used at doses appropriate for nasal polyp treatment. ○ Systemic corticosteroid treatment for nasal polyps at least once within the last 2 years or prior nasal polyp removal surgery. • Inhaled nasal corticosteroids will be used concomitantly with dupilumab (unless not tolerated or contraindicated). <p><u>Prurigo Nodularis</u></p> <ul style="list-style-type: none"> • Funded condition (as defined by guideline note 21 of the prioritized list) or age under 21. • Diagnosis of PN verified by a dermatologist and the patient has had the diagnosis for at least 3 months. • Severe or very severe itch (WI-NRS score \geq7) reported within the past week. • At least 20 PN lesions in total on both legs and/or both arms and/or trunk • Trial and failure (inadequate efficacy after 4-week trial, intolerable side effects) or contraindication to recommended first line agents for the treatment of PN including: <ul style="list-style-type: none"> ○ High potency topical steroids ○ Phototherapy ○ At least one systemic agent (immunosuppressant, gabapentinoid, or antidepressant). <p><u>Chronic Obstructive Pulmonary Disease (COPD)</u></p> <ul style="list-style-type: none"> • Diagnosis of COPD confirmed by post-bronchodilator FEV1/FVC $<$ 0.7 on spirometry.
--	---

	<ul style="list-style-type: none"> • Blood eosinophil count (BEC) ≥ 300 cells/μL within the past 3 months. • Chronic bronchitis, defined as a chronic productive cough for ≥ 3 months in the past year, in the absence of other known causes of chronic cough. • ≥ 2 moderate COPD exacerbation (defined as requiring treatment with either systemic corticosteroids and/or antibiotics) or ≥ 1 severe COPD exacerbation (defined as requiring hospitalization or observation for over 24 hours in emergency department or urgent care) within the past year despite the adherent use of inhaled LABA + LAMA+ ICS triple therapy [or LABA + LAMA dual therapy if ICS are contraindicated].
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> • Moderate to Severe Asthma: 6 years and older • Atopic Dermatitis: 6 months and older • Eosinophilic Esophagitis: 1 years and older • CRSwNP: 12 years and older • Prurigo Nodularis: 18 years and older • COPD: 18 years and older
Prescriber Restriction	<ul style="list-style-type: none"> • Atopic dermatitis: Dermatologist • Eosinophilic Esophagitis: Gastroenterologist or Immunologist • CRSwNP: ENT or Immunologist • Prurigo Nodularis: Dermatologist • Asthma, COPD: Pulmonologist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	3/1/2025
P&T Approval Date:	11/8/2022
P&T Revision Date:	1/14/2025, 1/9/2024, 11/8/2022

References

- Dupixent [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals Inc.; 2024.

Elagolix (Orilissa)

Products Affected

- Orilissa tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Moderate to Severe Pain Associated with Endometriosis
Required Medical Information and Criteria	<p><u>Moderate to Severe Pain Associated with Endometriosis</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of endometriosis with documentation of moderate to severe pain associated with the condition • Trial and failure of, or contraindication to, both of the following: <ul style="list-style-type: none"> ○ A 3-month trial of prescription strength NSAIDs ○ Two 3-month trials of hormonal therapies (e.g. combined oral contraceptives, progestins, or levonorgestrel IUD, etc.) • Additional info required for 200mg twice daily dosing: <ul style="list-style-type: none"> ○ Documentation of coexisting dyspareunia
Renewal Criteria	<ul style="list-style-type: none"> • Renewal only allowed for 150mg dose • Criteria requires documentation of: <ul style="list-style-type: none"> ○ A positive clinical response to therapy ○ Total therapy duration is 24 months or less
Age Restriction	<ul style="list-style-type: none"> • At least 18 years of age or older but not yet through menopause
Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Obstetrician or Gynecologist
Coverage Duration	<ul style="list-style-type: none"> • 150mg dose: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 18 months (maximum treatment duration of 24 months) • 200mg dose: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: no renewals allowed

Effective Date:	2/1/2023
P&T Approval Date:	1/10/2023

P&T Revision Date:	1/10/2023
-----------------------	-----------

References
<ul style="list-style-type: none"> Orilissa [package insert]. North Chicago, IL: Abbvie.; 2023.

Elefibranor (Iqirvo)

Products Affected

- Iqirvo tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Primary Biliary Cholangitis
Required Medical Information and Criteria	<p><u>Primary Biliary Cholangitis</u></p> <ul style="list-style-type: none">• Diagnosis of primary biliary cholangitis (PBC) confirmed by two of the following:<ul style="list-style-type: none">○ Biochemical evidence of cholestasis based on ALP elevation○ Presence of AMA or other PBC-specific autoantibodies○ Histology confirmation after biopsy• Trial and failure of 12 months of ursodiol.• No current decompensated cirrhosis.
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Primary Biliary Cholangitis: Gastroenterologist or Hepatologist
Coverage Duration	<ul style="list-style-type: none">• Primary Biliary Cholangitis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024

References

- Iqirvo [package insert]. Cambridge, MA: Ipsen Biopharmaceuticals Inc.; 2024.

Ellexacaftor-tezacaftor-ivacaftor (Trikafta)

Products Affected

- Trikafta

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design.<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<u>Cystic Fibrosis</u> <ul style="list-style-type: none">• Documentation of cystic fibrosis diagnosis with at least one F508del mutation• Not used in combination with other CFTR modulator treatments
Renewal Criteria	<ul style="list-style-type: none">• Documented clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 2 years of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Cystic Fibrosis: Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Cystic Fibrosis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	05/01/2025
P&T Approval Date:	05/01/2021
P&T Revision Date:	09/01/2021, 03/11/2025

References
<ul style="list-style-type: none">• Trikafta [package insert]. Boston, MA: Vertex Pharmaceuticals Incorporated; 2024.

Endothelin Receptor Antagonists

Products Affected

- Ambrisentan tablets
- Bosentan tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Pulmonary Arterial Hypertension
Required Medical Information and Criteria	<u>Pulmonary Arterial Hypertension</u> <ul style="list-style-type: none">• Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension)
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Ambrisentan: 18 years of age and over• Bosentan: 3 years and up
Prescriber Restriction	<ul style="list-style-type: none">• Pulmonary Arterial Hypertension: Cardiologist or Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Pulmonary Arterial Hypertension:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	7/13/2021
P&T Revision Date:	7/8/2025, 7/13/2021

References
<ul style="list-style-type: none">• Tracleer [package insert]. Titusville, NJ: Actelion Pharmaceuticals Inc; 2024.• Letairis [package insert]. Foster City, CA: Gilead Sciences Inc; 2025.

Erythromycin Gel and Solution

Products Affected

- Erythromycin 2% gel
- Erythromycin 2% solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<u>All Diagnoses</u> <ul style="list-style-type: none">• Documentation of trial and failure, intolerance, or contraindication to clindamycin 1% gel or solution
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
<ul style="list-style-type: none">• Acne clinical guideline (aad.org)

Erythropoietic Agents

Products Affected

- Retacrit

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Anemia Due to Chronic Kidney Disease (CKD) ○ Anemia Due to Chemotherapy ○ Anemia Associated with HIV Infection ○ Preoperative Use for Reduction of Allogeneic Blood Transfusion in Patients Undergoing Surgery ○ Anemia in Myelodysplastic Syndrome (MDS)
Required Medical Information and Criteria	<p><u>Anemia Due to Chronic Kidney Disease (CKD)</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of anemia with hematocrit less than 30% or hemoglobin less than 10g/dL within 30 days of request • Verification of iron evaluation for adequate iron stores • One of the following: <ul style="list-style-type: none"> ○ Patient is on dialysis ○ Patient is not on dialysis, but the rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. <p><u>Anemia Due to Chemotherapy</u></p> <ul style="list-style-type: none"> • Verification that other causes of anemia have been ruled out • Verification of anemia with hematocrit less than 30% or hemoglobin less than 10g/dL within the prior 2 weeks • Verification of iron evaluation for adequate iron stores • Verification that the cancer is a non-myeloid malignancy • Patient is receiving chemotherapy <p><u>Anemia Associated with HIV Infection</u></p> <ul style="list-style-type: none"> • Verification of anemia with hematocrit less than 36% or hemoglobin less than 12 g/dL within the prior 30 days • Verification of iron evaluation for adequate iron stores • Serum erythropoietin level less than or equal to 500 mU/mL • One of the following: <ul style="list-style-type: none"> ○ Patient is receiving zidovudine ○ Diagnosis of HIV infection

	<p><u>Preoperative use for reduction of allogeneic blood transfusion in patients undergoing surgery</u></p> <ul style="list-style-type: none"> • Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery • Hemoglobin (Hgb) is greater than 10 to less than or equal to 13 g/dL • Patient is at high risk for perioperative transfusions • Patient is unwilling or unable to donate autologous blood pre-operatively • Verification of iron evaluation for adequate iron stores <p><u>Anemia in Myelodysplastic Syndrome (MDS)</u></p> <ul style="list-style-type: none"> • Diagnosis of MDS • One of the following: <ul style="list-style-type: none"> ○ Serum erythropoietin level less than or equal to 500 mU/mL ○ Diagnosis of transfusion-dependent MDS • Verification of iron evaluation for adequate iron stores
<p>Renewal Criteria</p>	<p>Anemia Due to CKD</p> <ul style="list-style-type: none"> • There is a documented response to treatment • One of the following: <ul style="list-style-type: none"> ○ Patient is on dialysis and Hgb is 11g/dL or less ○ Patient is NOT on dialysis and Hgb is 10g/dL or less <p>Anemia Due to Chemotherapy</p> <ul style="list-style-type: none"> • There is a documented response to treatment as indicated by a decrease in the need to blood transfusion or Hgb increase from pre-treatment level • Documented Hct less than 30% or Hgb less than 10 g/dL • Patient is receiving chemotherapy <p>Anemia Associated with HIV Infection</p> <ul style="list-style-type: none"> • There is a documented response to treatment as indicated by a decrease in the need to blood transfusion or Hgb increase from pre-treatment level • Documented Hct less than 36% or Hgb less than 12 g/dL <p>Anemia in Myelodysplastic Syndrome</p> <ul style="list-style-type: none"> • There is a documented response to treatment as indicated by a decrease in the need to blood transfusion or Hgb increase from pre-treatment level • Documented Hct less than 36% or Hgb less than 12 g/dL

Coverage Duration	<ul style="list-style-type: none"> • Anemia due to CKD: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 12 months • Anemia Due to Chemotherapy: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 3 months • Anemia Associated with HIV: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 12 months • Anemia in MDS: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 12 months • Preoperative: <ul style="list-style-type: none"> ○ Initial: 1 months ○ Renewal: N/A

Effective Date:	9/1/2024
P&T Approval Date:	7/9/2024
P&T Revision Date:	7/9/2024

References
<ul style="list-style-type: none"> • Oregon Health Authority. (2024). <i>Guideline Note GN7: Erythropoiesis-Stimulating Agent (ESA) Guideline</i>. Health Evidence Review Commission.

Estradiol Ring (Estring)

Products Affected

- Estring 2mg vaginal ring

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<u>All FDA approved diagnoses</u> <ul style="list-style-type: none">• Trial and failure of estradiol vaginal cream and estradiol vaginal tablet
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	7/1/2023
P&T Approval Date:	5/9/2023
P&T Revision Date:	5/9/2023

References

- Estring [package insert]. New York, NY: Pfizer; 2023.

Etanercept (Enbrel)

Products Affected

- Enbrel Auto-Injector
- Enbrel Prefilled Syringe

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Plaque Psoriasis ○ Psoriatic Arthritis ○ Rheumatoid Arthritis ○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla. <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of both infliximab and adalimumab. <p><u>Plaque Psoriasis (PP):</u></p>

	<ul style="list-style-type: none"> • Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face, or mucous membrane involvement • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of all the following: <ul style="list-style-type: none"> ▪ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ▪ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ▪ PUVA or UVB Phototherapy ▪ Methotrexate ▪ At least 1 other second line systemic agent such as cyclosporine or acitretin. • Trial and failure of both infliximab and adalimumab. <p><u>Psoriatic Arthritis (PsA):</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of both infliximab and adalimumab.
--	---

	<p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab.
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Rheumatoid Arthritis: Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/1/2023
P&T Approval Date:	7/11/2023

P&T Revision Date:	7/11/2023
-----------------------	-----------

References
•

Etrasimod arginine (Velsipity)

Products Affected

- Velsipity tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>Ulcerative Colitis</u></p> <ul style="list-style-type: none">• Documentation of moderate-to-severe ulcerative colitis• The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following:<ul style="list-style-type: none">○ Mesalamine, sulfasalazine OR○ Mercaptopurine, azathioprine, OR○ Corticosteroids (prednisone, methylprednisolone)• Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none">• Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission
Age Restriction	<ul style="list-style-type: none">• 18 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• Ulcerative colitis: Gastroenterologist
Coverage Duration	<ul style="list-style-type: none">• Ulcerative colitis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	3/1/2024
P&T Approval Date:	1/9/2024
P&T Revision Date:	1/9/2024

References

- Velsipity [package insert]. New York, NY: Pfizer Inc.; 2024.

Ferrous Sulfate

Products Affected

- Ferrous Sulfate 300mg/5mL

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<u>All Diagnoses</u> <ul style="list-style-type: none">• Documentation of inability to use ferrous sulfate tabs or 200mg/5mL liquid
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	1/11/2024
P&T Approval Date:	11/14/2023
P&T Revision Date:	11/14/2023

Fezolinetant (Veozah)

Products Affected

- Veozah tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Vasomotor Symptoms Associated with Menopause
Required Medical Information and Criteria	<u>Vasomotor Symptoms Associated with Menopause</u> <ul style="list-style-type: none">• Diagnosis of moderate to severe vasomotor symptoms due to menopause.• Documented contraindication, intolerance, or inadequate response to at least 2 hormonal therapies.• Documented contraindication, intolerance, or inadequate response to two nonhormonal therapies (e.g., one SNRI and one SSRI).
Renewal Criteria	<ul style="list-style-type: none">• Documentation of at least 50% reduction in VMS from baseline.
Prescriber Restriction	<ul style="list-style-type: none">• All Diagnoses: Gynecologist
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
<ul style="list-style-type: none">• Veozah [package insert]. Northbrook, IL: Astellas Pharma US Inc.; 2023

Fingolimod

Products Affected

- Fingolimod 0.5mg capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Relapsing forms of Multiple Sclerosis
Required Medical Information and Criteria	<u>Multiple Sclerosis</u> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of a relapsing form of multiple sclerosis
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple Sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple Sclerosis:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
<ul style="list-style-type: none">• Gilenya [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Co.; 2023

Fluorouracil, topical

Products Affected

- Fluorouracil 5% cream
- Fluorouracil 2% solution
- Fluorouracil 5% solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Superficial Basal Cell Carcinoma
Required Medical Information and Criteria	<u>Basal Cell Carcinoma</u> <ul style="list-style-type: none">• Documentation confirming a diagnosis of superficial basal cell carcinoma with multiple lesions and/or difficult to treat areas.
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• Basal Cell Carcinoma:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">• Efudex [package insert]. Sugar Land, TX: Bausch Health Inc.; 2021

Glatiramer

Products Affected

- Glatiramer 20mg/mL inj
- Glatiramer 40mg/mL inj

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Relapsing forms of Multiple Sclerosis
Required Medical Information and Criteria	<u>Multiple Sclerosis</u> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of a relapsing form of multiple sclerosis
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple Sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple Sclerosis:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
<ul style="list-style-type: none">• Glatopa [package insert]. Princeton, NJ: Sandoz Inc.; 2023

Glucagon-Like Peptide-1 (GLP1) Receptor Agonist

Products Affected

- Trulicity Auto-Injector
- Liraglutide Pen-Injector

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Type 2 Diabetes
Required Medical Information and Criteria	<p><u>Type 2 Diabetes</u></p> <ul style="list-style-type: none"> • Documentation of clinically diagnosed Type 2 Diabetes • Documentation of adequate trial of maximally tolerated dose of metformin • Documentation of trial and failure of one of the following (or contraindication to both): <ul style="list-style-type: none"> ○ An SGLT-2 inhibitor (e.g. Farxiga, Jardiance, etc.) ○ A DPP-4 inhibitor (e.g. saxagliptin, Tradjenta, Januvia, etc.)
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy such as a decrease in A1c
Coverage Duration	<ul style="list-style-type: none"> • Type 2 Diabetes: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
•

Golimumab (Simponi)

Products Affected

- Simponi Auto-Injector
- Simponi Prefilled Syringe

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Psoriatic Arthritis ○ Rheumatoid Arthritis ○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla. <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of both infliximab and adalimumab. <p><u>Psoriatic Arthritis (PsA):</u></p>

	<ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of both infliximab and adalimumab. <p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab. <p><u>Ulcerative Colitis (UC):</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: mesalamine, sulfasalazine, mercaptopurine, azathioprine, or corticosteroids (prednisone, methylprednisolone).
--	---

	<ul style="list-style-type: none"> • Trial and failure of both infliximab and adalimumab.
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Ulcerative Colitis: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Rheumatoid Arthritis: Rheumatologist. • Ulcerative Colitis: Gastroenterologist.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/01/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
•

Gonadotropin-Releasing Hormone Agonists

Products Affected

- Lupron

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Endometriosis ○ Uterine Leiomyomata (Fibroids) – For the reduction of the size of fibroids ○ Uterine Leiomyomata (Fibroids) – Anemia ○ Central Precocious Puberty (CPP) ○ Prostate Cancer ○ Gender Dysphoria/Gender Incongruence
Required Medical Information and Criteria	<p><u>Endometriosis</u></p> <ul style="list-style-type: none"> • One of the following: <ul style="list-style-type: none"> ○ History of inadequate pain control response following and trial of at least 6 months or history of intolerance or contraindication to one of the following: <ul style="list-style-type: none"> ▪ Danazol ▪ Combination (estrogen/progestin) oral contraceptive ▪ Progestins ○ Patient has had surgical ablation to prevent recurrence <p><u>Uterine Leiomyomata (Fibroids) – For the reduction of the size of fibroids</u></p> <ul style="list-style-type: none"> • For use prior to surgery to reduce the size of fibroids to facilitate a surgical procedure (e.g., myomectomy, hysterectomy) <p><u>Uterine Leiomyomata (Fibroids) – Anemia</u></p> <ul style="list-style-type: none"> • Anemia is caused by uterine leiomyomata (fibroids) • Patient has tried and had an inadequate response to at least 1 month of monotherapy with iron • Used in combination with iron therapy • For use prior to surgery <p><u>Central Precocious Puberty (CPP)</u></p> <ul style="list-style-type: none"> • Diagnosis of central precocious puberty (idiopathic or neurogenic)

	<ul style="list-style-type: none"> • Early onset of secondary sexual characteristics in one of the following: <ul style="list-style-type: none"> ○ Females less than 8 years of age ○ Males less than 9 years of age • Advanced bone age of at least one year compared with chronological age • One of the following: <ul style="list-style-type: none"> ○ Patient has undergone gonadotropin-releasing hormone agonist (GnRHa) testing and peak luteinizing hormone (LH) level above pre-pubertal range ○ Patient has a random LH level in the pubertal range • One of the following: <ul style="list-style-type: none"> ○ Patient had one of the following diagnostic evaluations to rule out tumors, when suspected: <ul style="list-style-type: none"> ▪ Diagnostic imaging of the brain (MRI or CT scan) (in patients with symptoms suggestive of a brain tumor or in those 6 years of age or younger) ▪ Pelvic/testicular/adrenal ultrasound (if steroid levels suggest suspicion) ▪ Adrenal steroids to rule out congenital adrenal hyperplasia (when pubarche precedes thelarche or gonadarche) ○ Patient has no suspected tumors <p><u>Prostate Cancer</u></p> <ul style="list-style-type: none"> • Diagnosis of advanced or metastatic prostate cancer <p><u>Gender Dysphoria/Gender Incongruence</u></p> <ul style="list-style-type: none"> • Using gonadotropin for suppression of puberty • Diagnosis of gender dysphoria/gender incongruence
Renewal Criteria	<ul style="list-style-type: none"> • Endometriosis: Recurrence of symptoms following a trial of at least 6 months with leuprolide acetate AND used in combination with one of the following: Norethindrone 5mg daily, other “add-back” sex-hormones (e.g. estrogen, medroxyprogesterone), other bone-sparing agents (e.g., bisphosphonates). • Central Precocious Puberty (CPP): LH levels have been suppressed to pre-pubertal levels. • Prostate Cancer: Patient does not show evidence of progressive disease while on therapy.

Prescriber Restriction	<ul style="list-style-type: none"> • Central Precocious Puberty (CPP): Pediatric endocrinologist
Coverage Duration	<ul style="list-style-type: none"> • Endometriosis: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: up to 6 months • Uterine Leiomyomata (Fibroids Reduction): <ul style="list-style-type: none"> ○ Initial: 4 months • Uterine Leiomyomata (Fibroids Anemia): <ul style="list-style-type: none"> ○ Initial: 3 months • Central Precocious Puberty (CPP): <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: up to 12 months • Prostate Cancer: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: up to 12 months • Gender Dysphoria: <ul style="list-style-type: none"> ○ Initial: 12 months

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	5/14/2024, 1/10/2023

References
•

Grass Pollen Allergen Extract -Timothy Grass (Grastek)

Products Affected

- Grastek

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Grass pollen-induced allergic rhinitis
Required Medical Information and Criteria	<u>Grass Pollen-Induced Allergic Rhinitis</u> <ul style="list-style-type: none">• Confirmed diagnosis of grass pollen induced allergic rhinitis
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Grass Pollen-Induced Allergic Rhinitis: Allergist or Immunologist
Coverage Duration	<ul style="list-style-type: none">• Grass Pollen-Induced Allergic Rhinitis:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 3 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Hepatitis C Antivirals

Products Affected

- Mavyret
- Sofosbuvir-Velpatasvir
- Vosevi

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Treatment of Hepatitis C Infection
Required Medical Information and Criteria	<p><u>Hepatitis C Infection</u> *PA for Mavyret and Sofosbuvir-Velpatasvir is only required if there is a history of prior treatment (treatment experienced). Vosevi requires PA in all cases. *</p> <ul style="list-style-type: none"> • All appropriate pre-treatment testing has been done: <ul style="list-style-type: none"> ○ Genotype testing in the past 3 years is required if the patient has decompensated cirrhosis, prior treatment experience with DAA regimen, and if prescribed a regimen which is not pan-genotypic ○ History of previous HCV treatment, viral load after treatment, and outcome are required only if there is documentation of treatment experience • Documentation of prior treatment history • Documentation of whether the patient achieved a sustained virological response (SVR) at week 12 or longer following the completion of their last DAA regimen • Documentation if this is likely a reinfection, indicated by at least one of the following: <ul style="list-style-type: none"> ○ Does the patient have ongoing risk factors for hepatitis C reinfection (e.g. sexually active men who have sex with men, persons who inject drugs) ○ Is the hepatitis C infection due a different genotype than previous • If the request is for elbasvir/grazoprevir for GT 1a, ledipasvir/sofosbuvir for GT 1a treatment experienced or sofosbuvir for GT 3 with cirrhosis or treatment experience has the patient had a baseline NS5a resistance test. (Note: baseline NS5A resistance testing is required per OHA in this situation) • The prescribed drug regimen a recommended regimen based on the patient's genotype, age, treatment status (retreatment or treatment naïve) and cirrhosis status (see table 1 and 2)?

**Per OHA: treatment experienced are patients who received more than 4 weeks of HCV DAA therapy.*

Table 1. Recommended Regimens for Adults and adolescents 12 years of age and older

Treatment History	Cirrhosis Status	Recommended Regimen
Treatment Naïve (Genotype 1-6)		
Treatment naïve	Non-cirrhotic	SOF/VEL x 12 weeks G/P x 8 weeks
	Compensated cirrhosis	G/P x 8 weeks SOF/VEL x 12 weeks (baseline resistance testing recommended for GT3)
	Decompensated cirrhosis	SOF/VEL + RBV x 12 weeks SOF/VEL x 24 weeks (if RBV ineligible)
Treatment Experienced (Genotype 1-6)		
<u>Sofobuvir based regimen treatment failures, including:</u> sofosbuvir + ribavirin Ledipasvir/Sofobuvir Velpatasvir/sofobuvir	Non-cirrhotic or compensated cirrhosis	SOF/VEL/VO X x 12 weeks G/P x16 weeks (except GT3)
Elbasvir/grazoprevir treatment failures	Non-cirrhotic or compensated cirrhosis	SOF/VEL/VO X x 12 weeks
Glecaprevir/pibrentasvir treatment failures	Non-cirrhotic or compensated cirrhosis	G/P + SOF + RBV x 16 weeks SOF/VEL/VO X x 12 weeks (plus RBV if compensated cirrhosis)
<u>Multiple DAA Treatment Failures, including:</u> Sofobuvir/velpatasvir/voxilaprevir	Non-Cirrhotic or compensated cirrhosis	G/P + SOF + RBV x 16-24 weeks SOF/VEL/VO X x24 weeks

	Glecaprevir/pibrentasvir + sofosbuvir		
Table 2. Recommended Regimens for children 3-12 years of age			
	Treatment History	Cirrhosis Status	Recommended Regimen
	Treatment Naïve (Genotype 1-6)		
	Treatment Naïve	Non-cirrhotic or compensated cirrhosis	SOF/VEL x 12 weeks G/P x 8 weeks
		Decompensated cirrhosis	SOF/VEL + RBV x 12 weeks
	<p>Abbreviations: DAA = direct acting antiviral; EBV/GZR = elbasvir/grazoprevir; G/P = glecaprevir and pibrentasvir; PEG = pegylated interferon; RBV = ribavirin; SOF = sofosbuvir; SOF/VEL = sofosbuvir/velpatasvir; SOF/VEL/VOX = sofosbuvir/velpatasvir/voxilaprevir</p> <p>There is limited data supporting DAA regimens in treatment-experienced patients with decompensated cirrhosis. These patients should be handled on a case by case basis with the patient, prescriber, and CCO or FFS medical director</p> <p>Ribavirin ineligible/intolerance may include: 1) neutrophils < 750 mm³, 2) hemoglobin < 10 g/dl, 3) platelets <50,000 cells/mm³, autoimmune hepatitis or other autoimmune condition, hypersensitivity or allergy to ribavirin</p>		
Coverage Duration	<ul style="list-style-type: none"> • Hepatitis C Infection: <ul style="list-style-type: none"> ○ Initial: 8-24 weeks. Duration of approval depends on the specifics of the infection. 		

Effective Date:	12/1/2022
P&T Approval Date:	11/8/2022
P&T Revision Date:	11/8/2022

References
•

Inclisiran (Leqvio)

Products Affected

- Leqvio Prefilled Syringe

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Established clinical ASCVD ○ Primary or Familial Hyperlipidemia
Required Medical Information and Criteria	<p><u>Established Clinical ASCVD</u></p> <ul style="list-style-type: none"> • Documentation of very high risk ASCVD as evidenced by either: <ul style="list-style-type: none"> ○ History of multiple major ASCVD events OR ○ One major ASCVD event AND multiple high-risk conditions. • Documentation of a current LDL greater than or equal to 55 mg/dl. • Documentation of all the following: <ul style="list-style-type: none"> ○ Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins ○ Is receiving ezetimibe or has a documented intolerance to ezetimibe. • Documentation of failure of PCSK9 inhibitor (Repatha or Praluent). <p><u>Primary or Familial Hyperlipidemia</u></p> <ul style="list-style-type: none"> • Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL. • Documentation of current LDL greater than 100 mg/dL. • Documentation of all the following: <ul style="list-style-type: none"> ○ Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins ○ Is receiving ezetimibe or has a documented intolerance to ezetimibe. • Documentation of failure of PCSK9 inhibitor (Repatha or Praluent).
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy (significant decrease in lipid levels).

Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Cardiologist, Endocrinologist, or Lipid Specialist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024

References
•

Insulin Degludec

Products Affected

- Insulin Degludec Flextouch 100 unit/mL
- Insulin Degludec Flextouch 200 unit/mL

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Type 1 Diabetes○ Type 2 Diabetes
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none">• Trial and failure of Insulin Glargine or documented intolerance or contraindication to Insulin Glargine<ul style="list-style-type: none">○ Subset criteria 1○ Etc• Documentation of significant barriers to standardized administration requiring flexibility in dose timing<ul style="list-style-type: none">○ Subset 2○ Etc.• For U-200 concentration only:<ul style="list-style-type: none">○ Insulin requirements of greater than 160 units of insulin per dose○ Difficulty with multiple daily injections
Renewal Criteria	<ul style="list-style-type: none">• Renewal criteria
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References

•

Insulin U-500 (Humulin R U-500)

Products Affected

- Humulin R U-500 Pens
- Humulin R U-500 Vials

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Type 1 Diabetes○ Type 2 Diabetes
Required Medical Information and Criteria	<u>All Diagnoses</u> <ul style="list-style-type: none">• Attestation that the use of U-500 insulin is medically safe and appropriate
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	3/1/2025
P&T Approval Date:	1/14/2025
P&T Revision Date:	1/14/2025

References
<ul style="list-style-type: none">•

Interferon beta-1a (Avonex)

Products Affected

- Avonex Pen
- Avonex prefilled syringe

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Relapsing forms of Multiple Sclerosis
Required Medical Information and Criteria	<p><u>Multiple Sclerosis</u></p> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of a relapsing form of multiple sclerosis• Trial and failure of all the following:<ul style="list-style-type: none">○ Dimethyl fumarate○ Fingolimod○ Glatiramer
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple Sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple Sclerosis:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References

- Avonex [package insert]. Cambridge, MA: Biogen; 2023

Interferon beta-1a (Rebif)

Products Affected

- Rebif Inj.
- Rebif Rebido

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Relapsing forms of Multiple Sclerosis
Required Medical Information and Criteria	<p><u>Multiple Sclerosis</u></p> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of a relapsing form of multiple sclerosis• Trial and failure of all the following:<ul style="list-style-type: none">○ Dimethyl fumarate○ Fingolimod○ Glatiramer
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple Sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple Sclerosis:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
<ul style="list-style-type: none">• Rebif [package insert]. Rockland, MA: EMD Serono Inc.; 2023

Interferon beta-1b (Extavia)

Products Affected

- Extavia Inj.

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Relapsing forms of Multiple Sclerosis
Required Medical Information and Criteria	<p><u>Multiple Sclerosis</u></p> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of a relapsing form of multiple sclerosis• Trial and failure of all the following:<ul style="list-style-type: none">○ Dimethyl fumarate○ Fingolimod○ Glatiramer
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Multiple Sclerosis: Neurologist
Coverage Duration	<ul style="list-style-type: none">• Multiple Sclerosis:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
<ul style="list-style-type: none">• Extavia [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Co.; 2023

Isotretinoin

Products Affected

- Amnesteem capsules
- Isotretinoin capsules
- Myorisan capsules
- Claravis capsules
- Zenetane capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Severe nodulocystic acne
Required Medical Information and Criteria	<u>Severe Nodulocystic Acne</u> <ul style="list-style-type: none">• Confirmed diagnosis of severe nodulocystic acne• Documentation of trial and failure, intolerance, or contraindication to combined therapy with an oral antibiotic and topical therapy (benzoyl peroxide or retinoid)
Renewal Criteria	<ul style="list-style-type: none">• No Renewals
Age Restriction	<ul style="list-style-type: none">• 12 years and older
Coverage Duration	<ul style="list-style-type: none">• Severe Nodulocystic Acne:<ul style="list-style-type: none">○ Initial: 20 weeks○ Renewal: No Renewals

Effective Date:	2/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

Itraconazole

Products Affected

- Itraconazole Oral Solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Tinea Unguium
Required Medical Information and Criteria	<p><u>Tinea Unguium</u></p> <ul style="list-style-type: none">• Documentation of one of the following:<ul style="list-style-type: none">○ Patient is experiencing pain which limits normal activity (i.e, unable to wear shoes, difficulty walking, etc),○ Patient has diabetes,○ Patient has peripheral vascular disease,○ Patient is immunocompromised <p><u>All Other Diagnoses</u></p> <ul style="list-style-type: none">• Diagnosis is supported by compendia and is an above the line condition
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 3 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References
•

Ivacaftor (Kalydeco)

Products Affected

- Kalydeco Tablets
- Kalydeco Packets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design.<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<p><u>Cystic Fibrosis</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of cystic fibrosis with documentation showing at least one CFTR gene mutation that has shown to be responsive to Kalydeco• Not used in combination with other CFTR modulator treatments
Renewal Criteria	<ul style="list-style-type: none">• Documented clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 1 month of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Cystic Fibrosis: Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Cystic Fibrosis:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 6 months

Effective Date:	05/01/2025
P&T Approval Date:	05/01/2021
P&T Revision Date:	09/01/2021, 03/11/2025

References
<ul style="list-style-type: none">•

Ivermectin

Products Affected

- Ivermectin Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<u>All Diagnoses</u> <ul style="list-style-type: none">• Clinical documentation of a diagnosis for which ivermectin is FDA approved or compendia supported
Renewal Criteria	<ul style="list-style-type: none">• Documented reinfection
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 6 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Ixekizumab (Taltz)

Products Affected

- Taltz Auto-Injector
- Taltz Prefilled Syringe

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Plaque Psoriasis ○ Psoriatic Arthritis
Required Medical Information and Criteria	<p><u>All diagnoses:</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of infliximab and adalimumab <p><u>Plaque Psoriasis (PP):</u></p> <ul style="list-style-type: none"> • Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality

	<p>Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following:</p> <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face, or mucous membrane involvement • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of all the following: <ul style="list-style-type: none"> ▪ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ▪ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ▪ PUVA or UVB Phototherapy ▪ Methotrexate ▪ At least 1 other second line systemic agent such as cyclosporine or acitretin. • Trial and failure of infliximab and adalimumab <p><u>Psoriatic Arthritis (PsA):</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA

	<p>and/or functioning, such as ASAS40 or 2-point improvement in BASDAI</p> <ul style="list-style-type: none"> • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • Hidradenitis Suppurativa and Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Rheumatoid Arthritis, Juvenile Idiopathic Arthritis: Rheumatologist. • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist.

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
•

Lacosamide (Vimpat)

Products Affected

- Lacosamide tablets
- Lacosamide solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Focal Seizures○ Primary Generalized Tonic-Clonic Seizures
Required Medical Information and Criteria	<p><u>Focal Seizures</u></p> <ul style="list-style-type: none">• Documentation confirming epilepsy or seizure disorder• Solution only: Member under age 10 or unable to use tablets <p><u>Primary Generalized Tonic-Clonic Seizures</u></p> <ul style="list-style-type: none">• Documentation confirming epilepsy or seizure disorder• Solution only: Member under age 10 or unable to use tablets
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Solution only: Member under age 10 or unable to use tablets
Prescriber Restriction	<ul style="list-style-type: none">• All diagnoses: Neurologist
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: Lifetime

Effective Date:	05/01/2025
P&T Approval Date:	07/11/2023
P&T Revision Date:	03/12/2024, 03/11/2025

References
<ul style="list-style-type: none">• Vimpat [package insert]. Smyrna, GA: UCB Inc.; 2023.

Lanthanum Carbonate

Products Affected

- Lanthanum Carbonate Chew Tabs

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Hyperphosphatemia in Chronic Kidney Disease
Required Medical Information and Criteria	<u>Hyperphosphatemia in Chronic Kidney Disease</u> <ul style="list-style-type: none">• Confirmed diagnosis of hyperphosphatemia in chronic kidney disease• Trial and failure, contraindication, or intolerance (at least 6 weeks) to both of the following at maximally tolerated doses<ul style="list-style-type: none">○ Calcium acetate○ Sevelamer carbonate
Age Restriction	<ul style="list-style-type: none">• 6 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• Hyperphosphatemia in CKD: Nephrologist
Coverage Duration	<ul style="list-style-type: none">• Hyperphosphatemia in CKD:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	7/1/2023
P&T Approval Date:	5/9/2023
P&T Revision Date:	5/9/2023

References
<ul style="list-style-type: none">•

Lasmiditan (Reyvow)

Products Affected

- Reyvow tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of acute migraine
Required Medical Information and Criteria	<p><u>Acute Migraine</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of migraine• Documentation showing the member is currently on preventative therapy• Trial and failure (defined as at least 6 weeks per agent) of:<ul style="list-style-type: none">○ At least 3 oral formulary triptans used at up to the maximally indicated dosing and in combination with NSAID therapy (e.g. naproxen).
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 18 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• Acute Migraine: Neurologist or headache specialist
Coverage Duration	<ul style="list-style-type: none">• Acute Migraine:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	5/9/2023
P&T Revision Date:	7/8/2025, 5/9/2023

References

- Reyvow [package insert]. Indianapolis, IN: Eli Lilly and Co; 2022.

Lebrikizumab (Ebglyss)

Products Affected

- Ebglyss Prefilled Syringe
- Ebglyss Auto-Injector

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Atopic Dermatitis
Required Medical Information and Criteria	<p><u>Atopic Dermatitis</u></p> <ul style="list-style-type: none"> Diagnosed with severe atopic dermatitis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) ≥ 11 or Children's Dermatology Life Quality Index (CDLQI) ≥ 13 (or severe score on another validated tool) One or more of the following: <ul style="list-style-type: none"> At least 10% of body surface area involvement Hand, foot, or mucous membrane involvement Documented contraindication or failed trial to ALL of the following: <ul style="list-style-type: none"> Moderate-high potency corticosteroid (e.g., clobetasol, fluocinonide, fluticasone) Topical calcineurin inhibitor (e.g. tacrolimus) Oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil) OR the member is oral corticosteroid dependent.
Renewal Criteria	<ul style="list-style-type: none"> Documentation of positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none"> Atopic dermatitis: Dermatologist
Coverage Duration	<ul style="list-style-type: none"> All Diagnoses: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References
<ul style="list-style-type: none">•

Lenacapavir (Sunlenca)

Products Affected

- Sunlenca Therapy Pack
- Sunlenca Subcutaneous

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Multi-Drug-Resistant (MDR) HIV
Required Medical Information and Criteria	<p><u>Multi-Drug-Resistant HIV</u></p> <ul style="list-style-type: none">• Diagnosis of MDR HIV-1 infection with resistance to at least two drugs in each of at least three of the following classes: NRTIs, NNRTIs, PIs, and INSTIs• Will be used in combination with an optimized baseline regimen (OBR)• Current ARV regimen has been stable for at least 2 months• HIV-1 RNA is ≥ 400 copies/mL
Renewal Criteria	<ul style="list-style-type: none">• Continues to be used in combination with an optimized background regimen (OBR)• Provider states that patient continues to receive clinical benefit from the treatment
Age Restriction	<ul style="list-style-type: none">• Age 18 or older
Prescriber Restriction	<ul style="list-style-type: none">• MDR HIV Infection: HIV Specialist
Coverage Duration	<ul style="list-style-type: none">• MDR HIV Infection:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	4/1/2023
P&T Approval Date:	3/14/2023
P&T Revision Date:	3/14/2023

References
•

Lidocaine Patch

Products Affected

- Lidocaine External Patch

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Post-Herpetic Neuralgia
Required Medical Information and Criteria	<u>Post-Herpetic Neuralgia</u> <ul style="list-style-type: none">• Documentation confirming a diagnosis of post-herpetic neuralgia
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• Post-Herpetic Neuralgia:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 3 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Lisdexamfetamine

Products Affected

- Lisdexamfetamine Capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Attention Deficit Hyperactivity Disorder○ Binge Eating Disorder
Required Medical Information and Criteria	<p><u>Attention Deficit Hyperactivity Disorder (ADHD)</u></p> <ul style="list-style-type: none">• Prior trial of at least 30 days of both of the following:<ul style="list-style-type: none">○ An extended-release amphetamine product (amphetamine salts ER, dextroamphetamine ER, etc.)○ An extended-release methylphenidate product (dexmethylphenidate ER, methylphenidate ER, etc.) <p><u>Binge Eating Disorder (BED)</u></p> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of binge eating disorder per DSM-5 criteria• Trial and failure of at least 2 of the following therapeutic alternatives:<ul style="list-style-type: none">○ Selective serotonin reuptake inhibitors (SSRI)○ Topiramate○ Methylphenidate
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Binge Eating Disorder: 18 years of age or older
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2025
P&T Approval Date:	1/14/2025
P&T Revision Date:	1/14/2025

References
<ul style="list-style-type: none">• www.columbiapsychiatry.org/news/effort-underway-develop-first-u-s-guidelines-adhd-adults• www.cdc.gov/adhd/php/adults/index.html• www.cdc.gov/adhd/diagnosis/index.html

Long-Acting Opiates

Products Affected

- Fentanyl Patch
- Hydrocodone ER Capsule
- Hydrocodone ER Tablet
- Hydromorphone ER
- Methadone ER
- Morphine Sulfate ER Capsule
- Morphine Sulfate ER Tablets
- Nucynta ER
- Oxycodone ER
- Oxymorphone ER
- Xtampza ER

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Pain Related to Cancer, End of Life, or Palliative Care Pain in All Other Conditions
Required Medical Information and Criteria	<p><u>Pain Related to Cancer, End of Life, or Palliative Care</u></p> <ul style="list-style-type: none"> Confirmation of diagnosis <p><u>Other Pain</u></p> <ul style="list-style-type: none"> Documented use of current and/or recent usage of short-acting opioids for at least 15 days prior to long-acting opioids. For opioid naive (14 or fewer days filled in previous 120 days): 7-day maximum quantity limit; equal to or less than 50 MED [morphine equivalents per day]. For opioid experienced (greater than or equal to 15 days filled in previous 120 days): equal to or less than 90 MED [morphine equivalents per day]. Restricted to 2 fills in a 60-day period for both naive and experienced.
Renewal Criteria	<ul style="list-style-type: none"> Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> All Diagnoses: <ul style="list-style-type: none"> Initial: 12 months Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
•

Lotilaner (Xdemvy)

Products Affected

- Xdemvy 0.25% Ophthalmic solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Demodex Blepharitis
Required Medical Information and Criteria	<u>Demodex Blepharitis</u> <ul style="list-style-type: none">• Documentation of at least mild erythema of the upper eyelid margin• Presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination
Renewal Criteria	<ul style="list-style-type: none">• N/A
Age Restriction	<ul style="list-style-type: none">• 18 years of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Demodex Blepharitis: Optometrist or Ophthalmologist
Coverage Duration	<ul style="list-style-type: none">• Demodex Blepharitis:<ul style="list-style-type: none">○ Initial: 6 weeks○ Renewal: Renewals are not allowed

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References
<ul style="list-style-type: none">•

Lubiprostone

Products Affected

- Lubiprostone capsules

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Chronic idiopathic constipation in adults (in presence of above the line comorbid condition or if age under 21) ○ Irritable bowel syndrome in females (assigned at birth) > 18 YO (in presence of above the line comorbid condition or if age under 21) ○ Opioid induced constipation (in presence of above the line comorbid condition or if age under 21)
Required Medical Information and Criteria	<p><u>Chronic Idiopathic Constipation (CIC)</u></p> <ul style="list-style-type: none"> • Documentation of a funded comorbid condition that would be improved with the treatment of CIC • Documentation of failure to have benefit with recommended conventional first-line treatments (or contraindication to all first-line treatments): <ul style="list-style-type: none"> ○ Dietary Modification: Increased dietary fiber (25 grams/day) and increased fluid consumption ○ Bulk-forming Laxatives: Psyllium ○ Osmotic Laxatives: Polyethylene glycol, lactulose, magnesium hydroxide, milk of magnesia ○ Stool Softener: Docusate ○ Stimulant Laxatives: Senna, bisacodyl. <p><u>Irritable Bowel Syndrome with Constipation (IBS-C) in females (assigned at birth)</u></p> <ul style="list-style-type: none"> • Documentation of a funded comorbid condition that would be improved with the treatment of IBS-C • Documentation of failure to have benefit with recommended conventional first-line treatments (or contraindication to all first-line treatments): <ul style="list-style-type: none"> ○ Dietary Modification: Increased dietary fiber (25 grams/day) and increased fluid consumption ○ Bulk-forming Laxatives: Psyllium ○ Osmotic Laxatives: Polyethylene glycol, lactulose, magnesium hydroxide, milk of magnesia ○ Stool Softener: Docusate ○ Stimulant Laxatives: Senna, bisacodyl.

	<p><u>Opioid Induced Constipation (OIC)</u></p> <ul style="list-style-type: none"> • Documentation of a funded comorbid condition that would be improved with the treatment of OIC • Documentation of failure to have benefit with recommended conventional first-line treatments (or contraindication to all first-line treatments): <ul style="list-style-type: none"> ○ Dietary Modification: Increased dietary fiber (25 grams/day) and increased fluid consumption ○ Osmotic Laxatives: Polyethylene glycol, lactulose, magnesium hydroxide, milk of magnesia ○ Stool Softener: Docusate ○ Stimulant Laxatives: Senna, bisacodyl. • The patient is not using methadone.
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of efficacy and continued need
Age Restriction	<ul style="list-style-type: none"> • 18 years of age and older
Prescriber Restriction	<ul style="list-style-type: none"> • Chronic Idiopathic Constipation and Irritable Bowel Syndrome with Constipation: Gastroenterologist, other provider after patient consultation with a dietician • Opioid Induced Constipation: Gastroenterologist, pain management specialist, other provider after patient consultation with a dietician
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	7/8/2025
P&T Revision Date:	7/8/2025

References
<ul style="list-style-type: none"> • Change L, et al. American Gastroenterological Association-American College of Gastroenterology Clinical Practice Guideline: Pharmacological Management of Chronic Idiopathic Constipation. <i>Gastroenterology</i> 2023;164:1086–1106.

- Davids JS, et al. Clinical Practice Guidelines. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Anal Fissures. *Dis Colon Rectum* 2022; 66: 190–199.
- Paquette IM, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Evaluation and Management of Constipation. *Dis Colon Rectum* 2016; 59: 479–492.
- Tabbers MM, et al. Evaluation and Treatment of Functional Constipation in Infants and Children: Evidence-Based Recommendations from ESPGHAN and NASPGHAN. *JPGN* 2014;58: 258–274.
- Wald A, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. *Am J Gastroenterol* 2021;116:1987–2008.

Lumacaftor-Ivacaftor (Orkambi)

Products Affected

- Orkambi Granules
- Orkambi Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design.<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<p><u>Cystic Fibrosis</u></p> <ul style="list-style-type: none">• Documentation of cystic fibrosis diagnosis with homozygous F508del mutation• Not used in combination with other CFTR modulator treatments
Renewal Criteria	<ul style="list-style-type: none">• Documented clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 1 years of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Cystic Fibrosis: Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Cystic Fibrosis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	5/1/2025
P&T Approval Date:	5/1/2021
P&T Revision Date:	9/1/2021, 3/11/2025

References
<ul style="list-style-type: none">•

Mefloquine

Products Affected

- Mefloquine Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of Malaria
Required Medical Information and Criteria	<u>Malarial Treatment</u> <ul style="list-style-type: none">• Confirmed diagnosis of malaria
Renewal Criteria	<ul style="list-style-type: none">• N/A
Coverage Duration	<ul style="list-style-type: none">• Malaria:<ul style="list-style-type: none">○ Initial: 1 months○ Renewal: N/A

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Memantine

Products Affected

- Memantine 5mg Tablets
- Memantine 10mg Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Moderate to Severe Dementia of the Alzheimer's Type
Required Medical Information and Criteria	<p><u>Moderate to Severe Dementia of the Alzheimer's Type</u></p> <ul style="list-style-type: none">• Clinical documentation confirming a diagnosis of moderate to severe dementia of the Alzheimer's type. <p>Solution only: Documentation that the member is unable to use tablets</p>
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• Solution only: Age less than 10 (or can't swallow tablets)
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References
•

Mirabegron (Myrbetriq)

Products Affected

- Mirabegron ER tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Overactive Bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency ○ Neurogenic detrusor overactivity in pediatric members
Required Medical Information and Criteria	<p><u>Overactive Bladder with symptoms of urge urinary incontinence, urgency, and urinary frequency</u></p> <ul style="list-style-type: none"> • Documented trial and failure of at least 3 of the following, or contraindication to all: <ul style="list-style-type: none"> ○ Oxybutynin IR or ER ○ Fesoterodine ○ Solifenacin ○ Tolterodine IR or ER ○ Trospium IR or ER (requires step therapy through oxybutynin) <p><u>Neurogenic Detrusor Overactivity in pediatric members</u></p> <ul style="list-style-type: none"> • Is there documented trial and failure, intolerance, or contraindication to both of the following: <ul style="list-style-type: none"> ○ Oxybutynin IR or ER ○ Solifenacin
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> • 3 years of age and older
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

References

- Cameron AP, Chung DE, Dielubanza EJ, et al. The AUA/SUFU guideline on the diagnosis and treatment of idiopathic overactive bladder. J Urol. Published online April 23, 2024.
- Myrbetriq [package insert]. Northbrook, IL: Astellas.; 2021.

Mirikizumab (OmvoH)

Products Affected

- Omvoh 100mg/mL Auto-Injector
- Omvoh 100mg/mL Prefilled Syringe

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla <p><u>Ulcerative Colitis (UC)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following: <ul style="list-style-type: none"> ○ Mesalamine, sulfasalazine OR ○ Mercaptopurine, azathioprine, OR ○ Corticosteroids (prednisone, methylprednisolone) • Trial and failure of both infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Exclusion Criteria	<ul style="list-style-type: none"> • Not to be used in combination with other biologics for the same indication.
Prescriber Restriction	<ul style="list-style-type: none"> • Ulcerative Colitis: Gastroenterologist
Coverage Duration	<ul style="list-style-type: none"> • Ulcerative Colitis: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	7/1/2024
-----------------	----------

P&T Approval Date:	5/14/2024
P&T Revision Date:	5/14/2024

References
•

Mitapivat (Pyrukynd)

Products Affected

- Pyrukind Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Hemolytic Anemia
Required Medical Information and Criteria	<p><u>Hemolytic Anemia</u></p> <ul style="list-style-type: none">• Diagnosis of pyruvate kinase deficiency with at least two mutations within the PKLR gene, including a missense mutation• Confirmation of current hemoglobin $\leq 10\text{mg/dL}$• Patient is not homozygous for the R479H mutation• Patient does not have two non-missense variants in the PKLR gene, without the presence of another missense variant• Patient has had at least 6 RBC transfusions within the previous year for hemolytic anemia due to PKD• Prescriber confirmed concomitant use of daily folic acid• Confirmation that the patient does not have moderate or severe hepatic dysfunction.
Renewal Criteria	<ul style="list-style-type: none">• Clinical documentation showing an increase in Hb at least 1.5 mg/dL over baseline and/or a reduction in frequency of transfusions.
Age Restriction	<ul style="list-style-type: none">• 18 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• Hemolytic Anemia: Hematologist
Coverage Duration	<ul style="list-style-type: none">• Hemolytic Anemia:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 6 months

Effective Date:	8/01/2022
P&T Approval Date:	7/12/2022
P&T Revision Date:	7/12/2022

References

•

Mometasone Nasal Spray

Products Affected

- Mometasone Furoate Nasal Spray

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Nasal Congestion or Rhinitis
Required Medical Information and Criteria	<p><u>Nasal Congestion or Rhinitis</u></p> <ul style="list-style-type: none">• Diagnosis of asthma or an above the line comorbid condition that may worsen if not treated• Inadequate treatment response, intolerance, or contraindication to all the following<ul style="list-style-type: none">○ Fluticasone nasal spray○ Budesonide nasal spray○ Triamcinolone nasal spray
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	12/1/2022
P&T Approval Date:	11/8/2022
P&T Revision Date:	11/8/2022

References
•

Naltrexone (Vivitrol)

Products Affected

- Vivitrol Injection

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<u>All Diagnoses</u> <ul style="list-style-type: none">○ The drug will be dispensed directly to the provider and not the member.
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	7/12/2022

References
<ul style="list-style-type: none">•

Nasal Steroids

Products Affected

- Budesonide 32 mcg/act nasal spray
- Triamcinolone 55 mcg/act nasal spray

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Fluticasone 50 mcg/act nasal spray
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Nitroglycerin ointment

Products Affected

- Nitroglycerin 0.4% ointment

Covered Uses	<ul style="list-style-type: none">All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">Anal Fissure
Required Medical Information and Criteria	<p><u>Anal Fissure</u></p> <ul style="list-style-type: none">Diagnosis of moderate to severe pain associated with chronic anal fissure.
Renewal Criteria	<ul style="list-style-type: none">N/A
Prescriber Restriction	<ul style="list-style-type: none">None
Coverage Duration	<ul style="list-style-type: none">Anal Fissure:<ul style="list-style-type: none">Initial: 2 monthsRenewal: N/A

Effective Date:	9/1/2025
P&T Approval Date:	7/8/2025
P&T Revision Date:	7/8/2025

References
<ul style="list-style-type: none">Davids JS, et al. Clinical Practice Guidelines. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Anal Fissures. <i>Dis Colon Rectum</i> 2022; 66: 190–199.Wald A, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. <i>Am J Gastroenterol</i> 2021;116:1987–2008.

Non-Formulary Criteria

Products Affected

- All Non-Formulary Medications

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All FDA Approved Conditions</u></p> <ul style="list-style-type: none"> • One of the following: <ul style="list-style-type: none"> ○ The requested medication is being used to treat an above the line diagnosis on the OHA Prioritized list ○ The member has an above the line comorbid condition that will be treated indirectly by the requested medication ○ The member under the age of 21 • One of the following: <ul style="list-style-type: none"> ○ The requested medication is being used for an FDA-approved indication: ○ The requested medication is being used for an off-label indication with well-established, clinical evidence supporting its use. • All appropriate formulary alternatives have been tried and failed or are contraindicated. <p><u>Non-Formulary alternative dosage form (e.g. a request for NF oral solution when tablets are on formulary)</u></p> <ul style="list-style-type: none"> • One of the following: <ul style="list-style-type: none"> ○ Member is age 10 or younger ○ Member has clinically documented inability to swallow tablets/capsules • One of the following: <ul style="list-style-type: none"> ○ The formulary equivalent tablet/capsule does not require PA ○ The formulary equivalent tablet/capsule requires PA and the PA criteria has been met
Renewal Criteria	<ul style="list-style-type: none"> • Clinical documentation of follow-up indicating safety, efficacy and medication adherence over previous approval duration. • There are not any newly added formulary alternatives, or newly added formulary alternatives have been tried and failed or are contraindicated.

Age Restriction	<ul style="list-style-type: none"> • FDA labeled limits
Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Appropriate Specialist will vary by drug and condition
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months or shorter if appropriate for the drug/condition ○ Renewal: 12 months or shorter if appropriate for the drug/condition

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none"> •

Omalizumab (Xolair)

Products Affected

- Xolair Prefilled Syringe
- Xolair Auto-Injector

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Moderate to Severe Asthma ○ Chronic Rhinosinusitis with Nasal Polyps ○ Chronic Idiopathic Urticaria - Refractory ○ IgE-Mediated Food Allergy
Required Medical Information and Criteria	<p><u>Severe Asthma</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of moderate to severe persistent asthma. • Positive skin test or RAST to a perennial aeroallergen. • Baseline IgE serum level within FDA label. • Documentation of steps taken to avoid, within reason, environmental allergens and other triggers environmental allergens and other triggers. • Documented trial and failure, with claims history of adherence to: <ul style="list-style-type: none"> ○ High dose inhaled corticosteroid with a long-acting beta agonist (e.g., Advair), ○ Long acting anti-muscarinic (e.g., Spiriva), ○ Leukotriene Inhibitor (e.g., Singulair). • Documented trial and failure of, or contraindication to allergen immunotherapy. <p>Chronic Rhinosinusitis with Nasal Polyps (CRSwNP):</p> <ul style="list-style-type: none"> • Confirmed diagnosis of chronic rhinosinusitis with nasal polyps (CRSwNP). • Documentation of recurrent nasal polyps after prior sinus surgery. • Documented risk of another sinus surgery, or a statement why sinus surgery is not medically appropriate. • Documented trial and failure, with claims history of adherence to: <ul style="list-style-type: none"> ○ At least 2 intranasal corticosteroids (e.g., fluticasone, mometasone), ○ Sinuva. • Documentation that Xolair is intended as adjunct therapy with nasal corticosteroids.

	<p><u>Chronic Idiopathic Urticaria - refractory (CIU)</u></p> <ul style="list-style-type: none"> • Documentation of chronic spontaneous or idiopathic urticaria. • Age under 21, or a comorbid condition which would make chronic urticaria coverable under the prioritized list. • Documented trial and failure of at least 6 weeks of maximally tolerated doses of all the following: <ul style="list-style-type: none"> ○ 1st generation antihistamine – (e.g., doxepin, hydroxyzine) ○ 2nd generation antihistamine – (e.g., cetirizine, levocetirizine, fexofenadine, loratadine, desloratadine) ○ Histamine Type-2 Receptor Antagonists (e.g., famotidine, cimetidine) ○ Leukotriene inhibitor (e.g., montelukast, zafirlukast) <p><u>IgE-Mediated Food Allergy</u></p> <ul style="list-style-type: none"> • Diagnosis of IgE Mediated Food Allergy as evidenced by one of the following: <ul style="list-style-type: none"> ○ Positive skin prick test (defined as greater than or equal to 4 mm wheal greater than saline control) to food, ○ Positive food specific IgE (greater than or equal to 6 kUA/L), ○ Positive oral food challenge, defined as experiencing dose-limiting symptoms at a single dose of less than or equal to 300 mg of food protein. • Clinical history of IgE mediated food allergy. • Used in conjunction with food allergen avoidance. • Both of the following: <ul style="list-style-type: none"> ○ Baseline (pre-Xolair treatment) serum total IgE level is greater than or equal to 30 IU/mL and less than or equal to 1850 IU/mL, ○ Dosing is according to serum total IgE levels and body weight. • Xolair will not be used concomitantly with Palforzia. • Attestation that the member is co-prescribed epinephrine or has epinephrine at home.
Renewal Criteria	<p><u>IgE Mediated Food Allergy:</u></p> <ul style="list-style-type: none"> • Patient demonstrates positive clinical response to therapy (e.g., reduction of type 1 allergic reactions, including anaphylaxis, following accidental exposure to one or more foods). • Used in conjunction with food allergen avoidance.

	<ul style="list-style-type: none"> Dosing will continue to be based on body weight and pretreatment IgE serum levels. <p>All Other Diagnoses:</p> <ul style="list-style-type: none"> Documentation of clinically significant improvement in symptoms.
Exclusion Criteria	<ul style="list-style-type: none"> Specific criteria we want reviewed. This should be blank in many cases. Do not include things like “allergy to the drug you are asking for”; this is obvious and doesn’t need to be part of the criteria.
Age Restriction	<ul style="list-style-type: none"> Asthma: 6 years of age and older CIU: 12 years of age and older CRSwNP: 18 years of age and older IgE Mediated Food Allergy: 1 year of age and older
Prescriber Restriction	<ul style="list-style-type: none"> Asthma: Prescribed by or in consultation with a pulmonologist or immunologist. CIU: Prescribed by or in consultation with an immunologist. CRSwNP: Prescribed by or in consultation with an allergist or ENT. IgE Mediated Food Allergy: Prescribed by or in consultation with an allergist or immunologist.
Coverage Duration	<ul style="list-style-type: none"> Asthma: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months CRSwNP: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months CIU: <ul style="list-style-type: none"> Initial: 4 months Renewal: 6 months IgE Mediated Food Allergy: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months

Effective Date:	9/1/2024
-----------------	----------

P&T Approval Date:	7/9/2024
P&T Revision Date:	7/9/2024

References
•

General Oncology (Chemotherapy)

Products Affected

- Abemaciclib (Verzenio)
- Abiraterone
- Acalabrutinib (Calquence)
- Adagrasib (Krazati)
- Alectinib (Alecensa)
- Alpelisib (Piqray)
- Ascinimib (Scemblix)
- Asparaginase Erwinia (Rylaze)
- Avutometinib/Defactinib (Avmapi/Fazynja Pak)
- Belzutifan (Welireg)
- Binimetinib (Mektovi)
- Bosutinib (Bosulif)
- Brigatinib (Alunbrig)
- Busulfan (Myleran)
- Cabozantinib (Cabometyx)
- Capivasertib (Truqap)
- Capmatinib (Tabrecta)
- Ceritinib (Zykadia)
- Chlorambucil (Leukeran)
- Crizotinib (Xalkori)
- Dabrafenib (Tafinlar)
- Dasatinib
- Elacestrant (Orserdu)
- Encorafenib (Braftovi)
- Entrectinib (Rozlytrek)
- Erlotinib
- Estramustine (Emcyt)
- Everolimus
- Fruquintinib (Fruzaqla)
- Futibatinib (Lytgobi)
- Gefitinib
- Ibrutinib (Imbruvica)
- Inavolisib (Itovebi)
- Infigratinib (Truseltiq)
- Ivosidenib (Tibsovo)
- Lazertinib (Lazcluze)
- Lomustine (Gleostine)
- Lorlatinib (Lobrena)
- Midostaurin (Rydapt)
- Mobocertinib (Exkivity)
- Nilotinib (Tasigna)
- Niraparib/Abiraterone (Akeega)
- Nirogacestat (Ogsiveo)
- Olaparib (Lynparza)
- Olutasidenib (Rezlidhia)
- Osimertinib (Tagrisso)
- Pacritinib (Vonjo)
- Palbociclib (Ibrance)
- Pazopanib
- Pemigatinib (Pemazyre)
- Pexidartinib (Turalio)
- Pirtobrutinib (Jaypirca)
- Ponatinib (Iclusig)
- Quizartinib (Vanflyta)
- Repotrectinib (Augtyro)
- Ribociclib (Kisqali)
- Sargramostim (Leukine)
- Selpercatinib (Retevmo)
- Sorafenib
- Sotorasib (Lumakras)
- Sunitinib
- Talazoparib (Talzenna)
- Temozolomide
- Tepotinib (Tepmetko)
- Tivozanib (Fotivda)
- Topotecan (Hycamtin)
- Tovorafenib (Ojemda)
- Trametinib (Mekinist)
- Trifluridine/Tipiracil (Lonsurf)
- Tucatinib (Tukysa)
- Vimseltinib (Romvimza)
- Vorasidinib (Voranigo)

- Lenvatinib (Lenvima)
- Vorinostat (Zolinza)
- Zanubrutinib (Brukinsa)

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> • One of the following is true: <ul style="list-style-type: none"> ○ The requested drug is being used for an FDA approved indication. ○ The requested medication is being used according to National Comprehensive Cancer Network (NCCN) guidelines.
Renewal Criteria	<ul style="list-style-type: none"> • Submission of clinical documentation supporting provider follow-up that indicates safety and efficacy of the medication and adherence to treatment.
Age Restriction	<ul style="list-style-type: none"> • Refer to FDA indication and NCCN guidelines
Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Oncologist or hematologist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: up to 6 months

Effective Date:	9/1/2025
P&T Approval Date:	10/01/2022
P&T Revision Date:	7/8/2025, 5/13/2025, 3/11/2025, 5/14/2024, 03/01/2024, 01/09/2024, 11/1/2023, 09/01/2023, 7/11/2023, 05/09/2023, 03/14/2023, 01/10/2023, 10/01/2022

Oral Antifungal Agents

Products Affected

- Griseofulvin Ultramicrosize Tabs
- Griseofulvin Microsize Susp
- Ketoconazole Tabs

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> • The medication is being used for an FDA-approved indication OR being used for an off-label indication with well-established, clinical evidence supporting its use, including the dose being requested • For suspension only: <ul style="list-style-type: none"> ○ Is there a reason that a tablet form can't be used
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy • Evidence of continued need for the medication
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal (non-immunocompromised): 3 months ○ Renewal (immunocompromised): 6 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References
<ul style="list-style-type: none"> • Ketoconazole tablets [package insert]. Toronto, Ontario: Apotex Inc.; 2021. • Griseofulvin [package insert]. Bridgewater, GA: Valent LLC.; 2016.

Oral Nutrition Supplements

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> Documentation showing the prescribed oral nutritional formula and/or nutritional supplements are an integral part of treatment/medically necessary for a nutritional deficiency. Documentation including assessment by treating practitioner or registered dietitian that member is unable to meet their recommended caloric/protein or micronutrient needs through regular, liquified, blenderized, or pureed foods in any modified texture or form. <p>OR</p> <ul style="list-style-type: none"> For age under 21 <ul style="list-style-type: none"> Documented delayed growth or failure to thrive. Documentation showing the prescribed formula/nutritional supplement is for the prevention of nutritional deficiency or malnutrition.
Renewal Criteria	<ul style="list-style-type: none"> Documentation of continued positive response for the requested enteral nutrition/formula with a continued need for requested supplement.
Exclusion Criteria	<ul style="list-style-type: none"> Supplement is to be administered via enteral tube feeding (e.g. G-tube, NG-tube). For tube feedings please submit via a DME vendor through the DME benefit
Coverage Duration	<ul style="list-style-type: none"> All Diagnoses: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References
•

Pancrelipase (Creon, Pancreaze)

Products Affected

- Creon Capsules
- Pancreaze Capsules

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Exocrine Pancreatic Insufficiency
Required Medical Information and Criteria	<p><u>Exocrine Pancreatic Insufficiency</u></p> <ul style="list-style-type: none"> • Clinical documentation of one of the following: <ul style="list-style-type: none"> ○ Confirmed diagnosis of cystic fibrosis ○ History of pancreatectomy ○ Diagnosis of exocrine pancreatic cancer ○ Diagnosis of chronic pancreatitis confirmed by imaging ○ Confirmed diagnosis of pancreatic insufficiency confirmed with one of the following methods: <ul style="list-style-type: none"> ▪ Steatorrhea with fecal fat determination ▪ Measurement of fecal elastase ▪ Secretin or CCK pancreatic function testing ▪ Two of the following CFTR mutations (G542X, W1282X, R553X, 621+1G>T, 1717-1G>A, 3120+1G>A, R1162X, 3659delC, 1898+1G>A, 2184delA, 711+1G>T, F508del, I507del, G551D, N1303K, R560T)
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> • Exocrine Pancreatic Insufficiency: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	2/1/2022
P&T Approval Date:	1/11/2022
P&T Revision Date:	1/11/2022

References
•

Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) Inhibitors

Products Affected

- Repatha SureClick
- Repatha Prefilled Syringe
- Repatha Pushtronex
- Praluent Auto-Injector

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Established clinical ASCVD Primary or Familial Hyperlipidemia
Required Medical Information and Criteria	<p><u>Established Clinical ASCVD</u></p> <ul style="list-style-type: none"> Documentation of very high risk ASCVD as evidenced by either: <ul style="list-style-type: none"> History of multiple major ASCVD events One major ASCVD event AND multiple high-risk conditions. Documentation of a current LDL greater than or equal to 55 mg/dl. Documentation of all the following: <ul style="list-style-type: none"> Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins Is receiving ezetimibe or has a documented intolerance to ezetimibe. <p><u>Primary or Familial Hyperlipidemia</u></p> <ul style="list-style-type: none"> Documentation of an untreated (i.e., prior to lipid lowering therapy) LDL greater than 190 mg/dL. Documentation of current LDL greater than 100 mg/dL. Documentation of all the following: <ul style="list-style-type: none"> Patient is receiving maximally tolerated statin therapy (atorvastatin 40-80mg, rosuvastatin 20-40mg) or has a documented clinical intolerance to statins Is receiving ezetimibe or has a documented intolerance to ezetimibe.
Renewal Criteria	<ul style="list-style-type: none"> Documented positive clinical response to therapy (significant decrease in lipid levels).

Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Cardiologist, Endocrinologist, or Lipid Specialist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024, 1/11/2022

References
•

Peanut Powder (Palforzia)

Products Affected

- Palforzia Capsule and Powder

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Peanut Allergy
Required Medical Information and Criteria	<p><u>Peanut Allergy</u></p> <ul style="list-style-type: none">• Confirmed positive skin test or peanut-specific serum IgE greater than 0.35 kUA/L• Concurrent prescription with injectable epinephrine• Medical justification supports necessity for oral immunotherapy despite peanut avoidance.
Renewal Criteria	<ul style="list-style-type: none">• Currently receiving medication by way of previously approved SHP authorization or documents showing Initial approval criteria was or has been met.• For patients who required use of injectable epinephrine while on Palforzia, must have medical justification that supports continued need for Palforzia.• If 18 years or older, must have medical justification that supports continued need for oral immunotherapy despite peanut avoidance and documentation that Initial dose escalation happened between age 4 and 17.
Age Restriction	<ul style="list-style-type: none">• Patient must be between 4 and 17 at therapy initiation
Prescriber Restriction	<ul style="list-style-type: none">• Peanut Allergy: Allergist or Immunologist enrolled in Palforzia REMS program
Coverage Duration	<ul style="list-style-type: none">• Peanut Allergy:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Phosphodiesterase Type 5 (PDE5) inhibitors

Products Affected

- Sildenafil 20mg tablet
- Tadalafil 20mg tablet

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Pulmonary Arterial Hypertension
Required Medical Information and Criteria	<u>Pulmonary Arterial Hypertension</u> <ul style="list-style-type: none"> • Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension)
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Exclusion Criteria	<ul style="list-style-type: none"> • Treatment of erectile dysfunction
Age Restriction	<ul style="list-style-type: none"> • Tadalafil: 18 years of age and over
Prescriber Restriction	<ul style="list-style-type: none"> • Pulmonary Arterial Hypertension: Cardiologist or Pulmonologist
Coverage Duration	<ul style="list-style-type: none"> • Pulmonary Arterial Hypertension: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	7/13/2021
P&T Revision Date:	7/8/2025, 7/13/2021

References
<ul style="list-style-type: none"> • Adcirca (tadalafil). Prescribing information. Eli Lilly and Co. Indianapolis, IN 2020. • Revatio (sildenafil). Prescribing information. Viatris Specialty LLC. Morgantown, WV 2024

Phytonadione (Vitamin K) Step Therapy

Products Affected

- Phytonadione 5mg tab

Step Therapy Criteria	<ul style="list-style-type: none">• Concurrent use of warfarin
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Pitolisant (Wakix)

Products Affected

- Wakix Tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Cataplexy and Narcolepsy ○ Excessive Somnolence - Narcolepsy
Required Medical Information and Criteria	<p><u>Cataplexy and Narcolepsy</u></p> <ul style="list-style-type: none"> • Confirmation of diagnosis of narcolepsy based on both: <ul style="list-style-type: none"> ○ Polysomnography ○ A multiple sleep latency test • Documentation that CYP2D6 testing has been done, and the dosing will be adjusted if the patient is a poor metabolizer. • Diagnosis of Cataplexy confirmed by a specialist. • Trial and failure or contraindication to ALL the following: <ul style="list-style-type: none"> ○ SSRI antidepressant (e.g. fluoxetine) ○ SNRI antidepressant (e.g. venlafaxine and duloxetine) ○ Tricyclic antidepressant (e.g. clomipramine) ○ Sodium oxybate product titrated to maximally tolerated dose. <p><u>Excessive Somnolence - Narcolepsy</u></p> <ul style="list-style-type: none"> • Confirmation of diagnosis of narcolepsy based on both: <ul style="list-style-type: none"> ○ Polysomnography ○ A multiple sleep latency test • Documentation that CYP2D6 testing has been done, and the dosing will be adjusted if the patient is a poor metabolizer. • Documentation of fatigue severity using a validated measure (e.g. Epworth score, Brief Fatigue Inventory, or other validated measure) • Trial and failure or contraindication to ALL the following: <ul style="list-style-type: none"> ○ Modafinil (at least 200mg dose) AND armodafinil ○ Mixed amphetamine salts, methylphenidate or dexamethylphenidate, AND dextroamphetamine ○ Sunosi (solriamfetol) ○ A sodium oxybate product
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy

Exclusion Criteria	<ul style="list-style-type: none"> • Severe renal or hepatic impairment • Pregnant or actively trying to conceive
Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Sleep Specialist or Neurologist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 3 months ○ Renewal: 6 months

Effective Date:	3/1/2025
P&T Approval Date:	1/14/2025
P&T Revision Date:	1/14/2025

References
•

Potassium Binders

Products Affected

- Lokelma Powder
- Veltassa Packet

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Hyperkalemia
Required Medical Information and Criteria	<p><u>Hyperkalemia</u></p> <ul style="list-style-type: none">• Being used for the treatment of hyperkalemia based on current potassium labs• Failed all the following:<ul style="list-style-type: none">○ Dietary changes (potassium restriction)○ Dose modification/discontinuation of ACE-inhibitor, ARB, or other hyperkalemia causing agents○ Diuretics titrated to maximum tolerated dose• If the request is for Veltassa:<ul style="list-style-type: none">○ Trial and failure or contraindication to Lokelma
Renewal Criteria	<p>Documented positive clinical response to therapy by either of the following:</p> <ul style="list-style-type: none">• Potassium level returning to normal on therapy• Significant drop in potassium level from baseline on therapy
Coverage Duration	<ul style="list-style-type: none">• Hyperkalemia:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	9/13/2022

References
•

Pretomanid

Products Affected

- Pretomanid tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Pulmonary tuberculosis
Required Medical Information and Criteria	<p><u>Pulmonary tuberculosis</u></p> <ul style="list-style-type: none">• Evidence of extensively drug-resistant active pulmonary tuberculosis (XDR-TB) caused by mycobacterium tuberculosis.<ul style="list-style-type: none">○ XDR-TB is defined as TB that is resistant to rifampicin and isoniazid, at least one fluoroquinolone (levofloxacin or moxifloxacin) and a second-line injectable (amikacin, capreomycin, and kanamycin) OR Isoniazid, rifampin a fluoroquinolone AND bedaquiline or linezolid.• Pretomanid is prescribed as part of a guideline recommended multi-drug treatment regimen.
Renewal Criteria	<ul style="list-style-type: none">• N/A
Exclusion Criteria	<ul style="list-style-type: none">• Use outside of recognized treatment guidelines.• Medication is being received through a county clinic with a state funded TB program.
Age Restriction	<ul style="list-style-type: none">• Age of 14 or greater.
Prescriber Restriction	<ul style="list-style-type: none">• Infectious Disease
Coverage Duration	<ul style="list-style-type: none">• Pulmonary tuberculosis: 24 weeks

Effective Date:	5/1/2025
P&T Approval Date:	3/11/2025
P&T Revision Date:	3/11/2025

References

- Pretomanid [package insert]. Morgantown, WV: Mylan Specialty L.P.; 2024.

Prostacyclin Agonists

Products Affected

- Orenitram tablets
- Treprostinil
- Remodulin
- Tyvaso

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Pulmonary Arterial Hypertension
Required Medical Information and Criteria	<p><u>Pulmonary Arterial Hypertension</u></p> <ul style="list-style-type: none"> • Clinically documented diagnosis of Pulmonary Arterial Hypertension (WHO group 1 pulmonary hypertension) • Clinical documentation of WHO functional class III or IV • Evidence of an unfavorable response or intolerance to all of the following: <ul style="list-style-type: none"> ○ Phosphodiesterase Type 5 inhibitor (sildenafil, tadalafil) ○ Endothelin Receptor Antagonist (ambrisentan, bosentan) ○ Combination therapy with a Phosphodiesterase Type 5 inhibitor + Endothelin Receptor Antagonist (ambrisentan + tadalafil) • The requested medication will be an add on to an already established first line agent or agents (sildenafil, tadalafil, ambrisentan, bosentan)
Renewal Criteria	<ul style="list-style-type: none"> • Documentation of positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none"> • Age 16 or older
Prescriber Restriction	<ul style="list-style-type: none"> • Pulmonary Arterial Hypertension: Cardiologist or Pulmonologist
Coverage Duration	<ul style="list-style-type: none"> • Pulmonary Arterial Hypertension: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	7/13/2021
P&T Revision Date:	7/8/2025, 7/13/2021

References

- Orenitram [package insert]. Research Triangle Park, NC: United Therapeutics Corp; 2023.
- Remodulin [package insert]. Research Triangle Park, NC: United Therapeutics Corp; 2023.
- Tyvaso [package insert]. Research Triangle Park, NC: United Therapeutics Corp; 2022.

Quantity Limit Exception Criteria

Products Affected

- All Drugs with Quantity Limits

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All FDA Approved Conditions</u></p> <ul style="list-style-type: none"> • One of the following: <ul style="list-style-type: none"> ○ The requested medication is being used to treat an above the line diagnosis on the OHA Prioritized list ○ The member has an above the line comorbid condition that will be treated indirectly by the requested medication ○ The member under the age of 21 • One of the following: <ul style="list-style-type: none"> ○ The requested medication is being used for an FDA-approved indication: ○ The requested medication is being used for an off-label indication with well-established, clinical evidence supporting its use. • Documentation of failure of the requested medication within quantity limits, including failure of different strengths of the requested medication. • All appropriate formulary alternatives have been tried and failed or are contraindicated.
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Exclusion Criteria	<ul style="list-style-type: none"> • Specific criteria we want reviewed. This should be blank in many cases. Do not include things like “allergy to the drug you are asking for”; this is obvious and doesn’t need to be part of the criteria.
Age Restriction	<ul style="list-style-type: none"> • FDA labeled limits
Prescriber Restriction	<ul style="list-style-type: none"> • All Diagnoses: Appropriate Specialist will vary by drug and condition
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses:

	<ul style="list-style-type: none"> ○ Initial: 6 months or shorter if appropriate for the drug/condition ○ Renewal: 12 months or shorter if appropriate for the drug/condition
--	---

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
•

Resmetirom (Rezdiffra)

Products Affected

- Rezdiffra Tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Metabolic Dysfunction-Associated Steatohepatitis
Required Medical Information and Criteria	<p><u>Metabolic Dysfunction-Associated Steatohepatitis</u></p> <ul style="list-style-type: none"> • Clinical documentation confirming a diagnosis of metabolic dysfunction-associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH) • Patient does not have cirrhosis (e.g. decompensated cirrhosis) • Submission of medical records (e.g. chart notes) showing diagnosis has been confirmed by one of the following: <ul style="list-style-type: none"> ○ FibroScan-aspartate aminotransferase (FAST) ○ MRI-aspartate aminotransferase (MAST) ○ Liver biopsy • Submission of medical records (e.g. chart notes) showing disease is fibrosis stage F2 or F3 as confirmed by one of the following: <ul style="list-style-type: none"> ○ FibroScan ○ Fibrosis-4 index (FIB-4) ○ Magnetic resonance Elastography (MRE) • Presence of greater than or equal to 3 metabolic risk factors (e.g., Type 2 diabetes, hypertension, obesity) • Submission of medical records (e.g. chart notes) confirming drug is used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program)
Renewal Criteria	<ul style="list-style-type: none"> • Patient demonstrates positive response to therapy (e.g., MASH resolution, fibrosis stage improvement, etc.) • Submission of medical records (e.g., chart notes) confirming drug will continue to be used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community-based program)
Age Restriction	<ul style="list-style-type: none"> • ≥18 years of age or older
Prescriber Restriction	<ul style="list-style-type: none"> • Metabolic Dysfunction-Associated Steatohepatitis: Gastroenterologist, Hepatologist

Coverage Duration	<ul style="list-style-type: none"> • Metabolic Dysfunction-Associated Steatohepatitis: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months
--------------------------	---

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	5/14/2024

References
•

Rifapentine (Priftin)

Products Affected

- Priftin tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Latent tuberculosis○ Active tuberculosis
Required Medical Information and Criteria	<p><u>Latent tuberculosis</u></p> <ul style="list-style-type: none">• Used in combination with isoniazid (INH). <p><u>Active tuberculosis</u></p> <ul style="list-style-type: none">• Used as part of a multi-drug regimen.
Renewal Criteria	<ul style="list-style-type: none">• N/A
Exclusion Criteria	<ul style="list-style-type: none">• Medication is being received through a county clinic with a state funded TB program.
Age Restriction	<ul style="list-style-type: none">• Age \geq 2 years old with latent TB• Age \geq 12 years old with active TB
Prescriber Restriction	<ul style="list-style-type: none">• Latent TB: Not limited by specialty• Active TB: Infectious disease specialist required for multidrug resistant cases only
Coverage Duration	<ul style="list-style-type: none">• Latent TB: 3 months• Active TB: 6 months

Effective Date:	5/1/2025
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/11/2025, 3/12/2024

References
<ul style="list-style-type: none">• Priftin [package insert]. Bridgewater, NJ: Sanofi-Aventis.; 2021.

Rifaximin (Xifaxan)

Products Affected

- Xifaxan 550mg tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Hepatic Encephalopathy
Required Medical Information and Criteria	<p><u>Hepatic Encephalopathy</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of hepatic encephalopathy• One of the following:<ul style="list-style-type: none">○ Xifaxan will be used as add-on therapy to lactulose and there is documentation of failure to achieve optimal clinical response with lactulose alone○ Documentation of contraindication or intolerance to lactulose
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• Hepatic Encephalopathy:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Risdiplam (Evrysdi)

Products Affected

- Evrysdi Solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Spinal Muscular Atrophy
Required Medical Information and Criteria	<p><u>Spinal Muscular Atrophy</u></p> <ul style="list-style-type: none">• Confirmed (via genetic testing) diagnosis of 5q-autosomal recessive SMA (type 1, 2 or 3)• Patient is not dependent on invasive ventilation or tracheostomy or use of non-invasive ventilation beyond uses for sleeping• Patient is not receiving concomitant chronic SMN modifying therapy such as Spinraza• Patient has not previously received gene replacement therapy for the treatment of SMA (e.g. Zolgensma)
Renewal Criteria	<ul style="list-style-type: none">• Documentation of clinical improvement from baseline in motor functionality confirmed by standard exams (e.g. BSID-III, CHOP INTEND, HINE-2, RULM test)
Prescriber Restriction	<ul style="list-style-type: none">• Spinal Muscular Atrophy: Neurologist with expertise in the treatment of spinal muscular atrophy
Coverage Duration	<ul style="list-style-type: none">• Spinal Muscular Atrophy:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
<ul style="list-style-type: none">•

Roflumilast

Products Affected

- Product 1
- Product 2
- Product 3
- Etc.

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Chronic Obstructive Pulmonary Disorder (COPD)
Required Medical Information and Criteria	<u>Chronic Obstructive Pulmonary Disorder (COPD)</u> <ul style="list-style-type: none">• Diagnosis of moderate to severe COPD and chronic bronchitis• Trial and failure of at least 2 previous treatments for COPD
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to roflumilast therapy.
Exclusion Criteria	<ul style="list-style-type: none">• Specific criteria we want reviewed. This should be blank in many cases. Do not include things like “allergy to the drug you are asking for”; this is obvious and doesn’t need to be part of the criteria.
Coverage Duration	<ul style="list-style-type: none">• COPD:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal:12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Sacubitril/Valsartan (Entresto)

Products Affected

- Entresto tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Heart Failure
Required Medical Information and Criteria	<p><u>Heart Failure</u></p> <ul style="list-style-type: none">• Diagnosis of New York Heart Association class II to IV heart failure• The patient meets one of the following:<ul style="list-style-type: none">○ Receiving concomitant therapy with one of the following beta blockers: carvedilol, bisoprolol, sustained release metoprolol OR○ Beta-blockers are contraindicated or the member has been unable to tolerate them• The patient will discontinue use of any ACE inhibitor or ARB before initiating therapy with Entresto.
Renewal Criteria	<ul style="list-style-type: none">• Documented positive response to therapy.
Prescriber Restriction	<ul style="list-style-type: none">• Cardiologist, or in consultation with a cardiologist
Coverage Duration	<ul style="list-style-type: none">• Heart Failure:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	9/1/2025
P&T Approval Date:	4/25/2016
P&T Revision Date:	7/8/2025

References

- Entresto [package insert]. East Hanover, NJ: Novartis Pharmaceuticals; 2024.

Secukinumab (Cosentyx)

Products Affected

- Cosentyx Pen 300mg Dose
- Cosentyx Prefilled Syringe 300mg Dose

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Hidradenitis Suppurativa ○ Juvenile Idiopathic Arthritis ○ Plaque Psoriasis ○ Psoriatic Arthritis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla. <p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of infliximab and adalimumab

	<p><u>Hidradenitis Suppurativa (HS)</u></p> <ul style="list-style-type: none"> • Documentation of a diagnosis of moderate to severe hidradenitis suppurativa (Hurley Stage II or Hurley Stage III) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy (e.g. oral antibiotics). • Trial and failure of infliximab and adalimumab <p><u>Juvenile Idiopathic Arthritis (JIA)</u></p> <ul style="list-style-type: none"> • Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids • Documented intolerance or contraindication to DMARDs OR DMARD will be continued with Cosentyx • Trial and failure of infliximab and adalimumab <p><u>Plaque Psoriasis (PP):</u></p> <ul style="list-style-type: none"> • Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involved ○ Hand, foot, face, or mucous membrane involvement • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR
--	--

	<ul style="list-style-type: none"> ○ Documented trial and failure of all the following: <ul style="list-style-type: none"> ▪ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ▪ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ▪ PUVA or UVB Phototherapy ▪ Methotrexate ▪ At least 1 other second line systemic agent such as cyclosporine or acitretin. • Trial and failure of infliximab and adalimumab <p><u>Psoriatic Arthritis (PsA):</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of infliximab and adalimumab
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Juvenile Idiopathic Arthritis: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • Plaque Psoriasis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of

	<p>body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement.</p> <ul style="list-style-type: none"> • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count.
Prescriber Restriction	<ul style="list-style-type: none"> • Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Juvenile Idiopathic Arthritis: Rheumatologist. • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	9/1/2023, 07/11/2023, 01/11/2022

References
•

Seladelpar (Livdelzi)

Products Affected

- Livdelzi Capsules

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Primary Biliary Cholangitis
Required Medical Information and Criteria	<p><u>Primary Biliary Cholangitis</u></p> <ul style="list-style-type: none">• Diagnosis of primary biliary cholangitis (PBC) confirmed by two of the following:<ul style="list-style-type: none">○ Biochemical evidence of cholestasis based on ALP elevation○ Presence of AMA or other PBC-specific autoantibodies○ Histology confirmation after biopsy• Documentation of at least 12 months of inadequate response to ursodiol• No current decompensated cirrhosis
Renewal Criteria	<ul style="list-style-type: none">• Documented adherence to medication regimen and clinical benefit
Age Restriction	<ul style="list-style-type: none">• Age 18 or older
Prescriber Restriction	<ul style="list-style-type: none">• Primary Biliary Cholangitis: Hepatologist or Gastroenterologist
Coverage Duration	<ul style="list-style-type: none">• Primary Biliary Cholangitis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References
•

Semaglutide (Wegovy)

Products Affected

- Wegovy Injection

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Secondary Prevention of Major Adverse Events (MACE) ○ Obesity or Overweight (only applies to members under age 21)
Required Medical Information and Criteria	<p><u>Secondary Prevention of Major Adverse Events</u></p> <ul style="list-style-type: none"> • Wegovy is being used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) in adults with established cardiovascular disease and either obesity or overweight. • Wegovy is being used as adjunct to lifestyle modification (e.g., dietary, or caloric restriction, exercise, behavioral support, community-based program). • Patient has established cardiovascular disease as evidenced by one of the following: <ul style="list-style-type: none"> ○ Prior myocardial infarction ○ Prior stroke ○ Peripheral arterial disease (e.g., intermittent claudication with ankle-brachial index <0.85, peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease) • BMI greater than or equal to 27 kg/m² <p><u>Obesity or Overweight</u></p> <ul style="list-style-type: none"> • Age 20 or younger • BMI at or above the 95th percentile or 27kg/m². • Documentation of one of the following: <ul style="list-style-type: none"> ○ Comorbidities (e.g., hypertension, dyslipidemia, fatty liver disease, depression, or sleep apnea). ○ Trial and failure of at least 3 months of a diet/exercise plan administered by a health care provider in the last 6 months. <p>The patient is, or will be, engaged in a weight management lifestyle modification program in addition to pharmacotherapy.</p>
Renewal Criteria	<p><u>Secondary Prevention of Major Adverse Events (MACE):</u></p> <ul style="list-style-type: none"> • Documentation of treatment success (BMI reduction of 5% or

	<p>more).</p> <ul style="list-style-type: none"> Documentation of continuation of lifestyle modification program with reduced calorie diet and regular physical activity alongside continuous Wegovy use (80% adherence). <p>Obesity or overweight:</p> <ul style="list-style-type: none"> Patient is less than 21 years old. Documentation of at least a 1% decrease in BMI from baseline. Patient is continuing full weight loss plan (e.g., diet and exercise program, nutritional counseling).
Exclusion Criteria	<ul style="list-style-type: none"> Coverage for weight loss is excluded for members age 21 or older
Age Restriction	<ul style="list-style-type: none"> Secondary Prevention of Major Adverse Events (MACE): 12 years or older Obesity or overweight: Age 12 to under 21 years of age
Prescriber Restriction	<ul style="list-style-type: none"> Secondary Prevention of Major Adverse Events (MACE): Cardiologist Diagnosis 2: Specialist 2, etc. Etc.
Coverage Duration	<ul style="list-style-type: none"> Secondary Prevention of Major Adverse Events (MACE): <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months Obesity or overweight: <ul style="list-style-type: none"> Initial: 6 months Renewal: 12 months or until age 21, whichever is less

Effective Date:	9/1/2024
P&T Approval Date:	7/9/2024
P&T Revision Date:	7/9/2024, 5/13/2024

References
•

Sirolimus Solution

Products Affected

- Sirolimus Oral Solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Age Restriction	<ul style="list-style-type: none">• Patient is under age 10 or unable to use tablets
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References
<ul style="list-style-type: none">•

Sodium-Glucose Co-Transporter 2 (Sglt2) Inhibitors

Products Affected

- Farxiga Tablets
- Dapagliflozin Tablets
- Steglatro Tablets
- Jardiance Tablets
- Invokana Tablets

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> Type 2 Diabetes Chronic Kidney Disease Heart Failure
Required Medical Information and Criteria	<p><u>Type 2 Diabetes</u></p> <ul style="list-style-type: none"> Trial and failure of, or contraindication to metformin Trial and failure of sulfonylurea or pioglitazone or contraindication to both <p><u>Chronic Kidney Disease</u></p> <ul style="list-style-type: none"> Concurrent therapy with an ACEi or ARB at maximum tolerated doses, or documented contraindication to both Stage 2, 3, or 4 CKD or eGFR of 25 to 75 mL/min/1.73 m² No previous use of dialysis No history of polycystic kidney disease, type 1 diabetes, lupus nephritis, or antineutrophil cytoplasmic antibody-associate vasculitis. <p><u>Heart Failure</u></p> <ul style="list-style-type: none"> One of the following: <ul style="list-style-type: none"> All the following: <ul style="list-style-type: none"> Diagnosis of heart failure (NYHA class II-IV) with reduced ejection fraction (HFrEF) Documentation of concurrent use of ACEi, ARB, or ARNI, or contraindication to all Documentation of concurrent use of carvedilol, metoprolol succinate, or bisoprolol, or contraindication to all Documented eGFR >30 mL/min/1.73m² All of the following: <ul style="list-style-type: none"> Diagnosis of heart failure with mildly reduced ejection

	fraction (HFmrEF), or heart failure with preserved ejection fraction (HFpEF) <ul style="list-style-type: none"> ▪ eGFR >30 mL/min/1.73m²
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: Lifetime

Effective Date:	9/1/2024
P&T Approval Date:	3/8/2022
P&T Revision Date:	7/7/2024, 3/8/2022

References
•

Somatropin (Humatrope)

Products Affected

- Humatrope Cartridge

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Growth Hormone Deficiency (GHD) ○ Prader-Willi Syndrome ○ Noonan Syndrome ○ Turner Syndrome ○ Idiopathic Short Stature ○ Growth Failure secondary to chronic kidney disease (CKD) ○ Small for gestational age ○ Short stature homeobox-containing (SHOX) gene deficiency ○ HIV Associated Cachexia
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> • Documentation of goals of therapy and objective baseline assessment (e.g., quality of life, exercise capacity, height, body composition) <p><u>Patient Under Age 18</u></p> <ul style="list-style-type: none"> • Confirmation of one of the following: <ul style="list-style-type: none"> ○ Growth Hormone Deficiency (GHD) as noted by one of the following: <ul style="list-style-type: none"> ▪ Growth hormone deficiency is confirmed by a negative response to a growth hormone stimulation test (e.g., serum GH levels of <5 ng/mL on stimulation testing with either of the following: glucagon or insulin) ▪ Patient has had the pituitary removed or destroyed or has had panhypopituitarism since birth ○ Diagnosis of Prader-Willi Syndrome and patient does not have concurrent severe obesity nor history of upper airway obstruction nor sleep apnea nor severe respiratory impairment ○ Diagnosis of Noonan Syndrome ○ Diagnosis of Turner Syndrome ○ Diagnosis of Idiopathic Short Stature ○ Diagnosis of Growth Failure secondary to chronic kidney disease (CKD) ○ Diagnosis of Small for gestational age

	<ul style="list-style-type: none"> ○ Diagnosis of Short stature homeobox-containing (SHOX) gene deficiency ○ Diagnosis of HIV Associated Cachexia <p><u>Patients Aged 18 and Older</u></p> <ul style="list-style-type: none"> • Confirmation of one of the following: <ul style="list-style-type: none"> ○ Confirmation of all the following: <ul style="list-style-type: none"> ▪ Growth Hormone Deficiency (GHD) as noted by one of the following: <ul style="list-style-type: none"> ▪ Growth hormone deficiency is confirmed by a negative response to a growth hormone stimulation test (e.g., serum GH levels of <5 ng/mL on stimulation testing with either of the following: glucagon or insulin) ▪ Patient has had the pituitary removed or destroyed or has had panhypopituitarism since birth ▪ The prescriber certifies that the growth hormone is not being prescribed for anti-aging therapy or to enhance athletic ability or body building ○ HIV associated cachexia ○ Short Bowel Syndrome (SBS)
Renewal Criteria	<ul style="list-style-type: none"> • Treatment with agent initiated in a patient prior to reaching adulthood (<18 years of age) was to improve growth velocity or height and one of the following: <ul style="list-style-type: none"> ○ Growth velocity greater than 2.5 cm per year ○ Growth velocity less than 2.5 cm per year • Documentation that benefits of therapy continue to outweigh risks • Documentation of improvement from baseline as assessed by the prescribing provider
Prescriber Restriction	<ul style="list-style-type: none"> • Under 18: Pediatric Endocrinologist or Nephrologist • 18 and Older: Endocrinologist
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: up to 12 months

Effective Date:	4/1/2023
P&T Approval Date:	3/14/2023

P&T Revision Date:	3/14/2023
-----------------------	-----------

References
•

Sotatercept (Winrevair)

Products Affected

- Winrevair Injection

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Pulmonary Arterial Hypertension (PAH)
Required Medical Information and Criteria	<p><u>Pulmonary Arterial Hypertension (PAH)</u></p> <ul style="list-style-type: none">• Diagnosis of symptomatic PAH (WHO Group 1 PH) confirmed by right heart catheterization.• WHO functional class II or III symptoms.• On a stable dose of both<ul style="list-style-type: none">○ Endothelin-1 receptor antagonists (ERA) and○ Phosphodiesterase type 5 inhibitors or guanylate cyclase stimulant• Current PAH background therapies (ERA, PDE5i, etc.) will be continued unless not tolerated.• Baseline platelet count >500,000
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Pulmonary Arterial Hypertension: Pulmonologist or Cardiologist
Coverage Duration	<ul style="list-style-type: none">• Pulmonary Arterial Hypertension:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	11/1/2024
P&T Approval Date:	9/10/2024
P&T Revision Date:	9/10/2024

References
•

Sparsentan (Filspari)

Products Affected

- Filspari Tablets

Covered Uses	<ul style="list-style-type: none">All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">Primary Immunoglobulin A Nephropathy.
Required Medical Information and Criteria	<u>Primary Immunoglobulin A Nephropathy</u> <ul style="list-style-type: none">Urine protein-to-creatinine ratio (UPCR) ≥ 1.5 and eGFR ≥ 30 mL/min/1.73 m²Biopsy-verified primary IgA nephropathyNo history of kidney transplant and not currently receiving dialysisMember has failed to achieve a reduction in proteinuria to under 1 gram/day while receiving maximally tolerated doses of an ACE inhibitor or ARB for at least 12 weeks
Renewal Criteria	<ul style="list-style-type: none">Documentation of improved or stable kidney function compared to baseline or reduction in proteinuria
Age Restriction	<ul style="list-style-type: none">Age 18 or older
Prescriber Restriction	<ul style="list-style-type: none">Primary Immunoglobulin A Nephropathy: Nephrologist.
Coverage Duration	<ul style="list-style-type: none">Primary Immunoglobulin A Nephropathy:<ul style="list-style-type: none">Initial: 6 monthsRenewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023

References
<ul style="list-style-type: none">

Sumatriptan Nasal Spray Step Therapy

Products Affected

- Sumatriptan Nasal Spray

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Formulary triptan tablet
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Suzetrigine (Journavx)

Products Affected

- Journavx tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Acute, moderate to severe pain
Required Medical Information and Criteria	<p><u>Acute moderate to severe pain</u></p> <ul style="list-style-type: none"> • Clinical documentation of post-operative use following one of the following: <ul style="list-style-type: none"> ○ Abdominoplasty ○ Bunionectomy • Documentation of one of the following: <ul style="list-style-type: none"> ○ Diagnosis of opioid use disorder ○ Prescriber has a specific concern for opioid abuse.
Renewal Criteria	<ul style="list-style-type: none"> • Not eligible for renewal, patients will need to meet initial criteria with new surgery to be eligible for a new prescription.
Exclusion Criteria	<ul style="list-style-type: none"> • Use for more than 14 days. • Any use outside of acute post-procedural pain.
Age Restriction	<ul style="list-style-type: none"> • Age 18 or older.
Coverage Duration	<ul style="list-style-type: none"> • Acute moderate to severe pain: <ul style="list-style-type: none"> ○ Initial: 14 days ○ Renewal: N/A

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

References
<ul style="list-style-type: none"> • Chou, Roger et al. Management of Postoperative Pain: A Clinical Practice Guideline From the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists'

Committee on Regional Anesthesia, Executive Committee, and Administrative Council. The Journal of Pain, Volume 17, Issue 2, 131 - 157

- Dowel D, Ragan KR, Jones CM, Baldwin GT, Chou R. Center for Disease Control and Prevention. CDC Clinical Practice Guideline for Prescribing Opioids for Pain, United States, 2022. CDC Morbidity and Mortality Weekly Report. Recommendations and Reports; Nov 4, 2022: 71 (3); 1-95. Found at: CDC Clinical Practice Guideline for Prescribing Opioids for Pain — United States, 2022 | MMWR
- Hegmann, Kurt T. MD, MPH; Weiss, Michael S. MD, MPH; Bowden, Kirk PhD; Branco, Fernando MD; DuBrueier, Kimberly PharmD, RPh; Els, Charl MBChB, FCPsych, MMed Psych; Mandel, Steven MD; McKinney, David W. MD, MPH; Miguel, Rafael MD; Mueller, Kathryn L. MD, MPH; Nadig, Robert J. MD, MPH; Schaffer, Michael I. PhD, MS, DABFT, NRCC-TC; Studt, Larry MD; Talmage, James B. MD; Travis, Russell L. MD; Winters, Thomas MD; Thiese, Matthew S. PhD, MSPH; Harris, Jeffrey S. MD, MPH. ACOEM Practice Guidelines: Opioids for Treatment of Acute, Subacute, Chronic, and Postoperative Pain. Journal of Occupational and Environmental Medicine 56(12):p e143-e159, December 2014. | DOI: 10.1097/JOM.0000000000000352.
- Herzig SJA, Mosher HJR, Calcaterra SLN, Jena ABA, Nuckols TKL. Improving the Safety of Opioid Use for Acute Noncancer Pain in Hospitalized Adults: A Consensus Statement From the Society of Hospital Medicine. J Hosp Med. 2018 Apr;13(4):263-71. <https://pubmed.ncbi.nlm.nih.gov/29624189>. doi:10.12788/jhm.2980.
- Hsu, Joseph R. MD*; Mir, Hassan MD†; Wally, Meghan K. MSPH*; Seymour, Rachel B. PhD*; the Orthopaedic Trauma Association Musculoskeletal Pain Task Force. Clinical Practice Guidelines for Pain Management in Acute Musculoskeletal Injury. Journal of Orthopaedic Trauma 33(5):p e158-e182, May 2019. | DOI: 10.1097/BOT.0000000000001430
- Journavx (suzetrigine) [Package Insert]. Vertex Pharmaceuticals, Inc; Boston, MA: 2025. Accessed at: https://pi.vrtx.com/files/uspi_suzetrigine.pdf.
- Vertex Pharmaceuticals. Evaluation of Efficacy and Safety of VX-548 for Acute Pain After a Bunionectomy. NCT05553366. Updated March 26, 2025. Accessed March 26, 2025. <https://www.clinicaltrials.gov/study/NCT05553366>.
- Vertex Pharmaceuticals. Evaluation of Efficacy and Safety of VX-548 for Acute Pain After an Abdominoplasty. NCT05558410. Updated August 27, 2024, Accessed March 26, 2025. <https://www.clinicaltrials.gov/study/NCT05558410>.

Synthroid Step Therapy

Products Affected

- Synthroid

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Generic levothyroxine
------------------------------	---

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Tacrolimus Ointment

Products Affected

- Tacrolimus 0.1% Ointment
- Tacrolimus 0.03% Ointment

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Atopic Dermatitis○ Psoriasis
Required Medical Information and Criteria	<p><u>Atopic Dermatitis</u></p> <ul style="list-style-type: none">• Clinical documentation of diagnosis of moderate to severe atopic dermatitis<ul style="list-style-type: none">○ At least 10% of body surface area involved○ Hand, foot, face, or mucous membrane involvement• Trial and failure of all the following (or a reason they are not appropriate):<ul style="list-style-type: none">○ High potency topical steroids○ UVB phototherapy <p><u>Psoriasis</u></p> <ul style="list-style-type: none">• Diagnosis of moderate to severe psoriasis as indicated by a validated tool such as the Dermatology Life Quality Index and one of the following:<ul style="list-style-type: none">○ At least 10% of body surface area involved○ Hand, foot, face, or mucous membrane involvement• Trial and failure of all the following (or a reason they are not appropriate):<ul style="list-style-type: none">○ High potency topical steroids○ UVB phototherapy
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	

P&T Revision Date:	
-----------------------	--

References
•

Tazarotene

Products Affected

- Tazarotene 0.1% Cream

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Psoriasis○ Acne
Required Medical Information and Criteria	<p><u>Psoriasis</u></p> <ul style="list-style-type: none">• Diagnosis of moderate to severe psoriasis as indicated by a validated tool such as the Dermatology Life Quality Index and one of the following:<ul style="list-style-type: none">○ At least 10% of body surface area involved○ Hand, foot, face, or mucous membrane involvement <p><u>Acne</u></p> <ul style="list-style-type: none">• Confirmation that the diagnosis is one of the following:<ul style="list-style-type: none">○ Above the line (severe acne)○ The member is under the 21 and acne is substantially interfering with daily life• Trial and Failure of at least 2 other formulary alternatives used to treat acne
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	3/1/2023
P&T Approval Date:	1/10/2023
P&T Revision Date:	1/10/2023

References
•

Tenofovir Alafenamide (Descovy)

Products Affected

- Descovy Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of HIV Infection○ Pre-Exposure Prophylaxis of HIV Infection (PrEP)
Required Medical Information and Criteria	<p><u>Treatment of HIV Infection</u></p> <ul style="list-style-type: none">• Clinical contraindication to generic Truvada• Documentation that the drug will be used in combination with other HIV drugs as part of a complete treatment regimen <p><u>Pre-Exposure Prophylaxis of HIV Infection (PrEP)</u></p> <ul style="list-style-type: none">• Clinical contraindication to generic Truvada
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: Lifetime

Effective Date:	9/1/2024
P&T Approval Date:	7/11/2024
P&T Revision Date:	7/11/2024

References
<ul style="list-style-type: none">•

Testosterone

Products Affected

- Testosterone 1% Gel (25mg & 50mg)
- Testosterone 1% Gel pump

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Gender Dysphoria. ○ AIDS Wasting Syndrome. ○ Post-Menopausal Breast Cancer. ○ Hypogonadism.
Required Medical Information and Criteria	<p><u>Gender Dysphoria</u></p> <ul style="list-style-type: none"> • Diagnosis of gender dysphoria • Trial and failure or contraindication to injectable testosterone <p><u>AIDS Wasting Syndrome</u></p> <ul style="list-style-type: none"> • Diagnosis of AIDS wasting syndrome • Trial and failure or contraindication to injectable testosterone <p><u>Post-Menopausal Breast Cancer</u></p> <ul style="list-style-type: none"> • Diagnosis of post-menopausal breast cancer • Trial and failure or contraindication to injectable testosterone <p><u>Hypogonadism</u></p> <ul style="list-style-type: none"> • Diagnosis of hypogonadism • Trial and failure or contraindication to injectable testosterone
Renewal Criteria	<ul style="list-style-type: none"> • Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> • Gender Dysphoria, AIDS Wasting Syndrome, Post-Menopausal Breast Cancer: <ul style="list-style-type: none"> ○ Initial: Lifetime • Hypogonadism: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months

Effective Date:	7/1/2024
-----------------	----------

P&T Approval Date:	9/13/2022
P&T Revision Date:	5/14/2024, 9/13/2022

References
•

Tezacaftor-Ivacaftor (Symdeko)

Products Affected

- Symdeko Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design.<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<u>Cystic Fibrosis</u> <ul style="list-style-type: none">• Documentation of cystic fibrosis diagnosis with homozygous F508del mutation• Not used in combination with other CFTR modulator treatments
Renewal Criteria	<ul style="list-style-type: none">• Documented clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 1 years of age and older
Prescriber Restriction	<ul style="list-style-type: none">• Cystic Fibrosis: Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Cystic Fibrosis:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	05/01/2025
P&T Approval Date:	05/01/2021
P&T Revision Date:	09/01/2021, 03/11/2025

References
<ul style="list-style-type: none">•

Thickener (Thick-It)

Products Affected

- Thick-It
- Thick-It #2

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Swallowing Disorder
Required Medical Information and Criteria	<p><u>Swallowing Disorder</u></p> <ul style="list-style-type: none">• Documented swallowing study/evaluation for the treatment of swallowing disorder due to one of the following medical conditions:<ul style="list-style-type: none">○ Diagnosis of dysphagia which negatively impacts the ability to swallow○ Chronic diseases such as, but not limited to, Parkinson's, dementia, reflux disease, stroke, neuromuscular disease/disorder, and spinal cord injury○ Treatment of head, neck, or throat cancer○ Documented aspiration of food or liquid associated with chronic illness or disease
Renewal Criteria	<ul style="list-style-type: none">• Documentation of appointment in the last 12 months confirming effectiveness of the requested thickener and continued need
Coverage Duration	<ul style="list-style-type: none">• Swallowing Disorder:<ul style="list-style-type: none">○ Initial: 6 months○ Renewal: 12 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References
<ul style="list-style-type: none">• Oregon administrative rule 410-148-0260 (7)(a)(A-D)(b)

Tiagabine

Products Affected

- Tiagabine Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Partial (Focal) Seizures
Required Medical Information and Criteria	<u>Partial Seizures</u> <ul style="list-style-type: none">• Confirmation of diagnosis of Partial Seizures
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none">• Partial Seizures:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

References
<ul style="list-style-type: none">•

Ticagrelor

Products Affected

- Ticagrelor tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Acute Coronary Syndrome ○ Minor Ischemic Stroke ○ Etc.
Required Medical Information and Criteria	<p><u>Acute Coronary Syndrome</u></p> <ul style="list-style-type: none"> • Member has: <ul style="list-style-type: none"> ○ Either non-ST-elevation acute coronary syndrome (NSTEMI) or ST-elevation myocardial infarction (STEMI) AND ○ Has had percutaneous coronary intervention (PCI) AND ○ Has a contraindication to prasugrel <p>OR</p> <ul style="list-style-type: none"> • Member has NSTEMI and is treated with medical therapy alone (has not had PCI) <p><u>Minor Ischemic Stroke</u></p> <ul style="list-style-type: none"> • Member has had a minor non-cardioembolic ischemic stroke (NIHSS score ≤5) in the immediate past • Did not receive IV alteplase • Has a reason that clopidogrel can't be used
Renewal Criteria	<ul style="list-style-type: none"> • Acute Coronary Syndrome: Documentation of positive clinical response to therapy and continued need for treatment • Minor ischemic stroke: Renewal not appropriate
Age Restriction	<ul style="list-style-type: none"> • 18 years of age and older
Prescriber Restriction	<ul style="list-style-type: none"> • Acute Coronary Syndrome: Cardiologist • Minor Ischemic Stroke: Cardiologist, Neurologist
Coverage Duration	<ul style="list-style-type: none"> • Acute Coronary Syndrome: <ul style="list-style-type: none"> ○ Initial: 12 months ○ Renewal: 12 months • Minor Ischemic Stroke: <ul style="list-style-type: none"> ○ Initial: 1 month ○ Renewal: N/A

Effective Date:	9/1/2025
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/8/2025, 7/11/2023

<p>References</p> <ul style="list-style-type: none"> • Brilinta [package insert]. Wilmington, DE: AstraZeneca Pharmaceuticals LP; 2024. • Rao SV, et al. 2025 ACC/AHA/ ACEP/NAEMSP/SCAI guideline for the management of patients with acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. J Am Coll Cardiol. 2025;85(22):2135-2237. • Powers WJ,et al.; on behalf of the American Heart Association Stroke Council. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. Stroke. 2019;50:e344–e418
--

Tobramycin Solution

Products Affected

- Tobramycin Nebulization Solution

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<p><u>Cystic Fibrosis</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of cystic fibrosis <p><u>All Other Diagnoses</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of an FDA approved or compendia supported indication
Renewal Criteria	<ul style="list-style-type: none">• Confirmed diagnosis with clinical evidence supporting chronic use
Prescriber Restriction	<ul style="list-style-type: none">• Cystic Fibrosis: Prescribed by or in consultation with Infectious Disease Specialist, Pulmonologist, or Cystic Fibrosis Specialist• All Other Diagnoses: Prescribed by or in consultation with Infectious Disease Specialist, or Pulmonologist
Coverage Duration	<ul style="list-style-type: none">• Cystic Fibrosis:<ul style="list-style-type: none">○ Initial: Lifetime• All Other Diagnoses:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 12 months

Effective Date:	1/1/2025
P&T Approval Date:	11/12/2024
P&T Revision Date:	11/12/2024

References

- Mogayzel, Jr, PJ, et al. Cystic Fibrosis Foundation Pulmonary Guidelines. Pharmacologic Approaches to Prevention and Eradication of Initial *Pseudomonas aeruginosa* Infection. AnnalsATS 2014; 11(10): 1640-1650.
- Mogayzel, Jr, PJ, et al. Cystic Fibrosis Foundation Pulmonary Guidelines. Chronic Medications for Maintenance of Lung Health. Am J Respir Crit Care Med 2013; 18 (7): 680–689.

Tocilizumab (Actemra, Tyenne)

Products Affected

- Actemra Prefilled Syringe
- Actemra ActPen
- Tyenne Prefilled Syringe
- Tyenne Auto-Injector
- Tyenne IV Solution

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Juvenile Idiopathic Arthritis ○ Rheumatoid Arthritis ○ Giant Cell Arteritis ○ Systemic Sclerosis-Associated Interstitial Lung Disease ○ Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none"> • Initial testing for latent TB and treatment, if necessary, before starting treatment. • No current active infection at initiation of therapy. • Risks and benefits documented in cases of chronic or recurrent infection. • Will NOT be used in combination with another biologic or Otezla. <p><u>Juvenile Idiopathic Arthritis (JIA)</u></p> <ul style="list-style-type: none"> • Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids

	<ul style="list-style-type: none"> • Documented intolerance or contraindication to oral disease-modifying antirheumatic drugs (DMARD) such as methotrexate, leflunomide, etc. or DMARDs will be continued • Trial and failure of both infliximab and adalimumab <p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of both infliximab and adalimumab <p><u>Giant Cell Arteritis:</u></p> <ul style="list-style-type: none"> • Clinical documentation confirming diagnosis of Giant Cell Arteritis • Trial and failure of glucocorticoid treatment <p><u>Systemic Sclerosis-Associated Interstitial Lung Disease:</u></p> <ul style="list-style-type: none"> • Confirmed diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) • Trial and failure of mycophenolate mofetil <p><u>Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy:</u></p> <ul style="list-style-type: none"> • Member will receive or is receiving chimeric antigen receptor (CAR) T-cell immunotherapy
Renewal Criteria	<ul style="list-style-type: none"> • Juvenile Idiopathic Arthritis: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Giant Cell Arteritis: Demonstrated positive clinical response to therapy. • Systemic Sclerosis-Associated Interstitial Lung Disease: Demonstrated positive clinical response to therapy.

Prescriber Restriction	<ul style="list-style-type: none"> • Rheumatoid Arthritis, Juvenile Idiopathic Arthritis, Giant Cell Arteritis: Rheumatologist • Systemic Sclerosis-Associated Interstitial Lung Disease: Rheumatologist or Pulmonologist • Cytokine Release Syndrome (CRS) Risk due to CAR T-Cell Therapy: Oncologist or Hematologist
Coverage Duration	<ul style="list-style-type: none"> • Cytokine Release Syndrome: <ul style="list-style-type: none"> ○ Initial: 2 months ○ Renewal: N/A • All Other Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2024
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/9/2024, 7/11/2023

References
•

Tofacitinib (Xeljanz)

Products Affected

- Xeljanz XR tablets
- Xeljanz Tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Juvenile Idiopathic Arthritis ○ Psoriatic Arthritis ○ Rheumatoid Arthritis ○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of infliximab and adalimumab • Trial and failure of Cosentyx and Taltz <p><u>Juvenile Idiopathic Arthritis (JIA)</u></p> <ul style="list-style-type: none"> • Documentation of juvenile idiopathic arthritis with active systemic features of JIA, with a physician global assessment of 5 or higher (or any systemic activity in the absence of active joint involvement) or JIA without active systemic features • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR

	<ul style="list-style-type: none"> ○ Documented trial and failure of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ At least 6 months of 2 of Methotrexate, leflunomide, sulfasalazine, hydroxychloroquine, or systemic corticosteroids • Documented intolerance or contraindication to DMARDs OR DMARD will be continued with Xeljanz. • Trial and failure of infliximab and adalimumab • Trial and failure of Actemra and Orencia <p><u>Psoriatic Arthritis (PsA):</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of infliximab and adalimumab • Trial and failure of ustekinumab, Cosentyx, Otezla, Orencia, and Taltz <p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following:
--	---

	<ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of infliximab and adalimumab • Trial and failure of Actemra, Cimzia, Kineret, Orencia, and rituximab <p><u>Ulcerative Colitis (UC):</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: mesalamine, sulfasalazine, mercaptopurine, azathioprine, or corticosteroids (prednisone, methylprednisolone). • Trial and failure of infliximab and adalimumab • Trial and failure of ustekinumab and Entyvio
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Juvenile Idiopathic Arthritis: Evidence of 20% or greater improvement in tender joint count and swollen joint count or has there been an improvement in functional ability. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Ulcerative Colitis: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Juvenile Idiopathic Arthritis: Rheumatologist

	<ul style="list-style-type: none"> • Rheumatoid Arthritis: Rheumatologist. • Ulcerative Colitis: Gastroenterologist.
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023, 1/11/2022

References
•

Tolterodine Step Therapy

Products Affected

- Tolterodine ER capsule
- Tolterodine tablet

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Oxybutynin
------------------------------	--

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

Topical Antifungal Agents

Products Affected

- Ciclopirox 8% Solution
- Econazole 1% Cream
- Selenium Sulfide 2.5% Lotion
- Ketoconazole 2% Cream and Shampoo

Covered Uses	<ul style="list-style-type: none"> All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design
Required Medical Information and Criteria	<p><u>All Diagnoses</u></p> <ul style="list-style-type: none"> All the following have been tried and failed or are not appropriate for use: <ul style="list-style-type: none"> Clotrimazole 1% cream Miconazole 2% (cream, aerosol, or powder), Terbinafine 1%, cream, terbinafine tablets Nystatin 100,000 units/gram (ointment, cream, or powder)
Renewal Criteria	<ul style="list-style-type: none"> Documented positive clinical response to therapy
Coverage Duration	<ul style="list-style-type: none"> All Diagnoses: <ul style="list-style-type: none"> Initial: 3 months Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023, 1/11/2022

References
•

Toujeo Step Therapy

Products Affected

- Toujeo Max Solostar
- Toujeo Solostar

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Any non-concentrated basal insulin product
------------------------------	--

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Tranexamic Acid

Products Affected

- Tranexamic Acid Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Hemophilia – Hemorrhage Prophylaxis; Tooth Extraction○ Menorrhagia
Required Medical Information and Criteria	<p><u>Hemophilia – Hemorrhage Prophylaxis</u></p> <ul style="list-style-type: none">○ Documentation that use is intended for hemorrhage prophylaxis for tooth extraction <p><u>Menorrhagia</u></p> <ul style="list-style-type: none">• Documentation of abnormal uterine bleeding• Trail and failure of, current use of, or contraindication to all of the following:<ul style="list-style-type: none">○ Combined Oral Contraceptive therapy○ Progestin therapy (oral or LM) or Levonorgestrel IUD○ NSAID therapy
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to therapy
Prescriber Restriction	<ul style="list-style-type: none">• Hemorrhage Prophylaxis: Hematologist, Hemophilia Specialist, Dentist• Menorrhagia: Gynecologist
Coverage Duration	<ul style="list-style-type: none">• All Diagnoses:<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	7/1/2024
P&T Approval Date:	5/14/2024
P&T Revision Date:	5/14/2024

References

•

Tretinoin, Topical

Products Affected

- Tretinoin **0.025% cream**
- Tretinoin **0.05% cream**
- Tretinoin **0.1% cream**
- Tretinoin **0.01% gel**
- Tretinoin **0.025% gel**

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Acne
Required Medical Information and Criteria	<p><u>Acne</u></p> <ul style="list-style-type: none">• Documentation of trial and failure, intolerance, or contraindication to a topical product containing benzoyl peroxide.
Renewal Criteria	<ul style="list-style-type: none">• Documentation of positive clinical response to therapy.
Coverage Duration	<ul style="list-style-type: none">• All Diagnosis<ul style="list-style-type: none">○ Initial: 12 months○ Renewal: 12 months

Effective Date:	02/01/2023
P&T Approval Date:	01/10/2023
P&T Revision Date:	01/10/2023

Trospium Step Therapy

Products Affected

- Trospium IR tablets
- Trospium ER caps

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Oxybutynin IR or ER
------------------------------	---

Effective Date:	7/1/2025
P&T Approval Date:	5/13/2025
P&T Revision Date:	5/13/2025

Ubrogepant (Ubrelevy)

Products Affected

- Ubrelevy tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of acute migraine○
Required Medical Information and Criteria	<p><u>Acute Migraine</u></p> <ul style="list-style-type: none">• Confirmed diagnosis of migraine• Documentation showing the member is currently on preventative therapy• Trial and failure (defined as at least 6 weeks per agent) of:<ul style="list-style-type: none">○ At least 3 oral formulary triptans used at up to the maximally indicated dosing and in combination with NSAID therapy (e.g. naproxen).
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response to therapy
Age Restriction	<ul style="list-style-type: none">• 18 years of age or older
Prescriber Restriction	<ul style="list-style-type: none">• Acute Migraine: Neurologist or headache specialist
Coverage Duration	<ul style="list-style-type: none">• Acute Migraine:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 12 months

Effective Date:	9/1/2025
P&T Approval Date:	5/9/2023
P&T Revision Date:	7/8/2025, 5/9/2023

References

- Ubrelevy [package insert]. North Chicago, IL: Abbvie Inc; 2025.

Upadacitinib (Rinvoq)

Products Affected

- Rinvoq Tablets
- Rinvoq LQ Solution

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Ankylosing Spondylitis/Axial Spondyloarthritis ○ Atopic Dermatitis ○ Crohn's Disease ○ Psoriatic Arthritis ○ Rheumatoid Arthritis ○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>Ankylosing Spondylitis/Axial Spondyloarthritis (AS/SpA)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ankylosing spondylitis or axial spondyloarthritis as defined by: <ul style="list-style-type: none"> ○ Back pain and stiffness for more than 3 months AND ○ Signs of active inflammation on MRI OR radiological evidence of sacroiliitis OR HLA-B27 positive AND ○ BASDAI score of ≥ 4 • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ At least two NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated AND ▪ Physical therapy/exercise program • Trial and failure of infliximab and adalimumab • Trial and failure of Cosentyx and Taltz <p><u>Atopic Dermatitis</u></p> <ul style="list-style-type: none"> • Diagnosed with severe atopic dermatitis defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) ≥ 11 or Children's Dermatology Life Quality Index (CDLQI) ≥ 13 (or severe score on another validated tool) • One or more of the following: <ul style="list-style-type: none"> ○ At least 10% of body surface area involvement ○ Hand, foot, or mucous membrane involvement

	<ul style="list-style-type: none"> • Documented contraindication or failed trial to ALL of the following: <ul style="list-style-type: none"> ○ Moderate-high potency corticosteroid (e.g., clobetasol, fluocinonide, fluticasone) ○ Topical calcineurin inhibitor (e.g. tacrolimus) ○ Oral immunomodulator therapy (e.g. cyclosporine, methotrexate, azathioprine, mycophenolate mofetil) OR the member is oral corticosteroid dependent. • Failure of Dupixent <p><u>Crohn's Disease (CD)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe Crohn's Disease • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: 6-mercaptopurine, azathioprine, corticosteroid, methotrexate. • Trial and failure of infliximab and adalimumab • Trial and failure of ustekinumab, Cimzia, and Entyvio <p><u>Psoriatic Arthritis (PsA):</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current of history) ○ Radiological evidence of juxta-articular new bone formation • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ▪ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ▪ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine. • Trial and failure of infliximab and adalimumab
--	--

	<ul style="list-style-type: none"> • Trial and failure of ustekinumab, Cosentyx, Otezla, Orencia, and Taltz <p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • Documentation of a baseline of moderate to high disease activity of rheumatoid arthritis measured as such by an accepted assessment instrument (PAS, PASII, RAPID3, CDAI, DAS28, SDAI) • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of nonbiologic DMARD therapy: methotrexate (dosed at least 20mg per week for at least 8 weeks), leflunomide or sulfasalazine • Trial and failure of infliximab and adalimumab • Trial and failure of Actemra, Cimzia, Kineret, Orencia, and rituximab <p><u>Ulcerative Colitis (UC):</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • One of the following: <ul style="list-style-type: none"> ○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR ○ Documented trial and failure of at least 1 of the following: mesalamine, sulfasalazine, mercaptopurine, azathioprine, or corticosteroids (prednisone, methylprednisolone). • Trial and failure of infliximab and adalimumab • Trial and failure of ustekinumab and Entyvio
Renewal Criteria	<ul style="list-style-type: none"> • Ankylosing Spondylitis/Axial Spondyloarthritis: Evidence of significant improvement in signs and symptoms of AS/SpA and/or functioning, such as ASAS40 or 2-point improvement in BASDAI. • Atopic Dermatitis: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement.

	<ul style="list-style-type: none"> • Crohn's Disease: Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission. • Psoriatic Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Rheumatoid Arthritis: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • Ulcerative Colitis: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Prescriber Restriction	<ul style="list-style-type: none"> • Ankylosing Spondylitis, Axial Spondyloarthritis: Rheumatologist. • Atopic Dermatitis: Dermatologist. • Crohn's Disease and Ulcerative Colitis: Gastroenterologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist. • Rheumatoid Arthritis: Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All Diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	9/1/2023
P&T Approval Date:	7/11/2023
P&T Revision Date:	7/11/2023, 1/11/2022

References
•

Ustekinumab

Products Affected

- Selarsdi
- Yesintek
- Steqeyma

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Crohn's Disease○ Plaque Psoriasis○ Psoriatic Arthritis○ Ulcerative Colitis
Required Medical Information and Criteria	<p><u>All diagnoses</u></p> <ul style="list-style-type: none">• Initial testing for latent TB and treatment, if necessary, before starting treatment.• No current active infection at initiation of therapy.• Risks and benefits documented in cases of chronic or recurrent infection.• Will NOT be used in combination with another biologic or Otezla <p><u>Crohn's Disease (CD)</u></p> <ul style="list-style-type: none">• Documentation of moderate-to-severe Crohn's Disease• One of the following:<ul style="list-style-type: none">○ The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR○ Documented trial and failure of at least 1 of the following: 6-mercaptopurine, azathioprine, corticosteroid, methotrexate• Trial and failure of both infliximab and adalimumab <p><u>Plaque Psoriasis (PP)</u></p> <ul style="list-style-type: none">• Documentation of severe plaque psoriasis, defined as having functional impairment as indicated by Dermatology Life Quality Index (DLQI) = 11 or Children's Dermatology Life Quality Index (CDLQI) = 13 (or severe score on other validated tool) AND one or more of the following:<ul style="list-style-type: none">○ At least 10% of body surface area involved○ Hand, foot, face, or mucous membrane involvement

	<ul style="list-style-type: none"> • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of all the following: <ul style="list-style-type: none"> ○ High-potency topical corticosteroids (augmented betamethasone, clobetasol, etc.) ○ At least one other topical agent (calcipotriene, tazarotene, anthralin, tar, etc.) ○ PUVA or UVB Phototherapy ○ Methotrexate ○ At least 1 other second line systemic agent such as cyclosporine or acitretin • Trial and failure of both infliximab and adalimumab <p><u>Psoriatic Arthritis (PsA)</u></p> <ul style="list-style-type: none"> • Documentation of psoriatic arthritis based on at least 3 out of 5 of the following: <ul style="list-style-type: none"> ○ Psoriasis (1 point for personal or family history, 2 points for current) ○ Psoriatic nail dystrophy ○ Negative test result for RF ○ Dactylitis (current or history) ○ Radiological evidence of juxta-articular new bone formation • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of conventional therapy with both of the following: <ul style="list-style-type: none"> ○ NSAIDs for 3 months at maximum recommended or tolerated anti-inflammatory dose unless contraindicated, AND ○ Methotrexate or other DMARD such as leflunomide, sulfasalazine, or cyclosporine • Trial and failure of both infliximab and adalimumab <p><u>Ulcerative Colitis (UC)</u></p> <ul style="list-style-type: none"> • Documentation of moderate-to-severe ulcerative colitis • The member is transitioning to the requested treatment from a different biologic product previously approved by the plan OR documented failure of at least 1 of the following: <ul style="list-style-type: none"> ○ Mesalamine, sulfasalazine OR ○ Mercaptopurine, azathioprine, OR ○ Corticosteroids (prednisone, methylprednisolone) • Trial and failure of both infliximab and adalimumab
--	---

Renewal Criteria	<ul style="list-style-type: none"> • CD: Evidence of a decrease in symptoms, reduction in enterocutaneous fistulas or clinical remission. • PP: Evidence of positive clinical response to therapy as evidenced by ONE of the following: reduction of body surface area (BSA) involvement from baseline, improvement in symptoms (e.g. pruritus, inflammation) from baseline, or evidence of functional improvement. • PsA: Evidence of a 20% or greater improvement in tender joint count and swollen joint count. • UC: Evidence of a significant response such as a decrease in bloody stools per day or elimination of signs of toxicity.
Exclusion Criteria	<ul style="list-style-type: none"> • Not to be used in combination with other biologics for the same indication.
Prescriber Restriction	<ul style="list-style-type: none"> • Crohn's Disease and Ulcerative Colitis: Gastroenterologist. • Plaque Psoriasis: Dermatologist. • Psoriatic Arthritis: Dermatologist or Rheumatologist.
Coverage Duration	<ul style="list-style-type: none"> • All diagnoses: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: 12 months

Effective Date:	5/1/2025
P&T Approval Date:	7/11/2023
P&T Revision Date:	3/11/2025, 7/11/2023, 1/11/2022

References
<ul style="list-style-type: none"> • Selarsdi [package insert]. Parsippany, NJ: Teva Pharmaceuticals; 2025. • Yesintek [package insert]. Cambridge, MA: Biocon Biologics; 2024. • Steqeyma [package insert]. Jersey City, NJ: Celltrion USA Inc.; 2024.

Valacyclovir Step Therapy

Products Affected

- Valacyclovir tablets

Step Therapy Criteria	<ul style="list-style-type: none">• Trial and failure of<ul style="list-style-type: none">○ Acyclovir tablets
------------------------------	---

Effective Date:	
P&T Approval Date:	
P&T Revision Date:	

Vanzacaftor-tezacaftor-deutivacaftor (Alyftrek)

Products Affected

- Alyftrek Tablets

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design.<ul style="list-style-type: none">○ Cystic Fibrosis
Required Medical Information and Criteria	<p><u>Cystic Fibrosis</u></p> <ul style="list-style-type: none">• Presence of at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA):<ul style="list-style-type: none">○ F508del mutation○ A mutation in the CFTR gene that is responsive based on clinical, in vitro, or extrapolated data.• Not to be used in combination with other CFTR modulator treatments
Renewal Criteria	<ul style="list-style-type: none">• Demonstrated positive clinical response to therapy.
Age Restriction	<ul style="list-style-type: none">• Age 6 and older.
Prescriber Restriction	<ul style="list-style-type: none">• Pulmonologist or Specialist affiliated with a CF care center.
Coverage Duration	<ul style="list-style-type: none">• Initial: 6 months• Renewal: 12 months

Effective Date:	05/01/2025
P&T Approval Date:	03/11/2025
P&T Revision Date:	03/11/2025

References

- Alyftrek [package insert]. Boston, MA: Vexta Pharmaceuticals Inc.; 2025.

--

Vonoprazan (Voquezna)

Products Affected

- Voquezna 10mg and 20mg tablets

Covered Uses	<ul style="list-style-type: none"> • All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design <ul style="list-style-type: none"> ○ Erosive esophagitis ○ H. Pylori infection
Required Medical Information and Criteria	<p><u>Erosive Esophagitis</u></p> <ul style="list-style-type: none"> • Imaging confirmed LA Classification Grade C/D erosive esophagitis AND • Documented contraindication, intolerance, or inadequate response to 2 or more PPIs (i.e., lansoprazole, omeprazole, esomeprazole, etc.) at maximum tolerated twice-daily dosing for at least 8 weeks each. <p><u>H. Pylori Infection</u></p> <ul style="list-style-type: none"> • Confirmed H. pylori positive infection • Documented contraindication, intolerance, or inadequate response to: <ul style="list-style-type: none"> ○ Standard first-line therapies for H.pylori infection <ul style="list-style-type: none"> ▪ PPI + clarithromycin + (amoxicillin or metronidazole) ▪ Bismuth quadruple therapy • Co-prescribed with antibiotics
Renewal Criteria	<ul style="list-style-type: none"> • Renewals past the initial approved timelines are not allowed
Age Restriction	<ul style="list-style-type: none"> • 18 years of age or older.
Prescriber Restriction	<ul style="list-style-type: none"> • All diagnoses: Gastroenterologist or Infectious Disease specialist
Coverage Duration	<ul style="list-style-type: none"> • Initial healing of erosive esophagitis: <ul style="list-style-type: none"> ○ Initial: 2 months ○ Renewal: n/a • Maintenance of healing of erosive esophagitis: <ul style="list-style-type: none"> ○ Initial: 6 months ○ Renewal: n/a • H. Pylori infection:

	<ul style="list-style-type: none"> ○ Initial: 14 days ○ Renewal: n/a
--	--

Effective Date:	3/1/2024
P&T Approval Date:	1/9/2024
P&T Revision Date:	1/9/2024

References	
<ul style="list-style-type: none"> • Voquezna [package insert]. Buffalo Grove, IL: Phatham Pharmaceuticals; 2022. 	

Voriconazole

Products Affected

- Voriconazole tablets
- Voriconazole suspension

Covered Uses	<ul style="list-style-type: none">• All Food and Drug Administration (FDA)-approved indications not otherwise excluded by plan design<ul style="list-style-type: none">○ Treatment of invasive aspergillosis○ Treatment of serious fungal infections
Required Medical Information and Criteria	<u>All diagnoses</u> <ul style="list-style-type: none">• Documentation of one of the approved conditions
Renewal Criteria	<ul style="list-style-type: none">• Documented positive clinical response and continued need
Age Restriction	<ul style="list-style-type: none">• Suspension only: Member is under age 10 or unable to use tablets
Coverage Duration	<ul style="list-style-type: none">• All diagnoses:<ul style="list-style-type: none">○ Initial: 3 months○ Renewal: 3 months

Effective Date:	5/1/2024
P&T Approval Date:	3/12/2024
P&T Revision Date:	3/12/2024

References

- Vfend [package insert]. New York, NY: Pfizer.; 2025.

Zafirlukast Step Therapy

Products Affected

- Zafirlukast tablets

Step Therapy Criteria	<ul style="list-style-type: none">○ Trial and failure of montelukast
------------------------------	--

Effective Date:	9/1/2025
P&T Approval Date:	7/8/2025
P&T Revision Date:	7/8/2025