Medicare Part D: Fundamental Formulary 2025

Please click here.

For Medicare Part D: Prior Authorization Criteria

Please click here.

For Medicare Part D: Step Therapy Criteria

Please click here.

For more recent information or other questions, please contact:

Senior Blue HMO Freedom HMO Forever Blue PPO BlueSaver HMO Employer Group PDP

Pharmacy Service at 1-800-329-2792.

For TTY users, 711 National Relay Service, Oct. 1 – March 31, 8 a.m. – 8 p.m. ET, seven days a week, and April 1 – Aug. 30, 8 a.m. – 8 p.m. ET, Monday – Sunday.

Visit medicare.highmark.com.

Formulary ID: 25025 Version: 22

Updated: 12/2025

Note to existing members: This Formulary has changed since last year. Please review this document to make sure that it still contains the drugs you take.

When this Drug List (Formulary) refers to "we," "us," or "our," it means Highmark Blue Cross Blue Shield or Highmark Blue Cross.

When it refers to "plan" or "our plan," it means Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO.

This document includes a Drug List (formulary) for our plan, which is current as of January 1, 2025. For an updated Drug List (formulary), please contact us. Our contact information, along with the date we last updated the Drug List (formulary), appears on the front and back cover pages.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2026, and from time to time during the year.

What is the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO Formulary?

In this document, we use the terms Drug List and formulary to mean the same thing. A formulary is a list of covered drugs selected by our plan in consultation with a team of health care providers, which represents the prescription therapies believed to be a necessary part of a quality treatment program. Our plan will generally cover the drugs listed in our formulary as long as the drug is medically necessary, the prescription is filled at one of our plan's network pharmacies, and other plan rules are followed. For more information on how to fill your prescriptions, please review your Evidence of Coverage.

Can the formulary change?

Most changes in drug coverage happen on January 1, but we may add or remove drugs on the formulary during the year, move them to different cost-sharing tiers, or add new restrictions. We must follow the Medicare rules in making these changes. Updates to the formulary are posted monthly to our website here: **medicare.highmark.com/formulary**.

Changes that can affect you this year: In the below cases, you will be affected by coverage changes during the year:

• Immediate substitutions of certain new versions of brand name drugs and original biological products. We may immediately remove a drug from our formulary if we are replacing it with a certain new version of that drug that will appear on the same or lower cost-sharing tier and with the same or fewer restrictions. When we add a new version of a drug to our formulary, we may decide to keep the brand name drug or original biological product on our formulary, but immediately move it to a different cost-sharing tier or add new restrictions.

We can make these immediate changes only if we are adding a new generic version of a brand name drug, or adding certain new biosimilar versions of an original biological product, that was already on the formulary (for example, adding an interchangeable biosimilar that can be substituted for an original biological product by a pharmacy without a new prescription.)

If you are currently taking the brand name drug or original biological product, we may not tell you in advance before we make an immediate change, but we will later provide you with information about the specific change(s) we have made.

If we make such a change, you or your prescriber can ask us to make an exception and continue to cover for you the drug that is being changed. For more information, see the section below titled "How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO's Formulary?"

Some of these drug types may be new to you. For more information, see the section below titled "What are original biological products and how are they related to biosimilars?"

- **Drugs removed from the market.** If a drug is withdrawn from sale by the manufacturer or the Food and Drug Administration (FDA) determines to be withdrawn for safety or effectiveness reasons, we may immediately remove the drug from our formulary and later provide notice to members who take the drug.
- Other changes. We may make other changes that affect members currently taking a
 drug. For instance, we may remove a brand name drug from the formulary when adding
 a generic equivalent or remove an original biological product when adding a biosimilar.
 We may also apply new restrictions to the brand name drug or original biological product,
 or move it to a different cost-sharing tier, or both. We may make changes based on new
 clinical guidelines. If we remove drugs from our formulary, add prior authorization, quantity

limits and/or step therapy restrictions on a drug, or move a drug to a higher cost-sharing tier, we must notify affected members of the change at least 30 days before the change becomes effective. Alternatively, when a member requests a refill of the drug, they may receive a 31-day supply of the drug and notice of the change.

If we make these other changes, you or your prescriber can ask us to make an exception for you and continue to cover the drug you have been taking. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled "How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO's Formulary?"

Changes that will not affect you if you are currently taking the drug. Generally, if you are taking a drug on our 2025 formulary that was covered at the beginning of the year, we will not discontinue or reduce coverage of the drug during the 2025 coverage year except as described above. This means these drugs will remain available at the same cost-sharing and with no new restrictions for those members taking them for the remainder of the coverage year. You will not get direct notice this year about changes that do not affect you. However, on January 1 of the next year, such changes would affect you, and it is important to check the formulary for the new benefit year for any changes to drugs.

The enclosed formulary is current as of December 1, 2025. To get updated information about the dru gs covered by our plan, please contact us. Our contact information appears on the front and back cover pages. In the event of mid-year non-maintenance formulary changes, members will be notified by mail and prospective members will receive an update with this formulary. The most up-to-date formulary is available on our website at **medicare.highmark.com/formulary**.

How do I use the formulary?

There are two ways to find your drug within the formulary:

Medical Condition

The formulary begins on page 9. The drugs in this formulary are grouped into categories depending on the type of medical conditions that they are used to treat. For example, drugs used to treat a heart condition are listed under the category, "Cardiovascular – Hypertension & Lipids." If you know what your drug is used for, look for the category name in the list that begins on page 9. Then look under the category name for your drug.

Alphabetical Listing

If you are not sure what category to look under, you should look for your drug in the Index that begins on page 9. The Index provides an alphabetical list of all of the drugs included in this document. Both brand name drugs and generic drugs are listed in the Index. Look in the Index and find your drug. Next to your drug, you will see the page number where you can find coverage information. Turn to the page listed in the Index and find the name of your drug in the first column of the list.

What are generic drugs?

Our plan covers both brand name drugs and generic drugs. A generic drug is approved by the FDA as having the same active ingredient as the brand name drug. Generally, generic drugs work just as well as and usually cost less than brand name drugs. There are generic drug substitutes available for many brand name drugs. Generic drugs usually can be substituted for the brand name drug at the pharmacy without needing a new prescription, depending on state laws.

What are original biological products and how are they related to biosimilars?

On the formulary, when we refer to drugs, this could mean a drug or a biological product. Biological products are drugs that are more complex than typical drugs. Since biological products are more complex than typical drugs, instead of having a generic form, they have alternatives that are called biosimilars. Generally, biosimilars work just as well as the original biological product and may cost less. There are biosimilar alternatives for some original biological products. Some biosimilars are interchangeable biosimilars and, depending on state laws, may be substituted for the original biological product at the pharmacy without needing a new prescription, just like generic drugs can be substituted for brand name drugs.

• For discussion of drug types, please see the Evidence of Coverage, Chapters 5 for MA-PD insert and 3 for PDP insert, Section 3.1, "The 'Drug List' tells which Part D drugs are covered."

Are there any restrictions on my coverage?

Some covered drugs may have additional requirements or limits on coverage. These requirements and limits may include:

- Prior Authorization: Our plan requires you or your prescriber to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don't get approval, our plan may not cover the drug.
- Quantity Limits: For certain drugs, our plan limits the amount of the drug that we will cover. For example, our plan provides 31 tablets, per 31 days, per prescription for 100mg Losartan. This may be in addition to a standard one-month or three-month supply.
- Step Therapy: In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. For example, if Drug A and Drug B both treat your medical condition, our plan may not cover Drug B unless you try Drug A first. If Drug A does not work for you, our plan will then cover Drug B.

You can find out if your drug has any additional requirements or limits by looking in the formulary that begins on page 9. You can also get more information about the restrictions applied to specific covered drugs by visiting our website. We have posted online document(s) that

explain(s) our prior authorization and step therapy restrictions. You may also ask us to send you a copy. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You can ask our plan to make an exception to these restrictions or limits or for a list of other, similar drugs that may treat your health condition. See the section, "How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO's formulary?" on page 6 for information about how to request an exception.

What if my drug is not on the formulary?

If your drug is not included in this formulary (list of covered drugs), you should first contact Pharmacy Service and ask if your drug is covered.

If you learn that our plan does not cover your drug, you have two options:

- You can ask Pharmacy Service for a list of similar drugs that are covered by our plan.
 When you receive the list, show it to your doctor and ask them to prescribe a similar drug that is covered by our plan.
- You can ask our plan to make an exception and cover your drug. See below for information about how to request an exception.

How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO's formulary?

You can ask our plan to make an exception to our coverage rules. There are several types of exceptions that you can ask us to make.

- You can ask us to cover a drug even if it is not on our formulary. If approved, this drug will
 be covered at a pre-determined cost-sharing level, and you would not be able to ask us to
 provide the drug at a lower cost-sharing level.
- You can ask us to waive a coverage restriction including prior authorization, step therapy, or a quantity limit on your drug. For example, for certain drugs, our plan limits the amount of the drug that we will cover. If your drug has a quantity limit, you can ask us to waive the limit and cover a greater amount.
- You can ask us to cover a formulary drug at a lower cost-sharing level unless the drug is
 on the specialty tier. If approved, this would lower the amount you must pay for your drug.

Generally, our plan will only approve your request for an exception if the alternative drugs included on the plan's formulary, the lower cost-sharing drug, or applying the restriction, would not be as effective for you and/or would cause you to have adverse effects.

You or your prescriber should contact us to ask for a tiering or, formulary exception, including an exception to a coverage restriction. When you request an exception, your prescriber will need to explain the medical reasons why you need the exception. Generally, we must make our decision within 72 hours of getting your prescriber's supporting statement. You can ask for an expedited (fast) decision if you believe, and we agree, that your health could be seriously harmed by waiting up to 72 hours for a decision. If we agree, or if your prescriber asks for a fast decision, we must give you a decision no later than 24 hours after we get your prescriber's supporting statement.

What can I do if my drug is not on the formulary or has a restriction?

As a new or continuing member in our plan you may be taking drugs that are not on our formulary. Or, you may be taking a drug that is on our formulary but has a coverage restriction, such as prior authorization. You should talk to your prescriber about requesting a coverage decision to show that you meet the criteria for approval, switching to an alternative drug that we cover, or requesting a formulary exception so that we will cover the drug you take. While you and your doctor determine the right course of action for you, we may cover your drug in certain cases during the first 90 days you are a member of our plan.

For each of your drugs that is not on our formulary or has a coverage restriction, we will cover a temporary 31-day supply. If your prescription is written for fewer days, we'll allow refills to provide up to a maximum 31-day supply of medication. If your coverage is not approved after your first 31-day supply, we will not pay for these drugs, even if you have been a member of the plan less than 90 days.

If you are a resident of a long-term care facility and you need a drug that is not on our formulary, or if your ability to get your drugs is limited but you are past the first 90 days of membership in our plan, we will cover a 31-day emergency supply of that drug while you pursue a formulary exception.

The above transition process will be implemented to accommodate you if you have an immediate need for a non-formulary drug or a drug that requires prior authorization due to a change in your level of care while you are waiting for an exception request to be processed.

For more information

For more detailed information about your plan's prescription drug coverage, please review your Evidence of Coverage and other plan materials.

If you have questions about our plan, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

If you have general questions about Medicare prescription drug coverage, please call Medicare at 1-800-MEDICARE (1-800-633-4227) 24 hours a day, seven days a week. TTY users should call 1-877-486-2048, or visit http://www.medicare.gov.

Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO formulary

The formulary that begins on the next page provides coverage information about the drugs covered by your plan. If you have trouble finding your drug in the list, turn to the Index that begins on page 9.

The first column of the chart lists the drug name. Brand name drugs are capitalized (e.g., ABELCET) and generic drugs are listed in lower-case italics (e.g., abacavir).

The information in the Requirements/Limits column tells you if your plan has any special requirements for coverage of your drug.

The following is a Formulary Format Example Only:

Drug Name	Fundamental Drug Tier	Requirements/ Limits
Anti-Infectives		
XYZ DRUG	NF	QL-28

12/1/2025

Table of Contents

Anti - Infectives	3
Antineoplastic / Immunosuppressant Drugs	
Autonomic / Cns Drugs, Neurology / Psych	
Cardiovascular, Hypertension / Lipids	
Dermatologicals/Topical Therapy	
Diagnostics / Miscellaneous Agents	
Ear, Nose / Throat Medications	44
Endocrine/Diabetes	
Gastroenterology	
Immunology, Vaccines / Biotechnology	
Miscellaneous Supplies	54
Musculoskeletal / Rheumatology	54
Obstetrics / Gynecology	56
Ophthalmology	
Respiratory And Allergy	62
Respiratory And AllergyUrologicals	65
Vitamins, Hematinics / Electrolytes	

Drug Tier

lowercase italics = Generic drugs

UPPERCASE BOLD = Brand

name drugs

T1 = Cost-Sharing Tier 1 includes preferred generic drugs. This is the lowest cost-sharing tier.

T2 = Cost-Sharing Tier 2 includes generic drugs.

T3 = Cost-Sharing Tier 3 includes preferred brand name drugs and may include some single-sourced drugs (those generic drugs made by a single manufacturer).

T4 = Cost-Sharing Tier 4 includes non-preferred brand name drugs and may include some single-sourced generic drugs (those generic drugs made by a single manufacturer).

T5 = Cost-Sharing Tier 5 includes specialty drugs. This is the highest cost-sharing tier.

Requirements/Limits

LA = Limited access

PA = Prior authorization required **PA-BvD** = This drug may be covered under Medicare part B or D depending on the circumstance. Information may need to be submitted describing the use and

submitted describing the use an setting of the drug to make the determination.

PA-NS = Prior authorization required for new starts only

QL = Quantity limit applies. The quantity limit is noted for each drug. For example, if the quantity limit is QL (90 EA per 180 days), the quantity limit would be 90 units per 180-day supply.

ST = Step therapy applies ST-NS = Step therapy applies to new starts only

Drug Name	Drug Tier	Requirements/Limits
Anti - Infectives		
abacavir	T2	
abacavir-lamivudine	T2	
acyclovir oral capsule	T2	
acyclovir oral suspension 200 mg/5 ml	T2	
acyclovir oral tablet	T2	
acyclovir sodium intravenous solution	T4	PA-BvD
adefovir	T4	
albendazole	T4	
amantadine hcl oral capsule	T2	QL (124 EA per 31 days)
amantadine hcl oral solution	T2	
amantadine hcl oral tablet	T2	
amikacin injection solution 500 mg/2 ml	T4	
amoxicillin oral capsule	T2	
amoxicillin oral suspension for reconstitution	T2	
amoxicillin oral tablet	T2	
amoxicillin oral tablet,chewable 125 mg, 250 mg	T2	
amoxicillin-pot clavulanate oral suspension for reconstitution	T2	
amoxicillin-pot clavulanate oral tablet	T2	

Drug Name	Drug Tier	Requirements/Limits
amphotericin b	T4	PA-BvD
amphotericin b liposome	T5	PA-BvD
ampicillin oral capsule 500 mg	T2	
ampicillin sodium injection recon soln 1 gram, 10 gram	T4	
ampicillin-sulbactam injection	T4	
APTIVUS	T5	
ARIKAYCE	T5	PA
atazanavir	T4	
atovaquone	T4	
atovaquone-proguanil	Т3	
azithromycin intravenous	T4	
azithromycin oral tablet	T2	
aztreonam injection recon soln 1 gram	T4	
aztreonam injection recon soln 2 gram	T5	
BICILLIN C-R	Т3	
BICILLIN L-A INTRAMUSCULAR SYRINGE 1,200,000 UNIT/2 ML, 600,000 UNIT/ML	T4	
BIKTARVY	T5	QL (31 EA per 31 days)
caspofungin	T4	
CAYSTON	T5	PA
cefaclor oral capsule 500 mg	T2	
cefadroxil oral capsule	T2	
cefadroxil oral suspension for reconstitution 250 mg/5 ml, 500 mg/5 ml	T2	
cefadroxil oral tablet	T2	
cefazolin injection recon soln 1 gram, 10 gram, 500 mg	T4	
cefdinir oral capsule	T2	
cefepime injection	T4	
cefixime oral capsule	T4	
cefoxitin	T4	
cefpodoxime	T2	
cefprozil	T2	
ceftazidime	T4	

Drug Name	Drug Tier	Requirements/Limits
ceftriaxone injection recon soln 1 gram, 10 gram, 2 gram, 250 mg, 500 mg	T2	
cefuroxime axetil oral tablet	T2	
cefuroxime sodium injection recon soln 750 mg	T4	
cefuroxime sodium intravenous recon soln 1.5 gram	T4	
cephalexin oral capsule 250 mg, 500 mg	T2	
cephalexin oral suspension for reconstitution	T2	
chloroquine phosphate oral tablet 250 mg	Т3	QL (50 EA per 30 days)
chloroquine phosphate oral tablet 500 mg	Т3	QL (25 EA per 30 days)
CIMDUO	T5	QL (31 EA per 31 days)
ciprofloxacin hcl oral tablet 250 mg, 500 mg	T1	
ciprofloxacin hcl oral tablet 750 mg	T2	
ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml	T4	
clarithromycin	T2	
clindamycin hcl	T2	
clindamycin in 5 % dextrose	T4	
CLINDAMYCIN PEDIATRIC	T2	
clindamycin phosphate injection	T4	
clotrimazole mucous membrane	T2	
COARTEM	T4	
colistin (colistimethate na)	T4	
dapsone oral	Т3	
daptomycin	T4	
darunavir	T5	
DELSTRIGO	T5	QL (31 EA per 31 days)
DESCOVY	T5	QL (31 EA per 31 days)
dicloxacillin	T2	
DIFICID ORAL TABLET	T5	QL (20 EA per 10 days)
DOVATO	T5	QL (31 EA per 31 days)
DOXY-100	T4	
doxycycline hyclate intravenous	T4	
doxycycline hyclate oral capsule	T2	
doxycycline hyclate oral tablet 100 mg	T2	
doxycycline hyclate oral tablet,delayed release (dr/ec) 100 mg	Т4	

Drug Name	Drug Tier	Requirements/Limits
doxycycline monohydrate oral capsule 100 mg, 50 mg	T2	
doxycycline monohydrate oral capsule,ir - delay rel,biphase	T4	
doxycycline monohydrate oral tablet 100 mg, 50 mg	T2	
EDURANT	T5	
EDURANT PED	T5	
efavirenz oral tablet	T4	
efavirenz-emtricitabin-tenofov	T5	
efavirenz-lamivu-tenofov disop	T5	QL (31 EA per 31 days)
emtricitabine	T4	
emtricitabine-tenofovir (tdf) oral tablet 100-150 mg, 133-200 mg, 167-250 mg	T5	
emtricitabine-tenofovir (tdf) oral tablet 200-300 mg	T4	
emtricita-rilpivirine-tenof df	T5	
EMTRIVA ORAL SOLUTION	Т3	
EMVERM	T5	
entecavir	T4	
EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG	T5	PA; QL (28 EA per 28 days)
ertapenem	T4	
erythromycin ethylsuccinate oral tablet	T4	
erythromycin oral tablet	T4	
ethambutol	T2	
etravirine	T5	
EVOTAZ	T5	
famciclovir	T2	
fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml	T4	
fluconazole oral suspension for reconstitution	Т3	
fluconazole oral tablet	T2	
flucytosine oral capsule 250 mg	T4	
flucytosine oral capsule 500 mg	T5	
fosamprenavir	T5	

Drug Name	Drug Tier	Requirements/Limits
gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml	T4	
gentamicin injection	T4	
GENVOYA	T5	
griseofulvin microsize	T4	
griseofulvin ultramicrosize oral tablet 125 mg, 250 mg	T4	
hydroxychloroquine oral tablet 200 mg	T2	QL (93 EA per 31 days)
imipenem-cilastatin	T4	
INTELENCE ORAL TABLET 25 MG	T4	
ISENTRESS HD	T5	
ISENTRESS ORAL POWDER IN PACKET	T5	
ISENTRESS ORAL TABLET	T5	
ISENTRESS ORAL TABLET, CHEWABLE 100 MG	T5	
ISENTRESS ORAL TABLET, CHEWABLE 25 MG	Т3	
isoniazid oral	T2	
itraconazole oral capsule	T4	PA
ivermectin oral	T2	PA
JULUCA	T5	
KALETRA ORAL SOLUTION	T5	
ketoconazole oral	T2	
lamivudine	Т3	
lamivudine-zidovudine	Т3	
levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml	Т3	
levofloxacin oral	T2	
linezolid in dextrose 5%	T4	
linezolid oral tablet	T4	
LIVTENCITY	T5	PA; QL (372 EA per 31 days)
lopinavir-ritonavir oral tablet	T4	
maraviroc oral tablet 150 mg	T5	
maraviroc oral tablet 300 mg	T4	
MAVYRET ORAL PELLETS IN PACKET	T5	PA; QL (140 EA per 28 days)
MAVYRET ORAL TABLET	T5	PA; QL (84 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
mefloquine	T2	
meropenem intravenous recon soln 1 gram, 500 mg	Т3	
methenamine hippurate	T2	
metronidazole in nacl (iso-os)	T4	
metronidazole oral tablet 250 mg, 500 mg	T2	
micafungin	T4	
minocycline oral capsule	T2	
minocycline oral tablet	T4	
moxifloxacin oral	T2	
moxifloxacin-sod.chloride(iso)	T4	
nafcillin injection	T4	
neomycin	T2	
nevirapine oral suspension	T4	
nevirapine oral tablet	Т3	
nevirapine oral tablet extended release 24 hr 400 mg	T4	
nitazoxanide	T5	
nitrofurantoin macrocrystal oral capsule 100 mg	T2	QL (90 EA per 365 days)
nitrofurantoin macrocrystal oral capsule 50 mg	T2	QL (180 EA per 365 days)
nitrofurantoin monohyd/m-cryst	T2	QL (90 EA per 365 days)
nitrofurantoin oral suspension 25 mg/5 ml	T5	QL (1800 ML per 365 days)
NORVIR ORAL POWDER IN PACKET	T4	
nystatin oral	T2	
ODEFSEY	T5	QL (31 EA per 31 days)
ofloxacin oral tablet 300 mg, 400 mg	T4	
oseltamivir oral capsule 30 mg	T2	QL (170 EA per 365 days)
oseltamivir oral capsule 45 mg, 75 mg	T2	QL (90 EA per 365 days)
oseltamivir oral suspension for reconstitution	Т3	QL (1080 ML per 365 days)
oxacillin injection recon soln 1 gram	T4	
oxacillin injection recon soln 2 gram	T5	
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (10)- 100 MG (10)	Т3	QL (180 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (6)- 100 MG (5)	Т3	QL (99 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 300 MG (150 MG X 2)-100 MG	Т3	QL (270 EA per 365 days)

Drug Name	Drug Tier	Requirements/Limits
penicillin g pot in dextrose intravenous piggyback 2 million unit/50 ml, 3 million unit/50 ml	Т4	
penicillin g potassium injection recon soln 20 million unit	Т4	
penicillin v potassium	T2	
pentamidine inhalation	T4	PA-BvD
pentamidine injection	T4	
PIFELTRO	T5	QL (62 EA per 31 days)
piperacillin-tazobactam intravenous recon soln 2.25 gram, 3.375 gram, 4.5 gram, 40.5 gram	Т4	
posaconazole oral tablet,delayed release (dr/ec)	T5	PA
praziquantel	T4	
PREVYMIS ORAL PELLETS IN PACKET	T5	PA; QL (124 EA per 31 days)
PREVYMIS ORAL TABLET	T5	QL (31 EA per 31 days)
PREZCOBIX ORAL TABLET 800-150 MG-MG	T5	
PREZISTA ORAL SUSPENSION	T5	
PREZISTA ORAL TABLET 150 MG, 75 MG	T5	
PRIFTIN	Т3	
primaquine	Т3	
pyrazinamide	T4	
pyrimethamine	T5	PA
quinine sulfate	T4	PA; QL (42 EA per 28 days)
RELENZA DISKHALER	Т3	
REYATAZ ORAL POWDER IN PACKET	T5	
ribavirin oral capsule	Т3	
ribavirin oral tablet 200 mg	Т3	
rifabutin	T4	
rifampin intravenous	T5	
rifampin oral	Т3	
rimantadine	T2	
ritonavir	Т3	
RUKOBIA	T5	QL (62 EA per 31 days)
SELZENTRY ORAL SOLUTION	T5	
SIRTURO	T5	PA
sofosbuvir-velpatasvir	T5	PA; QL (28 EA per 28 days)
streptomycin	T5	

Drug Name	Drug Tier	Requirements/Limits
STRIBILD	T5	
sulfadiazine	T4	
sulfamethoxazole-trimethoprim oral suspension	T2	
sulfamethoxazole-trimethoprim oral tablet	T1	
SUNLENCA ORAL	T5	
SYMTUZA	T5	QL (31 EA per 31 days)
TEFLARO	T5	
tenofovir disoproxil fumarate	Т3	
terbinafine hcl oral	T2	QL (90 EA per 180 days)
tetracycline oral capsule	T4	
tigecycline	T5	
tinidazole	T2	
TIVICAY ORAL TABLET 50 MG	T5	
TIVICAY PD	T5	
TOBI PODHALER	T5	PA; QL (224 EA per 56 days)
tobramycin in 0.225 % nacl	T5	PA
tobramycin inhalation	T5	PA
tobramycin sulfate injection solution	T4	
trimethoprim	T2	
TRIUMEQ	T5	
TRIUMEQ PD	T4	QL (186 EA per 31 days)
TYBOST	Т3	
valacyclovir	T2	
valganciclovir oral recon soln	T4	
valganciclovir oral tablet	Т3	
vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg	T4	
vancomycin oral capsule 125 mg	T4	PA; QL (124 EA per 31 days)
vancomycin oral capsule 250 mg	T4	PA; QL (248 EA per 31 days)
VEMLIDY	T5	QL (31 EA per 31 days)
VIRACEPT ORAL TABLET	T5	
VIREAD ORAL POWDER	T5	
VIREAD ORAL TABLET 150 MG, 200 MG, 250 MG	T5	
VIVJOA	T4	PA; QL (18 EA per 84 days)
voriconazole intravenous	T5	PA

Drug Name	Drug Tier	Requirements/Limits
voriconazole oral	T4	
VOSEVI	T5	PA; QL (28 EA per 28 days)
XIFAXAN ORAL TABLET 200 MG	T4	QL (27 EA per 365 days)
XIFAXAN ORAL TABLET 550 MG	T5	PA; QL (62 EA per 31 days)
XOFLUZA ORAL TABLET 40 MG, 80 MG	Т3	QL (9 EA per 365 days)
zidovudine	T2	
Antineoplastic / Immunosuppressant		
Drugs		
abiraterone oral tablet 250 mg	T5	PA-NS; QL (124 EA per 31 days)
abiraterone oral tablet 500 mg	T5	PA-NS; QL (62 EA per 31 days)
ABIRTEGA	Т3	PA-NS; QL (124 EA per 31 days)
AKEEGA	T5	PA-NS; QL (62 EA per 31 days)
ALECENSA	T5	PA-NS; QL (248 EA per 31 days)
ALUNBRIG ORAL TABLET 180 MG, 90 MG	T5	PA-NS; QL (31 EA per 31 days)
ALUNBRIG ORAL TABLET 30 MG	T5	PA-NS; QL (186 EA per 31 days)
ALUNBRIG ORAL TABLETS,DOSE PACK	T5	PA-NS; QL (60 EA per 365 days)
anastrozole	T2	
AUGTYRO ORAL CAPSULE 160 MG	T5	PA-NS; QL (62 EA per 31 days)
AUGTYRO ORAL CAPSULE 40 MG	T5	PA-NS; QL (248 EA per 31 days)
AVMAPKI-FAKZYNJA	T5	PA-NS; QL (66 EA per 28 days)
AYVAKIT	T5	PA-NS; QL (31 EA per 31 days)
azathioprine oral tablet 50 mg	T2	PA-BvD
BALVERSA	T5	PA-NS
bexarotene oral	T5	PA-NS
bexarotene topical	T5	PA-NS; QL (60 GM per 28 days)
bicalutamide	T2	
BOSULIF ORAL CAPSULE 100 MG	T5	PA-NS; QL (186 EA per 31 days)
BOSULIF ORAL CAPSULE 50 MG	T5	PA-NS; QL (341 EA per 31 days)
BOSULIF ORAL TABLET 100 MG	T5	PA-NS; QL (93 EA per 31 days)
BOSULIF ORAL TABLET 400 MG, 500 MG	T5	PA-NS; QL (31 EA per 31 days)
BRAFTOVI	T5	PA-NS; QL (186 EA per 31 days)
BRUKINSA ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
CABOMETYX	T5	PA-NS; QL (31 EA per 31 days)
CALQUENCE (ACALABRUTINIB MAL)	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 300 MG	T5	PA-NS; QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
COMETRIQ ORAL CAPSULE 140 MG/DAY(80 MG X1-20 MG X3)	T5	PA-NS; QL (112 EA per 28 days)
COMETRIQ ORAL CAPSULE 60 MG/DAY (20 MG X 3/DAY)	T5	PA-NS; QL (84 EA per 28 days)
COPIKTRA	T5	PA-NS; QL (62 EA per 31 days)
COTELLIC	T5	PA-NS; LA; QL (63 EA per 28 days)
cyclophosphamide oral	Т3	PA-BvD
cyclosporine modified oral capsule	T2	PA-BvD
cyclosporine modified oral solution	T4	PA-BvD
cyclosporine oral capsule	T2	PA-BvD
DANZITEN	T5	PA-NS; QL (124 EA per 31 days)
dasatinib oral tablet 100 mg, 140 mg, 50 mg, 70 mg, 80 mg	T5	PA-NS; QL (31 EA per 31 days)
dasatinib oral tablet 20 mg	T5	PA-NS; QL (93 EA per 31 days)
DAURISMO ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
DAURISMO ORAL TABLET 25 MG	T5	PA-NS; QL (62 EA per 31 days)
ELIGARD	T4	ST-NS; QL (1 EA per 30 days)
ELIGARD (3 MONTH)	T4	ST-NS; QL (1 EA per 90 days)
ELIGARD (4 MONTH)	T4	ST-NS; QL (1 EA per 120 days)
ELIGARD (6 MONTH)	T4	ST-NS; QL (1 EA per 180 days)
ENVARSUS XR	T4	PA-BvD
ERIVEDGE	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 240 MG	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 60 MG	T5	PA-NS; QL (93 EA per 31 days)
erlotinib	T5	PA-NS; QL (31 EA per 31 days)
EULEXIN	T4	
everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 7.5 mg	T5	PA-NS; QL (31 EA per 31 days)
everolimus (antineoplastic) oral tablet 5 mg	T5	PA-NS; QL (62 EA per 31 days)
everolimus (antineoplastic) oral tablet for suspension 2 mg, 5 mg	Т5	PA-NS; QL (62 EA per 31 days)
everolimus (antineoplastic) oral tablet for suspension 3 mg	T5	PA-NS; QL (93 EA per 31 days)
everolimus (immunosuppressive)	T5	PA-BvD
exemestane	T4	

Drug Name	Drug Tier	Requirements/Limits
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 120 MG	Т5	PA-NS
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 80 MG	T4	PA-NS
FOTIVDA	T5	PA-NS; QL (21 EA per 28 days)
FRUZAQLA ORAL CAPSULE 1 MG	T5	PA-NS; QL (84 EA per 28 days)
FRUZAQLA ORAL CAPSULE 5 MG	T5	PA-NS; QL (21 EA per 28 days)
GAVRETO	T5	PA-NS; QL (124 EA per 31 days)
gefitinib	T5	PA-NS; QL (31 EA per 31 days)
GENGRAF ORAL CAPSULE	T2	PA-BvD
GILOTRIF	T5	PA-NS; QL (31 EA per 31 days)
GLEOSTINE ORAL CAPSULE 10 MG, 100 MG	Т5	PA-NS
GLEOSTINE ORAL CAPSULE 40 MG	T4	PA-NS
GOMEKLI ORAL CAPSULE 1 MG	T5	PA-NS; QL (126 EA per 28 days)
GOMEKLI ORAL CAPSULE 2 MG	T5	PA-NS; QL (84 EA per 28 days)
GOMEKLI ORAL TABLET FOR SUSPENSION	Т5	PA-NS; QL (168 EA per 28 days)
HERNEXEOS	T5	PA-NS; QL (93 EA per 31 days)
hydroxyurea	T2	
IBRANCE	T5	PA-NS; QL (21 EA per 28 days)
IBTROZI	T5	PA-NS; QL (93 EA per 31 days)
ICLUSIG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA ORAL TABLET 50 MG	T5	PA-NS; QL (62 EA per 31 days)
imatinib oral tablet 100 mg	T5	PA-NS; QL (93 EA per 31 days)
imatinib oral tablet 400 mg	T5	PA-NS; QL (62 EA per 31 days)
IMBRUVICA ORAL CAPSULE 140 MG	T5	PA-NS; QL (124 EA per 31 days)
IMBRUVICA ORAL CAPSULE 70 MG	T5	PA-NS; QL (31 EA per 31 days)
IMBRUVICA ORAL SUSPENSION	T5	PA-NS; QL (216 ML per 25 days)
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG	T5	PA-NS; QL (31 EA per 31 days)
IMKELDI	T5	PA-NS; QL (280 ML per 28 days)
INLYTA	T5	PA-NS; QL (124 EA per 31 days)
INQOVI	T5	PA-NS; QL (5 EA per 28 days)
INREBIC	T5	PA-NS; QL (124 EA per 31 days)
ITOVEBI ORAL TABLET 3 MG	T5	PA-NS; QL (62 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
ITOVEBI ORAL TABLET 9 MG	T5	PA-NS; QL (31 EA per 31 days)
IWILFIN	T5	PA-NS; QL (248 EA per 31 days)
JAKAFI	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG	Т5	PA-NS; QL (70 EA per 28 days)
KISQALI FEMARA CO-PACK ORAL TABLET 600 MG/DAY(200 MG X 3)-2.5 MG	T5	PA-NS; QL (91 EA per 28 days)
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (21 EA per 28 days)
KISQALI ORAL TABLET 400 MG/DAY (200 MG X 2)	T5	PA-NS; QL (42 EA per 28 days)
KISQALI ORAL TABLET 600 MG/DAY (200 MG X 3)	T5	PA-NS; QL (63 EA per 28 days)
KOSELUGO ORAL CAPSULE 10 MG	T5	PA-NS; QL (279 EA per 31 days)
KOSELUGO ORAL CAPSULE 25 MG	T5	PA-NS; QL (124 EA per 31 days)
KRAZATI	T5	PA-NS; QL (186 EA per 31 days)
lapatinib	T5	PA-NS; QL (186 EA per 31 days)
LAZCLUZE ORAL TABLET 240 MG	T5	PA-NS; QL (30 EA per 30 days)
LAZCLUZE ORAL TABLET 80 MG	T5	PA-NS; QL (60 EA per 30 days)
lenalidomide	T5	PA-NS; QL (21 EA per 28 days)
LENVIMA	T5	PA-NS
letrozole	T2	
leucovorin calcium oral	Т3	
LEUKERAN	T5	
leuprolide acetate (3 month)	T4	QL (1 EA per 84 days)
leuprolide subcutaneous kit	Т3	QL (2 EA per 28 days)
LONSURF	T5	PA-NS
LORBRENA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
LORBRENA ORAL TABLET 25 MG	T5	PA-NS; QL (93 EA per 31 days)
LUMAKRAS ORAL TABLET 120 MG	T5	PA-NS; QL (124 EA per 31 days)
LUMAKRAS ORAL TABLET 240 MG	T5	PA-NS; QL (62 EA per 31 days)
LUMAKRAS ORAL TABLET 320 MG	T5	PA-NS; QL (93 EA per 31 days)
LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	T5	QL (1 EA per 90 days)

Drug Name	Drug Tier	Requirements/Limits
LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 22.5 MG	T5	QL (1 EA per 84 days)
LUPRON DEPOT (4 MONTH)	T5	QL (1 EA per 112 days)
LUPRON DEPOT (6 MONTH)	T5	QL (1 EA per 168 days)
LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 3.75 MG	T5	QL (1 EA per 30 days)
LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 7.5 MG	T5	QL (1 EA per 28 days)
LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	Т5	PA; QL (1 EA per 90 days)
LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)	T5	PA; QL (1 EA per 30 days)
LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (1 EA per 168 days)
LYNPARZA	T5	PA-NS; QL (124 EA per 31 days)
LYSODREN	T5	
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3)	T5	PA-NS; QL (93 EA per 31 days)
LYTGOBI ORAL TABLET 16 MG/DAY (4 MG X 4)	T5	PA-NS; QL (124 EA per 31 days)
LYTGOBI ORAL TABLET 20 MG/DAY (4 MG X 5)	T5	PA-NS; QL (155 EA per 31 days)
MATULANE	T5	
megestrol oral suspension 400 mg/10 ml (40 mg/ml)	Т3	PA
megestrol oral suspension 625 mg/5 ml (125 mg/ml)	T4	PA
megestrol oral tablet	Т3	PA-NS
MEKINIST ORAL RECON SOLN	T5	PA-NS; QL (1260 ML per 31 days)
MEKINIST ORAL TABLET 0.5 MG	T5	PA-NS; QL (93 EA per 31 days)
MEKINIST ORAL TABLET 2 MG	T5	PA-NS; QL (31 EA per 31 days)
MEKTOVI	T5	PA-NS; QL (186 EA per 31 days)
mercaptopurine oral suspension	T5	
mercaptopurine oral tablet	T2	
mesna oral	T4	
methotrexate sodium	T2	PA-BvD
methotrexate sodium (pf) injection solution	T2	PA-BvD

Drug Name	Drug Tier	Requirements/Limits
MODEYSO	T5	PA-NS; QL (20 EA per 28 days)
mycophenolate mofetil oral capsule	T2	PA-BvD
mycophenolate mofetil oral suspension for reconstitution	T4	PA-BvD
mycophenolate mofetil oral tablet	T2	PA-BvD
mycophenolate sodium	T2	PA-BvD
NEMLUVIO	T5	PA; QL (2 EA per 28 days)
NERLYNX	T5	PA-NS; QL (186 EA per 31 days)
nilotinib hcl	T5	PA-NS; QL (124 EA per 31 days)
nilutamide	T5	
NINLARO	T5	PA-NS; QL (3 EA per 28 days)
NUBEQA	T5	PA-NS; QL (124 EA per 31 days)
octreotide acetate injection solution	T4	PA
ODOMZO	T5	PA-NS; LA; QL (31 EA per 31 days)
OGSIVEO ORAL TABLET 100 MG, 150 MG	T5	PA-NS; QL (62 EA per 31 days)
OGSIVEO ORAL TABLET 50 MG	T5	PA-NS; QL (186 EA per 31 days)
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	Т5	PA-NS; QL (96 ML per 28 days)
OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4)	Т5	PA-NS; QL (16 EA per 28 days)
OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)	T5	PA-NS; QL (20 EA per 28 days)
OJEMDA ORAL TABLET 600 MG/WEEK (100 MG X 6)	T5	PA-NS; QL (24 EA per 28 days)
OJJAARA	T5	PA-NS; QL (31 EA per 31 days)
ONUREG	T5	PA-NS; QL (14 EA per 28 days)
ORGOVYX	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 345 MG	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 86 MG	T5	PA-NS; QL (93 EA per 31 days)
pazopanib oral tablet 200 mg	T5	PA-NS; QL (124 EA per 31 days)
PEMAZYRE	T5	PA-NS; QL (14 EA per 21 days)
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (28 EA per 28 days)
PIQRAY ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	Т5	PA-NS; QL (56 EA per 28 days)
POMALYST	T5	PA-NS; QL (21 EA per 28 days)
L		

Drug Name	Drug Tier	Requirements/Limits
PROGRAF ORAL GRANULES IN PACKET	T4	PA-BvD
QINLOCK	T5	PA-NS; QL (93 EA per 31 days)
RETEVMO ORAL TABLET 120 MG, 160 MG, 80 MG	T5	PA-NS; QL (62 EA per 31 days)
RETEVMO ORAL TABLET 40 MG	T5	PA-NS; QL (93 EA per 31 days)
REVUFORJ ORAL TABLET 110 MG	T5	PA-NS; QL (124 EA per 31 days)
REVUFORJ ORAL TABLET 160 MG	T5	PA-NS; QL (62 EA per 31 days)
REVUFORJ ORAL TABLET 25 MG	T5	PA-NS; QL (248 EA per 31 days)
REZLIDHIA	T5	PA-NS; QL (62 EA per 31 days)
ROMVIMZA	T5	PA-NS; QL (8 EA per 28 days)
ROZLYTREK ORAL CAPSULE 100 MG	T5	PA-NS; QL (155 EA per 31 days)
ROZLYTREK ORAL CAPSULE 200 MG	T5	PA-NS; QL (93 EA per 31 days)
ROZLYTREK ORAL PELLETS IN PACKET	T5	PA-NS; QL (372 EA per 31 days)
RUBRACA	T5	PA-NS; QL (124 EA per 31 days)
RYDAPT	T5	PA-NS; QL (248 EA per 31 days)
SCEMBLIX ORAL TABLET 100 MG	T5	PA-NS; QL (124 EA per 31 days)
SCEMBLIX ORAL TABLET 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SCEMBLIX ORAL TABLET 40 MG	T5	PA-NS; QL (310 EA per 31 days)
SIGNIFOR	T5	PA
sirolimus oral solution	T5	PA-BvD
sirolimus oral tablet	T4	PA-BvD
SOLTAMOX	T4	
sorafenib	T5	PA-NS; QL (124 EA per 31 days)
STIVARGA	T5	PA-NS; QL (84 EA per 28 days)
sunitinib malate	T5	PA-NS; QL (31 EA per 31 days)
TABLOID	T4	
TABRECTA	T5	PA-NS; QL (124 EA per 31 days)
tacrolimus oral capsule	T2	PA-BvD
TAFINLAR ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
TAFINLAR ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (930 EA per 31 days)
TAGRISSO	T5	PA-NS; LA; QL (31 EA per 31 days)
TALZENNA	T5	PA-NS; QL (31 EA per 31 days)
tamoxifen	T2	
TAZVERIK	T5	PA-NS; QL (248 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
ТЕРМЕТКО	T5	PA-NS; QL (62 EA per 31 days)
THALOMID ORAL CAPSULE 100 MG	T5	PA-NS; QL (112 EA per 28 days)
THALOMID ORAL CAPSULE 50 MG	T5	PA-NS; QL (28 EA per 28 days)
TIBSOVO	T5	PA-NS; QL (62 EA per 31 days)
toremifene	T4	
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 11.25 MG	T4	ST-NS; QL (1 EA per 84 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 22.5 MG	T4	ST-NS; QL (1 EA per 168 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 3.75 MG	T4	ST-NS; QL (1 EA per 28 days)
tretinoin (antineoplastic)	T5	
TRUQAP	T5	PA-NS; QL (64 EA per 28 days)
TUKYSA ORAL TABLET 150 MG	T5	PA-NS; QL (124 EA per 31 days)
TUKYSA ORAL TABLET 50 MG	T5	PA-NS; QL (248 EA per 31 days)
TURALIO ORAL CAPSULE 125 MG	T5	PA-NS; QL (124 EA per 31 days)
VANFLYTA	T5	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 10 MG	Т3	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 100 MG	T5	PA-NS; QL (186 EA per 31 days)
VENCLEXTA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
VENCLEXTA STARTING PACK	T5	PA-NS; QL (84 EA per 365 days)
VERZENIO	T5	PA-NS; QL (62 EA per 31 days)
VIJOICE ORAL GRANULES IN PACKET	T5	PA-NS; QL (31 EA per 31 days)
VIJOICE ORAL TABLET 125 MG, 50 MG	T5	PA-NS; QL (28 EA per 28 days)
VIJOICE ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
VITRAKVI ORAL CAPSULE 100 MG	T5	PA-NS; QL (62 EA per 31 days)
VITRAKVI ORAL CAPSULE 25 MG	T5	PA-NS; QL (186 EA per 31 days)
VITRAKVI ORAL SOLUTION	T5	PA-NS; QL (310 ML per 31 days)
VIZIMPRO	T5	PA-NS; QL (31 EA per 31 days)
VONJO	T5	PA-NS; QL (124 EA per 31 days)
VORANIGO ORAL TABLET 10 MG	T5	PA-NS; QL (62 EA per 31 days)
VORANIGO ORAL TABLET 40 MG	T5	PA-NS; QL (31 EA per 31 days)
WELIREG	T5	PA-NS; QL (93 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
XALKORI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XALKORI ORAL PELLET 150 MG	T5	PA-NS; QL (186 EA per 31 days)
XALKORI ORAL PELLET 20 MG, 50 MG	T5	PA-NS; QL (124 EA per 31 days)
XATMEP	T4	PA-BvD
XERMELO	T5	PA; QL (93 EA per 31 days)
XGEVA	T5	PA-NS
XOSPATA	T5	PA-NS; QL (124 EA per 31 days)
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40MG TWICE WEEK (40 MG X 2), 80 MG/WEEK (40 MG X 2)	Т5	PA-NS; QL (8 EA per 28 days)
XPOVIO ORAL TABLET 40 MG/WEEK (10 MG X 4)	T5	PA-NS; QL (16 EA per 28 days)
XPOVIO ORAL TABLET 40 MG/WEEK (40 MG X 1), 60 MG/WEEK (60 MG X 1)	T5	PA-NS; QL (4 EA per 28 days)
XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK)	T5	PA-NS; QL (24 EA per 28 days)
XPOVIO ORAL TABLET 80MG TWICE WEEK (160 MG/WEEK)	T5	PA-NS; QL (32 EA per 28 days)
XTANDI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 40 MG	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 80 MG	T5	PA-NS; QL (62 EA per 31 days)
YONSA	T5	PA-NS; QL (124 EA per 31 days)
ZEJULA ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
ZELBORAF	T5	PA-NS; QL (248 EA per 31 days)
ZOLINZA	T5	PA-NS
ZYDELIG	T5	PA-NS; QL (62 EA per 31 days)
ZYKADIA	T5	PA-NS; QL (93 EA per 31 days)
Autonomic / Cns Drugs, Neurology / Psych		
ABILIFY MAINTENA	T5	QL (1 EA per 28 days)
acetaminophen-codeine oral solution 120-12 mg/5 ml	T2	PA; QL (5167 ML per 31 days)
acetaminophen-codeine oral tablet	T2	PA; QL (403 EA per 31 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML	Т3	PA; QL (1 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 70 MG/ML	Т3	PA; QL (2 ML per 28 days)
AJOVY AUTOINJECTOR	Т3	PA; QL (1.5 ML per 28 days)
AJOVY SYRINGE	Т3	PA; QL (1.5 ML per 28 days)
alprazolam oral tablet 0.25 mg, 0.5 mg	T2	PA; QL (93 EA per 31 days)
alprazolam oral tablet 1 mg, 2 mg	T2	PA; QL (155 EA per 31 days)
amitriptyline	T2	PA-NS
amoxapine	Т3	
aripiprazole oral solution	T4	PA-NS
aripiprazole oral tablet	T2	
aripiprazole oral tablet,disintegrating	T5	PA-NS
armodafinil	T4	PA; QL (31 EA per 31 days)
asenapine maleate	T4	PA-NS; QL (62 EA per 31 days)
atomoxetine oral capsule 10 mg, 25 mg, 40 mg	T4	QL (62 EA per 31 days)
atomoxetine oral capsule 100 mg, 60 mg, 80 mg	T4	QL (31 EA per 31 days)
atomoxetine oral capsule 18 mg	T4	QL (124 EA per 31 days)
AUVELITY	T5	PA-NS; QL (62 EA per 31 days)
baclofen oral tablet 10 mg, 20 mg, 5 mg	T2	
BAFIERTAM	T5	PA; QL (124 EA per 31 days)
benztropine oral	T1	PA
BRIVIACT ORAL SOLUTION	T5	QL (620 ML per 31 days)
BRIVIACT ORAL TABLET	T5	QL (62 EA per 31 days)
bromocriptine	T4	
buprenorphine	T4	PA; QL (4 EA per 28 days)
buprenorphine hcl sublingual tablet 2 mg	Т3	QL (93 EA per 31 days)
buprenorphine hcl sublingual tablet 8 mg	Т3	QL (62 EA per 31 days)
buprenorphine-naloxone sublingual film 12-3 mg, 4-1 mg, 8-2 mg	T2	QL (62 EA per 31 days)
buprenorphine-naloxone sublingual film 2-0.5 mg	T2	QL (93 EA per 31 days)
buprenorphine-naloxone sublingual tablet	T4	QL (93 EA per 31 days)
bupropion hcl oral tablet	T2	
bupropion hcl oral tablet extended release 24 hr 150 mg	T2	QL (93 EA per 31 days)
bupropion hcl oral tablet extended release 24 hr 300 mg	T2	QL (31 EA per 31 days)
bupropion hcl oral tablet sustained-release 12 hr	T2	QL (62 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
buspirone	T2	
butorphanol nasal	T2	QL (5 ML per 28 days)
CAPLYTA	T5	PA-NS; QL (31 EA per 31 days)
carbamazepine oral capsule, er multiphase 12 hr	T2	
carbamazepine oral suspension 100 mg/5 ml	T2	
carbamazepine oral tablet	T2	
carbamazepine oral tablet extended release 12 hr 100 mg, 200 mg	T2	
carbamazepine oral tablet extended release 12 hr 400 mg	Т3	
carbamazepine oral tablet,chewable 100 mg	T2	
carbidopa-levodopa oral tablet	T2	
carbidopa-levodopa oral tablet extended release	T2	
carbidopa-levodopa oral tablet,disintegrating	T2	
carbidopa-levodopa-entacapone	T4	
celecoxib oral capsule 100 mg, 200 mg, 50 mg	T2	QL (62 EA per 31 days)
celecoxib oral capsule 400 mg	Т3	QL (62 EA per 31 days)
chlorpromazine oral	T4	
citalopram oral solution	Т3	
citalopram oral tablet	T1	
clobazam oral suspension	T4	PA-NS; QL (496 ML per 31 days)
clobazam oral tablet	Т3	PA-NS; QL (62 EA per 31 days)
clomipramine	T4	PA-NS
clonazepam oral tablet 0.5 mg	T2	PA-NS; QL (93 EA per 31 days)
clonazepam oral tablet 1 mg	T2	PA-NS; QL (124 EA per 31 days)
clonazepam oral tablet 2 mg	T2	PA-NS; QL (310 EA per 31 days)
clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg	T2	PA-NS; QL (93 EA per 31 days)
clonazepam oral tablet,disintegrating 1 mg	T2	PA-NS; QL (124 EA per 31 days)
clonazepam oral tablet, disintegrating 2 mg	T2	PA-NS; QL (310 EA per 31 days)
clorazepate dipotassium oral tablet 15 mg	Т3	PA-NS; QL (186 EA per 31 days)
clorazepate dipotassium oral tablet 3.75 mg, 7.5 mg	Т3	PA-NS; QL (93 EA per 31 days)
clozapine oral tablet 100 mg, 25 mg	T2	QL (279 EA per 31 days)
clozapine oral tablet 200 mg	T2	QL (124 EA per 31 days)
clozapine oral tablet 50 mg	T2	QL (93 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
clozapine oral tablet, disintegrating 100 mg, 25 mg	T4	QL (279 EA per 31 days)
clozapine oral tablet,disintegrating 12.5 mg	T4	QL (93 EA per 31 days)
clozapine oral tablet,disintegrating 150 mg	T4	QL (186 EA per 31 days)
clozapine oral tablet,disintegrating 200 mg	T4	QL (124 EA per 31 days)
COBENFY	T5	PA-NS; QL (62 EA per 31 days)
COBENFY STARTER PACK	T5	PA-NS; QL (112 EA per 365 days)
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
COPAXONE SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
cyclobenzaprine oral tablet 10 mg	T2	QL (93 EA per 31 days)
cyclobenzaprine oral tablet 5 mg	T2	QL (155 EA per 31 days)
dalfampridine	T5	PA; QL (62 EA per 31 days)
dantrolene oral	T2	
DAYBUE	T5	PA; QL (3600 ML per 30 days)
desipramine	T2	
desvenlafaxine succinate	T2	QL (31 EA per 31 days)
dexmethylphenidate oral capsule,er biphasic 50- 50	T2	QL (31 EA per 31 days)
dexmethylphenidate oral tablet 10 mg	T2	QL (62 EA per 31 days)
dexmethylphenidate oral tablet 2.5 mg, 5 mg	T2	QL (93 EA per 31 days)
dextroamphetamine-amphetamine oral capsule,extended release 24hr	T2	QL (31 EA per 31 days)
dextroamphetamine-amphetamine oral tablet 10 mg, 12.5 mg, 15 mg, 30 mg, 5 mg, 7.5 mg	T2	QL (62 EA per 31 days)
dextroamphetamine-amphetamine oral tablet 20 mg	T2	QL (93 EA per 31 days)
DIACOMIT ORAL CAPSULE 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL CAPSULE 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIAZEPAM INTENSOL	T2	PA-NS; QL (248 ML per 31 days)
diazepam oral solution 5 mg/5 ml (1 mg/ml)	T2	PA-NS; QL (1500 ML per 31 days)
diazepam oral tablet	T2	PA-NS; QL (124 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
diazepam rectal	T4	
diclofenac potassium oral tablet 50 mg	T2	
diclofenac sodium oral	T2	
diclofenac sodium topical drops	T4	
diflunisal	T2	
dihydroergotamine nasal	T5	PA; QL (8 ML per 28 days)
DILANTIN	T3	
dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg (14)- 240 mg (46)	T5	PA; QL (120 EA per 365 days)
dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 240 mg	T5	PA; QL (62 EA per 31 days)
divalproex	T2	
donepezil oral tablet 10 mg, 5 mg	T1	
donepezil oral tablet 23 mg	T3	QL (31 EA per 31 days)
donepezil oral tablet, disintegrating	T2	
doxepin oral capsule	T2	PA-NS
doxepin oral concentrate	T2	PA-NS
doxepin oral tablet	T3	PA
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG	T4	PA-NS; QL (93 EA per 31 days)
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 30 MG, 60 MG	T4	PA-NS; QL (62 EA per 31 days)
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 40 MG	T4	PA-NS; QL (31 EA per 31 days)
duloxetine oral capsule,delayed release(dr/ec) 20 mg, 60 mg	T2	QL (62 EA per 31 days)
duloxetine oral capsule,delayed release(dr/ec) 30 mg	T2	QL (31 EA per 31 days)
EMGALITY PEN	Т3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML	Т3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 300 MG/3 ML (100 MG/ML X 3)	T5	PA; QL (3 ML per 28 days)
EMSAM	T5	QL (30 EA per 30 days)
ENDOCET	Т3	PA; QL (372 EA per 31 days)
entacapone	Т3	
EPIDIOLEX	T5	PA-NS
ergotamine-caffeine	T3	PA

Drug Name	Drug Tier	Requirements/Limits
escitalopram oxalate oral solution	T4	QL (620 ML per 31 days)
escitalopram oxalate oral tablet 10 mg	T1	QL (45 EA per 30 days)
escitalopram oxalate oral tablet 20 mg, 5 mg	T1	QL (30 EA per 30 days)
eslicarbazepine oral tablet 200 mg	T4	QL (186 EA per 31 days)
eslicarbazepine oral tablet 400 mg	T5	QL (93 EA per 31 days)
eslicarbazepine oral tablet 600 mg, 800 mg	T5	QL (62 EA per 31 days)
eszopiclone	T4	PA; QL (31 EA per 31 days)
ethosuximide	T2	
etodolac	T2	
EVRYSDI ORAL RECON SOLN	T5	PA; QL (240 ML per 31 days)
EVRYSDI ORAL TABLET	T5	PA; QL (31 EA per 31 days)
FANAPT ORAL TABLET 1 MG	T4	PA-NS; QL (62 EA per 31 days)
FANAPT ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG	T5	PA-NS; QL (62 EA per 31 days)
FANAPT TITRATION PACK A	T4	PA-NS; QL (16 EA per 365 days)
felbamate	T4	
fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 75 mcg/hr	T4	PA; QL (10 EA per 30 days)
fentanyl transdermal patch 72 hour 25 mcg/hr, 50 mcg/hr	T2	PA; QL (10 EA per 30 days)
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)	Т3	PA-NS; QL (56 EA per 365 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 40 MG, 80 MG	Т3	PA-NS; QL (31 EA per 31 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 20 MG	Т3	PA-NS; QL (93 EA per 31 days)
fingolimod	T5	PA; QL (31 EA per 31 days)
FINTEPLA	T5	PA-NS; QL (360 ML per 30 days)
FIRDAPSE	T5	PA; QL (248 EA per 31 days)
fluoxetine (pmdd)	T2	
fluoxetine oral capsule	T1	
fluoxetine oral solution	Т3	
fluoxetine oral tablet 10 mg, 20 mg	T2	
fluphenazine decanoate	T2	
fluphenazine hcl injection	T4	
fluphenazine hcl oral concentrate	T4	
fluphenazine hcl oral tablet	T4	

Drug Name	Drug Tier	Requirements/Limits
flurbiprofen oral tablet 100 mg	T2	
fluvoxamine oral tablet	T2	
FYCOMPA ORAL SUSPENSION	T5	QL (744 ML per 31 days)
gabapentin oral capsule 100 mg, 400 mg	T1	PA-NS; QL (270 EA per 30 days)
gabapentin oral capsule 300 mg	T1	PA-NS; QL (360 EA per 30 days)
gabapentin oral solution 250 mg/5 ml	T2	PA-NS; QL (2160 ML per 30 days)
gabapentin oral tablet 600 mg	T1	PA-NS; QL (180 EA per 30 days)
gabapentin oral tablet 800 mg	T1	PA-NS; QL (120 EA per 30 days)
galantamine oral capsule,ext rel. pellets 24 hr	Т3	
galantamine oral solution	T4	
galantamine oral tablet 12 mg, 8 mg	Т3	
galantamine oral tablet 4 mg	T2	
glatiramer subcutaneous syringe 20 mg/ml	T5	PA; QL (31 ML per 31 days)
glatiramer subcutaneous syringe 40 mg/ml	T5	PA; QL (12 ML per 28 days)
GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
GLATOPA SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
guanfacine oral tablet extended release 24 hr	T2	PA
haloperidol	T2	
haloperidol decanoate	T2	
haloperidol lactate injection	T2	
haloperidol lactate oral	T2	
hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg	T2	PA; QL (372 EA per 31 days)
hydromorphone oral liquid	T4	PA; QL (1240 ML per 31 days)
hydromorphone oral tablet 2 mg, 4 mg	T2	PA; QL (186 EA per 31 days)
hydromorphone oral tablet 8 mg	Т3	PA; QL (155 EA per 31 days)
IBU ORAL TABLET 600 MG, 800 MG	T1	
ibuprofen oral tablet 300 mg	T4	
ibuprofen oral tablet 400 mg, 600 mg, 800 mg	T1	
imipramine hcl	T4	PA-NS
indomethacin oral capsule	T2	
indomethacin oral capsule, extended release	T2	
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,092 MG/3.5 ML	Т5	QL (3.5 ML per 180 days)

Drug Name	Drug Tier	Requirements/Limits
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,560 MG/5 ML	T5	QL (5 ML per 180 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 117 MG/0.75 ML	T5	QL (0.75 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 156 MG/ML	T5	QL (1 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 234 MG/1.5 ML	T5	QL (1.5 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 39 MG/0.25 ML	Т3	QL (0.25 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 78 MG/0.5 ML	T5	QL (0.5 ML per 28 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 273 MG/0.88 ML	T5	QL (0.88 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 410 MG/1.32 ML	T5	QL (1.32 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 546 MG/1.75 ML	T5	QL (1.75 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 819 MG/2.63 ML	T5	QL (2.63 ML per 84 days)
KESIMPTA PEN	T5	PA; QL (0.4 ML per 28 days)
KLOXXADO	Т3	
lacosamide oral	T4	
lamotrigine oral tablet	T1	
lamotrigine oral tablet extended release 24hr	T4	
lamotrigine oral tablet, chewable dispersible	T2	
levetiracetam oral solution 100 mg/ml	T2	
levetiracetam oral tablet	T2	
levetiracetam oral tablet extended release 24 hr	T2	
lithium carbonate oral capsule	T1	
lithium carbonate oral tablet	T1	
lithium carbonate oral tablet extended release	T2	
lithium citrate	T2	
lofexidine	T5	
LORAZEPAM INTENSOL	T2	PA; QL (155 ML per 31 days)
lorazepam oral tablet 0.5 mg	T2	PA; QL (124 EA per 31 days)
lorazepam oral tablet 1 mg	T2	PA; QL (186 EA per 31 days)
lorazepam oral tablet 2 mg	T2	PA; QL (155 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
loxapine succinate	T2	
lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg	T4	PA-NS; QL (31 EA per 31 days)
lurasidone oral tablet 80 mg	T4	PA-NS; QL (62 EA per 31 days)
MARPLAN	T4	
meloxicam oral tablet	T1	
memantine oral capsule,sprinkle,er 24hr	T2	
memantine oral solution	T2	
memantine oral tablet	T2	
memantine-donepezil	Т3	PA
methadone oral solution 10 mg/5 ml	T2	PA; QL (620 ML per 31 days)
methadone oral solution 5 mg/5 ml	T2	PA; QL (1240 ML per 31 days)
methadone oral tablet 10 mg	T2	PA; QL (124 EA per 31 days)
methadone oral tablet 5 mg	T2	PA; QL (248 EA per 31 days)
methsuximide	T4	
methylphenidate hcl oral tablet	T2	QL (93 EA per 31 days)
mirtazapine oral tablet 15 mg, 30 mg, 45 mg	T1	
mirtazapine oral tablet 7.5 mg	T2	
mirtazapine oral tablet,disintegrating	T2	
modafinil	Т3	PA; QL (31 EA per 31 days)
molindone	T2	
morphine concentrate oral solution	T2	PA; QL (310 ML per 31 days)
morphine oral solution 10 mg/5 ml	T2	PA; QL (2800 ML per 31 days)
morphine oral solution 20 mg/5 ml (4 mg/ml)	T2	PA; QL (1400 ML per 31 days)
morphine oral tablet	T2	PA; QL (186 EA per 31 days)
morphine oral tablet extended release 100 mg	T2	PA; QL (62 EA per 31 days)
morphine oral tablet extended release 15 mg, 30 mg, 60 mg	T2	PA; QL (100 EA per 31 days)
morphine oral tablet extended release 200 mg	Т3	PA; QL (31 EA per 31 days)
nabumetone	T2	
naloxone injection solution	T2	
naloxone injection syringe	T2	
naltrexone	T2	
NAMZARIC ORAL CAPSULE,SPRINKLE,ER 24HR 7-10 MG	Т3	PA
naproxen oral tablet	T1	

Drug Name	Drug Tier	Requirements/Limits
naproxen oral tablet,delayed release (dr/ec) 375 mg	T2	
naproxen oral tablet,delayed release (dr/ec) 500 mg	T4	
naproxen sodium oral tablet 550 mg	T2	
naratriptan oral tablet 1 mg	Т3	QL (20 EA per 28 days)
naratriptan oral tablet 2.5 mg	Т3	QL (9 EA per 28 days)
NAYZILAM	T4	PA-NS; QL (10 EA per 30 days)
nefazodone	T2	
NEUPRO	T4	
nortriptyline	T2	
NUEDEXTA	T5	PA; QL (62 EA per 31 days)
NUPLAZID	T5	PA-NS; QL (31 EA per 31 days)
NURTEC ODT	Т3	PA; QL (18 EA per 28 days)
olanzapine intramuscular	T4	
olanzapine oral tablet	T2	QL (31 EA per 31 days)
olanzapine oral tablet, disintegrating	T4	QL (31 EA per 31 days)
OPIPZA	T5	PA-NS
oxaprozin oral tablet	T4	
oxcarbazepine oral suspension	T2	
oxcarbazepine oral tablet	T2	
oxycodone oral capsule	T2	PA; QL (186 EA per 31 days)
oxycodone oral concentrate	T4	PA; QL (180 ML per 31 days)
oxycodone oral tablet 10 mg, 15 mg, 20 mg, 5 mg	T2	PA; QL (186 EA per 31 days)
oxycodone oral tablet 30 mg	Т3	PA; QL (138 EA per 31 days)
oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg	Т3	PA; QL (372 EA per 31 days)
oxycodone-acetaminophen oral tablet 5-325 mg, 7.5-325 mg	T2	PA; QL (372 EA per 31 days)
paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 9 mg	T4	QL (31 EA per 31 days)
paliperidone oral tablet extended release 24hr 6 mg	T4	QL (62 EA per 31 days)
paroxetine hcl oral suspension	T4	
paroxetine hcl oral tablet	T1	
paroxetine hcl oral tablet extended release 24 hr	T4	

Drug Name	Drug Tier	Requirements/Limits
perampanel oral tablet 10 mg, 12 mg, 4 mg, 6 mg, 8 mg	T5	QL (31 EA per 31 days)
perampanel oral tablet 2 mg	T4	QL (31 EA per 31 days)
perphenazine	T2	
PERSERIS	T5	QL (1 EA per 28 days)
phenelzine	Т3	
phenobarbital	T2	PA-NS
PHENYTEK	T4	
phenytoin oral suspension 125 mg/5 ml	T2	
phenytoin oral tablet,chewable	T2	
phenytoin sodium extended oral capsule 100 mg	T2	
pimozide	T4	
piroxicam	T2	
pramipexole oral tablet	T2	
pregabalin oral capsule 100 mg, 150 mg, 200 mg, 25 mg, 50 mg, 75 mg	T2	PA-NS; QL (93 EA per 31 days)
pregabalin oral capsule 225 mg, 300 mg	T2	PA-NS; QL (62 EA per 31 days)
pregabalin oral solution	T2	PA-NS; QL (930 ML per 31 days)
primidone oral tablet 125 mg	T4	
primidone oral tablet 250 mg, 50 mg	T2	
protriptyline	T4	
pyridostigmine bromide oral tablet 60 mg	T2	
pyridostigmine bromide oral tablet extended release 180 mg	Т3	
quetiapine oral tablet 100 mg, 200 mg, 25 mg, 300 mg, 400 mg, 50 mg	T2	QL (62 EA per 31 days)
quetiapine oral tablet 150 mg	Т3	QL (62 EA per 31 days)
quetiapine oral tablet extended release 24 hr	T2	QL (62 EA per 31 days)
QULIPTA	Т3	PA; QL (31 EA per 31 days)
RADICAVA ORS STARTER KIT SUSP	T5	PA; QL (70 ML per 28 days)
RALDESY	T5	
ramelteon	T2	QL (31 EA per 31 days)
rasagiline	T4	
REXULTI ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
risperidone microspheres intramuscular suspension,extended rel recon 12.5 mg/2 ml, 25 mg/2 ml	Т3	QL (2 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
risperidone microspheres intramuscular suspension, extended rel recon 37.5 mg/2 ml	T4	QL (2 EA per 28 days)
risperidone microspheres intramuscular suspension, extended rel recon 50 mg/2 ml	T5	QL (2 EA per 28 days)
risperidone oral solution	T2	QL (496 ML per 31 days)
risperidone oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg	T1	QL (31 EA per 31 days)
risperidone oral tablet 3 mg	T1	QL (93 EA per 31 days)
risperidone oral tablet 4 mg	T1	QL (124 EA per 31 days)
risperidone oral tablet, disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg	T4	QL (31 EA per 31 days)
risperidone oral tablet, disintegrating 3 mg	T4	QL (93 EA per 31 days)
risperidone oral tablet, disintegrating 4 mg	T4	QL (124 EA per 31 days)
rivastigmine	Т3	QL (30 EA per 30 days)
rivastigmine tartrate	T2	
rizatriptan oral tablet 10 mg	T2	QL (12 EA per 28 days)
rizatriptan oral tablet 5 mg	T2	QL (24 EA per 28 days)
rizatriptan oral tablet,disintegrating 10 mg	T2	QL (12 EA per 28 days)
rizatriptan oral tablet, disintegrating 5 mg	T2	QL (24 EA per 28 days)
ropinirole oral tablet	T2	
ropinirole oral tablet extended release 24 hr	T4	
ROWEEPRA ORAL TABLET 500 MG	T2	
rufinamide oral suspension	T5	PA-NS
rufinamide oral tablet 200 mg	T4	PA-NS
rufinamide oral tablet 400 mg	T5	PA-NS
RYTARY	Т3	ST
SECUADO	T5	PA-NS; QL (31 EA per 31 days)
selegiline hcl	T2	
sertraline oral concentrate	T2	
sertraline oral tablet	T1	
SKYCLARYS	T5	PA; QL (93 EA per 31 days)
sodium oxybate	T5	PA; QL (540 ML per 30 days)
SPRITAM ORAL TABLET FOR SUSPENSION 250 MG, 500 MG	T4	
SUBVENITE ORAL TABLET	T2	
sulindac	T2	

Drug Name	Drug Tier	Requirements/Limits
sumatriptan nasal spray,non-aerosol 20 mg/actuation	T4	QL (8 EA per 28 days)
sumatriptan nasal spray,non-aerosol 5 mg/actuation	T4	QL (32 EA per 28 days)
sumatriptan succinate oral tablet 100 mg	T2	QL (9 EA per 28 days)
sumatriptan succinate oral tablet 25 mg	T2	QL (36 EA per 28 days)
sumatriptan succinate oral tablet 50 mg	T2	QL (18 EA per 28 days)
sumatriptan succinate subcutaneous cartridge 6 mg/0.5 ml	T4	QL (4 ML per 28 days)
sumatriptan succinate subcutaneous pen injector 4 mg/0.5 ml	T4	QL (6 ML per 28 days)
sumatriptan succinate subcutaneous pen injector 6 mg/0.5 ml	T4	QL (4 ML per 28 days)
sumatriptan succinate subcutaneous solution	T4	QL (4 ML per 28 days)
SYMPAZAN ORAL FILM 10 MG, 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SYMPAZAN ORAL FILM 5 MG	T4	PA-NS; QL (62 EA per 31 days)
TASCENSO ODT	T5	PA; QL (31 EA per 31 days)
tasimelteon	T5	PA; QL (31 EA per 31 days)
teriflunomide	T5	PA; QL (31 EA per 31 days)
tetrabenazine oral tablet 12.5 mg	T5	PA; QL (93 EA per 31 days)
tetrabenazine oral tablet 25 mg	T5	PA; QL (124 EA per 31 days)
thioridazine	Т3	
thiothixene	Т3	
tiagabine	T4	
tizanidine oral tablet	T2	
topiramate oral capsule, sprinkle 15 mg, 25 mg	T2	
topiramate oral solution	T4	PA-NS; QL (496 ML per 31 days)
topiramate oral tablet	T1	
tramadol oral tablet 50 mg	T2	PA; QL (240 EA per 30 days)
tramadol-acetaminophen	T2	PA; QL (372 EA per 31 days)
tranylcypromine	T4	
trazodone oral tablet 100 mg, 150 mg, 50 mg	T1	
trifluoperazine	T2	
trimipramine	T4	PA-NS
TRINTELLIX	T4	PA-NS
UBRELVY ORAL TABLET 100 MG	T5	PA; QL (17 EA per 28 days)
UBRELVY ORAL TABLET 50 MG	T5	PA; QL (34 EA per 28 days)

Drug Name	Drug Tier	Requirements/Limits
valproic acid	T2	
valproic acid (as sodium salt) oral solution 250 mg/5 ml	T2	
VALTOCO NASAL SPRAY,NON-AEROSOL 10 MG/SPRAY (0.1 ML), 5 MG/SPRAY (0.1 ML)	T4	PA-NS; QL (10 EA per 30 days)
VALTOCO NASAL SPRAY,NON-AEROSOL 15 MG/2 SPRAY (7.5/0.1ML X 2), 20 MG/2 SPRAY (10MG/0.1ML X2)	Т5	PA-NS; QL (10 EA per 30 days)
venlafaxine oral capsule,extended release 24hr 150 mg, 37.5 mg	T2	QL (31 EA per 31 days)
venlafaxine oral capsule,extended release 24hr 75 mg	T2	QL (93 EA per 31 days)
venlafaxine oral tablet	T2	
VERSACLOZ	T5	QL (558 ML per 31 days)
vigabatrin	T5	PA-NS
VIGADRONE	T5	PA-NS
VIGPODER	T5	PA-NS
vilazodone	Т3	PA-NS; QL (31 EA per 31 days)
VIVITROL	T5	
VRAYLAR ORAL CAPSULE	T5	PA-NS; QL (31 EA per 31 days)
VUMERITY	T5	PA; QL (124 EA per 31 days)
XCOPRI	T5	PA-NS
XCOPRI MAINTENANCE PACK	T5	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 12.5 MG (14)- 25 MG (14)	T4	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 150 MG (14)- 200 MG (14), 50 MG (14)- 100 MG (14)	Т5	PA-NS
XYREM	T5	PA; QL (540 ML per 30 days)
zaleplon oral capsule 10 mg	T4	PA; QL (62 EA per 31 days)
zaleplon oral capsule 5 mg	T4	PA; QL (93 EA per 31 days)
ZAVZPRET	T5	PA; QL (8 EA per 30 days)
ZEPOSIA	T5	PA; QL (31 EA per 31 days)
ZEPOSIA STARTER KIT (28-DAY)	T5	PA; QL (56 EA per 365 days)
ZEPOSIA STARTER PACK (7-DAY)	T5	PA; QL (14 EA per 365 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML	T5	PA; QL (11.648 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
ZILBRYSQ SUBCUTANEOUS SYRINGE 23 MG/0.574 ML	T5	PA; QL (16.072 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 32.4 MG/0.81 ML	T5	PA; QL (22.68 ML per 28 days)
ziprasidone hcl	T2	QL (62 EA per 31 days)
ziprasidone mesylate	T4	
zolpidem oral tablet	T2	PA; QL (31 EA per 31 days)
ZONISADE	T4	PA-NS; QL (930 ML per 31 days)
zonisamide	T2	
ZTALMY	T5	PA-NS; QL (1100 ML per 30 days)
ZUBSOLV SUBLINGUAL TABLET 0.7-0.18 MG, 2.9-0.71 MG, 8.6-2.1 MG	Т3	QL (62 EA per 31 days)
ZUBSOLV SUBLINGUAL TABLET 1.4-0.36 MG	Т3	QL (93 EA per 31 days)
ZUBSOLV SUBLINGUAL TABLET 11.4-2.9 MG, 5.7-1.4 MG	Т3	QL (31 EA per 31 days)
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG	T5	PA-NS; QL (28 EA per 180 days)
ZURZUVAE ORAL CAPSULE 30 MG	T5	PA-NS; QL (14 EA per 180 days)
Cardiovascular, Hypertension / Lipids		
acebutolol	T2	
aliskiren	T4	
amiloride	T2	
amiloride-hydrochlorothiazide	T2	
amiodarone oral	T2	
amlodipine	T1	
amlodipine-atorvastatin	T2	
amlodipine-benazepril	T1	
amlodipine-olmesartan	T2	QL (31 EA per 31 days)
amlodipine-valsartan	T1	
aspirin-dipyridamole	T4	
atenolol	T1	
atenolol-chlorthalidone	T2	
atorvastatin	T1	
benazepril	T1	
benazepril-hydrochlorothiazide	T1	

Drug Name	Drug Tier	Requirements/Limits
bisoprolol fumarate oral tablet 10 mg, 5 mg	T2	
bisoprolol fumarate oral tablet 2.5 mg	Т3	
bisoprolol-hydrochlorothiazide	T1	
bumetanide oral	T2	
CABLIVI INJECTION KIT	T5	PA; QL (31 EA per 31 days)
CAMZYOS	T5	PA; QL (31 EA per 31 days)
candesartan	T2	
candesartan-hydrochlorothiazid	T2	
captopril	T2	
CARTIA XT	T2	
carvedilol	T1	
chlorthalidone oral tablet 25 mg, 50 mg	T2	
cholestyramine (with sugar) oral powder in packet	T2	
CHOLESTYRAMINE LIGHT ORAL POWDER IN PACKET	T2	
cilostazol	T2	
clonidine	T4	
clonidine hcl oral tablet	T1	
clopidogrel oral tablet 75 mg	T1	
colesevelam	T4	
colestipol oral packet	T4	
colestipol oral tablet	Т3	
CORLANOR ORAL SOLUTION	Т3	PA; QL (420 ML per 28 days)
digoxin oral solution	Т3	QL (155 ML per 31 days)
digoxin oral tablet 125 mcg (0.125 mg)	T2	QL (62 EA per 31 days)
digoxin oral tablet 250 mcg (0.25 mg)	T2	QL (31 EA per 31 days)
diltiazem hcl oral capsule,extended release 12 hr	T2	
diltiazem hcl oral capsule,extended release 24 hr 360 mg, 420 mg	T2	
diltiazem hcl oral capsule,extended release 24hr 120 mg, 180 mg, 240 mg, 300 mg	T2	
diltiazem hcl oral tablet	T1	
diltiazem hcl oral tablet extended release 24 hr	T2	
DILT-XR	T2	
dofetilide	T4	
DOPTELET (10 TAB PACK)	T5	PA

Drug Name	Drug Tier	Requirements/Limits
DOPTELET (15 TAB PACK)	T5	PA
DOPTELET (30 TAB PACK)	T5	PA
doxazosin	T1	
ELIQUIS DVT-PE TREAT 30D START	Т3	QL (74 EA per 30 days)
ELIQUIS ORAL TABLET 2.5 MG	Т3	QL (60 EA per 30 days)
ELIQUIS ORAL TABLET 5 MG	Т3	QL (74 EA per 30 days)
eltrombopag olamine oral powder in packet 12.5 mg	T5	PA; QL (372 EA per 31 days)
eltrombopag olamine oral powder in packet 25 mg	T5	PA; QL (31 EA per 31 days)
eltrombopag olamine oral tablet 12.5 mg, 25 mg	T5	PA; QL (31 EA per 31 days)
eltrombopag olamine oral tablet 50 mg, 75 mg	T5	PA; QL (62 EA per 31 days)
enalapril maleate oral tablet	T1	
enalapril-hydrochlorothiazide	T1	
enoxaparin subcutaneous syringe	T4	
ENTRESTO ORAL TABLET 24-26 MG	Т3	QL (186 EA per 31 days)
ENTRESTO ORAL TABLET 49-51 MG	Т3	QL (93 EA per 31 days)
ENTRESTO ORAL TABLET 97-103 MG	Т3	QL (62 EA per 31 days)
eplerenone	T2	
ethacrynic acid	T4	
ezetimibe	T2	
ezetimibe-simvastatin	Т3	QL (31 EA per 31 days)
felodipine	T2	
fenofibrate micronized oral capsule 134 mg, 200 mg, 67 mg	T2	
fenofibrate nanocrystallized	T2	
fenofibrate oral tablet 160 mg, 54 mg	T2	
flecainide	T2	
fluvastatin oral capsule	T4	
fondaparinux subcutaneous syringe 10 mg/0.8 ml, 7.5 mg/0.6 ml	T5	
fondaparinux subcutaneous syringe 2.5 mg/0.5 ml, 5 mg/0.4 ml	T4	
fosinopril	T1	
fosinopril-hydrochlorothiazide	T2	
FUROSCIX	T5	PA; QL (8 EA per 30 days)
furosemide injection solution	T2	

Drug Name	Drug Tier	Requirements/Limits
furosemide oral solution 10 mg/ml, 40 mg/5 ml (8 mg/ml)	T2	
furosemide oral tablet	T1	
gemfibrozil	T1	
heparin (porcine) injection solution	Т3	PA-BvD
hydralazine oral	T2	
hydrochlorothiazide	T1	
icosapent ethyl oral capsule 0.5 gram	T2	QL (248 EA per 31 days)
icosapent ethyl oral capsule 1 gram	T2	QL (124 EA per 31 days)
indapamide	T1	
irbesartan	T1	QL (31 EA per 31 days)
irbesartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
isosorbide dinitrate oral tablet 10 mg, 20 mg, 30 mg, 5 mg	T2	
isosorbide mononitrate oral tablet	T2	
isosorbide mononitrate oral tablet extended release 24 hr	T1	
isradipine	T2	
ivabradine oral tablet 5 mg	Т3	PA; QL (93 EA per 31 days)
ivabradine oral tablet 7.5 mg	Т3	PA; QL (62 EA per 31 days)
JANTOVEN	T1	
KERENDIA	T4	PA; QL (31 EA per 31 days)
labetalol oral tablet 100 mg, 200 mg, 300 mg	T2	
lisinopril	T1	
lisinopril-hydrochlorothiazide	T1	
losartan oral tablet 100 mg	T1	QL (31 EA per 31 days)
losartan oral tablet 25 mg	T1	QL (93 EA per 31 days)
losartan oral tablet 50 mg	T1	QL (62 EA per 31 days)
losartan-hydrochlorothiazide	T1	
lovastatin	T1	
metolazone	T2	
metoprolol succinate	T1	
metoprolol ta-hydrochlorothiaz	T2	
metoprolol tartrate oral tablet 100 mg, 25 mg, 50 mg	T1	
metoprolol tartrate oral tablet 37.5 mg, 75 mg	T2	
metyrosine	T5	PA

Drug Name	Drug Tier	Requirements/Limits
mexiletine	Т3	
minoxidil oral	T2	
moexipril	T1	
MULPLETA	T5	PA
MULTAQ	T4	
nadolol	T2	
nebivolol oral tablet 10 mg, 2.5 mg	T2	QL (93 EA per 31 days)
nebivolol oral tablet 20 mg	T2	QL (62 EA per 31 days)
nebivolol oral tablet 5 mg	T2	QL (217 EA per 31 days)
NEXLETOL	Т3	PA; QL (31 EA per 31 days)
NEXLIZET	T4	PA; QL (31 EA per 31 days)
niacin oral tablet extended release 24 hr 1,000 mg, 750 mg	T2	
niacin oral tablet extended release 24 hr 500 mg	T2	QL (31 EA per 31 days)
nicardipine oral capsule 20 mg	T4	
nicardipine oral capsule 30 mg	T5	
nifedipine oral tablet extended release	T2	
nifedipine oral tablet extended release 24hr	T2	
nimodipine oral capsule	T4	
NITRO-BID	T2	
nitroglycerin sublingual	T2	
nitroglycerin transdermal patch 24 hour	T2	
nitroglycerin translingual	T4	
olmesartan oral tablet 20 mg, 40 mg	T1	QL (31 EA per 31 days)
olmesartan oral tablet 5 mg	T1	QL (93 EA per 31 days)
olmesartan-amlodipin-hcthiazid	Т3	
olmesartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
omega-3 acid ethyl esters	T2	QL (124 EA per 31 days)
PACERONE ORAL TABLET 100 MG, 200 MG, 400 MG	T2	
pentoxifylline	T2	
perindopril erbumine	T1	
pindolol	Т3	
pitavastatin calcium	Т3	
prasugrel hcl	T2	
pravastatin	T1	

Drug Name	Drug Tier	Requirements/Limits
prazosin	T2	
PREVALITE ORAL POWDER IN PACKET	T2	
propafenone oral capsule,extended release 12 hr	T4	
propafenone oral tablet	T2	
propranolol oral capsule,extended release 24 hr	T2	
propranolol oral solution	T2	
propranolol oral tablet	T1	
quinapril	T1	
quinapril-hydrochlorothiazide	T1	
quinidine sulfate oral tablet	T2	
ramipril	T1	
ranolazine	T4	QL (62 EA per 31 days)
REPATHA SURECLICK	Т3	PA; QL (3 ML per 28 days)
REPATHA SYRINGE	Т3	PA; QL (3 ML per 28 days)
rosuvastatin	T1	
sacubitril-valsartan oral tablet 24-26 mg	T2	QL (186 EA per 31 days)
sacubitril-valsartan oral tablet 49-51 mg	T2	QL (93 EA per 31 days)
sacubitril-valsartan oral tablet 97-103 mg	T2	QL (62 EA per 31 days)
simvastatin	T1	
SOTALOL AF	T2	
sotalol oral	T2	
spironolactone oral tablet	T1	
spironolacton-hydrochlorothiaz	T2	
telmisartan	T2	
telmisartan-amlodipine	T2	
telmisartan-hydrochlorothiazid	T2	
terazosin	T1	
TIADYLT ER	T2	
ticagrelor	T2	
timolol maleate oral	T2	
torsemide oral	T2	
trandolapril	T1	
triamterene-hydrochlorothiazid	T1	
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 400 MCG, 600 MCG, 800 MCG	Т5	PA; QL (62 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
UPTRAVI ORAL TABLET 200 MCG	T5	PA; QL (224 EA per 28 days)
UPTRAVI ORAL TABLETS,DOSE PACK	T5	PA; QL (400 EA per 365 days)
valsartan oral tablet 160 mg, 40 mg, 80 mg	T1	QL (62 EA per 31 days)
valsartan oral tablet 320 mg	T1	QL (31 EA per 31 days)
valsartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
verapamil oral capsule, 24 hr er pellet ct	T4	
verapamil oral capsule,ext rel. pellets 24 hr 120 mg, 180 mg, 240 mg	T2	
verapamil oral capsule,ext rel. pellets 24 hr 360 mg	T4	
verapamil oral tablet	T2	
verapamil oral tablet extended release	T2	
VERQUVO	T4	PA; QL (31 EA per 31 days)
VYNDAQEL	T5	PA; QL (124 EA per 31 days)
warfarin	T1	
XARELTO DVT-PE TREAT 30D START	Т3	QL (51 EA per 30 days)
XARELTO ORAL SUSPENSION FOR RECONSTITUTION	Т3	QL (930 ML per 31 days)
XARELTO ORAL TABLET 10 MG, 20 MG	Т3	QL (31 EA per 31 days)
XARELTO ORAL TABLET 15 MG	Т3	QL (52 EA per 31 days)
XARELTO ORAL TABLET 2.5 MG	Т3	QL (62 EA per 31 days)
Dermatologicals/Topical Therapy		
ACCUTANE ORAL CAPSULE 10 MG, 20 MG, 40 MG	T4	
acitretin	T4	PA
acyclovir topical ointment	T4	QL (30 GM per 30 days)
ADBRY	T5	PA; QL (4 ML per 28 days)
ALA-CORT TOPICAL CREAM 1 %	T2	
alclometasone	T2	
ammonium lactate	T2	
AMNESTEEM	T4	
ANZUPGO	T5	PA; QL (60 GM per 30 days)
betamethasone dipropionate	T2	
betamethasone valerate topical cream	T2	
betamethasone valerate topical lotion	T2	
betamethasone valerate topical ointment	T2	
betamethasone, augmented	T2	

Drug Name	Drug Tier	Requirements/Limits
calcipotriene scalp	Т3	QL (60 ML per 28 days)
calcipotriene topical cream	T4	QL (60 GM per 28 days)
calcipotriene topical ointment	Т3	QL (60 GM per 28 days)
CIBINQO	T5	PA; QL (31 EA per 31 days)
ciclopirox topical cream	T2	QL (90 GM per 28 days)
ciclopirox topical gel	T2	QL (45 GM per 28 days)
ciclopirox topical shampoo	T2	QL (120 ML per 28 days)
ciclopirox topical solution	T2	
ciclopirox topical suspension	T2	QL (60 ML per 28 days)
CLARAVIS	T4	
clindamycin phosphate topical gel	T2	QL (60 GM per 28 days)
clindamycin phosphate topical lotion	T2	QL (60 ML per 28 days)
clindamycin phosphate topical solution	T2	QL (60 ML per 28 days)
clobetasol topical cream 0.05 %	T4	QL (60 GM per 28 days)
clobetasol topical ointment	T4	QL (60 GM per 28 days)
clotrimazole topical cream	T2	QL (45 GM per 28 days)
clotrimazole topical solution	T2	QL (30 ML per 28 days)
clotrimazole-betamethasone topical cream	T2	QL (45 GM per 28 days)
clotrimazole-betamethasone topical lotion	Т3	QL (60 ML per 28 days)
COSENTYX (2 SYRINGES)	T5	PA; QL (2 ML per 28 days)
COSENTYX PEN (2 PENS)	T5	PA; QL (2 ML per 28 days)
COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
COSENTYX UNOREADY PEN	T5	PA; QL (2 ML per 28 days)
desoximetasone topical cream	T4	QL (100 GM per 28 days)
desoximetasone topical gel	T4	QL (60 GM per 28 days)
diclofenac sodium topical gel 3 %	T4	PA; QL (100 GM per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
econazole nitrate topical cream	T2	QL (85 GM per 28 days)
ERY PADS	T2	

Drug Name	Drug Tier	Requirements/Limits
erythromycin with ethanol topical solution	T2	QL (60 ML per 28 days)
FILSUVEZ	T5	PA
fluocinolone and shower cap	T2	QL (118.28 ML per 28 days)
fluocinolone topical cream 0.01 %	T2	QL (60 GM per 28 days)
fluocinolone topical cream 0.025 %	T2	QL (120 GM per 28 days)
fluocinolone topical ointment	T2	QL (120 GM per 28 days)
fluocinolone topical solution	T2	QL (90 ML per 28 days)
fluocinonide topical cream 0.05 %	T2	QL (60 GM per 28 days)
fluocinonide topical gel	Т3	QL (60 GM per 28 days)
fluocinonide topical ointment	T2	QL (60 GM per 28 days)
fluocinonide topical solution	T2	QL (60 ML per 28 days)
fluocinonide-emollient	T4	QL (60 GM per 28 days)
fluorouracil topical cream 5 %	T2	
fluorouracil topical solution	T2	
fluticasone propionate topical cream	T2	
gentamicin topical	T2	QL (60 GM per 28 days)
halobetasol propionate topical cream	T2	QL (50 GM per 28 days)
halobetasol propionate topical ointment	T2	QL (50 GM per 28 days)
hydrocortisone topical cream 1 %	T2	
hydrocortisone topical lotion 2.5 %	T2	QL (118 ML per 28 days)
hydrocortisone topical ointment 1 %, 2.5 %	T2	
imiquimod topical cream in packet 5 %	T2	
ketoconazole topical cream	T2	QL (60 GM per 28 days)
ketoconazole topical shampoo	T2	QL (120 ML per 28 days)
lidocaine hcl mucous membrane solution 4 % (40 mg/ml)	T2	PA; QL (50 ML per 28 days)
lidocaine topical adhesive patch,medicated 5 %	T2	PA; QL (93 EA per 31 days)
lidocaine topical ointment	T4	PA; QL (50 GM per 28 days)
LIDOCAINE VISCOUS	T2	
lidocaine-prilocaine topical cream	T2	PA; QL (30 GM per 28 days)
malathion	T2	
metronidazole topical cream	T2	
metronidazole topical gel	T2	
metronidazole topical lotion	T2	
mometasone topical	T2	
mupirocin	T2	

Drug Name	Drug Tier	Requirements/Limits
NYAMYC	T2	QL (60 GM per 28 days)
nystatin topical cream	T2	QL (30 GM per 28 days)
nystatin topical ointment	T2	QL (30 GM per 28 days)
nystatin topical powder	T2	QL (60 GM per 28 days)
nystatin-triamcinolone	T2	QL (60 GM per 28 days)
NYSTOP	T2	QL (60 GM per 28 days)
PANRETIN	T5	PA-NS
penciclovir	T4	QL (5 GM per 28 days)
permethrin	T2	
pimecrolimus	T4	QL (100 GM per 28 days)
podofilox topical solution	T2	
selenium sulfide topical lotion	T2	
silver sulfadiazine	T2	
SKYRIZI SUBCUTANEOUS PEN INJECTOR	T5	PA; QL (1 ML per 84 days)
SKYRIZI SUBCUTANEOUS SYRINGE	T5	PA; QL (1 ML per 84 days)
SSD	T4	
STELARA SUBCUTANEOUS SOLUTION	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	Т3	PA; QL (0.5 ML per 84 days)
STEQEYMA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
sulfacetamide sodium (acne)	T2	
SULFAMYLON TOPICAL CREAM	Т3	
tacrolimus topical	T2	QL (100 GM per 28 days)
TALTZ AUTOINJECTOR	T5	PA; QL (1 ML per 28 days)
TALTZ SYRINGE SUBCUTANEOUS SYRINGE 20 MG/0.25 ML	T5	PA; QL (0.25 ML per 28 days)
TALTZ SYRINGE SUBCUTANEOUS SYRINGE 40 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
TALTZ SYRINGE SUBCUTANEOUS SYRINGE 80 MG/ML	T5	PA; QL (1 ML per 28 days)
tazarotene topical cream	T4	PA; QL (60 GM per 28 days)
tretinoin topical cream	T2	PA; QL (45 GM per 28 days)

Drug Name	Drug Tier	Requirements/Limits
tretinoin topical gel 0.01 %, 0.025 %	Т3	PA; QL (45 GM per 28 days)
triamcinolone acetonide topical cream	T2	
triamcinolone acetonide topical lotion	T2	
triamcinolone acetonide topical ointment 0.025 %, 0.1 %, 0.5 %	T2	
TRIDERM TOPICAL CREAM 0.5 %	T4	
VALCHLOR	T5	PA-NS
YESINTEK SUBCUTANEOUS SOLUTION	Т3	PA; QL (0.5 ML per 84 days)
YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	Т3	PA; QL (0.5 ML per 84 days)
YESINTEK SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
ZELSUVMI	T5	PA; QL (31 GM per 28 days)
Diagnostics / Miscellaneous Agents		
acamprosate	T4	
anagrelide	T2	
bupropion hcl (smoking deter)	T2	QL (62 EA per 31 days)
carglumic acid	T5	PA
cevimeline	T2	
СНЕМЕТ	T4	
CLINIMIX 4.25%/D5W SULFIT FREE	T4	PA-BvD
d10 %-0.45 % sodium chloride	T2	
d2.5 %-0.45 % sodium chloride	T2	
d5 % and 0.9 % sodium chloride	T2	
d5 %-0.45 % sodium chloride	T2	
deferasirox oral granules in packet	T5	PA
deferasirox oral tablet 180 mg, 360 mg	T5	PA
deferasirox oral tablet 90 mg	Т3	PA
deferasirox oral tablet, dispersible 125 mg	T4	PA
deferasirox oral tablet, dispersible 250 mg, 500 mg	T5	PA
deferiprone	T5	PA
dextrose 10 % in water (d10w)	T2	
dextrose 5 % in water (d5w) intravenous parenteral solution	T2	
disulfiram	Т3	
droxidopa oral capsule 100 mg	T5	PA; QL (465 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
droxidopa oral capsule 200 mg, 300 mg	T5	PA; QL (186 EA per 31 days)
DUVYZAT	T5	PA; QL (420 ML per 35 days)
FABHALTA	T5	PA; QL (62 EA per 31 days)
INCRELEX	T5	PA
JOENJA	T5	PA; QL (60 EA per 30 days)
KIONEX (WITH SORBITOL)	T2	
levocarnitine (with sugar)	T2	PA-BvD
levocarnitine oral tablet	T2	PA-BvD
LITFULO	T5	PA; QL (28 EA per 28 days)
LOKELMA	Т3	PA; QL (93 EA per 31 days)
midodrine	T2	
NICOTROL NS	T4	
nitisinone	T5	PA
PHEBURANE	T5	PA; QL (620 GM per 31 days)
pilocarpine hcl oral	Т3	
PROLASTIN-C INTRAVENOUS SOLUTION	T5	PA
RAVICTI	T5	PA
REZDIFFRA	T5	PA; QL (31 EA per 31 days)
riluzole	Т3	
risedronate oral tablet 30 mg	T4	
sodium chloride 0.9 % intravenous parenteral solution	T2	
sodium chloride irrigation	T2	
sodium phenylbutyrate	T5	PA
sodium polystyrene sulfonate oral powder	T2	
SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 2.5 MG, 5 MG	T5	PA; QL (31 EA per 31 days)
SOHONOS ORAL CAPSULE 10 MG	T5	PA; QL (62 EA per 31 days)
SPS (WITH SORBITOL) ORAL	T2	
trientine oral capsule 250 mg	T5	QL (248 EA per 31 days)
varenicline tartrate oral tablet	T4	QL (60 EA per 30 days)
varenicline tartrate oral tablets,dose pack	T4	QL (106 EA per 365 days)
Ear, Nose / Throat Medications		
acetic acid otic (ear)	T2	
azelastine nasal spray,non-aerosol 137 mcg (0.1 %)	T2	QL (30 ML per 25 days)

Drug Name	Drug Tier	Requirements/Limits
chlorhexidine gluconate mucous membrane	T1	
ciprofloxacin-dexamethasone	Т3	
fluocinolone acetonide oil	T2	
hydrocortisone-acetic acid	T2	
ipratropium bromide nasal spray,non-aerosol 21 mcg (0.03 %)	T2	QL (30 ML per 28 days)
ipratropium bromide nasal spray,non-aerosol 42 mcg (0.06 %)	Т2	QL (15 ML per 28 days)
KOURZEQ	T2	
neomycin-polymyxin-hc otic (ear)	T2	
ofloxacin otic (ear)	T2	
olopatadine nasal	Т3	QL (30.5 GM per 30 days)
PERIOGARD	T1	
triamcinolone acetonide dental	T2	
Endocrine/Diabetes		
acarbose	T2	QL (93 EA per 31 days)
ALCOHOL PADS	T2	PA
BAQSIMI	Т3	
cabergoline	T2	
calcitonin (salmon) nasal	T2	PA-BvD
calcitriol oral	T2	PA-BvD
CERDELGA	T5	PA; QL (62 EA per 31 days)
cinacalcet oral tablet 30 mg, 60 mg	T4	PA-BvD; QL (62 EA per 31 days)
cinacalcet oral tablet 90 mg	T4	PA-BvD; QL (124 EA per 31 days)
danazol	T4	
desmopressin nasal spray,non-aerosol 10 mcg/spray (0.1 ml)	T2	
desmopressin oral	T2	
dexamethasone oral solution	T2	
dexamethasone oral tablet	T1	
diazoxide	T5	
doxercalciferol oral	T4	PA-BvD
FARXIGA	Т3	QL (31 EA per 31 days)
fludrocortisone	T1	
glimepiride oral tablet 1 mg, 2 mg, 4 mg	T1	
glipizide oral tablet 10 mg, 5 mg	T1	

Drug Name	Drug Tier	Requirements/Limits
glipizide oral tablet extended release 24hr	T1	
glipizide-metformin	T1	
GLUCAGON EMERGENCY KIT (HUMAN)	Т3	
GLYXAMBI	Т3	QL (31 EA per 31 days)
GVOKE	Т3	
GVOKE HYPOPEN 2-PACK	Т3	
GVOKE PFS 1-PACK SYRINGE SUBCUTANEOUS SYRINGE 1 MG/0.2 ML	Т3	
HUMALOG JUNIOR KWIKPEN U-100	Т3	
HUMALOG KWIKPEN INSULIN	Т3	
HUMALOG MIX 50-50 KWIKPEN	Т3	
HUMALOG MIX 75-25 KWIKPEN	Т3	
HUMALOG MIX 75-25(U-100)INSULN	Т3	
HUMALOG TEMPO PEN(U-100)INSULN	Т3	
HUMALOG U-100 INSULIN	Т3	
HUMULIN 70/30 U-100 INSULIN	Т3	
HUMULIN 70/30 U-100 KWIKPEN	Т3	
HUMULIN N NPH INSULIN KWIKPEN	Т3	
HUMULIN N NPH U-100 INSULIN	Т3	
HUMULIN R REGULAR U-100 INSULN	Т3	
HUMULIN R U-500 (CONC) INSULIN	Т3	
HUMULIN R U-500 (CONC) KWIKPEN	Т3	
hydrocortisone oral	T2	
insulin lispro	Т3	
insulin lispro protamin-lispro	Т3	
JANUMET	Т3	QL (62 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 100-1,000 MG, 50-500 MG	Т3	QL (31 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 50-1,000 MG	Т3	QL (62 EA per 31 days)
JANUVIA ORAL TABLET 100 MG, 50 MG	Т3	QL (31 EA per 31 days)
JANUVIA ORAL TABLET 25 MG	Т3	QL (93 EA per 31 days)
JARDIANCE ORAL TABLET 10 MG	Т3	QL (62 EA per 31 days)
JARDIANCE ORAL TABLET 25 MG	Т3	QL (31 EA per 31 days)
JAVYGTOR	T5	PA

Drug Name	Drug Tier	Requirements/Limits
JENTADUETO ORAL TABLET 2.5-1,000 MG, 2.5-500 MG	Т3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 5-1,000 MG	Т3	QL (31 EA per 31 days)
LANTUS SOLOSTAR U-100 INSULIN	Т3	
LANTUS U-100 INSULIN	Т3	
levothyroxine oral tablet	T1	
LEVOXYL ORAL TABLET 100 MCG, 112 MCG, 125 MCG, 137 MCG, 150 MCG, 175 MCG, 200 MCG, 25 MCG, 50 MCG, 75 MCG, 88 MCG	Т3	
liothyronine oral	T2	
metformin oral tablet 1,000 mg, 500 mg, 850 mg	T1	
metformin oral tablet extended release 24 hr	T1	
metformin oral tablet extended release 24hr	NF	
metformin oral tablet,er gast.retention 24 hr	NF	
methimazole oral tablet 10 mg, 5 mg	T1	
methylprednisolone	T2	
mifepristone oral tablet 300 mg	T5	PA; QL (124 EA per 31 days)
miglustat	T5	PA; QL (93 EA per 31 days)
MOUNJARO	Т3	PA; QL (2 ML per 28 days)
nateglinide	T2	QL (93 EA per 31 days)
OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)	Т3	PA; QL (3 ML per 28 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (15 ML per 30 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 2.5 MG/0.5 ML	T5	PA; QL (4 ML per 30 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (90 ML per 30 days)
paricalcitol oral	T4	PA-BvD
pioglitazone	T1	QL (31 EA per 31 days)
pioglitazone-metformin	T2	QL (93 EA per 31 days)
prednisolone oral solution	T2	

Drug Name	Drug Tier	Requirements/Limits
prednisolone sodium phosphate oral solution 25 mg/5 ml (5 mg/ml), 5 mg base/5 ml (6.7 mg/5 ml)	T2	
prednisone oral solution	Т3	
prednisone oral tablet	T1	
prednisone oral tablets,dose pack	T2	
propylthiouracil	T2	
RECORLEV	T5	PA; QL (248 EA per 31 days)
repaglinide oral tablet 0.5 mg, 1 mg	T2	QL (124 EA per 31 days)
repaglinide oral tablet 2 mg	T2	QL (248 EA per 31 days)
RYBELSUS	T3	PA; QL (31 EA per 31 days)
sapropterin	T5	PA
SOLIQUA 100/33	Т3	QL (18 ML per 30 days)
SOMAVERT	T5	PA
SYMLINPEN 120	T5	QL (10.8 ML per 28 days)
SYMLINPEN 60	T5	QL (6 ML per 28 days)
SYNAREL	T5	PA
SYNJARDY	Т3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 12.5-1,000 MG, 5-1,000 MG	Т3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 25-1,000 MG	Т3	QL (31 EA per 31 days)
SYNTHROID	Т3	
testosterone cypionate	T2	PA
testosterone enanthate	Т3	PA
testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)	Т3	PA
testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram)	T2	PA
testosterone transdermal gel in packet 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)	Т3	PA
testosterone transdermal solution in metered pump w/app	T2	PA
tolvaptan	T5	PA
TOUJEO MAX U-300 SOLOSTAR	Т3	
TOUJEO SOLOSTAR U-300 INSULIN	Т3	
TRADJENTA	T3	QL (31 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-5-1,000 MG, 25-5-1,000 MG	Т3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 12.5-2.5-1,000 MG, 5-2.5- 1,000 MG	Т3	QL (62 EA per 31 days)
TRULICITY	Т3	PA; QL (2 ML per 28 days)
UNITHROID	Т3	
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 10-500 MG, 5- 500 MG	Т3	QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG, 5-1,000 MG	Т3	QL (62 EA per 31 days)
XULTOPHY 100/3.6	Т3	QL (15 ML per 30 days)
YARGESA	T5	PA; QL (93 EA per 31 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML	T5	PA; QL (1.12 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 294 MCG/0.98 ML	T5	PA; QL (1.96 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 420 MCG/1.4 ML	T5	PA; QL (2.8 ML per 28 days)
ZEGALOGUE AUTOINJECTOR	Т3	
ZEGALOGUE SYRINGE	Т3	
Gastroenterology		
alosetron oral tablet 0.5 mg	T4	PA; QL (93 EA per 31 days)
alosetron oral tablet 1 mg	T5	PA; QL (62 EA per 31 days)
aprepitant oral capsule 125 mg	T5	PA-BvD
aprepitant oral capsule 40 mg, 80 mg	T4	PA-BvD
aprepitant oral capsule,dose pack	T4	PA-BvD
balsalazide	T2	
betaine	T5	
budesonide oral capsule,delayed,extend.release	T4	
budesonide oral tablet,delayed and ext.release	T5	
CIMZIA POWDER FOR RECONST	T5	PA; QL (2 EA per 28 days)
CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)	T5	PA; QL (2 EA per 28 days)
COMPRO	T4	
CONSTULOSE	T2	

Drug Name	Drug Tier	Requirements/Limits
CREON	T3	
cromolyn oral	T4	
dicyclomine oral capsule	T2	
dicyclomine oral solution	T2	
dicyclomine oral tablet 20 mg	T2	
diphenoxylate-atropine oral liquid	T4	
diphenoxylate-atropine oral tablet	T2	
dronabinol	T4	PA-BvD
ENULOSE	T2	
esomeprazole magnesium oral capsule,delayed release(dr/ec)	T2	QL (31 EA per 31 days)
famotidine oral suspension for reconstitution	T2	
famotidine oral tablet 20 mg, 40 mg	T1	
GATTEX 30-VIAL	T5	PA
GAVILYTE-C	T1	
GAVILYTE-G	T1	
GAVILYTE-N	T1	
GENERLAC	T2	
glycopyrrolate oral tablet 1 mg, 2 mg	T2	
granisetron hcl oral	T2	PA-BvD
hydrocortisone rectal	T4	
hydrocortisone topical cream with perineal applicator 2.5 %	T2	
IBSRELA	T5	PA; QL (62 EA per 31 days)
lactulose oral solution	T2	
lansoprazole oral capsule,delayed release(dr/ec) 15 mg	T2	QL (31 EA per 31 days)
lansoprazole oral capsule,delayed release(dr/ec) 30 mg	T2	QL (62 EA per 31 days)
LINZESS	Т3	QL (31 EA per 31 days)
loperamide oral capsule	T2	
lubiprostone	Т3	QL (62 EA per 31 days)
meclizine oral tablet 12.5 mg, 25 mg	T2	
mesalamine oral capsule (with del rel tablets)	T4	QL (186 EA per 31 days)
mesalamine oral capsule,extended release 24hr	T4	QL (124 EA per 31 days)
mesalamine oral tablet,delayed release (dr/ec) 1.2 gram	T4	QL (124 EA per 31 days)

Drug Name	Drug Tier	Requirements/Limits
mesalamine rectal enema	T4	QL (1860 ML per 31 days)
metoclopramide hcl oral solution	T2	
metoclopramide hcl oral tablet	T1	
misoprostol	T2	
MOVANTIK	Т3	QL (31 EA per 31 days)
nitroglycerin rectal	T4	
omeprazole oral capsule,delayed release(dr/ec)	T1	
ondansetron hcl oral solution	T2	PA-BvD
ondansetron hcl oral tablet 4 mg, 8 mg	T2	PA-BvD
ondansetron oral tablet, disintegrating 4 mg, 8 mg	T2	PA-BvD
pantoprazole oral tablet,delayed release (dr/ec)	T1	
peg 3350-electrolytes	T1	
peg3350-sod sul-nacl-kcl-asb-c	T4	
peg-electrolyte soln	T1	
prochlorperazine	T4	
prochlorperazine maleate	T1	
PROCTO-MED HC	T2	
PROCTOSOL HC TOPICAL	T2	
PROCTOZONE-HC	T2	
rabeprazole oral tablet,delayed release (dr/ec)	T2	QL (62 EA per 31 days)
scopolamine base	Т3	QL (10 EA per 30 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML)	T5	PA; QL (1.2 ML per 56 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 360 MG/2.4 ML (150 MG/ML)	T5	PA; QL (2.4 ML per 56 days)
sodium,potassium,mag sulfates	T4	
sucralfate oral suspension	T4	
sucralfate oral tablet	T2	
sulfasalazine	T2	
ursodiol oral capsule 300 mg	T2	
ursodiol oral tablet	Т3	
VIBERZI	T5	PA; QL (62 EA per 31 days)
VOWST	T5	PA; QL (12 EA per 14 days)
Immunology, Vaccines / Biotechnology		
ABRYSVO (PF)	Т3	QL (1 EA per 365 days)
ACTHIB (PF)	Т3	

Drug Name	Drug Tier	Requirements/Limits
ACTIMMUNE	T5	PA
ADACEL(TDAP ADOLESN/ADULT)(PF)	Т3	
AREXVY (PF)	Т3	QL (1 EA per 365 days)
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	T5	PA; QL (1 EA per 28 days)
AVONEX INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (1 EA per 28 days)
bcg vaccine, live (pf)	Т3	
BESREMI	T5	PA-NS; QL (2 ML per 28 days)
BETASERON SUBCUTANEOUS KIT	T5	PA; QL (14 EA per 28 days)
BEXSERO	Т3	
BIVIGAM	T5	PA
BOOSTRIX TDAP	Т3	
DAPTACEL (DTAP PEDIATRIC) (PF)	Т3	
ENGERIX-B (PF)	Т3	PA-BvD
ENGERIX-B PEDIATRIC (PF)	Т3	PA-BvD
FULPHILA	T5	
GAMMAGARD LIQUID	T5	PA
GAMMAGARD S-D (IGA < 1 MCG/ML)	T5	PA
GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GAMMAPLEX	T5	PA
GAMMAPLEX (WITH SORBITOL)	T5	PA
GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GARDASIL 9 (PF)	Т3	
HAVRIX (PF)	Т3	
HEPLISAV-B (PF)	Т3	PA-BvD
HIBERIX (PF)	Т3	
IMOVAX RABIES VACCINE (PF)	Т3	PA-BvD
INFANRIX (DTAP) (PF)	Т3	
IPOL	Т3	
IXIARO (PF)	Т3	
JYNNEOS (PF)	Т3	PA-BvD
KINRIX (PF)	Т3	
LEUKINE INJECTION RECON SOLN	T5	PA
MENQUADFI (PF)	Т3	

Drug Name	Drug Tier	Requirements/Limits
MENVEO A-C-Y-W-135-DIP (PF) INTRAMUSCULAR KIT	Т3	
M-M-R II (PF)	Т3	
MRESVIA (PF)	T3	QL (1 ML per 365 days)
NEULASTA	T5	
NIVESTYM	T5	
NORDITROPIN FLEXPRO	T5	PA
OCTAGAM	T5	PA
PANZYGA	T5	PA
PEDIARIX (PF)	Т3	
PEDVAX HIB (PF)	Т3	
PEGASYS	T5	PA
PENBRAYA (PF)	Т3	
PENMENVY MEN A-B-C-W-Y (PF)	Т3	
PENTACEL (PF)	Т3	
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	Т5	PA; QL (1 ML per 28 days)
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PRIORIX (PF)	T3	
PRIVIGEN	T5	PA
PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML	Т3	PA-BvD
PROCRIT INJECTION SOLUTION 20,000 UNIT/ML, 40,000 UNIT/ML	T5	PA-BvD
PROQUAD (PF)	Т3	
QUADRACEL (PF)	Т3	
RABAVERT (PF)	Т3	PA-BvD
RECOMBIVAX HB (PF)	Т3	PA-BvD
RETACRIT	Т3	PA-BvD
ROTARIX ORAL SUSPENSION	Т3	
ROTATEQ VACCINE	Т3	
SHINGRIX (PF)	Т3	QL (2 EA per 999 days)
TENIVAC (PF)	Т3	
TICOVAC	Т3	
TRUMENBA	Т3	

Drug Name	Drug Tier	Requirements/Limits
TWINRIX (PF)	Т3	
TYPHIM VI	Т3	
VAQTA (PF)	Т3	
VARIVAX (PF)	Т3	
VAXCHORA VACCINE	Т3	QL (200 ML per 365 days)
VIMKUNYA	T3	
VIVOTIF	Т3	
XOLREMDI	T5	PA; QL (124 EA per 31 days)
YF-VAX (PF)	Т3	
ZARXIO	T5	
ZIEXTENZO	T5	
Miscellaneous Supplies		
ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"	Т3	
GAUZE PAD TOPICAL BANDAGE 2 X 2 "	Т3	PA
insulin syringe-needle u-100 syringe 0.3 ml 29 gauge, 1 ml 29 gauge x 1/2", 1/2 ml 28 gauge	Т3	
pen needle, diabetic needle 29 gauge x 1/2"	Т3	
Musculoskeletal / Rheumatology		
ACTEMRA ACTPEN	T5	PA; QL (3.6 ML per 28 days)
ACTEMRA SUBCUTANEOUS	T5	PA; QL (3.6 ML per 28 days)
alendronate oral tablet 10 mg, 35 mg, 70 mg	T1	
allopurinol oral tablet 100 mg, 300 mg	T1	
BENLYSTA SUBCUTANEOUS	T5	PA; QL (4 ML per 28 days)
colchicine oral tablet	T2	QL (62 EA per 31 days)
CYLTEZO(CF)	T5	PA; QL (2 EA per 28 days)
CYLTEZO(CF) PEN	T5	PA; QL (2 EA per 28 days)
CYLTEZO(CF) PEN CROHN'S-UC-HS SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML	Т5	PA; QL (12 EA per 365 days)
CYLTEZO(CF) PEN PSORIASIS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML	T5	PA; QL (8 EA per 365 days)
ENBREL MINI	T5	PA; QL (8 ML per 28 days)
ENBREL SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5)	T5	PA; QL (4 ML per 28 days)

Drug Name	Drug Tier	Requirements/Limits
ENBREL SUBCUTANEOUS SYRINGE 50 MG/ML (1 ML)	T5	PA; QL (8 ML per 28 days)
ENBREL SURECLICK	T5	PA; QL (8 ML per 28 days)
EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML (105MG/1.17MLX2)	T5	PA; QL (2.34 ML per 28 days)
febuxostat	T2	PA
HADLIMA	T5	PA; QL (1.6 ML per 28 days)
HADLIMA PUSHTOUCH	T5	PA; QL (1.6 ML per 28 days)
HADLIMA(CF)	T5	PA; QL (0.8 ML per 28 days)
HADLIMA(CF) PUSHTOUCH	T5	PA; QL (0.8 ML per 28 days)
HUMIRA PEN	T5	PA; QL (2 EA per 28 days)
HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF)	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF) PEN	T5	PA; QL (2 EA per 28 days)
HUMIRA(CF) PEN CROHNS-UC-HS	T5	PA; QL (6 EA per 365 days)
HUMIRA(CF) PEN PSOR-UV-ADOL HS	T5	PA; QL (6 EA per 365 days)
ibandronate oral	T2	
KEVZARA	T5	PA; QL (2.28 ML per 28 days)
KINERET	T5	PA; QL (18.76 ML per 28 days)
leflunomide	T2	
OLUMIANT	T5	PA; QL (31 EA per 31 days)
ORENCIA CLICKJECT	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 50 MG/0.4 ML	T5	PA; QL (1.6 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 87.5 MG/0.7 ML	T5	PA; QL (2.8 ML per 28 days)
OTEZLA	T5	PA; QL (62 EA per 31 days)
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51)	T5	PA; QL (110 EA per 365 days)
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)	T5	PA; QL (55 EA per 28 days)
penicillamine oral tablet	T5	
probenecid	T2	
probenecid-colchicine	T2	
PROLIA	Т3	PA; QL (1 ML per 180 days)

Drug Name	Drug Tier	Requirements/Limits
raloxifene	T2	
RINVOQ LQ	T5	PA; QL (372 ML per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG	T5	PA; QL (31 EA per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 45 MG	T5	PA; QL (168 EA per 365 days)
risedronate oral tablet 150 mg, 35 mg, 35 mg (12 pack), 35 mg (4 pack), 5 mg	T4	
risedronate oral tablet,delayed release (dr/ec)	T4	
SAVELLA	T3	PA
SIMLANDI(CF)	T5	PA; QL (2 EA per 28 days)
SIMLANDI(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML	T5	PA; QL (1 ML per 28 days)
SIMPONI SUBCUTANEOUS PEN INJECTOR 50 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML	T5	PA; QL (1 ML per 28 days)
SIMPONI SUBCUTANEOUS SYRINGE 50 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)	T5	PA; QL (2.48 ML per 28 days)
TYMLOS	T5	PA; QL (1.56 ML per 30 days)
XELJANZ ORAL SOLUTION	T5	PA; QL (310 ML per 31 days)
XELJANZ ORAL TABLET	T5	PA; QL (62 EA per 31 days)
XELJANZ XR	T5	PA; QL (31 EA per 31 days)
YUFLYMA(CF)	T5	PA; QL (2 EA per 28 days)
YUFLYMA(CF) AI CROHN'S-UC-HS	T5	PA; QL (6 EA per 365 days)
YUFLYMA(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
Obstetrics / Gynecology		
ABIGALE	T2	
ALTAVERA (28)	T2	
ALYACEN 1/35 (28)	T2	
APRI	T2	
ARANELLE (28)	T2	
AVIANE	T2	
AZURETTE (28)	T2	
CAMILA	T2	

Drug Name	Drug Tier	Requirements/Limits
clindamycin phosphate vaginal	T2	
CRYSELLE (28)	T2	
CYRED EQ	T2	
DEPO-SUBQ PROVERA 104	Т3	
DOTTI	T2	
drospirenone-e.estradiol-lm.fa oral tablet 3-0.02-0.451 mg (24) (4)	T2	
drospirenone-ethinyl estradiol	T2	
ELURYNG	T4	
ENPRESSE	T2	
ENSKYCE	T2	
ERRIN	T2	
ESTARYLLA	T2	
estradiol oral	T2	
estradiol transdermal patch semiweekly	T2	
estradiol transdermal patch weekly	T2	
estradiol vaginal	T2	
estradiol-norethindrone acet	T2	
ethynodiol diac-eth estradiol oral tablet 1-35 mg- mcg	T2	
etonogestrel-ethinyl estradiol	T3	
FEIRZA	T2	
GALLIFREY	T2	
HALOETTE	T4	
HEATHER	T2	
ICLEVIA	T2	
IMVEXXY MAINTENANCE PACK	Т3	
IMVEXXY STARTER PACK	T3	
INCASSIA	T2	
INTROVALE	T2	
ISIBLOOM	T2	
JASMIEL (28)	T2	
JINTELI	T4	
JULEBER	T2	
KARIVA (28)	T2	
KELNOR 1/35 (28)	T2	

Drug Name	Drug Tier	Requirements/Limits
KURVELO (28)	T2	
l norgest/e.estradiol-e.estrad oral tablets,dose pack,3 month 0.1 mg-20 mcg (84)/10 mcg (7), 0.15 mg-30 mcg (84)/10 mcg (7)	T2	
LESSINA	T2	
LEVONEST (28)	T2	
levonorgestrel-ethinyl estrad oral tablet 0.1-20 mg-mcg, 0.15-0.03 mg	T2	
levonorgestrel-ethinyl estrad oral tablets,dose pack,3 month	T2	
levonorg-eth estrad triphasic	T2	
LEVORA-28	T2	
LILETTA	Т3	
LORYNA (28)	T2	
LOW-OGESTREL (28)	T2	
LUTERA (28)	T2	
LYLEQ	T2	
LYLLANA	T2	
LYZA	T2	
MARLISSA (28)	T2	
medroxyprogesterone	T2	
metronidazole vaginal gel 0.75 % (37.5mg/5 gram)	Т3	
MICONAZOLE-3 VAGINAL SUPPOSITORY	T2	
MICROGESTIN 1.5/30 (21)	T2	
MICROGESTIN 1/20 (21)	T2	
MICROGESTIN FE 1.5/30 (28)	T2	
MICROGESTIN FE 1/20 (28)	T2	
MILI	T2	
NEXPLANON	Т3	
norethindrone (contraceptive)	T2	
norethindrone acetate	T2	
norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg	T4	
norethindrone ac-eth estradiol oral tablet 1-20 mg-mcg	T2	
norgestimate-ethinyl estradiol	T2	

Drug Name	Drug Tier	Requirements/Limits
NORTREL 0.5/35 (28)	T2	
NORTREL 1/35 (21)	T2	
NORTREL 1/35 (28)	T2	
NORTREL 7/7/7 (28)	T2	
PIMTREA (28)	T2	
PORTIA 28	T2	
PREMARIN ORAL	Т3	
PREMARIN VAGINAL	Т3	
PREMPRO	Т3	
progesterone micronized oral	T2	
RECLIPSEN (28)	T2	
SETLAKIN	T2	
SPRINTEC (28)	T2	
SRONYX	T2	
SYEDA	T2	
terconazole	T2	
TILIA FE	T2	
tranexamic acid oral	Т3	
TRI-ESTARYLLA	T2	
TRI-LEGEST FE	T2	
TRI-LO-ESTARYLLA	T2	
TRI-LO-SPRINTEC	T2	
TRI-SPRINTEC (28)	T2	
TURQOZ (28)	T2	
VALTYA ORAL TABLET 1-50 MG-MCG	T2	
VELIVET TRIPHASIC REGIMEN (28)	T2	
VESTURA (28)	T2	
VIENVA	T2	
XELRIA FE	T2	
YUVAFEM	Т3	
ZAFEMY	T3	
ZOVIA 1-35 (28)	T2	
Ophthalmology		
acetazolamide	T2	
ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.1 %	Т3	

Drug Name	Drug Tier	Requirements/Limits
apraclonidine	Т3	
atropine ophthalmic (eye) drops 1 %	T2	
azelastine ophthalmic (eye)	T2	
bacitracin ophthalmic (eye)	T2	
bacitracin-polymyxin b	T2	
BESIVANCE	Т3	
betaxolol ophthalmic (eye)	T2	
bimatoprost ophthalmic (eye)	T2	
brimonidine ophthalmic (eye) drops 0.1 %	Т3	
brimonidine ophthalmic (eye) drops 0.15 %, 0.2 %	T2	
brimonidine-timolol	Т3	
brinzolamide	T4	
bromfenac ophthalmic (eye) drops 0.07 %, 0.075 %	Т3	
bromfenac ophthalmic (eye) drops 0.09 %	T2	
carteolol	T2	
ciprofloxacin hcl ophthalmic (eye)	T2	
COMBIGAN	Т3	
cromolyn ophthalmic (eye)	T2	
cyclosporine ophthalmic (eye)	Т3	QL (60 EA per 30 days)
CYSTARAN	T5	PA; QL (60 ML per 28 days)
dexamethasone sodium phosphate ophthalmic (eye)	T2	
diclofenac sodium ophthalmic (eye)	T2	
difluprednate	T4	
dorzolamide	T2	
dorzolamide-timolol	T2	
dorzolamide-timolol (pf) ophthalmic (eye) dropperette	T2	
erythromycin ophthalmic (eye)	T2	
fluorometholone	T2	
flurbiprofen sodium	T2	
gatifloxacin	Т3	
gentamicin ophthalmic (eye) drops	T2	
ketorolac ophthalmic (eye)	T2	
latanoprost	T1	

Drug Name	Drug Tier	Requirements/Limits
levobunolol ophthalmic (eye) drops 0.5 %	T2	
LUMIGAN OPHTHALMIC (EYE) DROPS 0.01 %	Т3	QL (5 ML per 31 days)
methazolamide	T4	
moxifloxacin ophthalmic (eye) drops	T2	
NATACYN	T4	
neomycin-bacitracin-poly-hc	T2	
neomycin-bacitracin-polymyxin	T2	
neomycin-polymyxin b-dexameth	T2	
neomycin-polymyxin-gramicidin	T2	
neomycin-polymyxin-hc ophthalmic (eye)	Т3	
NEO-POLYCIN	T2	
NEO-POLYCIN HC	T2	
ofloxacin ophthalmic (eye)	T2	
pilocarpine hcl ophthalmic (eye) drops 1 %, 2 %, 4 %	T2	
POLYCIN	T2	
polymyxin b sulf-trimethoprim	T2	
prednisolone acetate	T2	
prednisolone sodium phosphate ophthalmic (eye)	T2	
PROLENSA	Т3	
RESTASIS	Т3	QL (60 EA per 30 days)
RESTASIS MULTIDOSE	Т3	QL (5.5 ML per 27 days)
RHOPRESSA	T4	ST
ROCKLATAN	T4	ST
SIMBRINZA	T4	
sulfacetamide sodium ophthalmic (eye)	T2	
sulfacetamide-prednisolone	T2	
timolol maleate ophthalmic (eye) drops	T1	
timolol maleate ophthalmic (eye) gel forming solution	Т3	
tobramycin ophthalmic (eye)	T2	
tobramycin-dexamethasone	Т3	
travoprost	Т3	
trifluridine	Т3	
XDEMVY	T5	PA; QL (10 ML per 42 days)

Drug Name	Drug Tier	Requirements/Limits
XIIDRA	Т3	QL (60 EA per 30 days)
ZIRGAN	T4	ST
Respiratory And Allergy		
acetylcysteine	T2	PA-BvD
ADEMPAS	T5	PA; QL (93 EA per 31 days)
albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation	T2	QL (17 GM per 30 days)
albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020503)	T2	QL (13.4 GM per 30 days)
albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020983)	NF	
albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml	T2	PA-BvD
albuterol sulfate oral syrup	T2	
albuterol sulfate oral tablet	T4	
ALYQ	T5	PA; QL (62 EA per 31 days)
ambrisentan	T5	PA; QL (31 EA per 31 days)
ANORO ELLIPTA	Т3	QL (60 EA per 30 days)
arformoterol	Т3	PA-BvD
ASMANEX HFA	Т3	QL (13 GM per 30 days)
ASMANEX TWISTHALER INHALATION AEROSOL POWDR BREATH ACTIVATED 110 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (120), 220 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (60)	Т3	QL (1 EA per 30 days)
ATROVENT HFA	Т3	QL (25.8 GM per 30 days)
azelastine-fluticasone	T4	QL (23 GM per 30 days)
BREO ELLIPTA	Т3	QL (60 EA per 30 days)
BREYNA	Т3	QL (10.3 GM per 30 days)
BREZTRI AEROSPHERE	T3	QL (10.7 GM per 30 days)
BRINSUPRI	T5	PA; QL (31 EA per 31 days)
budesonide inhalation	T4	PA-BvD
budesonide-formoterol	T3	QL (10.2 GM per 30 days)
cetirizine oral solution 1 mg/ml	T2	QL (310 ML per 31 days)
CINRYZE	T5	PA; QL (20 EA per 28 days)
COMBIVENT RESPIMAT	T3	QL (4 GM per 30 days)
cromolyn inhalation	T4	PA-BvD

Drug Name	Drug Tier	Requirements/Limits
desloratadine oral tablet	T2	QL (31 EA per 31 days)
DULERA	Т3	QL (13 GM per 30 days)
epinephrine injection auto-injector 0.15 mg/0.3 ml, 0.3 mg/0.3 ml	Т3	
FASENRA PEN	T5	PA; QL (1 ML per 28 days)
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (0.5 ML per 56 days)
FASENRA SUBCUTANEOUS SYRINGE 30 MG/ML	T5	PA; QL (1 ML per 28 days)
flunisolide	T2	QL (50 ML per 25 days)
fluticasone propionate inhalation blister with device 100 mcg/actuation, 50 mcg/actuation	Т4	ST; QL (60 EA per 30 days)
fluticasone propionate inhalation blister with device 250 mcg/actuation	T4	ST; QL (240 EA per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 110 mcg/actuation	Т4	ST; QL (12 GM per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 220 mcg/actuation	T4	ST; QL (24 GM per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 44 mcg/actuation	T4	ST; QL (10.6 GM per 30 days)
fluticasone propionate nasal	T2	QL (16 GM per 30 days)
fluticasone propion-salmeterol inhalation aerosol powdr breath activated	Т3	QL (1 EA per 30 days)
fluticasone propion-salmeterol inhalation blister with device	Т3	QL (60 EA per 30 days)
hydroxyzine hcl oral tablet	T2	PA
icatibant	T5	PA; QL (18 ML per 30 days)
ipratropium bromide inhalation	T2	PA-BvD
ipratropium-albuterol	T2	PA-BvD
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 5.8 MG, 50 MG, 75 MG	T5	PA; QL (56 EA per 28 days)
KALYDECO ORAL GRANULES IN PACKET 25 MG	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL TABLET	T5	PA; QL (62 EA per 31 days)
levalbuterol hcl inhalation solution for nebulization 1.25 mg/3 ml	Т3	PA-BvD
levalbuterol tartrate	Т3	QL (30 GM per 30 days)
levocetirizine oral solution	T4	QL (310 ML per 31 days)
levocetirizine oral tablet	T2	QL (31 EA per 31 days)

T2	Drug Name	Drug Tier	Requirements/Limits
T2	mometasone nasal	T2	QL (34 GM per 30 days)
NUCALA SUBCUTANEOUS AUTO- INJECTOR	montelukast oral tablet	T2	QL (31 EA per 31 days)
NJECTOR 15	montelukast oral tablet,chewable	T2	QL (31 EA per 31 days)
NUCALA SUBCUTANEOUS SYRINGE 100 T5	NUCALA SUBCUTANEOUS AUTO- INJECTOR	T5	PA; QL (3 ML per 28 days)
MG/ML	NUCALA SUBCUTANEOUS RECON SOLN	T5	PA; QL (3 EA per 28 days)
MG/0.4 ML 15 PA; QL (0.4 ML per 28 days) OFEV T5 PA; QL (62 EA per 31 days) OPSUMIT T5 PA; QL (31 EA per 31 days) OPSYNVI T5 PA; QL (31 EA per 31 days) ORKAMBI ORAL GRANULES IN PACKET T5 PA; QL (62 EA per 31 days) ORKAMBI ORAL TABLET T5 PA; QL (124 EA per 31 days) pirfenidone oral capsule T5 PA; QL (279 EA per 31 days) pirfenidone oral tablet T5 PA; QL (93 EA per 31 days) promethazine oral T4 PA PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION T3 QL (21.2 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) SAAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SEREVENT DISKUS T3 QL (4 GM per 30 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM	NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML	T5	PA; QL (3 ML per 28 days)
OPSUMIT T5 PA; QL (31 EA per 31 days) OPSYNVI T5 PA; QL (31 EA per 31 days) ORKAMBI ORAL GRANULES IN PACKET T5 PA; QL (62 EA per 31 days) ORKAMBI ORAL TABLET T5 PA; QL (124 EA per 31 days) pirfenidone oral capsule T5 PA; QL (279 EA per 31 days) pirfenidone oral tablet T5 PA; QL (93 EA per 31 days) promethazine oral T4 PA PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION T3 QL (21.2 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) MCG/ACTUATION T5 PA; QL (18 ML per 30 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SIGHAMINATE T3 QL (4 GM per 30 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 3	NUCALA SUBCUTANEOUS SYRINGE 40 MG/0.4 ML	T5	PA; QL (0.4 ML per 28 days)
OPSYNVI T5 PA; QL (31 EA per 31 days) ORKAMBI ORAL GRANULES IN PACKET T5 PA; QL (62 EA per 31 days) ORKAMBI ORAL TABLET T5 PA; QL (124 EA per 31 days) pirfenidone oral capsule T5 PA; QL (279 EA per 31 days) pirfenidone oral tablet T5 PA; QL (93 EA per 31 days) promethazine oral T4 PA PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) MCG/ACTUATION T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SEREVENT DISKUS T3 QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (60 EA per 28 days) SYMDEKO T5 P	OFEV	T5	PA; QL (62 EA per 31 days)
ORKAMBI ORAL GRANULES IN PACKET T5 PA; QL (62 EA per 31 days) ORKAMBI ORAL TABLET T5 PA; QL (124 EA per 31 days) pirfenidone oral capsule T5 PA; QL (279 EA per 31 days) pirfenidone oral tablet T5 PA; QL (93 EA per 31 days) promethazine oral T4 PA PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) WCG/ACTUATION T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SEREVENT DISKUS T3 QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) <th< td=""><td>OPSUMIT</td><td>T5</td><td>PA; QL (31 EA per 31 days)</td></th<>	OPSUMIT	T5	PA; QL (31 EA per 31 days)
ORKAMBI ORAL TABLET ORKAMBI ORAL TABLET Description of the proof of	OPSYNVI	T5	PA; QL (31 EA per 31 days)
pirfenidone oral capsule pirfenidone oral tablet promethazine oral T5 PA; QL (279 EA per 31 days) promethazine oral T4 PA PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION T6 T1 QL (31 EA per 30 days) MCG/ACTUATION T6 SEREVENT DISKUS T3 QL (60 EA per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STIOLTO RESPIMAT T5 PA; QL (4 GM per 30 days) STIOLTO RESPIMAT T6 QL (4 GM per 30 days) T6 EA per 31 days) T7 PA; QL (56 EA per 28 days) T6 EA per 31 days) T7 PA; QL (56 EA per 28 days) T6 EA per 31 days) T7 PA; QL (66 EA per 31 days) T8 PA; QL (66 EA per 31 days) T8 PA; QL (66 EA per 31 days) T9 PA; QL (66 EA per 31 days) T7 PA; QL (66 EA per 31 days) T8 PA; QL (66 EA per 31 days)	ORKAMBI ORAL GRANULES IN PACKET	T5	PA; QL (62 EA per 31 days)
pirfenidone oral tablet promethazine oral PA; QL (93 EA per 31 days) POWAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION T3 QL (21.2 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) SIdenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STIOLTO RESPIMAT T5 PA; QL (66 EA per 28 days) T5 PA; QL (66 EA per 28 days) T5 PA; QL (66 EA per 31 days)	ORKAMBI ORAL TABLET	T5	PA; QL (124 EA per 31 days)
PULMOZYME QVAR REDIHALER INHALATION HFA AAROSOL BREATH ACTIVATED 40 MCG/ACTUATION QVAR REDIHALER INHALATION HFA AAROSOL BREATH ACTIVATED 80 MCG/ACTUATION PA QL (21.2 GM per 30 days) MCG/ACTUATION T3 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS Sildenafil (pulm.hypertension) oral tablet T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T5 PA; QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T5 PA; QL (62 EA per 28 days) T4 T5 PA; QL (62 EA per 31 days)	pirfenidone oral capsule	T5	PA; QL (279 EA per 31 days)
PULMOZYME T5 PA QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION T3 QL (21.2 GM per 30 days) QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION T4 QL (31 EA per 31 days) roflumilast T4 QL (18 ML per 30 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STIVERDI RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral T4 T4	pirfenidone oral tablet	T5	PA; QL (93 EA per 31 days)
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION T3 QL (21.2 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS Sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T5 PA; QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	promethazine oral	T4	PA
AEROSOL BREATH ACTIVATED 40 T3 QL (10.6 GM per 30 days) MCG/ACTUATION QVAR REDIHALER INHALATION HFA QL (21.2 GM per 30 days) AEROSOL BREATH ACTIVATED 80 T3 QL (21.2 GM per 30 days) MCG/ACTUATION T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) Sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (30 EA per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral T4	PULMOZYME	T5	PA
AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION T3 QL (21.2 GM per 30 days) roflumilast T4 QL (31 EA per 31 days) SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (4 GM per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral T4	QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION	Т3	QL (10.6 GM per 30 days)
SAJAZIR T5 PA; QL (18 ML per 30 days) SEREVENT DISKUS T3 QL (60 EA per 30 days) Sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T4 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION	Т3	QL (21.2 GM per 30 days)
SEREVENT DISKUS T3 QL (60 EA per 30 days) Sildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	roflumilast	T4	QL (31 EA per 31 days)
resildenafil (pulm.hypertension) oral tablet T3 PA; QL (372 EA per 31 days) SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	SAJAZIR	T5	PA; QL (18 ML per 30 days)
SPIRIVA RESPIMAT T3 QL (4 GM per 30 days) SPIRIVA WITH HANDIHALER T3 QL (30 EA per 30 days) STIOLTO RESPIMAT T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	SEREVENT DISKUS	Т3	QL (60 EA per 30 days)
SPIRIVA WITH HANDIHALERT3QL (30 EA per 30 days)STIOLTO RESPIMATT3QL (4 GM per 30 days)STRIVERDI RESPIMATT4QL (4 GM per 30 days)SYMDEKOT5PA; QL (56 EA per 28 days)tadalafil (pulm. hypertension)T5PA; QL (62 EA per 31 days)terbutaline oralT4	sildenafil (pulm.hypertension) oral tablet	Т3	PA; QL (372 EA per 31 days)
STIOLTO RESPIMAT T3 QL (4 GM per 30 days) STRIVERDI RESPIMAT T4 QL (4 GM per 30 days) SYMDEKO T5 PA; QL (56 EA per 28 days) tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	SPIRIVA RESPIMAT	Т3	QL (4 GM per 30 days)
STRIVERDI RESPIMATT4QL (4 GM per 30 days)SYMDEKOT5PA; QL (56 EA per 28 days)tadalafil (pulm. hypertension)T5PA; QL (62 EA per 31 days)terbutaline oralT4	SPIRIVA WITH HANDIHALER	Т3	QL (30 EA per 30 days)
SYMDEKOT5PA; QL (56 EA per 28 days)tadalafil (pulm. hypertension)T5PA; QL (62 EA per 31 days)terbutaline oralT4	STIOLTO RESPIMAT	Т3	QL (4 GM per 30 days)
tadalafil (pulm. hypertension) T5 PA; QL (62 EA per 31 days) terbutaline oral	STRIVERDI RESPIMAT	T4	QL (4 GM per 30 days)
terbutaline oral T4	SYMDEKO	T5	PA; QL (56 EA per 28 days)
	tadalafil (pulm. hypertension)	T5	PA; QL (62 EA per 31 days)
THEO-24 T3	terbutaline oral	T4	
	THEO-24	Т3	

Drug Name	Drug Tier	Requirements/Limits
theophylline oral solution	T2	
theophylline oral tablet extended release 12 hr	T2	
theophylline oral tablet extended release 24 hr	T2	
TRELEGY ELLIPTA	Т3	QL (60 EA per 30 days)
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	T5	PA; QL (56 EA per 28 days)
TRIKAFTA ORAL TABLETS, SEQUENTIAL	T5	PA; QL (84 EA per 28 days)
VENTOLIN HFA	Т3	QL (36 GM per 30 days)
WIXELA INHUB	Т3	QL (60 EA per 30 days)
XOLAIR	T5	PA
zafirlukast oral tablet 10 mg	T4	QL (93 EA per 31 days)
zafirlukast oral tablet 20 mg	T4	QL (62 EA per 31 days)
Urologicals		
alfuzosin	T2	QL (31 EA per 31 days)
bethanechol chloride oral tablet 10 mg, 25 mg, 5 mg	T2	
bethanechol chloride oral tablet 50 mg	T3	
CYSTAGON	T4	
dutasteride	T2	QL (31 EA per 31 days)
dutasteride-tamsulosin	T4	QL (31 EA per 31 days)
ELMIRON	T4	
finasteride oral tablet 5 mg	T2	
GEMTESA	T4	QL (31 EA per 31 days)
MYRBETRIQ ORAL SUSPENSION,EXTENDED REL RECON	Т3	QL (300 ML per 30 days)
MYRBETRIQ ORAL TABLET EXTENDED RELEASE 24 HR	Т3	QL (31 EA per 31 days)
oxybutynin chloride oral syrup	Т3	
oxybutynin chloride oral tablet 5 mg	Т3	
oxybutynin chloride oral tablet extended release 24hr 10 mg, 5 mg	Т3	QL (31 EA per 31 days)
oxybutynin chloride oral tablet extended release 24hr 15 mg	Т3	QL (62 EA per 31 days)
potassium citrate oral tablet extended release	T2	
RIVFLOZA SUBCUTANEOUS SOLUTION	T5	PA; QL (1 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML	Т5	PA; QL (0.8 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
RIVFLOZA SUBCUTANEOUS SYRINGE 160 MG/ML	T5	PA; QL (1 ML per 28 days)
silodosin	T4	
tadalafil oral tablet 2.5 mg	T4	PA; QL (62 EA per 31 days)
tadalafil oral tablet 5 mg	T4	PA; QL (31 EA per 31 days)
tamsulosin	T1	
tolterodine oral capsule, extended release 24hr	Т3	QL (31 EA per 31 days)
tolterodine oral tablet	T2	QL (62 EA per 31 days)
trospium oral capsule, extended release 24hr	Т3	QL (31 EA per 31 days)
trospium oral tablet	T2	QL (93 EA per 31 days)
Vitamins, Hematinics / Electrolytes		
CLINIMIX 5%/D15W SULFITE FREE	T4	PA-BvD
CLINIMIX 4.25%/D10W SULF FREE	T4	PA-BvD
CLINIMIX 5%-D20W(SULFITE-FREE)	T4	PA-BvD
fluoride (sodium) oral tablet	T2	
INTRALIPID INTRAVENOUS EMULSION 20 %	T4	PA-BvD
ISOLYTE S PH 7.4	Т3	PA-BvD
ISOLYTE-P IN 5 % DEXTROSE	T4	PA-BvD
KLOR-CON	T4	
KLOR-CON M10	T1	
KLOR-CON M15	T2	
KLOR-CON M20	T1	
magnesium sulfate injection	T2	
PLENAMINE	T4	PA-BvD
potassium chlorid-d5-0.45%nacl	T2	
potassium chloride in 0.9%nacl intravenous parenteral solution 20 meq/l, 40 meq/l	T2	
potassium chloride in 5 % dex intravenous parenteral solution 20 meq/l	T2	
potassium chloride in lr-d5 intravenous parenteral solution 20 meq/l	T2	
potassium chloride in water intravenous piggyback 10 meq/100 ml, 20 meq/100 ml, 40 meq/100 ml	T2	
potassium chloride intravenous	T2	
potassium chloride oral capsule, extended release	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
potassium chloride oral liquid	T2	
potassium chloride oral packet	T2	
potassium chloride oral tablet extended release 10 meq, 20 meq, 8 meq	T1	
potassium chloride oral tablet,er particles/crystals 10 meq, 20 meq	T1	
potassium chloride oral tablet,er particles/crystals 15 meq	T2	
potassium chloride-0.45 % nacl	T2	
potassium chloride-d5-0.2%nacl intravenous parenteral solution 20 meq/l	T2	
potassium chloride-d5-0.9%nacl	T2	
PRENATAL VITAMIN PLUS LOW IRON	T2	PA
sodium chloride 0.45 % intravenous	T2	
sodium chloride 3 % hypertonic	T2	
sodium chloride 5 % hypertonic	T2	
TRAVASOL 10 %	T4	PA-BvD
TROPHAMINE 10 %	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



Index of Drugs

<i>amikacin</i> 3	AVIANE 56
<i>amiloride</i> 33	AVMAPKI-FAKZYNJA11
amiloride-hydrochlorothiazide33	AVONEX 52
<i>amiodarone</i> 33	AYVAKIT 11
<i>amitriptyline</i> 20	<i>azathioprine</i> 11
	azelastine
	azelastine-fluticasone62
	azithromycin4
	aztreonam4
÷	AZURETTE (28) 56
-	<i>bacitracin</i> 60
	bacitracin-polymyxin b60
	baclofen20
	BAFIERTAM20
	balsalazide
*	BALVERSA11
•	BAQSIMI45
	bcg vaccine, live (pf)52
	benazepril
•	benazepril-hydrochlorothiazide33
	BENLYSTA54
	<i>benztropine</i>
	BESIVANCE 60
	BESREMI 52
	betaine
	betamethasone dipropionate39
	betamethasone valerate39
	betamethasone, augmented39
	BETASERON52
	betaxolol
	bethanechol chloride65
v	bexarotene11
* *	BEXSERO
· ·	bicalutamide11
-	BICILLIN C-R4
	BICILLIN L-A4
	BIKTARVY4
1 1 2	bimatoprost60
	bisoprolol fumarate34
	bisoprolol-hydrochlorothiazide34
	BIVIGAM 52
	BOOSTRIX TDAP 52
	BOSULIF 11
	BRAFTOVI 11
atorvastatin33	BREO ELLIPTA
	BREYNA
atovaquone-proguanil4	BREZTRI AEROSPHERE62
<i>atropine</i> 60	<i>brimonidine</i> 60
ATROVENT HFA	brimonidine-timolol60
AUGTYRO 11	BRINSUPRI
AUVELITY 20	brinzolamide60
	amiloride 33 amiloride-hydrochlorothiazide 33 amiodarone 33 amiodipine 20 amlodipine 33 amlodipine-atorvastatin 33 amlodipine-benazepril 33 amlodipine-olmesartan 33 amlodipine-valsartan 33 amnonium lactate 39 AMNESTEM 39 amoxapine 20 amoxicillin 3 amoxicillin-pot clavulanate 3 amphotericin b 4 amphotericin b liposome 4 ampicillin sodium 4 ampicillin sodium 4 ampicillin-sulbactam 4 anastrozole 11 ANORO ELLIPTA 62 ANZUPGO 39 apraclonidine 60 aprepitant 49 APRI 56 APTIVUS 4 ARANELLE (28) 56 AREXVY (PF) 52 arformoterol 62 ASMANEX TWISTHALER 62 <td< td=""></td<>

BRIVIACT	cephalexin5	COBENFY 22
<i>bromfenac</i> 60		COBENFY STARTER PACK.22
bromocriptine20	<i>cetirizine</i> 62	<i>colchicine</i> 54
BRUKINSA 11	<i>cevimeline</i> 43	colesevelam34
<i>budesonide</i> 49, 62	CHEMET 43	<i>colestipol</i> 34
budesonide-formoterol62	chlorhexidine gluconate45	colistin (colistimethate na) 5
bumetanide34	chloroquine phosphate5	COMBIGAN 60
buprenorphine20	chlorpromazine21	COMBIVENT RESPIMAT 62
buprenorphine hcl20		COMETRIQ 12
buprenorphine-naloxone20	cholestyramine (with sugar) 34	COMPRO 49
bupropion hcl20	CHOLESTYRAMINE	CONSTULOSE 49
bupropion hcl (smoking deter)43	LIGHT 34	COPAXONE
buspirone21	CIBINQO 40	COPIKTRA 12
butorphanol21	<i>ciclopirox</i> 40	CORLANOR 34
cabergoline45	<i>cilostazol</i> 34	COSENTYX 40
CABLIVI		COSENTYX (2 SYRINGES)40
CABOMETYX 11	CIMZIA 49	COSENTYX PEN (2 PENS) 40
calcipotriene40	CIMZIA POWDER FOR	COSENTYX UNOREADY
calcitonin (salmon)45		PEN 40
calcitriol45	<i>cinacalcet</i> .45	COTELLIC 12
CALQUENCE	CINRYZE 62	CREON 50
(ACALABRUTINIB MAL) 11	ciprofloxacin hcl5, 60	<i>cromolyn</i> 50, 60, 62
CAMILA 56	ciprofloxacin in 5 % dextrose5	CRYSELLE (28) 57
CAMZYOS 34	ciprofloxacin-dexamethasone45	cyclobenzaprine22
candesartan34	<i>citalopram</i> 21	cyclophosphamide12
candesartan-hydrochlorothiazid.34	CLARAVIS	cyclosporine12, 60
CAPLYTA 21	clarithromycin5	cyclosporine modified12
CAPRELSA11	clindamycin hcl5	CYLTEZO (CF)54
captopril34	clindamycin in 5 % dextrose5	CYLTEZO(CF) PEN 54
carbamazepine21		CYLTEZO(CF) PEN
carbidopa-levodopa21	PEDIATRIC5	CROHN'S-UC-HS 54
carbidopa-levodopa-entacapone 21	clindamycin phosphate 5, 40, 57	CYLTEZO(CF) PEN
carglumic acid43	CLINIMIX 5%/D15W	PSORIASIS-UV 54
carteolol60	SULFITE FREE66	CYRED EQ 57
CARTIA XT 34	CLINIMIX 4.25%/D10W	CYSTAGON 65
carvedilol34	SULF FREE	CYSTARAN 60
caspofungin4	CLINIMIX 4.25%/D5W	d10 %-0.45 % sodium chloride43
CAYSTON4	SULFIT FREE	d2.5 %-0.45 % sodium chloride43
cefaclor4	CLINIMIX 5%-	d5 % and 0.9 % sodium
cefadroxil4	· · · · · · · · · · · · · · · · · · ·	<i>chloride</i> 43
cefazolin4	<i>clobazam</i> 21	<i>d5</i> %-0.45 % sodium chloride43
cefdinir4	<i>clobetasol</i> 40	dalfampridine22
cefepime4	<i>clomipramine</i> 21	<i>danazol</i> 45
cefixime4	<i>clonazepam</i> 21	<i>dantrolene</i> 22
cefoxitin4	<i>clonidine</i> 34	DANZITEN 12
cefpodoxime4	clonidine hcl34	<i>dapsone</i> 5
cefprozil4	clopidogrel34	DAPTACEL (DTAP
ceftazidime4		PEDIATRIC) (PF) 52
ceftriaxone5	<i>clotrimazole</i> 5, 40	<i>daptomycin</i> 5
cefuroxime axetil5	clotrimazole-betame thas one40	darunavir5
cefuroxime sodium5	*	<i>dasatinib</i> 12
celecoxib21	COARTEM 5	DAURISMO 12

DAYBUE	22	DOXY-100	5	entecavir	6
deferasirox	43	doxycycline hyclate	5	ENTRESTO	
deferiprone		doxycycline monohydrate	6	ENULOSE	
DELSTRIGO	5	DRIZALMA SPRINKLE	23	ENVARSUS XR	12
DEPO-SUBQ PROVERA	104 . 57	dronabinol	50	EPCLUSA	6
DESCOVY	5	drospirenone-e.estradiol-lm.f	a 57	EPIDIOLEX	23
desipraminedesipramine	22	drospirenone-ethinyl estradio	l57	epinephrine	63
desloratadine	63	droxidopa	43, 44	eplerenone	
desmopressin	45	DULERA	63	ergotamine-caffeine	
desoximetasone	40	duloxetine	23	ERIVEDGE	12
desvenlafaxine succinate		DUPIXENT PEN	40	ERLEADA	12
dexamethasone	45	DUPIXENT SYRINGE		erlotinib	12
dexamethasone sodium		dutasteride	65	ERRIN	57
phosphate	60	dutasteride-tamsulosin	65	ertapenem	6
dexmethylphenidate		DUVYZAT		ERY PADS	
dextroamphetamine-		econazole nitrate	40	erythromycin	6, 60
amphetamine	22	EDURANT		erythromycin ethylsuccinate	
dextrose 10 % in water (d1		EDURANT PED		erythromycin with ethanol	
dextrose 5 % in water (d5v	,	efavirenz		escitalopram oxalate	
DIACOMIT	*	efavirenz-emtricitabin-tenofo		eslicarbazepine	
diazepam		efavirenz-lamivu-tenofov diso		esomeprazole magnesium	
DIAZEPAM INTENSOL		ELIGARD	•	ESTARYLLA	
diazoxide		ELIGARD (3 MONTH)		estradiol	
diclofenac potassium		ELIGARD (4 MONTH)		estradiol-norethindrone acet	
diclofenac sodium2		ELIGARD (6 MONTH)		eszopiclone	
dicloxacillin		ELIQUIS		ethacrynic acid	
dicyclomine		ELIQUIS DVT-PE TREAT		ethambutol	
DIFICID		30D START		ethosuximide	
diflunisal		ELMIRON		ethynodiol diac-eth estradiol	
difluprednate		eltrombopag olamine		etodolac	
digoxin		ELURYNG		etonogestrel-ethinyl estradiol	
dihydroergotamine		EMGALITY PEN		etravirine	
DILANTIN		EMGALITY SYRINGE		EULEXIN	
diltiazem hcl	34	EMSAM	23	EVENITY	55
DILT-XR		emtricitabine		everolimus (antineoplastic)	
dimethyl fumarate		emtricitabine-tenofovir (tdf)		everolimus	
diphenoxylate-atropine		emtricita-rilpivirine-tenof df		(immunosuppressive)	12
disulfiramdisulfiram		EMTRIVA		EVOTAZ	6
divalproex		EMVERM	6	EVRYSDI	
dofetilide		enalapril maleate	35	exemestane	12
donepezil		enalapril-hydrochlorothiazide		ezetimibe	
DOPTELET (10 TAB PA		ENBREL		ezetimibe-simvastatin	
DOPTELET (15 TAB PA		ENBREL MINI	54	FABHALTA	
DOPTELET (30 TAB PA		ENBREL SURECLICK	55	famciclovir	6
dorzolamide		ENDOCET	23	famotidine	
dorzolamide-timolol		ENGERIX-B (PF)		FANAPT	
dorzolamide-timolol (pf)		ENGERIX-B PEDIATRIC		FANAPT TITRATION	
DOTTI		(PF)	52	PACK A	24
DOVATO		enoxaparin		FARXIGA	
doxazosin		ENPRESSE		FASENRA	
doxepin		ENSKYCE		FASENRA PEN	
doxercalciferol		entacapone		febuxostat	

FEIRZA	. 57	GAMMAGARD LIQUID	.52	haloperidol decanoate2	5
felbamate	.24	GAMMAGARD S-D (IGA < 1		haloperidol lactate2	5
felodipine	35	MCG/ML)	.52	HAVRIX (PF)5	2
fenofibrate	.35	GAMMAKED	52	HEATHER 5	7
fenofibrate micronized	. 35	GAMMAPLEX	52	heparin (porcine)3	6
fenofibrate nanocrystallized	35	GAMMAPLEX (WITH		HEPLISAV-B (PF)5	2
fentanyl	24	SORBITOL)	52	HERNEXEOS1	3
FETZIMA	.24	GAMUNEX-C	.52	HIBERIX (PF) 5	2
FILSUVEZ	41	GARDASIL 9 (PF)	.52	HUMALOG JUNIOR	
finasteride	65	gatifloxacin		KWIKPEN U-100 4	6
fingolimod	. 24	GATTEX 30-VIAL	50	HUMALOG KWIKPEN	
FINTEPLA		GAUZE PAD	.54	INSULIN 4	6
FIRDAPSE	24	GAVILYTE-C	.50	HUMALOG MIX 50-50	
FIRMAGON KIT W		GAVILYTE-G	.50	KWIKPEN 4	6
DILUENT SYRINGE	. 13	GAVILYTE-N	.50	HUMALOG MIX 75-25	
flecainide	.35	GAVRETO	13	KWIKPEN4	6
fluconazole		gefitinib	13	HUMALOG MIX 75-25(U-	
fluconazole in nacl (iso-osm)		gemfibrozil		100)INSULN 4	6
flucytosine		GEMTESA		HUMALOG TEMPO PEN(U-	
fludrocortisone		GENERLAC	.50	100)INSULN 4	6
flunisolide		GENGRAF	13	HUMALOG U-100 INSULIN4	
fluocinolone		gentamicin7, 41,	60	HUMIRA 5	
fluocinolone acetonide oil		gentamicin in nacl (iso-osm)		HUMIRA PEN5	5
fluocinolone and shower cap		GENVOYA		HUMIRA(CF) 5	
fluocinonide		GILOTRIF		HUMIRA(CF) PEN5	
fluocinonide-emollient		glatiramer		HUMIRA(CF) PEN	
fluoride (sodium)		GLATOPA		CROHNS-UC-HS5	5
fluorometholone		GLEOSTINE		HUMIRA(CF) PEN PSOR-	
fluorouracil		glimepiride		UV-ADOL HS5	5
fluoxetine		glipizide45,		HUMULIN 70/30 U-100	
fluoxetine (pmdd)		glipizide-metformin		INSULIN4	6
fluphenazine decanoate		GLUCAGON EMERGENCY		HUMULIN 70/30 U-100	
fluphenazine hcl		KIT (HUMAN)	.46	KWIKPEN4	6
flurbiprofen		glycopyrrolate		HUMULIN N NPH INSULIN	
flurbiprofen sodium		GLYXAMBI		KWIKPEN 4	6
fluticasone propionate 41,		GOMEKLI	13	HUMULIN N NPH U-100	
fluticasone propion-salmeterol		granisetron hcl	.50	INSULIN4	6
fluvastatin		griseofulvin microsize		HUMULIN R REGULAR U-	
fluvoxamine		griseofulvin ultramicrosize		100 INSULN 4	6
fondaparinux		guanfacine		HUMULIN R U-500 (CONC)	
fosamprenavir		GVOKE		INSULIN4	6
fosinopril		GVOKE HYPOPEN 2-PACK.		HUMULIN R U-500 (CONC)	
fosinopril-hydrochlorothiazide		GVOKE PFS 1-PACK		KWIKPEN4	6
FOTIVDA		SYRINGE	46	hydralazine3	
FRUZAQLA		HADLIMA		hydrochlorothiazide3	
FULPHILA		HADLIMA PUSHTOUCH		hydrocodone-acetaminophen2	
FUROSCIX		HADLIMA(CF)		hydrocortisone 41, 46, 5	
furosemide35,		HADLIMA(CF)	-	hydrocortisone-acetic acid 4	
FYCOMPA		PUSHTOUCH	.55	hydromorphone2	
gabapentin		halobetasol propionate		hydroxychloroquine	
galantamine		HALOETTE		hydroxyurea1	
GALLIFREY		haloperidol		hydroxyzine hcl6	
		- T	-	y	-

ibandronate	55	isosorbide mononitrate	36	lactulose	50
IBRANCE	13	isradipine	36	lamivudine	7
IBSRELA	50	ITOVEBI	13, 14	lamivudine-zidovudine	7
IBTROZI	13	itraconazole	7	lamotrigine	
IBU	25	ivabradine	36	lansoprazole	
ibuprofen	25	ivermectin	7	LANTUS SOLOSTAR U-10	
icatibant		IWILFIN	14	INSULIN	47
ICLEVIA		IXIARO (PF)		LANTUS U-100 INSULIN	
ICLUSIG		JAKAFI		lapatinib	
icosapent ethyl	36	JANTOVEN		latanoprost	
IDHİFA		JANUMET		LAZCLUZE	
imatinib		JANUMET XR		leflunomide	
IMBRUVICA		JANUVIA		lenalidomide	
imipenem-cilastatin		JARDIANCE		LENVIMA	
imipramine hcl		JASMIEL (28)		LESSINA	
imiquimod		JAVYGTOR		letrozole	
IMKELDI		JAYPIRCA		leucovorin calcium	
IMOVAX RABIES	10	JENTADUETO		LEUKERAN	
VACCINE (PF)	52	JENTADUETO XR		LEUKINE	
IMVEXXY MAINTENAN		JINTELI		leuprolide	
PACK		JOENJA		leuprolide acetate (3 month)	
IMVEXXY STARTER PA		JULEBER		levalbuterol hcl	
INCASSIA		JULUCA		levalbuterol tartrate	
INCRELEX		JYNNEOS (PF)		levetiracetam	
indapamide		KALETRA		levobunolol	
indomethacin		KALYDECO		levocarnitine	
INFANRIX (DTAP) (PF)		KARIVA (28)		levocarnitine (with sugar)	
INLYTA		KELNOR 1/35 (28)		levocetirizine	
INQOVI		KERENDIA		levofloxacin	
INREBIC		KESIMPTA PEN		levofloxacin in d5w	······ 7
insulin lispro		ketoconazole		LEVONEST (28)	
insulin lispro protamin-lispro		ketorolac	,	levonorgestrel-ethinyl estrad.	
insulin syringe-needle u-100		KEVZARA		levonorg-eth estrad triphasic.	
INTELENCE		KINERET		LEVORA-28	
INTRALIPID		KINRIX (PF)		levothyroxine	
INTROVALE		KINKIX (IT) KIONEX (WITH		LEVOXYL	
INVEGA HAFYERA		SORBITOL)	11	lidocaine	
INVEGA HAFTERAINVEGA SUSTENNA	,	KISQALI		lidocaine hcl	
INVEGA SUSTENNAINVEGA TRINZA		KISQALI FEMARA CO		LIDOCAINE VISCOUS	
IPOL		PACK		lidocaine-prilocaine	
ipratropium bromide		KLOR-CON		LILETTA	
		KLOR-CON M10		linezolid	
ipratropium-albuterolirbesartan		KLOR-CON M15		linezolid in dextrose 5%	
		KLOR-CON M15KLOR-CON M20		LINZESS	
irbesartan-hydrochlorothiazi					
ISENTRESS		KLOXXADO		liothyronine	
ISENTRESS HD		KOSELUGO		lisinopril	
ISIBLOOM		KOURZEQ		lisinopril-hydrochlorothiazide	
ISOLYTE P IN 5 %	00	KRAZATI		LITFULO	
ISOLYTE-P IN 5 %	66	KURVELO (28)		lithium carbonate	
DEXTROSE		l norgest/e.estradiol-e.estr		lithium citrate	
isoniazid		labetalol		LIVTENCITY	
isosorbide dinitrate	36	lacosamide	26	lofexidine	26

LOKELMA	44	MENVEO A-C-Y-W-135-DIP	moxifloxacin-sod.chloride(iso)	8
LONSURF	14	(PF)53	•	
loperamide	50	mercaptopurine15	· · ·	
lopinavir-ritonavir		meropenem		37
lorazepam		mesalamine50, 53	<u> </u>	
LORAZEPAM INTENSOI		mesna1	-	
LORBRENA	14	metformin4	• •	
LORYNA (28)		methadone2		
losartan		methazolamide6		
losartan-hydrochlorothiazide		methenamine hippurate		
lovastatin		methimazole4		
LOW-OGESTREL (28)		methotrexate sodium15		
loxapine succinate		methotrexate sodium (pf)1		
lubiprostone		methsuximide2		
LUMAKRAS		methylphenidate hcl2		
LUMIGAN		methylprednisolone4	-	
LUPRON DEPOT		metoclopramide hcl5		
LUPRON DEPOT (3	13	metolazone		
MONTH)	14 15	metoprolol succinate30		
LUPRON DEPOT (4	17, 13	metoprolol ta-hydrochlorothiaz30	_	
MONTH)	15	metoprolol tartrate30		
LUPRON DEPOT (6	13	metronidazole		
MONTH)	15	metronidazole in nacl (iso-os)	·	
LUPRON DEPOT-PED		metyrosine30		
LUPRON DEPOT-PED (3	13	mexiletine3°		
MONTH)	15	micafungin		
MON111)	13	micajungin	s neomycin-bacuracin-potymyxino	, ,
lunggidana	27	MICONAZOLE 2 50) recommender reclamation le	
lurasidone		MICONAZOLE-3	, , , , ,	
LUTERA (28)	58	MICROGESTIN 1.5/30 (21) 58	3 dexameth6	51
LUTERA (28) LYLEQ	58 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58	3 dexameth68 3 neomycin-polymyxin-gramicidin 6	51 51
LYLEQLYLLANA	58 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30	dexameth6 neomycin-polymyxin-gramicidin 6 neomycin-polymyxin-hc45, 6	51 51 51
LUTERA (28)LYLEQLYLLANALYNPARZA	58 58 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 (28) 58	dexameth	51 51 51
LUTERA (28)LYLEQLYLLANALYNPARZALYSODREN	58 58 58 15	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 (28) 58 MICROGESTIN FE 1/20 (28) . 58	3 dexameth	51 51 51 51
LUTERA (28)LYLEQLYLLANALYNPARZALYSODRENLYTGOBI	58 58 15 15	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 (28) 58 MICROGESTIN FE 1/20 (28) . 58 midodrine 44	dexameth	51 51 51 51 51
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA	58 58 15 15 15 15	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 (28) 58 MICROGESTIN FE 1/20 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47	dexameth	51 51 51 51 51 66 53
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate	58 58 15 15 15 58 66	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47	3 dexameth	51 51 51 51 51 6 53
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion	58 58 15 15 15 58 66 41	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58	dexameth	51 51 51 51 66 53 28 8
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc	58 58 15 15 15 58 66 41	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58 minocycline 58	dexameth	51 51 51 51 51 53 28 8
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28)	58 58 15 15 15 58 66 41 7	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58 minocycline 8 minoxidil 37	dexameth 6 neomycin-polymyxin-gramicidin 6 neomycin-polymyxin-hc 45, 6 NEO-POLYCIN 6 NEO-POLYCIN HC 6 NERLYNX 1 NEULASTA 5 NEUPRO 2 nevirapine 7 NEXLETOL 3 NEXLETOL 3	51 51 51 51 51 65 53 28 8 87
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN	58 58 15 15 15 58 66 41 7	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 42 mifepristone 44 miglustat 47 MILI 58 minocycline 8 minoxidil 3 mirtazapine 2	dexameth	51 51 51 51 53 28 8 8 7 58
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE	58 58 15 15 15 58 66 41 7 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58 minocycline 37 minoxidil 37 mirtazapine 27 misoprostol 51	dexameth 6 neomycin-polymyxin-gramicidin 6 neomycin-polymyxin-hc 45, 6 NEO-POLYCIN 6 NEO-POLYCIN HC 6 NERLYNX 1 NEULASTA 5 NEUPRO 2 nevirapine 7 NEXPLANON 5 niacin 3	51 51 51 51 51 51 53 28 8 8 7 7 88
LUTERA (28) LYLEQ. LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET	58 58 15 15 58 66 41 7 58 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58 minocycline 37 minoxidil 37 mirtazapine 27 misoprostol 55 M-M-R II (PF) 52	dexameth 6 neomycin-polymyxin-gramicidin 6 neomycin-polymyxin-hc 45, 6 NEO-POLYCIN 6 NEO-POLYCIN HC 6 NERLYNX 1 NEULASTA 5 NEUPRO 2 nevirapine 7 NEXPLATOL 3 NEXLETOL 3 NEXLETOL 3 NEXPLANON 5 niacin 3 nicardipine 3	51 51 51 51 51 53 88 87 58 87
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine	58 58 15 15 15 58 66 41 7 58 27 55	MICROGESTIN 1.5/30 (21) 53 MICROGESTIN 1/20 (21) 53 MICROGESTIN FE 1.5/30 53 (28) 53 MICROGESTIN FE 1/20 (28) 53 midodrine 44 mifepristone 47 miglustat 47 MILI 53 minocycline 37 minoxidil 37 mirtazapine 27 misoprostol 55 M-M-R II (PF) 53 modafinil 27	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NERLYNX 1 1 NEULASTA 5 1 NEUPRO 2 1 NEXLETOL 3 1 NEXLIZET 3 1 NEXPLANON 5 1 niacin 3 1 NICOTROL NS 4	51 51 51 51 51 53 28 8 8 7 58 8 8 7 14
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone	58 58 15 15 58 66 41 7 58 27 15	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 44 mifepristone 47 miglustat 47 MILI 58 minocycline 37 minoxidil 37 mirtazapine 27 misoprostol 57 M-M-R II (PF) 57 modafinil 27 MODEYSO 16	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NERLYNX 1 1 NEULASTA 5 1 NEUPRO 2 2 nevirapine 3 3 NEXLETOL 3 4 NEXPLANON 5 5 niacin 3 6 nicardipine 3 NICOTROL NS 4 6 nifedipine 3	51 51 51 51 51 53 88 87 737 14
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine	58 58 15 15 58 66 41 7 58 27 58 27 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 42 mifepristone 47 miglustat 47 MILI 58 minocycline 30 minoxidil 37 mirtazapine 22 misoprostol 55 M-M-R II (PF) 52 modafinil 27 MODEYSO 16 moexipril 37	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NEULASTA 5 9 NEULASTA 5 9 NEUPRO 2 9 nevirapine 3 9 NEXLETOL 3 9 NEXLIZET 3 9 niacin 3 9 nicardipine 3 9 nifedipine 3 9 nilotinib hcl 1	51 51 51 51 51 53 88 87 58 837 84 437
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol	58 58 15 15 15 58 66 41 7 58 15 58 15	MICROGESTIN 1.5/30 (21) 53 MICROGESTIN 1/20 (21) 53 MICROGESTIN FE 1.5/30 53 (28) 53 MICROGESTIN FE 1/20 (28) 53 midodrine 44 mifepristone 47 MILI 53 minocycline 37 minoxidil 37 mirtazapine 27 modafinil 27 MODEYSO 16 moexipril 37 molindone 27	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 9 neomycin-polymyxin-hc 45, 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NELYNX 1 1 NEULASTA 5 NEUPRO 2 3 nevirapine 3 NEXLETOL 3 NEXLIZET 3 NEXPLANON 5 niacin 3 nicardipine 3 NICOTROL NS 4 nilotinib hcl 1 nilutamide 1	51 51 51 51 51 51 53 88 87 58 87 58 87 14
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST	5858585555	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 4e mifepristone 4f miglustat 4f MILI 58 minocycline 3f minoxidil 3f mirtazapine 2f misoprostol 5f M-M-R II (PF) 5f modafinil 2f moexipril 3f molindone 2f mometasone 41, 6f	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NELYNX 1 1 NEULASTA 5 1 NEULASTA 5 1 NEUPRO 2 2 nevirapine 3 3 NEXLETOL 3 4 NEXPLANON 5 5 niacin 3 8 NICOTROL NS 4 6 nifedipine 3 7 nilotinib hcl 1 8 nilotinib hcl 1 9 nilotinib incl 1 1 nilutamide 1 1 nimodipine 3	51 51 51 51 51 53 28 837 58 37 58 37 56 53
LUTERA (28) LYLEQ. LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST MEKTOVI	58 58 58 58 58 58 58 58 66 41 7 58 27 58 7 58 15 58 15 15	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 42 mifepristone 47 miglustat 47 MILI 58 minocycline 30 minoxidil 37 mirtazapine 22 misoprostol 55 M-M-R II (PF) 52 modafinil 27 MODEYSO 16 mometasone 41, 64 montelukast 64	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NEULASTA 5 9 NEULASTA 5 9 NEUPRO 2 9 nevirapine 3 9 NEXLIZET 3 9 NEXPLANON 5 9 nicardipine 3 9 nifedipine 3 9 nilotinib hcl 1 1 nilutamide 1 1 nimodipine 3 NINLARO 1	51 51 51 51 51 53 88 87 58 87 58 87 66 66 67 66
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST MEKTOVI meloxicam	58 58 15 15 58 66 41 7 58 27 50 	MICROGESTIN 1.5/30 (21) 53 MICROGESTIN 1/20 (21) 53 MICROGESTIN FE 1.5/30 53 (28) 53 MICROGESTIN FE 1/20 (28) 53 midodrine 44 mifepristone 4 miglustat 4 MILI 53 minocycline 3 minoxidil 3' mirtazapine 2' misoprostol 5 M-M-R II (PF) 5' modafinil 2' MODEYSO 10 moexipril 3' molindone 2' montelukast 64 morphine 2'	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 9 neomycin-polymyxin-hc 45, 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 9 NELYNX 1 1 NEULASTA 5 NEUPRO 2 3 nevirapine 3 NEXLETOL 3 NEXLIZET 3 NEXPLANON 5 niacin 3 nicardipine 3 NICOTROL NS 4 nilotinib hcl 1 nilutamide 1 nimodipine 3 NINLARO 1 nitazoxanide 1	51 51 51 51 51 53 88 7 37 58 87 58 87 56 66 87 66 87 66 88
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST MEKTOVI meloxicam memantine	58 58 58 15 15 58 66 41 7 58 15 50 50 58 50 58 50 58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 4e mifepristone 4f miglustat 4f MILI 58 minocycline 3f minoxidil 3f mirtazapine 2f misoprostol 5f M-M-R II (PF) 5f modafinil 2f MODEYSO 10 moexipril 3f molindone 2f montelukast 6f morphine 2f morphine 2f morphine 2f	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 8 NEVELYNX 1 9 NEULASTA 5 9 NEULASTA 5 9 NEVELASTA 3 9 NEXLETOL 3 10 NEXLIZET 3 11 NEXPLANON 5 12 NICOTROL NS 4 13 nilotinib hcl 1 14 nilotinib hcl 1 15 nilotinib hcl 1 16 nilotinib hcl 1 17 nitazoxanide 1 18 nitisinone 4	51 51 51 51 51 53 28 837 58 37 58 37 56 66 37 66 84
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST MEKTOVI meloxicam memantine memantine-donepezil	58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 42 mifepristone 47 miglustat 47 MILI 58 minocycline 30 minoxidil 31 mirtazapine 22 misoprostol 55 M-M-R II (PF) 52 modafinil 22 MODEYSO 16 monetasone 41, 64 montelukast 64 morphine 22 morphine concentrate 22 MOUNJARO 44	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 8 NEO-POLYCIN 6 8 NEO-POLYCIN HC 6 8 NEO-POLYCIN HC 6 9 NEULASTA 5 9 NEULASTA 5 9 nevirapine 3 9 NEXLETOL 3 9 NEXLIZET 3 9 NICOTROL NS 4 10 nifedipine 3 11 nilotinib hcl 1 12 nilotinib hcl 1 13 NINLARO 1 14 nitisinone 4 15 NITRO-BID 3	51 51 51 51 51 53 83 7 58 87 7 58 87 7 6 6 87 6 87 6 87 87 87 87 87 87 87 87 87 87 87 87 87
LUTERA (28) LYLEQ LYLLANA LYNPARZA LYSODREN LYTGOBI LYZA magnesium sulfate malathion maraviroc MARLISSA (28) MARPLAN MATULANE MAVYRET meclizine medroxyprogesterone mefloquine megestrol MEKINIST MEKTOVI meloxicam memantine	58	MICROGESTIN 1.5/30 (21) 58 MICROGESTIN 1/20 (21) 58 MICROGESTIN FE 1.5/30 58 (28) 58 MICROGESTIN FE 1/20 (28) 58 midodrine 4e mifepristone 4f miglustat 4f MILI 58 minocycline 3f minoxidil 3f mirtazapine 2f misoprostol 5f M-M-R II (PF) 5f modafinil 2f MODEYSO 10 moexipril 3f molindone 2f montelukast 6f morphine 2f morphine 2f morphine 2f	8 dexameth 6 8 neomycin-polymyxin-gramicidin 6 9 neomycin-polymyxin-hc 45, 6 10 NEO-POLYCIN 6 11 NEO-POLYCIN HC 6 12 NELYNX 1 13 NEULASTA 5 14 NEUPRO 2 15 nevirapine 3 16 NEXLIZET 3 17 NEXPLANON 5 18 nicardipine 3 19 nifedipine 3 10 nilotinib hcl 1 11 nilutamide 1 12 nitazoxanide 1 13 nitrofurantoin 3	51 51 51 51 51 51 53 88 7 37 58 87 37 56 66 87 66 87 88 87 87 88 87 87 88 87 88 87 88 87 88 88

nitrofurantoin monohyd/m-cryst8	OTEZLA	55	pimecrolimus	42
nitroglycerin37, 51	OTEZLA STARTER	55	pimozide	
NIVESTYM53	oxacillin	8	PIMTREA (28)	59
NORDITROPIN FLEXPRO 53	oxaprozin	28	pindolol	
norethindrone (contraceptive)58	oxcarbazepine	28	pioglitazone	47
norethindrone acetate58	_		pioglitazone-metformin	47
norethindrone ac-eth estradiol 58			piperacillin-tazobactam	
norgestimate-ethinyl estradiol58	oxycodone-acetaminophen	28	PIQRAY	16
NORTREL 0.5/35 (28)59		47	pirfenidone	64
NORTREL 1/35 (21)59	PACERONE	37	piroxicam	29
NORTREL 1/35 (28)59	paliperidone	28	pitavastatin calcium	37
NORTREL 7/7/7 (28)59	PALYNZIQ	47	PLEGRIDY	
nortriptyline28	PANRETIN	42	PLENAMINE	66
NORVIR	s pantoprazole	51	podofilox	42
NUBEQA 16	PANZYGA	53	POLYCIN	61
NUCALA64	paricalcitol	47	polymyxin b sulf-trimethoprim.	61
NUEDEXTA28	paroxetine hcl	28	POMALYST	16
NUPLAZID 28	PAXLOVID	8	PORTIA 28	59
NURTEC ODT 28	s pazopanib	16	posaconazole	9
NYAMYC 42	PEDIARIX (PF)	53	potassium chlorid-d5-	
<i>nystatin</i>	PEDVAX HIB (PF)	53	0.45%nacl	66
nystatin-triamcinolone42	peg 3350-electrolytes	51	potassium chloride66	5, 67
NYSTOP 42			potassium chloride in 0.9%naci	l. 66
OCTAGAM 53	PEGASYS	53	potassium chloride in 5 % dex	66
octreotide acetate16	peg-electrolyte soln	51	potassium chloride in lr-d5	66
ODEFSEY			potassium chloride in water	66
ODOMZO 16	pen needle, diabetic	54	potassium chloride-0.45 % nac	l. 67
OFEV 64	PENBRAYA (PF)	53	potassium chloride-d5-	
ofloxacin	penciclovir	42	0.2%nacl	67
OGSIVEO 16	penicillamine	55	potassium chloride-d5-	
OJEMDA 16	penicillin g pot in dextrose	9	0.9%nacl	
OJJAARA 16	penicillin g potassium	9	potassium citrate	65
olanzapine28	penicillin v potassium	9	pramipexole	29
olmesartan37	PENMENVY MEN A-B-C-V	W -	prasugrel hcl	37
olmesartan-amlodipin-hcthiazid.37	Y (PF)	53	pravastatin	37
olmesartan-hydrochlorothiazide.37	PENTACEL (PF)	53	praziquantel	9
olopatadine45	pentamidine	9	prazosin	38
OLUMIANT55	pentoxifylline	37	prednisolone	47
omega-3 acid ethyl esters 37	perampanel	29	prednisolone acetate	61
<i>omeprazole</i> 51	perindopril erbumine	37	prednisolone sodium phosphate	?
ondansetron51	PERIOGARD	45	48	3, 61
ondansetron hcl51	permethrin	42	prednisone	48
ONUREG16	perphenazine	29	pregabalin	
OPIPZA 28	PERSERIS	29	PREMARIN	59
OPSUMIT 64	PHEBURANE	44	PREMPRO	59
OPSYNVI 64	phenelzine	29	PRENATAL VITAMIN	
ORENCIA55	phenobarbital	29	PLUS LOW IRON	67
ORENCIA CLICKJECT 55	PHENYTEK	29	PREVALITE	38
ORGOVYX 16	1 2	29	PREVYMIS	
ORKAMBI 64	1 2		PREZCOBIX	
ORSERDU 16			PREZISTA	
oseltamivir	pilocarpine hcl	14, 61	PRIFTIN	9

primaquine	9	REPATHA SURECLICK	38	SEREVENT DISKUS64
primidone	29	REPATHA SYRINGE	38	<i>sertraline</i> 30
PRIORIX (PF)	53	RESTASIS	61	SETLAKIN 59
PRIVIGEN	53	RESTASIS MULTIDOSE	E61	SHINGRIX (PF) 53
probenecid	55	RETACRIT	53	SIGNIFOR 17
probenecid-colchicine	55	RETEVMO	17	sildenafil (pulm.hypertension) 64
prochlorperazine		REVUFORJ	17	silodosin66
prochlorperazine maleate	51	REXULTI	29	silver sulfadiazine42
PROCRIT	53	REYATAZ	9	SIMBRINZA 61
PROCTO-MED HC	51	REZDIFFRA	44	SIMLANDI(CF) 56
PROCTOSOL HC	51	REZLIDHIA	17	SIMLANDI(CF)
PROCTOZONE-HC	51	RHOPRESSA	61	AUTOINJECTOR56
progesterone micronized	59	ribavirin	9	SIMPONI 56
PROGRAF		rifabutin	9	<i>simvastatin</i> 38
PROLASTIN-C	44	rifampin	9	<i>sirolimus</i> 17
PROLENSA	61	riluzole	44	SIRTURO 9
PROLIA	55	rimantadine	9	SKYCLARYS 30
promethazine	64	RINVOQ	56	SKYRIZI
propafenone		RINVOQ LQ	56	sodium chloride
propranolol		risedronate		<i>sodium chloride 0.45 %</i> 67
propylthiouracil		risperidone		<i>sodium chloride</i> 0.9 %
PROQUAD (PF)		risperidone microspheres		sodium chloride 3 % hypertonic . 67
protriptyline		ritonavir		sodium chloride 5 % hypertonic . 67
PULMOZYME		rivastigmine		<i>sodium oxybate</i> 30
pyrazinamide		rivastigmine tartrate		sodium phenylbutyrate44
pyridostigmine bromide		RIVFLOZA		sodium polystyrene sulfonate44
pyrimethamine		rizatriptan		sodium,potassium,mag sulfates 51
QINLOCK		ROCKLATAN		sofosbuvir-velpatasvir9
QUADRACEL (PF)		roflumilast		SOHONOS44
quetiapine		ROMVIMZA		SOLIQUA 100/33 48
quinapril		ropinirole	30	SOLTAMOX 17
quinapril-hydrochlorothiazid		rosuvastatin		SOMAVERT 48
quinidine sulfate		ROTARIX		<i>sorafenib</i> 17
quinine sulfate		ROTATEQ VACCINE		sotalol38
QULIPTA		ROWEEPRA		SOTALOL AF 38
QVAR REDIHALER		ROZLYTREK		SPIRIVA RESPIMAT64
RABAVERT (PF)		RUBRACA		SPIRIVA WITH
rabeprazole		rufinamide		HANDIHALER 64
RADICAVA ORS STARTI		RUKOBIA		spironolactone38
KIT SUSP		RYBELSUS		spironolacton-hydrochlorothiaz. 38
RALDESY		RYDAPT		SPRINTEC (28)
raloxifene		RYTARY		SPRITAM 30
ramelteon		sacubitril-valsartan		SPS (WITH SORBITOL)44
ramipril		SAJAZIR		SRONYX
ranolazine		sapropterin		SSD
rasagiline		SAVELLA		STELARA 42
RAVICTI		SCEMBLIX		STEQEYMA
RECLIPSEN (28)		scopolamine base		STIOLTO RESPIMAT64
RECOMBIVAX HB (PF)		SECUADO		STIVARGA
RECORLEV		selegiline hcl		streptomycin9
RELENZA DISKHALER		selenium sulfide		STRIBILD 10
repaglinide		SELZENTRY		STRIVERDI RESPIMAT64
r -0		, ,		

SUBVENITE	30	teriparatide	56	TRIDERM	43
sucralfate	51	testosterone	48	trientine	
sulfacetamide sodium	61	testosterone cypionate	48	TRI-ESTARYLLA	59
sulfacetamide sodium (acne)	42	testosterone enanthate	48	trifluoperazine	31
sulfacetamide-prednisolone	61	tetrabenazine	31	trifluridine	
sulfadiazine		tetracycline	10	TRIJARDY XR	
sulfamethoxazole-trimethoprin		THALOMID		TRIKAFTA	65
SULFAMYLON		THEO-24		TRI-LEGEST FE	
sulfasalazine	51	theophylline		TRI-LO-ESTARYLLA	59
sulindac		thioridazine		TRI-LO-SPRINTEC	
sumatriptan		thiothixene		trimethoprim	
sumatriptan succinate		TIADYLT ER		trimipramine	
sunitinib malate		tiagabine		TRINTELLIX	
SUNLENCA		TIBSOVO		TRI-SPRINTEC (28)	
SYEDA		ticagrelor		TRIUMEQ	
SYMDEKO	64	TICOVAC		TRIUMEQ PD	
SYMLINPEN 120		tigecycline		TROPHAMINE 10 %	
SYMLINPEN 60		TILIA FE		trospium	
SYMPAZAN		timolol maleate3		TRULICITY	
SYMTUZA		tinidazole	,	TRUMENBA	
SYNAREL		TIVICAY		TRUQAP	
SYNJARDY		TIVICAY PD		TUKYSA	
SYNJARDY XR		tizanidine		TURALIO	
SYNTHROID		TOBI PODHALER		TURQOZ (28)	
TABLOID		tobramycin1		TWINRIX (PF)	
TABRECTA		tobramycin in 0.225 % nacl		TYBOST	
tacrolimus1		tobramycin sulfate		TYMLOS	
tadalafil	,	tobramycin-dexamethasone		TYPHIM VI	54
tadalafil (pulm. hypertension)		tolterodine		UBRELVY	31
TAFINLAR		tolvaptan		UNITHROID	49
TAGRISSO		topiramate		UPTRAVI 38	3, 39
TALTZ AUTOINJECTOR		toremifene		ursodiol	51
TALTZ SYRINGE		torsemide		valacyclovir	10
TALZENNA	17	TOUJEO MAX U-300		VALCHLOR	
tamoxifen	17	SOLOSTAR	48	valganciclovir	
tamsulosin		TOUJEO SOLOSTAR U-300)	valproic acid	
TASCENSO ODT		INSULIN	48	valproic acid (as sodium salt)	
tasimelteon	31	TRADJENTA		valsartan	
tazarotene		tramadol	31	valsartan-hydrochlorothiazide	39
TAZVERIK	17	tramadol-acetaminophen		VALTOCO	
TEFLARO	10	trandolapril		VALTYA	
telmisartan	38	tranexamic acid		vancomycin	10
telmisartan-amlodipine	38	tranylcypromine		VANFLYTA	
telmisartan-hydrochlorothiazi		TRAVASOL 10 %		VAQTA (PF)	54
TENIVAC (PF)		travoprost	61	varenicline tartrate	
tenofovir disoproxil fumarate.	10	trazodone		VARIVAX (PF)	54
TEPMETKO		TRELEGY ELLIPTA		VAXCHORA VACCINE	
terazosin	38	TRELSTAR	18	VELIVET TRIPHASIC	
terbinafine hcl		tretinoin4	2, 43	REGIMEN (28)	59
terbutaline		tretinoin (antineoplastic)		VEMLIDY	
terconazole	59	triamcinolone acetonide 4		VENCLEXTA	18
teriflunomide	31	triamterene-hydrochlorothiazid	d38		

VENCLEXTA STARTING	XIIDRA62
PACK 18	XOFLUZA 11
venlafaxine32	XOLAIR 65
VENTOLIN HFA65	XOLREMDI 54
verapamil39	XOSPATA 19
VERQUVO 39	XPOVIO 19
VERSACLOZ32	XTANDI 19
VERZENIO 18	XULTOPHY 100/3.6 49
VESTURA (28) 59	XYREM 32
VIBERZI 51	YARGESA49
VIENVA 59	YESINTEK 43
vigabatrin32	YF-VAX (PF) 54
VIGADRONE 32	YONSA
VIGPODER32	YORVIPATH 49
VIJOICE 18	YUFLYMA(CF) 56
vilazodone32	YUFLYMA(CF) AI
VIMKUNYA54	CROHN'S-UC-HS 56
VIRACEPT 10	YUFLYMA(CF)
VIREAD10	AUTOINJECTOR56
VITRAKVI18	YUVAFEM59
VIVITROL 32	ZAFEMY 59
VIVJOA 10	zafirlukast65
VIVOTIF 54	<i>zaleplon</i> 32
VIZIMPRO 18	ZARXIO
VONJO	ZAVZPRET32
VORANIGO 18	ZEGALOGUE
voriconazole10, 11	AUTOINJECTOR49
VOSEVI 11	ZEGALOGUE SYRINGE 49
VOWST 51	ZEJULA 19
VRAYLAR 32	ZELBORAF 19
VUMERITY 32	ZELSUVMI 43
VYNDAQEL	ZEPOSIA 32
warfarin39	ZEPOSIA STARTER KIT
WELIREG 18	(28-DAY)32
WIXELA INHUB 65	ZEPOSIA STARTER PACK
XALKORI 19	(7-DAY)32
XARELTO 39	zidovudine11
XARELTO DVT-PE TREAT	ZIEXTENZO 54
30D START 39	ZILBRYSQ 32, 33
XATMEP 19	<i>ziprasidone hcl.</i> 33
XCOPRI 32	ziprasidone mesylate33
XCOPRI MAINTENANCE	ZIRGAN
PACK 32	ZOLINZA 19
XCOPRI TITRATION PACK 32	<i>zolpidem</i> 33
XDEMVY 61	ZONISADE
XELJANZ 56	<i>zonisamide</i> 33
XELJANZ XR 56	ZOVIA 1-35 (28) 59
XELRIA FE 59	ZTALMY 33
XERMELO 19	ZUBSOLV 33
XGEVA 19	ZURZUVAE 33
XIFAXAN 11	ZYDELIG 19
XIGDUO XR 49	ZYKADIA 19

Section: Prior Authorization

Acitretin

Products Affected

• acitretin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Actemra

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide). For giant cell arteritis, patients must have therapeutic failure or intolerance to one systemic corticosteroid (e.g., prednisone). For polyarticular juvenile idiopathic arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For systemic sclerosis-associated interstitial lung disease (SSc-ILD), patients must have therapeutic failure or intolerance to one immunosuppressant (e.g., mycophenolate mofetil, corticosteroids, cyclophosphamide). Documentation of systemic juvenile idiopathic arthritis.
Age Restrictions	Deny if less than 18 years of age for systemic sclerosis-associated interstitial lung disease (SSc-ILD), Rheumatoid Arthritis, and Giant Cell Arteritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Actimmune

Products Affected

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Adbry

Products Affected

• ADBRY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: Dupixent and Rinvoq. For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADHD Drugs

Products Affected

• guanfacine oral tablet extended release 24 hr

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Afinitor

- everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg
- everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For everolimus only, documentation of advanced, hormone receptorpositive, HER2-negative breast cancer -AND- all of the following (1-3): 1) member is a postmenopausal woman 2) using in combination with exemestane 3) therapeutic failure or intolerance to prior treatment with letrozole or anastrozole. For everolimus only, documentation of nonfunctional neuroendocrine tumors -AND- all of the following (1-2): 1) disease is classified as progressive, well-differentiated, non-functional 2) disease is of gastrointestinal or lung origin that are unresectable, locally advanced or metastatic. For everolimus only, documentation of advanced renal cell carcinoma -AND- therapeutic failure or intolerance to prior treatment with sunitinib or sorafenib. For everolimus only, documentation of renal angiomyolipoma and tuberous sclerosis complex (TSC), member does not require immediate surgery. For everolimus and everolimus tablets for oral suspension, documentation of TSC with Subependymal Giant Cell Astrocytoma -AND- member is not a candidate for curative surgical resection. For everolimus only, documentation of progressive neuroendocrine tumors of pancreatic origin -AND- disease is unresectable, locally advanced or metastatic. For everolimus tablets for oral suspension only, documentation of use for adjunctive treatment of TSC-associated partial-onset seizures.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Aimovig

Products Affected

• AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML, 70 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ajovy

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-4). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) For episodic migraine, if greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 45 kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Akeega

Products Affected

AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) BRCA mutations, 2) Comcomitant therapy with prednisone, 3) Concomitant therapy with a gonadotropin-releasing hormone analog or a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alecensa

Products Affected

· ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test - AND- One of the following (1-2): 1) metastatic disease, 2) will be used as adjuvant treatment following tumor resection of node positive or greater than or equal to 4 cm tumor(s).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alpha1-Proteinase Inhibitors

Products Affected

• PROLASTIN-C INTRAVENOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of panacinar emphysema AND documentation of a decline in forced expiratory volume in 1 second (fev1) despite medical therapy (bronchodilators, corticosteroids) AND documentation of phenotype (pi*zz, pi*znull or pi*nullnull) associated with causing serum alpha 1-antitrypsin of less than 80 mg/dl AND documentation of an alpha 1-antitrypsin serum level below the value of 35% of normal (less than 80 mg/dl).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when furnished incident to a physician service and is not self-administered.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Alunbrig

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ampyra

Products Affected

• dalfampridine

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure disorder, Cr Cl less than 50ml/min
Required Medical Information	Documentation of diagnosis -AND- documentation that the patient is ambulatory and has walking impairment as evidenced by one of the following. 1. Functional status score (EDSS score). 2. Timed 25-foot Walk Test (T25W).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement in walking impairment from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Anzupgo

Products Affected

· ANZUPGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic hand eczema (CHE) -AND- symptoms persisting for more than 3 months or recurring at least twice within a 12-month timeframe -AND- therapeutic failure, contraindication, or intolerance to 1 generic, on-formulary, medium/high/super-high potency topical corticosteroid.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Arikayce

Products Affected

• ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Mycobacterium avium complex lung disease -AND-Attestation of not achieving negative sputum cultures despite at least 6 months with a multidrug background regimen containing 2 of the following: 1) macrolide 2) rifamycin or 3) ethambutal -AND- Arikayce will be used in conjunction with a background multidrug regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive sputum cultures -OR- negative sputum cultures for insufficient period of time (e.g. less than 12 months).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ATTR-CM drugs

Products Affected

• VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	Concomitantly with transthyretin-lowering agents
Required Medical Information	Documentation of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) with amyloid deposits on cardiac biopsy or scintigraphy with heart contralateral greater than 1.5 (Grade III) -AND- Cardiac involvement supported by cardiac magnetic resonance, echocardiography or serum cardiac biomarker (e.g. B-type natriuretic peptide, cardiac troponin) - AND- Primary (light chain) amyloidosis has been ruled out by immunohistochemistry, mass spectrometry or scintigraphy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement or delayed disease progression from baseline demonstrated by 6-minute walk test, cardiac function (e.g. LVEF, NYHA class), Kansas City Cardiomyopathy Questionnaire-Overall Summary, number of cardiovascular-related hospitalizations or serum cardiac biomarkers (e.g. B-type natriuretic peptide, cardiac troponin)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Atypical Antipsychotics

- aripiprazole oral solution aripiprazole oral tablet,disintegrating

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Aubagio

Products Affected

• teriflunomide

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as fingolimod, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Augtyro

Products Affected

• AUGTYRO ORAL CAPSULE 160 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive. Documentation of solid tumors - AND- disease harbors a NTRK gene fusion -AND- one of the following (1-2): 1) disease is locally advanced or metastatic, or 2) surgical resection is likely to result in severe morbidity -AND- one of the following (3-4): 3) disease has progressed following treatment, or 4) the member has no satisfactory alternative therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Auvelity

Products Affected

AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND-Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Avmapki-Fakzynja

Products Affected

• AVMAPKI-FAKZYNJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent low-grade serous ovarian cancer (LGSOC), disease is KRAS-mutated -AND- member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ayvakit

Products Affected

AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable or metastatic gastrointestinal stromal tumor -AND- tumors harbor a PDGFRA exon 18 mutation. Documentation of agressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm, mast cell leukemia, or indolent systemic mastocytosis -AND- platelet count greater than or equal to 50 x 10*9/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bafiertam

Products Affected

BAFIERTAM

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Balversa

Products Affected

• BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) FGFR3 mutation status as detected by an FDA approved test 2) disease progression on or after at least on prior line of systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Banzel

Products Affected

• rufinamide

PA Criteria	Criteria Details
Exclusion Criteria	Patients with familial short QT syndrome
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND-documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Benlysta

Products Affected

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of active systemic lupus erythematosus (SLE) -AND-documentation of positive anti-nuclear antibody (ANA) titer (greater than or equal to 1:80) or anti-double-stranded DNA antibody (anti-dsDNA) greater than or equal to 30IU/mL -AND- member will continue to receive concomitant standard of care treatment with use of at least one of the following (alone or in combination): 1.) corticosteroids (e.g. prednisone) 2.) antimalarials (e.g. hydroxychloroquine) 3.) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate) -OR-Documentation of active lupus nephritis -AND- documentation of positive ANA titer (greater than or equal to 1:80) or anti-dsDNA greater than or equal to 30 IU/mL -AND- member will continue to receive concomitant standard of care treatment which includes corticosteroids (e.g. prednisone) with at least one of the following: 1.) mycophenolate for induction followed by mycophenolate for maintenance 2.) cyclophosphamide for induction followed by azathioprine for maintenance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For SLE reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with corticosteroids, antimalarials, or immunosuppressives. For active LN reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with mycophenolate or azathioprine. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Besremi

Products Affected

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Bosulif

Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy. For pediatric patients 1 year of age and older, one of the following (3, 4): 3) newly diagnosed PH-positive CML in the chronic phase 4) diagnosis of Ph-positive CML in the chronic phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Bosutinib capsules and 18 years of age or older, inability to swallow tablets is required. For Bosutinib 100mg capsules and pediatric 1 year of age or older, inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Braftovi

Products Affected

• BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	Use in wild-type BRAF melanoma or wild-type BRAF CRC
Required Medical Information	Documentation of diagnosis. For metastatic colorectal cancer (mCRC) and using in combination with cetuximab, all of the following (1-3): 1) BRAF V600E mutation status 2) using in combination with cetuximab 3) member has received prior therapy for CRC. For mCRC and using in combination with cetuximab and modified FOLFOX6, all of the following (4-5): 4) BRAF V600E mutation status, as detected by an FDA-approved test 5) using in combination with cetuximab and modified FOLFOX6. For unresectable or metastatic melanoma, all of the following (6-7): 6) BRAF V600E or V600K mutation status 7) using in combination with binimetinib. For metastatic non-small cell lung cancer, all of the following (8-9): 8) BRAF V600E mutation status 9) using in combination with binimetinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Brinsupri

Products Affected

• BRINSUPRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of non-cystic fibrosis bronchiectasis (NCFB) confirmed by computed tomography (CT) -AND- at least 1 of the following symptoms consistent with bronchiectasis (1-8): 1) cough on most days of the week, 2) chronic sputum production, 3) history of recurrent respiratory infections, 4) dyspnea, 5) wheezing, 6) rhinosinusitis, 7) hemoptysis, 8) recurrent pleurisy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Brukinsa

Products Affected

• BRUKINSA ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma (MCL), previous treatment with at least 1 prior therapy. For marginal zone lymphoma (MZL), previous treatment with at least 1 anti-CD20-based regimen. For follicular lymphoma (FL), using in combination with obinutuzumab and previous treatment with at least 2 prior lines of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Buphenyl

Products Affected

• sodium phenylbutyrate

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cablivi

Products Affected

• CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of acquired thrombocytopenic purpura (aTTP) -AND- Used in conjunction with plasma exchange and immunosuppressive therapy (i.e. systemic corticosteroids or rituximab)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	75 days initial authorization, 28 days reauthorization
Other Criteria	For reauthorization, attestation of remaining signs and symptoms of persistent disease (e.g. suppressed ADAMTS 13 activity level remain present)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cabometyx

Products Affected

· CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-6): 1) diagnosis of advanced renal cell carcinoma (RCC), 2) diagnosis of advanced RCC and using as a first-line treatment in combination with nivolumab, 3) member has previously been treated with sorafenib for hepatocellular carcinoma, 4) member has experienced disease progression following prior VEGFR-targeted therapy and is either radioactive iodine-refractory or is ineligible for radioactive iodine therapy for locally advanced or metastatic differentiated thyroid cancer, 5) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated pancreatic neuroendocrine tumors (pNET), 6) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated extra-pancreatic neuroendocrine tumors (epNET)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Calquence

Products Affected

• CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma, member has received at least one prior therapy -OR- all of the following (1-3): 1) member has not received prior therapy for MCL, 2) member is ineligible for autologous hematopoietic stem cell transplantation (HSCT), and 3) using in combination with bendamustine and rituximab.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Camzyos

Products Affected

· CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy -AND- Left ventricular ejection fraction of greater than or equal to 55% -AND- Valsalva left ventricular outflow tract peak gradient of greater than or equal to 50 mmHg at rest or after provocation -AND- Therapeutic failure or intolerance to one of the following (1 or 2) or contraindication to all: 1) Non-vasodilating beta blocker (e.g. metoprolol) 2) non-dihydropyridine calcium channel blocker (e.g. diltiazem) -AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, prescriber attestation of no NYHA class worsening - AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Caplyta

Products Affected

· CAPLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Caprelsa

Products Affected

• CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic or progressive medullary thyroid cancer in patients with unresectable locally advanced or metastatic disease.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Carbaglu

Products Affected

• carglumic acid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use as an adjunct therapy for acute hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA) -OR-maintenance therapy for chronic hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Cayston

Products Affected

· CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cerdelga

Products Affected

· CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of type 1 Gaucher disease confirmed by the following A. or B. A) With one of the following symptoms (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm*3. 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia)OR- B) Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles -AND- Documentation of CYP2D6 metabolizer status (e.g. intermediate metabolizer).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CF drugs

Products Affected

- TOBI PODHALER
- tobramycin in 0.225 % nacl
- tobramycin inhalation

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization of tobramycin products, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cialis

Products Affected

• tadalafil oral tablet 2.5 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of benign prostatic hyperplasia (BPH) and trial/failure of at least two alternative medications in the following classes (alpha-1 adrenergic blockers and/or 5-alpha reductase inhibitors)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cibinqo

Products Affected

· CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: Dupixent and Rinvoq.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cimzia

Products Affected

- CIMZIA POWDER FOR RECONST
- CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide) or all nonbiologic DMARDs are contraindicated. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or all are contraindicated. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR-inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- all nonbiologic DMARDs are contraindicated -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity, or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age for juvenile idiopathic arthritis or less than 18 years of age for all other indications.
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For Crohn's disease, patients must have therapeutic failure or intolerance to 2 of the following preferred biologic products: a preferred adalimumab product, Stelara, Rinvoq, and Skyrizi SC. For Rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Xeljanz/Xeljanz XR and Rinvoq. For plaque psoriasis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Cosentyx, Otezla, Stelara SC, Skyrizi SC, and Enbrel. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For Psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cosentyx, Enbrel, a preferred adalimumab product, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For Juvenile Idiopathic Arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Rinvoz/Rinvoq LQ, Xeljanx/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For initial and induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended initial and induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cinryze

Products Affected

· CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Cobenfy

Products Affected

- · COBENFY
- COBENFY STARTER PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cometriq

Products Affected

 COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressive, metastatic medullary thyroid cancer
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Copiktra

Products Affected

· COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is no longer responding or is intolerant to at least 2 prior therapies for chronic lymphocytic leukemia and small lymphocytic leukemia.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Corlanor

Products Affected

- CORLANOR ORAL SOLUTION
- ivabradine oral tablet 5 mg, 7.5 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) Normal sinus rhythm, 2) Resting heart rate greater than or equal to 70 beats per minute, 3) Left ventricular ejection fraction less than or equal to 35 percent, when applicable, 4) In adult patients (greater than or equal to 18 years), concurrent use, therapeutic failure, or intolerance to the maximum tolerated dose of one beta-blocker used for treatment of heart failure (i.e. bisoprolol, carvedilol, metoprolol succinate), or contraindication to beta-blocker use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For oral solution, attestation of inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cosentyx

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML

• COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR-inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs or all are contraindicated.
Age Restrictions	Deny if less than 6 years of age for moderate to severe plaque psoriasis - OR- less than 2 years of age for psoriatic arthritis -OR- less than 4 years of age for enthesitis-related arthritis -OR- less than 18 years of age for all other indications
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication. For hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cotellic

Products Affected

· COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following for unresectable or metastatic melanoma (1-2): 1) BRAF V600E or V600K mutation status 2) Concomitant therapy with vemurafenib. For cobimetinib monotherapy, documentation of histiocytic neoplasms.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Cysteamine Ophthalmic Drops

Products Affected

CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystinosis -AND- Attestation of accumulation of corneal cystine crystals
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Danziten

Products Affected

• DANZITEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib -OR- member is newly diagnosed in the chronic phase.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daraprim

Products Affected

• pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For primary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 100 cells/mm*3 -AND- Toxoplasma IgG positive -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For secondary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 200 cells/mm*3. For secondary prophylaxis of cystoisosporiasis with CD4 count less than 200 cells/mm*3 or acute cystoisosporiasis infection: failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For primary prophylaxis of Pneumocystis jirovecii pneumonia: diagnosis of HIV -AND- CD4 count less than 200 cells/mm*3 -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Daurismo

Products Affected

• DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of newly diagnosed Acute Myeloid Leukemia -AND-Used in combination with cytarabine -AND- At least one comorbidity that preclude use of intensive induction chemotherapy defined as one of the following: 1) Age greater than or equal to 75 2) Severe cardiac or pulmonary comorbidity 3) Reduced renal function 4) Hepatic impairment 5.) Physician attests patient is not a candidate for intensive induction therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Daybue

Products Affected

• DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Rett syndrome confirmed by all of the following (1 thru 4): 1) Partial or complete loss of acquired purposeful hand skills, 2) Partial or complete loss of acquired spoken language, 3) Gait abnormalities: impaired or absence of ability, 4) Stereotypic hand movements such as hand wringing/squeezing, clapping/tapping, mouthing, and washing/rubbing automatisms
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Deferasirox

Products Affected

• deferasirox

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic iron overload due to blood transfusions, transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -And-history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw). For Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia (NTDT) Syndrome, LIC of at least 5 mg Fe/g dw -AND- serum ferritin greater than 300 mcg/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of chronic iron overload due to blood transfusion, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw. For reauthorization of chronic iron overload in NTDT syndrome, LIC greater than or equal to 3 mg Fe/g dw.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Diacomit

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravets syndrome - AND- Used in combination with clobazam
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dihydroergotamine

Products Affected

• dihydroergotamine nasal

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND-requires non-oral route of administration -AND- therapeutic failure or intolerance to generic sumatriptan nasal spray.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Doptelet

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND-beneficiary is scheduled to undergo a procedure -OR- Documentation of persistent or chronic immune thrombocytopenia -AND- Trial, intolerance, or inadequate response to corticosteroid therapy, immunoglobulin therapy or splenectomy -AND- One of the following (1 or 2): 1) Platelet count less than or equal to 50 x 10*9/L and has significant mucous member bleeding or at least one risk factor for bleeding (e.g. hypertension, peptic ulcer disease). 2) Platelets count of less than or equal to 30 x 10*9/L
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For thrombocytopenia with chronic liver disease- 1 mo. For immune thrombocytopenia- 12 mo.
Other Criteria	Platelet count is provided for applicable dosing.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Drizalma

Products Affected

• DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG, 30 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow tablets/capsules. For fibromyalgia, members must also have widespread bilateral pain above and below the waist for greater than 3 months duration -AND- At least 1 fibromyalgia-related symptom (e.g., cognitive impairment, fatigue, sleep disturbance, neurologic symptoms, exercise intolerance).
Age Restrictions	Deny if less than 18 years of age in the treatment of fibromyalgia, major depressive disorder, diabetic peripheral neuropathy and chronic musculoskeletal pain -OR- if less than 7 years of age in generalized anxiety disorder
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Dupixent

Products Affected

 DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML

• DUPIXENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Moderate/Severe atopic dermatitis (AD): failure/intolerance to 1 topical corticosteroid or, if 2 yrs or older, topical calcineurin inhibitor (e.g. tacrolimus, pimecrolimus) -OR- severe AD and unable to apply topical tx due to the extent of body surface area involvement or topical tx are contraindicated due to severely damaged skin. Moderate/severe asthma: history of 2 or more exacerbations requiring oral/injectable corticosteroids in past 12mos or 1 or more exacerbation requiring hospitalization in past 12mos -AND- blood eosinophils of 150cells/uL or higher or current daily/alternate-day oral corticosteroid (OCS) -AND- inadequate control despite regular medium/high-dose inhaled corticosteroids (ICS) + 1 or more add'l asthma controller (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline) +/- OCS, unless intolerant or contraindicated to all -AND- will continue medium/ high-dose ICS + 1 or more add'l asthma controller medication +/- OCS. Chronic rhinosinusitis with nasal polyposis: failure/intolerance/contraindication to intra-nasal corticosteroid + 14-day OCS. Eosinophilic esophagitis: esophageal eosinophils of 15eos/hpf or higher on esophageal biopsy -AND- clinical esophageal dysfunction symptoms. Prurigo nodularis. COPD: blood eosinophils of at least 300cells/uL or current daily/alternate-day OCS -AND- inadequate control despite 3 or more months of regular LAMA, LABA, and ICS tx, unless intolerant or contraindicated to all. Chronic Spontaneous Urticaria: failure/contraindication/intolerance to 1 second-generation non-sedating H1 antihistamine at max recommended dose (e.g. cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine). Bullous pemphigoid: failure/intolerance/contraindication on high/super high potency topical corticosteroid or oral corticosteroid.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For atopic dermatitis reauthorization, attestation of positive clinical response to therapy. For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For EoE reauthorization, attestation of histological remission (less than 15 eos/hpf) on esophageal biopsy or reduced severity or frequency of clinical symptoms of esophageal dysfunction. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline. For COPD reauthorization, attestation of one of the following is required (1-4): 1) reduction in COPD symptoms, 2) improvement in exercise tolerance, 3) delayed disease progression, or 4) reduction in the number of COPD exacerbations. For CSU reauthorization, improved CSU symptoms. For BP reauthorization, attestation of disease control, reduction in number of relapses, improvement in BP symptoms or reduction in oral corticosteroid use.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Duvyzat

Products Affected

• DUVYZAT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy with pathogenic mutation in the dystrophin gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EGFR Tyrosine Kinase Inhibitors

- erlotinib
- · GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	Afatinib products: tumors with resistant EGFR mutations. Erlotinib products: use in NSCLC tumors with mutations other than those in FDA-approved indications. Use in combination with platinum based chemotherapy.
Required Medical Information	For afatinib, documentation of metastatic non-small cell lung cancer (NSCLC) -AND- one of the following, as detected by an FDA-approved test (1-3): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutation 3) disease harbors non-resistant EGFR mutation (i.e., S768I, L861Q, G719X) -OR-documentation of squamous metastatic NSCLC and member has experienced progression on platinum-based chemotherapy. For erlotinib, documentation of metastatic NSCLC -AND- one of the following, as detected by an FDA-approved test (1-2): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutations -OR- documentation of locally advanced, unresectable or metastatic pancreatic cancer -AND- all of the following (1-2): 1) using erlotinib as first-line therapy 2) using in combination with gemcitabine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Emgality

- EMGALITY PEN
- EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For episodic cluster headaches, characterized by severe or very severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes when left untreated -AND-Attack frequency of at least one attack every other day during the cluster period.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency is required. For reauthorization of cluster headache, attestation of reduction in the number of mean weekly cluster headaches from baseline is required.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Enbrel

Products Affected

• ENBREL MINI

- ENBREL SURECLICK
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR-contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Ankylosing Spondylitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis or Less than 4 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epclusa

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG
- sofosbuvir-velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Epidiolex

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Lennox-Gastaut, Dravet syndromes or Tuberous Sclerosis Complex. For Lennox-Gastaut, trial and failure or intolerance of at least two standard of care treatments (e.g. lamotrigine, clobazam). For Lennox-Gastaut and Dravet syndromes, treatment is in combination with other conventional agents.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Eprontia

Products Affected

• topiramate oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow pills or food. For migraine, therapeutic failure, intolerance, or contraindication to two generic preventive migraine therapies. For partial onset seizures, primary generalized tonic-clonic seizures, or adjunctive treatment of Lennox-Gastaut Syndrome, therapeutic failure, contraindication, or intolerance to two generic anti-epileptic drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ergotamine

Products Affected

• ergotamine-caffeine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use to abort a vascular headache -AND- therapeutic failure or intolerance to a generic triptan -OR- documentation of use to prevent a vascular headache -AND- therapeutic failure or intolerance to generic prophylactic migraine medication
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Erivedge

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if disease is locally advanced all of the following: 1) disease has recurred following surgery, or is not a candidate for surgery, 2) is not a candidate for radiation.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Erleada

Products Affected

• ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member meets one of the following (1 or 2) 1. Documentation of use in combination with a GnRH analog -OR- 2. The member has had a bilateral orchiectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Evenity

Products Affected

• EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML (105MG/1.17MLX2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Postmenopausal woman at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug. A cumulative lifetime approval of romosozumab will be limited to a coverage duration of 12 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Evrysdi

- EVRYSDI ORAL RECON SOLN
- EVRYSDI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of spinal muscular atrophy (SMA) -AND- Baseline motor function test results (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT) -AND- Not using concomitantly with Spinraza (nusinersen) -AND- Molecular genetic testing of 5q SMA showing Homozygous gene deletion, Homozygous conversion mutation or Compound heterozygote -AND- No history of gene replacement therapy to treat of SMA or experienced a decline in clinical status following gene replacement therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	For reauthorization, attestation of stable or clinically significant improvement in SMA-associated symptoms (e.g. stabilization or decreased decline in motor function compared to the predicted natural history trajectory of disease) or stable or improved motor function results compared to baseline (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT) -AND- Not using concomitantly with Spinraza (nusinersen) - AND- No history of gene replacement therapy to treat of SMA or experienced a decline in clinical status following gene replacement therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fabhalta

Products Affected

• FABHALTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For paroxysmal nocturnal hemoglobinuria (PNH), meets one of the following (1 or 2): 1) PNH mutant clones confirmed by flow cytometry, or 2) glycosylphosphatidylinositol-anchored proteins (GPI-AP)-deficient polymorphonuclear cells (PMNs) confirmed by flow cytometry -AND- meets one of the following (3-6): 3) anemia secondary to PNH (e.g. hemoglobin less than 10.5 g/dL with symptoms of anemia), 4) elevated lactate dehydrogenase (LDH) greater than or equal to 1.5 times the upper limit of normal, 5) history of a thromboembolic event, or 6) clinical findings of systemic complications (e.g. fatigue, hemoglobinuria, abdominal pain, dyspnea, dysphagia, erectile dysfunction, history of blood cell transfusion due to PNH) -AND-will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)) unless initially cross-titrating. For diagnosis of primary immunoglobulin A nephrology (IgAN) confirmed by biopsy, member is at risk for rapid disease progression evidenced by one of the following (7 or 8): 7) Urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g or 8) Proteinuria greater than or equal to 1 g/day -AND- has experienced therapeutic failure, contraindication, or intolerance to a maximally tolerated dose of one of the following (a or b): a) Angiotensin converting enzyme (ACE) inhibitor, b) Angiotensin receptor blocker (ARB) -AND- experienced therapeutic failure, contraindication, or intolerance to one of the following (c or d): c) Filspari (sparsentan) or d) Tarpeyo (budesonide). For Complement 3 Glomerulopathy (C3G) confirmed by biopsy, meets all of the following (13-14): 13) UPCR greater than or equal to 1.0 g/g, 14) currently therapy of the member should be on the maximally tolerated dose of one of the following (e or f): e) ACE-inhibitor, f) ARB.
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of PNH, attestation of positive clinical response defined as one of the following (1-3): 1) hemoglobin stabilization or increase from baseline, 2) decrease in transfusions from baseline, or 3) decrease in LDH levels from baseline or reduction of hemolysis -AND-will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)). For reauthorization of IgAN, reduction in urine protein-to-creatinine ratio (UPCR) or proteinuria from baseline. For reauthorization of C3G, reduction in UPCR from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fanapt

- FANAPT
- FANAPT TITRATION PACK A

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fasenra

- FASENRA PEN
- FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe asthma and all of the following (1-4): 1) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos 2) blood eosinophils greater than or equal to 150cells/uL within the past 6 weeks or greater than or equal to 300cells/uL within the past 12 mos in without other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) 3) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), with or without oral corticosteroids (OCS), unless intolerant or contraindicated to all 4) will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, with or without OCS -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) and all of the following (5-6): 5) history of relapsing or refractory disease 6) will be receiving standard of care while on Fasenra therapy with glucocorticoid treatment (e.g. prednisone or prednisolone), with or without immunosuppressive therapy (e.g. cyclosporine, leflunomide, azathioprine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For severe asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EPGA reauthorization, attestation of one of the following is required (5-8): 5) reduction in the frequency and/or severity of relapses, 6) reduction or discontinuation of doses of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fecal Microbiota Products

Products Affected

· vowst

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a recent diagnosis of recurrent Clostridioides difficile infection (CDI) -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	For reauthorization, attestation of recurrent CDI episodes after administration of the initial fecal microbiota product -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ferriprox

Products Affected

• deferiprone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of transfusional iron overload due to thalassemia syndromes, sickle cell disease and other anemias -AND- transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -AND- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Initial authorization requires trial and failure of deferasirox (generic Exjade) and reauthorization requires failure of generic deferasirox (generic Exjade) if not previously trialed. For reauthorization, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Fetzima

Products Affected

• FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26) CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG

• FETZIMA ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder and trial and failure of two other generic antidepressants.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Filsuvez

Products Affected

• FILSUVEZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB) -AND- at least one open wound.
Age Restrictions	Deny if less than 6 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement in targeted wound(s) - AND- member requires additional courses of treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fintepla

Products Affected

• FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravet syndrome or Lennox-Gastaut syndrome. For Lennox-Gastaut syndrome, therapeutic failure, contraindication, or intolerance to at least 2 standard of care treatments (e.g. lamotrigine, clobazam).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Firazyr

- icatibant
- SAJAZIR

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Firdapse

Products Affected

• FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Firmagon

Products Affected

• FIRMAGON KIT W DILUENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Forteo

Products Affected

• teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of teriparatide will be limited to a coverage duration of 24 months in the absence of provider attestation that the member remains at or has returned to having a high risk for fracture
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fotivda

Products Affected

• FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received at least two prior systemic therapies.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Fruzaqla

Products Affected

• FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received previous threatment with a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy -AND- member has received previous treatment with an anti-VEGF therapy -AND- one of the following, if member is RAS wild-type (1-2): 1) member has received previous therapy with an anti-EGFR therapy 2) prescriber attests that treatment with an anti-EGFR therapy would not be medically appropriate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Furoscix

Products Affected

• FUROSCIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- medication regimen includes an oral loop diuretic (e.g. furosemide, bumetanide, torsemide) -AND- treatment with oral diuretics will replace the use of Furoscix as soon as practical.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gabapentin

- gabapentin oral capsule 100 mg, 300 mg, 400 mg
- gabapentin oral solution 250 mg/5 ml
- gabapentin oral tablet 600 mg, 800 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drugdrug interaction adverse events for opiate potentiators
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gattex

Products Affected

• GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of short bowel syndrome (SBS) having less than 200 cm of functional small bowel -AND- Dependence on parenteral/intravenous nutrition -AND- weight of at least 10 kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gavreto

Products Affected

• GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, disease is RET fusion-positive as detected by an FDA approved test. For advanced or metastatic thyroid cancer, all of the following (1-2): 1) disease is RET fusion-positive 2) if radioactive iodine is appropriate, the member is radioactive iodine-refractory.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gilenya

Products Affected

• fingolimod

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Glatiramer

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml
- GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gleevec

Products Affected

• imatinib oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult agressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvent treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Gleostine

Products Affected

• GLEOSTINE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or metastatic brain tumor(s) -AND- member has previously received surgical and/or radiotherapeutic procedure(s). Documentation of Hodgkin's lymphoma -AND- all of the following (1-2): 1) using lomustine in combination with other chemotherapies 2) member has experienced disease progression with initial chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLP1

Products Affected

• MOUNJARO MG/DOSE (8 MG/3 ML)

OZEMPIC SUBCUTANEOUS PEN
 RYBELSUS
 INJECTOR 0.25 MG OR 0.5 MG (2 MG/3
 TRULICITY
 ML), 1 MG/DOSE (4 MG/3 ML), 2

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Gomekli

- GOMEKLI ORAL CAPSULE 1 MG, 2 MG
- GOMEKLI ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) -AND- presence of symptomatic plexiform neurofibromas (PN) not amenable to complete resection.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Growth Hormone

Products Affected

NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, bone age, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria. Additionally for growth failure due to chronic kidney disease, glomerular filtration rate is less than 89ml/min per 1.73m*2. For HIV wasting and cachexia, Concurrent use of antiretroviral therapy -AND-weight loss of at least 10 percent from baseline. For short bowel syndrome, receiving management for short bowel syndrome, including specialized nutritional support -AND- less than 200 cm of functional small bowel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines. For reauthorization of HIV wasting and cachexia, attestation of increase in weight from start of therapy. For reauthorization of short bowel syndrome, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Hernexeos

Products Affected

• HERNEXEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For non-squamous non-small cell lung cancer (NSCLC), all of the following (1-3): 1) disease is unresectable or metastatic, 2) disease harbors HER2 (ERBB2) tyrosine kinase domain activating mutations, as detected by an FDA-approved test, and 3) member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

High-risk meds

- amitriptyline
- benztropine oral
- clomipramine
- doxepin oral capsule
- doxepin oral concentrate

- doxepin oral tablet
- hydroxyzine hcl oral tablet
- imipramine hcl
- promethazine oral
- trimipramine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For all medications subject to this PA group, the following information (1 through 3) is required: 1. Documentation of diagnosis 2. Explanation of risk-benefit profile favoring use of the high-risk medication 3. Attestation of an intent to monitor and address treatment-related adverse events. For target tricyclic antidepressants (TCAs), in addition to criteria 1 through 3 above, trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs for the same indication, if available, is required (e.g. SSRIs and SNRIs). If using a TCA for a medically-accepted indication not shared by the safer alternatives listed, then no trial of alternatives is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Doxepin doses less than or equal to 6 mg per day will receive automatic approval.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

High-risk meds phenobarbital

Products Affected

• phenobarbital

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia
Required Medical Information	For use in seizures the following are required: 1. Explanation of risk-benefit profile favoring use of the high-risk medication 2. Attestation of an intent to monitor and address treatment-related adverse events. 3. For new starts, the trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs used for seizures (e.g. carbamazepine, lamotrigine) is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Seizure disorders
Part B Prerequisite	No

Humira

- CYLTEZO(CF)
- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS SUBCUTANEOUS PEN INJECTOR KIT · HUMIRA(CF) 40 MG/0.8 ML
- CYLTEZO(CF) PEN PSORIASIS-UV SUBCUTANEOUS PEN INJECTOR KIT 40 MG/0.8 ML
- HADLIMA
- HADLIMA PUSHTOUCH
- HADLIMA(CF)
- HADLIMA(CF) PUSHTOUCH

- HUMIRA PEN
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PSOR-UV-ADOL HS
- SIMLANDI(CF)
- SIMLANDI(CF) AUTOINJECTOR
- YUFLYMA(CF)
- · YUFLYMA(CF) AI CROHN'S-UC-HS
- YUFLYMA(CF) AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 1 immunosuppressant or corticosteroid, or all are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	Plan Year
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. Please Note: This criteria is applicable to Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ibrance

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of HR-positive, HER2-negative advanced or metastatic breast cancer -AND- meets one of the following (1. or 2.): 1) documentation of use with an aromatase inhibitor as initial endocrine-based therapy -OR- 2) documentation of use with fulvestrant in patients with disease progression following endocrine therapy. Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND-all of the following (3-6): 3) disease is HR-positive, HER2-negative 4) disease is PIK3CA-mutated, as detected by an FDA-approved test 5) the member is using in combination with inavolisib and fulvestrant 6) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ibsrela

Products Affected

• IBSRELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of irritable bowel syndrome with constipation -AND-failure or intolerance to Linzess -AND- if member is female, failure or intolerance to lubiprostone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ibtrozi

Products Affected

• IBTROZI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Iclusig

Products Affected

· ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of newly-diagnosed chronic phase CML
Required Medical Information	Documentation of T3151+ chronic myeloid leukemia (CML) -OR-documentation of chronic phase CML and member has experienced resistance or intolerance to at least two prior kinase inhibitors -OR-documentation of accelerated phase or blast phase CML and no other kinase inhibitor is indicated -OR- member is using Iclusig as monotherapy and meets one of the following (1-2): 1) documentation of T3151+ acute lymphoblastic leukemia (ALL) 2) documentation of Ph+ ALL and no other tyrosine kinase inhibitor therapy is indicated -OR- member has newly diagnosed Ph+ ALL and is using Iclusig in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Idhifa

Products Affected

• IDHIFA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IG

- BIVIGAM
- GAMMAGARD LIQUID
- GAMMAGARD S-D (IGA
- GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)
- GAMMAPLEX

- GAMMAPLEX (WITH SORBITOL)
- GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)
- OCTAGAMPANZYGA
- PRIVIGEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For select diagnoses the following apply 1) For Myasthenia Gravis Syndrome, documentation that the patient is refractory to other standard therapies (e.g., cholinesterase inhibitors, corticosteroids, azathioprine) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 2) For Multiple Sclerosis, patient is refractory to other standard therapies (e.g., interferons) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 3) For B-cell CLL, associated with recurrent bacterial infections OR with Associated Hypogammaglobulinemia defined as IgG level less than 600mg/dL or evidence of a specific antibody deficiency. 4) For Bone Marrow Transplantation, the member is 20 years of age or older and within the first 100 days after transplantation. 5) For Dermatomyositis/Polymyositis, trial and failure, intolerance, or contraindication to standard fist line therapy (i.e. corticosteroids or immunosuppressants). 6) For Pediatric HIV, the patient is less than 13 y.o. who have entry CD4 lymphocyte count greater than or equal to 200/mcl and IgG less than 400mg/dL OR recurrent bacterial infections. 7) For Guillain-Barre syndrome, impaired function by objective assessment and/or objective findings on physical exam at the time of initial therapy and IVIG therapy must be initiated within 2 weeks of symptom onset. 8) For Autoimmune Mucocutaneous Blistering Diseases (e.g. Stevens-Johnson Syndrome), trial and failure, intolerance, or contraindication to conventional therapy (e.g. corticosteroids) or the patient has rapidly progressive disease in which a clinical response could not be affected quickly enough using conventional agents.
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when administered in the home to a member with a diagnosis of primary immunodeficiency disease
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Myasthenia Gravis syndrome, Multiple Sclerosis, Polymyositis, Bone Marrow Transplant, Pediatric HIV, Guillain-Barre syndrome, Autoimmune Mucocutaneous Blistering Diseases
Part B Prerequisite	No

Imbruvica

- IMBRUVICA ORAL CAPSULE 140 MG, 280 MG, 420 MG 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG,

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic graft versus host disease, previous treatment with at least 1 prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For suspension, inability to swallow oral tablets or oral capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Imkeldi

Products Affected

• IMKELDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult agressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvent treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of an inability to swallow oral tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Increlex

Products Affected

INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe primary IGF-1 deficiency and all of the following: 1) Normal or elevated response (greater than 10 ng/ml) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 2) Serum IGF-1 concentration that is less than or equal to three (3) standard deviations below the normal value based on laboratory reference range. 3) Height less than or equal to three (3) standard deviations below normal (at or below the third percentile for gender and age). 4) Member has open epiphysesOR- Documentation of diagnosis of growth hormone deficiency caused by gene deletion and all of the following: 1) Growth velocity at least 2 standard deviations below the age-appropriate mean or height at least 2.25 standard deviations below the age-appropriate mean. 2) Subnormal response (less than 10 ng/mL) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 3) Development of neutralizing antibodies to growth hormone product(s). 4) Member has open epiphyses
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation member has a growth velocity of at least 2 cm/year -AND- member has open epiphyses.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Inlyta

Products Affected

· INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced renal cell carcinoma -AND- one of the following (1-2): 1) if using axitinib as first line therapy, member is using axitinib in combination with avelumab or pembrolizumab 2) if using axitinib as a single-agent, member has been treated with at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inqovi

Products Affected

· INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation chronic myelomonocytic leukemia. Documentation of myelodysplastic syndrome -AND- One of the following (1 or 2): 1) French American-British MDS subtypes of refractory anemia, refractory anemia with ringed sideroblasts or refractory anemia with excess blasts. 2) International Prognostic Scoring System group of intermediate-1, intermediate-2 or high-risk.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Inrebic

Products Affected

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate-2 or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) -AND- If a new start, baseline platelet count of greater than or equal to $50x10*9/L$
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Insulin Supplies

Products Affected

- · ALCOHOL PADS
- GAUZE PAD TOPICAL BANDAGE 2 X 2 "

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation the product is being used for the delivery of insulin into the body.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Interferon Alfa

Products Affected

• PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Interferon Beta

Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT
- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IPF Agents

Products Affected

- OFEV
- pirfenidone oral capsule pirfenidone oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 50% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30%. For Ofev only, documentation of systemic sclerosis-associated interstitial lung disease -AND- baseline forced vital capacity (FVC) of at least 40% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- documentation of a chest computed tomography (CT) scan demonstrating greater than or equal to 10% pulmonary fibrosis. For Ofev only, documentation of chronic fibrosing interstitial lung disease with progressive phenotype -AND- chest computing tomography (CT) scan demonstrating greater than 10% fibrosing disease -AND- baseline forced vital capacity (FVC) of at least 45% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- disease progression in previous 24 months shown by one of the following: 1. Relative decline in FVC greater than or equal to 5% but less than 10% predicted and either worsening of respiratory symptoms or increased extent of fibrotic changes on CT 3. Worsening of respiratory symptoms and increasing extent of fibrotic changes on HRCT
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

Iressa

Products Affected

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	Use in tumors with EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.
Required Medical Information	Documentation of diagnosis -AND- the following: 1) EGFR exon 19 deletion mutations or exon 21 (L858R) mutations as detected by an FDA-approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Itovebi

Products Affected

• ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (1-4): 1) disease is HR-positive, HER2-negative 2) disease is PIK3CA-mutated, as detected by an FDA-approved test 3) the member is using in combination with palbociclib and fulvestrant 4) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Itraconazole

Products Affected

• itraconazole oral capsule

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If using for diagnosis of onychomycosis, confirmation through positive laboratory testing (e.g. KOH preparation, fungal culture, or nail biopsy) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Onychomycosis: 3 months. All other indications: 3 months initial, 12 months reauth
Other Criteria	Documentation of trial/failure or intolerance of amphotericin b must be provided for approval in patients with aspergillosis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ivermectin Oral

Products Affected

• ivermectin oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of strongyloidiasis of the intestinal tract (non-disseminated disease) or onchocerciasis -AND- Member weighs greater than or equal to 15kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Iwilfin

Products Affected

• IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of high-risk neuroblastoma (HRNB) -AND- partial response to anti-glycolipid disialoganglioside (GD2) immunotherapy (e.g., dinutuximab, naxitamab).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jakafi

Products Affected

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) and if a new start, baseline platelet count of greater than or equal to $50x10*9/L$ -OR- documentation of polycythemia vera and inadequate response or intolerance to hydroxyurea -OR-Documentation of steroid refractory acute graft-versus-host disease and prior therapy with at least one systemic corticosteroid -OR-Documentation of chronic graft-versus-host disease with prior failure of at least one systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Jaypirca

Products Affected

• JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory mantle cell lymphoma, member has received at least two (2) previous lines of systemic therapy, at least one (1) of which was a BTK inhibitor. For Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia, member has received at least two (2) prior lines of therapy, including at least one (1) from all of the following classes (1-2): 1) BTK inhibitor 2) BCL-2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Joenja

Products Affected

· JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of activated phosphoinositide 3-kinase delta syndrome (APDS) with genetic confirmation of variant in PIK3CD or PIK3R1 gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kalydeco

Products Affected

- KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG
- KALYDECO ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to ivacaftor based on clinical and or in vitro assay (e.g. G551D, G1244E, G1349D)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kerendia

Products Affected

KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than 25 mL/min/1.73 m*2, serum potassium greater than 5.5 mEq/L
Required Medical Information	Pending CMS Review
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For T2DM+CKD reauthorization, attestation that signs or symptoms of hyperkalemia are not present. For HF reauthorization, attestation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kesimpta

Products Affected

• KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kevzara

Products Affected

KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For polymyalgia rheumatica (PMR), one of the following (1-3): 1) inadequate response to corticosteroids, 2) intolerance to corticosteroid taper, or 3) used in combination with corticosteroid tapering course. For juvenile idiopathic arthritis, weight greater than or equal to 63 kg -AND- one of the following (4-5): 4) inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide), or 5) requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis and polymyalgia rheumatica.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Rinvoz/Rinvoq LQ, Xeljanx/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Kineret

Products Affected

KINERET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), therapeutic failure or intolerance to at least one (1) corticosteroid, or all corticosteroids are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kisqali

Products Affected

- KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
- KISQALI ORAL TABLET 200 MG/DAY

(200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative -AND- one of the following (1-3): 1) member is using ribociclib in combination with an aromatase inhibitor as initial endocrine-based therapy 2) member is using ribociclib in combination with fulvestrant and member is using fulvestrant as initial endocrine-based therapy or member has experienced disease progression on endocrine therapy. 3) disease is classified as stage II or stage III early breast cancer at high risk of recurrence, ribociclib is being used in combination with an aromatase inhibitor as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Korlym

Products Affected

• mifepristone oral tablet 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperglycemia secondary to hypercortisolism in patients with endogenous Cushing's syndrome -AND- Patient is not a candidate for surgery or where surgery has failed -AND- one of the following (1 or 2): 1) Diagnosis of diabetes with trial and failure, intolerance, or contraindication to one previous therapy for Type 2 Diabetes (e.g. metformin, sulfonylureas, insulin) or using in addition to a therapy for Type 2 diabetes. 2) Glucose intolerance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Koselugo

Products Affected

• KOSELUGO ORAL CAPSULE 10 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- for neurofibromatosis type 1 (NF1), documentation of symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	Deny if less than 1 years of age or older than 17 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Krazati

Products Affected

• KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of locally advanced or metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA- approved test -AND- using in combination with cetuximab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Kuvan

Products Affected

- JAVYGTOR
- sapropterin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of PKU -AND- documented baseline Phe level greater than 6 mg/dL -AND- clinical documentation of current weight -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, initial therapy has resulted in decrease in phenylalanine levels from baseline or current phenylalanine levels within the range of 120-360 micromol/L -AND- clinical documentation of current weight is required -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Latuda

Products Affected

• lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for bipolar 1 disorder, documentation of trial and failure or intolerance to one other formulary medication indicated in bipolar 1 disorder (e.g. quetiapine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lazcluze

Products Affected

• LAZCLUZE ORAL TABLET 240 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) -AND- all of the following (1-4): 1) disease harbors EGFR exon 19 deletions or EGFR exon 21 (L858R) substitution mutation, as detected by an FDA-approved test 2) disease is locally advanced or metastatic, 3) member is treatment naive for advanced disease, 4) using in combination with amivantinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lenvima

Products Affected

• LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of differentiated thyroid cancer -AND- meets all of the following (1-3): 1) disease is locally recurrent or metastatic 2) disease is progressive 3) disease is radioactive iodine refractory. Documentation of advanced renal cell carcinoma -AND- one of the following (4-5): 4) member is using lenvatinib in combination with pembrolizumab and is using lenvatinib and pembrolizumab as first-line treatment 5) member is using lenvatinib in combination with everolimus and has experienced therapeutic failure or intolerance to one prior anti-angiogenic therapy. Documentation of unresectable hepatocellular carcinoma -AND- member is using lenvatinib as first-line treatment. Documentation of endometrial cancer -AND- meets all of the following (6-10): 6) member is using lenvatinib in combination with pembrolizumab 7) disease is advanced 8) disease is not classified as microsatellite instability-high or disease is classified as mismatch repair proficient as determined by an FDA-approved test 9) member has experienced disease progression following prior systemic therapy 10) member is not a candidate for curative surgery or radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Leukine

Products Affected

• LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis: following induction chemotherapy in patients who are 55 years or older with acute myelogenous leukemia (AML) -OR- mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation -OR- acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation -OR- acceleration of myeloid reconstitution following allogeneic BMT -OR- treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic BMT -OR- following acute exposure to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lidoderm

Products Affected

• lidocaine topical adhesive patch,medicated 5 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) or diabetic peripheral neuropathy (DPN) -AND- One of the following (1-3): 1) trial and failure of 1 other agent used to treat diagnosis (e.g. gabapentin for PHN, duloxetine for DPN), 2) inability to swallow oral medication, 3) unable to take an oral medication due to potential adverse events (e.g. sedation).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Diabetic peripheral neuropathy
Part B Prerequisite	No

Litfulo

Products Affected

LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For severe alopecia areata, therapeutic failure/intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Livtencity

Products Affected

• LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of CMV prophylaxis.
Required Medical Information	Documentation of refractory cytomegalovirus infection or disease as evidenced by antigenemia or polymerase chain reaction (PCR) test -AND-all of the following (1-3): 1) member weighs at least 35 kg, 2) member is a recipient of hematopoietic stem cell transplant -OR- solid organ transplant. 3) member has experienced therapeutic failure to one of the following: ganciclovir, valganciclovir, cidofovir, or foscarnet.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of a previous reduction in CMV DNA level -AND- documentation of one of the following (1-3): 1) new onset symptomatic CMV infection, 2) virologic relapse with treatment-emergent maribavir resistance or 3) continued antiviral treatment is required to achieve virologic clearance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lokelma

Products Affected

LOKELMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 7.4 mmol/L on at least two (2) screenings -AND-Modification of medications to reduce serum potassium levels were not successful, when applicable
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in serum potassium levels following Lokelma administration and continued treatment for hyperkalemia is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lonsurf

Products Affected

• LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer as a single agent or in combination with bevacizumab in patients who have previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy -OR- documentation of metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/neutargeted therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lorbrena

Products Affected

• LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ALK-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lotronex

Products Affected

• alosetron oral tablet 0.5 mg, 1 mg

PA Criteria	Criteria Details
Exclusion Criteria	For irritable bowel syndrome (IBS): Exclude if male gender
Required Medical Information	Documentation of chronic severe diarrhea-predominant IBS -AND- trial and failure or intolerance to one anti-diarrheal (e.g. loperamide), antispasmodic, or tricyclic antidepressant, or contraindication to all
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that symptoms of IBS continue to persist AND positive clinical response.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lumakras

Products Affected

 LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- using in combination with panitumumab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lupron Depot Ped

Products Affected

- LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG
- · LUPRON DEPOT-PED

INTRAMUSCULAR KIT 7.5 MG (PED)
 LUPRON DEPOT-PED
 INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of central precocious puberty -AND- advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3 IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy defined as pre-pubertal slowing/decline, normalization of FSH, normalization LH, normalization of bone age, normalization of bone age, normalization of estradiol level or normalization of testosterone level
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lynparza

Products Affected

· LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) in complete or partial response to first-line platinum-based chemotherapy 2) diagnosis of deleterious or suspected deleterious BRCA mutated disease or disease is associated with homologous recombination deficiency (HRD) positive status with a deleterious or suspected deleterious BRCA mutation or genomic instability and will be using in combination with bevacizumab. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) deleterious or suspected deleterious germline or somatic BRCA mutation 2) in complete or partial response to platinum-based chemotherapy. For deleterious or suspected deleterious gBRCAm, HER2-negative breast cancer, 1 of the following (1-2): 1) classified as high-risk, early breast cancer and has been treated with neoadjuvant or adjuvant chemotherapy 2) has been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting and if hormone receptor (HR)-positive, has been previously treated with or considered inappropriate for treatment with endocrine therapy. For metastatic pancreatic adenocarcinoma, all of the following (1-2): 1) a deleterious or suspected deleterious gBRCA mutation 2) did not progress on at least 16 weeks of a first-line platinum-based chemotherapy regimen. For deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene mutation metastatic castration-resistant prostate cancer, all of the following (1-2): 1) progressed following prior treatment with enzalutamide or abiraterone 2) concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy. For deleterious or suspected deleterious BRCA mutation metastatic castration-resistant prostate cancer, using in combination with all of the following (1-2): 1) abiraterone 2) prednisone or prednisolone.
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lyrica

- pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg
- pregabalin oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of DPN and trial/failure or intolerance to duloxetine -OR-PHN and trial/failure or intolerance to gabapentin -OR- Partial-onset seizures and trial/failure or intolerance to two AEDS -OR- Neuropathic pain associated with spinal cord injury -OR- Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine. When using pregabalin products concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Lytgobi

Products Affected

• LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-2): 1) disease harbors FGFR2 fusions or other rearrangements 2) member has experienced therapeutic failure or intolerance to at least one prior therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mavyret

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Megace

- megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)
- megestrol oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mekinist

- MEKINIST ORAL RECON SOLN
- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For trametinib monotherapy, documentation of unresectable or metastatic melanoma -AND- meets all of the following (1-2): 1) member has a BRAF V600E or V600K mutation 2) member is BRAF inhibitor treatment naive. For use in combination with dabrafenib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with dabrafenib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA approved test when FDA indicated. For use in combination with dabrafenib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Mekinist powder for oral solution, attestation of inability to swallow Mekinist (trametinib) tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mektovi

Products Affected

MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, all of the following (1-2): 1) BRAF V600E or V600K mutation status 2) using in combination with encorafenib. For metastatic non-small cell lung cancer, all of the following (1-2): 1) BRAF V600E mutation status 2) using in combination with encorafenib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Metyrosine

Products Affected

• metyrosine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pheochromocytoma defined by 1 of the following (1 or 2): 1) Elevated metanephrines in plasma or urine, or 2) Tumor evidence from CT scan or MRI -AND- Documentation of 1 of the following (3., 4., or 5.): 3) Planned resection surgery, 4) Resection surgery is contraindicated, or 5) malignant pheochromocytomaAND- therapeutic failure, contraindication, or intolerance to a selective alpha-blocker (e.g., doxazosin, prazosin, terazosin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Modeyso

Products Affected

· MODEYSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of diffuse midline glioma -AND- disease harbors an H3 K27M mutation -AND- member has experienced progressive disease following prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Mulpleta

Products Affected

• MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND-beneficiary is scheduled to undergo a procedure -AND- trial and failure or intolerance to Doptelet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Myasthenia Gravis

Products Affected

• ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML, 23 MG/0.574 ML, 32.4 MG/0.81 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized myasthenia gravis (gMG) -AND- Antiacetylcholine receptor (AChR) antibody-positive -AND- Therapeutic failure, contraindication, or intolerance to generic pyridostigmine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of gMG signs/symptoms improvement (e.g., speech, swallowing, mobility, and/or respiratory function) -OR-decreased gMG exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Namzaric

- memantine-donepezil
- NAMZARIC ORAL CAPSULE,SPRINKLE,ER 24HR 7-10 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and intolerance to generic memantine and generic donepezil
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nayzilam

Products Affected

NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nemluvio

Products Affected

· NEMLUVIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of prurigo nodularis -OR- Documentation of moderate to severe atopic dermatitis and one of the following (1-3): 1) trial and failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus), 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement, or 3) severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For prurigo nodularis, must follow recommended dosing guidelines based upon weight. For atopic dermatitis, must follow recommended dosing based on FDA approved dosing guidelines. For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nerlynx

Products Affected

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of early-stage, HER2-positive breast cancer -AND- meets all of the following: 1)member has received adjuvant trastuzumab-based therapy 2) member is using neratinib as a single agent. Documentation of advanced HER-2 positive, or metastatic HER2-positive breast cancer and meets all of the following 1) using neratinib in combination with capecitabine 2) member has received two or more prior anti-HER2 based regimens in the metastatic setting.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nexavar

Products Affected

• sorafenib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally recurrent or metastatic, progressive, differentiated thyroid carcinoma, refractory to radioactive iodine treatment
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nexletol

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For HeFH, diagnosis supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite Simon Broome register criteria, or definite on the Make Early Diagnosis to Prevent Early Deaths tool -AND- LDL-C greater than 100 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND-Therapeutic failure, intolerance or contraindication to ezetimibe. For Hyperlipidemia with ASCVD or Hyperlipidemia with attestation of high risk for CVD, LDL-C greater than 70 mg/dL -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND-Therapeutic failure, intolerance or contraindication to ezetimibe. For Primary Hyperlipidemia not associated with ASCVD or HeFH, LDL-C greater than 70 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For reauthorization, documentation showing an LDL-C reduction from baseline -AND- for HeFH and Primary Hyperlipidemia (not associated with ASCVD or HeFH), attestation of continued use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statin which resolved upon discontinuation of statin or documentation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ninlaro

Products Affected

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- previous treatment with at least 1 prior therapy -AND- used in combination with lenalidomide and dexamethasone
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nitisinone

Products Affected

• nitisinone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary tyrosinemia type 1 (HT-1) confirmed by biochemical or genetic testing -AND- dietary restriction of tyrosine and phenylalanine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Northera

Products Affected

• droxidopa oral capsule 100 mg, 200 mg, 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of neurogenic orthostatic hypotension caused by primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency or non-diabetic autonomic neuropathy -AND- documentation of inadequate response, intolerance or contraindication to preferred generic alternative midodrine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of increase from baseline of systolic or diastolic blood pressure upon standing -OR- attestation of decrease from baseline of neurogenic orthostatic hypotension symptoms upon standing (e.g., dizziness, feeling faint, etc.).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nubeqa

Products Affected

• NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-metastatic castration-resistant prostate cancer - AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy. Documentation of metastatic castration-sensitive prostate cancer -AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nucala

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Severe asthma: history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12mos or at least 1 asthma exacerbation requiring hospitalization in past 12mos -AND- blood eosinophils of at least 150cells/uL within past 6wks or at least 300cells/uL within past 12mos w/o other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) -AND- inadequate symptom control despite regular treatment w/ medium or high dose inhaled corticosteroids (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), w/ or w/o oral corticosteroids (OCS) -AND- will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, w/ or w/o OCS. Eosinophilic granulomatosis with polyangiitis (EGPA): history of relapsing or refractory disease -AND- will be receiving concomitant glucocorticoid treatment w/ or w/o immunosuppressive therapy. Hypereosinophilic syndrome (HES) w/o identifiable non-hematologic secondary cause for at least 6mos: at least 2 HES flares (HES-related worsening of clinical symptoms or blood eosinophil counts requiring an escalation in therapy) within past 12mos - AND- Stable on HES therapy for at least 4wks (chronic or episodic OCS, immunosuppressive or cytotoxic therapy). Chronic rhinosinusitis with nasal polyps (CRSwNP): trial/failure, contraindication, or intolerance to intranasal corticosteroid. Chronic obstructive pulmonary disease: blood eosinophils of at least 300cells/uL or current daily or alternate-day OCS therapy -AND- inadequate symptom control despite regular treatment for at least 3 months with LAMA, LABA, and ICS, unless intolerant or contraindicated to all.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	For Severe Asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EGPA reauthorization, attestation of one of the following is required (5-8): 5) reduction in frequency and/or severity of relapses, 6) reduction or discontinuation of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms. For HES reauthorization, attestation of one of the following is required (9-10): 9) reduction in frequency of HES flares, or 10) maintenance or reduction in background HES therapy requirements. For CRSwNP reauthorization, attestation of one of the following is required (11-12): 11) decrease in nasal polyp score, or 12) reduction in nasal congestion/obstruction severity score. For COPD reauthorization, attestation of one of the following is required (13-16): 13) reduction in COPD symptoms, 14) improvement in exercise tolerance, 15) delayed disease progression, or 16) reduction in the number of COPD exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuedexta

Products Affected

NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting improvement in symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuplazid

Products Affected

NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hallucinations and delusions associated with Parkinson's disease psychosis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nurtec

Products Affected

NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine Prevention, defined as 4-14 migraine days per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For acute treatment of migraine with or without aura, trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency. For reauthorization of acute treatment of migraine, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Nuvigil

Products Affected

• armodafinil

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 173 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Octreotide

Products Affected

• octreotide acetate injection solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin- like growth factor-1 (IGF-1) based on laboratory reference range -AND- therapeutic failure or cannot be treated with surgical resection, pituitary irradiation or bromocriptine mesylate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Odomzo

Products Affected

· ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced basal cell carcinoma (laBCC) that has recurred following surgery or radiation therapy or for use in patients who are not candidates for surgery or radiation therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ogsiveo

Products Affected

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressing desmoid tumor(s) requiring systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ojemda

Products Affected

- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION
- OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500

MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory pediatric low-grade glioma - AND- one of the following (1-2): 1) BRAF fusion or rearrangement, or 2) BRAF V600K mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ojjaara

Products Affected

· OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND-attestation of anemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Olumiant

Products Affected

· OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis an inadequate response or intolerance to at least one non-biologic DMARD (e.g., methotrexate, leflunomide). For severe alopecia areata, inadequate response or intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Alopecia Areata
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR, with at least 1 being a tumor necrosis factor blocker. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Onfi

Products Affected

- clobazam oral suspension clobazam oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND-documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Onureg

Products Affected

· ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute myeloid leukemia that has achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy -AND- Inability to complete intensive curative therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Opipza

Products Affected

· OPIPZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- 1 of the following (1-2): 1) intolerance to generic aripiprazole tablets, 2) inability to swallow oral tablets. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orencia

Products Affected

- ORENCIA CLICKJECT
- ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients over 18 years of age must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Xeljanz/Xeljanz XR, Otezla and Stelara SC, Rinvoq, and Skyrizi SC. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Xeljanx/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Orgovyx

Products Affected

ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orkambi

Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis and homozygous F508del mutation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Orserdu

Products Affected

• ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-4): 1) member is male or a postmenopausal female 2) tumor status is ER-positive, HER2-negative 3) an ESR1 gene mutation is present in the tumor 4) member has experienced disease progression on or after an endocrine based regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OTEZLA

Products Affected

• OTEZLA (47)

 OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For plaque psoriasis, all of the following (1-2): 1) if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg and member has moderate-to-severe disease -AND- 2) inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For oral ulcers associated with Behcet's Disease, inadequate response or intolerance to one systemic therapy for prevention of recurrent oral ulcers. For psoriatic arthritis, if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg.
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Palynziq

Products Affected

• PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of phenylketonuria -AND- all of the following criteria (1-3): 1) Baseline Phe level greater than 600 micromoles/L, 2) Failure or intolerance to existing management (i.e. sapropterin dihydrochloride), 3) Has a prescription for epinephrine agent unless contraindicated.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in baseline pretreatment Phe levels -OR- blood Phe levels are less than or equal to 600 micromoles/L - OR- attestation that additional therapy with Palynziq is needed to allow adequate trial of maximum dose of 60mg per day for 16 weeks or member is not currently on 60mg per day dose.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Panretin

Products Affected

• PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions in patients with AIDS-related Kaposi Sarcoma (KS) who are not receiving systemic therapy for KS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pemazyre

Products Affected

• PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable locally advanced cholangiocarcinoma or metastatic cholangiocarcinoma -AND- all of the following (1-2): 1) disease harbors FGFR2 fusion or other rearrangement as detected by an FDA-approved test 2) member has experienced therapeutic failure or intolerance to at least on prior therapy. Documentation of relapsed or refractory myeloid/lymphoid neoplasms -AND- disease harbors an FGFR1 rearrangement.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pheburane

Products Affected

• PHEBURANE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure or intolerence to generic sodium phenylbutyrate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Piqray

Products Affected

 PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR-positive, HER2-negative tumor status, 2) PIK3CA mutation positive as detected by an FDA-approved test, 3) disease progression on or after an endocrine-based regimen, 4) Concomitant therapy with fulvestrant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pomalyst

Products Affected

· POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma, and combination use with dexamethasone, and previous trial of at least 2 therapies including lenalidomide and a proteasome inhibitor, and disease progression on or within 60 days of completion of the last therapy -OR- Documentation of AIDS-related Kaposi sarcoma (KS) after failure of highly active antiretroviral therapy (HAART) or in patients with KS who are HIV negative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Posaconazole Tablet

Products Affected

• posaconazole oral tablet,delayed release (dr/ec)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight greater than 40 kg. For invasive aspergillosis infection, trial/failure or contraindication to voriconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prenatal Vitamins

Products Affected

• PRENATAL VITAMIN PLUS LOW IRON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of nutritional supplementation required in a female of child-bearing potential during pre-conception, pregnancy, or lactation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prescription Drug Combo

Products Affected

- acetaminophen-codeine oral solution 120-12 mg/5 ml
- acetaminophen-codeine oral tablet
- alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg
- buprenorphine
- clonazepam oral tablet 0.5 mg, 1 mg, 2 mg
- clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg
- clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg
- DIAZEPAM INTENSOL
- diazepam oral solution 5 mg/5 ml (1 mg/ml)
- diazepam oral tablet
- ENDOCET
- eszopiclone
- fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr
- hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg
- hydromorphone oral liquid
- hydromorphone oral tablet 2 mg, 4 mg, 8 mg

- LORAZEPAM INTENSOL
- lorazepam oral tablet 0.5 mg, 1 mg, 2 mg
- methadone oral solution 10 mg/5 ml, 5 mg/5 ml
- methadone oral tablet 10 mg, 5 mg
- morphine concentrate oral solution
- morphine oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)
- morphine oral tablet
- morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg
- oxycodone oral capsule
- oxycodone oral concentrate
- oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg
- oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg
- tramadol oral tablet 50 mg
- tramadol-acetaminophen
- zaleplon oral capsule 10 mg, 5 mg
- zolpidem oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	For concomitant use of an opiate agonist and substance abuse therapy, documentation that the member has an acute pain condition (e.g. acute traumatic injury) in which treatment with other agents would cause insufficient pain control or if the member requires treatment for pain related to a terminal illness. For concomitant use of an opiate agonist, benzodiazepine, and a centrally acting skeletal muscle relaxant, documentation that the member has tried/failed at least 2 other skeletal muscle relaxants (e.g. methocarbamol, metaxalone), understanding these skeletal muscle relaxants are high-risk medications in geriatric patients, attestations that non-opiate therapies (e.g. NSAIDs) and non-benzodiazepine therapies (e.g. SSRI, SNRI) have been considered, AND attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For concomitant use of an opiate agonist and other opiate potentiators (e.g. gabapentinoids, sedative-hypnotics) attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For long acting (e.g. extended release) opioid medications, the following apply (1-5). 1)Pain is severe enough to require daily, around-the-clock, long-term opioid treatment. 2)Patient is not opioid naive. 3)Attestation that non-opiate alternative therapies have been explored (e.g. NSAIDs). 4)Attestation that controlled substance Rx history has been reviewed in the state Prescription Drug Monitoring Program. 5)Attestation of counseling on the potential adverse effects of opioid analgesics, including the risk of misuse, abuse, and addiction.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Opiate tx for pain+subs. abuse, approve opiate x 1mo. All other combos and dx approve x 12mo.
Other Criteria	Opiate agonists will receive automatic approval if no recent claims for a substance abuse therapy (e.g. buprenorphine-naloxone) OR a benzodiazepine with a centrally acting skeletal muscle relaxant (e.g., carisoprodol) OR a gabapentinoid OR a sedative-hypnotic. Benzodiazepines (e.g. triazolam, alprazolam) will receive automatic approval if no recent claims for an opiate agonist with a centrally acting skeletal muscle relaxant (e.g. carisoprodol). Sedative-hypnotics (e.g. zolpidem) will receive automatic approval if no recent claims for an opiate agonist.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Prevymis Oral Pellets

Products Affected

• PREVYMIS ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 6 months of age with HSCT. Deny if less than 12 years of age with kidney transplant.
Prescriber Restrictions	
Coverage Duration	7 months
Other Criteria	One of the following is required (1-2): 1) inability to swallow tablets, 2) unable to use Prevymis (letermovir) tablets due to body weight dosing limitations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Prolia

Products Affected

· PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -And- For osteoporosis at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 (i.e. osteopenia) -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For osteoporosis and osteopenia, documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to postmenopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Provigil

Products Affected

• modafinil

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 17/3 of the normal value based on laboratory reference range -OR-2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR-3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications. Diagnosis of fatigue associated with Multiple Sclerosis (MS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Fatigue associated with Multiple Sclerosis (MS)
Part B Prerequisite	No

Pulmonary Arterial Hypertension

Products Affected

- ADEMPAS
- ALYQ
- ambrisentan
- · OPSUMIT
- · OPSYNVI
- sildenafil (pulm.hypertension) oral tablet
- tadalafil (pulm. hypertension)
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group. For sildenafil in pediatric individuals, an exception to RHC will be allowed when the risk of RHC outweighs the benefit -AND- prescriber attests alternative studies have been completed (i.e. CT, MRI or specified test ruling out other causes of pulmonary hypertension). For Adempas, additional diagnosis of CTEPH as documented by right heart catheterization and V/Q scan substantiating mPAP greater than 20 mmHg at rest and PWP less than or equal to 15 mmHg and documented presence of occlusive thrombi within the pulmonary arteries will be approved.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Pulmozyme

Products Affected

PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis -AND- Used in conjunction with standard therapies for management of cystic fibrosis to improve pulmonary function.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization, attestation of increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Qinlock

Products Affected

· QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced gastrointestinal stromal tumor -AND- Prior treatment with imatinib and 2 other kinase inhibitors.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Quinine

Products Affected

• quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	Treatment or prevention of leg cramps
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	Doses for duration greater than 10 days will not be approved
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Qulipta

Products Affected

• QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, of which 8 or more are migraine days. The following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Radicava ORS

Products Affected

• RADICAVA ORS STARTER KIT SUSP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of amyotrophic lateral sclerosis (ALS) -AND- Therapeutic failure, intolerance, or contraindication to riluzole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of experiencing stability or improvement of symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ravicti

Products Affected

• RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	Urea cycle disorders due to N-acetylglutamate synthetase deficiency, Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of a urea cycle disorders (UCDs)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Repatha

Products Affected

- REPATHA SURECLICK
- REPATHA SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
Required Medical Information	Pending CMS Review
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation showing an LDL-C reduction on Repatha therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Retevmo

Products Affected

• RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is classified as RET gene fusion as detected by an FDA approved test. Documentation of advanced or metastatic medullary thyroid cancer -AND- disease is classified as RET mutation as detected by an FDA approved test. Documentation of advanced or metastatic thyroid cancer -AND- all of the following (1-2): 1) disease is classified as RET gene fusion as detected by an FDA approved test 2) if radioactive iodine is appropriate for the member, the member is radioactive iodine-refractory. Documentation of locally advanced or metastatic solid tumor(s) -AND-disease harbors a RET gene fusion, as detected by an FDA-approved test -AND- one of the following (1-2): 1) the member has no satisfactory alternative treatments 2) the member's tumors have progressed following prior systemic treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Revlimid

Products Affected

• lenalidomide

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of chronic lymphocytic leukemia outside of a controlled clinical trial
Required Medical Information	Diagnosis of multiple myeloma in combination with dexamethasone -OR-diagnosis of multiple myeloma, as maintenance following autologous hematopoietic stem cell transplant (auto-HSCT) -OR- diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities -OR-diagnosis of mantle cell lymphoma (MCL) in which disease has relapsed or progressed after two prior therapies, one of which included bortezomib -OR- diagnosis of follicular lymphoma in combination with a rituximab product after previous treatment -OR- diagnosis of marginal zone lymphoma in combination with a rituximab product after previous treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Revuforj

Products Affected

 REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute leukemia with a lysine methyltransferase 2A gene (KMT2A) translocation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rexulti

Products Affected

• REXULTI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI). For schizophrenia, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rezdiffra

Products Affected

REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in conjunction with diet and exercise. Diagnosis of NASH confirmed by a liver biopsy or non-invasive tests (NITs) performed within the previous 6 months indicating F2 or F3 fibrosis - AND- member does not have any evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of continued use in conjunction with diet and exercise -AND- member has experienced stabilization of fibrosis as demonstrated by NIT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rezlidhia

Products Affected

REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rinvoq

Products Affected

• RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate) or all non-biologic DMARDs are contraindicated. For moderate to severe refractory atopic dermatitis whose disease is not adequately controlled with other systemic drug products, documentation of one of the following (1 or 2): 1) trial & failure, or intolerance to at least one topical corticosteroid -OR- topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) 2) The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For Non-radiographic Axial Spondyloarthritis, trial & failure or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or contraindication to all. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For giant cell arteritis, patients must have therapeutic failure or intolerance to one systemic corticosteroid (e.g., prednisone).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For diagnoses in which tumor necrosis factor (TNF) blockers are also indicated (e.g., Rheumatoid Arthritis, Psoriatic Arthritis), the member has experienced therapeutic failure or intolerance to at least 1 TNF blocker.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rinvoq LQ

Products Affected

• RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	For PsA, deny if 18 years of age or older
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rivfloza

Products Affected

- RIVFLOZA SUBCUTANEOUS SOLUTION
- RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3): 1) genetically confirmed diagnosis of primary hyperoxaluria type 1 (PH1), 2) relatively preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2), and 3) at least two elevated urinary oxalate levels greater than 1.5 times the upper reference limit.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, continued preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2) -AND- reduction in urinary oxalate levels from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Romvimza

Products Affected

ROMVIMZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor -AND-attestation that surgical resection may cause one of the following (1-2): 1) worsening functional limitation, 2) severe morbidity
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rozlytrek

Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, the tumor status is ROS1-positive. For solid tumors with NTRK gene fusion without a known acquired resistance mutation, the tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or the tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rubraca

Products Affected

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) Disease harbors a deleterious BRCA mutation (germline or somatic) 2) member is in complete or partial response to platinum-based chemotherapy. For metastatic castration-resistant prostate cancer, all of the following (1-3): 1) disease harbors a deleterious BRCA mutation (germline and/or somatic) 2) member has been treated with androgen receptor-directed therapy and taxane-based chemotherapy 3) member is concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Rydapt

Products Affected

· RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Use as single agent induction therapy for AML
Required Medical Information	Documentation of diagnosis. For a new diagnosis of acute myeloid leukemia, member is using in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation chemotherapy regimens and member is FLT3 mutation positive as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sabril

Products Affected

- vigabatrinVIGADRONE
- VIGPODER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of refractory complex partial seizures -AND-documentation of adjunctive therapy -AND- therapeutic failure or intolerance to at least two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, tiagabine) -OR-documentation of use as monotherapy in treatment of infantile spasms
Age Restrictions	Deny if less than 2 years of age in treatment of refractory complex partial seizures -OR- if less than 1 month old and greater than 2 years of age in treatment of infantile spasms
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Samsca

Products Affected

• tolvaptan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic hypervolemic or euvolemic hyponatremia evidenced by (1. or 2.): 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	Doses must be initiated in the hospital setting to closely monitor serum sodium. Treatment should be limited to 30 days to minimize risk of liver injury. For reauthorization, treatment is for a new episode of a clinically significant euvolemic or hypervolemic hyponatremia -AND- on of the following (1. or 2.) 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Saphris

Products Affected

• asenapine maleate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Savella

Products Affected

• SAVELLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Scemblix

Products Affected

• SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-3): 1) the member is newly-diagnosed, 2) member has been previously treated for Ph+ CML in chronic phase, 3) disease is positive for the T3151 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Secuado

Products Affected

• SECUADO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure of generic asenapine sublingual tablets -AND- trial/failure or intolerance to 1 of the following or all are contraindicated (1-3): 1) olanzapine, 2) quetiapine, 3) risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Simponi

Products Affected

- SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML, 50 MG/0.5 ML
- SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML, 50 MG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Simponi 50mg: For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide) and Simponi will be used in combination with methotrexate. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). Diagnosis of psoriatic arthritis Simponi 100mg: Diagnosis of moderate to severe ulcerative colitis.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For ulcerative colitis, patients must have therapeutic failure or intolerance to the preferred products: a preferred adalimumab product, Stelara SC, Rinvoq, and Xeljanz/Xeljanz XR. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For ulcerative colitis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sirturo

Products Affected

· SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Criteria will be applied consistent with the current ATS/CDC/ERS/IDSA Clinical Practice Guideline for the Treatment of Drug-Susceptible and Drug-Resistant Tuberculosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of disease improvement -AND- member requires additional antimicrobial therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skyclarys

Products Affected

SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Friedreichs ataxia confirmed by genetic testing (i.e., FXN gene mutation).
Age Restrictions	Deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Skyrizi

Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE
- SKYRIZI SUBCUTANEOUS

WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohns disease, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC. For moderate to severe ulcerative colitis, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sohonos

Products Affected

• SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 2.5 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	Deny if female and less than 8 years of age -OR- if male and less than 10 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in volume of new heterotopic ossification from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Solaraze

Products Affected

• diclofenac sodium topical gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure, intolerance, or contraindication to one of the following (1 or 2): 1) topical fluorouracil solution or fluorouracil 5% cream 2) topical imiquimod 5% cream
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of 30 day washout period since optimal therapeutic effect may not be evident until 30 days following cessation of therapy AND attestation of previous response to diclofenac sodium 3% topical gel therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Somavert

Products Affected

· SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin- like growth factor-1 (IGF-1) based on laboratory reference range -AND- inadequate or partial response to surgery or radiotherapy or not a candidate for surgery or radiotherapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sprycel

Products Affected

• dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adults with Ph+ chronic myeloid leukemia, the member is newly diagnosed in the chronic phase -OR- the member is in chronic, accelerated, or myeloid or lymphoid blast phase and has resistance or intolerance to prior therapy including imatinib. For adults with Ph+ acute lymphocytic leukemia, member has had resistance or intolerance to prior therapy. For pediatric patients with Ph+ CML, the member is in the chronic phase. For pediatric patients with Ph+ acute lymphoblastic leukemia, the member is newly diagnosed and will be using in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Stelara

Products Affected

- STELARA SUBCUTANEOUS SOLUTION
- STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML
- STEQEYMA SUBCUTANEOUS
- SYRINGE 45 MG/0.5 ML, 90 MG/ML
- YESINTEK SUBCUTANEOUS SOLUTION
- YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR-inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease, attestation of receiving or currently undergoing a single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC. For Ulcerative colitis, attestation of receiving or currently undergoing a single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC
Age Restrictions	Deny if less than 18 years of age for Crohn's Disease and Ulcerative Colitis or less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	Must follow recommended dosing guidelines based upon weight. Subcutaneous induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Steroidogenesis Inhibitors

Products Affected

- RECORLEV
- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in urinary free cortisol levels from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Stivarga

Products Affected

STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer and trial of a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy, AND an anti-VEGF therapy AND if RAS wild-type, an anti-EGFR therapy -OR- documentation of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with both imatinib and sunitinib -OR- documentation of hepatocellular cancer AND previous treatment with sorafenib
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sutent

Products Affected

• sunitinib malate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For gastrointestinal stromal tumor (GIST), the member has experienced therapeutic failure, intolerance, or contraindication to imatinib. For a high risk of recurrent renal cell carcinoma, member has had a nephrectomy and sunitinib is to be used as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Symdeko

Products Affected

SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have either the homozygous F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to tezacaftor/ivacaftor based on clinical and/or in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Sympazan

Products Affected

· SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND-documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy -AND- unable to tolerate generic clobazam
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Synarel

Products Affected

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For central precocious puberty (CPP), advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L. For female with endometriosis, attestation of not pregnant if of childbearing age -AND-Therapeutic failure, contraindication or intolerance to 2 of the following standard of care treatments: NSAIDs, combination hormonal contraceptive, progestin (i.e. medroxyprogesterone injection), GnRH agonist (i.e. Leuprolide) or danazol
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty. Deny if less than 18 years of age for endometriosis.
Prescriber Restrictions	
Coverage Duration	Endometriosis: 6 months, CPP: 6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization for CPP, attestation of pubertal development slowing from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tabrecta

Products Affected

TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with MET exon 14 skipping mutation as detected by an FDA approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tafinlar

- TAFINLAR ORAL CAPSULE
- TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For dabrafenib monotherapy, documentation of unresectable or metastatic melanoma -AND- documentation of a BRAF V600E mutation. For use in combination with trametinib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with trametinib, documentation of metastatic nonsmall cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA-approved test when FDA indicated. For use in combination with trametinib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Tafinlar tablets for oral suspension, attestation of inability to swallow Tafinlar (dabrafenib) capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tagrisso

Products Affected

· TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) and one of the following (1-5): 1) Adjuvant therapy after tumor resection -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test 2) Locally advanced disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used in combination with pemetrexed and platinum-based chemotherapy, 3) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used with or without combination therapy of pemetrexed and platinum-based chemotherapy, 4) Metastatic disease -AND- disease harbors EGFR T790M mutations, as detected by an FDA-approved test -AND- has progressed on or after EGFR TKI therapy, 5) Locally advanced, unresectable (stage III) disease -AND-disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy -AND- disease harbors EGFR exon 19 deletions or EGFR exon 21 L858R mutations, as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

Taltz

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE SUBCUTANEOUS SYRINGE 20 MG/0.25 ML, 40 MG/0.5 ML, 80 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs.
Age Restrictions	Deny if less than 18 years of age for Psoriatic Arthritis, Ankylosing Spondylitis and non-radiographic axial spondyloarthritis or less than 6 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cosentyx, Enbrel, a preferred adalimumab product, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For plaque psoriasis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Cosentyx, Otezla, Stelara SC, Skyrizi SC, and Enbrel. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For non-radiographic axial spondyloarthritis patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cimzia, Rinvoq, Cosentyx. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Talzenna

Products Affected

• TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of deleterious or suspected deleterious gBRCAm, HER2- negative locally advanced or metastatic breast cancer as a single agent - OR- Documentation of HRR gene-mutated metastatic castration-resistant prostate cancer in combination with enzalutamide
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Targretin

- bexarotene oral
- bexarotene topical

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations of cutaneous T-cell lymphoma in patients who are refractory to at least one prior systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tascenso ODT

Products Affected

· TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Inability to swallow capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tasigna

Products Affected

• nilotinib hcl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adult patients with Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib - OR- member is newly diagnosed in the chronic phase. For pediatric patients, one of the following (1-2): 1) member has chronic phase or accelerated phase Ph+ CML and is is resistant or intolerant to prior tyrosine kinase inhibitor therapy 2) member is newly diagnosed with Ph+ CML in the chronic phase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tasimelteon

Products Affected

• tasimelteon

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tazorac

Products Affected

• tazarotene topical cream

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 of the following (A or B). A) Documentation of plaque psoriasis -AND- trial and failure or intolerance to at least one topical corticosteroid (e.g. fluocinonide, mometasone, triamcinolone, betamethasone). B) Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tazverik

Products Affected

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic epithelioid sarcoma - AND- Disease is not eligible for complete resection. Documentation of relapsed or refractory follicular lymphoma -AND-Tumors are EZH2 mutation positive, as detected by FDA approved test, in a member that has received at least 2 prior systemic therapies -OR- Prescriber attests there are no satisfactory alternative treatment options.
Age Restrictions	For epithelioid sarcoma, deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tecfidera

Products Affected

• dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tepmetko

Products Affected

TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with a MET exon 14 skipping alteration
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Testosterone (androgens)

- testosterone cypionate
- testosterone enanthate
- testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)
- testosterone transdermal gel in packet
- testosterone transdermal solution in metered pump w/app

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or secondary hypogonadism in males with testicular failure due to cryptorchidism, bilateral torsions, orchitis, vanishing testis syndrome, orchidectomy, Klinefelter's syndrome, chemotherapy, radiation or toxic damage -OR- documentation of primary or secondary hypogonadism in males with multiple symptoms of hypogonadism including at least one of the following specific symptoms: height loss due to vertebral fractures, low trauma fractures, low bone density, incomplete or delayed sexual development, breast discomfort, loss of axillar and/or pubic body hair, hot flushes -OR- documentation of HIV infection in men with weight loss -OR- documentation of chronic steroid treatment in men. In all previously noted indications, members must also have documented low total testosterone level below the normal range for the laboratory -OR- a total testosterone level near the lower limit of the normal range with a low free testosterone level which is less than normal based upon the laboratory reference range -OR- the member is not producing any testosterone. Additional approvable indications include female patients with metastatic breast cancer (testosterone enanthate only), primary or secondary hypogonadism in males with testicular failure due to double orchidectomy, and delayed puberty in males (testosterone enanthate only).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.

PA Criteria	Criteria Details
Off Label Uses	HIV Wasting
Part B Prerequisite	No

Thalomid

Products Affected

• THALOMID ORAL CAPSULE 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Use as monotherapy for ENL treatment in the presence of moderate to severe neuritis
Required Medical Information	Documentation of multiple myeloma in combination with dexamethasone -OR- documentation for use in the treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) -OR-documentation of therapy for prevention and suppression of the cutaneous manifestations of ENL recurrence
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Thrombopoiesis Stimulating Agents

- eltrombopag olamine oral powder in packet 12.5 mg, 25 mg
- eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of persistent or chronic immune idiopathic thrombocytopenia purpura and trial and failure of corticosteroid or immunoglobulin therapy or splenectomy -OR- documentation of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy -OR- severe aplastic anemia who have had an insufficient response to immunosuppressive therapy -OR- documentation of first line treatment for severe aplastic anemia and used in combination with at least two immunosuppressive therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Platelet count to be provided
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

Tibsovo

Products Affected

· TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is IDH1 mutation positive as detected by an FDA-approved test. For newly-diagnosed acute myeloid leukemia, member is using as monotherapy or in combination with azacitidine -AND- member meets one of the following (1-5): 1) age is greater than or equal to 75 years of age 2) severe cardiac or pulmonary comorbidity 3) reduced renal function 4) hepatic impairment 5) or prescriber attestation that member is not a candidate for intensive induction therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Topical Lidocaine

- lidocaine hcl mucous membrane solution 4 % (40 mg/ml)
- lidocaine topical ointment
- lidocaine-prilocaine topical cream

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tretinoin

- tretinoin topical creamtretinoin topical gel 0.01 %, 0.025 %

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two generic topical non-retinoid acne medications (e.g. clindamycin, sulfacetamide, erythromycin)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Trikafta

- TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL
- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have at least one F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to elexacaftor/tezacaftor/ivacaftor based on in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Trintellix

Products Affected

TRINTELLIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Truqap

Products Affected

• TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) HR mutation status, HER2 mutation status, and PIK3CA/AKT1/PTEN status 2) concomitant therapy with fulvestrant 3) disease progression on at least one endocrine-based regimen in the metastatic setting -OR- recurrence on or within 12 months of completing adjuvant therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tukysa

Products Affected

• TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced unresectable HER2-positive breast cancer or metastatic HER2-positive breast cancer, member will be using in combination with trastuzumab and capecitabine -AND- member has received one or more prior anti-HER2 based regimens in the metastatic setting. For RAS wild-type HER2-positive unresectable or metastatic colorectal cancer, member will be using in combination with trastuzumab -AND- member has experienced disease progression following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Turalio

Products Affected

• TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor associated with severe morbidity and functional limitations -AND- patient is not amenable to improvement with surgery or not a candidate for surgery
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tykerb

Products Affected

• lapatinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced HER2-positive, or metastatic HER2-positive breast cancer, the member has received prior therapy with an anthracycline, a taxane, and trastuzumab -AND- will be using in combination with capecitabine. For HR+, metastatic breast cancer, the member is post-menopausal -AND- the member's cancer over expresses the HER2 receptor -AND- the member will be using lapatinib in combination with letrozole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Tymlos

Products Affected

· TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 24 months.
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Additional documentation of trial/failure, intolerance or contraindication to preferred parathyroid hormone analog teriparatide. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Tymlos will be limited to a coverage duration of 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Ubrelvy

Products Affected

• UBRELVY ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of acute treatment of migraine with or without aura -AND- trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	Patients must have therapeutic failure, intolerance, or contraindication to Nurtec ODT. For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Uloric

Products Affected

• febuxostat

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic management of hyperuricemia due to gout -And-trial/failure, intolerance or contraindication to allopurinol.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Valchlor

Products Affected

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received at least one prior skin-directed therapy (e.g. topical corticosteroids, topical chemotherapy, local radiation and topical retinoids).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Valtoco

Products Affected

· VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vancomycin

Products Affected

• vancomycin oral capsule 125 mg, 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vanflyta

Products Affected

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is FLT3-ITD-positive as detected by an FDA-approved test -AND- one of the following (1-3): 1) member is receiving induction therapy and is using Vanflyta in combination with standard cytarabine and anthracycline induction therapy 2) member is receiving consolidation therapy and is using Vanflyta in combination with standard cytarabine consolidation therapy 3) member is receiving maintenance therapy and is using Vanflyta as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Venclexta

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
 • VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For newly-diagnosed AML, member is using in combination with azacitidine, decitabine, cytarabine -AND- age greater than or equal to 75 years or presence of at least one comorbidity that precludes use of intensive induction chemotherapy (i.e. severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) is required.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verquvo

Products Affected

· VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of heart failure (NYHA Class II to IV) -AND- Left ventricular ejection fraction less than 45% -AND- Hospitalization for heart failure or received outpatient IV diuretics for heart failure -AND- Used in combination with a angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker or Entresto -AND- Used in combination with bisoprolol, carvedilol IR/ER or metoprolol succinate ER.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Verzenio

Products Affected

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative. For early breast cancer that is at high risk of recurrence and is node-positive, all of the following (1-2): 1) used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) 2) used as adjuvant treatment. For advanced or metastatic breast cancer, used as initial endocrine-based therapy and used in combination with an aromatase inhibitor -OR- used after documented disease progression following endocrine therapy and used in combination with fulvestrant -OR- used after documented disease progression and used following endocrine therapy and prior chemotherapy in the metastatic setting and will be used as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Viberzi

Products Affected

VIBERZI

PA Criteria	Criteria Details
Exclusion Criteria	Severe (Child-Pugh C) hepatic impairment
Required Medical Information	Documentation of diarrhea predominant, irritable bowel syndrome (IBS-D) -AND- trial/failure or intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Viibryd

Products Affected

• vilazodone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vijoice

- VIJOICE ORAL GRANULES IN MG PACKET
- VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe manifestations of PIK3CA Related Overgrowth Spectrum (PROS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vitrakvi

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- · VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation -AND- Tumors are metastatic or surgical resection is likely to result in severe morbidity -AND- There are no satisfactory alternative treatments or tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vivjoa

Products Affected

· VIVJOA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of recurrent vulvovaginal candidiasis defined as at least 3 episodes of vulvovaginal candidiasis in less than one year -AND Documentation the member is NOT of reproductive potential defined as postmenopausal or another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy) -AND- the member has experienced therapeutic failure, contraindication, or intolerance to a sixmonth maintenance course of oral fluconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vizimpro

Products Affected

VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer -AND- one of the following, as detected by an FDA-approved test (1 or 2): 1) Epidermal growth factor (EGFR) exon 19 deletions, 2) Epidermal growth factor receptor (EGFR) exon 21 L858R substitution mutations.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vonjo

Products Affected

· VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND-attestation of a platelet count of less than 50 x 10*9/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voranigo

Products Affected

 VORANIGO ORAL TABLET 10 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) grade 2 astrocytoma or grade 2 oligodendroglioma, 2) disease harbors a susceptible isocitrate dehydrogenase (IDH)-1 or IDH-2 mutation, as detected by an FDA-approved test, 3) will be used following surgery including biopsy, sub-total resection, or gross total resection.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Voriconazole

Products Affected

• voriconazole intravenous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary cannot take oral voriconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of continued indicators of active disease (e.g. histopathology, positive cultures) is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vosevi

Products Affected

· VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Votrient

Products Affected

• pazopanib oral tablet 200 mg

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of adipocytic soft tissue sarcoma or gastrointestinal stromal tumor
Required Medical Information	Documentation of diagnosis. For advanced soft-tissue sarcoma, trial/failure of at least one prior chemotherapy regimen.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vraylar

Products Affected

• VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For a diagnosis of schizophrenia and bipolar I disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Vumerity

Products Affected

VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Welireg

Products Affected

WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For von Hippel Lindau (VHL) syndrome, one of the following diagnoses not requiring immediate surgery (1, 2, or 3): 1) Renal cell carcinoma (RCC) 2) CNS hemangioblastoma 3) Pancreatic neuroendocrine tumor. For advanced RCC with a clear cell component, prior treatment with a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) -AND- a programmed death receptor-1 (PD-1) or a programmed death-ligand (PD-L1) inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xalkori

- XALKORI ORAL CAPSULE
- XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer (NSCLC), disease is ALK-positive as detected by an FDA-approved test or ROS1-positive as detected by an FDA-approved test. For relapsed or refractory anaplastic large cell lymphoma (ALCL), disease is ALK-positive. For unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT), disease is ALK-positive.
Age Restrictions	Deny if less than 1 year of age or greater than 21 years of age for ALCL.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For crizotinib oral pellets and NSCLC, inability to swallow capsules is required. For crizotinib oral pellets and ALCL / IMT, inability to swallow oral capsules -OR- body surface area less than 1.34 m2 is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xcopri

- XCOPRI
- XCOPRI MAINTENANCE PACK
- XCOPRI TITRATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Therapeutic failure, intolerance or contraindication to 1 other anti-epileptic drug (e.g. carbamazepine, levetiracetam)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xdemvy

Products Affected

• XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis confirmed by identification of Demodex infection via microscopic examination of pulled eyelashes -OR-identification of collarettes via slit-lamp evaluation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xeljanz

- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis and an inadequate response or intolerance to methotrexate. Xeljanz immediate release for juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID).
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis, psoriatic arthritis, ulcerative colitis, ankylosing spondylitis. For Xeljanz regular release tablet, deny if less than 2 years of age for juvenile idiopathic arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced inadequate response or intolerance to at least 1 tumor necrosis factor (TNF) blocker. Doses greater than 10 mg per day for Xeljanz and 11 mg per day for Xeljanz XR will not be approved for rheumatoid arthritis and psoriatic arthritis. Doses greater than 20mg per day for Xeljanz and 22 mg per day for Xeljanz XR will not be approved for ulcerative colitis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xeljanz Solution

Products Affected

• XELJANZ ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of juvenile idiopathic arthritis -AND- Inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xenazine

Products Affected

• tetrabenazine oral tablet 12.5 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary is not actively suicidal
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	In patients with comorbid depression, attestation of adequate treatment for depression is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xermelo

Products Affected

• XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of carcinoid syndrome diarrhea AND used in combination with a somatostatin analog AND trial and failure of somatostatin analog monotherapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in average number of daily bowel movements -AND- the member will continue to use in combination with a somatostatin analog.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xgeva

Products Affected

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For hypercalcemia of malignancy, refractory to bisphosphonates. For giant cell tumor of bone, unresectable or surgical resection is likely to result in severe morbidity -AND- one of the following (1. or 2.)- 1.) the member is 18 years old or older -OR- 2.) the member is a skeletally mature adolescent (e.g. has at least one mature long bone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xifaxan

Products Affected

• XIFAXAN ORAL TABLET 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 or 2. 1) Diagnosis of hepatic encephalopathy AND trial/failure, intolerance, or contraindication to lactulose. 2) Diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D) AND trial/failure, intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Hepatic encephalopathy: 1 year. IBS-D: 14 days.
Other Criteria	No more than three courses of rifaximin for the treatment of IBS-D will be approved per lifetime.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xolair

Products Affected

XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of the following (1-2): 1) Chronic Spontaneous Urticaria, 2) trial/failure or intolerance of a second-generation non-sedating H1 antihistamine at the maximum recommended doses (e.g. cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine)OR-Documentation of the following (3-9): 3) moderate to severe persistent asthma, 4) a positive skin test or in vitro reactivity to a perennial aeroallergen, 5) Baseline IgE titer greater than or equal to 30 IU/mL, 6) documented pretreatment FEV1 less than 80% predicted in adults or less than 90% predicted in children and adolescents or FEV1 reversibility of at least 12% and 200mL after albuterol administration, 7) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos, 8) inadequately controlled symptoms despite a 3-month trial of both of the following (a-b) a) medium-dose inhaled corticosteroid or systemic steroid b) a long-acting beta-agonist or leukotriene antagonist, 9) currently on 1 of the following (c, d, e): c) a long-acting beta2-agonist, d) leukotriene modifier, or e) theophyllineOR- Documentation of the following (10-11): 10) chronic rhinosinusitis with nasal polyps (CRSwNP), 11) will use concomitantly with nasal corticosteroid maintenance treatment, -OR- Documentation of the following (12-17): 12) IgE mediated food allergy, 13) diagnosis confirmed by skin prick test or food-specific antibodies, 14) previous allergic reaction to food, 15) using for the reduction of allergic reactions (type 1), including anaphylaxis, 16) will be used in conjunction with food allergen avoidance, 17) member has a documented prescription for epinephrine.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CSU reauthorization, improved CSU symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For IgE-mediated food allergy reauthorization, member requires continuation of therapy and will continue food allergen avoidance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xolremdi

Products Affected

· XOLREMDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of WHIM syndrome (warts, hypogammaglobulinemia, infections, myelokathexis).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in incidence of infections is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xospata

Products Affected

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is FLT3 mutation-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xpovio

Products Affected

XPOVIO ORAL TABLET 100
 MG/WEEK (50 MG X 2), 40 MG/WEEK
 (10 MG X 4), 40 MG/WEEK (40 MG X 1),
 40MG TWICE WEEK (40 MG X 2), 60
 MG/WEEK (60 MG X 1), 60MG TWICE
 WEEK (120 MG/WEEK), 80 MG/WEEK
 (40 MG X 2), 80MG TWICE WEEK (160
 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use in combination with dexamethasone for relapse or refractory multiple myeloma with failure, intolerance or contraindication to 5 therapies (e.g. bortezomib, carfilzomib, lenalidomide, pomalidomide and daratumumab) -OR- Documentation of use in combination with both bortezomib and dexamethasone for multiple myeloma after receiving 1 prior multiple myeloma therapy -OR- Documentation of relapsing or refractory diffuse large B-cell lymphoma with failure, intolerance or contraindication to at least 2 lines of systemic therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xtandi

- · XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, $80 \ \text{MG}$

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For castration-resistant prostate cancer and metastatic castration sensitive-prostate cancer, the member is using in combination with a GnRH analog or the member has had a bilateral orchiectomy. For non-metastatic castration-sensitive prostate cancer, the member has biochemical recurrence at high risk for metastasis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Xyrem

- sodium oxybate XYREM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND-Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If no diagnosis of cataplexy- trial and failure, intolerance, or contraindication to generic modafinil AND a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Yonsa

Products Affected

· YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with methylprednisolone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Yorvipath

Products Affected

 YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML, 294 MCG/0.98 ML, 420 MCG/1.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hypoparathyroidism -AND- albumin-corrected serum calcium greater than or equal to 7.8 mg/dL -AND- trial and failure of calcium -AND- trial and failure of an active form of vitamin D (e.g. calcitriol, alfacalcidol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of an improvement in total serum calcium from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zavesca

- miglustatYARGESA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of mild to moderate type 1 Gaucher disease confirmed by the following (A. or B.) A. (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm*3 -OR- 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia)OR- B. Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation enzyme replacement therapy (e.g. Cerezyme, Elelyso, or VPRIV) is not a therapeutic option
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zavzpret

Products Affected

ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND-Therapeutic failure, contraindication or intolerance to one generic triptan -AND-Inability to swallow capsules/tablets
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zejula

Products Affected

• ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer, member is in complete or partial response to first-line platinum-based therapy -AND- disease is associated with homologous recombination deficiency-positive status, defined by at least one of the following (1 or 2): 1) a deleterious or suspected deleterious BRCA mutation or 2) genomic instability. For recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, all of the following (1-2): 1) disease harbors a deleterious or suspected deleterious germline BRCA mutation 2) member is in a complete or partial response to platinum-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zelboraf

Products Affected

· ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma and use in combination with cobimetinib, member has a BRAF V600E or V600K mutation. For unresectable or metastatic melanoma and use as monotherapy, member has a BRAF V600E mutation. For Erdheim-Chester Disease and use as monotherapy, member has a BRAF V600 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zelsuvmi

Products Affected

· ZELSUVMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of molluscum contagiosum
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	For reauthorization, attestation that the member has previously experienced complete or partial clearance of molluscum contagiosum lesions with Zelsuvmi -AND- additional course of therapy is required for recurrence of molluscum contagiosum
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zeposia

Products Affected

- · ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details	
Exclusion Criteria	Concomitant use of Zeposia and other disease modifying agents such as interferons, Copaxone, Tysabri.	
Required Medical Information	Documentation of diagnosis. For moderate to severe active ulcerative colitis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate, cyclosporine).	
Age Restrictions		
Prescriber Restrictions		
Coverage Duration	Ulcerative Colitis: 12 months, Multiple Sclerosis: 24 months	
Other Criteria	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq, Xeljanz/Xeljanz XR and Stelara SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.	
Indications	All FDA-approved Indications.	
Off Label Uses		
Part B Prerequisite	No	

Zolinza

Products Affected

· ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations in patients with cutaneous T-cell lymphoma (CTCL) who have progressive, persistent, or recurrent disease on or following 2 systemic therapies. Systemic therapies include bexarotene, interferon alpha, extracorporeal photochemotherapy, PUVA, single agent or combination chemotherapies (e.g. cyclophosphamide, vinblastine, romidepsin)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zonisade

Products Affected

· ZONISADE

PA Criteria	Criteria Details	
Exclusion Criteria		
Required Medical Information	Documentation of partial-onset seizures -AND- Documentation of adjunctive therapy -AND- Inability to swallow capsules -AND- Therapeutic failure/intolerance to 2 or contraindication to all of the following (1-6): 1) generic carbamazepine suspension/chewable tablet/extended-release capsule, 2) generic gabapentin capsules/solution, 3) generic lacosamide solution, 4) generic levetiracetam solution, 5) generic oxcarbazepine suspension, 6) generic pregabalin capsules/solution.	
Age Restrictions		
Prescriber Restrictions		
Coverage Duration	12 months	
Other Criteria		
Indications	All FDA-approved Indications.	
Off Label Uses		
Part B Prerequisite	No	

Ztalmy

Products Affected

• ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures associated with CDKL5 deficiency confirmed by genetic testing -AND- therapeutic failure or intolerance to 2 previous antiepileptic therapies
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zurzuvae

Products Affected

 ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of moderate to severe postpartum depression -AND- less than or equal to 12 months postpartum.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zydelig

Products Affected

· ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	First line treatment. Combination use with benadmustine and/or rituximab for the treatment of FL.
Required Medical Information	Documentation of relapsed chronic lymphocytic leukemia -AND- all of the following (1-2): 1) will be used in combination with rituximab 2) use of rituximab alone would be appropriate due to other due to other comorbidities.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zykadia

Products Affected

· ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- ALK mutations, as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Zytiga

Products Affected

- abiraterone oral tablet 250 mg, 500 mgABIRTEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with prednisone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

Index of Drugs	
abiraterone oral tablet 250 mg, 500 mg355	BALVERSA25
ABIRTEGA355	BENLYSTA SUBCUTANEOUS27
acetaminophen-codeine oral solution 120-	benztropine oral112
12 mg/5 ml	BESREMI29
acetaminophen-codeine oral tablet222	BETASERON SUBCUTANEOUS KIT. 133
acitretin1	bexarotene oral280
ACTEMRA ACTPEN2	bexarotene topical280
ACTEMRA SUBCUTANEOUS2	BIVIGAM 121
ACTIMMUNE3	BOSULIF ORAL CAPSULE 100 MG,
ADBRY 4	50 MG 30
ADEMPAS	BOSULIF ORAL TABLET 100 MG,
AIMOVIG AUTOINJECTOR	400 MG, 500 MG
SUBCUTANEOUS AUTO-INJECTOR	BRAFTOVI 31
140 MG/ML, 70 MG/ML 8	BRINSUPRI32
AJOVY AUTOINJECTOR9	BRUKINSA ORAL CAPSULE33
AJOVY SYRINGE9	buprenorphine222
AKEEGA 10	CABLIVI INJECTION KIT 35
ALCOHOL PADS 131	CABOMETYX 36
ALECENSA11	CALQUENCE (ACALABRUTINIB
alosetron oral tablet 0.5 mg, 1 mg 166	MAL)37
alprazolam oral tablet 0.25 mg, 0.5 mg, 1	CAMZYOS 38
<i>mg</i> , 2 <i>mg</i>	CAPLYTA 39
ALUNBRIG ORAL TABLET 180 MG,	CAPRELSA ORAL TABLET 100 MG,
30 MG, 90 MG	300 MG 40
ALUNBRIG ORAL TABLETS, DOSE	carglumic acid41
PACK 13	CAYSTON 42
ALYQ 229	CERDELGA 43
ambrisentan229	CIBINQO 46
amitriptyline 112	CIMZIA POWDER FOR RECONST47
ANZUPGO	CIMZIA SUBCUTANEOUS SYRINGE
ARIKAYCE 16	KIT 400 MG/2 ML (200 MG/ML X 2) 47
aripiprazole oral solution18	CINRYZE 49
aripiprazole oral tablet, disintegrating 18	clobazam oral suspension205
<i>armodafinil</i> 197	clobazam oral tablet205
asenapine maleate253	<i>clomipramine</i> 112
AUGTYRO ORAL CAPSULE 160 MG,	clonazepam oral tablet 0.5 mg, 1 mg, 2 mg 222
40 MG	clonazepam oral tablet, disintegrating
AUVELITY 21	0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg222
AVMAPKI-FAKZYNJA22	clorazepate dipotassium oral tablet 15 mg,
AVONEX INTRAMUSCULAR PEN	3.75 mg, 7.5 mg
INJECTOR KIT 133	COBENFY 51
AVONEX INTRAMUSCULAR	COBENFY STARTER PACK51
SYRINGE KIT 133	
AYVAKIT	
BAFIERTAM 24	

COMETRIQ ORAL CAPSULE 100	DOPTELET (15 TAB PACK) 65
MG/DAY(80 MG X1-20 MG X1), 140	DOPTELET (30 TAB PACK) 65
MG/DAY(80 MG X1-20 MG X3), 60	doxepin oral capsule112
MG/DAY (20 MG X 3/DAY) 52	doxepin oral concentrate112
COPAXONE SUBCUTANEOUS	doxepin oral tablet112
SYRINGE 20 MG/ML, 40 MG/ML 104	DRIZALMA SPRINKLE ORAL
COPIKTRA 53	CAPSULE, DELAYED REL
CORLANOR ORAL SOLUTION54	SPRINKLE 20 MG, 30 MG, 40 MG, 60
COSENTYX (2 SYRINGES)55	MG 66
COSENTYX PEN (2 PENS)55	droxidopa oral capsule 100 mg, 200 mg,
COSENTYX SUBCUTANEOUS	<i>300 mg</i>
SYRINGE 75 MG/0.5 ML	DUPIXENT PEN SUBCUTANEOUS
COSENTYX UNOREADY PEN55	PEN INJECTOR 200 MG/1.14 ML, 300
COTELLIC 56	MG/2 ML67
CYLTEZO(CF) 114	DUPIXENT SYRINGE
CYLTEZO(CF) PEN 114	SUBCUTANEOUS SYRINGE 200
CYLTEZO(CF) PEN CROHN'S-UC-HS	MG/1.14 ML, 300 MG/2 ML67
SUBCUTANEOUS PEN INJECTOR	DUVYZAT 69
KIT 40 MG/0.8 ML 114	eltrombopag olamine oral powder in
CYLTEZO(CF) PEN PSORIASIS-UV	packet 12.5 mg, 25 mg291
SUBCUTANEOUS PEN INJECTOR	eltrombopag olamine oral tablet 12.5 mg,
KIT 40 MG/0.8 ML 114	25 mg, 50 mg, 75 mg291
CYSTARAN57	EMGALITY PEN71
dalfampridine14	EMGALITY SYRINGE
DANZITEN58	SUBCUTANEOUS SYRINGE 120
dasatinib oral tablet 100 mg, 140 mg, 20	MG/ML, 300 MG/3 ML (100 MG/ML X
mg, 50 mg, 70 mg, 80 mg265	3)71
DAURISMO ORAL TABLET 100 MG,	ENBREL MINI73
25 MG	ENBREL SUBCUTANEOUS
DAYBUE 61	SOLUTION 73
<i>deferasirox</i>	ENBREL SUBCUTANEOUS SYRINGE
deferiprone88	25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)73
DIACOMIT ORAL CAPSULE 250 MG,	ENBREL SURECLICK73
500 MG	ENDOCET 222
DIACOMIT ORAL POWDER IN	EPCLUSA ORAL PELLETS IN
PACKET 250 MG, 500 MG63	PACKET 150-37.5 MG74
DIAZEPAM INTENSOL222	EPIDIOLEX 75
diazepam oral solution 5 mg/5 ml (1	ergotamine-caffeine77
<i>mg/ml</i>)222	ERIVEDGE78
diazepam oral tablet222	ERLEADA ORAL TABLET 240 MG,
diclofenac sodium topical gel 3 %263	60 MG 79
dihydroergotamine nasal64	erlotinib70
dimethyl fumarate oral capsule,delayed	eszopiclone222
release(dr/ec) 120 mg, 120 mg (14)- 240	EVENITY SUBCUTANEOUS
<i>mg</i> (46), 240 <i>mg</i> 286	SYRINGE 210MG/2.34ML (
DOPTELET (10 TAB PACK) 65	105MG/1.17MLX2)80

everolimus (antineoplastic) oral tablet 10	GAUZE PAD TOPICAL BANDAGE 2
<i>mg</i> , 2.5 <i>mg</i> , 5 <i>mg</i> , 7.5 <i>mg</i> 6	X 2 "
everolimus (antineoplastic) oral tablet for	GAVRETO 102
suspension 2 mg, 3 mg, 5 mg6	<i>gefitinib</i> 136
EVRYSDI ORAL RECON SOLN 81	GILOTRIF 70
EVRYSDI ORAL TABLET 81	glatiramer subcutaneous syringe 20 mg/ml,
FABHALTA 82	<i>40 mg/ml</i> 104
FANAPT 84	GLATOPA SUBCUTANEOUS
FANAPT TITRATION PACK A84	SYRINGE 20 MG/ML, 40 MG/ML 104
FASENRA PEN 85	GLEOSTINE 107
FASENRA SUBCUTANEOUS	GOMEKLI ORAL CAPSULE 1 MG, 2
SYRINGE 10 MG/0.5 ML, 30 MG/ML 85	MG 109
<i>febuxostat</i> 303	GOMEKLI ORAL TABLET FOR
fentanyl transdermal patch 72 hour 100	SUSPENSION 109
mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr,	guanfacine oral tablet extended release 24
75 mcg/hr222	<i>hr</i> 5
FETZIMA ORAL CAPSULE,EXT REL	HADLIMA 114
24HR DOSE PACK 20 MG (2)- 40 MG	HADLIMA PUSHTOUCH114
(26)89	HADLIMA(CF) 114
FETZIMA ORAL	HADLIMA(CF) PUSHTOUCH114
CAPSULE,EXTENDED RELEASE 24	HERNEXEOS111
HR 120 MG, 20 MG, 40 MG, 80 MG89	HUMIRA PEN114
FILSUVEZ 90	HUMIRA SUBCUTANEOUS
fingolimod103	SYRINGE KIT 40 MG/0.8 ML 114
FINTEPLA 91	HUMIRA(CF) 114
FIRDAPSE 94	HUMIRA (CF) PEN 114
FIRMAGON KIT W DILUENT	HUMIRA(CF) PEN CROHNS-UC-HS 114
SYRINGE 95	HUMIRA(CF) PEN PSOR-UV-ADOL
FOTIVDA97	HS 114
FRUZAQLA ORAL CAPSULE 1 MG, 5	hydrocodone-acetaminophen oral tablet
MG98	10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325
FUROSCIX99	<i>mg</i>
gabapentin oral capsule 100 mg, 300 mg,	hydromorphone oral liquid222
400 mg100	hydromorphone oral tablet 2 mg, 4 mg, 8
gabapentin oral solution 250 mg/5 ml100	mg222
gabapentin oral tablet 600 mg, 800 mg100	hydroxyzine hcl oral tablet112
GAMMAGARD LIQUID121	IBRANCE 116
GAMMAGARD S-D (IGA < 1	IBSRELA 117
MCG/ML)121	IBTROZI 118
GAMMAKED INJECTION	icatibant 92
SOLUTION 1 GRAM/10 ML (10 %) 121	ICLUSIG
GAMMAPLEX121	IDHIFA ORAL TABLET 100 MG, 50
GAMMAPLEX (WITH SORBITOL) 121	MG
GAMUNEX-C INJECTION	<i>imatinib oral tablet 100 mg, 400 mg</i> 105
SOLUTION 1 GRAM/10 ML (10 %) 121	IMBRUVICA ORAL CAPSULE 140
GATTEX 30-VIAL101	
GATTLA 50-VIAL101	MG, 70 MG 123

IMBRUVICA ORAL SUSPENSION		LEUKINE INJECTION RECON SOL	
IMBRUVICA ORAL TABLET 140 M	,		
280 MG, 420 MG	123	lidocaine hcl mucous membrane solution	4
imipramine hcl	112	% (40 mg/ml)	293
IMKELDI	124	lidocaine topical adhesive patch, medicate	ed –
INCRELEX	126	5 %	160
INLYTA	128	lidocaine topical ointment	293
INQOVI	129	lidocaine-prilocaine topical cream	293
INREBIC	130	LITFULO	
ITOVEBI ORAL TABLET 3 MG, 9 N	ЛG	LIVTENCITY	162
	137	LOKELMA	163
itraconazole oral capsule	138	LONSURF	164
ivabradine oral tablet 5 mg, 7.5 mg		LORAZEPAM INTENSOL	222
ivermectin oral	139	lorazepam oral tablet 0.5 mg, 1 mg, 2 mg	222
IWILFIN	140	LORBRENA ORAL TABLET 100 MG	
JAKAFI	141	25 MG	165
JAVYGTOR	154	LUMAKRAS ORAL TABLET 120 MO	J,
JAYPIRCA ORAL TABLET 100 MG	r,	240 MG, 320 MG	167
50 MG	142	LUPRON DEPOT-PED (3 MONTH)	
JOENJA	143	INTRAMUSCULAR SYRINGE KIT	
KALYDECO ORAL GRANULES IN		11.25 MG	168
PACKET 13.4 MG, 25 MG, 5.8 MG, 5	50	LUPRON DEPOT-PED	
MG, 75 MG	144	INTRAMUSCULAR KIT 7.5 MG (PE	D)
KALYDECO ORAL TABLET	144		168
KERENDIA		LUPRON DEPOT-PED	
KESIMPTA PEN	146	INTRAMUSCULAR SYRINGE KIT	168
KEVZARA	147	lurasidone oral tablet 120 mg, 20 mg, 40	
KINERET	149	mg, 60 mg, 80 mg	155
KISQALI FEMARA CO-PACK ORA	\mathbf{L}	LYNPARZA	169
TABLET 400 MG/DAY(200 MG X 2)	-	LYTGOBI ORAL TABLET 12	
2.5 MG, 600 MG/DAY(200 MG X 3)-2	2.5	MG/DAY (4 MG X 3), 16 MG/DAY (4	
MG	150	MG X 4), 20 MG/DAY (4 MG X 5)	172
KISQALI ORAL TABLET 200		MAVYRET ORAL PELLETS IN	
MG/DAY (200 MG X 1), 400 MG/DAY	Y	PACKET	173
(200 MG X 2), 600 MG/DAY (200 MG	X	MAVYRET ORAL TABLET	173
3)		megestrol oral suspension 400 mg/10 ml	
KOSELUGO ORAL CAPSULE 10 M	G,	(40 mg/ml), 625 mg/5 ml (125 mg/ml)	174
25 MG	152	megestrol oral tablet	174
KRAZATI	153	MEKINIST ORAL RECON SOLN	175
lapatinib	300	MEKINIST ORAL TABLET 0.5 MG,	2
LAZCLUZE ORAL TABLET 240 MO	G,	MG	
80 MG	156	MEKTOVI	176
lenalidomide	238	memantine-donepezil	181
LENVIMA	157	methadone oral solution 10 mg/5 ml, 5	
		mg/5 ml	222
		methadone oral tablet 10 mg, 5 mg	222

metyrosine17°	OJEMDA ORAL TABLET 400
mifepristone oral tablet 300 mg 15	MG/WEEK (100 MG X 4), 500
miglustat343	MG/WEEK (100 MG X 5), 600
modafinil22	7 MG/WEEK (100 MG X 6)202
MODEYSO 178	
morphine concentrate oral solution 222	2 OLUMIANT 204
morphine oral solution 10 mg/5 ml, 20	ONUREG 206
$mg/5 ml (4 mg/ml) \dots 222$	
morphine oral tablet222	
morphine oral tablet extended release 100	OPSYNVI 229
mg, 15 mg, 200 mg, 30 mg, 60 mg222	
MOUNJARO 108	
MULPLETA 179	
NAMZARIC ORAL	87.5 MG/0.7 ML 208
CAPSULE,SPRINKLE,ER 24HR 7-10	ORGOVYX 210
MG 183	
NAYZILAM 182	
NEMLUVIO 183	
NERLYNX 184	
NEXLETOL180	
NEXLIZET 180	5 OTEZLA
nilotinib hcl282	
NINLARO 188	TABLETS,DOSE PACK 10 MG (4)- 20
nitisinone189	MG (51), 10 MG (4)-20 MG (4)-30 MG
NORDITROPIN FLEXPRO110) (47)213
NUBEQA 192	oxycodone oral capsule222
NUCALA SUBCUTANEOUS AUTO-	oxycodone oral concentrate222
INJECTOR 192	2 oxycodone oral tablet 10 mg, 15 mg, 20
NUCALA SUBCUTANEOUS RECON	mg, 30 mg, 5 mg222
SOLN 192	2 oxycodone-acetaminophen oral tablet 10-
NUCALA SUBCUTANEOUS	325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg
SYRINGE 100 MG/ML, 40 MG/0.4 ML 192	2
NUEDEXTA 194	4 OZEMPIC SUBCUTANEOUS PEN
NUPLAZID 195	
NURTEC ODT 196	6 MG/3 ML), 1 MG/DOSE (4 MG/3 ML),
OCTAGAM12	
octreotide acetate injection solution199	PALYNZIQ SUBCUTANEOUS
ODOMZO 200	,
OFEV 134	4 ML, 20 MG/ML 214
OGSIVEO ORAL TABLET 100 MG,	PANRETIN
150 MG, 50 MG	
OJEMDA ORAL SUSPENSION FOR	pazopanib oral tablet 200 mg321
RECONSTITUTION 202	
	PEMAZYRE 216
	PHEBURANE217
	phenobarbital113

PIQRAY ORAL TABLET 200		RINVOQ ORAL TABLET EXTENDED)
MG/DAY (200 MG X 1), 250 MG/DAY		RELEASE 24 HR 15 MG, 30 MG, 45	
(200 MG X1-50 MG X1), 300 MG/DAY		MG	. 243
(150 MG X 2)	.218	RIVFLOZA SUBCUTANEOUS	
pirfenidone oral capsule	.134	SOLUTION	. 246
pirfenidone oral tablet	.134	RIVFLOZA SUBCUTANEOUS	
PLEGRIDY SUBCUTANEOUS PEN		SYRINGE 128 MG/0.8 ML, 160	
INJECTOR 125 MCG/0.5 ML	. 133	MG/ML	.246
PLEGRIDY SUBCUTANEOUS		ROMVIMZA	247
SYRINGE 125 MCG/0.5 ML	133	ROZLYTREK ORAL CAPSULE 100	
POMALYST	. 219	MG, 200 MG	. 248
posaconazole oral tablet,delayed release		ROZLYTREK ORAL PELLETS IN	
(dr/ec)	. 220	PACKET	. 248
pregabalin oral capsule 100 mg, 150 mg,		RUBRACA	. 249
200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 7	5	rufinamide	26
<i>mg</i>	. 171	RYBELSUS	. 108
pregabalin oral solution	. 171	RYDAPT	250
PRENATAL VITAMIN PLUS LOW		SAJAZIR	92
IRON	.221	sapropterin	. 154
PREVYMIS ORAL PELLETS IN		SAVELLA	
PACKET	. 225	SCEMBLIX ORAL TABLET 100 MG,	
PRIVIGEN	. 121	20 MG, 40 MG	. 255
PROLASTIN-C INTRAVENOUS		SECUADO	.256
SOLUTION	12	SIGNIFOR	. 267
PROLIA	226	sildenafil (pulm.hypertension) oral tablet.	. 229
promethazine oral		SIMLANDI(CF)	
PULMOZYME	. 230	SIMLANDI(CF) AUTOINJECTOR	.114
pyrimethamine	59	SIMPONI SUBCUTANEOUS PEN	
QINLOCK		INJECTOR 100 MG/ML, 50 MG/0.5	
quinine sulfate		ML	.257
QULIPTA	.233	SIMPONI SUBCUTANEOUS	
RADICAVA ORS STARTER KIT		SYRINGE 100 MG/ML, 50 MG/0.5 ML	, 257
SUSP		SIRTURO	
RAVICTI		SKYCLARYS	. 260
RECORLEV		SKYRIZI SUBCUTANEOUS PEN	
REPATHA SURECLICK		INJECTOR	. 261
REPATHA SYRINGE	. 236	SKYRIZI SUBCUTANEOUS	
RETEVMO ORAL TABLET 120 MG,		SYRINGE	.261
160 MG, 40 MG, 80 MG		SKYRIZI SUBCUTANEOUS	
REVUFORJ ORAL TABLET 110 MG,		WEARABLE INJECTOR 180 MG/1.2	
160 MG, 25 MG		ML (150 MG/ML), 360 MG/2.4 ML (15	
REXULTI ORAL TABLET		MG/ML)	
REZDIFFRA		sodium oxybate	
REZLIDHIA		sodium phenylbutyrate	
RINVOQ LQ	. 245	sofosbuvir-velpatasvir	74

SOHONOS ORAL CAPSULE 1 MG, 1	l . 5	TOBI PODHALER	44
MG, 10 MG, 2.5 MG, 5 MG	262	tobramycin in 0.225 % nacl	44
SOMAVERT	264	tobramycin inhalation	44
sorafenib	185	tolvaptan	. 252
STELARA SUBCUTANEOUS		topiramate oral solution	76
SOLUTION	266	tramadol oral tablet 50 mg	
STELARA SUBCUTANEOUS		tramadol-acetaminophen	.222
SYRINGE 45 MG/0.5 ML, 90 MG/ML	266	tretinoin topical cream	
STEQEYMA SUBCUTANEOUS		tretinoin topical gel 0.01 %, 0.025 %	.294
SYRINGE 45 MG/0.5 ML, 90 MG/ML	266	TRIKAFTA ORAL GRANULES IN	
STIVARGA	268	PACKET, SEQUENTIAL	295
sunitinib malate	269	TRIKAFTA ORAL TABLETS,	
SYMDEKO	270	SEQUENTIAL	.295
SYMPAZAN	271	trimipramine	. 112
SYNAREL	272	TRINTELLIX	. 296
TABRECTA	273	TRULICITY	108
tadalafil (pulm. hypertension)	229	TRUQAP	297
tadalafil oral tablet 2.5 mg, 5 mg	45	TUKYSA ORAL TABLET 150 MG, 50	
TAFINLAR ORAL CAPSULE	274	MG	298
TAFINLAR ORAL TABLET FOR		TURALIO ORAL CAPSULE 125 MG	299
SUSPENSION	274	TYMLOS	.301
TAGRISSO	275	UBRELVY ORAL TABLET 100 MG,	
TALTZ AUTOINJECTOR	277	50 MG	302
TALTZ SYRINGE SUBCUTANEOUS	5	UPTRAVI ORAL TABLET 1,000 MCG	r,
SYRINGE 20 MG/0.25 ML, 40 MG/0.5	5	1,200 MCG, 1,400 MCG, 1,600 MCG,	
ML, 80 MG/ML	277	200 MCG, 400 MCG, 600 MCG, 800	
TALZENNA	279	MCG	229
TASCENSO ODT	281	UPTRAVI ORAL TABLETS,DOSE	
tasimelteon	283	PACK	.229
tazarotene topical cream	284	VALCHLOR	. 304
TAZVERIK		VALTOCO	.305
TEPMETKO	287	vancomycin oral capsule 125 mg, 250 mg.	306
teriflunomide	19	VANFLYTA	.307
teriparatide subcutaneous pen injector 20)	VENCLEXTA ORAL TABLET 10 MG	,
mcg/dose (560mcg/2.24ml)	96	100 MG, 50 MG	308
testosterone cypionate	288	VENCLEXTA STARTING PACK	.308
testosterone enanthate	288	VERQUVO	.309
testosterone transdermal gel in metered-		VERZENIO	.310
dose pump 20.25 mg/1.25 gram (1.62 %)	288	VIBERZI	311
testosterone transdermal gel in packet		vigabatrin	251
testosterone transdermal solution in		VIGADRONE	
metered pump w/app	288	VIGPODER	.251
tetrabenazine oral tablet 12.5 mg, 25 mg.		VIJOICE ORAL GRANULES IN	
THALOMID ORAL CAPSULE 100		PACKET	313
MG, 50 MG	290		
TIRSOVO	292		

VIJUICE URAL TABLET 125 MIG, 250	,	ATANDI OKAL TADLET 40 MG, 80	
MG/DAY (200 MG X1-50 MG X1), 50		MG	339
MG	313	XYREM	340
vilazodone	312	YARGESA	.343
VITRAKVI ORAL CAPSULE 100 MG,		YESINTEK SUBCUTANEOUS	
25 MG		SOLUTION	266
VITRAKVI ORAL SOLUTION		YESINTEK SUBCUTANEOUS	
VIVJOA		SYRINGE 45 MG/0.5 ML, 90 MG/ML	266
VIZIMPRO		YONSA	
VONJO		YORVIPATH SUBCUTANEOUS PEN	511
VORANIGO ORAL TABLET 10 MG,	517	INJECTOR 168 MCG/0.56 ML, 294	
40 MG	318	MCG/0.98 ML, 420 MCG/1.4 ML	342
voriconazole intravenous		YUFLYMA(CF)	
VOSEVI		YUFLYMA(CF) AI CROHN'S-UC-HS.	
VOWST		YUFLYMA(CF) AUTOINJECTOR	
VRAYLAR ORAL CAPSULE		zaleplon oral capsule 10 mg, 5 mg	
VUMERITY		ZAVZPRET	
VYNDAQEL		ZEJULA ORAL TABLET	
WELIREG		ZELBORAF	
XALKORI ORAL CAPSULE	_	ZELSUVMI	
XALKORI ORAL PELLET 150 MG, 20		ZEPOSIA	
MG, 50 MG		ZEPOSIA STARTER KIT (28-DAY)	
XCOPRI		ZEPOSIA STARTER PACK (7-DAY)	
XCOPRI MAINTENANCE PACK		ZILBRYSQ SUBCUTANEOUS	
XCOPRI TITRATION PACK		SYRINGE 16.6 MG/0.416 ML, 23	
XDEMVY		MG/0.574 ML, 32.4 MG/0.81 ML	180
XELJANZ ORAL SOLUTION		ZOLINZA	
XELJANZ ORAL TABLET		zolpidem oral tablet	
XELJANZ XR		ZONISADE	
XERMELO	331	ZTALMY	
XGEVA		ZURZUVAE ORAL CAPSULE 20 MG,	,
XIFAXAN ORAL TABLET 550 MG	333	25 MG, 30 MG	
XOLAIR	334	ZYDELIG	.353
XOLREMDI	336	ZYKADIA	354
XOSPATA	337		
XPOVIO ORAL TABLET 100			
MG/WEEK (50 MG X 2), 40			
MG/WEEK (10 MG X 4), 40			
MG/WEEK (40 MG X 1), 40MG			
TWICE WEEK (40 MG X 2), 60			
MG/WEEK (60 MG X 1), 60MG			
TWICE WEEK (120 MG/WEEK), 80			
MG/WEEK (40 MG X 2), 80MG			
TWICE WEEK (160 MG/WEEK)	338		
XTANDI ORAL CAPSULE			

Section: Step Therapy

GnRH Agonists

Products Affected

- ELIGARD 22.5 MG (3 MONTH) SUBCUTANEOUS SYRINGE
- ELIGARD 30 MG (4 MONTH) SUBCUTANEOUS SYRINGE
- ELIGARD 45 MG (6 MONTH) SUBCUTANEOUS SYRINGE
- ELIGARD 7.5 MG (1 MONTH) SUBCUTANEOUS SYRINGE
- TRELSTAR 11.25 MG IM SUSPENSION
- TRELSTAR 22.5 MG IM SUSPENSION
- TRELSTAR 3.75 MG IM SUSPENSION

Criteria	Require a trial of Lupron Depot (Step 1 drug) in the last 180 days when being utilized for the same medically accepted indication
----------	---

Herpetic Keratitis

Products Affected

· ZIRGAN 0.15 % EYE GEL

Criteria Require a 1 month trial of generic trifluridine eye drops (Step 1 drug) in the last 90 days	1
--	---

Pulmonary Antiinflammatory

Products Affected

- fluticasone propionate 100 mcg/actuation blister powder for inhalation
- fluticasone propionate 110 mcg/actuation hfa aerosol inhaler
- fluticasone propionate 220 mcg/actuation hfa aerosol inhaler
- fluticasone propionate 250 mcg/actuation blister powder for inhalation
- fluticasone propionate 44 mcg/actuation hfa aerosol inhaler
- fluticasone propionate 50 mcg/actuation blister powder for inhalation

Criteria	Require a 1 month trial of Qvar and Asmanex/Asmanex HFA (Step 1 drugs) in the last 180 days
----------	---

Rho Kinase Inhibitors

Products Affected

• RHOPRESSA 0.02 % EYE DROPS

DROPS

• ROCKLATAN 0.02 %-0.005 % EYE

	Require a 1 month trial of one preferred glaucoma drug (Step 1 drug) in the last 120 days
--	---

Rytary

Products Affected

- RYTARY 23.75 MG-95 MG CAPSULE,EXTENDED RELEASE
- RYTARY 36.25 MG-145 MG CAPSULE,EXTENDED RELEASE
- RYTARY 48.75 MG-195 MG CAPSULE,EXTENDED RELEASE
- RYTARY 61.25 MG-245 MG CAPSULE,EXTENDED RELEASE

Criteria	Require a trial of generic carbidopa/levodopa product (Step 1 drug) in the last 90 days
----------	---

Index of Drugs	
ELIGARD 22.5 MG (3 MONTH)	
SUBCUTANEOUS SYRINGE	1
ELIGARD 30 MG (4 MONTH)	
SUBCUTANEOUS SYRINGE	1
ELIGARD 45 MG (6 MONTH)	
SUBCUTANEOUS SYRINGE	1
ELIGARD 7.5 MG (1 MONTH)	
SUBCUTANEOUS SYRINGE	1
fluticasone propionate 100 mcg/actuation	
blister powder for inhalation	3
fluticasone propionate 110 mcg/actuation	
hfa aerosol inhaler	3
fluticasone propionate 220 mcg/actuation	
hfa aerosol inhaler	3
fluticasone propionate 250 mcg/actuation	
blister powder for inhalation	3
fluticasone propionate 44 mcg/actuation	
hfa aerosol inhaler	3
fluticasone propionate 50 mcg/actuation	
blister powder for inhalation	3
RHOPRESSA 0.02 % EYE DROPS	
ROCKLATAN 0.02 %-0.005 % EYE	
DROPS	4
RYTARY 23.75 MG-95 MG	
CAPSULE, EXTENDED RELEASE	5
RYTARY 36.25 MG-145 MG	
CAPSULE, EXTENDED RELEASE	5
RYTARY 48.75 MG-195 MG	
CAPSULE, EXTENDED RELEASE	5
RYTARY 61.25 MG-245 MG	
CAPSULE, EXTENDED RELEASE	5
TRELSTAR 11.25 MG IM	9
SUSPENSION	1
TRELSTAR 22.5 MG IM SUSPENSION	
TRELSTAR 22.5 MG IM SUSPENSION TRELSTAR 3.75 MG IM SUSPENSION	
ZIRGAN 0.15 % EYE GEL	
LINGAN U.15 % EXE GEL	2