

## **Medicare Part D: Fundamental Formulary 2025**

Please click [here](#).

## **For Medicare Part D: Prior Authorization Criteria**

Please click [here](#).

## **For Medicare Part D: Step Therapy Criteria**

Please click [here](#).

For more recent information or other questions, please contact:

Senior Blue HMO  
Freedom HMO  
Forever Blue PPO  
BlueSaver HMO  
Employer Group PDP

Pharmacy Service at 1-800-329-2792.

For TTY users, 711 National Relay Service, Oct. 1 – March 31, 8 a.m. – 8 p.m. ET, seven days a week, and April 1 – Aug. 30, 8 a.m. – 8 p.m. ET, Monday – Sunday.

Visit **[medicare.highmark.com](https://www.medicare.highmark.com)**.

Formulary ID: 25025 Version: 14

Updated: 4/2025

**Note to existing members:** This Formulary has changed since last year. Please review this document to make sure that it still contains the drugs you take.

When this Drug List (Formulary) refers to “we,” “us,” or “our,” it means Highmark Blue Cross Blue Shield.

When it refers to “plan” or “our plan,” it means Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO.

This document includes a Drug List (formulary) for our plan, which is current as of January 1, 2025. For an updated Drug List (formulary), please contact us. Our contact information, along with the date we last updated the Drug List (formulary), appears on the front and back cover pages.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2026, and from time to time during the year.

## **What is the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO Formulary?**

In this document, we use the terms Drug List and formulary to mean the same thing. A formulary is a list of covered drugs selected by our plan in consultation with a team of health care providers, which represents the prescription therapies believed to be a necessary part of a quality treatment program. Our plan will generally cover the drugs listed in our formulary as long as the drug is medically necessary, the prescription is filled at one of our plan’s network pharmacies, and other plan rules are followed. For more information on how to fill your prescriptions, please review your Evidence of Coverage.

## Can the formulary change?

Most changes in drug coverage happen on January 1, but we may add or remove drugs on the formulary during the year, move them to different cost-sharing tiers, or add new restrictions. We must follow the Medicare rules in making these changes. Updates to the formulary are posted monthly to our website here: [medicare.highmark.com/formulary](https://www.medicare.highmark.com/formulary).

**Changes that can affect you this year:** In the below cases, you will be affected by coverage changes during the year:

- **Immediate substitutions of certain new versions of brand name drugs and original biological products.** We may immediately remove a drug from our formulary if we are replacing it with a certain new version of that drug that will appear on the same or lower cost-sharing tier and with the same or fewer restrictions. When we add a new version of a drug to our formulary, we may decide to keep the brand name drug or original biological product on our formulary, but immediately move it to a different cost-sharing tier or add new restrictions.

We can make these immediate changes only if we are adding a new generic version of a brand name drug, or adding certain new biosimilar versions of an original biological product, that was already on the formulary (for example, adding an interchangeable biosimilar that can be substituted for an original biological product by a pharmacy without a new prescription.)

If you are currently taking the brand name drug or original biological product, we may not tell you in advance before we make an immediate change, but we will later provide you with information about the specific change(s) we have made.

If we make such a change, you or your prescriber can ask us to make an exception and continue to cover for you the drug that is being changed. For more information, see the section below titled “How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO’s Formulary?”

Some of these drug types may be new to you. For more information, see the section below titled “What are original biological products and how are they related to biosimilars?”

- **Drugs removed from the market.** If a drug is withdrawn from sale by the manufacturer or the Food and Drug Administration (FDA) determines to be withdrawn for safety or effectiveness reasons, we may immediately remove the drug from our formulary and later provide notice to members who take the drug.
- **Other changes.** We may make other changes that affect members currently taking a drug. For instance, we may remove a brand name drug from the formulary when adding a generic equivalent or remove an original biological product when adding a biosimilar. We may also apply new restrictions to the brand name drug or original biological product, or move it to a different cost-sharing tier, or both. We may make changes based on new clinical guidelines. If we remove drugs from our formulary, add prior authorization, quantity

limits and/or step therapy restrictions on a drug, or move a drug to a higher cost-sharing tier, we must notify affected members of the change at least 30 days before the change becomes effective. Alternatively, when a member requests a refill of the drug, they may receive a 31-day supply of the drug and notice of the change.

If we make these other changes, you or your prescriber can ask us to make an exception for you and continue to cover the drug you have been taking. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled “How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO’s Formulary?”

**Changes that will not affect you if you are currently taking the drug.** Generally, if you are taking a drug on our 2025 formulary that was covered at the beginning of the year, we will not discontinue or reduce coverage of the drug during the 2025 coverage year except as described above. This means these drugs will remain available at the same cost-sharing and with no new restrictions for those members taking them for the remainder of the coverage year. You will not get direct notice this year about changes that do not affect you. However, on January 1 of the next year, such changes would affect you, and it is important to check the formulary for the new benefit year for any changes to drugs.

The enclosed formulary is current as of April 1, 2025. To get updated information about the drugs covered by our plan, please contact us. Our contact information appears on the front and back cover pages. In the event of mid-year non-maintenance formulary changes, members will be notified by mail and prospective members will receive an update with this formulary. The most up-to-date formulary is available on our website at [medicare.highmark.com/formulary](https://www.medicare.highmark.com/formulary).

## How do I use the formulary?

There are two ways to find your drug within the formulary:

### Medical Condition

The formulary begins on page 9. The drugs in this formulary are grouped into categories depending on the type of medical conditions that they are used to treat. For example, drugs used to treat a heart condition are listed under the category, “Cardiovascular – Hypertension & Lipids.” If you know what your drug is used for, look for the category name in the list that begins on page 9. Then look under the category name for your drug.

### Alphabetical Listing

If you are not sure what category to look under, you should look for your drug in the Index that begins on page 9. The Index provides an alphabetical list of all of the drugs included in this document. Both brand name drugs and generic drugs are listed in the Index. Look in the Index and find your drug. Next to your drug, you will see the page number where you can find coverage information. Turn to the page listed in the Index and find the name of your drug in the first column of the list.

## What are generic drugs?

Our plan covers both brand name drugs and generic drugs. A generic drug is approved by the FDA as having the same active ingredient as the brand name drug. Generally, generic drugs work just as well as and usually cost less than brand name drugs. There are generic drug substitutes available for many brand name drugs. Generic drugs usually can be substituted for the brand name drug at the pharmacy without needing a new prescription, depending on state laws.

## What are original biological products and how are they related to biosimilars?

On the formulary, when we refer to drugs, this could mean a drug or a biological product. Biological products are drugs that are more complex than typical drugs. Since biological products are more complex than typical drugs, instead of having a generic form, they have alternatives that are called biosimilars. Generally, biosimilars work just as well as the original biological product and may cost less. There are biosimilar alternatives for some original biological products. Some biosimilars are interchangeable biosimilars and, depending on state laws, may be substituted for the original biological product at the pharmacy without needing a new prescription, just like generic drugs can be substituted for brand name drugs.

- For discussion of drug types, please see the Evidence of Coverage, Chapters 5 for MA-PD insert and 3 for PDP insert, Section 3.1, “The ‘Drug List’ tells which Part D drugs are covered.”

## Are there any restrictions on my coverage?

Some covered drugs may have additional requirements or limits on coverage. These requirements and limits may include:

- **Prior Authorization:** Our plan requires you or your prescriber to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don’t get approval, our plan may not cover the drug.
- **Quantity Limits:** For certain drugs, our plan limits the amount of the drug that we will cover. For example, our plan provides 31 tablets, per 31 days, per prescription for 100mg Losartan. This may be in addition to a standard one-month or three-month supply.
- **Step Therapy:** In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. For example, if Drug A and Drug B both treat your medical condition, our plan may not cover Drug B unless you try Drug A first. If Drug A does not work for you, our plan will then cover Drug B.

You can find out if your drug has any additional requirements or limits by looking in the formulary that begins on page 9. You can also get more information about the restrictions applied to specific covered drugs by visiting our website. We have posted online document(s) that

explain(s) our prior authorization and step therapy restrictions. You may also ask us to send you a copy. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You can ask our plan to make an exception to these restrictions or limits or for a list of other, similar drugs that may treat your health condition. See the section, “How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO’s formulary?” on page 6 for information about how to request an exception.

## **What if my drug is not on the formulary?**

If your drug is not included in this formulary (list of covered drugs), you should first contact Pharmacy Service and ask if your drug is covered.

If you learn that our plan does not cover your drug, you have two options:

- You can ask Pharmacy Service for a list of similar drugs that are covered by our plan. When you receive the list, show it to your doctor and ask them to prescribe a similar drug that is covered by our plan.
- You can ask our plan to make an exception and cover your drug. See below for information about how to request an exception.

## **How do I request an exception to the Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO’s formulary?**

You can ask our plan to make an exception to our coverage rules. There are several types of exceptions that you can ask us to make.

- You can ask us to cover a drug even if it is not on our formulary. If approved, this drug will be covered at a pre-determined cost-sharing level, and you would not be able to ask us to provide the drug at a lower cost-sharing level.
- You can ask us to waive a coverage restriction including prior authorization, step therapy, or a quantity limit on your drug. For example, for certain drugs, our plan limits the amount of the drug that we will cover. If your drug has a quantity limit, you can ask us to waive the limit and cover a greater amount.
- You can ask us to cover a formulary drug at a lower cost-sharing level unless the drug is on the specialty tier. If approved, this would lower the amount you must pay for your drug.

Generally, our plan will only approve your request for an exception if the alternative drugs included on the plan’s formulary, the lower cost-sharing drug, or applying the restriction, would not be as effective for you and/or would cause you to have adverse effects.

You or your prescriber should contact us to ask for a tiering or, formulary exception, including an exception to a coverage restriction. **When you request an exception, your prescriber will need to explain the medical reasons why you need the exception.** Generally, we must make our decision within 72 hours of getting your prescriber's supporting statement. You can ask for an expedited (fast) decision if you believe, and we agree, that your health could be seriously harmed by waiting up to 72 hours for a decision. If we agree, or if your prescriber asks for a fast decision, we must give you a decision no later than 24 hours after we get your prescriber's supporting statement.

## **What can I do if my drug is not on the formulary or has a restriction?**

As a new or continuing member in our plan you may be taking drugs that are not on our formulary. Or, you may be taking a drug that is on our formulary but has a coverage restriction, such as prior authorization. You should talk to your prescriber about requesting a coverage decision to show that you meet the criteria for approval, switching to an alternative drug that we cover, or requesting a formulary exception so that we will cover the drug you take. While you and your doctor determine the right course of action for you, we may cover your drug in certain cases during the first 90 days you are a member of our plan.

For each of your drugs that is not on our formulary or has a coverage restriction, we will cover a temporary 31-day supply. If your prescription is written for fewer days, we'll allow refills to provide up to a maximum 31-day supply of medication. If your coverage is not approved after your first 31-day supply, we will not pay for these drugs, even if you have been a member of the plan less than 90 days.

If you are a resident of a long-term care facility and you need a drug that is not on our formulary, or if your ability to get your drugs is limited but you are past the first 90 days of membership in our plan, we will cover a 31-day emergency supply of that drug while you pursue a formulary exception.

The above transition process will be implemented to accommodate you if you have an immediate need for a non-formulary drug or a drug that requires prior authorization due to a change in your level of care while you are waiting for an exception request to be processed.

## **For more information**

For more detailed information about your plan's prescription drug coverage, please review your Evidence of Coverage and other plan materials.

If you have questions about our plan, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

If you have general questions about Medicare prescription drug coverage, please call Medicare at 1-800-MEDICARE (1-800-633-4227) 24 hours a day, seven days a week. TTY users should call 1-877-486-2048, or visit <http://www.medicare.gov>.

## Senior Blue HMO, BlueSaver HMO, Forever Blue PPO, Employer Group PDP, or Freedom PPO formulary

The formulary that begins on the next page provides coverage information about the drugs covered by your plan. If you have trouble finding your drug in the list, turn to the Index that begins on page 9.

The first column of the chart lists the drug name. Brand name drugs are capitalized (e.g., ABELCET) and generic drugs are listed in lower-case italics (e.g., *abacavir*).

The information in the Requirements/Limits column tells you if your plan has any special requirements for coverage of your drug.

*The following is a Formulary Format Example Only:*

Drug Name	Fundamental Drug Tier	Requirements/Limits
Anti-Infectives		
XYZ DRUG	NF	QL-28



## Table of Contents

Anti - Infectives.....	3
Antineoplastic / Immunosuppressant Drugs.....	11
Autonomic / Cns Drugs, Neurology / Psych.....	19
Cardiovascular, Hypertension / Lipids.....	32
Dermatologicals/Topical Therapy.....	38
Diagnostics / Miscellaneous Agents.....	42
Ear, Nose / Throat Medications.....	44
Endocrine/Diabetes.....	44
Gastroenterology.....	48
Immunology, Vaccines / Biotechnology.....	51
Miscellaneous Supplies.....	53
Musculoskeletal / Rheumatology.....	53
Obstetrics / Gynecology.....	55
Ophthalmology.....	58
Respiratory And Allergy.....	60
Urologicals.....	64
Vitamins, Hematinics / Electrolytes.....	65



<p><b>lowercase italics</b> = Generic drugs  <b>UPPERCASE BOLD</b> = Brand name drugs</p>	<p><b>Drug Tier</b>  <b>T1</b> = Cost-Sharing Tier 1 includes preferred generic drugs. This is the lowest cost-sharing tier.  <b>T2</b> = Cost-Sharing Tier 2 includes generic drugs.  <b>T3</b> = Cost-Sharing Tier 3 includes preferred brand name drugs and may include some single-sourced drugs (those generic drugs made by a single manufacturer).  <b>T4</b> = Cost-Sharing Tier 4 includes non-preferred brand name drugs and may include some single-sourced generic drugs (those generic drugs made by a single manufacturer).  <b>T5</b> = Cost-Sharing Tier 5 includes specialty drugs. This is the highest cost-sharing tier.</p>	<p><b>Requirements/Limits</b>  <b>LA</b> = Limited access  <b>PA</b> = Prior authorization required  <b>PA-BvD</b> = This drug may be covered under Medicare part B or D depending on the circumstance. Information may need to be submitted describing the use and setting of the drug to make the determination.  <b>PA-NS</b> = Prior authorization required for new starts only  <b>QL</b> = Quantity limit applies. The quantity limit is noted for each drug. For example, if the quantity limit is QL (90 EA per 180 days), the quantity limit would be 90 units per 180-day supply.  <b>ST</b> = Step therapy applies  <b>ST-NS</b> = Step therapy applies to new starts only</p>
-----------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Drug Name	Drug Tier	Requirements/Limits
<b>Anti - Infectives</b>		
<i>abacavir</i>	T2	
<i>abacavir-lamivudine</i>	T2	
<i>acyclovir oral capsule</i>	T2	
<i>acyclovir oral suspension 200 mg/5 ml</i>	T2	
<i>acyclovir oral tablet</i>	T2	
<i>acyclovir sodium intravenous solution</i>	T4	PA-BvD
<i>adefovir</i>	T4	
<i>albendazole</i>	T4	
<i>amantadine hcl oral capsule</i>	T2	QL (124 EA per 31 days)
<i>amantadine hcl oral solution</i>	T2	
<i>amantadine hcl oral tablet</i>	T2	
<i>amikacin injection solution 500 mg/2 ml</i>	T4	
<i>amoxicillin oral capsule</i>	T2	
<i>amoxicillin oral suspension for reconstitution</i>	T2	
<i>amoxicillin oral tablet</i>	T2	
<i>amoxicillin oral tablet, chewable 125 mg, 250 mg</i>	T2	
<i>amoxicillin-pot clavulanate oral suspension for reconstitution</i>	T2	
<i>amoxicillin-pot clavulanate oral tablet</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>amoxicillin-pot clavulanate oral tablet, chewable 400-57 mg</i>	T2	
<i>amphotericin b</i>	T4	PA-BvD
<i>amphotericin b liposome</i>	T5	PA-BvD
<i>ampicillin oral capsule 500 mg</i>	T2	
<i>ampicillin sodium injection recon soln 1 gram, 10 gram, 125 mg</i>	T4	
<i>ampicillin-sulbactam injection</i>	T4	
<b>APTIVUS</b>	T5	
<b>ARIKAYCE</b>	T5	PA
<i>atazanavir</i>	T4	
<i>atovaquone</i>	T4	
<i>atovaquone-proguanil</i>	T3	
<i>azithromycin intravenous</i>	T4	
<i>azithromycin oral tablet</i>	T2	
<i>aztreonam injection recon soln 1 gram</i>	T4	
<i>aztreonam injection recon soln 2 gram</i>	T5	
<b>BICILLIN C-R</b>	T3	
<b>BICILLIN L-A INTRAMUSCULAR SYRINGE 1,200,000 UNIT/2 ML, 600,000 UNIT/ML</b>	T4	
<b>BIKTARVY</b>	T5	QL (31 EA per 31 days)
<i>caspofungin</i>	T4	
<b>CAYSTON</b>	T5	PA
<i>cefaclor oral capsule 500 mg</i>	T2	
<i>cefadroxil oral capsule</i>	T2	
<i>cefadroxil oral suspension for reconstitution 250 mg/5 ml, 500 mg/5 ml</i>	T2	
<i>cefadroxil oral tablet</i>	T2	
<i>cefazolin injection recon soln 1 gram, 10 gram, 500 mg</i>	T4	
<i>cefdinir oral capsule</i>	T2	
<i>cefepime injection</i>	T4	
<i>cefixime oral capsule</i>	T4	
<i>cefoxitin</i>	T4	
<i>cefpodoxime</i>	T2	
<i>cefprozil</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>ceftazidime</i>	T4	
<i>ceftriaxone injection recon soln 1 gram, 10 gram, 2 gram, 250 mg, 500 mg</i>	T2	
<i>cefuroxime axetil oral tablet</i>	T2	
<i>cefuroxime sodium injection recon soln 750 mg</i>	T4	
<i>cefuroxime sodium intravenous recon soln 1.5 gram</i>	T4	
<i>cephalexin oral capsule 250 mg, 500 mg</i>	T2	
<i>cephalexin oral suspension for reconstitution</i>	T2	
<i>chloroquine phosphate oral tablet 250 mg</i>	T3	QL (50 EA per 30 days)
<i>chloroquine phosphate oral tablet 500 mg</i>	T3	QL (25 EA per 30 days)
<b>CIMDUO</b>	T5	QL (31 EA per 31 days)
<i>ciprofloxacin hcl oral tablet 250 mg, 500 mg</i>	T1	
<i>ciprofloxacin hcl oral tablet 750 mg</i>	T2	
<i>ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml</i>	T4	
<i>clarithromycin</i>	T2	
<i>clindamycin hcl</i>	T2	
<i>clindamycin in 5 % dextrose</i>	T4	
<b>CLINDAMYCIN PEDIATRIC</b>	T2	
<i>clindamycin phosphate injection</i>	T4	
<i>clotrimazole mucous membrane</i>	T2	
<b>COARTEM</b>	T4	
<i>colistin (colistimethate na)</i>	T4	
<b>COMPLERA</b>	T5	
<i>dapsone oral</i>	T3	
<i>daptomycin</i>	T4	
<i>darunavir</i>	T5	
<b>DELSTRIGO</b>	T5	QL (31 EA per 31 days)
<b>DESCOVY</b>	T5	QL (31 EA per 31 days)
<i>dicloxacillin</i>	T2	
<b>DIFICID ORAL TABLET</b>	T5	QL (20 EA per 10 days)
<b>DOVATO</b>	T5	QL (31 EA per 31 days)
<b>DOXY-100</b>	T4	
<i>doxycycline hyclate oral capsule</i>	T2	
<i>doxycycline hyclate oral tablet 100 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>doxycycline hyclate oral tablet, delayed release (dr/ec) 100 mg</i>	T4	
<i>doxycycline monohydrate oral capsule 100 mg, 50 mg</i>	T2	
<i>doxycycline monohydrate oral capsule, ir - delay rel, biphasic</i>	T4	
<i>doxycycline monohydrate oral tablet 100 mg, 50 mg</i>	T2	
<b>EDURANT</b>	T5	
<i>efavirenz oral tablet</i>	T4	
<i>efavirenz-emtricitabin-tenofovir</i>	T5	
<i>efavirenz-lamivudine-tenofovir disoproxil fumarate</i>	T5	QL (31 EA per 31 days)
<i>emtricitabine</i>	T4	
<i>emtricitabine-tenofovir (tdf) oral tablet 100-150 mg, 133-200 mg, 167-250 mg</i>	T5	
<i>emtricitabine-tenofovir (tdf) oral tablet 200-300 mg</i>	T4	
<b>EMTRIVA ORAL SOLUTION</b>	T3	
<b>EMVERM</b>	T5	
<i>entecavir</i>	T4	
<b>EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG</b>	T5	PA; QL (28 EA per 28 days)
<i>ertapenem</i>	T4	
<b>ERY-TAB ORAL TABLET, DELAYED RELEASE (DR/EC) 250 MG, 333 MG</b>	T4	
<i>erythromycin ethylsuccinate oral tablet</i>	T4	
<i>erythromycin oral tablet</i>	T4	
<i>ethambutol</i>	T2	
<i>etravirine</i>	T5	
<b>EVOTAZ</b>	T5	
<i>famciclovir</i>	T2	
<i>fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml, 400 mg/200 ml</i>	T4	
<i>fluconazole oral suspension for reconstitution</i>	T3	
<i>fluconazole oral tablet</i>	T2	
<i>flucytosine oral capsule 250 mg</i>	T4	
<i>flucytosine oral capsule 500 mg</i>	T5	
<i>fosamprenavir</i>	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>FUZEON SUBCUTANEOUS RECON SOLN</b>	T5	
<i>gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml</i>	T4	
<i>gentamicin injection solution 40 mg/ml</i>	T4	
<b>GENVOYA</b>	T5	
<i>griseofulvin microsize</i>	T4	
<i>griseofulvin ultramicrosize oral tablet 125 mg, 250 mg</i>	T4	
<i>hydroxychloroquine oral tablet 200 mg</i>	T2	QL (93 EA per 31 days)
<i>imipenem-cilastatin</i>	T4	
<b>INTELENCE ORAL TABLET 25 MG</b>	T4	
<b>ISENTRESS HD</b>	T5	
<b>ISENTRESS ORAL POWDER IN PACKET</b>	T5	
<b>ISENTRESS ORAL TABLET</b>	T5	
<b>ISENTRESS ORAL TABLET,CHEWABLE 100 MG</b>	T5	
<b>ISENTRESS ORAL TABLET,CHEWABLE 25 MG</b>	T3	
<i>isoniazid oral</i>	T2	
<i>itraconazole oral capsule</i>	T4	PA
<i>ivermectin oral tablet 3 mg</i>	T2	PA
<b>JULUCA</b>	T5	
<i>ketoconazole oral</i>	T2	
<i>lamivudine</i>	T3	
<i>lamivudine-zidovudine</i>	T3	
<i>levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml</i>	T3	
<i>levofloxacin oral</i>	T2	
<i>linezolid in dextrose 5%</i>	T4	
<i>linezolid oral tablet</i>	T4	
<b>LIVTENCITY</b>	T5	PA; QL (372 EA per 31 days)
<i>lopinavir-ritonavir</i>	T4	
<i>maraviroc oral tablet 150 mg</i>	T5	
<i>maraviroc oral tablet 300 mg</i>	T4	
<b>MAVYRET ORAL PELLETS IN PACKET</b>	T5	PA; QL (140 EA per 28 days)
<b>MAVYRET ORAL TABLET</b>	T5	PA; QL (84 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>mefloquine</i>	T2	
<i>meropenem intravenous recon soln 1 gram, 500 mg</i>	T3	
<i>methenamine hippurate</i>	T2	
<i>metronidazole in nacl (iso-os)</i>	T4	
<i>metronidazole oral tablet 250 mg, 500 mg</i>	T2	
<i>micafungin</i>	T4	
<i>minocycline oral capsule</i>	T2	
<i>minocycline oral tablet</i>	T4	
<i>moxifloxacin oral</i>	T2	
<i>moxifloxacin-sod.chloride(iso)</i>	T4	
<i>nafcillin injection</i>	T4	
<i>neomycin</i>	T2	
<i>nevirapine oral suspension</i>	T4	
<i>nevirapine oral tablet</i>	T3	
<i>nevirapine oral tablet extended release 24 hr 400 mg</i>	T4	
<i>nitazoxanide</i>	T5	
<i>nitrofurantoin macrocrystal oral capsule 100 mg</i>	T2	QL (90 EA per 365 days)
<i>nitrofurantoin macrocrystal oral capsule 50 mg</i>	T2	QL (180 EA per 365 days)
<i>nitrofurantoin monohyd/m-cryst</i>	T2	QL (90 EA per 365 days)
<i>nitrofurantoin oral suspension 25 mg/5 ml</i>	T5	QL (1800 ML per 365 days)
<b>NORVIR ORAL POWDER IN PACKET</b>	T4	
<i>nystatin oral</i>	T2	
<b>ODEFSEY</b>	T5	QL (31 EA per 31 days)
<i>ofloxacin oral tablet 300 mg, 400 mg</i>	T4	
<i>oseltamivir oral capsule 30 mg</i>	T2	QL (170 EA per 365 days)
<i>oseltamivir oral capsule 45 mg, 75 mg</i>	T2	QL (90 EA per 365 days)
<i>oseltamivir oral suspension for reconstitution</i>	T3	QL (1080 ML per 365 days)
<i>oxacillin injection recon soln 1 gram</i>	T4	
<i>oxacillin injection recon soln 2 gram</i>	T5	
<b>PAXLOVID ORAL TABLETS,DOSE PACK 150-100 MG</b>	T3	QL (180 EA per 365 days)
<b>PAXLOVID ORAL TABLETS,DOSE PACK 300 MG (150 MG X 2)-100 MG</b>	T3	QL (270 EA per 365 days)
<i>penicillin g pot in dextrose intravenous piggyback 2 million unit/50 ml, 3 million unit/50 ml</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>penicillin g potassium injection recon soln 20 million unit</i>	T4	
<i>penicillin v potassium</i>	T2	
<i>pentamidine inhalation</i>	T4	PA-BvD
<i>pentamidine injection</i>	T4	
<b>PIFELTRO</b>	T5	QL (62 EA per 31 days)
<i>piperacillin-tazobactam intravenous recon soln 2.25 gram, 3.375 gram, 4.5 gram, 40.5 gram</i>	T4	
<i>posaconazole oral tablet, delayed release (dr/ec)</i>	T5	PA
<i>praziquantel</i>	T4	
<b>PREVYMIS ORAL PELLETS IN PACKET</b>	T5	PA; QL (124 EA per 31 days)
<b>PREVYMIS ORAL TABLET</b>	T5	QL (31 EA per 31 days)
<b>PREZCOBIX</b>	T5	
<b>PREZISTA ORAL SUSPENSION</b>	T5	
<b>PREZISTA ORAL TABLET 150 MG, 75 MG</b>	T5	
<b>PRIFTIN</b>	T3	
<i>primaquine</i>	T3	
<i>pyrazinamide</i>	T4	
<i>pyrimethamine</i>	T5	PA
<i>quinine sulfate</i>	T4	PA; QL (42 EA per 28 days)
<b>RELENZA DISKHALER</b>	T3	
<b>REYATAZ ORAL POWDER IN PACKET</b>	T5	
<i>ribavirin oral capsule</i>	T3	
<i>ribavirin oral tablet 200 mg</i>	T3	
<i>rifabutin</i>	T4	
<i>rifampin intravenous</i>	T5	
<i>rifampin oral</i>	T3	
<i>rimantadine</i>	T2	
<i>ritonavir</i>	T3	
<b>RUKOBIA</b>	T5	QL (62 EA per 31 days)
<b>SELZENTRY ORAL SOLUTION</b>	T5	
<b>SIRTURO</b>	T5	PA
<i>sofosbuvir-velpatasvir</i>	T5	PA; QL (28 EA per 28 days)
<i>streptomycin</i>	T5	
<b>STRIBILD</b>	T5	
<i>sulfadiazine</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>sulfamethoxazole-trimethoprim oral suspension</i>	T2	
<i>sulfamethoxazole-trimethoprim oral tablet</i>	T1	
<b>SUNLENCA ORAL</b>	T5	
<b>SYMTUZA</b>	T5	QL (31 EA per 31 days)
<b>TEFLARO</b>	T5	
<i>tenofovir disoproxil fumarate</i>	T3	
<i>terbinafine hcl oral</i>	T2	QL (90 EA per 180 days)
<i>tetracycline oral capsule</i>	T4	
<i>tigecycline</i>	T5	
<i>tinidazole</i>	T2	
<b>TIVICAY ORAL TABLET 50 MG</b>	T5	
<b>TIVICAY PD</b>	T5	
<b>TOBI PODHALER</b>	T5	PA; QL (224 EA per 56 days)
<i>tobramycin in 0.225 % nacl</i>	T5	PA
<i>tobramycin inhalation</i>	T5	PA
<i>tobramycin sulfate injection solution</i>	T4	
<b>TRECATOR</b>	T4	
<i>trimethoprim</i>	T2	
<b>TRIUMEQ</b>	T5	
<b>TRIUMEQ PD</b>	T4	QL (186 EA per 31 days)
<b>TYBOST</b>	T3	
<i>valacyclovir</i>	T2	
<i>valganciclovir oral recon soln</i>	T4	
<i>valganciclovir oral tablet</i>	T3	
<i>vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg</i>	T4	
<i>vancomycin oral capsule 125 mg</i>	T4	PA; QL (124 EA per 31 days)
<i>vancomycin oral capsule 250 mg</i>	T4	PA; QL (248 EA per 31 days)
<b>VEMLIDY</b>	T5	QL (31 EA per 31 days)
<b>VIRACEPT ORAL TABLET</b>	T5	
<b>VIREAD ORAL POWDER</b>	T5	
<b>VIREAD ORAL TABLET 150 MG, 200 MG, 250 MG</b>	T5	
<b>VIVJOA</b>	T4	PA; QL (18 EA per 84 days)
<i>voriconazole intravenous</i>	T5	PA
<i>voriconazole oral</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>VOSEVI</b>	T5	PA; QL (28 EA per 28 days)
<b>XIFAXAN ORAL TABLET 200 MG</b>	T4	QL (27 EA per 365 days)
<b>XIFAXAN ORAL TABLET 550 MG</b>	T5	PA; QL (62 EA per 31 days)
<b>XOFLUZA ORAL TABLET 40 MG, 80 MG</b>	T3	QL (9 EA per 365 days)
<i>zidovudine</i>	T2	
<b>Antineoplastic / Immunosuppressant Drugs</b>		
<i>abiraterone oral tablet 250 mg</i>	T5	PA-NS; QL (124 EA per 31 days)
<i>abiraterone oral tablet 500 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<b>AKEEGA</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>ALECENSA</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>ALUNBRIG ORAL TABLET 180 MG, 90 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ALUNBRIG ORAL TABLET 30 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>ALUNBRIG ORAL TABLETS,DOSE PACK</b>	T5	PA-NS; QL (60 EA per 365 days)
<i>anastrozole</i>	T2	
<b>AUGTYRO ORAL CAPSULE 160 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>AUGTYRO ORAL CAPSULE 40 MG</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>AYVAKIT</b>	T5	PA-NS; QL (31 EA per 31 days)
<i>azathioprine oral tablet 50 mg</i>	T2	PA-BvD
<b>BALVERSA</b>	T5	PA-NS
<i>bexarotene oral</i>	T5	PA-NS
<i>bexarotene topical</i>	T5	PA-NS; QL (60 GM per 28 days)
<i>bicalutamide</i>	T2	
<b>BOSULIF ORAL CAPSULE 100 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>BOSULIF ORAL CAPSULE 50 MG</b>	T5	PA-NS; QL (341 EA per 31 days)
<b>BOSULIF ORAL TABLET 100 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>BOSULIF ORAL TABLET 400 MG, 500 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>BRAFTOVI</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>BRUKINSA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>CABOMETYX</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>CALQUENCE</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>CALQUENCE (ACALABRUTINIB MAL)</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>CAPRELSA ORAL TABLET 100 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>CAPRELSA ORAL TABLET 300 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1)</b>	T5	PA-NS; QL (56 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>COMETRIQ ORAL CAPSULE 140 MG/DAY(80 MG X1-20 MG X3)</b>	T5	PA-NS; QL (112 EA per 28 days)
<b>COMETRIQ ORAL CAPSULE 60 MG/DAY (20 MG X 3/DAY)</b>	T5	PA-NS; QL (84 EA per 28 days)
<b>COPIKTRA</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>COTELLIC</b>	T5	PA-NS; LA; QL (63 EA per 28 days)
<i>cyclophosphamide oral</i>	T3	PA-BvD
<i>cyclosporine modified oral capsule</i>	T2	PA-BvD
<i>cyclosporine modified oral solution</i>	T4	PA-BvD
<i>cyclosporine oral capsule</i>	T2	PA-BvD
<b>DANZITEN</b>	T5	PA-NS; QL (124 EA per 31 days)
<i>dasatinib</i>	T5	PA-NS; QL (31 EA per 31 days)
<b>DAURISMO ORAL TABLET 100 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>DAURISMO ORAL TABLET 25 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>ELIGARD</b>	T4	ST-NS; QL (1 EA per 30 days)
<b>ELIGARD (3 MONTH)</b>	T4	ST-NS; QL (1 EA per 90 days)
<b>ELIGARD (4 MONTH)</b>	T4	ST-NS; QL (1 EA per 120 days)
<b>ELIGARD (6 MONTH)</b>	T4	ST-NS; QL (1 EA per 180 days)
<b>ENVARUSUS XR</b>	T4	PA-BvD
<b>ERIVEDGE</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ERLEADA ORAL TABLET 240 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ERLEADA ORAL TABLET 60 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<i>erlotinib</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 7.5 mg</i>	T5	PA-NS; QL (31 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet 5 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 2 mg, 5 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<i>everolimus (antineoplastic) oral tablet for suspension 3 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>everolimus (immunosuppressive)</i>	T5	PA-BvD
<i>exemestane</i>	T4	
<b>FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 120 MG</b>	T5	PA-NS
<b>FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 80 MG</b>	T4	PA-NS
<b>FOTIVDA</b>	T5	PA-NS; QL (21 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>FRUZAQLA ORAL CAPSULE 1 MG</b>	T5	PA-NS; QL (84 EA per 28 days)
<b>FRUZAQLA ORAL CAPSULE 5 MG</b>	T5	PA-NS; QL (21 EA per 28 days)
<b>GAVRETO</b>	T5	PA-NS; QL (124 EA per 31 days)
<i>gefitinib</i>	T5	PA-NS; QL (31 EA per 31 days)
<b>GENGRAF ORAL CAPSULE</b>	T2	PA-BvD
<b>GENGRAF ORAL SOLUTION</b>	T5	PA-BvD
<b>GILOTRIF</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>GLEOSTINE ORAL CAPSULE 10 MG, 100 MG</b>	T5	PA-NS
<b>GLEOSTINE ORAL CAPSULE 40 MG</b>	T4	PA-NS
<i>hydroxyurea</i>	T2	
<b>IBRANCE</b>	T5	PA-NS; QL (21 EA per 28 days)
<b>ICLUSIG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>IDHIFA ORAL TABLET 100 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>IDHIFA ORAL TABLET 50 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<i>imatinib oral tablet 100 mg</i>	T5	PA-NS; QL (93 EA per 31 days)
<i>imatinib oral tablet 400 mg</i>	T5	PA-NS; QL (62 EA per 31 days)
<b>IMBRUVICA ORAL CAPSULE 140 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>IMBRUVICA ORAL CAPSULE 70 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>IMBRUVICA ORAL SUSPENSION</b>	T5	PA-NS; QL (216 ML per 25 days)
<b>IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>IMKELDI</b>	T5	PA-NS; QL (280 ML per 28 days)
<b>INLYTA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>INQOVI</b>	T5	PA-NS; QL (5 EA per 28 days)
<b>INREBIC</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>ITOVEBI ORAL TABLET 3 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>ITOVEBI ORAL TABLET 9 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>IWILFIN</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>JAKAFI</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>JAYPIRCA ORAL TABLET 100 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>JAYPIRCA ORAL TABLET 50 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG</b>	T5	PA-NS; QL (70 EA per 28 days)
<b>KISQALI FEMARA CO-PACK ORAL TABLET 600 MG/DAY(200 MG X 3)-2.5 MG</b>	T5	PA-NS; QL (91 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1)</b>	T5	PA-NS; QL (21 EA per 28 days)
<b>KISQALI ORAL TABLET 400 MG/DAY (200 MG X 2)</b>	T5	PA-NS; QL (42 EA per 28 days)
<b>KISQALI ORAL TABLET 600 MG/DAY (200 MG X 3)</b>	T5	PA-NS; QL (63 EA per 28 days)
<b>KOSELUGO ORAL CAPSULE 10 MG</b>	T5	PA-NS; QL (279 EA per 31 days)
<b>KOSELUGO ORAL CAPSULE 25 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>KRAZATI</b>	T5	PA-NS; QL (186 EA per 31 days)
<i>lapatinib</i>	T5	PA-NS; QL (186 EA per 31 days)
<b>LAZCLUZE ORAL TABLET 240 MG</b>	T5	PA-NS; QL (30 EA per 30 days)
<b>LAZCLUZE ORAL TABLET 80 MG</b>	T5	PA-NS; QL (60 EA per 30 days)
<i>lenalidomide</i>	T5	PA-NS; QL (21 EA per 28 days)
<b>LENVIMA</b>	T5	PA-NS
<i>letrozole</i>	T2	
<i>leucovorin calcium oral</i>	T3	
<i>leuprolide (3 month)</i>	T4	QL (1 EA per 84 days)
<i>leuprolide subcutaneous kit</i>	T3	QL (2 EA per 28 days)
<b>LONSURF</b>	T5	PA-NS
<b>LORBRENA ORAL TABLET 100 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>LORBRENA ORAL TABLET 25 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>LUMAKRAS ORAL TABLET 120 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>LUMAKRAS ORAL TABLET 240 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>LUMAKRAS ORAL TABLET 320 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG</b>	T5	QL (1 EA per 90 days)
<b>LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 22.5 MG</b>	T5	QL (1 EA per 84 days)
<b>LUPRON DEPOT (4 MONTH)</b>	T5	QL (1 EA per 112 days)
<b>LUPRON DEPOT (6 MONTH)</b>	T5	QL (1 EA per 168 days)
<b>LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 3.75 MG</b>	T5	QL (1 EA per 30 days)
<b>LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 7.5 MG</b>	T5	QL (1 EA per 28 days)
<b>LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG</b>	T5	PA; QL (1 EA per 90 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)</b>	T5	PA; QL (1 EA per 30 days)
<b>LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT</b>	T5	PA; QL (1 EA per 168 days)
<b>LYNPARZA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>LYSODREN</b>	T5	
<b>LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3)</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>LYTGOBI ORAL TABLET 16 MG/DAY (4 MG X 4)</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>LYTGOBI ORAL TABLET 20 MG/DAY (4 MG X 5)</b>	T5	PA-NS; QL (155 EA per 31 days)
<b>MATULANE</b>	T5	
<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml)</i>	T3	PA
<i>megestrol oral suspension 625 mg/5 ml (125 mg/ml)</i>	T4	PA
<i>megestrol oral tablet</i>	T3	PA-NS
<b>MEKINIST ORAL RECON SOLN</b>	T5	PA-NS; QL (1260 ML per 31 days)
<b>MEKINIST ORAL TABLET 0.5 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>MEKINIST ORAL TABLET 2 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>MEKTOVI</b>	T5	PA-NS; QL (186 EA per 31 days)
<i>mercaptopurine oral tablet</i>	T2	
<i>mesna oral</i>	T4	
<i>methotrexate sodium</i>	T2	PA-BvD
<i>methotrexate sodium (pf) injection solution</i>	T2	PA-BvD
<i>mycophenolate mofetil oral capsule</i>	T2	PA-BvD
<i>mycophenolate mofetil oral suspension for reconstitution</i>	T4	PA-BvD
<i>mycophenolate mofetil oral tablet</i>	T2	PA-BvD
<i>mycophenolate sodium</i>	T2	PA-BvD
<b>NEMLUVIO</b>	T5	PA; QL (2 EA per 28 days)
<b>NERLYNX</b>	T5	PA-NS; QL (186 EA per 31 days)
<i>nilutamide</i>	T5	
<b>NINLARO</b>	T5	PA-NS; QL (3 EA per 28 days)
<b>NUBEQA</b>	T5	PA-NS; QL (124 EA per 31 days)
<i>octreotide acetate injection solution</i>	T4	PA

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>ODOMZO</b>	T5	PA-NS; LA; QL (31 EA per 31 days)
<b>OGSIVEO ORAL TABLET 100 MG, 150 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>OGSIVEO ORAL TABLET 50 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>OJEMDA ORAL SUSPENSION FOR RECONSTITUTION</b>	T5	PA-NS; QL (96 ML per 28 days)
<b>OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4)</b>	T5	PA-NS; QL (16 EA per 28 days)
<b>OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)</b>	T5	PA-NS; QL (20 EA per 28 days)
<b>OJEMDA ORAL TABLET 600 MG/WEEK (100 MG X 6)</b>	T5	PA-NS; QL (24 EA per 28 days)
<b>OJJAARA</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ONUREG</b>	T5	PA-NS; QL (14 EA per 28 days)
<b>ORGOVYX</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ORSERDU ORAL TABLET 345 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ORSERDU ORAL TABLET 86 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<i>pazopanib</i>	T5	PA-NS; QL (124 EA per 31 days)
<b>PEMAZYRE</b>	T5	PA-NS; QL (14 EA per 21 days)
<b>PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1)</b>	T5	PA-NS; QL (28 EA per 28 days)
<b>PIQRAY ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)</b>	T5	PA-NS; QL (56 EA per 28 days)
<b>POMALYST</b>	T5	PA-NS; QL (21 EA per 28 days)
<b>PROGRAF ORAL GRANULES IN PACKET</b>	T4	PA-BvD
<b>PURIXAN</b>	T5	
<b>QINLOCK</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>RETEVMO ORAL CAPSULE 40 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>RETEVMO ORAL CAPSULE 80 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>RETEVMO ORAL TABLET 120 MG, 160 MG, 80 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>RETEVMO ORAL TABLET 40 MG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>REVUFORJ ORAL TABLET 110 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>REVUFORJ ORAL TABLET 160 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>REZLIDHIA</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>ROZLYTREK ORAL CAPSULE 100 MG</b>	T5	PA-NS; QL (155 EA per 31 days)
<b>ROZLYTREK ORAL CAPSULE 200 MG</b>	T5	PA-NS; QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>ROZLYTREK ORAL PELLETS IN PACKET</b>	T5	PA-NS; QL (372 EA per 31 days)
<b>RUBRACA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>RYDAPT</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>SCEMBLIX ORAL TABLET 100 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>SCEMBLIX ORAL TABLET 20 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>SCEMBLIX ORAL TABLET 40 MG</b>	T5	PA-NS; QL (310 EA per 31 days)
<b>SIGNIFOR</b>	T5	PA
<i>sirolimus oral solution</i>	T5	PA-BvD
<i>sirolimus oral tablet</i>	T4	PA-BvD
<b>SOLTAMOX</b>	T4	
<i>sorafenib</i>	T5	PA-NS; QL (124 EA per 31 days)
<b>STIVARGA</b>	T5	PA-NS; QL (84 EA per 28 days)
<i>sunitinib malate</i>	T5	PA-NS; QL (31 EA per 31 days)
<b>TABRECTA</b>	T5	PA-NS; QL (124 EA per 31 days)
<i>tacrolimus oral capsule</i>	T2	PA-BvD
<b>TAFINLAR ORAL CAPSULE</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>TAFINLAR ORAL TABLET FOR SUSPENSION</b>	T5	PA-NS; QL (930 EA per 31 days)
<b>TAGRISSO</b>	T5	PA-NS; LA; QL (31 EA per 31 days)
<b>TALZENNA</b>	T5	PA-NS; QL (31 EA per 31 days)
<i>tamoxifen</i>	T2	
<b>TASIGNA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>TAZVERIK</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>TEPMETKO</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>THALOMID ORAL CAPSULE 100 MG, 50 MG</b>	T5	PA-NS; QL (28 EA per 28 days)
<b>TIBSOVO</b>	T5	PA-NS; QL (62 EA per 31 days)
<i>toremifene</i>	T4	
<b>TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 11.25 MG</b>	T4	ST-NS; QL (1 EA per 84 days)
<b>TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 22.5 MG</b>	T4	ST-NS; QL (1 EA per 168 days)
<b>TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 3.75 MG</b>	T4	ST-NS; QL (1 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>tretinoin (antineoplastic)</i>	T5	
<b>TRUQAP</b>	T5	PA-NS; QL (64 EA per 28 days)
<b>TUKYSA ORAL TABLET 150 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>TUKYSA ORAL TABLET 50 MG</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>TURALIO ORAL CAPSULE 125 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>VANFLYTA</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>VENCLEXTA ORAL TABLET 10 MG</b>	T3	PA-NS; QL (62 EA per 31 days)
<b>VENCLEXTA ORAL TABLET 100 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>VENCLEXTA ORAL TABLET 50 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>VENCLEXTA STARTING PACK</b>	T5	PA-NS; QL (84 EA per 365 days)
<b>VERZENIO</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>VIJOICE ORAL GRANULES IN PACKET</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>VIJOICE ORAL TABLET 125 MG, 50 MG</b>	T5	PA-NS; QL (28 EA per 28 days)
<b>VIJOICE ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1)</b>	T5	PA-NS; QL (56 EA per 28 days)
<b>VITRAKVI ORAL CAPSULE 100 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>VITRAKVI ORAL CAPSULE 25 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>VITRAKVI ORAL SOLUTION</b>	T5	PA-NS; QL (310 ML per 31 days)
<b>VIZIMPRO</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>VONJO</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>VORANIGO ORAL TABLET 10 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>VORANIGO ORAL TABLET 40 MG</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>WELIREG</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>XALKORI ORAL CAPSULE</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>XALKORI ORAL PELLET 150 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>XALKORI ORAL PELLET 20 MG, 50 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>XATMEP</b>	T4	PA-BvD
<b>XERMELO</b>	T5	PA; QL (93 EA per 31 days)
<b>XGEVA</b>	T5	PA-NS
<b>XOSPATA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40MG TWICE WEEK (40 MG X 2), 80 MG/WEEK (40 MG X 2)</b>	T5	PA-NS; QL (8 EA per 28 days)
<b>XPOVIO ORAL TABLET 40 MG/WEEK (40 MG X 1), 60 MG/WEEK (60 MG X 1)</b>	T5	PA-NS; QL (4 EA per 28 days)
<b>XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK)</b>	T5	PA-NS; QL (24 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<b>XPOVIO ORAL TABLET 80MG TWICE WEEK (160 MG/WEEK)</b>	T5	PA-NS; QL (32 EA per 28 days)
<b>XTANDI ORAL CAPSULE</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>XTANDI ORAL TABLET 40 MG</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>XTANDI ORAL TABLET 80 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>YONSA</b>	T5	PA-NS; QL (124 EA per 31 days)
<b>ZEJULA ORAL TABLET</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>ZELBORAF</b>	T5	PA-NS; QL (248 EA per 31 days)
<b>ZOLINZA</b>	T5	PA-NS
<b>ZYDELIG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>ZYKADIA</b>	T5	PA-NS; QL (93 EA per 31 days)
<b>Autonomic / Cns Drugs, Neurology / Psych</b>		
<b>ABILIFY MAINTENA</b>	T5	QL (1 EA per 28 days)
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i>	T2	PA; QL (5167 ML per 31 days)
<i>acetaminophen-codeine oral tablet</i>	T2	PA; QL (403 EA per 31 days)
<b>AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML</b>	T3	PA; QL (1 ML per 28 days)
<b>AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 70 MG/ML</b>	T3	PA; QL (2 ML per 28 days)
<b>AJOVY AUTOINJECTOR</b>	T3	PA; QL (1.5 ML per 28 days)
<b>AJOVY SYRINGE</b>	T3	PA; QL (1.5 ML per 28 days)
<i>alprazolam oral tablet 0.25 mg, 0.5 mg</i>	T2	PA; QL (93 EA per 31 days)
<i>alprazolam oral tablet 1 mg, 2 mg</i>	T2	PA; QL (155 EA per 31 days)
<i>amitriptyline</i>	T2	PA-NS
<i>amoxapine</i>	T3	
<b>APTIOM ORAL TABLET 200 MG</b>	T4	QL (186 EA per 31 days)
<b>APTIOM ORAL TABLET 400 MG</b>	T5	QL (93 EA per 31 days)
<b>APTIOM ORAL TABLET 600 MG, 800 MG</b>	T5	QL (62 EA per 31 days)
<i>aripiprazole oral solution</i>	T4	PA-NS
<i>aripiprazole oral tablet</i>	T2	
<i>aripiprazole oral tablet, disintegrating</i>	T5	PA-NS
<i>armodafinil</i>	T4	PA; QL (31 EA per 31 days)
<i>asenapine maleate</i>	T4	PA-NS; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>atomoxetine oral capsule 10 mg, 25 mg, 40 mg</i>	T4	QL (62 EA per 31 days)
<i>atomoxetine oral capsule 100 mg, 60 mg, 80 mg</i>	T4	QL (31 EA per 31 days)
<i>atomoxetine oral capsule 18 mg</i>	T4	QL (124 EA per 31 days)
<b>AUVELITY</b>	T5	PA-NS; QL (62 EA per 31 days)
<i>baclofen oral tablet 10 mg, 20 mg, 5 mg</i>	T2	
<b>BAFIERTAM</b>	T5	PA; QL (124 EA per 31 days)
<i>benztropine oral</i>	T1	PA
<b>BRIVIACT ORAL SOLUTION</b>	T5	QL (620 ML per 31 days)
<b>BRIVIACT ORAL TABLET</b>	T5	QL (62 EA per 31 days)
<i>bromocriptine</i>	T4	
<i>buprenorphine</i>	T4	PA; QL (4 EA per 28 days)
<i>buprenorphine hcl sublingual tablet 2 mg</i>	T3	QL (93 EA per 31 days)
<i>buprenorphine hcl sublingual tablet 8 mg</i>	T3	QL (62 EA per 31 days)
<i>buprenorphine-naloxone sublingual film 12-3 mg, 4-1 mg, 8-2 mg</i>	T2	QL (62 EA per 31 days)
<i>buprenorphine-naloxone sublingual film 2-0.5 mg</i>	T2	QL (93 EA per 31 days)
<i>buprenorphine-naloxone sublingual tablet</i>	T4	QL (93 EA per 31 days)
<i>bupropion hcl oral tablet</i>	T2	
<i>bupropion hcl oral tablet extended release 24 hr 150 mg</i>	T2	QL (93 EA per 31 days)
<i>bupropion hcl oral tablet extended release 24 hr 300 mg</i>	T2	QL (31 EA per 31 days)
<i>bupropion hcl oral tablet sustained-release 12 hr</i>	T2	QL (62 EA per 31 days)
<i>buspirone</i>	T2	
<i>butorphanol nasal</i>	T2	QL (5 ML per 28 days)
<b>CAPLYTA</b>	T5	PA-NS; QL (31 EA per 31 days)
<i>carbamazepine oral capsule, er multiphase 12 hr</i>	T2	
<i>carbamazepine oral suspension 100 mg/5 ml</i>	T2	
<i>carbamazepine oral tablet</i>	T2	
<i>carbamazepine oral tablet extended release 12 hr 100 mg, 200 mg</i>	T2	
<i>carbamazepine oral tablet extended release 12 hr 400 mg</i>	T3	
<i>carbamazepine oral tablet, chewable 100 mg</i>	T2	
<i>carbidopa-levodopa</i>	T2	
<i>carbidopa-levodopa-entacapone</i>	T4	
<i>celecoxib oral capsule 100 mg, 200 mg, 50 mg</i>	T2	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>celecoxib oral capsule 400 mg</i>	T3	QL (62 EA per 31 days)
<i>chlorpromazine oral</i>	T4	
<i>citalopram oral solution</i>	T3	
<i>citalopram oral tablet</i>	T1	
<i>clobazam oral suspension</i>	T4	PA-NS; QL (496 ML per 31 days)
<i>clobazam oral tablet</i>	T3	PA-NS; QL (62 EA per 31 days)
<i>clomipramine</i>	T4	PA-NS
<i>clonazepam oral tablet 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>clonazepam oral tablet 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>clonazepam oral tablet,disintegrating 1 mg</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>clonazepam oral tablet,disintegrating 2 mg</i>	T2	PA-NS; QL (310 EA per 31 days)
<i>clorazepate dipotassium oral tablet 15 mg</i>	T3	PA-NS; QL (186 EA per 31 days)
<i>clorazepate dipotassium oral tablet 3.75 mg, 7.5 mg</i>	T3	PA-NS; QL (93 EA per 31 days)
<i>clozapine oral tablet 100 mg, 25 mg</i>	T2	QL (279 EA per 31 days)
<i>clozapine oral tablet 200 mg</i>	T2	QL (124 EA per 31 days)
<i>clozapine oral tablet 50 mg</i>	T2	QL (93 EA per 31 days)
<i>clozapine oral tablet,disintegrating 100 mg, 25 mg</i>	T4	QL (279 EA per 31 days)
<i>clozapine oral tablet,disintegrating 12.5 mg</i>	T4	QL (93 EA per 31 days)
<i>clozapine oral tablet,disintegrating 150 mg</i>	T4	QL (186 EA per 31 days)
<i>clozapine oral tablet,disintegrating 200 mg</i>	T4	QL (124 EA per 31 days)
<b>COBENFY</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>COBENFY STARTER PACK</b>	T5	PA-NS; QL (112 EA per 365 days)
<b>COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML</b>	T5	PA; QL (31 ML per 31 days)
<b>COPAXONE SUBCUTANEOUS SYRINGE 40 MG/ML</b>	T5	PA; QL (12 ML per 28 days)
<i>cyclobenzaprine oral tablet 10 mg</i>	T2	QL (93 EA per 31 days)
<i>cyclobenzaprine oral tablet 5 mg</i>	T2	QL (155 EA per 31 days)
<i>dalfampridine</i>	T5	PA; QL (62 EA per 31 days)
<i>dantrolene oral</i>	T2	
<b>DAYBUE</b>	T5	PA; QL (3600 ML per 30 days)
<i>desipramine</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>desvenlafaxine succinate</i>	T2	QL (31 EA per 31 days)
<i>dexmethylphenidate oral capsule,er biphasic 50-50</i>	T2	QL (31 EA per 31 days)
<i>dexmethylphenidate oral tablet 10 mg</i>	T2	QL (62 EA per 31 days)
<i>dexmethylphenidate oral tablet 2.5 mg, 5 mg</i>	T2	QL (93 EA per 31 days)
<i>dextroamphetamine-amphetamine oral capsule,extended release 24hr</i>	T2	QL (31 EA per 31 days)
<i>dextroamphetamine-amphetamine oral tablet 10 mg, 12.5 mg, 15 mg, 30 mg, 5 mg, 7.5 mg</i>	T2	QL (62 EA per 31 days)
<i>dextroamphetamine-amphetamine oral tablet 20 mg</i>	T2	QL (93 EA per 31 days)
<b>DIACOMIT ORAL CAPSULE 250 MG</b>	T5	PA-NS; QL (341 EA per 31 days)
<b>DIACOMIT ORAL CAPSULE 500 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>DIACOMIT ORAL POWDER IN PACKET 250 MG</b>	T5	PA-NS; QL (341 EA per 31 days)
<b>DIACOMIT ORAL POWDER IN PACKET 500 MG</b>	T5	PA-NS; QL (186 EA per 31 days)
<b>DIAZEPAM INTENSOL</b>	T2	PA-NS; QL (248 ML per 31 days)
<i>diazepam oral solution 5 mg/5 ml (1 mg/ml)</i>	T2	PA-NS; QL (1500 ML per 31 days)
<i>diazepam oral tablet</i>	T2	PA-NS; QL (124 EA per 31 days)
<i>diazepam rectal</i>	T4	
<i>diclofenac potassium oral tablet 50 mg</i>	T2	
<i>diclofenac sodium oral</i>	T2	
<i>diclofenac sodium topical drops</i>	T4	
<i>diflunisal</i>	T2	
<i>dihydroergotamine nasal</i>	T5	PA; QL (8 ML per 28 days)
<b>DILANTIN</b>	T3	
<i>dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg (14)- 240 mg (46)</i>	T5	PA; QL (120 EA per 365 days)
<i>dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 240 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>divalproex</i>	T2	
<i>donepezil oral tablet 10 mg, 5 mg</i>	T1	
<i>donepezil oral tablet 23 mg</i>	T3	QL (31 EA per 31 days)
<i>donepezil oral tablet,disintegrating</i>	T2	
<i>doxepin oral capsule</i>	T2	PA-NS
<i>doxepin oral concentrate</i>	T2	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>doxepin oral tablet</i>	T3	PA
<b>DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG</b>	T4	PA-NS; QL (93 EA per 31 days)
<b>DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 30 MG, 60 MG</b>	T4	PA-NS; QL (62 EA per 31 days)
<b>DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 40 MG</b>	T4	PA-NS; QL (31 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 20 mg, 60 mg</i>	T2	QL (62 EA per 31 days)
<i>duloxetine oral capsule, delayed release(dr/ec) 30 mg</i>	T2	QL (31 EA per 31 days)
<b>EMGALITY PEN</b>	T3	PA; QL (1 ML per 28 days)
<b>EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML</b>	T3	PA; QL (1 ML per 28 days)
<b>EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 300 MG/3 ML (100 MG/ML X 3)</b>	T5	PA; QL (3 ML per 28 days)
<b>EMSAM</b>	T5	QL (30 EA per 30 days)
<b>ENDOCET</b>	T3	PA; QL (372 EA per 31 days)
<i>entacapone</i>	T3	
<b>EPIDIOLEX</b>	T5	PA-NS
<b>EPITOL</b>	T2	
<b>EPRONTIA</b>	T4	PA-NS; QL (496 ML per 31 days)
<i>ergotamine-caffeine</i>	T3	PA
<i>escitalopram oxalate oral solution</i>	T4	QL (620 ML per 31 days)
<i>escitalopram oxalate oral tablet 10 mg</i>	T1	QL (45 EA per 30 days)
<i>escitalopram oxalate oral tablet 20 mg, 5 mg</i>	T1	QL (30 EA per 30 days)
<i>eszopiclone</i>	T4	PA; QL (31 EA per 31 days)
<i>ethosuximide</i>	T2	
<i>etodolac</i>	T2	
<b>EVRYSDI ORAL RECON SOLN</b>	T5	PA; QL (240 ML per 31 days)
<b>FANAPT ORAL TABLET 1 MG</b>	T4	PA-NS; QL (62 EA per 31 days)
<b>FANAPT ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>FANAPT ORAL TABLETS,DOSE PACK</b>	T4	PA-NS; QL (16 EA per 365 days)
<i>felbamate</i>	T4	
<i>fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 75 mcg/hr</i>	T4	PA; QL (10 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>fentanyl transdermal patch 72 hour 25 mcg/hr, 50 mcg/hr</i>	T2	PA; QL (10 EA per 30 days)
<b>FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)</b>	T3	PA-NS; QL (56 EA per 365 days)
<b>FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 40 MG, 80 MG</b>	T3	PA-NS; QL (31 EA per 31 days)
<b>FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 20 MG</b>	T3	PA-NS; QL (93 EA per 31 days)
<i>fingolimod</i>	T5	PA; QL (31 EA per 31 days)
<b>FINTEPLA</b>	T5	PA-NS; QL (360 ML per 30 days)
<b>FIRDAPSE</b>	T5	PA; QL (248 EA per 31 days)
<i>fluoxetine (pmd)</i>	T2	
<i>fluoxetine oral capsule</i>	T1	
<i>fluoxetine oral solution</i>	T3	
<i>fluoxetine oral tablet 10 mg, 20 mg</i>	T2	
<i>fluphenazine decanoate</i>	T2	
<i>fluphenazine hcl injection</i>	T4	
<i>fluphenazine hcl oral concentrate</i>	T4	
<i>fluphenazine hcl oral tablet</i>	T4	
<i>flurbiprofen oral tablet 100 mg</i>	T2	
<i>fluvoxamine oral tablet</i>	T2	
<b>FYCOMPA ORAL SUSPENSION</b>	T5	QL (744 ML per 31 days)
<b>FYCOMPA ORAL TABLET 10 MG, 12 MG, 4 MG, 6 MG, 8 MG</b>	T5	QL (31 EA per 31 days)
<b>FYCOMPA ORAL TABLET 2 MG</b>	T4	QL (31 EA per 31 days)
<i>gabapentin oral capsule 100 mg, 400 mg</i>	T1	PA-NS; QL (270 EA per 30 days)
<i>gabapentin oral capsule 300 mg</i>	T1	PA-NS; QL (360 EA per 30 days)
<i>gabapentin oral solution 250 mg/5 ml</i>	T2	PA-NS; QL (2160 ML per 30 days)
<i>gabapentin oral tablet 600 mg</i>	T1	PA-NS; QL (180 EA per 30 days)
<i>gabapentin oral tablet 800 mg</i>	T1	PA-NS; QL (120 EA per 30 days)
<i>galantamine oral capsule,ext rel. pellets 24 hr</i>	T3	
<i>galantamine oral solution</i>	T4	
<i>galantamine oral tablet 12 mg, 8 mg</i>	T3	
<i>galantamine oral tablet 4 mg</i>	T2	
<i>glatiramer subcutaneous syringe 20 mg/ml</i>	T5	PA; QL (31 ML per 31 days)
<i>glatiramer subcutaneous syringe 40 mg/ml</i>	T5	PA; QL (12 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML</b>	T5	PA; QL (31 ML per 31 days)
<b>GLATOPA SUBCUTANEOUS SYRINGE 40 MG/ML</b>	T5	PA; QL (12 ML per 28 days)
<i>guanfacine oral tablet extended release 24 hr</i>	T2	PA
<i>haloperidol</i>	T2	
<i>haloperidol decanoate</i>	T2	
<i>haloperidol lactate injection</i>	T2	
<i>haloperidol lactate oral</i>	T2	
<i>hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i>	T2	PA; QL (372 EA per 31 days)
<i>hydromorphone oral liquid</i>	T4	PA; QL (1240 ML per 31 days)
<i>hydromorphone oral tablet 2 mg, 4 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>hydromorphone oral tablet 8 mg</i>	T3	PA; QL (155 EA per 31 days)
<b>IBU ORAL TABLET 600 MG, 800 MG</b>	T1	
<i>ibuprofen oral tablet 400 mg, 600 mg, 800 mg</i>	T1	
<i>imipramine hcl</i>	T4	PA-NS
<i>indomethacin oral capsule</i>	T2	
<i>indomethacin oral capsule, extended release</i>	T2	
<b>INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,092 MG/3.5 ML</b>	T5	QL (3.5 ML per 180 days)
<b>INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,560 MG/5 ML</b>	T5	QL (5 ML per 180 days)
<b>INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 117 MG/0.75 ML</b>	T5	QL (0.75 ML per 28 days)
<b>INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 156 MG/ML</b>	T5	QL (1 ML per 28 days)
<b>INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 234 MG/1.5 ML</b>	T5	QL (1.5 ML per 28 days)
<b>INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 39 MG/0.25 ML</b>	T3	QL (0.25 ML per 28 days)
<b>INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 78 MG/0.5 ML</b>	T5	QL (0.5 ML per 28 days)
<b>INVEGA TRINZA INTRAMUSCULAR SYRINGE 273 MG/0.88 ML</b>	T5	QL (0.88 ML per 84 days)
<b>INVEGA TRINZA INTRAMUSCULAR SYRINGE 410 MG/1.32 ML</b>	T5	QL (1.32 ML per 84 days)
<b>INVEGA TRINZA INTRAMUSCULAR SYRINGE 546 MG/1.75 ML</b>	T5	QL (1.75 ML per 84 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>INVEGA TRINZA INTRAMUSCULAR SYRINGE 819 MG/2.63 ML</b>	T5	QL (2.63 ML per 84 days)
<b>KESIMPTA PEN</b>	T5	PA; QL (0.4 ML per 28 days)
<b>KLOXXADO</b>	T3	
<i>lacosamide oral</i>	T4	
<i>lamotrigine oral tablet</i>	T1	
<i>lamotrigine oral tablet extended release 24hr</i>	T4	
<i>lamotrigine oral tablet, chewable dispersible</i>	T2	
<i>levetiracetam oral solution 100 mg/ml</i>	T2	
<i>levetiracetam oral tablet</i>	T2	
<i>levetiracetam oral tablet extended release 24 hr</i>	T2	
<b>LIBERVANT</b>	T5	PA-NS; QL (10 EA per 30 days)
<i>lithium carbonate oral capsule</i>	T1	
<i>lithium carbonate oral tablet</i>	T1	
<i>lithium carbonate oral tablet extended release</i>	T2	
<i>lithium citrate</i>	T2	
<i>lofexidine</i>	T5	
<b>LORAZEPAM INTENSOL</b>	T2	PA; QL (155 ML per 31 days)
<i>lorazepam oral tablet 0.5 mg</i>	T2	PA; QL (124 EA per 31 days)
<i>lorazepam oral tablet 1 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>lorazepam oral tablet 2 mg</i>	T2	PA; QL (155 EA per 31 days)
<i>loxapine succinate</i>	T2	
<i>lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg</i>	T4	PA-NS; QL (31 EA per 31 days)
<i>lurasidone oral tablet 80 mg</i>	T4	PA-NS; QL (62 EA per 31 days)
<b>MARPLAN</b>	T4	
<i>meloxicam oral tablet</i>	T1	
<i>memantine oral capsule, sprinkle, er 24hr</i>	T2	
<i>memantine oral solution</i>	T2	
<i>memantine oral tablet</i>	T2	
<i>methadone oral solution 10 mg/5 ml</i>	T2	PA; QL (620 ML per 31 days)
<i>methadone oral solution 5 mg/5 ml</i>	T2	PA; QL (1240 ML per 31 days)
<i>methadone oral tablet 10 mg</i>	T2	PA; QL (124 EA per 31 days)
<i>methadone oral tablet 5 mg</i>	T2	PA; QL (248 EA per 31 days)
<i>methsuximide</i>	T4	
<i>methylphenidate hcl oral tablet</i>	T2	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>mirtazapine oral tablet 15 mg, 30 mg, 45 mg</i>	T1	
<i>mirtazapine oral tablet 7.5 mg</i>	T2	
<i>mirtazapine oral tablet,disintegrating</i>	T2	
<i>modafinil</i>	T3	PA; QL (31 EA per 31 days)
<i>molindone</i>	T2	
<i>morphine concentrate oral solution</i>	T2	PA; QL (310 ML per 31 days)
<i>morphine oral solution 10 mg/5 ml</i>	T2	PA; QL (2800 ML per 31 days)
<i>morphine oral solution 20 mg/5 ml (4 mg/ml)</i>	T2	PA; QL (1400 ML per 31 days)
<i>morphine oral tablet</i>	T2	PA; QL (186 EA per 31 days)
<i>morphine oral tablet extended release 100 mg</i>	T2	PA; QL (62 EA per 31 days)
<i>morphine oral tablet extended release 15 mg, 30 mg, 60 mg</i>	T2	PA; QL (100 EA per 31 days)
<i>morphine oral tablet extended release 200 mg</i>	T3	PA; QL (31 EA per 31 days)
<i>nabumetone</i>	T2	
<i>naloxone injection solution</i>	T2	
<i>naloxone injection syringe</i>	T2	
<i>naltrexone</i>	T2	
<b>NAMZARIC</b>	T3	PA
<i>naproxen oral tablet</i>	T1	
<i>naproxen oral tablet,delayed release (dr/ec) 375 mg</i>	T2	
<i>naproxen oral tablet,delayed release (dr/ec) 500 mg</i>	T4	
<i>naproxen sodium oral tablet 550 mg</i>	T2	
<i>naratriptan oral tablet 1 mg</i>	T3	QL (20 EA per 28 days)
<i>naratriptan oral tablet 2.5 mg</i>	T3	QL (9 EA per 28 days)
<b>NAYZILAM</b>	T4	PA-NS; QL (10 EA per 30 days)
<i>nefazodone</i>	T2	
<b>NEUPRO</b>	T4	
<i>nortriptyline</i>	T2	
<b>NUEDEXTA</b>	T5	PA; QL (62 EA per 31 days)
<b>NUPLAZID</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>NURTEC ODT</b>	T3	PA; QL (18 EA per 28 days)
<i>olanzapine intramuscular</i>	T4	
<i>olanzapine oral tablet</i>	T2	QL (31 EA per 31 days)
<i>olanzapine oral tablet,disintegrating</i>	T4	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>oxaprozin oral tablet</i>	T4	
<i>oxcarbazepine oral suspension</i>	T2	
<i>oxcarbazepine oral tablet</i>	T2	
<i>oxycodone oral capsule</i>	T2	PA; QL (186 EA per 31 days)
<i>oxycodone oral concentrate</i>	T4	PA; QL (180 ML per 31 days)
<i>oxycodone oral tablet 10 mg, 15 mg, 20 mg, 5 mg</i>	T2	PA; QL (186 EA per 31 days)
<i>oxycodone oral tablet 30 mg</i>	T3	PA; QL (138 EA per 31 days)
<i>oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg</i>	T3	PA; QL (372 EA per 31 days)
<i>oxycodone-acetaminophen oral tablet 5-325 mg, 7.5-325 mg</i>	T2	PA; QL (372 EA per 31 days)
<i>paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 9 mg</i>	T4	QL (31 EA per 31 days)
<i>paliperidone oral tablet extended release 24hr 6 mg</i>	T4	QL (62 EA per 31 days)
<i>paroxetine hcl oral suspension</i>	T4	
<i>paroxetine hcl oral tablet</i>	T1	
<i>paroxetine hcl oral tablet extended release 24 hr</i>	T4	
<i>perphenazine</i>	T2	
<b>PERSERIS</b>	T5	QL (1 EA per 28 days)
<i>phenelzine</i>	T3	
<i>phenobarbital</i>	T2	PA-NS
<b>PHENYTEK</b>	T4	
<i>phenytoin oral suspension 125 mg/5 ml</i>	T2	
<i>phenytoin oral tablet, chewable</i>	T2	
<i>phenytoin sodium extended oral capsule 100 mg</i>	T2	
<i>pimozide</i>	T4	
<i>piroxicam</i>	T2	
<i>pramipexole oral tablet</i>	T2	
<i>pregabalin oral capsule 100 mg, 150 mg, 200 mg, 25 mg, 50 mg, 75 mg</i>	T2	PA-NS; QL (93 EA per 31 days)
<i>pregabalin oral capsule 225 mg, 300 mg</i>	T2	PA-NS; QL (62 EA per 31 days)
<i>pregabalin oral solution</i>	T2	PA-NS; QL (930 ML per 31 days)
<i>primidone oral tablet 125 mg</i>	T4	
<i>primidone oral tablet 250 mg, 50 mg</i>	T2	
<i>protriptyline</i>	T4	
<i>pyridostigmine bromide oral tablet 60 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>pyridostigmine bromide oral tablet extended release</i>	T3	
<i>quetiapine oral tablet 100 mg, 200 mg, 25 mg, 300 mg, 400 mg, 50 mg</i>	T2	QL (62 EA per 31 days)
<i>quetiapine oral tablet 150 mg</i>	T3	QL (62 EA per 31 days)
<i>quetiapine oral tablet extended release 24 hr</i>	T2	QL (62 EA per 31 days)
<b>QULIPTA</b>	T3	PA; QL (31 EA per 31 days)
<b>RADICAVA ORS STARTER KIT SUSP</b>	T5	PA; QL (70 ML per 28 days)
<i>ramelteon</i>	T2	QL (31 EA per 31 days)
<i>rasagiline</i>	T4	
<b>REXULTI ORAL TABLET</b>	T5	PA-NS; QL (31 EA per 31 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 12.5 mg/2 ml, 25 mg/2 ml</i>	T3	QL (2 EA per 28 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 37.5 mg/2 ml</i>	T4	QL (2 EA per 28 days)
<i>risperidone microspheres intramuscular suspension,extended rel recon 50 mg/2 ml</i>	T5	QL (2 EA per 28 days)
<i>risperidone oral solution</i>	T2	QL (496 ML per 31 days)
<i>risperidone oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T1	QL (31 EA per 31 days)
<i>risperidone oral tablet 3 mg</i>	T1	QL (93 EA per 31 days)
<i>risperidone oral tablet 4 mg</i>	T1	QL (124 EA per 31 days)
<i>risperidone oral tablet,disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg</i>	T4	QL (31 EA per 31 days)
<i>risperidone oral tablet,disintegrating 3 mg</i>	T4	QL (93 EA per 31 days)
<i>risperidone oral tablet,disintegrating 4 mg</i>	T4	QL (124 EA per 31 days)
<i>rivastigmine</i>	T3	QL (30 EA per 30 days)
<i>rivastigmine tartrate</i>	T2	
<i>rizatriptan oral tablet 10 mg</i>	T2	QL (12 EA per 28 days)
<i>rizatriptan oral tablet 5 mg</i>	T2	QL (24 EA per 28 days)
<i>rizatriptan oral tablet,disintegrating 10 mg</i>	T2	QL (12 EA per 28 days)
<i>rizatriptan oral tablet,disintegrating 5 mg</i>	T2	QL (24 EA per 28 days)
<i>ropinirole oral tablet</i>	T2	
<i>ropinirole oral tablet extended release 24 hr</i>	T4	
<b>ROWEEPRA ORAL TABLET 500 MG</b>	T2	
<i>rufinamide oral suspension</i>	T5	PA-NS
<i>rufinamide oral tablet 200 mg</i>	T4	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>rufinamide oral tablet 400 mg</i>	T5	PA-NS
<b>RYTARY</b>	T3	ST
<b>SECUADO</b>	T5	PA-NS; QL (31 EA per 31 days)
<i>selegiline hcl</i>	T2	
<i>sertraline oral concentrate</i>	T2	
<i>sertraline oral tablet</i>	T1	
<b>SKYCLARYS</b>	T5	PA; QL (93 EA per 31 days)
<i>sodium oxybate</i>	T5	PA; QL (540 ML per 30 days)
<b>SPRITAM</b>	T4	
<b>SUBVENITE</b>	T2	
<i>sulindac</i>	T2	
<i>sumatriptan nasal spray,non-aerosol 20 mg/actuation</i>	T4	QL (8 EA per 28 days)
<i>sumatriptan nasal spray,non-aerosol 5 mg/actuation</i>	T4	QL (32 EA per 28 days)
<i>sumatriptan succinate oral tablet 100 mg</i>	T2	QL (9 EA per 28 days)
<i>sumatriptan succinate oral tablet 25 mg</i>	T2	QL (36 EA per 28 days)
<i>sumatriptan succinate oral tablet 50 mg</i>	T2	QL (18 EA per 28 days)
<i>sumatriptan succinate subcutaneous cartridge 6 mg/0.5 ml</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan succinate subcutaneous pen injector 4 mg/0.5 ml</i>	T4	QL (6 ML per 28 days)
<i>sumatriptan succinate subcutaneous pen injector 6 mg/0.5 ml</i>	T4	QL (4 ML per 28 days)
<i>sumatriptan succinate subcutaneous solution</i>	T4	QL (4 ML per 28 days)
<b>SYMPAZAN ORAL FILM 10 MG, 20 MG</b>	T5	PA-NS; QL (62 EA per 31 days)
<b>SYMPAZAN ORAL FILM 5 MG</b>	T4	PA-NS; QL (62 EA per 31 days)
<b>TASCENSO ODT</b>	T5	PA; QL (31 EA per 31 days)
<i>tasimelteon</i>	T5	PA; QL (31 EA per 31 days)
<i>teriflunomide</i>	T5	PA; QL (31 EA per 31 days)
<i>tetrabenazine oral tablet 12.5 mg</i>	T5	PA; QL (93 EA per 31 days)
<i>tetrabenazine oral tablet 25 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>thioridazine</i>	T3	
<i>thiothixene</i>	T3	
<i>tiagabine</i>	T4	
<i>tizanidine oral tablet</i>	T2	
<i>topiramate oral capsule, sprinkle 15 mg, 25 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>topiramate oral tablet</i>	T1	
<i>tramadol oral tablet 50 mg</i>	T2	PA; QL (240 EA per 30 days)
<i>tramadol-acetaminophen</i>	T2	PA; QL (372 EA per 31 days)
<i>tranylcypromine</i>	T4	
<i>trazodone oral tablet 100 mg, 150 mg, 50 mg</i>	T1	
<i>trifluoperazine</i>	T2	
<i>trimipramine</i>	T4	PA-NS
<b>TRINTELLIX</b>	T4	PA-NS
<b>UBRELVY ORAL TABLET 100 MG</b>	T5	PA; QL (17 EA per 28 days)
<b>UBRELVY ORAL TABLET 50 MG</b>	T5	PA; QL (34 EA per 28 days)
<i>valproic acid</i>	T2	
<i>valproic acid (as sodium salt) oral solution 250 mg/5 ml</i>	T2	
<b>VALTOCO NASAL SPRAY, NON-AEROSOL 10 MG/SPRAY (0.1 ML), 5 MG/SPRAY (0.1 ML)</b>	T4	PA-NS; QL (10 EA per 30 days)
<b>VALTOCO NASAL SPRAY, NON-AEROSOL 15 MG/2 SPRAY (7.5/0.1ML X 2), 20 MG/2 SPRAY (10MG/0.1ML X2)</b>	T5	PA-NS; QL (10 EA per 30 days)
<i>venlafaxine oral capsule, extended release 24hr 150 mg, 37.5 mg</i>	T2	QL (31 EA per 31 days)
<i>venlafaxine oral capsule, extended release 24hr 75 mg</i>	T2	QL (93 EA per 31 days)
<i>venlafaxine oral tablet</i>	T2	
<b>VERSACLOZ</b>	T5	QL (558 ML per 31 days)
<i>vigabatrin</i>	T5	PA-NS
<b>VIGADRONE</b>	T5	PA-NS
<b>VIGPODER</b>	T5	PA-NS
<i>vilazodone</i>	T3	PA-NS; QL (31 EA per 31 days)
<b>VIVITROL</b>	T5	
<b>VRAYLAR ORAL CAPSULE</b>	T5	PA-NS; QL (31 EA per 31 days)
<b>VUMERITY</b>	T5	PA; QL (124 EA per 31 days)
<b>XCOPRI</b>	T5	PA-NS
<b>XCOPRI MAINTENANCE PACK</b>	T5	PA-NS
<b>XCOPRI TITRATION PACK ORAL TABLETS, DOSE PACK 12.5 MG (14)- 25 MG (14)</b>	T4	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



Drug Name	Drug Tier	Requirements/Limits
<b>XCOPRI TITRATION PACK ORAL TABLETS, DOSE PACK 150 MG (14)- 200 MG (14), 50 MG (14)- 100 MG (14)</b>	T5	PA-NS
<b>XYREM</b>	T5	PA; QL (540 ML per 30 days)
<i>zaleplon oral capsule 10 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>zaleplon oral capsule 5 mg</i>	T4	PA; QL (93 EA per 31 days)
<b>ZAVZPRET</b>	T5	PA; QL (8 EA per 30 days)
<b>ZEPOSIA</b>	T5	PA; QL (31 EA per 31 days)
<b>ZEPOSIA STARTER KIT (28-DAY)</b>	T5	PA; QL (56 EA per 365 days)
<b>ZEPOSIA STARTER PACK (7-DAY)</b>	T5	PA; QL (14 EA per 365 days)
<b>ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML</b>	T5	PA; QL (11.648 ML per 28 days)
<b>ZILBRYSQ SUBCUTANEOUS SYRINGE 23 MG/0.574 ML</b>	T5	PA; QL (16 ML per 28 days)
<b>ZILBRYSQ SUBCUTANEOUS SYRINGE 32.4 MG/0.81 ML</b>	T5	PA; QL (22.68 ML per 28 days)
<i>ziprasidone hcl</i>	T2	QL (62 EA per 31 days)
<i>ziprasidone mesylate</i>	T4	
<i>zolpidem oral tablet</i>	T2	PA; QL (31 EA per 31 days)
<b>ZONISADE</b>	T4	PA-NS; QL (930 ML per 31 days)
<i>zonisamide</i>	T2	
<b>ZTALMY</b>	T5	PA-NS; QL (1100 ML per 30 days)
<b>ZUBSOLV SUBLINGUAL TABLET 0.7-0.18 MG, 2.9-0.71 MG, 8.6-2.1 MG</b>	T3	QL (62 EA per 31 days)
<b>ZUBSOLV SUBLINGUAL TABLET 1.4-0.36 MG</b>	T3	QL (93 EA per 31 days)
<b>ZUBSOLV SUBLINGUAL TABLET 11.4-2.9 MG, 5.7-1.4 MG</b>	T3	QL (31 EA per 31 days)
<b>ZURZUVAE ORAL CAPSULE 20 MG, 25 MG</b>	T5	PA-NS; QL (28 EA per 180 days)
<b>ZURZUVAE ORAL CAPSULE 30 MG</b>	T5	PA-NS; QL (14 EA per 180 days)
<b>Cardiovascular, Hypertension / Lipids</b>		
<i>acebutolol</i>	T2	
<i>aliskiren</i>	T4	
<i>amiloride</i>	T2	
<i>amiloride-hydrochlorothiazide</i>	T2	
<i>amiodarone oral</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>amlodipine</i>	T1	
<i>amlodipine-atorvastatin</i>	T2	
<i>amlodipine-benazepril</i>	T1	
<i>amlodipine-olmesartan</i>	T2	QL (31 EA per 31 days)
<i>amlodipine-valsartan</i>	T1	
<i>aspirin-dipyridamole</i>	T4	
<i>atenolol</i>	T1	
<i>atenolol-chlorthalidone</i>	T2	
<i>atorvastatin</i>	T1	
<i>benazepril</i>	T1	
<i>benazepril-hydrochlorothiazide</i>	T1	
<i>bisoprolol fumarate</i>	T2	
<i>bisoprolol-hydrochlorothiazide</i>	T1	
<b>BRILINTA</b>	T3	
<i>bumetanide oral</i>	T2	
<b>CABLIVI INJECTION KIT</b>	T5	PA; QL (31 EA per 31 days)
<b>CAMZYOS</b>	T5	PA; QL (31 EA per 31 days)
<i>candesartan</i>	T2	
<i>candesartan-hydrochlorothiazid</i>	T2	
<i>captopril</i>	T2	
<b>CARTIA XT</b>	T2	
<i>carvedilol</i>	T1	
<i>chlorthalidone oral tablet 25 mg, 50 mg</i>	T2	
<i>cholestyramine (with sugar) oral powder in packet</i>	T2	
<b>CHOLESTYRAMINE LIGHT ORAL POWDER IN PACKET</b>	T2	
<i>cilostazol</i>	T2	
<i>clonidine</i>	T4	
<i>clonidine hcl oral tablet</i>	T1	
<i>clopidogrel oral tablet 75 mg</i>	T1	
<i>colesevelam</i>	T4	
<i>colestipol oral packet</i>	T4	
<i>colestipol oral tablet</i>	T3	
<b>CORLANOR ORAL SOLUTION</b>	T3	PA; QL (420 ML per 28 days)
<i>digoxin oral solution</i>	T3	QL (155 ML per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>digoxin oral tablet 125 mcg (0.125 mg)</i>	T2	QL (62 EA per 31 days)
<i>digoxin oral tablet 250 mcg (0.25 mg)</i>	T2	QL (31 EA per 31 days)
<i>diltiazem hcl oral capsule,extended release 12 hr</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24 hr 360 mg, 420 mg</i>	T2	
<i>diltiazem hcl oral capsule,extended release 24hr 120 mg, 180 mg, 240 mg, 300 mg</i>	T2	
<i>diltiazem hcl oral tablet</i>	T1	
<i>diltiazem hcl oral tablet extended release 24 hr</i>	T2	
<b>DILT-XR</b>	T2	
<i>dofetilide</i>	T4	
<b>DOPTELET (10 TAB PACK)</b>	T5	PA
<b>DOPTELET (15 TAB PACK)</b>	T5	PA
<b>DOPTELET (30 TAB PACK)</b>	T5	PA
<i>doxazosin</i>	T1	
<b>ELIQUIS DVT-PE TREAT 30D START</b>	T3	QL (74 EA per 30 days)
<b>ELIQUIS ORAL TABLET 2.5 MG</b>	T3	QL (60 EA per 30 days)
<b>ELIQUIS ORAL TABLET 5 MG</b>	T3	QL (74 EA per 30 days)
<i>enalapril maleate oral tablet</i>	T1	
<i>enalapril-hydrochlorothiazide oral tablet 5-12.5 mg</i>	T1	
<i>enoxaparin subcutaneous syringe</i>	T4	
<b>ENTRESTO ORAL TABLET 24-26 MG</b>	T3	QL (186 EA per 31 days)
<b>ENTRESTO ORAL TABLET 49-51 MG</b>	T3	QL (93 EA per 31 days)
<b>ENTRESTO ORAL TABLET 97-103 MG</b>	T3	QL (62 EA per 31 days)
<i>eplerenone</i>	T2	
<i>ethacrynic acid</i>	T4	
<i>ezetimibe</i>	T2	
<i>ezetimibe-simvastatin</i>	T3	QL (31 EA per 31 days)
<i>felodipine</i>	T2	
<i>fenofibrate micronized oral capsule 134 mg, 200 mg, 67 mg</i>	T2	
<i>fenofibrate nanocrystallized</i>	T2	
<i>fenofibrate oral tablet 160 mg, 54 mg</i>	T2	
<i>flecainide</i>	T2	
<i>fluvastatin oral capsule</i>	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>fondaparinux subcutaneous syringe 10 mg/0.8 ml, 7.5 mg/0.6 ml</i>	T5	
<i>fondaparinux subcutaneous syringe 2.5 mg/0.5 ml, 5 mg/0.4 ml</i>	T4	
<i>fosinopril</i>	T1	
<i>fosinopril-hydrochlorothiazide</i>	T2	
<b>FUROSCIX</b>	T5	PA; QL (8 EA per 30 days)
<i>furosemide injection solution</i>	T2	
<i>furosemide oral solution 10 mg/ml, 40 mg/5 ml (8 mg/ml)</i>	T2	
<i>furosemide oral tablet</i>	T1	
<i>gemfibrozil</i>	T1	
<i>heparin (porcine) injection solution</i>	T3	PA-BvD
<i>hydralazine oral</i>	T2	
<i>hydrochlorothiazide</i>	T1	
<i>icosapent ethyl oral capsule 0.5 gram</i>	T2	QL (248 EA per 31 days)
<i>icosapent ethyl oral capsule 1 gram</i>	T2	QL (124 EA per 31 days)
<i>indapamide</i>	T1	
<i>irbesartan</i>	T1	QL (31 EA per 31 days)
<i>irbesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>isosorbide dinitrate oral tablet 10 mg, 20 mg, 30 mg, 5 mg</i>	T2	
<i>isosorbide mononitrate oral tablet extended release 24 hr</i>	T1	
<i>isradipine</i>	T2	
<i>ivabradine oral tablet 5 mg</i>	T3	PA; QL (93 EA per 31 days)
<i>ivabradine oral tablet 7.5 mg</i>	T3	PA; QL (62 EA per 31 days)
<b>JANTOVEN</b>	T1	
<b>KERENDIA</b>	T4	PA; QL (31 EA per 31 days)
<i>labetalol oral tablet 100 mg, 200 mg, 300 mg</i>	T2	
<i>lisinopril</i>	T1	
<i>lisinopril-hydrochlorothiazide</i>	T1	
<i>losartan oral tablet 100 mg</i>	T1	QL (31 EA per 31 days)
<i>losartan oral tablet 25 mg</i>	T1	QL (93 EA per 31 days)
<i>losartan oral tablet 50 mg</i>	T1	QL (62 EA per 31 days)
<i>losartan-hydrochlorothiazide</i>	T1	
<i>lovastatin</i>	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>metolazone</i>	T2	
<i>metoprolol succinate</i>	T1	
<i>metoprolol ta-hydrochlorothiaz</i>	T2	
<i>metoprolol tartrate oral tablet 100 mg, 25 mg, 50 mg</i>	T1	
<i>metoprolol tartrate oral tablet 37.5 mg, 75 mg</i>	T2	
<i>metyrosine</i>	T5	PA
<i>mexiletine</i>	T3	
<i>minoxidil oral</i>	T2	
<i>moexipril</i>	T1	
<b>MULPLETA</b>	T5	PA
<b>MULTAQ</b>	T4	
<i>nadolol</i>	T2	
<i>nebivolol oral tablet 10 mg, 2.5 mg</i>	T2	QL (93 EA per 31 days)
<i>nebivolol oral tablet 20 mg</i>	T2	QL (62 EA per 31 days)
<i>nebivolol oral tablet 5 mg</i>	T2	QL (217 EA per 31 days)
<b>NEXLETOL</b>	T3	PA; QL (31 EA per 31 days)
<b>NEXLIZET</b>	T4	PA; QL (31 EA per 31 days)
<i>niacin oral tablet extended release 24 hr 1,000 mg, 750 mg</i>	T2	
<i>niacin oral tablet extended release 24 hr 500 mg</i>	T2	QL (31 EA per 31 days)
<i>nicardipine oral capsule 20 mg</i>	T4	
<i>nicardipine oral capsule 30 mg</i>	T5	
<i>nifedipine oral tablet extended release</i>	T2	
<i>nifedipine oral tablet extended release 24hr</i>	T2	
<i>nimodipine oral capsule</i>	T4	
<b>NITRO-BID</b>	T2	
<i>nitroglycerin sublingual</i>	T2	
<i>nitroglycerin transdermal patch 24 hour</i>	T2	
<i>nitroglycerin translingual</i>	T4	
<i>olmesartan oral tablet 20 mg, 40 mg</i>	T1	QL (31 EA per 31 days)
<i>olmesartan oral tablet 5 mg</i>	T1	QL (93 EA per 31 days)
<i>olmesartan-amlodipin-hcthiazyd</i>	T3	
<i>olmesartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>omega-3 acid ethyl esters</i>	T2	QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>PACERONE ORAL TABLET 100 MG, 200 MG, 400 MG</b>	T2	
<i>pentoxifylline</i>	T2	
<i>perindopril erbumine</i>	T1	
<i>pindolol</i>	T3	
<i>pitavastatin calcium</i>	T3	
<i>prasugrel hcl</i>	T2	
<i>pravastatin</i>	T1	
<i>prazosin</i>	T2	
<b>PREVALITE ORAL POWDER IN PACKET</b>	T2	
<b>PROMACTA ORAL POWDER IN PACKET 12.5 MG</b>	T5	PA; QL (372 EA per 31 days)
<b>PROMACTA ORAL POWDER IN PACKET 25 MG</b>	T5	PA; QL (31 EA per 31 days)
<b>PROMACTA ORAL TABLET 12.5 MG, 25 MG</b>	T5	PA; QL (31 EA per 31 days)
<b>PROMACTA ORAL TABLET 50 MG, 75 MG</b>	T5	PA; QL (62 EA per 31 days)
<i>propafenone oral capsule,extended release 12 hr</i>	T4	
<i>propafenone oral tablet</i>	T2	
<i>propranolol oral capsule,extended release 24 hr</i>	T2	
<i>propranolol oral solution</i>	T2	
<i>propranolol oral tablet</i>	T1	
<i>quinapril</i>	T1	
<i>quinapril-hydrochlorothiazide</i>	T1	
<i>quinidine sulfate oral tablet</i>	T2	
<i>ramipril</i>	T1	
<i>ranolazine</i>	T4	QL (62 EA per 31 days)
<b>REPATHA PUSHTRONEX</b>	T3	PA; QL (7 ML per 28 days)
<b>REPATHA SURECLICK</b>	T3	PA; QL (3 ML per 28 days)
<b>REPATHA SYRINGE</b>	T3	PA; QL (3 ML per 28 days)
<i>rosuvastatin</i>	T1	
<i>simvastatin</i>	T1	
<b>SOTALOL AF</b>	T2	
<i>sotalol oral</i>	T2	
<i>spironolactone oral tablet</i>	T1	
<i>spironolacton-hydrochlorothiaz</i>	T2	
<i>telmisartan</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>telmisartan-amlodipine</i>	T2	
<i>telmisartan-hydrochlorothiazid</i>	T2	
<i>terazosin</i>	T1	
<b>TIADYLT ER</b>	T2	
<i>timolol maleate oral</i>	T2	
<i>toremide oral</i>	T2	
<i>trandolapril</i>	T1	
<i>triamterene-hydrochlorothiazid</i>	T1	
<b>UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 400 MCG, 600 MCG, 800 MCG</b>	T5	PA; QL (62 EA per 31 days)
<b>UPTRAVI ORAL TABLET 200 MCG</b>	T5	PA; QL (224 EA per 28 days)
<b>UPTRAVI ORAL TABLETS,DOSE PACK</b>	T5	PA; QL (400 EA per 365 days)
<i>valsartan oral tablet 160 mg, 40 mg, 80 mg</i>	T1	QL (62 EA per 31 days)
<i>valsartan oral tablet 320 mg</i>	T1	QL (31 EA per 31 days)
<i>valsartan-hydrochlorothiazide</i>	T1	QL (31 EA per 31 days)
<i>verapamil oral capsule, 24 hr er pellet ct</i>	T4	
<i>verapamil oral capsule,ext rel. pellets 24 hr 120 mg, 180 mg, 240 mg</i>	T2	
<i>verapamil oral capsule,ext rel. pellets 24 hr 360 mg</i>	T4	
<i>verapamil oral tablet</i>	T2	
<i>verapamil oral tablet extended release</i>	T2	
<b>VERQUVO</b>	T4	PA; QL (31 EA per 31 days)
<b>VYNDAQEL</b>	T5	PA; QL (124 EA per 31 days)
<i>warfarin</i>	T1	
<b>XARELTO DVT-PE TREAT 30D START</b>	T3	QL (51 EA per 30 days)
<b>XARELTO ORAL SUSPENSION FOR RECONSTITUTION</b>	T3	QL (930 ML per 31 days)
<b>XARELTO ORAL TABLET 10 MG, 20 MG</b>	T3	QL (31 EA per 31 days)
<b>XARELTO ORAL TABLET 15 MG</b>	T3	QL (52 EA per 31 days)
<b>XARELTO ORAL TABLET 2.5 MG</b>	T3	QL (62 EA per 31 days)
<b>Dermatologicals/Topical Therapy</b>		
<b>ACUTANE ORAL CAPSULE 10 MG, 20 MG, 40 MG</b>	T4	
<i>acitretin</i>	T4	PA
<i>acyclovir topical ointment</i>	T4	QL (30 GM per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>ADBRY</b>	T5	PA; QL (4 ML per 28 days)
<b>ALA-CORT TOPICAL CREAM 1 %</b>	T2	
<i>alclometasone</i>	T2	
<i>ammonium lactate</i>	T2	
<b>AMNESTEEM</b>	T4	
<i>betamethasone dipropionate</i>	T2	
<i>betamethasone valerate topical cream</i>	T2	
<i>betamethasone valerate topical lotion</i>	T2	
<i>betamethasone valerate topical ointment</i>	T2	
<i>betamethasone, augmented</i>	T2	
<i>calcipotriene scalp</i>	T3	QL (60 ML per 28 days)
<i>calcipotriene topical cream</i>	T4	QL (60 GM per 28 days)
<i>calcipotriene topical ointment</i>	T3	QL (60 GM per 28 days)
<b>CIBINQO</b>	T5	PA; QL (31 EA per 31 days)
<i>ciclopirox topical cream</i>	T2	QL (90 GM per 28 days)
<i>ciclopirox topical gel</i>	T2	QL (45 GM per 28 days)
<i>ciclopirox topical shampoo</i>	T2	QL (120 ML per 28 days)
<i>ciclopirox topical solution</i>	T2	
<i>ciclopirox topical suspension</i>	T2	QL (60 ML per 28 days)
<b>CLARAVIS</b>	T4	
<i>clindamycin phosphate topical gel</i>	T2	QL (60 GM per 28 days)
<i>clindamycin phosphate topical lotion</i>	T2	QL (60 ML per 28 days)
<i>clindamycin phosphate topical solution</i>	T2	QL (60 ML per 28 days)
<i>clobetasol topical cream 0.05 %</i>	T4	QL (60 GM per 28 days)
<i>clobetasol topical ointment</i>	T4	QL (60 GM per 28 days)
<i>clotrimazole topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole topical solution</i>	T2	QL (30 ML per 28 days)
<i>clotrimazole-betamethasone topical cream</i>	T2	QL (45 GM per 28 days)
<i>clotrimazole-betamethasone topical lotion</i>	T3	QL (60 ML per 28 days)
<b>COSENTYX (2 SYRINGES)</b>	T5	PA; QL (2 ML per 28 days)
<b>COSENTYX PEN (2 PENS)</b>	T5	PA; QL (2 ML per 28 days)
<b>COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 28 days)
<b>COSENTYX UNOREADY PEN</b>	T5	PA; QL (2 ML per 28 days)
<i>desoximetasone topical cream</i>	T4	QL (100 GM per 28 days)
<i>desoximetasone topical gel</i>	T4	QL (60 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>diclofenac sodium topical gel 3 %</i>	T4	PA; QL (100 GM per 28 days)
<b>DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML</b>	T5	PA; QL (2.28 ML per 28 days)
<b>DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML</b>	T5	PA; QL (8 ML per 28 days)
<b>DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML</b>	T5	PA; QL (2.28 ML per 28 days)
<b>DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 300 MG/2 ML</b>	T5	PA; QL (8 ML per 28 days)
<i>econazole nitrate</i>	T2	QL (85 GM per 28 days)
<b>ERY PADS</b>	T2	
<i>erythromycin with ethanol topical solution</i>	T2	QL (60 ML per 28 days)
<b>FILSUEVZ</b>	T5	PA
<i>fluocinolone and shower cap</i>	T2	QL (118.28 ML per 28 days)
<i>fluocinolone topical cream 0.01 %</i>	T2	QL (60 GM per 28 days)
<i>fluocinolone topical cream 0.025 %</i>	T2	QL (120 GM per 28 days)
<i>fluocinolone topical ointment</i>	T2	QL (120 GM per 28 days)
<i>fluocinolone topical solution</i>	T2	QL (90 ML per 28 days)
<i>fluocinonide topical cream 0.05 %</i>	T2	QL (60 GM per 28 days)
<i>fluocinonide topical gel</i>	T3	QL (60 GM per 28 days)
<i>fluocinonide topical ointment</i>	T2	QL (60 GM per 28 days)
<i>fluocinonide topical solution</i>	T2	QL (60 ML per 28 days)
<i>fluocinonide-emollient</i>	T4	QL (60 GM per 28 days)
<i>fluorouracil topical cream 5 %</i>	T2	
<i>fluorouracil topical solution</i>	T2	
<i>fluticasone propionate topical cream</i>	T2	
<i>gentamicin topical</i>	T2	QL (60 GM per 28 days)
<i>halobetasol propionate topical cream</i>	T2	QL (50 GM per 28 days)
<i>halobetasol propionate topical ointment</i>	T2	QL (50 GM per 28 days)
<i>hydrocortisone topical cream 1 %</i>	T2	
<i>hydrocortisone topical lotion 2.5 %</i>	T2	QL (118 ML per 28 days)
<i>hydrocortisone topical ointment 1 %, 2.5 %</i>	T2	
<i>imiquimod topical cream in packet 5 %</i>	T2	
<i>ketoconazole topical cream</i>	T2	QL (60 GM per 28 days)
<i>ketoconazole topical shampoo</i>	T2	QL (120 ML per 28 days)
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i>	T2	PA; QL (50 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>lidocaine topical adhesive patch,medicated 5 %</i>	T2	PA; QL (93 EA per 31 days)
<i>lidocaine topical ointment</i>	T4	PA; QL (50 GM per 28 days)
<b>LIDOCAINE VISCOUS</b>	T2	
<i>lidocaine-prilocaine topical cream</i>	T2	PA; QL (30 GM per 28 days)
<i>malathion</i>	T2	
<i>metronidazole topical cream</i>	T2	
<i>metronidazole topical gel</i>	T2	
<i>metronidazole topical lotion</i>	T2	
<i>mometasone topical</i>	T2	
<i>mupirocin</i>	T2	
<b>NYAMYC</b>	T2	QL (60 GM per 28 days)
<i>nystatin topical cream</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical ointment</i>	T2	QL (30 GM per 28 days)
<i>nystatin topical powder</i>	T2	QL (60 GM per 28 days)
<i>nystatin-triamcinolone</i>	T2	QL (60 GM per 28 days)
<b>NYSTOP</b>	T2	QL (60 GM per 28 days)
<b>PANRETIN</b>	T5	PA-NS
<i>penciclovir</i>	T4	QL (5 GM per 28 days)
<i>permethrin</i>	T2	
<i>pimecrolimus</i>	T4	QL (100 GM per 28 days)
<i>podofilox topical solution</i>	T2	
<b>REGRANEX</b>	T5	PA
<i>selenium sulfide topical lotion</i>	T2	
<i>silver sulfadiazine</i>	T2	
<b>SKYRIZI SUBCUTANEOUS PEN INJECTOR</b>	T5	PA; QL (1 ML per 84 days)
<b>SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML</b>	T5	PA; QL (1 ML per 84 days)
<b>SSD</b>	T4	
<b>STELARA SUBCUTANEOUS SOLUTION</b>	T5	PA; QL (0.5 ML per 84 days)
<b>STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 84 days)
<b>STELARA SUBCUTANEOUS SYRINGE 90 MG/ML</b>	T5	PA; QL (1 ML per 56 days)
<i>sulfacetamide sodium (acne)</i>	T2	
<b>SULFAMYLON TOPICAL CREAM</b>	T3	
<i>tacrolimus topical</i>	T2	QL (100 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>TALTZ AUTOINJECTOR</b>	T5	PA; QL (1 ML per 28 days)
<b>TALTZ SYRINGE SUBCUTANEOUS SYRINGE 20 MG/0.25 ML</b>	T5	PA; QL (0.25 ML per 28 days)
<b>TALTZ SYRINGE SUBCUTANEOUS SYRINGE 40 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 28 days)
<b>TALTZ SYRINGE SUBCUTANEOUS SYRINGE 80 MG/ML</b>	T5	PA; QL (1 ML per 28 days)
<i>tazarotene topical cream</i>	T4	PA; QL (60 GM per 28 days)
<i>tretinoin topical cream</i>	T2	PA; QL (45 GM per 28 days)
<i>tretinoin topical gel 0.01 %, 0.025 %</i>	T3	PA; QL (45 GM per 28 days)
<i>triamcinolone acetonide topical cream</i>	T2	
<i>triamcinolone acetonide topical lotion</i>	T2	
<i>triamcinolone acetonide topical ointment 0.025 %, 0.1 %, 0.5 %</i>	T2	
<b>TRIDERM TOPICAL CREAM 0.5 %</b>	T4	
<b>VALCHLOR</b>	T5	PA-NS
<b>Diagnostics / Miscellaneous Agents</b>		
<i>acamprosate</i>	T4	
<i>anagrelide</i>	T2	
<i>bupropion hcl (smoking deter)</i>	T2	QL (62 EA per 31 days)
<i>carglumic acid</i>	T5	PA
<i>cevimeline</i>	T2	
<b>CHEMET</b>	T4	
<b>CLINIMIX 4.25%/D5W SULFIT FREE</b>	T4	PA-BvD
<i>d10 %-0.45 % sodium chloride</i>	T2	
<i>d2.5 %-0.45 % sodium chloride</i>	T2	
<i>d5 % and 0.9 % sodium chloride</i>	T2	
<i>d5 %-0.45 % sodium chloride</i>	T2	
<i>deferasirox oral granules in packet</i>	T5	PA
<i>deferasirox oral tablet 180 mg, 360 mg</i>	T5	PA
<i>deferasirox oral tablet 90 mg</i>	T3	PA
<i>deferasirox oral tablet, dispersible 125 mg</i>	T4	PA
<i>deferasirox oral tablet, dispersible 250 mg, 500 mg</i>	T5	PA
<i>deferiprone</i>	T5	PA
<i>dextrose 10 % in water (d10w)</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>dextrose 5 % in water (d5w) intravenous parenteral solution</i>	T2	
<i>disulfiram</i>	T3	
<i>droxidopa oral capsule 100 mg</i>	T5	PA; QL (465 EA per 31 days)
<i>droxidopa oral capsule 200 mg, 300 mg</i>	T5	PA; QL (186 EA per 31 days)
<b>DUVYZAT</b>	T5	PA; QL (420 ML per 35 days)
<b>FABHALTA</b>	T5	PA; QL (62 EA per 31 days)
<b>INCRELEX</b>	T5	PA
<b>JOENJA</b>	T5	PA; QL (60 EA per 30 days)
<b>KIONEX (WITH SORBITOL)</b>	T2	
<i>levocarnitine (with sugar)</i>	T2	PA-BvD
<i>levocarnitine oral tablet</i>	T2	PA-BvD
<b>LITFULO</b>	T5	PA; QL (28 EA per 28 days)
<b>LOKELMA</b>	T3	PA; QL (93 EA per 31 days)
<i>midodrine</i>	T2	
<b>NICOTROL NS</b>	T4	
<i>nitisinone</i>	T5	PA
<b>PHEBURANE</b>	T5	PA; QL (620 GM per 31 days)
<i>pilocarpine hcl oral</i>	T3	
<b>PROLASTIN-C INTRAVENOUS SOLUTION</b>	T5	PA
<b>RAVICTI</b>	T5	PA
<b>REZDIFFRA</b>	T5	PA; QL (31 EA per 31 days)
<i>riluzole</i>	T3	
<i>risedronate oral tablet 30 mg</i>	T4	
<i>sodium chloride 0.9 % intravenous parenteral solution</i>	T2	
<i>sodium chloride irrigation</i>	T2	
<i>sodium phenylbutyrate</i>	T5	PA
<i>sodium polystyrene sulfonate oral powder</i>	T2	
<b>SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 2.5 MG, 5 MG</b>	T5	PA; QL (31 EA per 31 days)
<b>SOHONOS ORAL CAPSULE 10 MG</b>	T5	PA; QL (62 EA per 31 days)
<b>SPS (WITH SORBITOL) ORAL</b>	T2	
<i>trientine oral capsule 250 mg</i>	T5	QL (248 EA per 31 days)
<i>varenicline tartrate oral tablet</i>	T4	QL (60 EA per 30 days)
<i>varenicline tartrate oral tablets,dose pack</i>	T4	QL (106 EA per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<b>Ear, Nose / Throat Medications</b>		
<i>acetic acid otic (ear)</i>	T2	
<i>azelastine nasal spray,non-aerosol 137 mcg (0.1 %)</i>	T2	QL (30 ML per 25 days)
<i>chlorhexidine gluconate mucous membrane</i>	T1	
<i>ciprofloxacin-dexamethasone</i>	T3	
<i>fluocinolone acetonide oil</i>	T2	
<i>hydrocortisone-acetic acid</i>	T2	
<i>ipratropium bromide nasal spray,non-aerosol 21 mcg (0.03 %)</i>	T2	QL (30 ML per 28 days)
<i>ipratropium bromide nasal spray,non-aerosol 42 mcg (0.06 %)</i>	T2	QL (15 ML per 28 days)
<b>KOURZEQ</b>	T2	
<i>neomycin-polymyxin-hc otic (ear)</i>	T2	
<i>ofloxacin otic (ear)</i>	T2	
<i>olopatadine nasal</i>	T3	QL (30.5 GM per 30 days)
<b>PERIOGARD</b>	T1	
<i>triamcinolone acetonide dental</i>	T2	
<b>Endocrine/Diabetes</b>		
<i>acarbose</i>	T2	QL (93 EA per 31 days)
<b>ALCOHOL PADS</b>	T2	PA
<b>BAQSIMI</b>	T3	
<i>cabergoline</i>	T2	
<i>calcitonin (salmon) nasal</i>	T2	PA-BvD
<i>calcitriol oral</i>	T2	PA-BvD
<b>CERDELGA</b>	T5	PA; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 30 mg, 60 mg</i>	T4	PA-BvD; QL (62 EA per 31 days)
<i>cinacalcet oral tablet 90 mg</i>	T4	PA-BvD; QL (124 EA per 31 days)
<i>danazol</i>	T4	
<i>desmopressin nasal spray,non-aerosol 10 mcg/spray (0.1 ml)</i>	T2	
<i>desmopressin oral</i>	T2	
<i>dexamethasone oral solution</i>	T2	
<i>dexamethasone oral tablet</i>	T1	
<i>diazoxide</i>	T5	
<i>doxercalciferol oral</i>	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>EUTHYROX</b>	T3	
<b>FARXIGA</b>	T3	QL (31 EA per 31 days)
<i>fludrocortisone</i>	T1	
<i>glimepiride oral tablet 1 mg, 2 mg, 4 mg</i>	T1	
<i>glipizide oral tablet 10 mg, 5 mg</i>	T1	
<i>glipizide oral tablet extended release 24hr</i>	T1	
<i>glipizide-metformin</i>	T1	
<b>GLUCAGON EMERGENCY KIT (HUMAN)</b>	T3	
<b>GLYXAMBI</b>	T3	QL (31 EA per 31 days)
<b>GVOKE</b>	T3	
<b>GVOKE HYPOPEN 2-PACK</b>	T3	
<b>GVOKE PFS 1-PACK SYRINGE SUBCUTANEOUS SYRINGE 1 MG/0.2 ML</b>	T3	
<b>HUMALOG JUNIOR KWIKPEN U-100</b>	T3	
<b>HUMALOG KWIKPEN INSULIN</b>	T3	
<b>HUMALOG MIX 50-50 KWIKPEN</b>	T3	
<b>HUMALOG MIX 75-25 KWIKPEN</b>	T3	
<b>HUMALOG MIX 75-25(U-100)INSULN</b>	T3	
<b>HUMALOG TEMPO PEN(U-100)INSULN</b>	T3	
<b>HUMALOG U-100 INSULIN</b>	T3	
<b>HUMULIN 70/30 U-100 INSULIN</b>	T3	
<b>HUMULIN 70/30 U-100 KWIKPEN</b>	T3	
<b>HUMULIN N NPH INSULIN KWIKPEN</b>	T3	
<b>HUMULIN N NPH U-100 INSULIN</b>	T3	
<b>HUMULIN R REGULAR U-100 INSULN</b>	T3	
<b>HUMULIN R U-500 (CONC) INSULIN</b>	T3	
<b>HUMULIN R U-500 (CONC) KWIKPEN</b>	T3	
<i>hydrocortisone oral</i>	T2	
<i>insulin lispro</i>	T3	
<i>insulin lispro protamin-lispro</i>	T3	
<b>JANUMET</b>	T3	QL (62 EA per 31 days)
<b>JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 100-1,000 MG, 50-500 MG</b>	T3	QL (31 EA per 31 days)
<b>JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 50-1,000 MG</b>	T3	QL (62 EA per 31 days)
<b>JANUVIA ORAL TABLET 100 MG, 50 MG</b>	T3	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>JANUVIA ORAL TABLET 25 MG</b>	T3	QL (93 EA per 31 days)
<b>JARDIANCE ORAL TABLET 10 MG</b>	T3	QL (62 EA per 31 days)
<b>JARDIANCE ORAL TABLET 25 MG</b>	T3	QL (31 EA per 31 days)
<b>JAVYGTOR</b>	T5	PA
<b>JENTADUETO ORAL TABLET 2.5-1,000 MG, 2.5-500 MG</b>	T3	QL (62 EA per 31 days)
<b>JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG</b>	T3	QL (62 EA per 31 days)
<b>JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 5-1,000 MG</b>	T3	QL (31 EA per 31 days)
<b>LANTUS SOLOSTAR U-100 INSULIN</b>	T3	
<b>LANTUS U-100 INSULIN</b>	T3	
<i>levothyroxine oral tablet</i>	T1	
<b>LEVOXYL ORAL TABLET 100 MCG, 112 MCG, 125 MCG, 137 MCG, 150 MCG, 175 MCG, 200 MCG, 25 MCG, 50 MCG, 75 MCG, 88 MCG</b>	T3	
<i>liothyronine oral</i>	T2	
<i>metformin oral tablet 1,000 mg, 500 mg, 850 mg</i>	T1	
<i>metformin oral tablet extended release 24 hr</i>	T1	
<i>metformin oral tablet extended release 24hr</i>	NF	
<i>metformin oral tablet,er gast.retention 24 hr</i>	NF	
<i>methimazole oral tablet 10 mg, 5 mg</i>	T1	
<i>methylprednisolone</i>	T2	
<i>mifepristone oral tablet 300 mg</i>	T5	PA; QL (124 EA per 31 days)
<i>miglustat</i>	T5	PA; QL (93 EA per 31 days)
<b>MOUNJARO</b>	T3	PA; QL (2 ML per 28 days)
<i>nateglinide</i>	T2	QL (93 EA per 31 days)
<b>OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)</b>	T3	PA; QL (3 ML per 28 days)
<b>PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML</b>	T5	PA; QL (15 ML per 30 days)
<b>PALYNZIQ SUBCUTANEOUS SYRINGE 2.5 MG/0.5 ML</b>	T5	PA; QL (4 ML per 30 days)
<b>PALYNZIQ SUBCUTANEOUS SYRINGE 20 MG/ML</b>	T5	PA; QL (90 ML per 30 days)
<i>paricalcitol oral</i>	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>pioglitazone</i>	T1	QL (31 EA per 31 days)
<i>pioglitazone-metformin</i>	T2	QL (93 EA per 31 days)
<i>prednisolone oral solution</i>	T2	
<i>prednisolone sodium phosphate oral solution 25 mg/5 ml (5 mg/ml), 5 mg base/5 ml (6.7 mg/5 ml)</i>	T2	
<i>prednisone oral solution</i>	T3	
<i>prednisone oral tablet</i>	T1	
<i>prednisone oral tablets,dose pack</i>	T2	
<i>propylthiouracil</i>	T2	
<b>RECORLEV</b>	T5	PA; QL (248 EA per 31 days)
<i>repaglinide oral tablet 0.5 mg, 1 mg</i>	T2	QL (124 EA per 31 days)
<i>repaglinide oral tablet 2 mg</i>	T2	QL (248 EA per 31 days)
<b>RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG</b>	T3	PA; QL (31 EA per 31 days)
<i>sapropterin</i>	T5	PA
<b>SOLQUA 100/33</b>	T3	QL (18 ML per 30 days)
<b>SOMAVERT</b>	T5	PA
<b>SYMLINPEN 120</b>	T5	QL (10.8 ML per 28 days)
<b>SYMLINPEN 60</b>	T5	QL (6 ML per 28 days)
<b>SYNAREL</b>	T5	PA
<b>SYNJARDY</b>	T3	QL (62 EA per 31 days)
<b>SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 12.5-1,000 MG, 5-1,000 MG</b>	T3	QL (62 EA per 31 days)
<b>SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 25-1,000 MG</b>	T3	QL (31 EA per 31 days)
<b>SYNTHROID</b>	T3	
<i>testosterone cypionate</i>	T2	PA
<i>testosterone enanthate</i>	T3	PA
<i>testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)</i>	T3	PA
<i>testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram)</i>	T2	PA
<i>testosterone transdermal gel in packet 1.62 % (20.25 mg/1.25 gram), 1.62 % (40.5 mg/2.5 gram)</i>	T3	PA
<i>testosterone transdermal solution in metered pump w/app</i>	T2	PA
<i>tolvaptan</i>	T5	PA

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TOUJEO MAX U-300 SOLOSTAR	T3	
TOUJEO SOLOSTAR U-300 INSULIN	T3	
TRADJENTA	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-5-1,000 MG, 25-5-1,000 MG	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 12.5-2.5-1,000 MG, 5-2.5-1,000 MG	T3	QL (62 EA per 31 days)
TRULICITY	T3	PA; QL (2 ML per 28 days)
UNITHROID	T3	
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 10-500 MG, 5-500 MG	T3	QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
XULTOPHY 100/3.6	T3	QL (15 ML per 30 days)
YARGESA	T5	PA; QL (93 EA per 31 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML	T5	PA; QL (1.12 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 294 MCG/0.98 ML	T5	PA; QL (1.96 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 420 MCG/1.4 ML	T5	PA; QL (2.8 ML per 28 days)
ZEGALOGUE AUTOINJECTOR	T3	
ZEGALOGUE SYRINGE	T3	
<b>Gastroenterology</b>		
<i>alosetron oral tablet 0.5 mg</i>	T4	PA; QL (93 EA per 31 days)
<i>alosetron oral tablet 1 mg</i>	T5	PA; QL (62 EA per 31 days)
<i>aprepitant oral capsule 125 mg</i>	T5	PA-BvD
<i>aprepitant oral capsule 40 mg, 80 mg</i>	T4	PA-BvD
<i>aprepitant oral capsule, dose pack</i>	T4	PA-BvD
<i>balsalazide</i>	T2	
<i>betaine</i>	T5	
<i>budesonide oral capsule, delayed, extend. release</i>	T4	
<i>budesonide oral tablet, delayed and ext. release</i>	T5	
CIMZIA POWDER FOR RECONST	T5	PA; QL (2 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)</b>	T5	PA; QL (2 EA per 28 days)
<b>COMPRO</b>	T4	
<b>CONSTULOSE</b>	T2	
<b>CREON</b>	T3	
<i>cromolyn oral</i>	T4	
<i>dicyclomine oral capsule</i>	T2	
<i>dicyclomine oral solution</i>	T2	
<i>dicyclomine oral tablet</i>	T2	
<i>diphenoxylate-atropine oral liquid</i>	T4	
<i>diphenoxylate-atropine oral tablet</i>	T2	
<i>dronabinol</i>	T4	PA-BvD
<b>ENULOSE</b>	T2	
<i>esomeprazole magnesium oral capsule, delayed release(dr/ec)</i>	T2	QL (31 EA per 31 days)
<i>famotidine oral suspension for reconstitution</i>	T2	
<i>famotidine oral tablet 20 mg, 40 mg</i>	T1	
<b>GATTEX 30-VIAL</b>	T5	PA
<b>GAVILYTE-C</b>	T1	
<b>GAVILYTE-G</b>	T1	
<b>GAVILYTE-N</b>	T1	
<b>GENERLAC</b>	T2	
<i>glycopyrrolate oral tablet 1 mg, 2 mg</i>	T2	
<i>granisetron hcl oral</i>	T2	PA-BvD
<i>hydrocortisone rectal</i>	T4	
<i>hydrocortisone topical cream with perineal applicator 2.5 %</i>	T2	
<b>IBSRELA</b>	T5	PA; QL (62 EA per 31 days)
<i>lactulose oral solution</i>	T2	
<i>lansoprazole oral capsule, delayed release(dr/ec) 15 mg</i>	T2	QL (31 EA per 31 days)
<i>lansoprazole oral capsule, delayed release(dr/ec) 30 mg</i>	T2	QL (62 EA per 31 days)
<b>LINZESS</b>	T3	QL (31 EA per 31 days)
<i>loperamide oral capsule</i>	T2	
<i>lubiprostone</i>	T3	QL (62 EA per 31 days)
<i>meclizine oral tablet 12.5 mg, 25 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>mesalamine oral capsule (with del rel tablets)</i>	T4	QL (186 EA per 31 days)
<i>mesalamine oral capsule,extended release 24hr</i>	T4	QL (124 EA per 31 days)
<i>mesalamine oral tablet,delayed release (dr/ec) 1.2 gram</i>	T4	QL (124 EA per 31 days)
<i>mesalamine rectal enema</i>	T4	QL (1860 ML per 31 days)
<i>metoclopramide hcl oral solution</i>	T2	
<i>metoclopramide hcl oral tablet</i>	T1	
<i>misoprostol</i>	T2	
<b>MOVANTI</b>	T3	QL (31 EA per 31 days)
<i>nitroglycerin rectal</i>	T4	
<b>OALIVA</b>	T5	PA; QL (31 EA per 31 days)
<i>omeprazole oral capsule,delayed release(dr/ec)</i>	T1	
<i>ondansetron hcl oral solution</i>	T2	PA-BvD
<i>ondansetron hcl oral tablet 4 mg, 8 mg</i>	T2	PA-BvD
<i>ondansetron oral tablet,disintegrating 4 mg, 8 mg</i>	T2	PA-BvD
<i>pantoprazole oral tablet,delayed release (dr/ec)</i>	T1	
<i>peg 3350-electrolytes</i>	T1	
<i>peg3350-sod sul-nacl-kcl-asb-c</i>	T4	
<i>peg-electrolyte soln</i>	T1	
<i>prochlorperazine</i>	T4	
<i>prochlorperazine maleate</i>	T1	
<b>PROCTO-MED HC</b>	T2	
<b>PROCTOSOL HC TOPICAL</b>	T2	
<b>PROCTOZONE-HC</b>	T2	
<i>rabeprazole oral tablet,delayed release (dr/ec)</i>	T2	QL (62 EA per 31 days)
<i>scopolamine base</i>	T3	QL (10 EA per 30 days)
<b>SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML)</b>	T5	PA; QL (1.2 ML per 56 days)
<b>SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 360 MG/2.4 ML (150 MG/ML)</b>	T5	PA; QL (2.4 ML per 56 days)
<i>sodium,potassium,mag sulfates</i>	T4	
<i>sucralfate oral suspension</i>	T4	
<i>sucralfate oral tablet</i>	T2	
<i>sulfasalazine</i>	T2	
<i>ursodiol oral capsule 300 mg</i>	T2	
<i>ursodiol oral tablet</i>	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
VIBERZI	T5	PA; QL (62 EA per 31 days)
VOWST	T5	PA; QL (12 EA per 14 days)
<b>Immunology, Vaccines / Biotechnology</b>		
ABRYSVO (PF)	T3	QL (1 EA per 365 days)
ACTHIB (PF)	T3	
ACTIMMUNE	T5	PA
ADACEL(TDAP ADOLESN/ADULT)(PF)	T3	
AREXVY (PF)	T3	QL (1 EA per 365 days)
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	T5	PA; QL (1 EA per 28 days)
AVONEX INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (1 EA per 28 days)
<i>bcg vaccine, live (pf)</i>	T3	
BESREMI	T5	PA-NS; QL (2 ML per 28 days)
BETASERON SUBCUTANEOUS KIT	T5	PA; QL (14 EA per 28 days)
BEXSERO	T3	
BIVIGAM	T5	PA
BOOSTRIX TDAP	T3	
DAPTACEL (DTAP PEDIATRIC) (PF)	T3	
ENGERIX-B (PF)	T3	PA-BvD
ENGERIX-B PEDIATRIC (PF)	T3	PA-BvD
FULPHILA	T5	
GAMMAGARD LIQUID	T5	PA
GAMMAGARD S-D (IGA < 1 MCG/ML)	T5	PA
GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GAMMAPLEX	T5	PA
GAMMAPLEX (WITH SORBITOL)	T5	PA
GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)	T5	PA
GARDASIL 9 (PF)	T3	
HAVRIX (PF)	T3	
HEPLISAV-B (PF)	T3	PA-BvD
HIBERIX (PF)	T3	
IMOVAX RABIES VACCINE (PF)	T3	PA-BvD
INFANRIX (DTAP) (PF)	T3	
IPOL	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>IXCHIQ (PF)</b>	T3	
<b>IXIARO (PF)</b>	T3	
<b>JYNNEOS (PF)</b>	T3	PA-BvD
<b>KINRIX (PF)</b>	T3	
<b>LEUKINE INJECTION RECON SOLN</b>	T5	PA
<b>MENACTRA (PF) INTRAMUSCULAR SOLUTION</b>	T3	
<b>MENQUADFI (PF)</b>	T3	
<b>MENVEO A-C-Y-W-135-DIP (PF) INTRAMUSCULAR KIT</b>	T3	
<b>M-M-R II (PF)</b>	T3	
<b>MRESVIA (PF)</b>	T3	QL (1 ML per 365 days)
<b>NEULASTA</b>	T5	
<b>NIVESTYM</b>	T5	
<b>NORDITROPIN FLEXPPO</b>	T5	PA
<b>OCTAGAM</b>	T5	PA
<b>PANZYGA</b>	T5	PA
<b>PEDIARIX (PF)</b>	T3	PA-BvD
<b>PEDVAX HIB (PF)</b>	T3	
<b>PEGASYS</b>	T5	PA
<b>PENBRAYA (PF)</b>	T3	
<b>PENTACEL (PF) INTRAMUSCULAR KIT 15LF-20MCG-5LF- 62 DU/0.5 ML</b>	T3	
<b>PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML</b>	T5	PA; QL (1 ML per 28 days)
<b>PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML</b>	T5	PA; QL (1 ML per 28 days)
<b>PRIORIX (PF)</b>	T3	
<b>PRIVIGEN</b>	T5	PA
<b>PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML</b>	T3	PA-BvD
<b>PROCRIT INJECTION SOLUTION 20,000 UNIT/ML, 40,000 UNIT/ML</b>	T5	PA-BvD
<b>PROQUAD (PF)</b>	T3	
<b>QUADRACEL (PF)</b>	T3	
<b>RABAVERT (PF)</b>	T3	PA-BvD
<b>RECOMBIVAX HB (PF)</b>	T3	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
RETACRIT	T3	PA-BvD
ROTARIX ORAL SUSPENSION	T3	
ROTATEQ VACCINE	T3	
SHINGRIX (PF)	T3	QL (2 EA per 999 days)
TENIVAC (PF)	T3	
TICOVAC	T3	
TRUMENBA	T3	
TWINRIX (PF)	T3	
TYPHIM VI	T3	
VAQTA (PF)	T3	
VARIVAX (PF)	T3	
VAXCHORA VACCINE	T3	QL (200 ML per 365 days)
XOLREMDI	T5	PA; QL (124 EA per 31 days)
YF-VAX (PF)	T3	
ZARXIO	T5	
ZIEXTENZO	T5	
<b>Miscellaneous Supplies</b>		
ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"	T3	
GAUZE PAD TOPICAL BANDAGE 2 X 2 "	T3	PA
<i>insulin syringe-needle u-100 syringe 0.3 ml 29 gauge, 1 ml 29 gauge x 1/2", 1/2 ml 28 gauge</i>	T3	
<i>pen needle, diabetic needle 29 gauge x 1/2"</i>	T3	
<b>Musculoskeletal / Rheumatology</b>		
ACTEMRA ACTPEN	T5	PA; QL (3.6 ML per 28 days)
ACTEMRA SUBCUTANEOUS	T5	PA; QL (3.6 ML per 28 days)
<i>alendronate oral tablet 10 mg, 35 mg, 70 mg</i>	T1	
<i>allopurinol oral tablet 100 mg, 300 mg</i>	T1	
BENLYSTA SUBCUTANEOUS	T5	PA; QL (4 ML per 28 days)
<i>colchicine oral tablet</i>	T2	QL (62 EA per 31 days)
CYLTEZO(CF)	T5	PA; QL (2 EA per 28 days)
CYLTEZO(CF) PEN	T5	PA; QL (2 EA per 28 days)
CYLTEZO(CF) PEN CROHN'S-UC-HS	T5	PA; QL (12 EA per 365 days)
CYLTEZO(CF) PEN PSORIASIS-UV	T5	PA; QL (8 EA per 365 days)
ENBREL MINI	T5	PA; QL (8 ML per 28 days)
ENBREL SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5)</b>	T5	PA; QL (4 ML per 28 days)
<b>ENBREL SUBCUTANEOUS SYRINGE 50 MG/ML (1 ML)</b>	T5	PA; QL (8 ML per 28 days)
<b>ENBREL SURECLICK</b>	T5	PA; QL (8 ML per 28 days)
<b>EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML ( 105MG/1.17MLX2)</b>	T5	PA; QL (2.34 ML per 28 days)
<i>febuxostat</i>	T2	PA
<b>HUMIRA PEN</b>	T5	PA; QL (2 EA per 28 days)
<b>HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML</b>	T5	PA; QL (2 EA per 28 days)
<b>HUMIRA(CF)</b>	T5	PA; QL (2 EA per 28 days)
<b>HUMIRA(CF) PEN</b>	T5	PA; QL (2 EA per 28 days)
<b>HUMIRA(CF) PEN CROHNS-UC-HS</b>	T5	PA; QL (6 EA per 365 days)
<b>HUMIRA(CF) PEN PSOR-UV-ADOL HS</b>	T5	PA; QL (6 EA per 365 days)
<i>ibandronate oral</i>	T2	
<b>KEVZARA</b>	T5	PA; QL (2.28 ML per 28 days)
<b>KINERET</b>	T5	PA; QL (18.76 ML per 28 days)
<i>leflunomide</i>	T2	
<b>OLUMIANT</b>	T5	PA; QL (31 EA per 31 days)
<b>ORENCIA CLICKJECT</b>	T5	PA; QL (4 ML per 28 days)
<b>ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML</b>	T5	PA; QL (4 ML per 28 days)
<b>ORENCIA SUBCUTANEOUS SYRINGE 50 MG/0.4 ML</b>	T5	PA; QL (1.6 ML per 28 days)
<b>ORENCIA SUBCUTANEOUS SYRINGE 87.5 MG/0.7 ML</b>	T5	PA; QL (2.8 ML per 28 days)
<b>OTEZLA</b>	T5	PA; QL (62 EA per 31 days)
<b>OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51)</b>	T5	PA; QL (110 EA per 365 days)
<b>OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)</b>	T5	PA; QL (55 EA per 28 days)
<i>penicillamine oral tablet</i>	T5	
<i>probenecid</i>	T2	
<i>probenecid-colchicine</i>	T2	
<b>PROLIA</b>	T3	PA; QL (1 ML per 180 days)
<i>raloxifene</i>	T2	
<b>RINVOQ LQ</b>	T5	PA; QL (372 ML per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<b>RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG</b>	T5	PA; QL (31 EA per 31 days)
<b>RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 45 MG</b>	T5	PA; QL (168 EA per 365 days)
<i>risedronate oral tablet 150 mg, 35 mg, 35 mg (12 pack), 35 mg (4 pack), 5 mg</i>	T4	
<i>risedronate oral tablet, delayed release (dr/ec)</i>	T4	
<b>SAVELLA</b>	T3	PA
<b>SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML</b>	T5	PA; QL (1 ML per 28 days)
<b>SIMPONI SUBCUTANEOUS PEN INJECTOR 50 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 28 days)
<b>SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML</b>	T5	PA; QL (1 ML per 28 days)
<b>SIMPONI SUBCUTANEOUS SYRINGE 50 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 28 days)
<i>teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)</i>	T5	PA; QL (2.48 ML per 28 days)
<b>TYMLOS</b>	T5	PA; QL (1.56 ML per 30 days)
<b>XELJANZ ORAL SOLUTION</b>	T5	PA; QL (310 ML per 31 days)
<b>XELJANZ ORAL TABLET</b>	T5	PA; QL (62 EA per 31 days)
<b>XELJANZ XR</b>	T5	PA; QL (31 EA per 31 days)
<b>YUFLYMA(CF)</b>	T5	PA; QL (2 EA per 28 days)
<b>YUFLYMA(CF) AI CROHN'S-UC-HS</b>	T5	PA; QL (6 EA per 365 days)
<b>YUFLYMA(CF) AUTOINJECTOR</b>	T5	PA; QL (2 EA per 28 days)
<b>Obstetrics / Gynecology</b>		
<b>ALTAVERA (28)</b>	T2	
<b>ALYACEN 1/35 (28)</b>	T2	
<b>APRI</b>	T2	
<b>ARANELLE (28)</b>	T2	
<b>AVIANE</b>	T2	
<b>AZURETTE (28)</b>	T2	
<b>CAMILA</b>	T2	
<i>clindamycin phosphate vaginal</i>	T2	
<b>CRYSELLE (28)</b>	T2	
<b>CYRED EQ</b>	T2	
<b>DEPO-SUBQ PROVERA 104</b>	T3	
<b>DOTTI</b>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>drospirenone-e.estradiol-lm.fa oral tablet 3-0.02-0.451 mg (24) (4)</i>	T2	
<i>drospirenone-ethinyl estradiol</i>	T2	
<b>ELURYNG</b>	T4	
<b>ENPRESSE</b>	T2	
<b>ENSKYCE</b>	T2	
<b>ERRIN</b>	T2	
<b>ESTARYLLA</b>	T2	
<i>estradiol oral</i>	T2	
<i>estradiol transdermal patch semiweekly</i>	T2	
<i>estradiol transdermal patch weekly</i>	T2	
<i>estradiol vaginal</i>	T2	
<i>estradiol-norethindrone acet</i>	T2	
<i>ethynodiol diac-eth estradiol</i>	T2	
<i>etonogestrel-ethinyl estradiol</i>	T3	
<b>GALLIFREY</b>	T2	
<b>HALOETTE</b>	T4	
<b>HEATHER</b>	T2	
<b>ICLEVIA</b>	T2	
<b>IMVEXXY MAINTENANCE PACK</b>	T3	
<b>IMVEXXY STARTER PACK</b>	T3	
<b>INCASSIA</b>	T2	
<b>INTROVALE</b>	T2	
<b>ISIBLOOM</b>	T2	
<b>JASMIEL (28)</b>	T2	
<b>JINTELI</b>	T4	
<b>JULEBER</b>	T2	
<b>KARIVA (28)</b>	T2	
<b>KELNOR 1/35 (28)</b>	T2	
<b>KELNOR 1/50 (28)</b>	T2	
<b>KURVELO (28)</b>	T2	
<i>l norgest/e.estradiol-e.estradiol</i>	T2	
<b>LESSINA</b>	T2	
<b>LEVONEST (28)</b>	T2	
<i>levonorgestrel-ethinyl estradiol oral tablet 0.1-20 mg-mcg, 0.15-0.03 mg</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>levonorgestrel-ethinyl estrad oral tablets,dose pack,3 month</i>	T2	
<i>levonorg-eth estrad triphasic</i>	T2	
<b>LEVORA-28</b>	T2	
<b>LILETTA</b>	T3	
<b>LORYNA (28)</b>	T2	
<b>LOW-OGESTREL (28)</b>	T2	
<b>LUTERA (28)</b>	T2	
<b>LYLEQ</b>	T2	
<b>LYLLANA</b>	T2	
<b>LYZA</b>	T2	
<b>MARLISSA (28)</b>	T2	
<i>medroxyprogesterone</i>	T2	
<i>metronidazole vaginal gel 0.75 % (37.5mg/5 gram)</i>	T3	
<b>MICONAZOLE-3 VAGINAL SUPPOSITORY</b>	T2	
<b>MICROGESTIN 1.5/30 (21)</b>	T2	
<b>MICROGESTIN 1/20 (21)</b>	T2	
<b>MICROGESTIN FE 1.5/30 (28)</b>	T2	
<b>MICROGESTIN FE 1/20 (28)</b>	T2	
<b>MILI</b>	T2	
<b>NEXPLANON</b>	T3	
<i>norethindrone (contraceptive)</i>	T2	
<i>norethindrone acetate</i>	T2	
<i>norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg</i>	T4	
<i>norethindrone ac-eth estradiol oral tablet 1-20 mg-mcg</i>	T2	
<i>norethindrone-e.estradiol-iron oral tablet 1 mg-20 mcg (21)/75 mg (7), 1-20(5)/1-30(7) /1mg-35mcg (9)</i>	T2	
<i>norgestimate-ethinyl estradiol</i>	T2	
<b>NORTREL 0.5/35 (28)</b>	T2	
<b>NORTREL 1/35 (21)</b>	T2	
<b>NORTREL 1/35 (28)</b>	T2	
<b>NORTREL 7/7/7 (28)</b>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<b>PIMTREA (28)</b>	T2	
<b>PORTIA 28</b>	T2	
<b>PREMARIN ORAL</b>	T3	
<b>PREMARIN VAGINAL</b>	T3	
<b>PREMPRO</b>	T3	
<i>progesterone micronized</i>	T2	
<b>RECLIPSEN (28)</b>	T2	
<b>SETLAKIN</b>	T2	
<b>SPRINTEC (28)</b>	T2	
<b>SRONYX</b>	T2	
<b>SYEDA</b>	T2	
<i>terconazole</i>	T2	
<b>TILIA FE</b>	T2	
<i>tranexamic acid oral</i>	T3	
<b>TRI-ESTARYLLA</b>	T2	
<b>TRI-LEGEST FE</b>	T2	
<b>TRI-LO-ESTARYLLA</b>	T2	
<b>TRI-LO-SPRINTEC</b>	T2	
<b>TRI-SPRINTEC (28)</b>	T2	
<b>TRIVORA (28)</b>	T2	
<b>TURQOZ (28)</b>	T2	
<b>VELIVET TRIPHASIC REGIMEN (28)</b>	T2	
<b>VESTURA (28)</b>	T2	
<b>VIENVA</b>	T2	
<b>YUVAFEM</b>	T3	
<b>ZAFEMY</b>	T3	
<b>ZOVIA 1-35 (28)</b>	T2	
<b>Ophthalmology</b>		
<i>acetazolamide</i>	T2	
<b>ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.1 %</b>	T3	
<i>apraclonidine</i>	T3	
<i>atropine ophthalmic (eye) drops 1 %</i>	T2	
<i>azelastine ophthalmic (eye)</i>	T2	
<i>bacitracin ophthalmic (eye)</i>	T2	
<i>bacitracin-polymyxin b</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>BESIVANCE</b>	T3	
<i>betaxolol ophthalmic (eye)</i>	T2	
<i>bimatoprost ophthalmic (eye)</i>	T2	
<i>brimonidine ophthalmic (eye) drops 0.1 %</i>	T3	
<i>brimonidine ophthalmic (eye) drops 0.15 %, 0.2 %</i>	T2	
<i>brimonidine-timolol</i>	T3	
<i>brinzolamide</i>	T4	
<i>bromfenac ophthalmic (eye) drops 0.07 %, 0.075 %</i>	T3	
<i>bromfenac ophthalmic (eye) drops 0.09 %</i>	T2	
<i>carteolol</i>	T2	
<i>ciprofloxacin hcl ophthalmic (eye)</i>	T2	
<b>COMBIGAN</b>	T3	
<i>cromolyn ophthalmic (eye)</i>	T2	
<i>cyclosporine ophthalmic (eye)</i>	T3	QL (60 EA per 30 days)
<b>CYSTARAN</b>	T5	PA; QL (60 ML per 28 days)
<i>dexamethasone sodium phosphate ophthalmic (eye)</i>	T2	
<i>diclofenac sodium ophthalmic (eye)</i>	T2	
<i>difluprednate</i>	T4	
<i>dorzolamide</i>	T2	
<i>dorzolamide-timolol</i>	T2	
<i>dorzolamide-timolol (pf) ophthalmic (eye) dropperette</i>	T2	
<i>erythromycin ophthalmic (eye)</i>	T2	
<i>fluorometholone</i>	T2	
<i>flurbiprofen sodium</i>	T2	
<i>gatifloxacin</i>	T3	
<i>gentamicin ophthalmic (eye) drops</i>	T2	
<i>ketorolac ophthalmic (eye)</i>	T2	
<i>latanoprost</i>	T1	
<i>levobunolol ophthalmic (eye) drops 0.5 %</i>	T2	
<b>LUMIGAN OPHTHALMIC (EYE) DROPS 0.01 %</b>	T3	QL (5 ML per 31 days)
<i>methazolamide</i>	T4	
<i>moxifloxacin ophthalmic (eye) drops</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<i>neomycin-bacitracin-poly-hc</i>	T2	
<i>neomycin-bacitracin-polymyxin</i>	T2	
<i>neomycin-polymyxin b-dexameth</i>	T2	
<i>neomycin-polymyxin-gramicidin</i>	T2	
<i>neomycin-polymyxin-hc ophthalmic (eye)</i>	T3	
<b>NEO-POLYCIN</b>	T2	
<b>NEO-POLYCIN HC</b>	T2	
<i>ofloxacin ophthalmic (eye)</i>	T2	
<i>pilocarpine hcl ophthalmic (eye) drops 1 %, 2 %, 4 %</i>	T2	
<b>POLYCIN</b>	T2	
<i>polymyxin b sulf-trimethoprim</i>	T2	
<i>prednisolone acetate</i>	T2	
<i>prednisolone sodium phosphate ophthalmic (eye)</i>	T2	
<b>PROLENSA</b>	T3	
<b>RESTASIS</b>	T3	QL (60 EA per 30 days)
<b>RESTASIS MULTIDOSE</b>	T3	QL (5.5 ML per 27 days)
<b>SIMBRINZA</b>	T4	
<i>sulfacetamide sodium ophthalmic (eye)</i>	T2	
<i>sulfacetamide-prednisolone</i>	T2	
<i>timolol maleate ophthalmic (eye) drops</i>	T1	
<i>timolol maleate ophthalmic (eye) gel forming solution</i>	T3	
<i>tobramycin ophthalmic (eye)</i>	T2	
<i>tobramycin-dexamethasone</i>	T3	
<i>travoprost</i>	T3	
<i>trifluridine</i>	T3	
<b>XDEMVY</b>	T5	PA; QL (10 ML per 42 days)
<b>XIIDRA</b>	T3	QL (60 EA per 30 days)
<b>ZIRGAN</b>	T4	ST
<b>Respiratory And Allergy</b>		
<i>acetylcysteine</i>	T2	PA-BvD
<b>ADEMPAS</b>	T5	PA; QL (93 EA per 31 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation</i>	T2	QL (17 GM per 30 days)
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020503)</i>	T2	QL (13.4 GM per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.



<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020983)</i>	NF	
<i>albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg/3 ml (0.083 %), 2.5 mg/0.5 ml</i>	T2	PA-BvD
<i>albuterol sulfate oral syrup</i>	T2	
<i>albuterol sulfate oral tablet</i>	T4	
<b>ALYQ</b>	T5	PA; QL (62 EA per 31 days)
<i>ambrisentan</i>	T5	PA; QL (31 EA per 31 days)
<b>ANORO ELLIPTA</b>	T3	QL (60 EA per 30 days)
<i>arformoterol</i>	T3	PA-BvD
<b>ASMANEX HFA</b>	T3	QL (13 GM per 30 days)
<b>ASMANEX TWISTHALER INHALATION AEROSOL POWDR BREATH ACTIVATED 110 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (120), 220 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (60)</b>	T3	QL (1 EA per 30 days)
<b>ATROVENT HFA</b>	T3	QL (25.8 GM per 30 days)
<i>azelastine-fluticasone</i>	T4	QL (23 GM per 30 days)
<b>BREO ELLIPTA</b>	T3	QL (60 EA per 30 days)
<b>BREYNA</b>	T3	QL (10.3 GM per 30 days)
<b>BREZTRI AEROSPHERE</b>	T3	QL (10.7 GM per 30 days)
<i>budesonide inhalation</i>	T4	PA-BvD
<i>budesonide-formoterol</i>	T3	QL (10.2 GM per 30 days)
<i>cetirizine oral solution 1 mg/ml</i>	T2	QL (310 ML per 31 days)
<b>CINRYZE</b>	T5	PA; QL (20 EA per 28 days)
<b>COMBIVENT RESPIMAT</b>	T3	QL (4 GM per 30 days)
<i>cromolyn inhalation</i>	T4	PA-BvD
<i>desloratadine oral tablet</i>	T2	QL (31 EA per 31 days)
<b>DULERA</b>	T3	QL (13 GM per 30 days)
<i>epinephrine injection auto-injector 0.15 mg/0.3 ml, 0.3 mg/0.3 ml</i>	T3	
<b>FASENRA PEN</b>	T5	PA; QL (1 ML per 56 days)
<b>FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML</b>	T5	PA; QL (0.5 ML per 56 days)
<b>FASENRA SUBCUTANEOUS SYRINGE 30 MG/ML</b>	T5	PA; QL (1 ML per 56 days)
<i>flunisolide</i>	T2	QL (50 ML per 25 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>fluticasone propionate inhalation blister with device 100 mcg/actuation, 50 mcg/actuation</i>	T4	ST; QL (60 EA per 30 days)
<i>fluticasone propionate inhalation blister with device 250 mcg/actuation</i>	T4	ST; QL (240 EA per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 110 mcg/actuation</i>	T4	ST; QL (12 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 220 mcg/actuation</i>	T4	ST; QL (24 GM per 30 days)
<i>fluticasone propionate inhalation hfa aerosol inhaler 44 mcg/actuation</i>	T4	ST; QL (10.6 GM per 30 days)
<i>fluticasone propionate nasal</i>	T2	QL (16 GM per 30 days)
<i>fluticasone propion-salmeterol inhalation aerosol powdr breath activated</i>	T3	QL (1 EA per 30 days)
<i>fluticasone propion-salmeterol inhalation blister with device</i>	T3	QL (60 EA per 30 days)
<i>hydroxyzine hcl oral tablet</i>	T2	PA
<i>icatibant</i>	T5	PA; QL (18 ML per 30 days)
<i>ipratropium bromide inhalation</i>	T2	PA-BvD
<i>ipratropium-albuterol</i>	T2	PA-BvD
<b>KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 5.8 MG, 50 MG, 75 MG</b>	T5	PA; QL (56 EA per 28 days)
<b>KALYDECO ORAL GRANULES IN PACKET 25 MG</b>	T5	PA; QL (62 EA per 31 days)
<b>KALYDECO ORAL TABLET</b>	T5	PA; QL (62 EA per 31 days)
<i>levalbuterol hcl inhalation solution for nebulization 1.25 mg/3 ml</i>	T3	PA-BvD
<i>levalbuterol tartrate</i>	T3	QL (30 GM per 30 days)
<i>levocetirizine oral solution</i>	T4	QL (310 ML per 31 days)
<i>levocetirizine oral tablet</i>	T2	QL (31 EA per 31 days)
<i>mometasone nasal</i>	T2	QL (34 GM per 30 days)
<i>montelukast oral tablet</i>	T2	QL (31 EA per 31 days)
<i>montelukast oral tablet,chewable</i>	T2	QL (31 EA per 31 days)
<b>NUCALA SUBCUTANEOUS AUTO-INJECTOR</b>	T5	PA; QL (3 ML per 28 days)
<b>NUCALA SUBCUTANEOUS RECON SOLN</b>	T5	PA; QL (3 EA per 28 days)
<b>NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML</b>	T5	PA; QL (3 ML per 28 days)
<b>NUCALA SUBCUTANEOUS SYRINGE 40 MG/0.4 ML</b>	T5	PA; QL (0.4 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>OFEV</b>	T5	PA; QL (62 EA per 31 days)
<b>OPSUMIT</b>	T5	PA; QL (31 EA per 31 days)
<b>OPSYNVI</b>	T5	PA; QL (31 EA per 31 days)
<b>ORKAMBI ORAL GRANULES IN PACKET</b>	T5	PA; QL (62 EA per 31 days)
<b>ORKAMBI ORAL TABLET</b>	T5	PA; QL (124 EA per 31 days)
<i>pirfenidone oral capsule</i>	T5	PA; QL (279 EA per 31 days)
<i>pirfenidone oral tablet</i>	T5	PA; QL (93 EA per 31 days)
<i>promethazine oral</i>	T4	PA
<b>PULMOZYME</b>	T5	PA
<b>QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION</b>	T3	QL (10.6 GM per 30 days)
<b>QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION</b>	T3	QL (21.2 GM per 30 days)
<i>roflumilast</i>	T4	QL (31 EA per 31 days)
<b>SAJAZIR</b>	T5	PA; QL (18 ML per 30 days)
<b>SEREVENT DISKUS</b>	T3	QL (60 EA per 30 days)
<i>sildenafil (pulm.hypertension) oral tablet</i>	T3	PA; QL (372 EA per 31 days)
<b>SPIRIVA RESPIMAT</b>	T3	QL (4 GM per 30 days)
<b>SPIRIVA WITH HANDIHALER</b>	T3	QL (30 EA per 30 days)
<b>STIOLTO RESPIMAT</b>	T3	QL (4 GM per 30 days)
<b>STRIVERDI RESPIMAT</b>	T4	QL (4 GM per 30 days)
<b>SYMDEKO</b>	T5	PA; QL (56 EA per 28 days)
<i>tadalafil (pulm. hypertension)</i>	T5	PA; QL (62 EA per 31 days)
<i>terbutaline oral</i>	T4	
<b>THEO-24</b>	T3	
<i>theophylline oral solution</i>	T2	
<i>theophylline oral tablet extended release 12 hr</i>	T2	
<i>theophylline oral tablet extended release 24 hr</i>	T2	
<b>TRELEGY ELLIPTA</b>	T3	QL (60 EA per 30 days)
<b>TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL</b>	T5	PA; QL (56 EA per 28 days)
<b>TRIKAFTA ORAL TABLETS, SEQUENTIAL</b>	T5	PA; QL (84 EA per 28 days)
<b>VENTOLIN HFA</b>	T3	QL (36 GM per 30 days)
<b>WIXELA INHUB</b>	T3	QL (60 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<b>XOLAIR</b>	T5	PA
<i>zafirlukast oral tablet 10 mg</i>	T4	QL (93 EA per 31 days)
<i>zafirlukast oral tablet 20 mg</i>	T4	QL (62 EA per 31 days)
<b>Urologicals</b>		
<i>alfuzosin</i>	T2	QL (31 EA per 31 days)
<i>bethanechol chloride oral tablet 10 mg, 25 mg, 5 mg</i>	T2	
<i>bethanechol chloride oral tablet 50 mg</i>	T3	
<b>CYSTAGON</b>	T4	
<i>dutasteride</i>	T2	QL (31 EA per 31 days)
<i>dutasteride-tamsulosin</i>	T4	QL (31 EA per 31 days)
<b>ELMIRON</b>	T4	
<i>finasteride oral tablet 5 mg</i>	T2	
<b>MYRBETRIQ ORAL SUSPENSION,EXTENDED REL RECON</b>	T3	QL (300 ML per 30 days)
<b>MYRBETRIQ ORAL TABLET EXTENDED RELEASE 24 HR</b>	T3	QL (31 EA per 31 days)
<i>oxybutynin chloride oral syrup</i>	T3	
<i>oxybutynin chloride oral tablet 5 mg</i>	T3	
<i>oxybutynin chloride oral tablet extended release 24hr 10 mg, 5 mg</i>	T3	QL (31 EA per 31 days)
<i>oxybutynin chloride oral tablet extended release 24hr 15 mg</i>	T3	QL (62 EA per 31 days)
<i>potassium citrate oral tablet extended release</i>	T2	
<b>RIVFLOZA SUBCUTANEOUS SOLUTION</b>	T5	PA; QL (1 ML per 28 days)
<b>RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML</b>	T5	PA; QL (0.8 ML per 28 days)
<b>RIVFLOZA SUBCUTANEOUS SYRINGE 160 MG/ML</b>	T5	PA; QL (1 ML per 28 days)
<i>silodosin</i>	T4	
<i>tadalafil oral tablet 2.5 mg</i>	T4	PA; QL (62 EA per 31 days)
<i>tadalafil oral tablet 5 mg</i>	T4	PA; QL (31 EA per 31 days)
<i>tamsulosin</i>	T1	
<i>tolterodine oral capsule,extended release 24hr</i>	T3	QL (31 EA per 31 days)
<i>tolterodine oral tablet</i>	T2	QL (62 EA per 31 days)
<i>trospium oral capsule,extended release 24hr</i>	T3	QL (31 EA per 31 days)
<i>trospium oral tablet</i>	T2	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
<b>Vitamins, Hematinics / Electrolytes</b>		
<b>CLINIMIX 5%/D15W SULFITE FREE</b>	T4	PA-BvD
<b>CLINIMIX 4.25%/D10W SULF FREE</b>	T4	PA-BvD
<b>CLINIMIX 5%-D20W(SULFITE-FREE)</b>	T4	PA-BvD
<i>fluoride (sodium) oral tablet</i>	T2	
<b>INTRALIPID INTRAVENOUS EMULSION 20 %</b>	T4	PA-BvD
<b>ISOLYTE S PH 7.4</b>	T3	PA-BvD
<b>ISOLYTE-P IN 5 % DEXTROSE</b>	T4	PA-BvD
<b>KLOR-CON</b>	T4	
<b>KLOR-CON M10</b>	T1	
<b>KLOR-CON M15</b>	T2	
<b>KLOR-CON M20</b>	T1	
<i>magnesium sulfate injection</i>	T2	
<b>PLENAMINE</b>	T4	PA-BvD
<i>potassium chlorid-d5-0.45%nacl</i>	T2	
<i>potassium chloride in 0.9%nacl intravenous parenteral solution 20 meq/l, 40 meq/l</i>	T2	
<i>potassium chloride in 5 % dex intravenous parenteral solution 20 meq/l</i>	T2	
<i>potassium chloride in lr-d5 intravenous parenteral solution 20 meq/l</i>	T2	
<i>potassium chloride in water intravenous piggyback 10 meq/100 ml, 20 meq/100 ml, 40 meq/100 ml</i>	T2	
<i>potassium chloride intravenous</i>	T2	
<i>potassium chloride oral capsule, extended release</i>	T1	
<i>potassium chloride oral liquid</i>	T2	
<i>potassium chloride oral packet</i>	T2	
<i>potassium chloride oral tablet extended release 10 meq, 20 meq, 8 meq</i>	T1	
<i>potassium chloride oral tablet,er particles/crystals 10 meq, 20 meq</i>	T1	
<i>potassium chloride oral tablet,er particles/crystals 15 meq</i>	T2	
<i>potassium chloride-0.45 % nacl</i>	T2	
<i>potassium chloride-d5-0.2%nacl intravenous parenteral solution 20 meq/l</i>	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

<b>Drug Name</b>	<b>Drug Tier</b>	<b>Requirements/Limits</b>
<i>potassium chloride-d5-0.9%nacl</i>	T2	
<b>PRENATAL VITAMIN PLUS LOW IRON</b>	T2	PA
<i>sodium chloride 0.45 % intravenous</i>	T2	
<i>sodium chloride 3 % hypertonic</i>	T2	
<i>sodium chloride 5 % hypertonic</i>	T2	
<b>TRAVASOL 10 %</b>	T4	PA-BvD
<b>TROPHAMINE 10 %</b>	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

## Index of Drugs

<i>abacavir</i> .....	3	<i>amiodarone</i> .....	32	<i>azathioprine</i> .....	11
<i>abacavir-lamivudine</i> .....	3	<i>amitriptyline</i> .....	19	<i>azelastine</i> .....	44, 58
<b>ABILIFY MAINTENA</b> .....	19	<i>amlodipine</i> .....	33	<i>azelastine-fluticasone</i> .....	61
<i>abiraterone</i> .....	11	<i>amlodipine-atorvastatin</i> .....	33	<i>azithromycin</i> .....	4
<b>ABRYSVO (PF)</b> .....	51	<i>amlodipine-benazepril</i> .....	33	<i>aztreonam</i> .....	4
<i>acamprosate</i> .....	42	<i>amlodipine-olmesartan</i> .....	33	<b>AZURETTE (28)</b> .....	55
<i>acarbose</i> .....	44	<i>amlodipine-valsartan</i> .....	33	<i>bacitracin</i> .....	58
<b>ACCUTANE</b> .....	38	<i>ammonium lactate</i> .....	39	<i>bacitracin-polymyxin b</i> .....	58
<i>acebutolol</i> .....	32	<b>AMNESTEEM</b> .....	39	<i>baclofen</i> .....	20
<i>acetaminophen-codeine</i> .....	19	<i>amoxapine</i> .....	19	<b>BAFIERTAM</b> .....	20
<i>acetazolamide</i> .....	58	<i>amoxicillin</i> .....	3	<i>balsalazide</i> .....	48
<i>acetic acid</i> .....	44	<i>amoxicillin-pot clavulanate</i> .....	3, 4	<b>BALVERSA</b> .....	11
<i>acetylcysteine</i> .....	60	<i>amphotericin b</i> .....	4	<b>BAQSIMI</b> .....	44
<i>acitretin</i> .....	38	<i>amphotericin b liposome</i> .....	4	<i>bcg vaccine, live (pf)</i> .....	51
<b>ACTEMRA</b> .....	53	<i>ampicillin</i> .....	4	<i>benazepril</i> .....	33
<b>ACTEMRA ACTPEN</b> .....	53	<i>ampicillin sodium</i> .....	4	<i>benazepril-hydrochlorothiazide</i> .....	33
<b>ACTHIB (PF)</b> .....	51	<i>ampicillin-sulbactam</i> .....	4	<b>BENLYSTA</b> .....	53
<b>ACTIMMUNE</b> .....	51	<i>anagrelide</i> .....	42	<i>benztropine</i> .....	20
<i>acyclovir</i> .....	3, 38	<i>anastrozole</i> .....	11	<b>BESIVANCE</b> .....	59
<i>acyclovir sodium</i> .....	3	<b>ANORO ELLIPTA</b> .....	61	<b>BESREMI</b> .....	51
<b>ADACEL(TDAP</b>		<i>apraclonidine</i> .....	58	<i>betaine</i> .....	48
<b>ADOLESN/ADULT)(PF)</b> .....	51	<i>aprepitant</i> .....	48	<i>betamethasone dipropionate</i> .....	39
<b>ADBRY</b> .....	39	<b>APRI</b> .....	55	<i>betamethasone valerate</i> .....	39
<i>adefovir</i> .....	3	<b>APTiom</b> .....	19	<i>betamethasone, augmented</i> .....	39
<b>ADEMPAS</b> .....	60	<b>APTIVUS</b> .....	4	<b>BETASERON</b> .....	51
<b>AIMOVIG AUTOINJECTOR</b> .....	19	<b>ARANELLE (28)</b> .....	55	<i>betaxolol</i> .....	59
<b>AJOVY AUTOINJECTOR</b> .....	19	<b>AREXVY (PF)</b> .....	51	<i>bethanechol chloride</i> .....	64
<b>AJOVY SYRINGE</b> .....	19	<i>arformoterol</i> .....	61	<i>bexarotene</i> .....	11
<b>AKEEGA</b> .....	11	<b>ARIKAYCE</b> .....	4	<b>BEXSERO</b> .....	51
<b>ALA-CORT</b> .....	39	<i>aripiprazole</i> .....	19	<i>bicalutamide</i> .....	11
<i>albendazole</i> .....	3	<i>armodafinil</i> .....	19	<b>BICILLIN C-R</b> .....	4
<i>albuterol sulfate</i> .....	60, 61	<i>asenapine maleate</i> .....	19	<b>BICILLIN L-A</b> .....	4
<i>alclometasone</i> .....	39	<b>ASMANEX HFA</b> .....	61	<b>BIKTARVY</b> .....	4
<b>ALCOHOL PADS</b> .....	44	<b>ASMANEX TWISTHALER</b> .....	61	<i>bimatoprost</i> .....	59
<b>ALECENSA</b> .....	11	<i>aspirin-dipyridamole</i> .....	33	<i>bisoprolol fumarate</i> .....	33
<i>alendronate</i> .....	53	<b>ASSURE ID INSULIN</b>		<i>bisoprolol-hydrochlorothiazide</i> .....	33
<i>alfuzosin</i> .....	64	<b>SAFETY</b> .....	53	<b>BIVIGAM</b> .....	51
<i>aliskiren</i> .....	32	<i>atazanavir</i> .....	4	<b>BOOSTRIX TDAP</b> .....	51
<i>allopurinol</i> .....	53	<i>atenolol</i> .....	33	<b>BOSULIF</b> .....	11
<i>alosetron</i> .....	48	<i>atenolol-chlorthalidone</i> .....	33	<b>BRAFTOVI</b> .....	11
<b>ALPHAGAN P</b> .....	58	<i>atomoxetine</i> .....	20	<b>BREO ELLIPTA</b> .....	61
<i>alprazolam</i> .....	19	<i>atorvastatin</i> .....	33	<b>BREYNA</b> .....	61
<b>ALTAVERA (28)</b> .....	55	<i>atovaquone</i> .....	4	<b>BREZTRI AEROSPHERE</b> .....	61
<b>ALUNBRIG</b> .....	11	<i>atovaquone-proguanil</i> .....	4	<b>BRILINTA</b> .....	33
<b>ALYACEN 1/35 (28)</b> .....	55	<i>atropine</i> .....	58	<i>brimonidine</i> .....	59
<b>ALYQ</b> .....	61	<b>ATROVENT HFA</b> .....	61	<i>brimonidine-timolol</i> .....	59
<i>amantadine hcl</i> .....	3	<b>AUGTYRO</b> .....	11	<i>brinzolamide</i> .....	59
<i>ambrisentan</i> .....	61	<b>AUVELITY</b> .....	20	<b>BRIVIACT</b> .....	20
<i>amikacin</i> .....	3	<b>AVIANE</b> .....	55	<i>bromfenac</i> .....	59
<i>amiloride</i> .....	32	<b>AVONEX</b> .....	51	<i>bromocriptine</i> .....	20
<i>amiloride-hydrochlorothiazide</i> .....	32	<b>AYVAKIT</b> .....	11	<b>BRUKINSA</b> .....	11



<i>budesonide</i> .....	48, 61	<i>cevimeline</i> .....	42	<i>colesevelam</i> .....	33
<i>budesonide-formoterol</i> .....	61	<b>CHEMET</b> .....	42	<i>colestipol</i> .....	33
<i>bumetanide</i> .....	33	<i>chlorhexidine gluconate</i> .....	44	<i>colistin (colistimethate na)</i> .....	5
<i>buprenorphine</i> .....	20	<i>chloroquine phosphate</i> .....	5	<b>COMBIGAN</b> .....	59
<i>buprenorphine hcl</i> .....	20	<i>chlorpromazine</i> .....	21	<b>COMBIVENT RESPIMAT</b> ....	61
<i>buprenorphine-naloxone</i> .....	20	<i>chlorthalidone</i> .....	33	<b>COMETRIQ</b> .....	11, 12
<i>bupropion hcl</i> .....	20	<i>cholestyramine (with sugar)</i> .....	33	<b>COMPLERA</b> .....	5
<i>bupropion hcl (smoking deter)</i> ....	42	<b>CHOLESTYRAMINE</b>		<b>COMPRO</b> .....	49
<i>buspirone</i> .....	20	<b>LIGHT</b> .....	33	<b>CONSTULOSE</b> .....	49
<i>butorphanol</i> .....	20	<b>CIBINQO</b> .....	39	<b>COPAXONE</b> .....	21
<i>cabergoline</i> .....	44	<i>ciclopirox</i> .....	39	<b>COPIKTRA</b> .....	12
<b>CABLIVI</b> .....	33	<i>cilostazol</i> .....	33	<b>CORLANOR</b> .....	33
<b>CABOMETYX</b> .....	11	<b>CIMDUO</b> .....	5	<b>COSENTYX</b> .....	39
<i>calcipotriene</i> .....	39	<b>CIMZIA</b> .....	49	<b>COSENTYX (2 SYRINGES)</b> ... 39	
<i>calcitonin (salmon)</i> .....	44	<b>CIMZIA POWDER FOR</b>		<b>COSENTYX PEN (2 PENS)</b> .... 39	
<i>calcitriol</i> .....	44	<b>RECONST</b> .....	48	<b>COSENTYX UNOREADY</b>	
<b>CALQUENCE</b> .....	11	<i>cinacalcet</i> .....	44	<b>PEN</b> .....	39
<b>CALQUENCE</b>		<b>CINRYZE</b> .....	61	<b>COTELLIC</b> .....	12
<b>(ACALABRUTINIB MAL)</b> ....	11	<i>ciprofloxacin hcl</i> .....	5, 59	<b>CREON</b> .....	49
<b>CAMILA</b> .....	55	<i>ciprofloxacin in 5 % dextrose</i> .....	5	<i>cromolyn</i> .....	49, 59, 61
<b>CAMZYOS</b> .....	33	<i>ciprofloxacin-dexamethasone</i> ....	44	<b>CRYSSELLE (28)</b> .....	55
<i>candesartan</i> .....	33	<i>citalopram</i> .....	21	<i>cyclobenzaprine</i> .....	21
<i>candesartan-hydrochlorothiazid</i> ..	33	<b>CLARAVIS</b> .....	39	<i>cyclophosphamide</i> .....	12
<b>CAPLYTA</b> .....	20	<i>clarithromycin</i> .....	5	<i>cyclosporine</i> .....	12, 59
<b>CAPRELSA</b> .....	11	<i>clindamycin hcl</i> .....	5	<i>cyclosporine modified</i> .....	12
<i>captopril</i> .....	33	<i>clindamycin in 5 % dextrose</i> .....	5	<b>CYLTEZO(CF)</b> .....	53
<i>carbamazepine</i> .....	20	<b>CLINDAMYCIN</b>		<b>CYLTEZO(CF) PEN</b> .....	53
<i>carbidopa-levodopa</i> .....	20	<b>PEDIATRIC</b> .....	5	<b>CYLTEZO(CF) PEN</b>	
<i>carbidopa-levodopa-entacapone</i> ..	20	<i>clindamycin phosphate</i> .....	5, 39, 55	<b>CROHN'S-UC-HS</b> .....	53
<i>carglumic acid</i> .....	42	<b>CLINIMIX 5%/D15W</b>		<b>CYLTEZO(CF) PEN</b>	
<i>carteolol</i> .....	59	<b>SULFITE FREE</b> .....	65	<b>PSORIASIS-UV</b> .....	53
<b>CARTIA XT</b> .....	33	<b>CLINIMIX 4.25%/D10W</b>		<b>CYRED EQ</b> .....	55
<i>carvedilol</i> .....	33	<b>SULF FREE</b> .....	65	<b>CYSTAGON</b> .....	64
<i>caspofungin</i> .....	4	<b>CLINIMIX 4.25%/D5W</b>		<b>CYSTARAN</b> .....	59
<b>CAYSTON</b> .....	4	<b>SULFIT FREE</b> .....	42	<i>d10 %-0.45 % sodium chloride</i> ... 42	
<i>cefaclor</i> .....	4	<b>CLINIMIX 5%-</b>		<i>d2.5 %-0.45 % sodium chloride</i> .. 42	
<i>cefadroxil</i> .....	4	<b>D20W(SULFITE-FREE)</b> .....	65	<i>d5 % and 0.9 % sodium</i>	
<i>cefazolin</i> .....	4	<i>clobazam</i> .....	21	<i>chloride</i> .....	42
<i>cefdinir</i> .....	4	<i>clobetasol</i> .....	39	<i>d5 %-0.45 % sodium chloride</i> .... 42	
<i>cefepime</i> .....	4	<i>clomipramine</i> .....	21	<i>dalfampridine</i> .....	21
<i>cefixime</i> .....	4	<i>clonazepam</i> .....	21	<i>danazol</i> .....	44
<i>cefoxitin</i> .....	4	<i>clonidine</i> .....	33	<i>dantrolene</i> .....	21
<i>cefpodoxime</i> .....	4	<i>clonidine hcl</i> .....	33	<b>DANZITEN</b> .....	12
<i>cefprozil</i> .....	4	<i>clopidogrel</i> .....	33	<i>dapsone</i> .....	5
<i>ceftazidime</i> .....	5	<i>clorazepate dipotassium</i> .....	21	<b>DAPTACEL (DTAP</b>	
<i>ceftriaxone</i> .....	5	<i>clotrimazole</i> .....	5, 39	<b>PEDIATRIC) (PF)</b> .....	51
<i>cefuroxime axetil</i> .....	5	<i>clotrimazole-betamethasone</i> .....	39	<i>daptomycin</i> .....	5
<i>cefuroxime sodium</i> .....	5	<i>clozapine</i> .....	21	<i>darunavir</i> .....	5
<i>celecoxib</i> .....	20, 21	<b>COARTEM</b> .....	5	<i>dasatinib</i> .....	12
<i>cephalexin</i> .....	5	<b>COBENFY</b> .....	21	<b>DAURISMO</b> .....	12
<b>CERDELGA</b> .....	44	<b>COBENFY STARTER PACK</b> ..	21	<b>DAYBUE</b> .....	21
<i>cetirizine</i> .....	61	<i>colchicine</i> .....	53	<i>deferasirox</i> .....	42

<i>deferiprone</i> .....	42	<i>doxycycline monohydrate</i> .....	6	<b>EPIDIOLEX</b> .....	23
<b>DELSTRIGO</b> .....	5	<b>DRIZALMA SPRINKLE</b> .....	23	<i>epinephrine</i> .....	61
<b>DEPO-SUBQ PROVERA 104</b> .....	55	<i>dronabinol</i> .....	49	<b>EPITOL</b> .....	23
<b>DESCOVY</b> .....	5	<i>drospirenone-e.estradiol-lm.fa</i> ...	56	<i>eplerenone</i> .....	34
<i>desipramine</i> .....	21	<i>drospirenone-ethinyl estradiol</i> ...	56	<b>EPRONTIA</b> .....	23
<i>desloratadine</i> .....	61	<i>droxidopa</i> .....	43	<i>ergotamine-caffeine</i> .....	23
<i>desmopressin</i> .....	44	<b>DULERA</b> .....	61	<b>ERIVEDGE</b> .....	12
<i>desoximetasone</i> .....	39	<i>duloxetine</i> .....	23	<b>ERLEADA</b> .....	12
<i>desvenlafaxine succinate</i> .....	22	<b>DUPIXENT PEN</b> .....	40	<i>erlotinib</i> .....	12
<i>dexamethasone</i> .....	44	<b>DUPIXENT SYRINGE</b> .....	40	<b>ERRIN</b> .....	56
<i>dexamethasone sodium</i>		<i>dutasteride</i> .....	64	<i>ertapenem</i> .....	6
<i>phosphate</i> .....	59	<i>dutasteride-tamsulosin</i> .....	64	<b>ERY PADS</b> .....	40
<i>dexmethylphenidate</i> .....	22	<b>DUVYZAT</b> .....	43	<b>ERY-TAB</b> .....	6
<i>dextroamphetamine-</i>		<i>econazole nitrate</i> .....	40	<i>erythromycin</i> .....	6, 59
<i>amphetamine</i> .....	22	<b>EDURANT</b> .....	6	<i>erythromycin ethylsuccinate</i> .....	6
<i>dextrose 10 % in water (d10w)</i> ...	42	<i>efavirenz</i> .....	6	<i>erythromycin with ethanol</i> .....	40
<i>dextrose 5 % in water (d5w)</i> .....	43	<i>efavirenz-emtricitabin-tenofov</i> .....	6	<i>escitalopram oxalate</i> .....	23
<b>DIACOMIT</b> .....	22	<i>efavirenz-lamivu-tenofov disop</i> .....	6	<i>esomeprazole magnesium</i> .....	49
<i>diazepam</i> .....	22	<b>ELIGARD</b> .....	12	<b>ESTARYLLA</b> .....	56
<b>DIAZEPAM INTENSOL</b> .....	22	<b>ELIGARD (3 MONTH)</b> .....	12	<i>estradiol</i> .....	56
<i>diazoxide</i> .....	44	<b>ELIGARD (4 MONTH)</b> .....	12	<i>estradiol-norethindrone acet</i> .....	56
<i>diclofenac potassium</i> .....	22	<b>ELIGARD (6 MONTH)</b> .....	12	<i>eszopiclone</i> .....	23
<i>diclofenac sodium</i> .....	22, 40, 59	<b>ELIQUIS</b> .....	34	<i>ethacrynic acid</i> .....	34
<i>dicloxacillin</i> .....	5	<b>ELIQUIS DVT-PE TREAT</b>		<i>ethambutol</i> .....	6
<i>dicyclomine</i> .....	49	<b>30D START</b> .....	34	<i>ethosuximide</i> .....	23
<b>DIFICID</b> .....	5	<b>ELMIRON</b> .....	64	<i>ethynodiol diac-eth estradiol</i> .....	56
<i>diflunisal</i> .....	22	<b>ELURYNG</b> .....	56	<i>etodolac</i> .....	23
<i>difluprednate</i> .....	59	<b>EMGALITY PEN</b> .....	23	<i>etonogestrel-ethinyl estradiol</i> .....	56
<i>digoxin</i> .....	33, 34	<b>EMGALITY SYRINGE</b> .....	23	<i>etravirine</i> .....	6
<i>dihydroergotamine</i> .....	22	<b>EMSAM</b> .....	23	<b>EUTHYROX</b> .....	45
<b>DILANTIN</b> .....	22	<i>emtricitabine</i> .....	6	<b>EVENITY</b> .....	54
<i>diltiazem hcl</i> .....	34	<i>emtricitabine-tenofovir (tdf)</i> .....	6	<i>everolimus (antineoplastic)</i> .....	12
<b>DILT-XR</b> .....	34	<b>EMTRIVA</b> .....	6	<i>everolimus</i>	
<i>dimethyl fumarate</i> .....	22	<b>EMVERM</b> .....	6	<i>(immunosuppressive)</i> .....	12
<i>diphenoxylate-atropine</i> .....	49	<i>enalapril maleate</i> .....	34	<b>EVOTAZ</b> .....	6
<i>disulfiram</i> .....	43	<i>enalapril-hydrochlorothiazide</i> ...	34	<b>EVRYSDI</b> .....	23
<i>divalproex</i> .....	22	<b>ENBREL</b> .....	53, 54	<i>exemestane</i> .....	12
<i>dofetilide</i> .....	34	<b>ENBREL MINI</b> .....	53	<i>ezetimibe</i> .....	34
<i>donepezil</i> .....	22	<b>ENBREL SURECLICK</b> .....	54	<i>ezetimibe-simvastatin</i> .....	34
<b>DOPTELET (10 TAB PACK)</b> .....	34	<b>ENDOCET</b> .....	23	<b>FABHALTA</b> .....	43
<b>DOPTELET (15 TAB PACK)</b> .....	34	<b>ENERGIX-B (PF)</b> .....	51	<i>famciclovir</i> .....	6
<b>DOPTELET (30 TAB PACK)</b> .....	34	<b>ENERGIX-B PEDIATRIC</b>		<i>famotidine</i> .....	49
<i>dorzolamide</i> .....	59	<b>(PF)</b> .....	51	<b>FANAPT</b> .....	23
<i>dorzolamide-timolol</i> .....	59	<i>enoxaparin</i> .....	34	<b>FARXIGA</b> .....	45
<i>dorzolamide-timolol (pf)</i> .....	59	<b>ENPRESSE</b> .....	56	<b>FASENRA</b> .....	61
<b>DOTTI</b> .....	55	<b>ENSKYCE</b> .....	56	<b>FASENRA PEN</b> .....	61
<b>DOVATO</b> .....	5	<i>entacapone</i> .....	23	<i>febuxostat</i> .....	54
<i>doxazosin</i> .....	34	<i>entecavir</i> .....	6	<i>felbamate</i> .....	23
<i>doxepin</i> .....	22, 23	<b>ENTRESTO</b> .....	34	<i>felodipine</i> .....	34
<i>doxercalciferol</i> .....	44	<b>ENULOSE</b> .....	49	<i>fenofibrate</i> .....	34
<b>DOXY-100</b> .....	5	<b>ENVARUS XR</b> .....	12	<i>fenofibrate micronized</i> .....	34
<i>doxycycline hyclate</i> .....	5, 6	<b>EPCLUSA</b> .....	6	<i>fenofibrate nanocrystallized</i> .....	34

<i>fentanyl</i> .....	23, 24
<b>FETZIMA</b> .....	24
<b>FILSUVEZ</b> .....	40
<i>finasteride</i> .....	64
<i> fingolimod</i> .....	24
<b>FINTEPLA</b> .....	24
<b>FIRDAPSE</b> .....	24
<b>FIRMAGON KIT W</b>	
<b>DILUENT SYRINGE</b> .....	12
<i>flecainide</i> .....	34
<i>fluconazole</i> .....	6
<i>fluconazole in nacl (iso-osm)</i> .....	6
<i>flucytosine</i> .....	6
<i>fludrocortisone</i> .....	45
<i>flunisolide</i> .....	61
<i>fluocinolone</i> .....	40
<i>fluocinolone acetonide oil</i> .....	44
<i>fluocinolone and shower cap</i> .....	40
<i>fluocinonide</i> .....	40
<i>fluocinonide-emollient</i> .....	40
<i>fluoride (sodium)</i> .....	65
<i>fluorometholone</i> .....	59
<i>fluorouracil</i> .....	40
<i>fluoxetine</i> .....	24
<i>fluoxetine (pmdd)</i> .....	24
<i>fluphenazine decanoate</i> .....	24
<i>fluphenazine hcl</i> .....	24
<i>flurbiprofen</i> .....	24
<i>flurbiprofen sodium</i> .....	59
<i>fluticasone propionate</i> .....	40, 62
<i>fluticasone propion-salmeterol</i> ...	62
<i>fluvastatin</i> .....	34
<i>fluvoxamine</i> .....	24
<i>fondaparinux</i> .....	35
<i>fosamprenavir</i> .....	6
<i>fosinopril</i> .....	35
<i>fosinopril-hydrochlorothiazide</i> ...	35
<b>FOTIVDA</b> .....	12
<b>FRUZAQLA</b> .....	13
<b>FULPHILA</b> .....	51
<b>FUROSCIX</b> .....	35
<i>furosemide</i> .....	35
<b>FUZEON</b> .....	7
<b>FYCOMPA</b> .....	24
<i>gabapentin</i> .....	24
<i>galantamine</i> .....	24
<b>GALLIFREY</b> .....	56
<b>GAMMAGARD LIQUID</b> .....	51
<b>GAMMAGARD S-D (IGA &lt; 1</b>	
<b>MCG/ML)</b> .....	51
<b>GAMMAKED</b> .....	51
<b>GAMMAPLEX</b> .....	51

<b>GAMMAPLEX (WITH</b>	
<b>SORBITOL)</b> .....	51
<b>GAMUNEX-C</b> .....	51
<b>GARDASIL 9 (PF)</b> .....	51
<i>gatifloxacin</i> .....	59
<b>GATTEX 30-VIAL</b> .....	49
<b>GAUZE PAD</b> .....	53
<b>GAVILYTE-C</b> .....	49
<b>GAVILYTE-G</b> .....	49
<b>GAVILYTE-N</b> .....	49
<b>GAVRETO</b> .....	13
<i>gefitinib</i> .....	13
<i>gemfibrozil</i> .....	35
<b>GENERLAC</b> .....	49
<b>GENGRAF</b> .....	13
<i>gentamicin</i> .....	7, 40, 59
<i>gentamicin in nacl (iso-osm)</i> .....	7
<b>GENVOYA</b> .....	7
<b>GILOTRIF</b> .....	13
<i>glatiramer</i> .....	24
<b>GLATOPA</b> .....	25
<b>GLEOSTINE</b> .....	13
<i>glimepiride</i> .....	45
<i>glipizide</i> .....	45
<i>glipizide-metformin</i> .....	45
<b>GLUCAGON EMERGENCY</b>	
<b>KIT (HUMAN)</b> .....	45
<i>glycopyrrolate</i> .....	49
<b>GLYXAMBI</b> .....	45
<i>granisetron hcl</i> .....	49
<i>griseofulvin microsize</i> .....	7
<i>griseofulvin ultramicrosize</i> .....	7
<i>guanfacine</i> .....	25
<b>GVOKE</b> .....	45
<b>GVOKE HYPOPEN 2-PACK</b> .....	45
<b>GVOKE PFS 1-PACK</b>	
<b>SYRINGE</b> .....	45
<i>halobetasol propionate</i> .....	40
<b>HALOETTE</b> .....	56
<i>haloperidol</i> .....	25
<i>haloperidol decanoate</i> .....	25
<i>haloperidol lactate</i> .....	25
<b>HAVRIX (PF)</b> .....	51
<b>HEATHER</b> .....	56
<i>heparin (porcine)</i> .....	35
<b>HEPLISAV-B (PF)</b> .....	51
<b>HIBERIX (PF)</b> .....	51
<b>HUMALOG JUNIOR</b>	
<b>KWIKPEN U-100</b> .....	45
<b>HUMALOG KWIKPEN</b>	
<b>INSULIN</b> .....	45

<b>HUMALOG MIX 50-50</b>	
<b>KWIKPEN</b> .....	45
<b>HUMALOG MIX 75-25</b>	
<b>KWIKPEN</b> .....	45
<b>HUMALOG MIX 75-25(U-</b>	
<b>100)INSULN</b> .....	45
<b>HUMALOG TEMPO PEN(U-</b>	
<b>100)INSULN</b> .....	45
<b>HUMALOG U-100 INSULIN</b> ..	45
<b>HUMIRA</b> .....	54
<b>HUMIRA PEN</b> .....	54
<b>HUMIRA(CF)</b> .....	54
<b>HUMIRA(CF) PEN</b> .....	54
<b>HUMIRA(CF) PEN</b>	
<b>CROHNS-UC-HS</b> .....	54
<b>HUMIRA(CF) PEN PSOR-</b>	
<b>UV-ADOL HS</b> .....	54
<b>HUMULIN 70/30 U-100</b>	
<b>INSULIN</b> .....	45
<b>HUMULIN 70/30 U-100</b>	
<b>KWIKPEN</b> .....	45
<b>HUMULIN N NPH INSULIN</b>	
<b>KWIKPEN</b> .....	45
<b>HUMULIN N NPH U-100</b>	
<b>INSULIN</b> .....	45
<b>HUMULIN R REGULAR U-</b>	
<b>100 INSULN</b> .....	45
<b>HUMULIN R U-500 (CONC)</b>	
<b>INSULIN</b> .....	45
<b>HUMULIN R U-500 (CONC)</b>	
<b>KWIKPEN</b> .....	45
<i>hydralazine</i> .....	35
<i>hydrochlorothiazide</i> .....	35
<i>hydrocodone-acetaminophen</i> .....	25
<i>hydrocortisone</i> .....	40, 45, 49
<i>hydrocortisone-acetic acid</i> .....	44
<i>hydromorphone</i> .....	25
<i>hydroxychloroquine</i> .....	7
<i>hydroxyurea</i> .....	13
<i>hydroxyzine hcl</i> .....	62
<i>ibandronate</i> .....	54
<b>IBRANCE</b> .....	13
<b>IBSRELA</b> .....	49
<b>IBU</b> .....	25
<i>ibuprofen</i> .....	25
<i>icatibant</i> .....	62
<b>ICLEVIA</b> .....	56
<b>ICLUSIG</b> .....	13
<i>icosapent ethyl</i> .....	35
<b>IDHIFA</b> .....	13
<i>imatinib</i> .....	13
<b>IMBRUVICA</b> .....	13

<i>imipenem-cilastatin</i> .....	7	<b>JANUVIA</b> .....	45, 46	<i>lenalidomide</i> .....	14
<i>imipramine hcl</i> .....	25	<b>JARDIANCE</b> .....	46	<b>LENVIMA</b> .....	14
<i>imiquimod</i> .....	40	<b>JASMIEL (28)</b> .....	56	<b>LESSINA</b> .....	56
<b>IMKELDI</b> .....	13	<b>JAVYGTOR</b> .....	46	<i>letrozole</i> .....	14
<b>IMOVAX RABIES</b>		<b>JAYPIRCA</b> .....	13	<i>leucovorin calcium</i> .....	14
<b>VACCINE (PF)</b> .....	51	<b>JENTADUETO</b> .....	46	<b>LEUKINE</b> .....	52
<b>IMVEXXY MAINTENANCE</b>		<b>JENTADUETO XR</b> .....	46	<i>leuprolide</i> .....	14
<b>PACK</b> .....	56	<b>JINTELI</b> .....	56	<i>leuprolide (3 month)</i> .....	14
<b>IMVEXXY STARTER PACK</b> .....	56	<b>JOENJA</b> .....	43	<i>levalbuterol hcl</i> .....	62
<b>INCASSIA</b> .....	56	<b>JULEBER</b> .....	56	<i>levalbuterol tartrate</i> .....	62
<b>INCRELEX</b> .....	43	<b>JULUCA</b> .....	7	<i>levetiracetam</i> .....	26
<i>indapamide</i> .....	35	<b>JYNNEOS (PF)</b> .....	52	<i>levobunolol</i> .....	59
<i>indomethacin</i> .....	25	<b>KALYDECO</b> .....	62	<i>levocarnitine</i> .....	43
<b>INFANRIX (DTAP) (PF)</b> .....	51	<b>KARIVA (28)</b> .....	56	<i>levocarnitine (with sugar)</i> .....	43
<b>INLYTA</b> .....	13	<b>KELNOR 1/35 (28)</b> .....	56	<i>levocetirizine</i> .....	62
<b>INQOVI</b> .....	13	<b>KELNOR 1/50 (28)</b> .....	56	<i>levofloxacin</i> .....	7
<b>INREBIC</b> .....	13	<b>KERENDIA</b> .....	35	<i>levofloxacin in d5w</i> .....	7
<i>insulin lispro</i> .....	45	<b>KESIMPTA PEN</b> .....	26	<b>LEVONEST (28)</b> .....	56
<i>insulin lispro protamin-lispro</i> .....	45	<i>ketoconazole</i> .....	7, 40	<i>levonorgestrel-ethinyl estrad</i> .....	56, 57
<i>insulin syringe-needle u-100</i> .....	53	<i>ketorolac</i> .....	59	<i>levonorg-eth estrad triphasic</i> .....	57
<b>INTELENCE</b> .....	7	<b>KEVZARA</b> .....	54	<b>LEVORA-28</b> .....	57
<b>INTRALIPID</b> .....	65	<b>KINERET</b> .....	54	<i>levothyroxine</i> .....	46
<b>INTROVALE</b> .....	56	<b>KINRIX (PF)</b> .....	52	<b>LEVOXYL</b> .....	46
<b>INVEGA HAFYERA</b> .....	25	<b>KIONEX (WITH</b>		<b>LIBERVANT</b> .....	26
<b>INVEGA SUSTENNA</b> .....	25	<b>SORBITOL)</b> .....	43	<i>lidocaine</i> .....	41
<b>INVEGA TRINZA</b> .....	25, 26	<b>KISQALI</b> .....	14	<i>lidocaine hcl</i> .....	40
<b>IPOL</b> .....	51	<b>KISQALI FEMARA CO-</b>		<b>LIDOCAINE VISCOUS</b> .....	41
<i>ipratropium bromide</i> .....	44, 62	<b>PACK</b> .....	13	<i>lidocaine-prilocaine</i> .....	41
<i>ipratropium-albuterol</i> .....	62	<b>KLOR-CON</b> .....	65	<b>LILETTA</b> .....	57
<i>irbesartan</i> .....	35	<b>KLOR-CON M10</b> .....	65	<i>linezolid</i> .....	7
<i>irbesartan-hydrochlorothiazide</i> .....	35	<b>KLOR-CON M15</b> .....	65	<i>linezolid in dextrose 5%</i> .....	7
<b>ISENTRESS</b> .....	7	<b>KLOR-CON M20</b> .....	65	<b>LINZESS</b> .....	49
<b>ISENTRESS HD</b> .....	7	<b>KLOXXADO</b> .....	26	<i>liothyronine</i> .....	46
<b>ISIBLOOM</b> .....	56	<b>KOSELUGO</b> .....	14	<i>lisinopril</i> .....	35
<b>ISOLYTE S PH 7.4</b> .....	65	<b>KOURZEQ</b> .....	44	<i>lisinopril-hydrochlorothiazide</i> .....	35
<b>ISOLYTE-P IN 5 %</b>		<b>KRAZATI</b> .....	14	<b>LITFULO</b> .....	43
<b>DEXTROSE</b> .....	65	<b>KURVELO (28)</b> .....	56	<i>lithium carbonate</i> .....	26
<i>isoniazid</i> .....	7	<i>l norgest/e.estradiol-e.estrad</i> .....	56	<i>lithium citrate</i> .....	26
<i>isosorbide dinitrate</i> .....	35	<i>labetalol</i> .....	35	<b>LIVTENCITY</b> .....	7
<i>isosorbide mononitrate</i> .....	35	<i>lacosamide</i> .....	26	<i>lofexidine</i> .....	26
<i>isradipine</i> .....	35	<i>lactulose</i> .....	49	<b>LOKELMA</b> .....	43
<b>ITOVEBI</b> .....	13	<i>lamivudine</i> .....	7	<b>LONSURF</b> .....	14
<i>itraconazole</i> .....	7	<i>lamivudine-zidovudine</i> .....	7	<i>loperamide</i> .....	49
<i>ivabradine</i> .....	35	<i>lamotrigine</i> .....	26	<i>lopinavir-ritonavir</i> .....	7
<i>ivermectin</i> .....	7	<i>lansoprazole</i> .....	49	<i>lorazepam</i> .....	26
<b>IWILFIN</b> .....	13	<b>LANTUS SOLOSTAR U-100</b>		<b>LORAZEPAM INTENSOL</b> .....	26
<b>IXCHIQ (PF)</b> .....	52	<b>INSULIN</b> .....	46	<b>LORBRENA</b> .....	14
<b>IXIARO (PF)</b> .....	52	<b>LANTUS U-100 INSULIN</b> .....	46	<b>LORYNA (28)</b> .....	57
<b>JAKAFI</b> .....	13	<i>lapatinib</i> .....	14	<i>losartan</i> .....	35
<b>JANTOVEN</b> .....	35	<i>latanoprost</i> .....	59	<i>losartan-hydrochlorothiazide</i> .....	35
<b>JANUMET</b> .....	45	<b>LAZCLUZE</b> .....	14	<i>lovastatin</i> .....	35
<b>JANUMET XR</b> .....	45	<i>leflunomide</i> .....	54	<b>LOW-OGESTREL (28)</b> .....	57



<i>loxapine succinate</i> .....	26	<i>methsuximide</i> .....	26	<i>naproxen</i> .....	27
<i>lubiprostone</i> .....	49	<i>methylphenidate hcl</i> .....	26	<i>naproxen sodium</i> .....	27
<b>LUMAKRAS</b> .....	14	<i>methylprednisolone</i> .....	46	<i>naratriptan</i> .....	27
<b>LUMIGAN</b> .....	59	<i>metoclopramide hcl</i> .....	50	<i>nateglinide</i> .....	46
<b>LUPRON DEPOT</b> .....	14	<i>metolazone</i> .....	36	<b>NAYZILAM</b> .....	27
<b>LUPRON DEPOT (3</b>		<i>metoprolol succinate</i> .....	36	<i>nebivolol</i> .....	36
<b>MONTH)</b> .....	14	<i>metoprolol ta-hydrochlorothiaz</i> ..	36	<i>nefazodone</i> .....	27
<b>LUPRON DEPOT (4</b>		<i>metoprolol tartrate</i> .....	36	<b>NEMLUVIO</b> .....	15
<b>MONTH)</b> .....	14	<i>metronidazole</i> .....	8, 41, 57	<i>neomycin</i> .....	8
<b>LUPRON DEPOT (6</b>		<i>metronidazole in nacl (iso-os)</i> .....	8	<i>neomycin-bacitracin-poly-hc</i> .....	60
<b>MONTH)</b> .....	14	<i>metyrosine</i> .....	36	<i>neomycin-bacitracin-polymyxin</i> ..	60
<b>LUPRON DEPOT-PED</b> .....	15	<i>mexiletine</i> .....	36	<i>neomycin-polymyxin b-</i>	
<b>LUPRON DEPOT-PED (3</b>		<i>micafungin</i> .....	8	<i>dexameth</i> .....	60
<b>MONTH)</b> .....	14	<b>MICONAZOLE-3</b> .....	57	<i>neomycin-polymyxin-gramicidin</i> ..	60
<i>lurasidone</i> .....	26	<b>MICROGESTIN 1.5/30 (21)</b> ....	57	<i>neomycin-polymyxin-hc</i> .....	44, 60
<b>LUTERA (28)</b> .....	57	<b>MICROGESTIN 1/20 (21)</b> .....	57	<b>NEO-POLYCIN</b> .....	60
<b>LYLEQ</b> .....	57	<b>MICROGESTIN FE 1.5/30</b>		<b>NEO-POLYCIN HC</b> .....	60
<b>LYLLANA</b> .....	57	<b>(28)</b> .....	57	<b>NERLYNX</b> .....	15
<b>LYNPARZA</b> .....	15	<b>MICROGESTIN FE 1/20 (28)</b> ..	57	<b>NEULASTA</b> .....	52
<b>LYSODREN</b> .....	15	<i>midodrine</i> .....	43	<b>NEUPRO</b> .....	27
<b>LYTGOBI</b> .....	15	<i>mifepristone</i> .....	46	<i>nevirapine</i> .....	8
<b>LYZA</b> .....	57	<i>miglustat</i> .....	46	<b>NEXLETOL</b> .....	36
<i>magnesium sulfate</i> .....	65	<b>MILI</b> .....	57	<b>NEXLIZET</b> .....	36
<i>malathion</i> .....	41	<i>minocycline</i> .....	8	<b>NEXPLANON</b> .....	57
<i>maraviroc</i> .....	7	<i>minoxidil</i> .....	36	<i>niacin</i> .....	36
<b>MARLISSA (28)</b> .....	57	<i>mirtazapine</i> .....	27	<i>nicardipine</i> .....	36
<b>MARPLAN</b> .....	26	<i>misoprostol</i> .....	50	<b>NICOTROL NS</b> .....	43
<b>MATULANE</b> .....	15	<b>M-M-R II (PF)</b> .....	52	<i>nifedipine</i> .....	36
<b>MAVYRET</b> .....	7	<i>modafinil</i> .....	27	<i>nilutamide</i> .....	15
<i>meclizine</i> .....	49	<i>moexipril</i> .....	36	<i>nimodipine</i> .....	36
<i>medroxyprogesterone</i> .....	57	<i>molindone</i> .....	27	<b>NINLARO</b> .....	15
<i>mefloquine</i> .....	8	<i>mometasone</i> .....	41, 62	<i>nitazoxanide</i> .....	8
<i>megestrol</i> .....	15	<i>montelukast</i> .....	62	<i>nitisinone</i> .....	43
<b>MEKINIST</b> .....	15	<i>morphine</i> .....	27	<b>NITRO-BID</b> .....	36
<b>MEKTOVI</b> .....	15	<i>morphine concentrate</i> .....	27	<i>nitrofurantoin</i> .....	8
<i>meloxicam</i> .....	26	<b>MOUNJARO</b> .....	46	<i>nitrofurantoin macrocrystal</i> .....	8
<i>memantine</i> .....	26	<b>MOVANTIK</b> .....	50	<i>nitrofurantoin monohyd/m-cryst</i> ...8	
<b>MENACTRA (PF)</b> .....	52	<i>moxifloxacin</i> .....	8, 59	<i>nitroglycerin</i> .....	36, 50
<b>MENQUADFI (PF)</b> .....	52	<i>moxifloxacin-sod.chloride(iso)</i> .....	8	<b>NIVESTYM</b> .....	52
<b>MENVEO A-C-Y-W-135-DIP</b>		<b>MRESVIA (PF)</b> .....	52	<b>NORDITROPIN FLEXP</b> RO... 52	
<b>(PF)</b> .....	52	<b>MULPLETA</b> .....	36	<i>norethindrone (contraceptive)</i> ....	57
<i>mercaptopurine</i> .....	15	<b>MULTAQ</b> .....	36	<i>norethindrone acetate</i> .....	57
<i>meropenem</i> .....	8	<i>mupirocin</i> .....	41	<i>norethindrone ac-eth estradiol</i> ... 57	
<i>mesalamine</i> .....	50	<i>mycophenolate mofetil</i> .....	15	<i>norethindrone-e.estradiol-iron</i> ... 57	
<i>mesna</i> .....	15	<i>mycophenolate sodium</i> .....	15	<i>norgestimate-ethinyl estradiol</i> ... 57	
<i>metformin</i> .....	46	<b>MYRBETRIQ</b> .....	64	<b>NORTREL 0.5/35 (28)</b> .....	57
<i>methadone</i> .....	26	<i>nabumetone</i> .....	27	<b>NORTREL 1/35 (21)</b> .....	57
<i>methazolamide</i> .....	59	<i>nadolol</i> .....	36	<b>NORTREL 1/35 (28)</b> .....	57
<i>methenamine hippurate</i> .....	8	<i>nafcillin</i> .....	8	<b>NORTREL 7/7/7 (28)</b> .....	57
<i>methimazole</i> .....	46	<i>naloxone</i> .....	27	<i>nortriptyline</i> .....	27
<i>methotrexate sodium</i> .....	15	<i>naltrexone</i> .....	27	<b>NORVIR</b> .....	8
<i>methotrexate sodium (pf)</i> .....	15	<b>NAMZARIC</b> .....	27	<b>NUBEQA</b> .....	15

<b>NUCALA</b> .....	62	<i>paricalcitol</i> .....	46	<i>posaconazole</i> .....	9
<b>NUDEXTA</b> .....	27	<i>paroxetine hcl</i> .....	28	<i>potassium chlorid-d5-</i>	
<b>NUPLAZID</b> .....	27	<b>PAXLOVID</b> .....	8	<i>0.45%nacl</i> .....	65
<b>NURTEC ODT</b> .....	27	<i>pazopanib</i> .....	16	<i>potassium chloride</i> .....	65
<b>NYAMYC</b> .....	41	<b>PEDIARIX (PF)</b> .....	52	<i>potassium chloride in 0.9%nacl</i> .....	65
<i>nystatin</i> .....	8, 41	<b>PEDVAX HIB (PF)</b> .....	52	<i>potassium chloride in 5 % dex</i> .....	65
<i>nystatin-triamcinolone</i> .....	41	<i>peg 3350-electrolytes</i> .....	50	<i>potassium chloride in lr-d5</i> .....	65
<b>NYSTOP</b> .....	41	<i>peg3350-sod sul-nacl-kcl-asb-c</i> ..	50	<i>potassium chloride in water</i> .....	65
<b>OALIVA</b> .....	50	<b>PEGASYS</b> .....	52	<i>potassium chloride-0.45 % nacl</i> .....	65
<b>OCTAGAM</b> .....	52	<i>peg-electrolyte soln</i> .....	50	<i>potassium chloride-d5-</i>	
<i>octreotide acetate</i> .....	15	<b>PEMAZYRE</b> .....	16	<i>0.2%nacl</i> .....	65
<b>ODEFSEY</b> .....	8	<i>pen needle, diabetic</i> .....	53	<i>potassium chloride-d5-</i>	
<b>ODOMZO</b> .....	16	<b>PENBRAYA (PF)</b> .....	52	<i>0.9%nacl</i> .....	66
<b>OFEV</b> .....	63	<i>penciclovir</i> .....	41	<i>potassium citrate</i> .....	64
<i>ofloxacin</i> .....	8, 44, 60	<i>penicillamine</i> .....	54	<i>pramipexole</i> .....	28
<b>OGSIVEO</b> .....	16	<i>penicillin g pot in dextrose</i> .....	8	<i>prasugrel hcl</i> .....	37
<b>OJEMDA</b> .....	16	<i>penicillin g potassium</i> .....	9	<i>pravastatin</i> .....	37
<b>OJJAARA</b> .....	16	<i>penicillin v potassium</i> .....	9	<i>praziquantel</i> .....	9
<i>olanzapine</i> .....	27	<b>PENTACEL (PF)</b> .....	52	<i>prazosin</i> .....	37
<i>olmesartan</i> .....	36	<i>pentamidine</i> .....	9	<i>prednisolone</i> .....	47
<i>olmesartan-amlodipin-hcthiazid</i> .....	36	<i>pentoxifylline</i> .....	37	<i>prednisolone acetate</i> .....	60
<i>olmesartan-hydrochlorothiazide</i> .....	36	<i>perindopril erbumine</i> .....	37	<i>prednisolone sodium phosphate</i>	
<i>olopatadine</i> .....	44	<b>PERIOGARD</b> .....	44	.....	47, 60
<b>OLUMIANT</b> .....	54	<i>permethrin</i> .....	41	<i>prednisone</i> .....	47
<i>omega-3 acid ethyl esters</i> .....	36	<i>perphenazine</i> .....	28	<i>pregabalin</i> .....	28
<i>omeprazole</i> .....	50	<b>PERSERIS</b> .....	28	<b>PREMARIN</b> .....	58
<i>ondansetron</i> .....	50	<b>PHEBURANE</b> .....	43	<b>PREMPRO</b> .....	58
<i>ondansetron hcl</i> .....	50	<i>phenelzine</i> .....	28	<b>PRENATAL VITAMIN</b>	
<b>ONUREG</b> .....	16	<i>phenobarbital</i> .....	28	<b>PLUS LOW IRON</b> .....	66
<b>OPSUMIT</b> .....	63	<b>PHENYTEK</b> .....	28	<b>PREVALITE</b> .....	37
<b>OPSYNVI</b> .....	63	<i>phenytoin</i> .....	28	<b>PREVYMIS</b> .....	9
<b>ORENCIA</b> .....	54	<i>phenytoin sodium extended</i> .....	28	<b>PREZCOBIX</b> .....	9
<b>ORENCIA CLICKJECT</b> .....	54	<b>PIFELTRO</b> .....	9	<b>PREZISTA</b> .....	9
<b>ORGOVYX</b> .....	16	<i>pilocarpine hcl</i> .....	43, 60	<b>PRIFTIN</b> .....	9
<b>ORKAMBI</b> .....	63	<i>pimecrolimus</i> .....	41	<i>primaquine</i> .....	9
<b>ORSERDU</b> .....	16	<i>pimozide</i> .....	28	<i>primidone</i> .....	28
<i>oseltamivir</i> .....	8	<b>PIMTREA (28)</b> .....	58	<b>PRIORIX (PF)</b> .....	52
<b>OTEZLA</b> .....	54	<i>pindolol</i> .....	37	<b>PRIVIGEN</b> .....	52
<b>OTEZLA STARTER</b> .....	54	<i>pioglitazone</i> .....	47	<i>probenecid</i> .....	54
<i>oxacillin</i> .....	8	<i>pioglitazone-metformin</i> .....	47	<i>probenecid-colchicine</i> .....	54
<i>oxaprozin</i> .....	28	<i>piperacillin-tazobactam</i> .....	9	<i>prochlorperazine</i> .....	50
<i>oxcarbazepine</i> .....	28	<b>PIQRAY</b> .....	16	<i>prochlorperazine maleate</i> .....	50
<i>oxybutynin chloride</i> .....	64	<i>pirfenidone</i> .....	63	<b>PROCRIT</b> .....	52
<i>oxycodone</i> .....	28	<i>piroxicam</i> .....	28	<b>PROCTO-MED HC</b> .....	50
<i>oxycodone-acetaminophen</i> .....	28	<i>pitavastatin calcium</i> .....	37	<b>PROCTOSOL HC</b> .....	50
<b>OZEMPIC</b> .....	46	<b>PLEGRIDY</b> .....	52	<b>PROCTOZONE-HC</b> .....	50
<b>PACERONE</b> .....	37	<b>PLENAMINE</b> .....	65	<i>progesterone micronized</i> .....	58
<i>paliperidone</i> .....	28	<i>podofilox</i> .....	41	<b>PROGRAF</b> .....	16
<b>PALYNZIQ</b> .....	46	<b>POLYCIN</b> .....	60	<b>PROLASTIN-C</b> .....	43
<b>PANRETIN</b> .....	41	<i>polymyxin b sulf-trimethoprim</i> ....	60	<b>PROLENSA</b> .....	60
<i>pantoprazole</i> .....	50	<b>POMALYST</b> .....	16	<b>PROLIA</b> .....	54
<b>PANZYGA</b> .....	52	<b>PORTIA 28</b> .....	58	<b>PROMACTA</b> .....	37

<i>promethazine</i> .....	63	<i>rimantadine</i> .....	9	<i>sodium chloride 5 % hypertonic</i> ..	66
<i>propafenone</i> .....	37	<b>RINVOQ</b> .....	55	<i>sodium oxybate</i> .....	30
<i>propranolol</i> .....	37	<b>RINVOQ LQ</b> .....	54	<i>sodium phenylbutyrate</i> .....	43
<i>propylthiouracil</i> .....	47	<i>risedronate</i> .....	43, 55	<i>sodium polystyrene sulfonate</i> .....	43
<b>PROQUAD (PF)</b> .....	52	<i>risperidone</i> .....	29	<i>sodium,potassium,mag sulfates</i> ..	50
<i>protriptyline</i> .....	28	<i>risperidone microspheres</i> .....	29	<i>sofosbuvir-velpatasvir</i> .....	9
<b>PULMOZYME</b> .....	63	<i>ritonavir</i> .....	9	<b>SOHONOS</b> .....	43
<b>PURIXAN</b> .....	16	<i>rivastigmine</i> .....	29	<b>SOLQUA 100/33</b> .....	47
<i>pyrazinamide</i> .....	9	<i>rivastigmine tartrate</i> .....	29	<b>SOLTAMOX</b> .....	17
<i>pyridostigmine bromide</i> .....	28, 29	<b>RIVFLOZA</b> .....	64	<b>SOMAVERT</b> .....	47
<i>pyrimethamine</i> .....	9	<i>rizatriptan</i> .....	29	<i>sorafenib</i> .....	17
<b>QINLOCK</b> .....	16	<i>roflumilast</i> .....	63	<i>sotalol</i> .....	37
<b>QUADRACEL (PF)</b> .....	52	<i>ropinirole</i> .....	29	<b>SOTALOL AF</b> .....	37
<i>quetiapine</i> .....	29	<i>rosuvastatin</i> .....	37	<b>SPIRIVA RESPIMAT</b> .....	63
<i>quinapril</i> .....	37	<b>ROTARIX</b> .....	53	<b>SPIRIVA WITH</b>	
<i>quinapril-hydrochlorothiazide</i> ....	37	<b>ROTATEQ VACCINE</b> .....	53	<b>HANDIHALER</b> .....	63
<i>quinidine sulfate</i> .....	37	<b>ROWEEPRA</b> .....	29	<i>spironolactone</i> .....	37
<i>quinine sulfate</i> .....	9	<b>ROZLYTREK</b> .....	16, 17	<i>spironolacton-hydrochlorothiaz</i> ..	37
<b>QULIPTA</b> .....	29	<b>RUBRACA</b> .....	17	<b>SPRINTEC (28)</b> .....	58
<b>QVAR REDHALER</b> .....	63	<i>rufinamide</i> .....	29, 30	<b>SPRITAM</b> .....	30
<b>RABAVERT (PF)</b> .....	52	<b>RUKOBIA</b> .....	9	<b>SPS (WITH SORBITOL)</b> .....	43
<i>rabeprazole</i> .....	50	<b>RYBELSUS</b> .....	47	<b>SRONYX</b> .....	58
<b>RADICAVA ORS STARTER</b>		<b>RYDAPT</b> .....	17	<b>SSD</b> .....	41
<b>KIT SUSP</b> .....	29	<b>RYTARY</b> .....	30	<b>STELARA</b> .....	41
<i>raloxifene</i> .....	54	<b>SAJAZIR</b> .....	63	<b>STIOLTO RESPIMAT</b> .....	63
<i>ramelteon</i> .....	29	<i>sapropterin</i> .....	47	<b>STIVARGA</b> .....	17
<i>ramipril</i> .....	37	<b>SAVELLA</b> .....	55	<i>streptomycin</i> .....	9
<i>ranolazine</i> .....	37	<b>SCSEMBLIX</b> .....	17	<b>STRIBILD</b> .....	9
<i>rasagiline</i> .....	29	<i>scopolamine base</i> .....	50	<b>STRIVERDI RESPIMAT</b> .....	63
<b>RAVICTI</b> .....	43	<b>SECUADO</b> .....	30	<b>SUBVENITE</b> .....	30
<b>RECLIPSEN (28)</b> .....	58	<i>selegiline hcl</i> .....	30	<i>sucralfate</i> .....	50
<b>RECOMBIVAX HB (PF)</b> .....	52	<i>selenium sulfide</i> .....	41	<i>sulfacetamide sodium</i> .....	60
<b>RECORLEV</b> .....	47	<b>SELZENTRY</b> .....	9	<i>sulfacetamide sodium (acne)</i> .....	41
<b>REGRANEX</b> .....	41	<b>SEREVENT DISKUS</b> .....	63	<i>sulfacetamide-prednisolone</i> .....	60
<b>RELENZA DISKHALER</b> .....	9	<i>sertraline</i> .....	30	<i>sulfadiazine</i> .....	9
<i>repaglinide</i> .....	47	<b>SETLAKIN</b> .....	58	<i>sulfamethoxazole-trimethoprim</i> ..	10
<b>REPATHA PUSHTRONEX</b> ....	37	<b>SHINGRIX (PF)</b> .....	53	<b>SULFAMYLON</b> .....	41
<b>REPATHA SURECLICK</b> .....	37	<b>SIGNIFOR</b> .....	17	<i>sulfasalazine</i> .....	50
<b>REPATHA SYRINGE</b> .....	37	<i>sildenafil (pulm.hypertension)</i> ....	63	<i>sulindac</i> .....	30
<b>RESTASIS</b> .....	60	<i>silodosin</i> .....	64	<i>sumatriptan</i> .....	30
<b>RESTASIS MULTIDOSE</b> .....	60	<i>silver sulfadiazine</i> .....	41	<i>sumatriptan succinate</i> .....	30
<b>RETACRIT</b> .....	53	<b>SIMBRINZA</b> .....	60	<i>sunitinib malate</i> .....	17
<b>RETEVMO</b> .....	16	<b>SIMPONI</b> .....	55	<b>SUNLENCA</b> .....	10
<b>REVUFORJ</b> .....	16	<i>simvastatin</i> .....	37	<b>SYEDA</b> .....	58
<b>REXULTI</b> .....	29	<i>sirolimus</i> .....	17	<b>SYMDEKO</b> .....	63
<b>REYATAZ</b> .....	9	<b>SIRTURO</b> .....	9	<b>SYMLINPEN 120</b> .....	47
<b>REZDIFFRA</b> .....	43	<b>SKYCLARYS</b> .....	30	<b>SYMLINPEN 60</b> .....	47
<b>REZLIDHIA</b> .....	16	<b>SKYRIZI</b> .....	41, 50	<b>SYMPAZAN</b> .....	30
<i>ribavirin</i> .....	9	<i>sodium chloride</i> .....	43	<b>SYMTUZA</b> .....	10
<i>rifabutin</i> .....	9	<i>sodium chloride 0.45 %</i> .....	66	<b>SYNAREL</b> .....	47
<i>rifampin</i> .....	9	<i>sodium chloride 0.9 %</i> .....	43	<b>SYNJARDY</b> .....	47
<i>riluzole</i> .....	43	<i>sodium chloride 3 % hypertonic</i> ..	66	<b>SYNJARDY XR</b> .....	47



<b>SYNTHROID</b> .....	47	<i>tobramycin</i> .....	10, 60	<b>TURALIO</b> .....	18
<b>TABRECTA</b> .....	17	<i>tobramycin in 0.225 % nacl</i> .....	10	<b>TURQOZ (28)</b> .....	58
<i>tacrolimus</i> .....	17, 41	<i>tobramycin sulfate</i> .....	10	<b>TWINRIX (PF)</b> .....	53
<i>tadalafil</i> .....	64	<i>tobramycin-dexamethasone</i> .....	60	<b>TYBOST</b> .....	10
<i>tadalafil (pulm. hypertension)</i> ....	63	<i>tolterodine</i> .....	64	<b>TYMLOS</b> .....	55
<b>TAFINLAR</b> .....	17	<i>tolvaptan</i> .....	47	<b>TYPHIM VI</b> .....	53
<b>TAGRISSO</b> .....	17	<i>topiramate</i> .....	30, 31	<b>UBRELVY</b> .....	31
<b>TALTZ AUTOINJECTOR</b> .....	42	<i>toremifene</i> .....	17	<b>UNITHROID</b> .....	48
<b>TALTZ SYRINGE</b> .....	42	<i>torsemide</i> .....	38	<b>UPTRAVI</b> .....	38
<b>TALZENNA</b> .....	17	<b>TOUJEO MAX U-300</b>		<i>ursodiol</i> .....	50
<i>tamoxifen</i> .....	17	<b>SOLOSTAR</b> .....	48	<i>valacyclovir</i> .....	10
<i>tamsulosin</i> .....	64	<b>TOUJEO SOLOSTAR U-300</b>		<b>VALCHLOR</b> .....	42
<b>TASCENSO ODT</b> .....	30	<b>INSULIN</b> .....	48	<i>valganciclovir</i> .....	10
<b>TASIGNA</b> .....	17	<b>TRADJENTA</b> .....	48	<i>valproic acid</i> .....	31
<i>tasimelteon</i> .....	30	<i>tramadol</i> .....	31	<i>valproic acid (as sodium salt)</i> ....	31
<i>tazarotene</i> .....	42	<i>tramadol-acetaminophen</i> .....	31	<i>valsartan</i> .....	38
<b>TAZVERIK</b> .....	17	<i>trandolapril</i> .....	38	<i>valsartan-hydrochlorothiazide</i> ...	38
<b>TEFLARO</b> .....	10	<i>tranexamic acid</i> .....	58	<b>VALTOCO</b> .....	31
<i>telmisartan</i> .....	37	<i>tranylcypromine</i> .....	31	<i>vancomycin</i> .....	10
<i>telmisartan-amlodipine</i> .....	38	<b>TRAVASOL 10 %</b> .....	66	<b>VANFLYTA</b> .....	18
<i>telmisartan-hydrochlorothiazid</i> ..	38	<i>travoprost</i> .....	60	<b>VAQTA (PF)</b> .....	53
<b>TENIVAC (PF)</b> .....	53	<i>trazodone</i> .....	31	<i>varenicline tartrate</i> .....	43
<i>tenofovir disoproxil fumarate</i> .....	10	<b>TRECATOR</b> .....	10	<b>VARIVAX (PF)</b> .....	53
<b>TEPMETKO</b> .....	17	<b>TRELEGY ELLIPTA</b> .....	63	<b>VAXCHORA VACCINE</b> .....	53
<i>terazosin</i> .....	38	<b>TRELSTAR</b> .....	17	<b>VELIVET TRIPHASIC</b>	
<i>terbinafine hcl</i> .....	10	<i>tretinoin</i> .....	42	<b>REGIMEN (28)</b> .....	58
<i>terbutaline</i> .....	63	<i>tretinoin (antineoplastic)</i> .....	18	<b>VEMLIDY</b> .....	10
<i>terconazole</i> .....	58	<i>triamcinolone acetonide</i> .....	42, 44	<b>VENCLEXTA</b> .....	18
<i>teriflunomide</i> .....	30	<i>triamterene-hydrochlorothiazid</i> ..	38	<b>VENCLEXTA STARTING</b>	
<i>teriparatide</i> .....	55	<b>TRIDERM</b> .....	42	<b>PACK</b> .....	18
<i>testosterone</i> .....	47	<i>trientine</i> .....	43	<i>venlafaxine</i> .....	31
<i>testosterone cypionate</i> .....	47	<b>TRI-ESTARYLLA</b> .....	58	<b>VENTOLIN HFA</b> .....	63
<i>testosterone enanthate</i> .....	47	<i>trifluoperazine</i> .....	31	<i>verapamil</i> .....	38
<i>tetrabenazine</i> .....	30	<i>trifluridine</i> .....	60	<b>VERQUVO</b> .....	38
<i>tetracycline</i> .....	10	<b>TRIJARDY XR</b> .....	48	<b>VERSACLOZ</b> .....	31
<b>THALOMID</b> .....	17	<b>TRIKAFTA</b> .....	63	<b>VERZENIO</b> .....	18
<b>THEO-24</b> .....	63	<b>TRI-LEGEST FE</b> .....	58	<b>VESTURA (28)</b> .....	58
<i>theophylline</i> .....	63	<b>TRI-LO-ESTARYLLA</b> .....	58	<b>VIBERZI</b> .....	51
<i>thioridazine</i> .....	30	<b>TRI-LO-SPRINTEC</b> .....	58	<b>VIENVA</b> .....	58
<i>thiothixene</i> .....	30	<i>trimethoprim</i> .....	10	<i>vigabatrin</i> .....	31
<b>TIADYL T ER</b> .....	38	<i>trimipramine</i> .....	31	<b>VIGADRONE</b> .....	31
<i>tiagabine</i> .....	30	<b>TRINTELLIX</b> .....	31	<b>VIGPODER</b> .....	31
<b>TIBSOVO</b> .....	17	<b>TRI-SPRINTEC (28)</b> .....	58	<b>VIJOICE</b> .....	18
<b>TICOVAC</b> .....	53	<b>TRIUMEQ</b> .....	10	<i>vilazodone</i> .....	31
<i>tigecycline</i> .....	10	<b>TRIUMEQ PD</b> .....	10	<b>VIRACEPT</b> .....	10
<b>TILIA FE</b> .....	58	<b>TRIVORA (28)</b> .....	58	<b>VIREAD</b> .....	10
<i>timolol maleate</i> .....	38, 60	<b>TROPHAMINE 10 %</b> .....	66	<b>VITRAKVI</b> .....	18
<i>tinidazole</i> .....	10	<i>trospium</i> .....	64	<b>VIVITROL</b> .....	31
<b>TIVICAY</b> .....	10	<b>TRULICITY</b> .....	48	<b>VIVJOA</b> .....	10
<b>TIVICAY PD</b> .....	10	<b>TRUMENBA</b> .....	53	<b>VIZIMPRO</b> .....	18
<i>tizanidine</i> .....	30	<b>TRUQAP</b> .....	18	<b>VONJO</b> .....	18
<b>TOBI PODHALER</b> .....	10	<b>TUKYSA</b> .....	18	<b>VORANIGO</b> .....	18

<i>voriconazole</i> .....	10	<b>ZEGALOGUE SYRINGE</b> .....	48
<b>VOSEVI</b> .....	11	<b>ZEJULA</b> .....	19
<b>VOWST</b> .....	51	<b>ZELBORAF</b> .....	19
<b>VRAYLAR</b> .....	31	<b>ZEPOSIA</b> .....	32
<b>VUMERITY</b> .....	31	<b>ZEPOSIA STARTER KIT</b>	
<b>VYNDAQEL</b> .....	38	<b>(28-DAY)</b> .....	32
<i>warfarin</i> .....	38	<b>ZEPOSIA STARTER PACK</b>	
<b>WELIREG</b> .....	18	<b>(7-DAY)</b> .....	32
<b>WIXELA INHUB</b> .....	63	<i>zidovudine</i> .....	11
<b>XALKORI</b> .....	18	<b>ZIEXTENZO</b> .....	53
<b>XARELTO</b> .....	38	<b>ZILBRYSQ</b> .....	32
<b>XARELTO DVT-PE TREAT</b>		<i>ziprasidone hcl</i> .....	32
<b>30D START</b> .....	38	<i>ziprasidone mesylate</i> .....	32
<b>XATMEP</b> .....	18	<b>ZIRGAN</b> .....	60
<b>XCOPRI</b> .....	31	<b>ZOLINZA</b> .....	19
<b>XCOPRI MAINTENANCE</b>		<i>zolpidem</i> .....	32
<b>PACK</b> .....	31	<b>ZONISADE</b> .....	32
<b>XCOPRI TITRATION PACK</b>		<i>zonisamide</i> .....	32
.....	31, 32	<b>ZOVIA 1-35 (28)</b> .....	58
<b>XDEMVY</b> .....	60	<b>ZTALMY</b> .....	32
<b>XELJANZ</b> .....	55	<b>ZUBSOLV</b> .....	32
<b>XELJANZ XR</b> .....	55	<b>ZURZUVAE</b> .....	32
<b>XERMELO</b> .....	18	<b>ZYDELIG</b> .....	19
<b>XGEVA</b> .....	18	<b>ZYKADIA</b> .....	19
<b>XIFAXAN</b> .....	11		
<b>XIGDUO XR</b> .....	48		
<b>XIIDRA</b> .....	60		
<b>XOFLUZA</b> .....	11		
<b>XOLAIR</b> .....	64		
<b>XOLREMDI</b> .....	53		
<b>XOSPATA</b> .....	18		
<b>XPOVIO</b> .....	18, 19		
<b>XTANDI</b> .....	19		
<b>XULTOPHY 100/3.6</b> .....	48		
<b>XYREM</b> .....	32		
<b>YARGESA</b> .....	48		
<b>YF-VAX (PF)</b> .....	53		
<b>YONSA</b> .....	19		
<b>YORVIPATH</b> .....	48		
<b>YUFLYMA(CF)</b> .....	55		
<b>YUFLYMA(CF) AI</b>			
<b>CROHN'S-UC-HS</b> .....	55		
<b>YUFLYMA(CF)</b>			
<b>AUTOINJECTOR</b> .....	55		
<b>YUVAFEM</b> .....	58		
<b>ZAFEMY</b> .....	58		
<i>zafirlukast</i> .....	64		
<i>zaleplon</i> .....	32		
<b>ZARXIO</b> .....	53		
<b>ZAVZPRET</b> .....	32		
<b>ZEGALOGUE</b>			
<b>AUTOINJECTOR</b> .....	48		

# Acitretin

---

## Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Actemra

## Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide). For giant cell arteritis, patients must have therapeutic failure or intolerance to one systemic corticosteroid (e.g., prednisone). For polyarticular juvenile idiopathic arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For systemic sclerosis-associated interstitial lung disease (SSc-ILD), patients must have therapeutic failure or intolerance to one immunosuppressant (e.g., mycophenolate mofetil, corticosteroids, cyclophosphamide). Documentation of systemic juvenile idiopathic arthritis.
Age Restrictions	Deny if less than 18 years of age for systemic sclerosis-associated interstitial lung disease (SSc-ILD), Rheumatoid Arthritis, and Giant Cell Arteritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Actimmune

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Adbry

## Products Affected

- ADBRY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: Dupixent and Rinvoq. For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ADHD Drugs

---

## Products Affected

- *guanfacine oral tablet extended release 24 hr*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Afinitor

## Products Affected

- *everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For everolimus only, documentation of advanced, hormone receptor-positive, HER2-negative breast cancer -AND- all of the following (1-3): 1) member is a postmenopausal woman 2) using in combination with exemestane 3) therapeutic failure or intolerance to prior treatment with letrozole or anastrozole. For everolimus only, documentation of non-functional neuroendocrine tumors -AND- all of the following (1-2): 1) disease is classified as progressive, well-differentiated, non-functional 2) disease is of gastrointestinal or lung origin that are unresectable, locally advanced or metastatic. For everolimus only, documentation of advanced renal cell carcinoma -AND- therapeutic failure or intolerance to prior treatment with sunitinib or sorafenib. For everolimus only, documentation of renal angiomyolipoma and tuberous sclerosis complex (TSC), member does not require immediate surgery. For everolimus and everolimus tablets for oral suspension, documentation of TSC with Subependymal Giant Cell Astrocytoma -AND- member is not a candidate for curative surgical resection. For everolimus only, documentation of progressive neuroendocrine tumors of pancreatic origin. For everolimus tablets for oral suspension only, documentation of use for adjunctive treatment of TSC-associated partial-onset seizures.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Aimovig

## Products Affected

- **AIMOVIG AUTOINJECTOR**  
**SUBCUTANEOUS AUTO-INJECTOR**  
**140 MG/ML, 70 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ajovy

## Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Akeega

---

## Products Affected

- AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) BRCA mutations, 2) Comcomitant therapy with prednisone, 3) Concomitant therapy with a gonadotropin-releasing hormone analog or a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Alecensa

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive -AND- One of the following (1-2): 1) metastatic disease, 2) will be used as adjuvant treatment following tumor resection of node positive or greater than or equal to 4 cm tumor(s).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Alpha1-Proteinase Inhibitors

## Products Affected

- **PROLASTIN-C INTRAVENOUS SOLUTION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of panacinar emphysema AND documentation of a decline in forced expiratory volume in 1 second (fev1) despite medical therapy (bronchodilators, corticosteroids) AND documentation of phenotype (pi*zz, pi*znull or pi*nullnull) associated with causing serum alpha 1-antitrypsin of less than 80 mg/dl AND documentation of an alpha 1-antitrypsin serum level below the value of 35% of normal (less than 80 mg/dl).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when furnished incident to a physician service and is not self-administered.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Alunbrig

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ampyra

## Products Affected

- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure disorder, Cr Cl less than 50ml/min
Required Medical Information	Documentation of diagnosis -AND- documentation that the patient is ambulatory and has walking impairment as evidenced by one of the following. 1. Functional status score (EDSS score). 2. Timed 25-foot Walk Test (T25W).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement in walking impairment from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Arikayce

## Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Mycobacterium avium complex lung disease -AND- Attestation of not achieving negative sputum cultures despite at least 6 months with a multidrug background regimen containing 2 of the following: 1) macrolide 2) rifamycin or 3) ethambutal -AND- Arikayce will be used in conjunction with a background multidrug regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive sputum cultures -OR- negative sputum cultures for insufficient period of time (e.g. less than 12 months).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ATTR-CM drugs

## Products Affected

- VYNDAQEL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitantly with transthyretin-lowering agents
<b>Required Medical Information</b>	Documentation of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) with amyloid deposits on cardiac biopsy or scintigraphy with heart contralateral greater than 1.5 (Grade III) -AND- Cardiac involvement supported by cardiac magnetic resonance, echocardiography or serum cardiac biomarker (e.g. B-type natriuretic peptide, cardiac troponin) - AND- Primary (light chain) amyloidosis has been ruled out by immunohistochemistry, mass spectrometry or scintigraphy.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For reauthorization, attestation of improvement or delayed disease progression from baseline demonstrated by 6-minute walk test, cardiac function (e.g. LVEF, NYHA class), Kansas City Cardiomyopathy Questionnaire-Overall Summary, number of cardiovascular-related hospitalizations or serum cardiac biomarkers (e.g. B-type natriuretic peptide, cardiac troponin)
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Atypical Antipsychotics

---

## Products Affected

- *aripiprazole oral solution*
- *aripiprazole oral tablet, disintegrating*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Aubagio

## Products Affected

- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as fingolimod, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Augtyro

---

## Products Affected

- **AUGTYRO ORAL CAPSULE 160 MG, 40 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive. Documentation of solid tumors - AND- disease harbors a NTRK gene fusion -AND- one of the following (1-2): 1) disease is locally advanced or metastatic, or 2) surgical resection is likely to result in severe morbidity -AND- one of the following (3-4): 3) disease has progressed following treatment, or 4) the member has no satisfactory alternative therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Auvelity

## Products Affected

- AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ayvakit

## Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable or metastatic gastrointestinal stromal tumor -AND- tumors harbor a PDGFRA exon 18 mutation. Documentation of aggressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm, mast cell leukemia, or indolent systemic mastocytosis -AND- platelet count greater than or equal to $50 \times 10^9/L$ .
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Bafiertam

## Products Affected

- **BAFIERTAM**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Balversa

## Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) FGFR3 mutation status as detected by an FDA approved test 2) disease progression on or after at least on prior line of systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Banzel

## Products Affected

- *rufinamide*

PA Criteria	Criteria Details
Exclusion Criteria	Patients with familial short QT syndrome
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Benlysta

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of active systemic lupus erythematosus (SLE) -AND- documentation of positive anti-nuclear antibody (ANA) titer (greater than or equal to 1:80) or anti-double-stranded DNA antibody (anti-dsDNA) greater than or equal to 30IU/mL -AND- member will continue to receive concomitant standard of care treatment with use of at least one of the following (alone or in combination): 1.) corticosteroids (e.g. prednisone) 2.) antimalarials (e.g. hydroxychloroquine) 3.) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate) -OR- Documentation of active lupus nephritis -AND- documentation of positive ANA titer (greater than or equal to 1:80) or anti-dsDNA greater than or equal to 30 IU/mL -AND- member will continue to receive concomitant standard of care treatment which includes corticosteroids (e.g. prednisone) with at least one of the following: 1.) mycophenolate for induction followed by mycophenolate for maintenance 2.) cyclophosphamide for induction followed by azathioprine for maintenance
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For SLE reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with corticosteroids, antimalarials, or immunosuppressives. For active LN reauthorization, attestation of disease stability or improvement - AND- attestation the member will continue to receive standard of care therapy with mycophenolate or azathioprine. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Besremi

## Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Bosulif

## Products Affected

- **BOSULIF ORAL CAPSULE 100 MG, 50 MG**
- **BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy. For pediatric patients 1 year of age and older, one of the following (3, 4): 3) newly diagnosed PH-positive CML in the chronic phase 4) diagnosis of Ph-positive CML in the chronic phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Bosutinib capsules and 18 years of age or older, inability to swallow tablets is required. For Bosutinib 100mg capsules and pediatric 1 year of age or older, inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Braftovi

## Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	Use in wild-type BRAF melanoma or wild-type BRAF CRC
Required Medical Information	Documentation of diagnosis. For metastatic colorectal cancer (mCRC) and using in combination with cetuximab, all of the following (1-3): 1) BRAF V600E mutation status 2) using in combination with cetuximab 3) member has received prior therapy for CRC. For mCRC and using in combination with cetuximab and modified FOLFOX6, all of the following (4-5): 4) BRAF V600E mutation status, as detected by an FDA-approved test 5) using in combination with cetuximab and modified FOLFOX6. For unresectable or metastatic melanoma, all of the following (6-7): 6) BRAF V600E or V600K mutation status 7) using in combination with binimetinib. For metastatic non-small cell lung cancer, all of the following (8-9): 8) BRAF V600E mutation status 9) using in combination with binimetinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Brukinsa

## Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma (MCL), previous treatment with at least 1 prior therapy. For marginal zone lymphoma (MZL), previous treatment with at least 1 anti-CD20-based regimen. For follicular lymphoma (FL), using in combination with obinutuzumab and previous treatment with at least 2 prior lines of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Buphenyl

## Products Affected

- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cablivi

## Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of acquired thrombocytopenic purpura (aTTP) -AND- Used in conjunction with plasma exchange and immunosuppressive therapy (i.e. systemic corticosteroids or rituximab)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	75 days initial authorization, 28 days reauthorization
Other Criteria	For reauthorization, attestation of remaining signs and symptoms of persistent disease (e.g. suppressed ADAMTS 13 activity level remain present)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cabometyx

## Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-4): 1) diagnosis of advanced renal cell carcinoma (RCC), 2) diagnosis of advanced RCC and using as a first-line treatment in combination with nivolumab, 3) member has previously been treated with sorafenib for hepatocellular carcinoma, 4) member has experienced disease progression following prior VEGFR-targeted therapy and is either radioactive iodine-refractory or is ineligible for radioactive iodine therapy for locally advanced or metastatic differentiated thyroid cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Calquence

## Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma, member has received at least one prior therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Camzyos

## Products Affected

- CAMZYOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy -AND- Left ventricular ejection fraction of greater than or equal to 55% -AND- Valsalva left ventricular outflow tract peak gradient of greater than or equal to 50 mmHg at rest or after provocation -AND- Therapeutic failure or intolerance to one of the following (1 or 2) or contraindication to all: 1) Non-vasodilating beta blocker (e.g. metoprolol) 2) non-dihydropyridine calcium channel blocker (e.g. diltiazem) -AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For reauthorization, prescriber attestation of no NYHA class worsening - AND- Not currently treated with and attestation Camzyos will not be used concomitantly with disopyramide, ranolazine, or combination therapy of beta-blocker and calcium channel blocker
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Caplyta

## Products Affected

- CAPLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Caprelsa

## Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic or progressive medullary thyroid cancer in patients with unresectable locally advanced or metastatic disease.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Carbaglu

## Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use as an adjunct therapy for acute hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA) -OR- maintenance therapy for chronic hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# Cayston

## Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cerdelga

## Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of type 1 Gaucher disease confirmed by the following A. or B. A) With one of the following symptoms (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm <sup>3</sup> . 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B) Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles -AND- Documentation of CYP2D6 metabolizer status (e.g. intermediate metabolizer).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CF drugs

## Products Affected

- **TOBI PODHALER**
- *tobramycin in 0.225 % nacl*
- *tobramycin inhalation*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization of tobramycin products, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cialis

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of benign prostatic hyperplasia (BPH) and trial/failure of at least two alternative medications in the following classes (alpha-1 adrenergic blockers and/or 5-alpha reductase inhibitors)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cibinqo

## Products Affected

- CIBINQO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe atopic dermatitis, documentation of one of the following (1 or 2): 1) trial & failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) -OR- 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement or severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Atopic Dermatitis, patients must have a trial/failure, intolerance, or contraindication to both preferred products: Dupixent and Rinvoq.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Cimzia

## Products Affected

- **CIMZIA POWDER FOR RECONST**
- **CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide) or all nonbiologic DMARDs are contraindicated. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or all are contraindicated. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- all nonbiologic DMARDs are contraindicated -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity, or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age for juvenile idiopathic arthritis or less than 18 years of age for all other indications.
Prescriber Restrictions	
Coverage Duration	12 months

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>For Crohn's disease, patients must have therapeutic failure or intolerance to 2 of the following preferred biologic products: a preferred adalimumab product, Stelara , Rinvoq, and Skyrizi SC. For Rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Xeljanz/Xeljanz XR and Rinvoq. For plaque psoriasis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Cosentyx, Otezla, Stelara SC, Skyrizi SC, and Enbrel. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For Psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cosentyx, Enbrel, a preferred adalimumab product, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For Juvenile Idiopathic Arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Rinvoz/Rinvoq LQ, Xeljanz/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For initial and induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended initial and induction therapy dosing regimens per indication.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Cinryze

## Products Affected

- CINRYZE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Member should not be on two prophylactic therapies simultaneously.
<b>Required Medical Information</b>	For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Cobenfy

---

## Products Affected

- **COBENFY**
- **COBENFY STARTER PACK**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Cometriq

## Products Affected

- **COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressive, metastatic medullary thyroid cancer
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Copiktra

## Products Affected

- COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is no longer responding or is intolerant to at least 2 prior therapies for chronic lymphocytic leukemia and small lymphocytic leukemia.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Corlanor

## Products Affected

- **CORLANOR ORAL SOLUTION**
- *ivabradine oral tablet 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) Normal sinus rhythm, 2) Resting heart rate greater than or equal to 70 beats per minute, 3) Left ventricular ejection fraction less than or equal to 35 percent, when applicable, 4) In adult patients (greater than or equal to 18 years), concurrent use, therapeutic failure, or intolerance to the maximum tolerated dose of one beta-blocker used for treatment of heart failure (i.e. bisoprolol, carvedilol, metoprolol succinate), or contraindication to beta-blocker use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For oral solution, attestation of inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Cosentyx

## Products Affected

- **COSENTYX (2 SYRINGES)**
- **COSENTYX PEN (2 PENS)**
- **COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML**
- **COSENTYX UNOREADY PEN**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs or all are contraindicated.
<b>Age Restrictions</b>	Deny if less than 6 years of age for moderate to severe plaque psoriasis - OR- less than 2 years of age for psoriatic arthritis -OR- less than 4 years of age for enthesitis-related arthritis -OR- less than 18 years of age for all other indications
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication. For hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Cotellic

## Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following for unresectable or metastatic melanoma (1-2): 1) BRAF V600E or V600K mutation status 2) Concomitant therapy with vemurafenib. For cobimetinib monotherapy, documentation of histiocytic neoplasms.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Cysteamine Ophthalmic Drops

---

## Products Affected

- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystinosis -AND- Attestation of accumulation of corneal cystine crystals
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Danziten

---

## Products Affected

- DANZITEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib -OR- member is newly diagnosed in the chronic phase.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Daraprim

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For primary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 100 cells/mm <sup>3</sup> -AND- Toxoplasma IgG positive -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For secondary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 200 cells/mm <sup>3</sup> . For secondary prophylaxis of cystoisosporiasis with CD4 count less than 200 cells/mm <sup>3</sup> or acute cystoisosporiasis infection: failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For primary prophylaxis of Pneumocystis jirovecii pneumonia: diagnosis of HIV - AND- CD4 count less than 200 cells/mm <sup>3</sup> -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# Daurismo

---

## Products Affected

- **DAURISMO ORAL TABLET 100 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of newly diagnosed Acute Myeloid Leukemia -AND- Used in combination with cytarabine -AND- At least one comorbidity that preclude use of intensive induction chemotherapy defined as one of the following: 1) Age greater than or equal to 75 2) Severe cardiac or pulmonary comorbidity 3) Reduced renal function 4) Hepatic impairment 5.) Physician attests patient is not a candidate for intensive induction therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Daybue

## Products Affected

- DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Rett syndrome confirmed by all of the following (1 thru 4): 1) Partial or complete loss of acquired purposeful hand skills, 2) Partial or complete loss of acquired spoken language, 3) Gait abnormalities: impaired or absence of ability, 4) Stereotypic hand movements such as hand wringing/squeezing, clapping/tapping, mouthing, and washing/rubbing automatisms
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Deferasirox

## Products Affected

- *deferasirox*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For chronic iron overload due to blood transfusions, transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -And- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw). For Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia (NTDT) Syndrome, LIC of at least 5 mg Fe/g dw -AND- serum ferritin greater than 300 mcg/L.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For reauthorization of chronic iron overload due to blood transfusion, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw. For reauthorization of chronic iron overload in NTDT syndrome, LIC greater than or equal to 3 mg Fe/g dw.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Diacomit

---

## Products Affected

- **DIACOMIT ORAL CAPSULE 250 MG, 500 MG**
- **DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravets syndrome - AND- Used in combination with clobazam
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Dihydroergotamine

## Products Affected

- *dihydroergotamine nasal*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- requires non-oral route of administration -AND- therapeutic failure or intolerance to generic sumatriptan nasal spray.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Doptelet

## Products Affected

- **DOPTelet (10 TAB PACK)**
- **DOPTelet (15 TAB PACK)**
- **DOPTelet (30 TAB PACK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -OR- Documentation of chronic immune thrombocytopenia -AND- Trial, intolerance, or inadequate response to corticosteroid therapy, immunoglobulin therapy or splenectomy -AND- One of the following (1 or 2): 1) Platelet count less than or equal to $50 \times 10^9/L$ and has significant mucous member bleeding or at least one risk factor for bleeding (e.g. hypertension, peptic ulcer disease). 2) Platelets count of less than or equal to $30 \times 10^9/L$
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For thrombocytopenia with chronic liver disease- 1 mo. For chronic immune thrombocytopenia- 12 mo.
Other Criteria	Platelet count is provided for applicable dosing.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Drizalma

## Products Affected

- **DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG, 30 MG, 40 MG, 60 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow tablets/capsules. For fibromyalgia, members must also have widespread bilateral pain above and below the waist for greater than 3 months duration -AND- At least 1 fibromyalgia-related symptom (e.g., cognitive impairment, fatigue, sleep disturbance, neurologic symptoms, exercise intolerance).
Age Restrictions	Deny if less than 18 years of age in the treatment of fibromyalgia, major depressive disorder, diabetic peripheral neuropathy and chronic musculoskeletal pain -OR- if less than 7 years of age in generalized anxiety disorder
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Dupixent

## Products Affected

- **DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML**
- **DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1-2: 1) moderate to severe atopic dermatitis 2) trial/failure, or intolerance to at least 1 topical corticosteroid -OR- If 2 years of age or older, topical calcineurin inhibitor (e.g. tacrolimus, pimecrolimus) -OR- The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin -OR- Documentation of 3-8: 3) moderate-to-severe asthma 4) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12mos or at least 1 asthma exacerbation requiring hospitalization in past 12mos 5) blood eosinophils greater than or equal to 150cells/uL or current daily or alternate-day oral corticosteroid (OCS) therapy 6) inadequate symptom control despite regular treatment with medium- or high-dose inhaled corticosteroids (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline) w/ or w/o OCS, unless intolerant or contraindicated to all 7) will continue treatment with a medium- or high-dose ICS and at least 1 add'l asthma controller medication w/ or w/o OCS -OR- Documentation of 8-9: 8) chronic rhinosinusitis with nasal polyposis 9) trial/ failure, intolerance or contraindication to intra-nasal corticosteroid and 14 day course of OCS. -OR- Documentation of 10-12: 10) eosinophilic esophagitis 11) esophageal eosinophil count greater than or equal to 15 eos/hpf on esophageal biopsy 12) clinical symptoms of esophageal dysfunction -OR- Documentation of 13: 13) prurigo nodularis -OR- Documentation of 14-16: 14) chronic obstructive pulmonary disease 15) blood eosinophils greater than or equal to 300cells/uL 16) inadequate symptom control despite regular treatment for at least 3 months with LAMA, LABA, and ICS, unless intolerant of or has contraindications to these agents.
Age Restrictions	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For atopic dermatitis reauthorization, attestation of positive clinical response to therapy. For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For EoE reauthorization, attestation of histological remission (less than 15 eos/hpf) on esophageal biopsy or reduced severity or frequency of clinical symptoms of esophageal dysfunction. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline. For COPD reauthorization, attestation of one of the following is required (1-4): 1) reduction in COPD symptoms, 2) improvement in exercise tolerance, 3) delayed disease progression, or 4) reduction in the number of COPD exacerbations.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Duvyzat

## Products Affected

- DUVYZAT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy with pathogenic mutation in the dystrophin gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EGFR Tyrosine Kinase Inhibitors

## Products Affected

- *erlotinib*
- **GILOTRIF**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Afatinib products: tumors with resistant EGFR mutations. Erlotinib products: use in NSCLC tumors with mutations other than those in FDA-approved indications. Use in combination with platinum based chemotherapy.
<b>Required Medical Information</b>	For afatinib, documentation of metastatic non-small cell lung cancer (NSCLC) -AND- one of the following (1-3): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutation 3) disease harbors non-resistant EGFR mutation (i.e., S768I, L861Q, G719X) -OR- documentation of squamous metastatic NSCLC and member has experienced progression on platinum-based chemotherapy. For erlotinib, documentation of metastatic NSCLC -AND- one of the following (1-2): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutations -OR- documentation of locally advanced, unresectable or metastatic pancreatic cancer -AND- all of the following (1-2): 1) using erlotinib as first-line therapy 2) using in combination with gemcitabine.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Emgality

## Products Affected

- **EMGALITY PEN**
- **EMGALITY SYRINGE**  
**SUBCUTANEOUS SYRINGE 120**  
**MG/ML, 300 MG/3 ML (100 MG/ML X 3)**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For episodic cluster headaches, characterized by severe or very severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes when left untreated -AND- Attack frequency of at least one attack every other day during the cluster period.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency is required. For reauthorization of cluster headache, attestation of reduction in the number of mean weekly cluster headaches from baseline is required.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Enbrel

## Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE  
25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Ankylosing Spondylitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis or Less than 4 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Epclusa

## Products Affected

- **EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG**
- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Epidiolex

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Lennox-Gastaut, Dravet syndromes or Tuberous Sclerosis Complex. For Lennox-Gastaut, trial and failure or intolerance of at least two standard of care treatments (e.g. lamotrigine, clobazam). For Lennox-Gastaut and Dravet syndromes, treatment is in combination with other conventional agents.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Eprontia

## Products Affected

- EPRONTIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow pills or food. For migraine, therapeutic failure, intolerance, or contraindication to two generic preventive migraine therapies. For partial onset seizures, primary generalized tonic-clonic seizures, or adjunctive treatment of Lennox-Gastaut Syndrome, therapeutic failure, contraindication, or intolerance to two generic anti-epileptic drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ergotamine

## Products Affected

- *ergotamine-caffeine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use to abort a vascular headache -AND- therapeutic failure or intolerance to a generic triptan -OR- documentation of use to prevent a vascular headache -AND- therapeutic failure or intolerance to generic prophylactic migraine medication
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Erivedge

## Products Affected

- **ERIVEDGE**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- if disease is locally advanced all of the following: 1) disease has recurred following surgery, or is not a candidate for surgery, 2) is not a candidate for radiation.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Erleada

## Products Affected

- **ERLEADA ORAL TABLET 240 MG, 60 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- the member meets one of the following (1 or 2) 1. Documentation of use in combination with a GnRH analog -OR- 2. The member has had a bilateral orchiectomy
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Evenity

## Products Affected

- **EVENITY SUBCUTANEOUS SYRINGE**  
**210MG/2.34ML ( 105MG/1.17MLX2)**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- Postmenopausal woman at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug. A cumulative lifetime approval of romosozumab will be limited to a coverage duration of 12 months.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Evrysdi

## Products Affected

- EVRYSDI ORAL RECON SOLN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of spinal muscular atrophy (SMA) -AND- Baseline motor function test results (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT) -AND- Not using concomitantly with Spinraza (nusinersen) -AND- Molecular genetic testing of 5q SMA showing Homozygous gene deletion, Homozygous conversion mutation or Compound heterozygote - AND- No history of gene replacement therapy to treat of SMA or experienced a decline in clinical status following gene replacement therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	For reauthorization, attestation of stable or clinically significant improvement in SMA-associated symptoms (e.g. stabilization or decreased decline in motor function compared to the predicted natural history trajectory of disease) or stable or improved motor function results compared to baseline (e.g. MFM, CHOP, HINE, RULM, HFMSE, 6MWT) -AND- Not using concomitantly with Spinraza (nusinersen) - AND- No history of gene replacement therapy to treat of SMA or experienced a decline in clinical status following gene replacement therapy.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Fabhalta

## Products Affected

- **FABHALTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For paroxysmal nocturnal hemoglobinuria (PNH), meets one of the following (1 or 2): 1) PNH mutant clones confirmed by flow cytometry, or 2) glycosylphosphatidylinositol-anchored proteins (GPI-AP)-deficient polymorphonuclear cells (PMNs) confirmed by flow cytometry -AND- meets one of the following (3-6): 3) anemia secondary to PNH (e.g. hemoglobin less than 10.5 g/dL with symptoms of anemia), 4) elevated lactate dehydrogenase (LDH) greater than or equal to 1.5 times the upper limit of normal, 5) history of a thromboembolic event, or 6) clinical findings of systemic complications (e.g. fatigue, hemoglobinuria, abdominal pain, dyspnea, dysphagia, erectile dysfunction, history of blood cell transfusion due to PNH) -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)) unless initially cross-titrating. For diagnosis of primary immunoglobulin A nephropathy (IgAN) confirmed by biopsy, member is at risk for rapid disease progression evidenced by one of the following (7 or 8): 7) Urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g or 8) Proteinuria greater than or equal to 1 g/day -AND- has experienced therapeutic failure, contraindication, or intolerance to a maximally tolerated dose of one of the following (9 or 10): 9) Angiotensin converting enzyme (ACE) inhibitor, or 10) Angiotensin receptor blocker (ARB) -AND- experienced therapeutic failure, contraindication, or intolerance to one of the following (1 or 2): 1) Filspari (sparsentan) or 2) Tarpeyo (budesonide)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For reauthorization of PNH, attestation of positive clinical response defined as one of the following (1-3): 1) hemoglobin stabilization or increase from baseline, 2) decrease in transfusions from baseline, or 3) decrease in LDH levels from baseline or reduction of hemolysis -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)). For reauthorization for IgAN, reduction in urine protein-to-creatinine (UPCR) or proteinuria from baseline.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Fanapt

---

## Products Affected

- FANAPT ORAL TABLET
- FANAPT ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Fasenra

## Products Affected

- **FASENRA PEN**
- **FASENRA SUBCUTANEOUS SYRINGE**  
10 MG/0.5 ML, 30 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe asthma and all of the following (1-4): 1) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos 2) blood eosinophils greater than or equal to 150cells/uL within the past 6 weeks or greater than or equal to 300cells/uL within the past 12 mos in without other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) 3) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), with or without oral corticosteroids (OCS), unless intolerant or contraindicated to all 4) will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, with or without OCS -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) and all of the following (5-6): 5) history of relapsing or refractory disease 6) will be receiving standard of care while on Fasenra therapy with glucocorticoid treatment (e.g. prednisone or prednisolone), with or without immunosuppressive therapy (e.g. cyclosporine, leflunomide, azathioprine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For severe asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EPGA reauthorization, attestation of one of the following is required (5-8): 5) reduction in the frequency and/or severity of relapses, 6) reduction or discontinuation of doses of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Fecal Microbiota Products

---

## Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a recent diagnosis of recurrent <i>Clostridioides difficile</i> infection (CDI) -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	For reauthorization, attestation of recurrent CDI episodes after administration of the initial fecal microbiota product -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ferriprox

## Products Affected

- *deferiprone*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of transfusional iron overload due to thalassemia syndromes, sickle cell disease and other anemias -AND- transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -AND- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw)
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Initial authorization requires trial and failure of deferasirox (generic Exjade) and reauthorization requires failure of generic deferasirox (generic Exjade) if not previously trialed. For reauthorization, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Fetzima

## Products Affected

- **FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)**      **CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG**
- **FETZIMA ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder and trial and failure of two other generic antidepressants.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Filsuvez

## Products Affected

- FILSUEZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB) -AND- at least one open wound.
Age Restrictions	Deny if less than 6 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement in targeted wound(s) - AND- member requires additional courses of treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Fintepla

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravet syndrome or Lennox-Gastaut syndrome. For Lennox-Gastaut syndrome, therapeutic failure, contraindication, or intolerance to at least 2 standard of care treatments (e.g. lamotrigine, clobazam).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Firazyr

## Products Affected

- *icatibant*
- **SAJAZIR**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
<b>Required Medical Information</b>	For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type I & II with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) type III with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE or FXII mutation 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Firdapse

## Products Affected

- **FIRDAPSE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Firmagon

---

## Products Affected

- **FIRMAGON KIT W DILUENT SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Forteo

## Products Affected

- *teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 months
<b>Other Criteria</b>	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of teriparatide will be limited to a coverage duration of 24 months in the absence of provider attestation that the member remains at or has returned to having a high risk for fracture
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Fotivda

## Products Affected

- FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received at least two prior systemic therapies.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Fruzaqla

## Products Affected

- **FRUZAQLA ORAL CAPSULE 1 MG, 5 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- member has received previous threatment with a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy -AND- member has received previous treatment with an anti-VEGF therapy -AND- one of the following, if member is RAS wild-type (1-2): 1) member has received previous therapy with an anti-EGFR therapy 2) prescriber attests that treatment with an anti-EGFR therapy would not be medically appropriate.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Furoscix

## Products Affected

- FUROSCIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of heart failure -AND- chronic heart failure medication regimen includes an oral loop diuretic (e.g. furosemide, bumetanide, torsemide) -AND- treatment with oral diuretics will replace the use of Furoscix as soon as practical.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Gabapentin

## Products Affected

- *gabapentin oral capsule 100 mg, 300 mg, 400 mg*
- *gabapentin oral solution 250 mg/5 ml*
- *gabapentin oral tablet 600 mg, 800 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. When using concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Gattex

## Products Affected

- GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of short bowel syndrome (SBS) having less than 200 cm of functional small bowel -AND- Dependence on parenteral/intravenous nutrition -AND- weight of at least 10 kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Gavreto

## Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, disease is RET fusion-positive as detected by an FDA approved test. For advanced or metastatic thyroid cancer, all of the following (1-2): 1) disease is RET fusion-positive 2) if radioactive iodine is appropriate, the member is radioactive iodine-refractory.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Gilenya

## Products Affected

- *fingolimod*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Glatiramer

## Products Affected

- **COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- **GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Gleevec

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult aggressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvant treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Gleostine

## Products Affected

- GLEOSTINE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or metastatic brain tumor(s) -AND- member has previously received surgical and/or radiotherapeutic procedure(s). Documentation of Hodgkin's lymphoma -AND- all of the following (1-2): 1) using lomustine in combination with other chemotherapies 2) member has experienced disease progression with initial chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLP1

## Products Affected

- MOUNJARO
- OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2
- MG/DOSE (8 MG/3 ML)
- RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG
- TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Growth Hormone

## Products Affected

- NORDITROPIN FLEXPPO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, bone age, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria. Additionally for growth failure due to chronic kidney disease, glomerular filtration rate is less than 89ml/min per 1.73m*2. For HIV wasting and cachexia, Concurrent use of antiretroviral therapy -AND- weight loss of at least 10 percent from baseline. For short bowel syndrome, receiving management for short bowel syndrome, including specialized nutritional support -AND- less than 200 cm of functional small bowel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and bone age, as applicable to meet standard continuation of therapy guidelines. For reauthorization of HIV wasting and cachexia, attestation of increase in weight from start of therapy. For reauthorization of short bowel syndrome, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# High-risk meds

## Products Affected

- *amitriptyline*
- *benztropine oral*
- *clomipramine*
- *doxepin oral capsule*
- *doxepin oral concentrate*
- *doxepin oral tablet*
- *hydroxyzine hcl oral tablet*
- *imipramine hcl*
- *promethazine oral*
- *trimipramine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	For all medications subject to this PA group, the following information (1 through 3) is required: 1. Documentation of diagnosis 2. Explanation of risk-benefit profile favoring use of the high-risk medication 3. Attestation of an intent to monitor and address treatment-related adverse events. For target tricyclic antidepressants (TCAs), in addition to criteria 1 through 3 above, trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs for the same indication, if available, is required (e.g. SSRIs and SNRIs). If using a TCA for a medically-accepted indication not shared by the safer alternatives listed, then no trial of alternatives is required.
<b>Age Restrictions</b>	Automatic approval if less than 65 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Doxepin doses less than or equal to 6 mg per day will receive automatic approval.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# High-risk meds phenobarbital

## Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coverage is not provided for use in sedation/insomnia
<b>Required Medical Information</b>	For use in seizures the following are required: 1. Explanation of risk-benefit profile favoring use of the high-risk medication 2. Attestation of an intent to monitor and address treatment-related adverse events. 3. For new starts, the trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs used for seizures (e.g. carbamazepine, lamotrigine) is required.
<b>Age Restrictions</b>	Automatic approval if less than 65 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off Label Uses</b>	Seizure disorders
<b>Part B Prerequisite</b>	No

# Humira

## Products Affected

- **CYLTEZO(CF)**
- **CYLTEZO(CF) PEN**
- **CYLTEZO(CF) PEN CROHN'S-UC-HS**
- **CYLTEZO(CF) PEN PSORIASIS-UV**
- **HUMIRA PEN**
- **HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML**
- **HUMIRA(CF)**
- **HUMIRA(CF) PEN**
- **HUMIRA(CF) PEN CROHNS-UC-HS**
- **HUMIRA(CF) PEN PSOR-UV-ADOL HS**
- **YUFLYMA(CF)**
- **YUFLYMA(CF) AI CROHN'S-UC-HS**
- **YUFLYMA(CF) AUTOINJECTOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 2 immunosuppressants.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	
Coverage Duration	12 months

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. Please Note: Covered preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Ibrance

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of HR-positive, HER2-negative advanced or metastatic breast cancer -AND- meets one of the following (1. or 2.): 1) documentation of use with an aromatase inhibitor as initial endocrine-based therapy -OR- 2) documentation of use with fulvestrant in patients with disease progression following endocrine therapy. Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (3-6): 3) disease is HR-positive, HER2-negative 4) disease is PIK3CA-mutated, as detected by an FDA-approved test 5) the member is using in combination with inavolisib and fulvestrant 6) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# Ibsrela

## Products Affected

- IBSRELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of irritable bowel syndrome with constipation -AND- failure or intolerance to Linzess -AND- if member is female, failure or intolerance to lubiprostone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Iclusig

## Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of newly-diagnosed chronic phase CML
Required Medical Information	Documentation of T3151+ chronic myeloid leukemia (CML) -OR- documentation of chronic phase CML and member has experienced resistance or intolerance to at least two prior kinase inhibitors -OR- documentation of accelerated phase or blast phase CML and no other kinase inhibitor is indicated -OR- member is using Iclusig as monotherapy and meets one of the following (1-2): 1) documentation of T3151+ acute lymphoblastic leukemia (ALL) 2) documentation of Ph+ ALL and no other tyrosine kinase inhibitor therapy is indicated -OR- member has newly diagnosed Ph+ ALL and is using Iclusig in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Idhifa

## Products Affected

- **IDHIFA ORAL TABLET 100 MG, 50 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA approved test
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IG

## Products Affected

- **BIVIGAM**
- **GAMMAGARD LIQUID**
- **GAMMAGARD S-D (IGA**
- **GAMMAKED INJECTION SOLUTION**
- **GAMMAPLEX**
- **GAMMAPLEX (WITH SORBITOL)**
- **GAMUNEX-C INJECTION SOLUTION**
- **1 GRAM/10 ML (10 %)**
- **OCTAGAM**
- **PANZYGA**
- **PRIVIGEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For select diagnoses the following apply 1) For Myasthenia Gravis Syndrome, documentation that the patient is refractory to other standard therapies (e.g., cholinesterase inhibitors, corticosteroids, azathioprine) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 2) For Multiple Sclerosis, patient is refractory to other standard therapies (e.g., interferons) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 3) For B-cell CLL, associated with recurrent bacterial infections OR with Associated Hypogammaglobulinemia defined as IgG level less than 600mg/dL or evidence of a specific antibody deficiency. 4) For Bone Marrow Transplantation, the member is 20 years of age or older and within the first 100 days after transplantation. 5) For Dermatomyositis/Polymyositis, trial and failure, intolerance, or contraindication to standard fist line therapy (i.e. corticosteroids or immunosuppressants). 6) For Pediatric HIV, the patient is less than 13 y.o. who have entry CD4 lymphocyte count greater than or equal to 200/mcl and IgG less than 400mg/dL OR recurrent bacterial infections. 7) For Guillain-Barre syndrome, impaired function by objective assessment and/or objective findings on physical exam at the time of initial therapy and IVIG therapy must be initiated within 2 weeks of symptom onset. 8) For Autoimmune Mucocutaneous Blistering Diseases (e.g. Stevens-Johnson Syndrome), trial and failure, intolerance, or contraindication to conventional therapy (e.g. corticosteroids) or the patient has rapidly progressive disease in which a clinical response could not be affected quickly enough using conventional agents.
Age Restrictions	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Covered under Part B when administered in the home to a member with a diagnosis of primary immunodeficiency disease
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off Label Uses</b>	Myasthenia Gravis syndrome, Multiple Sclerosis, Polymyositis, Bone Marrow Transplant, Pediatric HIV, Guillain-Barre syndrome, Autoimmune Mucocutaneous Blistering Diseases
<b>Part B Prerequisite</b>	No

# Imbruvica

## Products Affected

- **IMBRUVICA ORAL CAPSULE 140 MG, 280 MG, 420 MG**
- **IMBRUVICA ORAL SUSPENSION**
- **IMBRUVICA ORAL TABLET 140 MG,**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic graft versus host disease, previous treatment with at least 1 prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For suspension, inability to swallow oral tablets or oral capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Imkeldi

## Products Affected

- **IMKELDI**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult aggressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvant treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Documentation of an inability to swallow oral tablets is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Increlex

## Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe primary IGF-1 deficiency and all of the following: 1) Normal or elevated response (greater than 10 ng/ml) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 2) Serum IGF-1 concentration that is less than or equal to three (3) standard deviations below the normal value based on laboratory reference range. 3) Height less than or equal to three (3) standard deviations below normal (at or below the third percentile for gender and age). 4) Member has open epiphyses. -OR- Documentation of diagnosis of growth hormone deficiency caused by gene deletion and all of the following: 1) Growth velocity at least 2 standard deviations below the age-appropriate mean or height at least 2.25 standard deviations below the age-appropriate mean. 2) Subnormal response (less than 10 ng/mL) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 3) Development of neutralizing antibodies to growth hormone product(s). 4) Member has open epiphyses
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation member has a growth velocity of at least 2 cm/year -AND- member has open epiphyses.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Inlyta

## Products Affected

- INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-2): 1) if using axitinib as first line therapy for advanced renal cell carcinoma, member is using axitinib in combination with avelumab or pembrolizumab 2) member has been treated with at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Inqovi

## Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation chronic myelomonocytic leukemia. Documentation of myelodysplastic syndrome -AND- One of the following (1 or 2): 1) French American-British MDS subtypes of refractory anemia, refractory anemia with ringed sideroblasts or refractory anemia with excess blasts. 2) International Prognostic Scoring System group of intermediate-1, intermediate-2 or high-risk.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Inrebic

## Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate-2 or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) -AND- If a new start, baseline platelet count of greater than or equal to $50 \times 10^9/L$
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Insulin Supplies

## Products Affected

- ALCOHOL PADS
- GAUZE PAD TOPICAL BANDAGE 2 X 2 "

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation the product is being used for the delivery of insulin into the body.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Interferon Alfa

## Products Affected

- PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Interferon Beta

## Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT
- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IPF Agents

## Products Affected

- **OFEV**
- *pirfenidone oral capsule*
- *pirfenidone oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 50% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30%. For Ofev only, documentation of systemic sclerosis-associated interstitial lung disease -AND- baseline forced vital capacity (FVC) of at least 40% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- documentation of a high-resolution chest computed tomography (CT) scan demonstrating greater than or equal to 10% pulmonary fibrosis. For Ofev only, documentation of chronic fibrosing interstitial lung disease with progressive phenotype -AND- high resolution chest computing tomography (HRCT) scan demonstrating greater than 10% fibrosing disease -AND- baseline forced vital capacity (FVC) of at least 45% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- disease progression in previous 24 months shown by one of the following : 1. Relative decline in FVC greater than or equal to 10% predicted 2. Relative decline in FVC greater than or equal to 5% but less than 10% predicted and either worsening of respiratory symptoms or increased extent of fibrotic changes on HRCT 3. Worsening of respiratory symptoms and increasing extent of fibrotic changes on HRCT
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Iressa

## Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	Use in tumors with EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.
Required Medical Information	Documentation of diagnosis -AND- the following: 1) EGFR exon 19 deletion mutations or exon 21 (L858R) mutations as detected by an FDA-approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Itovebi

## Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (1-4): 1) disease is HR-positive, HER2-negative 2) disease is PIK3CA-mutated, as detected by an FDA-approved test 3) the member is using in combination with palbociclib and fulvestrant 4) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Itraconazole

## Products Affected

- *itraconazole oral capsule*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. If using for diagnosis of onychomycosis, confirmation through positive laboratory testing (e.g. KOH preparation, fungal culture, or nail biopsy) is required.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Onychomycosis: 3 months. All other indications: 3 months initial, 12 months reauth
<b>Other Criteria</b>	Documentation of trial/failure or intolerance of amphotericin b must be provided for approval in patients with aspergillosis.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Ivermectin Oral

## Products Affected

- *ivermectin oral tablet 3 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of strongyloidiasis of the intestinal tract (non-disseminated disease) or onchocerciasis -AND- Member weighs greater than or equal to 15kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Iwilfin

## Products Affected

- IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of high-risk neuroblastoma (HRNB) -AND- partial response to anti-glycolipid disialoganglioside (GD2) immunotherapy (e.g., dinutuximab, naxitamab).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Jakafi

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) and if a new start, baseline platelet count of greater than or equal to $50 \times 10^9/L$ -OR- documentation of polycythemia vera and inadequate response or intolerance to hydroxyurea -OR- Documentation of steroid refractory acute graft-versus-host disease and prior therapy with at least one systemic corticosteroid -OR- Documentation of chronic graft-versus-host disease with prior failure of at least one systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Jaypirca

## Products Affected

- **JAYPIRCA ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory mantle cell lymphoma, member has received at least two (2) previous lines of systemic therapy, at least one (1) of which was a BTK inhibitor. For Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia, member has received at least two (2) prior lines of therapy, including at least one (1) from all of the following classes (1-2): 1) BTK inhibitor 2) BCL-2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Joenja

## Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of activated phosphoinositide 3-kinase delta syndrome (APDS) with genetic confirmation of variant in PIK3CD or PIK3R1 gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Kalydeco

## Products Affected

- **KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG**
- **KALYDECO ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to ivacaftor based on clinical and or in vitro assay (e.g. G551D, G1244E, G1349D)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Kerendia

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than or equal to 25 mL/min/1.73 m*2, serum potassium greater than 5.5 mEq/L
Required Medical Information	Documentation of type 2 diabetes mellitus with chronic kidney disease - AND- one of the following (1. 2. or 3.): 1) concomitant use of a sodium-glucose Cotransporter-2 (SGLT2) inhibitor 2) therapeutic failure to at least one SGLT2 inhibitor or 3) contraindication or intolerance to at least one SGLT2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that signs or symptoms of hyperkalemia are not present.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Kesimpta

## Products Affected

- KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Kevzara

## Products Affected

- KEVZARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For polymyalgia rheumatica (PMR), one of the following (1-3): 1) inadequate response to corticosteroids, 2) intolerance to corticosteroid taper, or 3) used in combination with corticosteroid tapering course. For juvenile idiopathic arthritis, weight greater than or equal to 63 kg -AND- one of the following (4-5): 4) inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide), or 5) requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
<b>Age Restrictions</b>	Deny if less than 18 years of age for rheumatoid arthritis and polymyalgia rheumatica.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Rinvoq/Rinvoq LQ, Xeljanz/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Kineret

## Products Affected

- KINERET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), therapeutic failure or intolerance to at least one (1) corticosteroid, or all corticosteroids are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Kisqali

## Products Affected

- **KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG** (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)
- **KISQALI ORAL TABLET 200 MG/DAY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative -AND- one of the following (1-3): 1) member is using ribociclib in combination with an aromatase inhibitor as initial endocrine-based therapy 2) member is using ribociclib in combination with fulvestrant and member is using fulvestrant as initial endocrine-based therapy or member has experienced disease progression on endocrine therapy. 3) disease is classified as stage II or stage III early breast cancer at high risk of recurrence, ribociclib is being used in combination with an aromatase inhibitor as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Korlym

## Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperglycemia secondary to hypercortisolism in patients with endogenous Cushing's syndrome -AND- Patient is not a candidate for surgery or where surgery has failed -AND- one of the following (1 or 2): 1) Diagnosis of diabetes with trial and failure, intolerance, or contraindication to one previous therapy for Type 2 Diabetes (e.g. metformin, sulfonylureas, insulin) or using in addition to a therapy for Type 2 diabetes. 2) Glucose intolerance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Koselugo

---

## Products Affected

- **KOSELUGO ORAL CAPSULE 10 MG, 25 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- for neurofibromatosis type 1 (NF1), documentation of symptomatic, inoperable plexiform neurofibromas (PN)
<b>Age Restrictions</b>	Deny if less than 2 years of age or older than 17 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Krazati

## Products Affected

- KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of locally advanced or metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA-approved test -AND- using in combination with cetuximab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Kuvan

## Products Affected

- **JAVYGTOR**
- *sapropterin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of PKU -AND- documented baseline Phe level greater than 6 mg/dL -AND- clinical documentation of current weight -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, initial therapy has resulted in decrease in phenylalanine levels from baseline or current phenylalanine levels within the range of 120-360 micromol/L -AND- clinical documentation of current weight is required -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Latuda

## Products Affected

- *lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for bipolar 1 disorder, documentation of trial and failure or intolerance to one other formulary medication indicated in bipolar 1 disorder (e.g. quetiapine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lazcluze

## Products Affected

- **LAZCLUZE ORAL TABLET 240 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) -AND- all of the following (1-4): 1) disease harbors EGFR exon 19 deletions or EGFR exon 21 (L858R) substitution mutation, 2) disease is locally advanced or metastatic, 3) member is treatment naive for advanced disease, 4) using in combination with amivantinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lenvima

## Products Affected

- LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of differentiated thyroid cancer -AND- meets all of the following (1-3): 1) disease is locally recurrent or metastatic 2) disease is progressive 3) disease is radioactive iodine refractory. Documentation of advanced renal cell carcinoma -AND- one of the following (1-2): 1) member is using lenvatinib in combination with pembrolizumab and is using lenvatinib and pembrolizumab as first-line treatment 2) member is using lenvatinib in combination with everolimus and has experienced therapeutic failure or intolerance to one prior anti-angiogenic therapy. Documentation of unresectable hepatocellular carcinoma. Documentation of endometrial cancer -AND- meets all of the following (1-5): 1) member is using lenvatinib in combination with pembrolizumab 2) disease is advanced 3) disease is not classified as microsatellite instability-high or disease is classified as mismatch repair proficient as determined by an FDA-approved test 4) member has experienced disease progression following prior systemic therapy 5) member is not a candidate for curative surgery or radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Leukine

## Products Affected

- LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis: following induction chemotherapy in patients who are 55 years or older with acute myelogenous leukemia (AML) -OR- mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation -OR- acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation -OR- acceleration of myeloid reconstitution following allogeneic BMT -OR- treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic BMT -OR- following acute exposure to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Libervant

## Products Affected

- LIBERVANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 2 years of age or greater than 5 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lidoderm

## Products Affected

- *lidocaine topical adhesive patch, medicated 5 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) or diabetic peripheral neuropathy (DPN) -AND- One of the following (1-3): 1) trial and failure of 1 other agent used to treat diagnosis (e.g. gabapentin for PHN, duloxetine for DPN), 2) inability to swallow oral medication, 3) unable to take an oral medication due to potential adverse events (e.g. sedation).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Diabetic peripheral neuropathy
Part B Prerequisite	No

# Litfulo

## Products Affected

- LITFULO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For severe alopecia areata, therapeutic failure/intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Livtency

## Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of CMV prophylaxis.
Required Medical Information	Documentation of refractory cytomegalovirus infection or disease as evidenced by antigenemia or polymerase chain reaction (PCR) test -AND- all of the following (1-3): 1) member weighs at least 35 kg, 2) member is a recipient of hematopoietic stem cell transplant -OR- solid organ transplant. 3) member has experienced therapeutic failure to one of the following: ganciclovir, valganciclovir, cidofovir, or foscarnet.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of a previous reduction in CMV DNA level -AND- documentation of one of the following (1-3): 1) new onset symptomatic CMV infection, 2) virologic relapse with treatment-emergent maribavir resistance or 3) continued antiviral treatment is required to achieve virologic clearance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lokelma

## Products Affected

- **LOKELMA**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 7.4 mmol/L on at least two (2) screenings -AND- Modification of medications to reduce serum potassium levels were not successful, when applicable
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For reauthorization, attestation of reduction in serum potassium levels following Lokelma administration and continued treatment for hyperkalemia is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Lonsurf

## Products Affected

- LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer as a single agent or in combination with bevacizumab in patients who have previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy -OR- documentation of metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/neu-targeted therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lorbrena

## Products Affected

- **LORBRENA ORAL TABLET 100 MG, 25 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of ALK-positive metastatic non-small cell lung cancer (NSCLC)
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Lotronex

## Products Affected

- *alosetron oral tablet 0.5 mg, 1 mg*

PA Criteria	Criteria Details
Exclusion Criteria	For irritable bowel syndrome (IBS): Exclude if male gender
Required Medical Information	Documentation of chronic severe diarrhea-predominant IBS -AND- trial and failure or intolerance to one anti-diarrheal (e.g. loperamide), anti-spasmodic, or tricyclic antidepressant, or contraindication to all
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that symptoms of IBS continue to persist AND positive clinical response.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lumakras

## Products Affected

- LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lupron Depot Ped

## Products Affected

- **LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG**
- **LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)**
- **LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of central precocious puberty -AND- advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3 IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response to therapy defined as pre-pubertal slowing/decline, normalization of FSH, normalization LH, normalization of bone age, normalization of bone age, normalization of estradiol level or normalization of testosterone level
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lynparza

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) in complete or partial response to first-line platinum-based chemotherapy 2) diagnosis of deleterious or suspected deleterious BRCA mutated disease or disease is associated with homologous recombination deficiency (HRD) positive status with a deleterious or suspected deleterious BRCA mutation or genomic instability and will be using in combination with bevacizumab. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) deleterious or suspected deleterious germline or somatic BRCA mutation 2) in complete or partial response to platinum-based chemotherapy. For deleterious or suspected deleterious gBRCAm, HER2-negative breast cancer, 1 of the following (1-2): 1) classified as high-risk, early breast cancer and has been treated with neoadjuvant or adjuvant chemotherapy 2) has been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting and if hormone receptor (HR)-positive, has been previously treated with or considered inappropriate for treatment with endocrine therapy. For metastatic pancreatic adenocarcinoma, all of the following (1-2): 1) a deleterious or suspected deleterious gBRCA mutation 2) did not progress on at least 16 weeks of a first-line platinum-based chemotherapy regimen. For deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene mutation metastatic castration-resistant prostate cancer, all of the following (1-2): 1) progressed following prior treatment with enzalutamide or abiraterone 2) concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy. For deleterious or suspected deleterious BRCA mutation metastatic castration-resistant prostate cancer, using in combination with all of the following (1-2): 1) abiraterone 2) prednisone or prednisolone.
Age Restrictions	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Lyrica

## Products Affected

- *pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg*
- *pregabalin oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of DPN and trial/failure or intolerance to duloxetine -OR- PHN and trial/failure or intolerance to gabapentin -OR- Partial-onset seizures and trial/failure or intolerance to two AEDS -OR- Neuropathic pain associated with spinal cord injury -OR- Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine. When using pregabalin products concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Lytgobi

## Products Affected

- **LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-2): 1) disease harbors FGFR2 fusions or other rearrangements 2) member has experienced therapeutic failure or intolerance to at least one prior therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Mavyret

---

## Products Affected

- MAVYRET ORAL PELLETS IN PACKET
- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Megace

## Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Mekinist

## Products Affected

- MEKINIST ORAL RECON SOLN
- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For trametinib monotherapy, documentation of unresectable or metastatic melanoma -AND- meets all of the following (1-2): 1) member has a BRAF V600E or V600K mutation 2) member is BRAF inhibitor treatment naive. For use in combination with dabrafenib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with dabrafenib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation. For use in combination with dabrafenib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Mekinist powder for oral solution, attestation of inability to swallow Mekinist (trametinib) tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Mektovi

## Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, all of the following (1-2): 1) BRAF V600E or V600K mutation status 2) using in combination with encorafenib. For metastatic non-small cell lung cancer, all of the following (1-2): 1) BRAF V600E mutation status 2) using in combination with encorafenib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Metyrosine

## Products Affected

- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pheochromocytoma defined by 1 of the following (1 or 2): 1) Elevated metanephrines in plasma or urine, or 2) Tumor evidence from CT scan or MRI -AND- Documentation of 1 of the following (3., 4., or 5.): 3) Planned resection surgery, 4) Resection surgery is contraindicated, or 5) malignant pheochromocytoma. -AND- therapeutic failure, contraindication, or intolerance to a selective alpha-blocker (e.g., doxazosin, prazosin, terazosin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Mulpleta

## Products Affected

- MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -AND- trial and failure or intolerance to Doptelet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Myasthenia Gravis

## Products Affected

- **ZILBRYSQ SUBCUTANEOUS  
SYRINGE 16.6 MG/0.416 ML, 23  
MG/0.574 ML, 32.4 MG/0.81 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized myasthenia gravis (gMG) -AND- Anti-acetylcholine receptor (AChR) antibody-positive -AND- Therapeutic failure, contraindication, or intolerance to generic pyridostigmine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of gMG signs/symptoms improvement (e.g., speech, swallowing, mobility, and/or respiratory function) -OR- decreased gMG exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Namzarin

## Products Affected

- NAMZARIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and trial/failure of generic memantine and generic donepezil
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nayzilam

## Products Affected

- NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nemluvio

## Products Affected

- NEMLUVIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of prurigo nodularis -OR- Documentation of moderate to severe atopic dermatitis and one of the following (1-3): 1) trial and failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus), 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement, or 3) severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For prurigo nodularis, must follow recommended dosing guidelines based upon weight. For atopic dermatitis, must follow recommended dosing based on FDA approved dosing guidelines. For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Nerlynx

## Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of early-stage, HER2-positive breast cancer -AND- meets all of the following: 1)member has received adjuvant trastuzumab-based therapy 2) member is using neratinib as a single agent. Documentation of advanced HER-2 positive, or metastatic HER2-positive breast cancer and meets all of the following 1) using neratinib in combination with capecitabine 2) member has received two or more prior anti-HER2 based regimens in the metastatic setting.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nexavar

## Products Affected

- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally recurrent or metastatic, progressive, differentiated thyroid carcinoma, refractory to radioactive iodine treatment
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nexletol

## Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For HeFH, diagnosis supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite Simon Broome register criteria, or definite on the Make Early Diagnosis to Prevent Early Deaths tool -AND- LDL-C greater than 100 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe. For Hyperlipidemia with ASCVD or Hyperlipidemia with attestation of high risk for CVD, LDL-C greater than 70 mg/dL -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe. For Primary Hyperlipidemia not associated with ASCVD or HeFH, LDL-C greater than 70 mg/dL -AND- Attestation of use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible -AND- Therapeutic failure to a maximally tolerated statin or documentation of statin intolerance -AND- Therapeutic failure, intolerance or contraindication to ezetimibe.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For reauthorization, documentation showing an LDL-C reduction from baseline -AND- for HeFH and Primary Hyperlipidemia (not associated with ASCVD or HeFH), attestation of continued use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statin which resolved upon discontinuation of statin or documentation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Ninlaro

## Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- previous treatment with at least 1 prior therapy -AND- used in combination with lenalidomide and dexamethasone
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nitisinone

## Products Affected

- *nitisinone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary tyrosinemia type 1 (HT-1) confirmed by biochemical or genetic testing -AND- dietary restriction of tyrosine and phenylalanine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Northera

## Products Affected

- *droxidopa oral capsule 100 mg, 200 mg, 300 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of neurogenic orthostatic hypotension caused by primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency or non-diabetic autonomic neuropathy -AND- documentation of inadequate response, intolerance or contraindication to preferred generic alternative midodrine.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	For reauthorization, attestation of increase from baseline of systolic or diastolic blood pressure upon standing -OR- attestation of decrease from baseline of neurogenic orthostatic hypotension symptoms upon standing (e.g., dizziness, feeling faint, etc.).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Nubeqa

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-metastatic castration-resistant prostate cancer -AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy. Documentation of metastatic hormone-sensitive prostate cancer -AND- will be using in combination with docetaxel -AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nucala

## Products Affected

- **NUCALA SUBCUTANEOUS AUTO-INJECTOR**
- **NUCALA SUBCUTANEOUS RECON SOLN**
- **NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Diagnosis of severe asthma and all of the following (1-4): 1) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos 2) Blood eosinophils greater than or equal to 150cells/uL within past 6 wks or greater than or equal to 300cells/uL within past 12 mos without other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) 3) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), with or without oral corticosteroids (OCS) 4) will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, with or without OCS -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) with history of relapsing or refractory disease and will be receiving concomitant glucocorticoid treatment with or without immunosuppressive therapy -OR- Diagnosis of hypereosinophilic syndrome (HES) without identifiable non-hematologic secondary cause for greater than or equal to 6 mos and both of the following (5-6): 5) At least 2 HES flares (HES-related worsening of clinical symptoms or blood eosinophil counts requiring an escalation in therapy) within past 12 mos 6) Stable on HES therapy for at least 4 wks (chronic or episodic OCS, immunosuppressive or cytotoxic therapy) -OR- Diagnosis of chronic rhinosinusitis with nasal polyps (CRSwNP) and trial/failure, contraindication, or intolerance to intranasal corticosteroid.</p>
Age Restrictions	
Prescriber Restrictions	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For Severe Asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EGPA reauthorization, attestation of one of the following is required (5-8): 5) reduction in frequency and/or severity of relapses, 6) reduction or discontinuation of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms. For HES reauthorization, attestation of one of the following is required (9-10): 9) reduction in frequency of HES flares, or 10) maintenance or reduction in background HES therapy requirements. For CRSwNP reauthorization, attestation of one of the following is required (11-12): 11) decrease in nasal polyp score, or 12) reduction in nasal congestion/obstruction severity score.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Nuedexta

## Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting improvement in symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Nuplazid

## Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hallucinations and delusions associated with Parkinson's disease psychosis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nurtec

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine Prevention, defined as 4-14 migraine days per month, the following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes. For acute treatment of migraine with or without aura, trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency. For reauthorization of acute treatment of migraine, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Nuvigil

## Products Affected

- *armodafinil*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For reauthorization, provider attestation of improvement in daytime sleepiness is required.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Ocaliva

## Products Affected

- OCALIVA

PA Criteria	Criteria Details
Exclusion Criteria	Compensated cirrhosis with evidence of portal hypertension
Required Medical Information	Documentation of primary biliary cholangitis -AND- trial and failure, contraindication, or intolerance to ursodiol monotherapy -AND- will use concomitantly with ursodiol unless contraindicated or intolerant.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Octreotide

## Products Affected

- *octreotide acetate injection solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- therapeutic failure or cannot be treated with surgical resection, pituitary irradiation or bromocriptine mesylate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Odomzo

## Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced basal cell carcinoma (laBCC) that has recurred following surgery or radiation therapy or for use in patients who are not candidates for surgery or radiation therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ogsiveo

## Products Affected

- **OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of progressing desmoid tumor(s) requiring systemic therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Ojemda

## Products Affected

- **OJEMDA ORAL SUSPENSION FOR RECONSTITUTION** **MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)**
- **OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory pediatric low-grade glioma - AND- one of the following (1-2): 1) BRAF fusion or rearrangement, or 2) BRAF V600K mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ojjaara

## Products Affected

- OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of anemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Olumiant

## Products Affected

- OLUMIANT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Part A covered for Covid-19 in hospitalized patients
<b>Required Medical Information</b>	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis an inadequate response or intolerance to at least one non-biologic DMARD (e.g., methotrexate, leflunomide). For severe alopecia areata, inadequate response or intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
<b>Age Restrictions</b>	Deny if less than 18 years of age for Rheumatoid Arthritis and Alopecia Areata
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR, with at least 1 being a tumor necrosis factor blocker. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Onfi

## Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Onureg

## Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute myeloid leukemia that has achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy -AND- Inability to complete intensive curative therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Orencia

## Products Affected

- **ORENCIA CLICKJECT**
- **ORENCIA SUBCUTANEOUS SYRINGE**  
125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For juvenile idiopathic rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
<b>Age Restrictions</b>	Deny if less than 18 years of age for Rheumatoid Arthritis or less than 2 years of age for Juvenile Idiopathic Arthritis and Psoriatic Arthritis
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients over 18 years of age must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Xeljanz/Xeljanz XR, Otezla and Stelara SC, Rinvoq, and Skyrizi SC. For juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Xeljanz/Xeljanz solution and Actemra SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Orgovyx

## Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Orkambi

## Products Affected

- **ORKAMBI ORAL GRANULES IN PACKET**
- **ORKAMBI ORAL TABLET**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of cystic fibrosis and homozygous F508del mutation
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Orserdu

## Products Affected

- **ORSERDU ORAL TABLET 345 MG, 86 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-4): 1) member is male or a postmenopausal female 2) tumor status is ER-positive, HER2-negative 3) an ESR1 gene mutation is present in the tumor 4) member has experienced disease progression on or after an endocrine based regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OTEZLA

## Products Affected

- **OTEZLA** (47)
- **OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For plaque psoriasis, all of the following (1-2): 1) if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg and member has moderate-to-severe disease -AND- 2) inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For oral ulcers associated with Behcet's Disease, inadequate response or intolerance to one systemic therapy for prevention of recurrent oral ulcers
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Palynziq

## Products Affected

- **PALYNZIQ SUBCUTANEOUS  
SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5  
ML, 20 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of phenylketonuria -AND- all of the following criteria (1-3): 1) Baseline Phe level greater than 600 micromoles/L, 2) Failure or intolerance to existing management (i.e. sapropterin dihydrochloride), 3) Has a prescription for epinephrine agent unless contraindicated.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in baseline pretreatment Phe levels -OR- blood Phe levels are less than or equal to 600 micromoles/L -OR- attestation that additional therapy with Palynziq is needed to allow adequate trial of maximum dose of 60mg per day for 16 weeks or member is not currently on 60mg per day dose.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Panretin

## Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions in patients with AIDS-related Kaposi Sarcoma (KS) who are not receiving systemic therapy for KS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Pemazyre

## Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable locally advanced cholangiocarcinoma or metastatic cholangiocarcinoma -AND- all of the following (1-2): 1) disease harbors FGFR2 fusion or other rearrangement as detected by an FDA-approved test 2) member has experienced therapeutic failure or intolerance to at least one prior therapy. Documentation of relapsed or refractory myeloid/lymphoid neoplasms -AND- disease harbors an FGFR1 rearrangement.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Pheburane

## Products Affected

- PHEBURANE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure or intolerance to generic sodium phenylbutyrate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Piqray

## Products Affected

- **PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR-positive, HER2-negative tumor status, 2) PIK3CA mutation positive as detected by an FDA-approved test, 3) disease progression on or after an endocrine-based regimen, 4) Concomitant therapy with fulvestrant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Pomalyst

## Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma, and combination use with dexamethasone, and previous trial of at least 2 therapies including lenalidomide and a proteasome inhibitor, and disease progression on or within 60 days of completion of the last therapy -OR- Documentation of AIDS-related Kaposi sarcoma (KS) after failure of highly active antiretroviral therapy (HAART) or in patients with KS who are HIV negative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Posaconazole Tablet

## Products Affected

- *posaconazole oral tablet, delayed release (dr/ec)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight greater than 40 kg. For invasive aspergillosis infection, trial/failure or contraindication to voriconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Prenatal Vitamins

## Products Affected

- **PRENATAL VITAMIN PLUS LOW IRON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of nutritional supplementation required in a female of child-bearing potential during pre-conception, pregnancy, or lactation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Prescription Drug Combo

## Products Affected

- *acetaminophen-codeine oral solution 120-12 mg/5 ml*
- *acetaminophen-codeine oral tablet*
- *alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *buprenorphine*
- *clonazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- *clonazepam oral tablet, disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg*
- **DIAZEPAM INTENSOL**
- *diazepam oral solution 5 mg/5 ml (1 mg/ml)*
- *diazepam oral tablet*
- **ENDOCET**
- *eszopiclone*
- *fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr*
- *hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg*
- *hydromorphone oral liquid*
- *hydromorphone oral tablet 2 mg, 4 mg, 8 mg*
- **LORAZEPAM INTENSOL**
- *lorazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- *methadone oral solution 10 mg/5 ml, 5 mg/5 ml*
- *methadone oral tablet 10 mg, 5 mg*
- *morphine concentrate oral solution*
- *morphine oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml)*
- *morphine oral tablet*
- *morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg*
- *oxycodone oral capsule*
- *oxycodone oral concentrate*
- *oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg*
- *oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg*
- *tramadol oral tablet 50 mg*
- *tramadol-acetaminophen*
- *zaleplon oral capsule 10 mg, 5 mg*
- *zolpidem oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
<b>Required Medical Information</b>	<p>For concomitant use of an opiate agonist and substance abuse therapy, documentation that the member has an acute pain condition (e.g. acute traumatic injury) in which treatment with other agents would cause insufficient pain control or if the member requires treatment for pain related to a terminal illness. For concomitant use of an opiate agonist, benzodiazepine, and a centrally acting skeletal muscle relaxant, documentation that the member has tried/failed at least 2 other skeletal muscle relaxants (e.g. methocarbamol, metaxalone), understanding these skeletal muscle relaxants are high-risk medications in geriatric patients, attestations that non-opiate therapies (e.g. NSAIDs) and non-benzodiazepine therapies (e.g. SSRI, SNRI) have been considered, AND attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For concomitant use of an opiate agonist and other opiate potentiators (e.g. gabapentinoids, sedative-hypnotics) attestation of an intent to monitor and address concomitant drug-drug interaction adverse events. For long acting (e.g. extended release) opioid medications, the following apply (1-5). 1)Pain is severe enough to require daily, around-the-clock, long-term opioid treatment. 2)Patient is not opioid naive. 3)Attestation that non-opiate alternative therapies have been explored (e.g. NSAIDs). 4)Attestation that controlled substance Rx history has been reviewed in the state Prescription Drug Monitoring Program. 5)Attestation of counseling on the potential adverse effects of opioid analgesics, including the risk of misuse, abuse, and addiction.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Opiate tx for pain+subs. abuse, approve opiate x 1mo. All other combos and dx approve x 12mo.
<b>Other Criteria</b>	<p>Opiate agonists will receive automatic approval if no recent claims for a substance abuse therapy (e.g. buprenorphine-naloxone) OR a benzodiazepine with a centrally acting skeletal muscle relaxant (e.g., carisoprodol) OR a gabapentinoid OR a sedative-hypnotic. Benzodiazepines (e.g. triazolam, alprazolam) will receive automatic approval if no recent claims for an opiate agonist with a centrally acting skeletal muscle relaxant (e.g. carisoprodol). Sedative-hypnotics (e.g. zolpidem) will receive automatic approval if no recent claims for an opiate agonist.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# Prevymis Oral Pellets

## Products Affected

- **PREVYMIS ORAL PELLETS IN PACKET**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis.
<b>Age Restrictions</b>	Deny if less than 6 months of age with HSCT. Deny if less than 12 years of age with kidney transplant.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	7 months
<b>Other Criteria</b>	Pending CMS Review
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Prolia

## Products Affected

- PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -And- For osteoporosis at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 (i.e. osteopenia) -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For osteoporosis and osteopenia, documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Provigil

## Products Affected

- *modafinil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography as established in accordance with ICSD or DSM V criteria acceptable for all indications. Diagnosis of fatigue associated with Multiple Sclerosis (MS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.
<b>Off Label Uses</b>	Fatigue associated with Multiple Sclerosis (MS)
<b>Part B Prerequisite</b>	No

# Pulmonary Arterial Hypertension

## Products Affected

- **ADEMPAS**
- **ALYQ**
- *ambrisentan*
- **OPSUMIT**
- **OPSYNVI**
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*
- **UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG**
- **UPTRAVI ORAL TABLETS,DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group. For sildenafil in pediatric individuals, an exception to RHC will be allowed when the risk of RHC outweighs the benefit -AND- prescriber attests alternative studies have been completed (i.e. CT, MRI or specified test ruling out other causes of pulmonary hypertension). For Adempas, additional diagnosis of CTEPH as documented by right heart catheterization and V/Q scan substantiating mPAP greater than 20 mmHg at rest and PWP less than or equal to 15 mmHg and documented presence of occlusive thrombi within the pulmonary arteries will be approved.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Pulmozyme

## Products Affected

- PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis -AND- Used in conjunction with standard therapies for management of cystic fibrosis to improve pulmonary function.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization, attestation of increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Qinlock

## Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced gastrointestinal stromal tumor -AND- Prior treatment with imatinib and 2 other kinase inhibitors.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Quinine

## Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment or prevention of leg cramps
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	Doses for duration greater than 10 days will not be approved
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# Qulipta

## Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, of which 8 or more are migraine days. The following criteria will apply (1-3). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). 3) Trial and failure or intolerance to one agent from 2 unique prophylactic migraine medication classes: e.g. Anti-epileptic drugs (e.g. topiramate), beta-blockers (e.g. propranolol), calcium-channel blockers (e.g. verapamil), tricyclic antidepressants (e.g. amitriptyline) -OR- contraindication to all prophylactic medication classes.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Radicava ORS

## Products Affected

- RADICAVA ORS STARTER KIT SUSP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of amyotrophic lateral sclerosis (ALS) -AND- Therapeutic failure, intolerance, or contraindication to riluzole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of experiencing stability or improvement of symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Ravicti

## Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	Urea cycle disorders due to N-acetylglutamate synthetase deficiency, Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of a urea cycle disorders (UCDs)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Regranex

## Products Affected

- REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of lower-extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond and have an adequate blood supply -AND- being used as an adjunct to standard ulcer care practices (e.g. sharp debridement, non-weight bearing regimen, infection control) -AND- attestation of a wound care plan.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	20 weeks
Other Criteria	For reauthorization, one of the following (1-2): 1) documentation of decrease in ulcer(s) size without complete ulcer(s) closure -OR- 2) documentation of new lower extremity diabetic neuropathic ulcer(s) that extends into the subcutaneous tissue or beyond with an adequate blood supply, attestation of being used as an adjunct to standard ulcer care practices, and attestation of a wound care plan.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Repatha

## Products Affected

- **REPATHA PUSHTRONEX**
- **REPATHA SURECLICK**
- **REPATHA SYRINGE**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
<b>Required Medical Information</b>	1.HoFH supported by genetic confirmation of two mutant alleles at LDLR, APOB, PCSK9, or LDLRAP1 gene OR untreated LDL-C greater than 400mg/dL or TC greater than 500mg/dl with cutaneous or tendon xanthoma before age 10 yrs or HeFH in both parents AND The member has a current LDL-C of greater than 135 mg/dL (if 17 years of age or younger) or greater than 100mg/dL (18 years of age or older) despite use of maximally tolerated statin or statin intolerance AND The member will continue to receive concurrent lipid-lowering therapies for the treatment of HoFH. 2.HeFH supported by presence of causal mutation of FH by genetic testing OR untreated LDL-C greater than or equal to 190 mg/dL or untreated LDL-C greater than or equal to 160 mg/dL before 20 years of age with physical signs of FH (e.g. xanthomas, xanthelasma) OR diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria with score greater than 8 points, or definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool, AND LDL-C greater than 100 mg/dL if 18 and older or LDL-C greater than 130 if 17 and younger despite use of maximally tolerated statin or statin intolerance. If 17 and younger will continue to receive concurrent lipid-lowering therapies. 3.Hypercholesterolemia ASCVD or Primary Hyperlipidemia AND LDL-C greater than 70 mg/dL despite use of maximally tolerated statin or statin intolerance
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	For reauthorization, documentation showing an LDL-C reduction on Repatha therapy from baseline must be provided. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Retevmo

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG
- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is classified as RET gene fusion as detected by an FDA approved test. Documentation of advanced or metastatic medullary thyroid cancer -AND- disease is classified as RET mutation as detected by an FDA approved test. Documentation of advanced or metastatic thyroid cancer -AND- all of the following (1-2): 1) disease is classified as RET gene fusion as detected by an FDA approved test 2) if radioactive iodine is appropriate for the member, the member is radioactive iodine-refractory. Documentation of locally advanced or metastatic solid tumor(s) -AND- disease harbors a RET gene fusion, as detected by an FDA-approved test -AND- one of the following (1-2): 1) the member has no satisfactory alternative treatments 2) the member's tumors have progressed following prior systemic treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Revlimid

## Products Affected

- *lenalidomide*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Documentation of chronic lymphocytic leukemia outside of a controlled clinical trial
<b>Required Medical Information</b>	Diagnosis of multiple myeloma in combination with dexamethasone -OR- diagnosis of multiple myeloma, as maintenance following autologous hematopoietic stem cell transplant (auto-HSCT) -OR- diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities -OR- diagnosis of mantle cell lymphoma (MCL) in which disease has relapsed or progressed after two prior therapies, one of which included bortezomib -OR- diagnosis of follicular lymphoma in combination with a rituximab product -OR- diagnosis of marginal zone lymphoma in combination with a rituximab product after previous treatment.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Revuforj

## Products Affected

- **REVUFORJ ORAL TABLET 110 MG, 160 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute leukemia with a lysine methyltransferase 2A gene (KMT2A) translocation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rexulti

## Products Affected

- REXULTI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRI, TCA, MAOI). For schizophrenia, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Rezdiffra

## Products Affected

- REZDIFFRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in conjunction with diet and exercise. Diagnosis of NASH confirmed by a liver biopsy or non-invasive tests (NITs) performed within the previous 6 months indicating F2 or F3 fibrosis - AND- member does not have any evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of continued use in conjunction with diet and exercise -AND- member has experienced stabilization of fibrosis as demonstrated by NIT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rezlidhia

## Products Affected

- **REZLIDHIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rinvoq

## Products Affected

- **RINVOQ ORAL TABLET EXTENDED  
RELEASE 24 HR 15 MG, 30 MG, 45 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate) or all non-biologic DMARDs are contraindicated. For moderate to severe refractory atopic dermatitis whose disease is not adequately controlled with other systemic drug products, documentation of one of the following (1 or 2): 1) trial & failure, or intolerance to at least one topical corticosteroid -OR- topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) 2) The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For Non-radiographic Axial Spondyloarthritis, trial & failure or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or contraindication to all. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For diagnoses in which tumor necrosis factor (TNF) blockers are also indicated (e.g., Rheumatoid Arthritis, Psoriatic Arthritis), the member has experienced therapeutic failure or intolerance to at least 1 TNF blocker.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Rinvoq LQ

## Products Affected

- RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	For PsA, deny if 18 years of age or older
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rivfloza

## Products Affected

- **RIVFLOZA SUBCUTANEOUS SOLUTION**
- **RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3): 1) genetically confirmed diagnosis of primary hyperoxaluria type 1 (PH1), 2) relatively preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2), and 3) at least two elevated urinary oxalate levels greater than 1.5 times the upper reference limit.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, continued preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2) -AND- reduction in urinary oxalate levels from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rozlytrek

## Products Affected

- **ROZLYTREK ORAL CAPSULE 100 MG, 200 MG**
- **ROZLYTREK ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, the tumor status is ROS1-positive. For solid tumors with NTRK gene fusion without a known acquired resistance mutation, the tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or the tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Rubraca

## Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) Disease harbors a deleterious BRCA mutation (germline or somatic) 2) member is in complete or partial response to platinum-based chemotherapy. For metastatic castration-resistant prostate cancer, all of the following (1-3): 1) disease harbors a deleterious BRCA mutation (germline and/or somatic) 2) member has been treated with androgen receptor-directed therapy and taxane-based chemotherapy 3) member is concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Rydapt

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Use as single agent induction therapy for AML
Required Medical Information	Documentation of diagnosis. For a new diagnosis of acute myeloid leukemia, member is using in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation chemotherapy regimens and member is FLT3 mutation positive as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Sabril

## Products Affected

- *vigabatrin*
- **VIGADRONE**
- **VIGPODER**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of refractory complex partial seizures -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance to at least two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, tiagabine) -OR- documentation of use as monotherapy in treatment of infantile spasms
Age Restrictions	Deny if less than 2 years of age in treatment of refractory complex partial seizures -OR- if less than 1 month old and greater than 2 years of age in treatment of infantile spasms
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Samsca

## Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic hypervolemic or euvolemic hyponatremia evidenced by (1. or 2.): 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	Doses must be initiated in the hospital setting to closely monitor serum sodium. Treatment should be limited to 30 days to minimize risk of liver injury. For reauthorization, treatment is for a new episode of a clinically significant euvolemic or hypervolemic hyponatremia -AND- on of the following (1. or 2.) 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Saphris

## Products Affected

- *asenapine maleate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Savella

## Products Affected

- SAVELLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to duloxetine
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Scemblix

## Products Affected

- **SCSEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-3): 1) the member is newly-diagnosed, 2) member has been previously treated for Ph+ CML in chronic phase, 3) disease is positive for the T3151 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Secuado

## Products Affected

- SECUADO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure of generic asenapine sublingual tablets -AND- trial/failure or intolerance to 1 of the following or all are contraindicated (1-3): 1) olanzapine, 2) quetiapine, 3) risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Simponi

## Products Affected

- **SIMPONI SUBCUTANEOUS PEN INJECTOR 100 MG/ML, 50 MG/0.5 ML**
- **SIMPONI SUBCUTANEOUS SYRINGE 100 MG/ML, 50 MG/0.5 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Simponi 50mg: For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. leflunomide) and Simponi will be used in combination with methotrexate. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). Diagnosis of psoriatic arthritis Simponi 100mg: Diagnosis of moderate to severe ulcerative colitis.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Actemra SC, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For ulcerative colitis, patients must have therapeutic failure or intolerance to the preferred products: a preferred adalimumab product, Stelara SC, Rinvoq, and Xeljanz/Xeljanz XR. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For ulcerative colitis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Sirturo

## Products Affected

- **SIRTURO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of extensively drug resistant tuberculosis, pre-extensively drug-resistant tuberculosis, treatment-intolerant tuberculosis, or nonresponsive multidrug-resistant tuberculosis -AND- Therapeutic failure, contraindication, or intolerance to 1 of the following: isoniazid, rifampin, rifabutin, or rifapentine -AND- Used in combination with at least 3 other drugs for tuberculosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 weeks
Other Criteria	For reauthorization, additional therapy required due to doses of the regimen being missed for safety reasons
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Skyclarys

## Products Affected

- SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Friedreichs ataxia confirmed by genetic testing (i.e., FXN gene mutation).
Age Restrictions	Deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Skyrizi

## Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML
- SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohns disease, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC. For moderate to severe ulcerative colitis, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Sohonos

## Products Affected

- SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 2.5 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	Deny if female and less than 8 years of age -OR- if male and less than 10 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in volume of new heterotopic ossification from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Solaraze

## Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure, intolerance, or contraindication to one of the following (1 or 2): 1) topical fluorouracil solution or fluorouracil 5% cream 2) topical imiquimod 5% cream
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of 30 day washout period since optimal therapeutic effect may not be evident until 30 days following cessation of therapy AND attestation of previous response to diclofenac sodium 3% topical gel therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Somavert

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- inadequate or partial response to surgery or radiotherapy or not a candidate for surgery or radiotherapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Sprycel

## Products Affected

- *dasatinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adults with Ph+ chronic myeloid leukemia, the member is newly diagnosed in the chronic phase -OR- the member is in chronic, accelerated, or myeloid or lymphoid blast phase and has resistance or intolerance to prior therapy including imatinib. For adults with Ph+ acute lymphocytic leukemia, member has had resistance or intolerance to prior therapy. For pediatric patients with Ph+ CML, the member is in the chronic phase. For pediatric patients with Ph+ acute lymphoblastic leukemia, the member is newly diagnosed and will be using in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Stelara

## Products Affected

- **STELARA SUBCUTANEOUS SOLUTION**
- **STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease, attestation of clinical response or remission following single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC. For Ulcerative colitis, attestation of clinical response or remission following single induction dose of IV Stelara within 2 months of initiating therapy with Stelara SC
<b>Age Restrictions</b>	Deny if less than 18 years of age for Crohn's Disease and Ulcerative Colitis or less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Must follow recommended dosing guidelines based upon weight. Induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Steroidogenesis Inhibitors

## Products Affected

- RECORLEV
- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- patient is not a candidate for pituitary surgery or surgery has not been curative
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in urinary free cortisol levels from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Stivarga

## Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer and trial of a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy, AND an anti-VEGF therapy AND if RAS wild-type, an anti-EGFR therapy -OR- documentation of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with both imatinib and sunitinib -OR- documentation of hepatocellular cancer AND previous treatment with sorafenib
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Sutent

## Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For gastrointestinal stromal tumor (GIST), the member has experienced therapeutic failure, intolerance, or contraindication to imatinib. For a high risk of recurrent renal cell carcinoma, member has had a nephrectomy and sunitinib is to be used as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Symdeko

## Products Affected

- **SYMDEKO**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of cystic fibrosis (CF) in patients who have either the homozygous F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to tezacaftor/ivacaftor based on clinical and/or in vitro assay (e.g. E56K, R117C, A455E)
<b>Age Restrictions</b>	Deny if less than 6 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Sympazan

## Products Affected

- SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy -AND- unable to tolerate generic clobazam
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Synarel

## Products Affected

- SYNAREL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For central precocious puberty (CPP), advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L. For female with endometriosis, attestation of not pregnant if of childbearing age -AND- Therapeutic failure, contraindication or intolerance to 2 of the following standard of care treatments: NSAIDs, combination hormonal contraceptive, progestin (i.e. medroxyprogesterone injection), GnRH agonist (i.e. Leuprolide) or danazol
<b>Age Restrictions</b>	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty. Deny if less than 18 years of age for endometriosis.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Endometriosis: 6 months, CPP: 6 months initial authorization, 12 months reauthorization
<b>Other Criteria</b>	For reauthorization for CPP, attestation of pubertal development slowing from baseline.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Tabrecta

## Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with MET exon 14 skipping mutation as detected by an FDA approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Tafinlar

## Products Affected

- TAFINLAR ORAL CAPSULE
- TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For dabrafenib monotherapy, documentation of unresectable or metastatic melanoma -AND- documentation of a BRAF V600E mutation. For use in combination with trametinib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with trametinib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation. For use in combination with trametinib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Tafinlar tablets for oral suspension, attestation of inability to swallow Tafinlar (dabrafenib) capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tagrisso

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) and one of the following (1-5): 1) Adjuvant therapy after tumor resection -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, 2) Locally advanced disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations -AND- used in combination with pemetrexed and platinum-based chemotherapy, 3) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations -AND- used with or without combination therapy of pemetrexed and platinum-based chemotherapy, 4) Metastatic disease -AND- disease harbors EGFR T790M mutations -AND- has progressed on or after EGFR TKI therapy, 5) Locally advanced, unresectable (stage III) disease -AND- disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy -AND- disease harbors EGFR exon 19 deletions or EGFR exon 21 L858R mutations, as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Taltz

## Products Affected

- **TALTZ AUTOINJECTOR**
- **TALTZ SYRINGE SUBCUTANEOUS SYRINGE 20 MG/0.25 ML, 40 MG/0.5 ML, 80 MG/ML**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs.
<b>Age Restrictions</b>	Deny if less than 18 years of age for Psoriatic Arthritis, Ankylosing Spondylitis and non-radiographic axial spondyloarthritis or less than 6 years of age for Plaque Psoriasis
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>For psoriatic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cosentyx, Enbrel, a preferred adalimumab product, Xeljanz/Xeljanz XR, Otezla, Stelara SC, Rinvoq, and Skyrizi SC. For plaque psoriasis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Cosentyx, Otezla, Stelara SC, Skyrizi SC, and Enbrel. For ankylosing spondylitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Enbrel, Cosentyx, Rinvoq, and Xeljanz/Xeljanz XR. For non-radiographic axial spondyloarthritis patients must have therapeutic failure or intolerance to 2 of the following preferred products: Cimzia, Rinvoq, Cosentyx. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606. For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Talzenna

## Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of deleterious or suspected deleterious gBRCAm, HER2-negative locally advanced or metastatic breast cancer as a single agent - OR- Documentation of HRR gene-mutated metastatic castration-resistant prostate cancer in combination with enzalutamide
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Targretin

## Products Affected

- *bexarotene oral*
- *bexarotene topical*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations of cutaneous T-cell lymphoma in patients who are refractory to at least one prior systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tascenso ODT

## Products Affected

- TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Inability to swallow capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tasigna

## Products Affected

- TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adult patients with Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib - OR- member is newly diagnosed in the chronic phase. For pediatric patients, one of the following (1-2): 1) member has chronic phase or accelerated phase Ph+ CML and is resistant or intolerant to prior tyrosine kinase inhibitor therapy 2) member is newly diagnosed with Ph+ CML in the chronic phase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Tasimelteon

## Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tazorac

## Products Affected

- *tazarotene topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 of the following (A or B). A) Documentation of plaque psoriasis -AND- trial and failure or intolerance to at least one topical corticosteroid (e.g. fluocinonide, mometasone, triamcinolone, betamethasone). B) Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tazverik

## Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic epithelioid sarcoma - AND- Disease is not eligible for complete resection. Documentation of relapsed or refractory follicular lymphoma -AND-Tumors are EZH2 mutation positive, as detected by FDA approved test, in a member that has received at least 2 prior systemic therapies -OR- Prescriber attests there are no satisfactory alternative treatment options.
Age Restrictions	For epithelioid sarcoma, deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tecfidera

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
<b>Required Medical Information</b>	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	5 years
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Tepmetko

## Products Affected

- **TEPMETKO**

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of metastatic non-small cell lung cancer with a MET exon 14 skipping alteration
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Testosterone (androgens)

## Products Affected

- *testosterone cypionate*
- *testosterone enanthate*
- *testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)*
- *testosterone transdermal gel in packet*
- *testosterone transdermal solution in metered pump w/app*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Documentation of primary or secondary hypogonadism in males with testicular failure due to cryptorchidism, bilateral torsions, orchitis, vanishing testis syndrome, orchidectomy, Klinefelter's syndrome, chemotherapy, radiation or toxic damage -OR- documentation of primary or secondary hypogonadism in males with multiple symptoms of hypogonadism including at least one of the following specific symptoms: height loss due to vertebral fractures, low trauma fractures, low bone density, incomplete or delayed sexual development, breast discomfort, loss of axillar and/or pubic body hair, hot flushes -OR- documentation of HIV infection in men with weight loss -OR- documentation of chronic steroid treatment in men. In all previously noted indications, members must also have documented low total testosterone level below the normal range for the laboratory -OR- a total testosterone level near the lower limit of the normal range with a low free testosterone level which is less than normal based upon the laboratory reference range -OR- the member is not producing any testosterone. Additional approvable indications include female patients with metastatic breast cancer (testosterone enanthate only), primary or secondary hypogonadism in males with testicular failure due to double orchidectomy, and delayed puberty in males (testosterone enanthate only).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications, Some Medically-accepted Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	HIV Wasting
<b>Part B Prerequisite</b>	No

# Thalomid

## Products Affected

- **THALOMID ORAL CAPSULE 100 MG, 50 MG**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Use as monotherapy for ENL treatment in the presence of moderate to severe neuritis
<b>Required Medical Information</b>	Documentation of multiple myeloma in combination with dexamethasone -OR- documentation for use in the treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) -OR- documentation of therapy for prevention and suppression of the cutaneous manifestations of ENL recurrence
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# Thrombopoiesis Stimulating Agents

## Products Affected

- **PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG**
- **PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of persistent or chronic immune idiopathic thrombocytopenia purpura and trial and failure of corticosteroid or immunoglobulin therapy or splenectomy -OR- documentation of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy -OR- severe aplastic anemia who have had an insufficient response to immunosuppressive therapy -OR- documentation of first line treatment for severe aplastic anemia and used in combination with at least two immunosuppressive therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Platelet count to be provided
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tibsovo

## Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is IDH1 mutation positive as detected by an FDA-approved test. For newly-diagnosed acute myeloid leukemia, member is using as monotherapy or in combination with azacitidine -AND- member meets one of the following (1-5): 1) age is greater than or equal to 75 years of age 2) severe cardiac or pulmonary comorbidity 3) reduced renal function 4) hepatic impairment 5) or prescriber attestation that member is not a candidate for intensive induction therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Topical Lidocaine

## Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical ointment*
- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tretinoin

## Products Affected

- *tretinoin topical cream*
- *tretinoin topical gel 0.01 %, 0.025 %*

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two generic topical non-retinoid acne medications (e.g. clindamycin, sulfacetamide, erythromycin)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Trikafta

## Products Affected

- **TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL**
- **TRIKAFTA ORAL TABLETS, SEQUENTIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have at least one F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to elexacaftor/tezacaftor/ivacaftor based on in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Trintellix

## Products Affected

- **TRINTELLIX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Truqap

## Products Affected

- TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) HR mutation status, HER2 mutation status, and PIK3CA/AKT1/PTEN status 2) concomitant therapy with fulvestrant 3) disease progression on at least one endocrine-based regimen in the metastatic setting -OR- recurrence on or within 12 months of completing adjuvant therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tukysa

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced unresectable HER2-positive breast cancer or metastatic HER2-positive breast cancer, member will be using in combination with trastuzumab and capecitabine -AND- member has received one or more prior anti-HER2 based regimens in the metastatic setting. For RAS wild-type HER2-positive unresectable or metastatic colorectal cancer, member will be using in combination with trastuzumab -AND- member has experienced disease progression following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Turalio

## Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor associated with severe morbidity and functional limitations -AND- patient is not amenable to improvement with surgery or not a candidate for surgery
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tykerb

## Products Affected

- *lapatinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced HER2-positive, or metastatic HER2-positive breast cancer, the member has received prior therapy with an anthracycline, a taxane, and trastuzumab -AND- will be using in combination with capecitabine. For HR+, metastatic breast cancer, the member is post-menopausal -AND- the member's cancer over expresses the HER2 receptor -AND- the member will be using lapatinib in combination with letrozole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Tymlos

## Products Affected

- TYMLOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Treatment duration greater than 24 months.
<b>Required Medical Information</b>	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 months
<b>Other Criteria</b>	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Additional documentation of trial/failure, intolerance or contraindication to preferred parathyroid hormone analog teriparatide. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Tymlos will be limited to a coverage duration of 24 months.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Ubrelvy

## Products Affected

- **UBRELVY ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of acute treatment of migraine with or without aura -AND- trial and failure of one generic triptan.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	Patients must have therapeutic failure, intolerance, or contraindication to Nurtec ODT. For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Uloric

## Products Affected

- *febuxostat*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic management of hyperuricemia due to gout -And- trial/failure, intolerance or contraindication to allopurinol.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Valchlor

## Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received at least one prior skin-directed therapy (e.g. topical corticosteroids, topical chemotherapy, local radiation and topical retinoids).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Valtoco

## Products Affected

- VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	Deny if less than 6 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vancomycin

---

## Products Affected

- *vancomycin oral capsule 125 mg, 250 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Vanflyta

## Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is FLT3-ITD-positive as detected by an FDA-approved test -AND- one of the following (1-3): 1) member is receiving induction therapy and is using Vanflyta in combination with standard cytarabine and anthracycline induction therapy 2) member is receiving consolidation therapy and is using Vanflyta in combination with standard cytarabine consolidation therapy 3) member is receiving maintenance therapy and is using Vanflyta as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Venclexta

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For newly-diagnosed AML, member is using in combination with azacitidine, decitabine, cytarabine -AND- age greater than or equal to 75 years or presence of at least one comorbidity that precludes use of intensive induction chemotherapy (i.e. severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) is required.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Verquvo

## Products Affected

- VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of heart failure (NYHA Class II to IV) -AND- Left ventricular ejection fraction less than 45% -AND- Hospitalization for heart failure or received outpatient IV diuretics for heart failure -AND- Used in combination with a angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker or Entresto -AND- Used in combination with bisoprolol, carvedilol IR/ER or metoprolol succinate ER.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Verzenio

## Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative. For early breast cancer that is at high risk of recurrence and is node-positive, all of the following (1-2): 1) used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) 2) used as adjuvant treatment. For advanced or metastatic breast cancer, used as initial endocrine-based therapy and used in combination with an aromatase inhibitor -OR- used after documented disease progression following endocrine therapy and used in combination with fulvestrant -OR- used after documented disease progression and used following endocrine therapy and prior chemotherapy in the metastatic setting and will be used as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Viberzi

## Products Affected

- VIBERZI

PA Criteria	Criteria Details
Exclusion Criteria	Severe (Child-Pugh C) hepatic impairment
Required Medical Information	Documentation of diarrhea predominant, irritable bowel syndrome (IBS-D) -AND- trial/failure or intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Viibryd

## Products Affected

- *vilazodone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vioice

## Products Affected

- **VIJOICE ORAL GRANULES IN                      MG  
PACKET**
- **VIJOICE ORAL TABLET 125 MG, 250  
MG/DAY (200 MG X1-50 MG X1), 50**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe manifestations of PIK3CA Related Overgrowth Spectrum (PROS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vitrakvi

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation -AND- Tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Vivjoa

## Products Affected

- VIVJOA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of recurrent vulvovaginal candidiasis defined as at least 3 episodes of vulvovaginal candidiasis in less than one year -AND Documentation the member is NOT of reproductive potential defined as postmenopausal or another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy) -AND- the member has experienced therapeutic failure, contraindication, or intolerance to a six-month maintenance course of oral fluconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vizimpro

## Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer -AND- one of the following (1 or 2): 1. Epidermal growth factor (EGFR) exon 19 deletions - OR- 2. Epidermal growth factor receptor (EGFR) exon 21 L858R substitution mutations.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vonjo

## Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of a platelet count of less than $50 \times 10^9/L$ .
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Voranigo

## Products Affected

- **VORANIGO ORAL TABLET 10 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) grade 2 astrocytoma or grade 2 oligodendroglioma, 2) disease harbors a susceptible isocitrate dehydrogenase (IDH)-1 or IDH-2 mutation, 3) will be used following surgery including biopsy, sub-total resection, or gross total resection.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Voriconazole

## Products Affected

- *voriconazole intravenous*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary cannot take oral voriconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of continued indicators of active disease (e.g. histopathology, positive cultures) is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vosevi

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Votrient

## Products Affected

- *pazopanib*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of adipocytic soft tissue sarcoma or gastrointestinal stromal tumor
Required Medical Information	Documentation of diagnosis. For advanced soft-tissue sarcoma, trial/failure of at least one prior chemotherapy regimen.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Vraylar

## Products Affected

- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For a diagnosis of schizophrenia and bipolar I disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Vumerity

## Products Affected

- VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Welireg

## Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For von Hippel Lindau (VHL) syndrome, one of the following diagnoses not requiring immediate surgery (1, 2, or 3): 1) Renal cell carcinoma (RCC) 2) CNS hemangioblastoma 3) Pancreatic neuroendocrine tumor. For advanced RCC, prior treatment with a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) -AND- a programmed death receptor-1 (PD-1) or a programmed death-ligand (PD-L1) inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xalkori

## Products Affected

- **XALKORI ORAL CAPSULE**
- **XALKORI ORAL PELLETT 150 MG, 20 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer (NSCLC), disease is ALK-positive or ROS1-positive. For relapsed or refractory anaplastic large cell lymphoma (ALCL), disease is ALK-positive. For unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT), disease is ALK-positive.
Age Restrictions	Deny if less than 18 years of age for NSCLC. Deny if less than 1 year of age or greater than 21 years of age for ALCL.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For crizotinib oral pellets and NSCLC, inability to swallow capsules is required. For crizotinib oral pellets and ALCL / IMT, inability to swallow oral capsules -OR- body surface area less than 1.34 m <sup>2</sup> is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xcopri

## Products Affected

- **XCOPRI**
- **XCOPRI MAINTENANCE PACK**
- **XCOPRI TITRATION PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Therapeutic failure, intolerance or contraindication to 1 other anti-epileptic drug (e.g. carbamazepine, levetiracetam)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xdemvy

## Products Affected

- XDEMVEY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis confirmed by identification of Demodex infection via microscopic examination of pulled eyelashes -OR- identification of collarettes via slit-lamp evaluation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xeljanz

## Products Affected

- XELJANZ ORAL TABLET
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severely active rheumatoid arthritis and an inadequate response or intolerance to methotrexate. Xeljanz immediate release for juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID).
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis, psoriatic arthritis, ulcerative colitis, ankylosing spondylitis. For Xeljanz regular release tablet, deny if less than 2 years of age for juvenile idiopathic arthritis
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced inadequate response or intolerance to at least 1 tumor necrosis factor (TNF) blocker. Doses greater than 10 mg per day for Xeljanz and 11 mg per day for Xeljanz XR will not be approved for rheumatoid arthritis and psoriatic arthritis. Doses greater than 20mg per day for Xeljanz and 22 mg per day for Xeljanz XR will not be approved for ulcerative colitis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xeljanz Solution

## Products Affected

- XELJANZ ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of juvenile idiopathic arthritis -AND- Inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xenazine

## Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary is not actively suicidal
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	In patients with comorbid depression, attestation of adequate treatment for depression is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Xermelo

## Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of carcinoid syndrome diarrhea AND used in combination with a somatostatin analog AND trial and failure of somatostatin analog monotherapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in average number of daily bowel movements -AND- the member will continue to use in combination with a somatostatin analog.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xgeva

## Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For hypercalcemia of malignancy, refractory to bisphosphonates. For giant cell tumor of bone, unresectable or surgical resection is likely to result in severe morbidity -AND- one of the following (1. or 2.)- 1.) the member is 18 years old or older -OR- 2.) the member is a skeletally mature adolescent (e.g. has at least one mature long bone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xifaxan

## Products Affected

- XIFAXAN ORAL TABLET 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 or 2. 1) Diagnosis of hepatic encephalopathy AND trial/failure, intolerance, or contraindication to lactulose. 2) Diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D) AND trial/failure, intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Hepatic encephalopathy: 1 year. IBS-D: 14 days.
Other Criteria	No more than three courses of rifaximin for the treatment of IBS-D will be approved per lifetime.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xolair

## Products Affected

- XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	<p>Documentation of the following (1-2): 1) Chronic Spontaneous Urticaria, 2) trial/failure or intolerance of a second-generation non-sedating H1 antihistamine at the maximum recommended doses (e.g. cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine). -OR- Documentation of the following (3-9): 3) moderate to severe persistent asthma, 4) a positive skin test or in vitro reactivity to a perennial aeroallergen, 5) Baseline IgE titer greater than or equal to 30 IU/mL, 6) documented pretreatment FEV1 less than 80% predicted in adults or less than 90% predicted in children and adolescents or FEV1 reversibility of at least 12% and 200mL after albuterol administration, 7) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos, 8) inadequately controlled symptoms despite a 3-month trial of both of the following (a-b) a) medium-dose inhaled corticosteroid or systemic steroid b) a long-acting beta-agonist or leukotriene antagonist, 9) currently on 1 of the following (c, d, e): c) a long-acting beta2-agonist, d) leukotriene modifier, or e) theophylline. -OR- Documentation of the following (10-11): 10) chronic rhinosinusitis with nasal polyps (CRSwNP), 11) will use concomitantly with nasal corticosteroid maintenance treatment, -OR- Documentation of the following (12-17): 12) IgE mediated food allergy, 13) diagnosis confirmed by skin prick test or food-specific antibodies, 14) previous allergic reaction to food, 15) using for the reduction of allergic reactions (type 1), including anaphylaxis, 16) will be used in conjunction with food allergen avoidance, 17) member has a documented prescription for epinephrine.</p>
Age Restrictions	
Prescriber Restrictions	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CSU reauthorization, improved CSU symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For IgE-mediated food allergy reauthorization, member requires continuation of therapy and will continue food allergen avoidance.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Xolremdi

## Products Affected

- **XOLREMDI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of WHIM syndrome (warts, hypogammaglobulinemia, infections, myelokathexis).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in incidence of infections is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xospata

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is FLT3 mutation-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xpovio

## Products Affected

- **XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use in combination with dexamethasone for relapse or refractory multiple myeloma with failure, intolerance or contraindication to 5 therapies (e.g. bortezomib, carfilzomib, lenalidomide, pomalidomide and daratumumab) -OR- Documentation of use in combination with both bortezomib and dexamethasone for multiple myeloma after receiving 1 prior multiple myeloma therapy -OR- Documentation of relapsing or refractory diffuse large B-cell lymphoma with failure, intolerance or contraindication to at least 2 lines of systemic therapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Xtandi

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For castration-resistant prostate cancer and metastatic castration sensitive-prostate cancer, the member is using in combination with a GnRH analog or the member has had a bilateral orchiectomy. For non-metastatic castration-sensitive prostate cancer, the member has biochemical recurrence at high risk for metastasis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Xyrem

## Products Affected

- *sodium oxybate*
- **XYREM**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If no diagnosis of cataplexy- trial and failure, intolerance, or contraindication to generic modafinil AND a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Yonsa

## Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with methylprednisolone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Yorvipath

## Products Affected

- **YORVIPATH SUBCUTANEOUS PEN  
INJECTOR 168 MCG/0.56 ML, 294  
MCG/0.98 ML, 420 MCG/1.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hypoparathyroidism -AND- albumin-corrected serum calcium greater than or equal to 7.8 mg/dL -AND- trial and failure of calcium -AND- trial and failure of an active form of vitamin D (e.g. calcitriol, alfacalcidol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of an improvement in total serum calcium from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zavesca

## Products Affected

- *miglustat*
- YARGESA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of mild to moderate type 1 Gaucher disease confirmed by the following (A. or B.) A. (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm <sup>3</sup> -OR- 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B. Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation enzyme replacement therapy (e.g. Cerezyme, Elelyso, or VPRIV) is not a therapeutic option
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zavzpret

## Products Affected

- ZAVZPRET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- Therapeutic failure, contraindication or intolerance to one generic triptan - AND- Inability to swallow capsules/tablets
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zejula

## Products Affected

- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer, member is in complete or partial response to first-line platinum-based therapy. For recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, all of the following (1-2): 1) disease harbors a deleterious or suspected deleterious germline BRCA mutation 2) member is in a complete or partial response to platinum-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zelboraf

## Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma and use in combination with cobimetinib, member has a BRAF V600E or V600K mutation. For unresectable or metastatic melanoma and use as monotherapy, member has a BRAF V600E mutation. For Erdheim-Chester Disease and use as monotherapy, member has a BRAF V600 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# Zeposia

## Products Affected

- ZEPOSIA
- ZEPOSIA STARTER KIT (28-DAY)
- ZEPOSIA STARTER PACK (7-DAY)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use of Zeposia and other disease modifying agents such as interferons, Copaxone, Tysabri.
<b>Required Medical Information</b>	Documentation of diagnosis. For moderate to severe active ulcerative colitis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate, cyclosporine).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Ulcerative Colitis: 12 months, Multiple Sclerosis: 24 months
<b>Other Criteria</b>	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq, Xeljanz/Xeljanz XR and Stelara SC. Please Note: preferred adalimumab products include Humira with NDC starting with 00074, Cyltezo with NDC starting 00597 and Yuflyma with NDC starting 72606.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Zolinza

## Products Affected

- ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations in patients with cutaneous T-cell lymphoma (CTCL) who have progressive, persistent, or recurrent disease on or following 2 systemic therapies. Systemic therapies include bexarotene, interferon alpha, extracorporeal photochemotherapy, PUVA, single agent or combination chemotherapies (e.g. cyclophosphamide, vinblastine, romidepsin)
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zonisade

## Products Affected

- ZONISADE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Documentation of adjunctive therapy -AND- Inability to swallow capsules -AND- Therapeutic failure/intolerance to 2 or contraindication to all of the following (1-6): 1) generic carbamazepine suspension/chewable tablet/extended-release capsule, 2) generic gabapentin capsules/solution, 3) generic lacosamide solution, 4) generic levetiracetam solution, 5) generic oxcarbazepine suspension, 6) generic pregabalin capsules/solution.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Ztalmy

## Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures associated with CDKL5 deficiency confirmed by genetic testing -AND- therapeutic failure or intolerance to 2 previous antiepileptic therapies
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zurzuvae

## Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of moderate to severe postpartum depression -AND- less than or equal to 12 months postpartum.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zydelig

## Products Affected

- ZYDELIG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	First line treatment. Combination use with benadmustine and/or rituximab for the treatment of FL.
<b>Required Medical Information</b>	Documentation of relapsed chronic lymphocytic leukemia -AND- all of the following (1-2): 1) will be used in combination with rituximab 2) use of rituximab alone would be appropriate due to other due to other comorbidities.
<b>Age Restrictions</b>	Deny if less than 18 years of age
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# Zykadia

## Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- ALK mutations, as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# Zytiga

## Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with prednisone.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No





## Index of Drugs

<i>abiraterone oral tablet 250 mg, 500 mg</i> .....	347	<b>BESREMI</b> .....	27
<i>acetaminophen-codeine oral solution 120-12 mg/5 ml</i> .....	215	<b>BETASERON SUBCUTANEOUS KIT</b> .....	127
<i>acetaminophen-codeine oral tablet</i> .....	215	<i>bexarotene oral</i> .....	273
<i>acitretin</i> .....	1	<i>bexarotene topical</i> .....	273
<b>ACTEMRA ACTPEN</b> .....	2	<b>BIVIGAM</b> .....	115
<b>ACTEMRA SUBCUTANEOUS</b> .....	2	<b>BOSULIF ORAL CAPSULE 100 MG, 50 MG</b> .....	28
<b>ACTIMMUNE</b> .....	3	<b>BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG</b> .....	28
<b>ADBRY</b> .....	4	<b>BRAFTOVI</b> .....	29
<b>ADEMPAS</b> .....	222	<b>BRUKINSA</b> .....	30
<b>AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML, 70 MG/ML</b> .....	8	<i>buprenorphine</i> .....	215
<b>AJOVY AUTOINJECTOR</b> .....	9	<b>CABLIVI INJECTION KIT</b> .....	32
<b>AJOVY SYRINGE</b> .....	9	<b>CABOMETYX</b> .....	33
<b>AKEEGA</b> .....	10	<b>CALQUENCE</b> .....	34
<b>ALCOHOL PADS</b> .....	125	<b>CALQUENCE (ACALABRUTINIB MAL)</b> .....	34
<b>ALECENSA</b> .....	11	<b>CAMZYOS</b> .....	35
<i>alosetron oral tablet 0.5 mg, 1 mg</i> .....	160	<b>CAPLYTA</b> .....	36
<i>alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg</i> .....	215	<b>CAPRELSA ORAL TABLET 100 MG, 300 MG</b> .....	37
<b>ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG</b> .....	13	<i>carglumic acid</i> .....	38
<b>ALUNBRIG ORAL TABLETS,DOSE PACK</b> .....	13	<b>CAYSTON</b> .....	39
<b>ALYQ</b> .....	222	<b>CERDELGA</b> .....	40
<i>ambrisentan</i> .....	222	<b>CIBINQO</b> .....	43
<i>amitriptyline</i> .....	107	<b>CIMZIA POWDER FOR RECONST</b> .....	44
<b>ARIKAYCE</b> .....	15	<b>CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)</b> .....	44
<i>aripiprazole oral solution</i> .....	17	<b>CINRYZE</b> .....	46
<i>aripiprazole oral tablet,disintegrating</i> .....	17	<i>clobazam oral suspension</i> .....	199
<i>armodafinil</i> .....	190	<i>clobazam oral tablet</i> .....	199
<i>asenapine maleate</i> .....	247	<i>clomipramine</i> .....	107
<b>AUGTYRO ORAL CAPSULE 160 MG, 40 MG</b> .....	19	<i>clonazepam oral tablet 0.5 mg, 1 mg, 2 mg</i> .....	215
<b>AUVELITY</b> .....	20	<i>clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg</i> .....	215
<b>AVONEX INTRAMUSCULAR PEN INJECTOR KIT</b> .....	127	<i>clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg</i> .....	215
<b>AVONEX INTRAMUSCULAR SYRINGE KIT</b> .....	127	<b>COBENFY</b> .....	48
<b>AYVAKIT</b> .....	21	<b>COBENFY STARTER PACK</b> .....	48
<b>BAFIERTAM</b> .....	22	<b>COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)</b> .....	49
<b>BALVERSA</b> .....	23	<b>COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML</b> .....	101
<b>BENLYSTA SUBCUTANEOUS</b> .....	25		
<i>benztropine oral</i> .....	107		

<b>COPIKTRA</b> .....	50	<i>droxidopa oral capsule 100 mg, 200 mg, 300 mg</i> .....	183
<b>CORLANOR ORAL SOLUTION</b> .....	51	<b>DUPIXENT PEN SUBCUTANEOUS</b>	
<b>COSENTYX (2 SYRINGES)</b> .....	52	<b>PEN INJECTOR 200 MG/1.14 ML, 300</b>	
<b>COSENTYX PEN (2 PENS)</b> .....	52	<b>MG/2 ML</b> .....	64
<b>COSENTYX SUBCUTANEOUS</b>		<b>DUPIXENT SYRINGE</b>	
<b>SYRINGE 75 MG/0.5 ML</b> .....	52	<b>SUBCUTANEOUS SYRINGE 200</b>	
<b>COSENTYX UNOREADY PEN</b> .....	52	<b>MG/1.14 ML, 300 MG/2 ML</b> .....	64
<b>COTELLIC</b> .....	53	<b>DUVYZAT</b> .....	66
<b>CYLTEZO(CF)</b> .....	109	<b>EMGALITY PEN</b> .....	68
<b>CYLTEZO(CF) PEN</b> .....	109	<b>EMGALITY SYRINGE</b>	
<b>CYLTEZO(CF) PEN CROHN'S-UC-HS</b>		<b>SUBCUTANEOUS SYRINGE 120</b>	
.....	109	<b>MG/ML, 300 MG/3 ML (100 MG/ML X</b>	
<b>CYLTEZO(CF) PEN PSORIASIS-UV</b> ..	109	<b>3)</b> .....	68
<b>CYSTARAN</b> .....	54	<b>ENBREL MINI</b> .....	70
<i>dalfampridine</i> .....	14	<b>ENBREL SUBCUTANEOUS</b>	
<b>DANZITEN</b> .....	55	<b>SOLUTION</b> .....	70
<i>dasatinib</i> .....	259	<b>ENBREL SUBCUTANEOUS SYRINGE</b>	
<b>DAURISMO ORAL TABLET 100 MG,</b>		<b>25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)</b> ..	70
<b>25 MG</b> .....	57	<b>ENBREL SURECLICK</b> .....	70
<b>DAYBUE</b> .....	58	<b>ENDOCET</b> .....	215
<i>deferasirox</i> .....	59	<b>EPCLUSA ORAL PELLETS IN</b>	
<i>deferiprone</i> .....	85	<b>PACKET 150-37.5 MG</b> .....	71
<b>DIACOMIT ORAL CAPSULE 250 MG,</b>		<b>EPIDIOLEX</b> .....	72
<b>500 MG</b> .....	60	<b>EPRONTIA</b> .....	73
<b>DIACOMIT ORAL POWDER IN</b>		<i>ergotamine-caffeine</i> .....	74
<b>PACKET 250 MG, 500 MG</b> .....	60	<b>ERIVEDGE</b> .....	75
<b>DIAZEPAM INTENSOL</b> .....	215	<b>ERLEADA ORAL TABLET 240 MG,</b>	
<i>diazepam oral solution 5 mg/5 ml (1</i>		<b>60 MG</b> .....	76
<i>mg/ml)</i> .....	215	<i>erlotinib</i> .....	67
<i>diazepam oral tablet</i> .....	215	<i>eszopiclone</i> .....	215
<i>diclofenac sodium topical gel 3 %</i> .....	257	<b>EVENITY SUBCUTANEOUS</b>	
<i>dihydroergotamine nasal</i> .....	61	<b>SYRINGE 210MG/2.34ML (</b>	
<i>dimethyl fumarate oral capsule, delayed</i>		<b>105MG/1.17MLX2)</b> .....	77
<i>release(dr/ec) 120 mg, 120 mg (14)- 240</i>		<i>everolimus (antineoplastic) oral tablet 10</i>	
<i>mg (46), 240 mg</i> .....	279	<i>mg, 2.5 mg, 5 mg, 7.5 mg</i> .....	6
<b>DOPTELET (10 TAB PACK)</b> .....	62	<i>everolimus (antineoplastic) oral tablet for</i>	
<b>DOPTELET (15 TAB PACK)</b> .....	62	<i>suspension 2 mg, 3 mg, 5 mg</i> .....	6
<b>DOPTELET (30 TAB PACK)</b> .....	62	<b>EVRYSDI ORAL RECON SOLN</b> .....	78
<i>doxepin oral capsule</i> .....	107	<b>FABHALTA</b> .....	79
<i>doxepin oral concentrate</i> .....	107	<b>FANAPT ORAL TABLET</b> .....	81
<i>doxepin oral tablet</i> .....	107	<b>FANAPT ORAL TABLETS,DOSE</b>	
<b>DRIZALMA SPRINKLE ORAL</b>		<b>PACK</b> .....	81
<b>CAPSULE, DELAYED REL</b>		<b>FASENRA PEN</b> .....	82
<b>SPRINKLE 20 MG, 30 MG, 40 MG, 60</b>		<b>FASENRA SUBCUTANEOUS</b>	
<b>MG</b> .....	63	<b>SYRINGE 10 MG/0.5 ML, 30 MG/ML</b> ....	82

<i>febuxostat</i> .....	296	<b>HUMIRA PEN</b> .....	109
<i>fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr</i> .....	215	<b>HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML</b> .....	109
<b>FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)</b> .....	86	<b>HUMIRA(CF)</b> .....	109
<b>FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG</b> .....	86	<b>HUMIRA(CF) PEN</b> .....	109
<b>FILSUEVZ</b> .....	87	<b>HUMIRA(CF) PEN CROHNS-UC-HS..</b>	109
<i>fingolimod</i> .....	100	<b>HUMIRA(CF) PEN PSOR-UV-ADOL HS</b> .....	109
<b>FINTEPLA</b> .....	88	<i>hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i> .....	215
<b>FIRDAPSE</b> .....	91	<i>hydromorphone oral liquid</i> .....	215
<b>FIRMAGON KIT W DILUENT SYRINGE</b> .....	92	<i>hydromorphone oral tablet 2 mg, 4 mg, 8 mg</i> .....	215
<b>FOTIVDA</b> .....	94	<i>hydroxyzine hcl oral tablet</i> .....	107
<b>FRUZAQLA ORAL CAPSULE 1 MG, 5 MG</b> .....	95	<b>IBRANCE</b> .....	111
<b>FUROSCIX</b> .....	96	<b>IBSRELA</b> .....	112
<i>gabapentin oral capsule 100 mg, 300 mg, 400 mg</i> .....	97	<i>icatibant</i> .....	89
<i>gabapentin oral solution 250 mg/5 ml</i> .....	97	<b>ICLUSIG</b> .....	113
<i>gabapentin oral tablet 600 mg, 800 mg</i> .....	97	<b>IDHIFA ORAL TABLET 100 MG, 50 MG</b> .....	114
<b>GAMMAGARD LIQUID</b> .....	115	<i>imatinib oral tablet 100 mg, 400 mg</i> .....	102
<b>GAMMAGARD S-D (IGA &lt; 1 MCG/ML)</b> .....	115	<b>IMBRUVICA ORAL CAPSULE 140 MG, 70 MG</b> .....	117
<b>GAMMAKED INJECTION SOLUTION 1 GRAM/10 ML (10 %)</b> .....	115	<b>IMBRUVICA ORAL SUSPENSION....</b>	117
<b>GAMMAPLEX</b> .....	115	<b>IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG</b> .....	117
<b>GAMMAPLEX (WITH SORBITOL)</b> .....	115	<i>imipramine hcl</i> .....	107
<b>GAMUNEX-C INJECTION SOLUTION 1 GRAM/10 ML (10 %)</b> .....	115	<b>IMKELDI</b> .....	118
<b>GATTEX 30-VIAL</b> .....	98	<b>INCRELEX</b> .....	120
<b>GAUZE PAD TOPICAL BANDAGE 2 X 2 "</b> .....	125	<b>INLYTA</b> .....	122
<b>GAVRETO</b> .....	99	<b>INQOVI</b> .....	123
<i>gefitinib</i> .....	130	<b>INREBIC</b> .....	124
<b>GILOTRIF</b> .....	67	<b>ITOVEBI ORAL TABLET 3 MG, 9 MG</b> .....	131
<i>glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml</i> .....	101	<i>itraconazole oral capsule</i> .....	132
<b>GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML</b> .....	101	<i>ivabradine oral tablet 5 mg, 7.5 mg</i> .....	51
<b>GLEOSTINE</b> .....	104	<i>ivermectin oral tablet 3 mg</i> .....	133
<i>guanfacine oral tablet extended release 24 hr</i> .....	5	<b>IWILFIN</b> .....	134
		<b>JAKAFI</b> .....	135
		<b>JAVYGTOR</b> .....	148
		<b>JAYPIRCA ORAL TABLET 100 MG, 50 MG</b> .....	136
		<b>JOENJA</b> .....	137

<b>KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG</b> .....	138	<b>LUPRON DEPOT-PED INTRAMUSCULAR KIT 7.5 MG (PED)</b> .....	162
<b>KALYDECO ORAL TABLET</b> .....	138	<b>LUPRON DEPOT-PED</b>	
<b>KERENDIA</b> .....	139	<b>INTRAMUSCULAR SYRINGE KIT</b> .....	162
<b>KESIMPTA PEN</b> .....	140	<i>lurasidone oral tablet 120 mg, 20 mg, 40</i>	
<b>KEVZARA</b> .....	141	<i>mg, 60 mg, 80 mg</i> .....	149
<b>KINERET</b> .....	143	<b>LYNPARZA</b> .....	163
<b>KISQALI FEMARA CO-PACK ORAL TABLET 400 MG/DAY(200 MG X 2)- 2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG</b> .....	144	<b>LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)</b> .....	166
<b>KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)</b> .....	144	<b>MAVYRET ORAL PELLETS IN PACKET</b> .....	167
<b>KOSELUGO ORAL CAPSULE 10 MG, 25 MG</b> .....	146	<b>MAVYRET ORAL TABLET</b> .....	167
<b>KRAZATI</b> .....	147	<i>megestrol oral suspension 400 mg/10 ml</i>	
<i>lapatinib</i> .....	293	<i>(40 mg/ml), 625 mg/5 ml (125 mg/ml)</i> .....	168
<b>LAZCLUZE ORAL TABLET 240 MG, 80 MG</b> .....	150	<i>megestrol oral tablet</i> .....	168
<i>lenalidomide</i> .....	233	<b>MEKINIST ORAL RECON SOLN</b> .....	169
<b>LENVIMA</b> .....	151	<b>MEKINIST ORAL TABLET 0.5 MG, 2 MG</b> .....	169
<b>LEUKINE INJECTION RECON SOLN</b> .....	152	<b>MEKTOVI</b> .....	170
<b>LIBERVANT</b> .....	153	<i>methadone oral solution 10 mg/5 ml, 5</i>	
<i>lidocaine hcl mucous membrane solution 4</i>		<i>mg/5 ml</i> .....	215
<i>% (40 mg/ml)</i> .....	286	<i>methadone oral tablet 10 mg, 5 mg</i> .....	215
<i>lidocaine topical adhesive patch,medicated</i>		<i>metyrosine</i> .....	171
<i>5 %</i> .....	154	<i>mifepristone oral tablet 300 mg</i> .....	145
<i>lidocaine topical ointment</i> .....	286	<i>miglustat</i> .....	336
<i>lidocaine-prilocaine topical cream</i> .....	286	<i>modafinil</i> .....	220
<b>LITFULO</b> .....	155	<i>morphine concentrate oral solution</i> .....	215
<b>LIVTENCITY</b> .....	156	<i>morphine oral solution 10 mg/5 ml, 20</i>	
<b>LOKELMA</b> .....	157	<i>mg/5 ml (4 mg/ml)</i> .....	215
<b>LONSURF</b> .....	158	<i>morphine oral tablet</i> .....	215
<b>LORAZEPAM INTENSOL</b> .....	215	<i>morphine oral tablet extended release 100</i>	
<i>lorazepam oral tablet 0.5 mg, 1 mg, 2 mg</i> ..	215	<i>mg, 15 mg, 200 mg, 30 mg, 60 mg</i> .....	215
<b>LORBRENA ORAL TABLET 100 MG, 25 MG</b> .....	159	<b>MOUNJARO</b> .....	105
<b>LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG</b> .....	161	<b>MULPLETA</b> .....	172
<b>LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT</b>		<b>NAMZARIC</b> .....	174
<b>11.25 MG</b> .....	162	<b>NAYZILAM</b> .....	175
		<b>NEMLUVIO</b> .....	176
		<b>NERLYNX</b> .....	177
		<b>NEXLETOL</b> .....	179
		<b>NEXLIZET</b> .....	179
		<b>NINLARO</b> .....	181
		<i>nitisinone</i> .....	182
		<b>NORDITROPIN FLEXPOR</b> .....	106
		<b>NUBEQA</b> .....	184

<b>NUCALA SUBCUTANEOUS AUTO-INJECTOR</b> .....	185	<i>oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg</i> .....	215
<b>NUCALA SUBCUTANEOUS RECON SOLN</b> .....	185	<b>OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)</b> .....	105
<b>NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML</b>	185	<b>PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML</b> .....	207
<b>NUEDEXTA</b> .....	187	<b>PANRETIN</b> .....	208
<b>NUPLAZID</b> .....	188	<b>PANZYGA</b> .....	115
<b>NURTEC ODT</b> .....	189	<i>pazopanib</i> .....	314
<b>OALIVA</b> .....	192	<b>PEGASYS</b> .....	126
<b>OCTAGAM</b> .....	115	<b>PEMAZYRE</b> .....	209
<i>octreotide acetate injection solution</i> .....	193	<b>PHEBURANE</b> .....	210
<b>ODOMZO</b> .....	194	<i>phenobarbital</i> .....	108
<b>OFEV</b> .....	128	<b>PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)</b> .....	211
<b>OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG</b> .....	195	<i>pirfenidone oral capsule</i> .....	128
<b>OJEMDA ORAL SUSPENSION FOR RECONSTITUTION</b> .....	196	<i>pirfenidone oral tablet</i> .....	128
<b>OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500 MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)</b> .....	196	<b>PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML</b> .....	127
<b>OJJAARA</b> .....	197	<b>PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML</b> .....	127
<b>OLUMIANT</b> .....	198	<b>POMALYST</b> .....	212
<b>ONUREG</b> .....	200	<i>posaconazole oral tablet, delayed release (dr/ec)</i> .....	213
<b>OPSUMIT</b> .....	222	<i>pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg</i> .....	165
<b>OPSYNVI</b> .....	222	<i>pregabalin oral solution</i> .....	165
<b>ORENCIA CLICKJECT</b> .....	201	<b>PRENATAL VITAMIN PLUS LOW IRON</b> .....	214
<b>ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML</b> .....	201	<b>PREVYMIS ORAL PELLETS IN PACKET</b> .....	218
<b>ORGOVYX</b> .....	203	<b>PRIVIGEN</b> .....	115
<b>ORKAMBI ORAL GRANULES IN PACKET</b> .....	204	<b>PROLASTIN-C INTRAVENOUS SOLUTION</b> .....	12
<b>ORKAMBI ORAL TABLET</b> .....	204	<b>PROLIA</b> .....	219
<b>ORSERDU ORAL TABLET 345 MG, 86 MG</b> .....	205	<b>PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG</b> .....	284
<b>OTEZLA</b> .....	206	<b>PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG</b> .....	284
<b>OTEZLA STARTER ORAL TABLETS, DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47)</b> .....	206		
<i>oxycodone oral capsule</i> .....	215		
<i>oxycodone oral concentrate</i> .....	215		
<i>oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg</i> .....	215		



<i>promethazine oral</i> .....	107	<b>SECUADO</b> .....	250
<b>PULMOZYME</b> .....	223	<b>SIGNIFOR</b> .....	261
<i>pyrimethamine</i> .....	56	<i>sildenafil (pulm.hypertension) oral tablet</i> ..	222
<b>QINLOCK</b> .....	224	<b>SIMPONI SUBCUTANEOUS PEN</b>	
<i>quinine sulfate</i> .....	225	<b>INJECTOR 100 MG/ML, 50 MG/0.5</b>	
<b>QULIPTA</b> .....	226	<b>ML</b> .....	251
<b>RADICAVA ORS STARTER KIT</b>		<b>SIMPONI SUBCUTANEOUS</b>	
<b>SUSP</b> .....	227	<b>SYRINGE 100 MG/ML, 50 MG/0.5 ML</b>	251
<b>RAVICTI</b> .....	228	<b>SIRTURO</b> .....	253
<b>RECORLEV</b> .....	261	<b>SKYCLARYS</b> .....	254
<b>REGRANEX</b> .....	229	<b>SKYRIZI SUBCUTANEOUS PEN</b>	
<b>REPATHA PUSHTRONEX</b> .....	230	<b>INJECTOR</b> .....	255
<b>REPATHA SURECLICK</b> .....	230	<b>SKYRIZI SUBCUTANEOUS</b>	
<b>REPATHA SYRINGE</b> .....	230	<b>SYRINGE 150 MG/ML</b> .....	255
<b>RETEVMO ORAL CAPSULE 40 MG,</b>		<b>SKYRIZI SUBCUTANEOUS</b>	
<b>80 MG</b> .....	232	<b>WEARABLE INJECTOR 180 MG/1.2</b>	
<b>RETEVMO ORAL TABLET 120 MG,</b>		<b>ML (150 MG/ML), 360 MG/2.4 ML (150</b>	
<b>160 MG, 40 MG, 80 MG</b> .....	232	<b>MG/ML)</b> .....	255
<b>REVUFORJ ORAL TABLET 110 MG,</b>		<i>sodium oxybate</i> .....	333
<b>160 MG</b> .....	234	<i>sodium phenylbutyrate</i> .....	31
<b>REXULTI ORAL TABLET</b> .....	235	<i>sofosbuvir-velpatasvir</i> .....	71
<b>REZDIFFRA</b> .....	236	<b>SOHONOS ORAL CAPSULE 1 MG, 1.5</b>	
<b>REZLIDHIA</b> .....	237	<b>MG, 10 MG, 2.5 MG, 5 MG</b> .....	256
<b>RINVOQ LQ</b> .....	240	<b>SOMAVERT</b> .....	258
<b>RINVOQ ORAL TABLET EXTENDED</b>		<i>sorafenib</i> .....	178
<b>RELEASE 24 HR 15 MG, 30 MG, 45</b>		<b>STELARA SUBCUTANEOUS</b>	
<b>MG</b> .....	238	<b>SOLUTION</b> .....	260
<b>RIVFLOZA SUBCUTANEOUS</b>		<b>STELARA SUBCUTANEOUS</b>	
<b>SOLUTION</b> .....	241	<b>SYRINGE 45 MG/0.5 ML, 90 MG/ML</b> ..	260
<b>RIVFLOZA SUBCUTANEOUS</b>		<b>STIVARGA</b> .....	262
<b>SYRINGE 128 MG/0.8 ML, 160</b>		<i>sunitinib malate</i> .....	263
<b>MG/ML</b> .....	241	<b>SYMDEKO</b> .....	264
<b>ROZLYTREK ORAL CAPSULE 100</b>		<b>SYMPAZAN</b> .....	265
<b>MG, 200 MG</b> .....	242	<b>SYNAREL</b> .....	266
<b>ROZLYTREK ORAL PELLETS IN</b>		<b>TABRECTA</b> .....	267
<b>PACKET</b> .....	242	<i>tadalafil (pulm. hypertension)</i> .....	222
<b>RUBRACA</b> .....	243	<i>tadalafil oral tablet 2.5 mg, 5 mg</i> .....	42
<i>rufinamide</i> .....	24	<b>TAFINLAR ORAL CAPSULE</b> .....	268
<b>RYBELSUS ORAL TABLET 14 MG, 3</b>		<b>TAFINLAR ORAL TABLET FOR</b>	
<b>MG, 7 MG</b> .....	105	<b>SUSPENSION</b> .....	268
<b>RYDAPT</b> .....	244	<b>TAGRISSO</b> .....	269
<b>SAJAZIR</b> .....	89	<b>TALTZ AUTOINJECTOR</b> .....	270
<i>sapropterin</i> .....	148	<b>TALTZ SYRINGE SUBCUTANEOUS</b>	
<b>SAVELLA</b> .....	248	<b>SYRINGE 20 MG/0.25 ML, 40 MG/0.5</b>	
<b>SCEMBLIX ORAL TABLET 100 MG,</b>		<b>ML, 80 MG/ML</b> .....	270
<b>20 MG, 40 MG</b> .....	249	<b>TALZENNA</b> .....	272

<b>TASCENSO ODT</b> .....	274	<b>UPTRAVI ORAL TABLETS,DOSE</b>	
<b>TASIGNA</b> .....	275	<b>PACK</b> .....	222
<i>tasimelteon</i> .....	276	<b>VALCHLOR</b> .....	297
<i>tazarotene topical cream</i> .....	277	<b>VALTOCO</b> .....	298
<b>TAZVERIK</b> .....	278	<i>vancomycin oral capsule 125 mg, 250 mg</i> .....	299
<b>TEPMETKO</b> .....	280	<b>VANFLYTA</b> .....	300
<i>teriflunomide</i> .....	18	<b>VENCLEXTA ORAL TABLET 10 MG,</b>	
<i>teriparatide subcutaneous pen injector 20</i>		<b>100 MG, 50 MG</b> .....	301
<i>mcg/dose (620mcg/2.48ml)</i> .....	93	<b>VENCLEXTA STARTING PACK</b> .....	301
<i>testosterone cypionate</i> .....	281	<b>VERQUVO</b> .....	302
<i>testosterone enanthate</i> .....	281	<b>VERZENIO</b> .....	303
<i>testosterone transdermal gel in metered-</i>		<b>VIBERZI</b> .....	304
<i>dose pump 20.25 mg/1.25 gram (1.62 %)</i> .....	281	<i>vigabatrin</i> .....	245
<i>testosterone transdermal gel in packet</i> .....	281	<b>VIGADRONE</b> .....	245
<i>testosterone transdermal solution in</i>		<b>VIGPODER</b> .....	245
<i>metered pump w/app</i> .....	281	<b>VIJOICE ORAL GRANULES IN</b>	
<i>tetrabenazine oral tablet 12.5 mg, 25 mg</i> .....	323	<b>PACKET</b> .....	306
<b>THALOMID ORAL CAPSULE 100</b>		<b>VIJOICE ORAL TABLET 125 MG, 250</b>	
<b>MG, 50 MG</b> .....	283	<b>MG/DAY (200 MG X1-50 MG X1), 50</b>	
<b>TIBSOVO</b> .....	285	<b>MG</b> .....	306
<b>TOBI PODHALER</b> .....	41	<i>vilazodone</i> .....	305
<i>tobramycin in 0.225 % nacl</i> .....	41	<b>VITRAKVI ORAL CAPSULE 100 MG,</b>	
<i>tobramycin inhalation</i> .....	41	<b>25 MG</b> .....	307
<i>tolvaptan</i> .....	246	<b>VITRAKVI ORAL SOLUTION</b> .....	307
<i>tramadol oral tablet 50 mg</i> .....	215	<b>VIVJOA</b> .....	308
<i>tramadol-acetaminophen</i> .....	215	<b>VIZIMPRO</b> .....	309
<i>tretinoin topical cream</i> .....	287	<b>VONJO</b> .....	310
<i>tretinoin topical gel 0.01 %, 0.025 %</i> .....	287	<b>VORANIGO ORAL TABLET 10 MG,</b>	
<b>TRIKAFTA ORAL GRANULES IN</b>		<b>40 MG</b> .....	311
<b>PACKET, SEQUENTIAL</b> .....	288	<i>voriconazole intravenous</i> .....	312
<b>TRIKAFTA ORAL TABLETS,</b>		<b>VOSEVI</b> .....	313
<b>SEQUENTIAL</b> .....	288	<b>VOWST</b> .....	84
<i>trimipramine</i> .....	107	<b>VRAYLAR ORAL CAPSULE</b> .....	315
<b>TRINTELLIX</b> .....	289	<b>VUMERITY</b> .....	316
<b>TRULICITY</b> .....	105	<b>VYNDAQEL</b> .....	16
<b>TRUQAP</b> .....	290	<b>WELIREG</b> .....	317
<b>TUKYSA ORAL TABLET 150 MG, 50</b>		<b>XALKORI ORAL CAPSULE</b> .....	318
<b>MG</b> .....	291	<b>XALKORI ORAL PELLET 150 MG, 20</b>	
<b>TURALIO ORAL CAPSULE 125 MG</b> .....	292	<b>MG, 50 MG</b> .....	318
<b>TYMLOS</b> .....	294	<b>XCOPRI</b> .....	319
<b>UBRELVY ORAL TABLET 100 MG,</b>		<b>XCOPRI MAINTENANCE PACK</b> .....	319
<b>50 MG</b> .....	295	<b>XCOPRI TITRATION PACK</b> .....	319
<b>UPTRAVI ORAL TABLET 1,000 MCG,</b>		<b>XDEMZY</b> .....	320
<b>1,200 MCG, 1,400 MCG, 1,600 MCG,</b>		<b>XELJANZ ORAL SOLUTION</b> .....	322
<b>200 MCG, 400 MCG, 600 MCG, 800</b>		<b>XELJANZ ORAL TABLET</b> .....	321
<b>MCG</b> .....	222	<b>XELJANZ XR</b> .....	321



<b>XERMELO</b> .....	324
<b>XGEVA</b> .....	325
<b>XIFAXAN ORAL TABLET 550 MG</b> .....	326
<b>XOLAIR</b> .....	327
<b>XOLREMDI</b> .....	329
<b>XOSPATA</b> .....	330
<b>XPOVIO ORAL TABLET 100</b> <b>MG/WEEK (50 MG X 2), 40</b> <b>MG/WEEK (40 MG X 1), 40MG</b> <b>TWICE WEEK (40 MG X 2), 60</b> <b>MG/WEEK (60 MG X 1), 60MG</b> <b>TWICE WEEK (120 MG/WEEK), 80</b> <b>MG/WEEK (40 MG X 2), 80MG</b> <b>TWICE WEEK (160 MG/WEEK)</b> .....	331
<b>XTANDI ORAL CAPSULE</b> .....	332
<b>XTANDI ORAL TABLET 40 MG, 80</b> <b>MG</b> .....	332
<b>XYREM</b> .....	333
<b>YARGESA</b> .....	336
<b>YONSA</b> .....	334
<b>YORVIPATH SUBCUTANEOUS PEN</b> <b>INJECTOR 168 MCG/0.56 ML, 294</b> <b>MCG/0.98 ML, 420 MCG/1.4 ML</b> .....	335
<b>YUFLYMA(CF)</b> .....	109
<b>YUFLYMA(CF) AI CROHN'S-UC-HS</b> .....	109
<b>YUFLYMA(CF) AUTOINJECTOR</b> .....	109
<i>zaleplon oral capsule 10 mg, 5 mg</i> .....	215
<b>ZAVZPRET</b> .....	337
<b>ZEJULA ORAL TABLET</b> .....	338
<b>ZELBORAF</b> .....	339
<b>ZEPOSIA</b> .....	340
<b>ZEPOSIA STARTER KIT (28-DAY)</b> .....	340
<b>ZEPOSIA STARTER PACK (7-DAY)</b> ...	340
<b>ZILBRYSQ SUBCUTANEOUS</b> <b>SYRINGE 16.6 MG/0.416 ML, 23</b> <b>MG/0.574 ML, 32.4 MG/0.81 ML</b> .....	173
<b>ZOLINZA</b> .....	341
<i>zolpidem oral tablet</i> .....	215
<b>ZONISADE</b> .....	342
<b>ZTALMY</b> .....	343
<b>ZURZUVAE ORAL CAPSULE 20 MG,</b> <b>25 MG, 30 MG</b> .....	344
<b>ZYDELIG</b> .....	345
<b>ZYKADIA</b> .....	346

# GnRH Agonists

---

## Products Affected

- **ELIGARD 22.5 MG (3 MONTH) SUBCUTANEOUS SYRINGE**
- **ELIGARD 30 MG (4 MONTH) SUBCUTANEOUS SYRINGE**
- **ELIGARD 45 MG (6 MONTH) SUBCUTANEOUS SYRINGE**
- **ELIGARD 7.5 MG (1 MONTH) SUBCUTANEOUS SYRINGE**
- **TRELSTAR 11.25 MG IM SUSPENSION**
- **TRELSTAR 22.5 MG IM SUSPENSION**
- **TRELSTAR 3.75 MG IM SUSPENSION**

## Details

<b>Criteria</b>	Require a trial of Lupron Depot (Step 1 drug) in the last 180 days when being utilized for the same medically accepted indication
-----------------	-----------------------------------------------------------------------------------------------------------------------------------

# Herpetic Keratitis

---

## Products Affected

- **ZIRGAN 0.15 % EYE GEL**

## Details

<b>Criteria</b>	Require a 1 month trial of generic trifluridine eye drops (Step 1 drug) in the last 90 days
-----------------	---------------------------------------------------------------------------------------------

# Pulmonary Antiinflammatory

---

## Products Affected

- *fluticasone propionate 100 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 110 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 220 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 250 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 44 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 50 mcg/actuation blister powder for inhalation*

## Details

<b>Criteria</b>	Require a 1 month trial of Qvar and Asmanex/Asmanex HFA (Step 1 drugs) in the last 180 days
-----------------	---------------------------------------------------------------------------------------------

# Rytary

---

## Products Affected

- **RYTARY 23.75 MG-95 MG  
CAPSULE,EXTENDED RELEASE**
- **RYTARY 36.25 MG-145 MG  
CAPSULE,EXTENDED RELEASE**
- **RYTARY 48.75 MG-195 MG  
CAPSULE,EXTENDED RELEASE**
- **RYTARY 61.25 MG-245 MG  
CAPSULE,EXTENDED RELEASE**

## Details

<b>Criteria</b>	Require a trial of generic carbidopa/levodopa product (Step 1 drug) in the last 90 days
-----------------	-----------------------------------------------------------------------------------------

# Índice de drogas

<b>ELIGARD 22.5 MG (3 MONTH)</b>	
<b>SUBCUTANEOUS SYRINGE .....</b>	<b>1</b>
<b>ELIGARD 30 MG (4 MONTH)</b>	
<b>SUBCUTANEOUS SYRINGE .....</b>	<b>1</b>
<b>ELIGARD 45 MG (6 MONTH)</b>	
<b>SUBCUTANEOUS SYRINGE .....</b>	<b>1</b>
<b>ELIGARD 7.5 MG (1 MONTH)</b>	
<b>SUBCUTANEOUS SYRINGE .....</b>	<b>1</b>
<i>fluticasone propionate 100 mcg/actuation</i>	
<i>blister powder for inhalation .....</i>	<i>3</i>
<i>fluticasone propionate 110 mcg/actuation</i>	
<i>hfa aerosol inhaler .....</i>	<i>3</i>
<i>fluticasone propionate 220 mcg/actuation</i>	
<i>hfa aerosol inhaler .....</i>	<i>3</i>
<i>fluticasone propionate 250 mcg/actuation</i>	
<i>blister powder for inhalation .....</i>	<i>3</i>
<i>fluticasone propionate 44 mcg/actuation</i>	
<i>hfa aerosol inhaler .....</i>	<i>3</i>
<i>fluticasone propionate 50 mcg/actuation</i>	
<i>blister powder for inhalation .....</i>	<i>3</i>
<b>RYTARY 23.75 MG-95 MG</b>	
<b>CAPSULE,EXTENDED RELEASE .....</b>	<b>4</b>
<b>RYTARY 36.25 MG-145 MG</b>	
<b>CAPSULE,EXTENDED RELEASE .....</b>	<b>4</b>
<b>RYTARY 48.75 MG-195 MG</b>	
<b>CAPSULE,EXTENDED RELEASE .....</b>	<b>4</b>
<b>RYTARY 61.25 MG-245 MG</b>	
<b>CAPSULE,EXTENDED RELEASE .....</b>	<b>4</b>
<b>TRELSTAR 11.25 MG IM</b>	
<b>SUSPENSION .....</b>	<b>1</b>
<b>TRELSTAR 22.5 MG IM SUSPENSION..</b>	<b>1</b>
<b>TRELSTAR 3.75 MG IM SUSPENSION..</b>	<b>1</b>
<b>ZIRGAN 0.15 % EYE GEL .....</b>	<b>2</b>