Medicare Part D: 5 Tier Closed Venture Formulary 2026

Please click here.

For Medicare Part D: Prior Authorization Criteria

Please click here.

For Medicare Part D: Step Therapy Criteria

Please click here.

For more recent information or other questions, please contact:

Blue Rx PDP Complete
Blue Rx PDP Plus
Freedom Blue PPO Classic (PPO)
Freedom Blue PPO Select (PPO)
Freedom Blue PPO Standard (PPO)
Security Blue HMO-POS Deluxe (HMO-POS)
Security Blue HMO-POS Standard (HMO-POS)

Freedom Blue PPO (PA) Pharmacy Service at 1-800-550-8722
Security Blue HMO-POS Pharmacy Service at 1-800-935-2583
Community Blue Medicare HMO Pharmacy Service at 1-888-234-5397
Blue Rx PDP Pharmacy Service at 1-800-290-3914

For TTY users, *711 National Relay Service*, Monday through Sunday, 8:00 a.m. to 8:00 p.m.
Visit medicare.highmark.com.

Notice: Oral Antivirals for COVID-19 Coverage

Paxlovid is covered with $0 copay when dispensed at a preferred or non-preferred pharmacy.

Formulary ID: 26017 Version: 20

Updated: 11/2026

Note to existing members: This Formulary has changed since last year. Please review this document to make sure that it still contains the drugs you take.

When this Drug List (Formulary) refers to “we,” “us,” or “our,” it means Highmark Senior Health Company, Highmark Choice Company, or Highmark Health Insurance Company..

When it refers to “plan” or “our plan,” it means Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP.

This document includes a Drug List (formulary) for our plan, which is current as of January 1, 2026. For an updated Drug List (formulary), please contact us. Our contact information, along with the date we last updated the Drug List (formulary), appears on the front and back cover pages.

You must generally use network pharmacies to use your prescription drug benefit. Benefits, formulary, pharmacy network, and/or copayments/coinsurance may change on January 1, 2027, and from time to time during the year.

What is the Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP formulary?

In this document, we use the terms Drug List and formulary to mean the same thing. A formulary is a list of covered drugs selected by our plan in consultation with a team of health care providers, which represents the prescription therapies believed to be a necessary part of a quality treatment program. Our plan will generally cover the drugs listed in our formulary as long as the drug is medically necessary, the prescription is filled at a Freedom Blue PPO, Security Blue HMO-POS, or Blue Rx PDP network pharmacy, and other plan rules are followed. For more information on how to fill your prescriptions, please review your Evidence of Coverage.

Can the formulary change?

Most changes in drug coverage happen on January 1, but we may add or remove drugs on the formulary during the year, move them to different cost-sharing tiers, or add new restrictions. We must follow the Medicare rules in making these changes. Updates to the formulary are posted monthly to our website, **medicare.highmark.com/formulary**.

Changes that can affect you this year: In the below cases, you will be affected by coverage changes during the year:

- **Immediate substitutions of certain new versions of brand name drugs and original biological products.** We may immediately remove a drug from our formulary if we are replacing it with a certain new version of that drug that will appear on the same or lower cost-sharing tier and with the same or fewer restrictions. When we add a new version of a drug to our formulary, we may decide to keep the brand name drug or original biological product on our formulary, but immediately move it to a different cost-sharing tier or add new restrictions.

  We can make these immediate changes only if we are adding a new generic version of a brand name drug, or adding certain new biosimilar versions of an original biological product, that was already on the formulary (for example, adding an interchangeable biosimilar that can be substituted for an original biological product by a pharmacy without a new prescription.)

  If you are currently taking the brand name drug or original biological product, we may not tell you in advance before we make an immediate change, but we will later provide you with information about the specific change(s) we have made.

  If we make such a change, you or your prescriber can ask us to make an exception and continue to cover for you the drug that is being changed. For more information, see the section below titled “How do I request an exception to the Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP’s Formulary?”

  Some of these drug types may be new to you. For more information, see the section below titled “What are original biological products and how are they related to biosimilars?”

- **Drugs removed from the market.** If a drug is withdrawn from sale by the manufacturer or the Food and Drug Administration (FDA) determines to be withdrawn for safety or effectiveness reasons, we may immediately remove the drug from our formulary and later provide notice to members who take the drug.

- **Other changes.** We may make other changes that affect members currently taking a drug. For instance, we may remove a brand name drug from the formulary when adding a generic equivalent or remove an original biological product when adding a biosimilar. We may also apply new restrictions to the brand name drug or original biological product, or move it to a different cost-sharing tier, or both. We may make changes based on new clinical guidelines. If we remove drugs from our formulary, add prior authorization, quantity limits and/or step therapy restrictions on a drug, or move a drug to a higher cost-sharing tier, we must notify affected members of the change at least 30 days before the change becomes effective. Alternatively, when a member requests a refill of the drug, they may receive a 31-day supply of the drug and notice of the change.

If we make these other changes, you or your prescriber can ask us to make an exception for you and continue to cover the drug you have been taking. The notice we provide you will also include information on how to request an exception, and you can also find information in the section below entitled “How do I request an exception to the Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP’s Formulary?”

Changes that will not affect you if you are currently taking the drug. Generally, if you are taking a drug on our 2026 formulary that was covered at the beginning of the year, we will not discontinue or reduce coverage of the drug during the 2026 coverage year except as described above. This means these drugs will remain available at the same cost-sharing and with no new restrictions for those members taking them for the remainder of the coverage year. You will not get direct notice this year about changes that do not affect you. However, on January 1 of the next year, such changes would affect you, and it is important to check the formulary for the new benefit year for any changes to drugs.

The enclosed formulary is current as of November 1, 2026. To get updated information about the drugs covered by our plan, please contact us. Our contact information appears on the front and back cover pages. In the event of mid-year non-maintenance formulary changes, members will be notified by mail and prospective members will receive an update with this formulary. The most up-to-date formulary is available on our website, **medicare.highmark.com/formulary**.

How do I use the formulary?

There are two ways to find your drug within the formulary:

Medical Condition

The formulary begins on page 9. The drugs in this formulary are grouped into categories depending on the type of medical conditions that they are used to treat. For example, drugs used to treat a heart condition are listed under the category, “Cardiovascular – Hypertension & Lipids.” If you know what your drug is used for, look for the category name in the list that begins on page 9. Then look under the category name for your drug.

Alphabetical Listing

If you are not sure what category to look under, you should look for your drug in the Index that begins on page 9. The Index provides an alphabetical list of all of the drugs included in this document. Both brand name drugs and generic drugs are listed in the Index. Look in the Index and find your drug. Next to your drug, you will see the page number where you can find coverage information. Turn to the page listed in the Index and find the name of your drug in the first column of the list.

What are generic drugs?

Our plan covers both brand name drugs and generic drugs. A generic drug is approved by the FDA as having the same active ingredient as the brand name drug. Generally, generic drugs work just as well as and usually cost less than brand name drugs. There are generic drug substitutes available for many brand name drugs. Generic drugs usually can be substituted for the brand name drug at the pharmacy without needing a new prescription, depending on state laws.

What are original biological products and how are they related to biosimilars?

On the formulary, when we refer to drugs, this could mean a drug or a biological product. Biological products are drugs that are more complex than typical drugs. Since biological products are more complex than typical drugs, instead of having a generic form, they have alternatives that are called biosimilars. Generally, biosimilars work just as well as the original biological product and may cost less. There are biosimilar alternatives for some original biological products. Some biosimilars are interchangeable biosimilars and, depending on state laws, may be substituted for the original biological product at the pharmacy without needing a new prescription, just like generic drugs can be substituted for brand name drugs.

For discussion of drug types, please see the Evidence of Coverage, Chapter 3, Section 3.1, “The ‘Drug List’ tells which Part D drugs are covered.”

Are there any restrictions on my coverage?

Some covered drugs may have additional requirements or limits on coverage. These requirements and limits may include:

- **Prior Authorization:** Our plan requires you or your prescriber to get prior authorization for certain drugs. This means that you will need to get approval from our plan before you fill your prescriptions. If you don’t get approval, our plan may not cover the drug.
- **Quantity Limits:** For certain drugs, our plan limits the amount of the drug that we will cover. For example, our plan provides 31 tablets, per 31 days, for 100mg Drug A. This may be in addition to a standard one-month or three-month supply.
- **Step Therapy:** In some cases, our plan requires you to first try certain drugs to treat your medical condition before we will cover another drug for that condition. For example, if Drug A and Drug B both treat your medical condition, our plan may not cover Drug B unless you try Drug A first. If Drug A does not work for you, our plan will then cover Drug B.

You can find out if your drug has any additional requirements or limits by looking in the formulary that begins on page 9. You can also get more information about the restrictions applied to specific covered drugs by visiting our website. We have posted online document(s) that explain(s) our prior authorization and step therapy restrictions. You may also ask us to send you a copy. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

You can ask our plan to make an exception to these restrictions or limits or for a list of other, similar drugs that may treat your health condition. See the section, “How do I request an exception to the Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP’s Formulary?” on page 6 for information about how to request an exception.

What if my drug is not on the formulary?

If your drug is not included in this formulary (list of covered drugs), you should first contact Pharmacy Service and ask if your drug is covered.

If you learn that our plan does not cover your drug, you have two options:

- You can ask Pharmacy Service for a list of similar drugs that are covered by our plan. When you receive the list, show it to your doctor and ask them to prescribe a similar drug that is covered by our plan.
- You can ask our plan to make an exception and cover your drug. See below for information about how to request an exception.

How do I request an exception to the Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP’s formulary?

You can ask our plan to make an exception to our coverage rules. There are several types of exceptions that you can ask us to make.

- You can ask us to cover a drug even if it is not on our formulary. If approved, this drug will be covered at a pre-determined cost-sharing level, and you would not be able to ask us to provide the drug at a lower cost-sharing level.
- You can ask us to waive a coverage restriction including prior authorization, step therapy, or a quantity limit on your drug. For example, for certain drugs, our plan limits the amount of the drug that we will cover. If your drug has a quantity limit, you can ask us to waive the limit and cover a greater amount.
- You can ask us to cover a formulary drug at a lower cost-sharing level unless the drug is on the specialty tier. If approved, this would lower the amount you must pay for your drug.

Generally, our plan will only approve your request for an exception if the alternative drugs included on the plan’s formulary, the lower cost-sharing drug, or applying the restriction would not be as effective for you and/or would cause you to have adverse effects.

You or your prescriber should contact us to ask for a tiering or, formulary exception, including an exception to a coverage restriction. ***When you request an exception, your prescriber will need to explain the medical reasons why you need the exception.*** Generally, we must make our decision within 72 hours of getting your prescriber’s supporting statement. You can ask for an expedited (fast) decision if you believe, and we agree, that your health could be seriously harmed by waiting up to 72 hours for a decision. If we agree, or if your prescriber asks for a for a fast decision, we must give you a decision no later than 24 hours after we get your prescriber’s supporting statement.

What can I do if my drug is not on the formulary or has a restriction?

As a new or continuing member in our plan you may be taking drugs that are not on our formulary. Or, you may be taking a drug that is on our formulary but has a coverage restriction, such as prior authorization. You should talk to your prescriber about requesting a coverage decision to show that you meet the criteria for approval, switching to an alternative drug that we cover, or requesting a formulary exception so that we will cover the drug you take. While you and your doctor determine the right course of action for you, we may cover your drug in certain cases during the first 90 days you are a member of our plan.

For each of your drugs that is not on our formulary or has a coverage restriction, we will cover a temporary 31-day supply. If your prescription is written for fewer days, we'll allow refills to provide up to a maximum 31-day supply of medication. If coverage is not approved, after your first 31-day supply, we will not pay for these drugs, even if you have been a member of the plan less than 90 days.

If you are a resident of a long-term care facility and you need a drug that is not on our formulary or if your ability to get your drugs is limited, but you are past the first 90 days of membership in our plan, we will cover a 31-day emergency supply of that drug while you pursue a formulary exception.

The above transition process will be implemented to accommodate you if you have an immediate need for a non-formulary drug or a drug that requires prior authorization due to a change in your level of care while you are waiting for an exception request to be processed.

For more information

For more detailed information about your plan's prescription drug coverage, please review your Evidence of Coverage and other plan materials.

If you have questions about your plan, please contact us. Our contact information, along with the date we last updated the formulary, appears on the front and back cover pages.

If you have general questions about Medicare prescription drug coverage, please call Medicare at 1-800-MEDICARE (1-800-633-4227) 24 hours a day/ seven days a week. TTY users should call 1-877-486-2048. Or visit **http://www.medicare.gov**.

Freedom Blue PPO, Security Blue HMO-POS, and Blue Rx PDP formulary

The formulary that begins on the next page provides coverage information about the drugs covered by your plan. If you have trouble finding your drug in the list, turn to the Index that begins on page 9.

The first column of the chart lists the drug name. Brand name drugs are capitalized (e.g., ABELCET) and generic drugs are listed in lower-case italics (e.g., *abacavir*).

The information in the Requirements/Limits column tells you if your plan has any special requirements for coverage of your drug.

The following is a Formulary Format Example Only:

Drug Name	Performance Drug Tier	Requirements/ Limits
Anti-Infectives		
XYZ DRUG	NF	QL-28

Table of Contents

Anti - Infectives 3
Antineoplastic / Immunosuppressant Drugs 12
Autonomic / Cns Drugs, Neurology / Psych 20
Cardiovascular, Hypertension / Lipids 37
Dermatologicals/Topical Therapy 43
Diagnostics / Miscellaneous Agents 49
Ear, Nose / Throat Medications 50
Endocrine/Diabetes 51
Gastroenterology 56
Immunology, Vaccines / Biotechnology 59
Miscellaneous Supplies 62
Musculoskeletal / Rheumatology 62
Obstetrics / Gynecology 64
Ophthalmology 67
Respiratory And Allergy 69
Urologicals 73
Vitamins, Hematinics / Electrolytes 74

lowercase italics = Generic drugs
UPPERCASE BOLD = Brand name drugs

Drug Tier
T1 = Cost-Sharing Tier 1 includes preferred generic drugs. This is the lowest cost-sharing tier.
T2 = Cost-Sharing Tier 2 includes generic drugs.
T3 = Cost-Sharing Tier 3 includes preferred brand name drugs and may include some single-sourced drugs (those generic drugs made by a single manufacturer).
T4 = Cost-Sharing Tier 4 includes non-preferred brand name drugs and may include some single-sourced generic drugs (those generic drugs made by a single manufacturer).
T5 = Cost-Sharing Tier 5 includes specialty drugs. This is the highest cost-sharing tier.

Requirements/Limits
LA = Limited access
PA = Prior authorization required
PA-BvD = This drug may be covered under Medicare part B or D depending on the circumstance. Information may need to be submitted describing the use and setting of the drug to make the determination.
PA-NS = Prior authorization required for new starts only
QL = Quantity limit applies. The quantity limit is noted for each drug. For example, if the quantity limit is QL (90 EA per 180 days), the quantity limit would be 90 units per 180-day supply.
ST = Step therapy applies
ST-NS = Step therapy applies to new starts only

Drug Name	Drug Tier	Requirements/Limits
Anti - Infectives		
abacavir oral solution	T4	
abacavir oral tablet	T3	
abacavir-lamivudine	T4	
acyclovir oral capsule	T2	
acyclovir oral suspension 200 mg/5 ml	T2	
acyclovir oral tablet	T2	
acyclovir sodium intravenous solution	T2	PA-BvD
adefovir	T4	
albendazole	T4	
amantadine hcl oral capsule	T2	QL (124 EA per 31 days)
amantadine hcl oral solution	T2	
amantadine hcl oral tablet	T2	
amikacin injection solution 500 mg/2 ml	T4	
amoxicillin oral capsule	T1	
amoxicillin oral suspension for reconstitution	T1	
amoxicillin oral tablet	T1	
amoxicillin oral tablet,chewable 125 mg, 250 mg	T1	
amoxicillin-pot clavulanate oral suspension for reconstitution	T2	
amoxicillin-pot clavulanate oral tablet	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
amoxicillin-pot clavulanate oral tablet,chewable	T2	
amphotericin b	T4	PA-BvD
amphotericin b liposome	T5	PA-BvD
ampicillin oral capsule 500 mg	T2	
ampicillin sodium injection recon soln 1 gram, 10 gram, 2 gram	T2	
ampicillin-sulbactam injection	T2	
APTIVUS	T5	
ARIKAYCE	T5	PA
atazanavir	T4	
atovaquone	T4	
atovaquone-proguanil	T2	
AVYCAZ	T5	
azithromycin intravenous	T2	
azithromycin oral tablet	T2	
aztreonam	T4	
BICILLIN C-R	T3	
BICILLIN L-A INTRAMUSCULAR SYRINGE 600,000 UNIT/ML	T3	
BIKTARVY	T5	QL (31 EA per 31 days)
BLUJEPA	T4	QL (20 EA per 5 days)
caspofungin	T4	
CAYSTON	T5	PA
cefaclor oral capsule 500 mg	T2	
cefadroxil oral capsule	T2	
cefadroxil oral suspension for reconstitution 250 mg/5 ml	T2	
cefadroxil oral suspension for reconstitution 500 mg/5 ml	T3	
cefadroxil oral tablet	T3	
cefazolin injection recon soln 1 gram, 500 mg	T2	
cefazolin intravenous recon soln 10 gram	T2	
cefdinir oral capsule	T2	
cefepime injection	T4	
cefixime oral capsule	T2	
cefixime oral suspension for reconstitution	T2	
cefotetan injection	T2	
cefoxitin	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
cefpodoxime	T2	
cefprozil	T2	
ceftaroline fosamil	T5	
ceftazidime	T4	
ceftriaxone injection recon soln 1 gram, 10 gram, 2 gram, 250 mg, 500 mg	T2	
cefuroxime axetil oral tablet	T2	
cefuroxime sodium injection recon soln 750 mg	T2	
cefuroxime sodium intravenous recon soln 1.5 gram	T2	
cephalexin	T2	
chloroquine phosphate oral tablet 250 mg	T3	QL (50 EA per 30 days)
chloroquine phosphate oral tablet 500 mg	T3	QL (25 EA per 30 days)
CIMDUO	T5	QL (31 EA per 31 days)
ciprofloxacin hcl oral tablet 250 mg, 500 mg, 750 mg	T1	
ciprofloxacin in 5 % dextrose intravenous piggyback 200 mg/100 ml	T2	
clarithromycin oral suspension for reconstitution 125 mg/5 ml	T2	
clarithromycin oral suspension for reconstitution 250 mg/5 ml	T4	
clarithromycin oral tablet	T2	
clarithromycin oral tablet extended release 24 hr	T2	
clindamycin hcl	T2	
clindamycin in 5 % dextrose	T2	
CLINDAMYCIN PEDIATRIC	T2	
clindamycin phosphate injection	T2	
clotrimazole mucous membrane	T2	
COARTEM	T4	
colistin (colistimethate na)	T4	
CRESEMBA ORAL	T5	
cycloserine	T4	
dalbavancin	T5	
dapsone oral	T3	
daptomycin intravenous recon soln 350 mg	T4	
daptomycin intravenous recon soln 500 mg	T5	
darunavir oral tablet 600 mg	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
darunavir oral tablet 800 mg	T5	
DELSTRIGO	T5	QL (31 EA per 31 days)
demeclocycline oral tablet 150 mg	T2	
demeclocycline oral tablet 300 mg	T4	
DESCOVY	T5	QL (31 EA per 31 days)
dicloxacillin	T2	
DIFICID ORAL SUSPENSION FOR RECONSTITUTION	T5	QL (136 ML per 12 days)
DOVATO	T5	QL (31 EA per 31 days)
DOXY-100	T3	
doxycycline hyclate intravenous	T2	
doxycycline hyclate oral capsule 100 mg, 50 mg	T2	
doxycycline hyclate oral tablet 100 mg	T2	
doxycycline hyclate oral tablet 20 mg	T1	
doxycycline hyclate oral tablet 50 mg	T4	
doxycycline hyclate oral tablet,delayed release (dr/ec) 100 mg	T2	
doxycycline hyclate oral tablet,delayed release (dr/ec) 150 mg, 200 mg, 50 mg	T3	
doxycycline hyclate oral tablet,delayed release (dr/ec) 75 mg	T1	
doxycycline monohydrate oral capsule 100 mg, 50 mg	T2	
doxycycline monohydrate oral suspension for reconstitution	T4	
doxycycline monohydrate oral tablet 100 mg, 50 mg, 75 mg	T2	
EDURANT PED	T5	
efavirenz oral tablet	T3	
efavirenz-emtricitabin-tenofov	T5	
efavirenz-lamivu-tenofov disop	T5	QL (31 EA per 31 days)
emtricitabine	T4	
emtricitabine-tenofovir (tdf) oral tablet 100-150 mg, 167-250 mg, 200-300 mg	T4	
emtricitabine-tenofovir (tdf) oral tablet 133-200 mg	T5	
emtricita-rilpivirine-tenof df	T5	
EMTRIVA ORAL SOLUTION	T3	
entecavir	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ERAXIS(WATER DILUENT) INTRAVENOUS RECON SOLN 100 MG	T5	
ERAXIS(WATER DILUENT) INTRAVENOUS RECON SOLN 50 MG	T4	
ertapenem	T4	
erythromycin oral tablet	T4	
ethambutol	T2	
etravirine	T5	
EVOTAZ	T5	
famciclovir	T2	
fidaxomicin	T5	QL (20 EA per 10 days)
fluconazole in nacl (iso-osm) intravenous piggyback 200 mg/100 ml	T2	
fluconazole in nacl (iso-osm) intravenous piggyback 400 mg/200 ml	T3	
fluconazole oral suspension for reconstitution	T3	
fluconazole oral tablet	T2	
flucytosine	T5	
fosamprenavir	T5	
fosfomycin tromethamine	T4	
gentamicin in nacl (iso-osm) intravenous piggyback 100 mg/100 ml, 60 mg/50 ml, 80 mg/100 ml, 80 mg/50 ml	T2	
gentamicin injection	T1	
GENVOYA	T5	
griseofulvin microsize oral suspension	T2	
griseofulvin microsize oral tablet	T4	
griseofulvin ultramicrosize oral tablet 125 mg, 250 mg	T4	
HEPCLUDEX	T5	PA; QL (31 EA per 31 days)
hydroxychloroquine oral tablet 100 mg	T4	QL (93 EA per 31 days)
hydroxychloroquine oral tablet 200 mg	T2	QL (93 EA per 31 days)
hydroxychloroquine oral tablet 300 mg	T4	QL (62 EA per 31 days)
hydroxychloroquine oral tablet 400 mg	T4	QL (31 EA per 31 days)
IDVYNSO	T5	QL (31 EA per 31 days)
imipenem-cilastatin	T2	
INTELENCE ORAL TABLET 25 MG	T4	
ISENTRESS HD	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ISENTRESS ORAL POWDER IN PACKET	T5	
ISENTRESS ORAL TABLET	T5	
ISENTRESS ORAL TABLET,CHEWABLE 100 MG	T5	
ISENTRESS ORAL TABLET,CHEWABLE 25 MG	T3	
isoniazid oral solution	T2	
isoniazid oral tablet	T1	
itraconazole oral capsule	T2	PA
ivermectin oral	T2	PA
JULUCA	T5	
ketoconazole oral	T2	
LAGEVRIO (EUA)	T3	
lamivudine	T2	
lamivudine-zidovudine	T2	
levofloxacin in d5w intravenous piggyback 500 mg/100 ml, 750 mg/150 ml	T2	
levofloxacin oral solution	T3	
levofloxacin oral tablet	T2	
linezolid in dextrose 5%	T4	
linezolid oral tablet	T4	
LIVTENCITY	T5	PA; QL (372 EA per 31 days)
lopinavir-ritonavir oral tablet	T4	
maraviroc oral tablet 150 mg	T5	
maraviroc oral tablet 300 mg	T3	
MAVYRET ORAL PELLETS IN PACKET	T5	PA; QL (140 EA per 28 days)
MAVYRET ORAL TABLET	T5	PA; QL (84 EA per 28 days)
mefloquine	T2	
meropenem intravenous recon soln 1 gram, 500 mg	T3	
methenamine hippurate	T2	
metronidazole in nacl (iso-os)	T2	
metronidazole oral tablet 250 mg, 500 mg	T1	
micafungin	T4	
minocycline oral capsule	T2	
minocycline oral tablet	T2	
moxifloxacin oral	T3	
moxifloxacin-sod.chloride(iso)	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
nafcillin injection recon soln 1 gram	T2	
nafcillin injection recon soln 10 gram	T5	
nafcillin injection recon soln 2 gram	T4	
neomycin	T2	
nevirapine oral suspension	T2	
nevirapine oral tablet	T2	
nevirapine oral tablet extended release 24 hr	T4	
nitazoxanide	T5	
nitrofurantoin macrocrystal oral capsule 100 mg	T2	QL (90 EA per 365 days)
nitrofurantoin macrocrystal oral capsule 25 mg	T2	QL (360 EA per 365 days)
nitrofurantoin macrocrystal oral capsule 50 mg	T2	QL (180 EA per 365 days)
nitrofurantoin monohyd/m-cryst	T2	QL (90 EA per 365 days)
NUZYRA	T5	
nystatin oral suspension	T2	
nystatin oral tablet	T2	
ODEFSEY	T5	QL (31 EA per 31 days)
ofloxacin oral tablet 400 mg	T2	
ORLYNVAH	T4	QL (10 EA per 5 days)
oseltamivir oral capsule 30 mg	T2	QL (170 EA per 365 days)
oseltamivir oral capsule 45 mg, 75 mg	T2	QL (90 EA per 365 days)
oseltamivir oral suspension for reconstitution	T3	QL (1080 ML per 365 days)
oxacillin injection recon soln 1 gram, 2 gram	T2	
oxacillin injection recon soln 10 gram	T4	
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (10)- 100 MG (10)	T2	QL (180 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 150 MG (6)- 100 MG (5)	T2	QL (99 EA per 365 days)
PAXLOVID ORAL TABLETS,DOSE PACK 300 MG (150 MG X 2)-100 MG	T2	QL (270 EA per 365 days)
penicillin g pot in dextrose intravenous piggyback 2 million unit/50 ml, 3 million unit/50 ml	T4	
penicillin g potassium injection recon soln 20 million unit	T2	
penicillin g sodium	T5	
penicillin v potassium	T1	
pentamidine inhalation	T4	PA-BvD
pentamidine injection	T4	
PIFELTRO	T5	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
piperacillin-tazobactam intravenous recon soln 2.25 gram, 3.375 gram, 4.5 gram, 40.5 gram	T4	
polymyxin b sulfate	T2	
posaconazole oral tablet,delayed release (dr/ec)	T5	PA
praziquantel	T4	
PREVYMIS ORAL PELLETS IN PACKET	T5	PA; QL (124 EA per 31 days)
PREVYMIS ORAL TABLET	T5	QL (31 EA per 31 days)
PREZCOBIX	T5	
PREZISTA ORAL SUSPENSION	T5	
PREZISTA ORAL TABLET 150 MG	T5	
PREZISTA ORAL TABLET 75 MG	T4	
primaquine	T3	
pyrazinamide	T4	
pyrimethamine	T5	PA
quinine sulfate	T3	PA; QL (42 EA per 28 days)
RELENZA DISKHALER	T3	
REYATAZ ORAL POWDER IN PACKET	T5	
ribavirin oral capsule	T3	
ribavirin oral tablet 200 mg	T2	
rifabutin	T4	
rifampin intravenous	T4	
rifampin oral	T2	
rifapentine	T3	
rilpivirine hcl	T5	
rimantadine	T2	
ritonavir	T3	
RUKOBIA	T5	QL (62 EA per 31 days)
SIRTURO	T5	PA
sofosbuvir-velpatasvir	T5	PA; QL (28 EA per 28 days)
streptomycin	T5	
STRIBILD	T5	
sulfadiazine	T2	
sulfamethoxazole-trimethoprim oral	T1	
SUNLENCA	T5	
SYMTUZA	T5	QL (31 EA per 31 days)
tenofovir disoproxil fumarate	T3	
terbinafine hcl oral	T1	QL (90 EA per 180 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
tetracycline oral capsule	T4	
tigecycline	T4	
tinidazole	T2	
TIVICAY ORAL TABLET 50 MG	T5	
TIVICAY PD	T5	
TOBI PODHALER	T5	PA; QL (224 EA per 56 days)
tobramycin in 0.225 % nacl	T5	PA
tobramycin inhalation	T5	PA
tobramycin sulfate injection solution	T1	
trimethoprim	T2	
TRIUMEQ	T5	
TRIUMEQ PD	T4	QL (186 EA per 31 days)
valacyclovir	T2	
valganciclovir oral recon soln	T4	
valganciclovir oral tablet	T3	
vancomycin intravenous recon soln 1,000 mg, 10 gram, 500 mg, 750 mg	T4	
vancomycin oral capsule 125 mg	T4	QL (124 EA per 31 days)
vancomycin oral capsule 250 mg	T4	QL (248 EA per 31 days)
VEMLIDY	T5	QL (31 EA per 31 days)
VIRACEPT ORAL TABLET	T5	
VIREAD ORAL POWDER	T5	
VIREAD ORAL TABLET 150 MG, 200 MG, 250 MG	T5	
VIVJOA	T4	PA; QL (18 EA per 84 days)
voriconazole intravenous recon soln	T5	PA
voriconazole oral	T4	
VOSEVI	T5	PA; QL (28 EA per 28 days)
XIFAXAN ORAL TABLET 200 MG	T4	QL (27 EA per 365 days)
XIFAXAN ORAL TABLET 550 MG	T5	PA; QL (62 EA per 31 days)
XOCOVA	T5	QL (63 EA per 365 days)
XOFLUZA ORAL TABLET 40 MG, 80 MG	T3	QL (9 EA per 365 days)
zidovudine	T2	
ZOSYN IN DEXTROSE (ISO-OSM) INTRAVENOUS PIGGYBACK 2.25 GRAM/50 ML	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
Antineoplastic / Immunosuppressant Drugs		
abiraterone oral tablet 250 mg	T5	PA-NS; QL (124 EA per 31 days)
abiraterone oral tablet 500 mg	T5	PA-NS; QL (62 EA per 31 days)
ABIRTEGA	T3	PA-NS; QL (124 EA per 31 days)
AKEEGA	T5	PA-NS; QL (62 EA per 31 days)
ALECENSA	T5	PA-NS; QL (248 EA per 31 days)
ALUNBRIG ORAL TABLET 180 MG, 90 MG	T5	PA-NS; QL (31 EA per 31 days)
ALUNBRIG ORAL TABLET 30 MG	T5	PA-NS; QL (186 EA per 31 days)
ALUNBRIG ORAL TABLETS,DOSE PACK	T5	PA-NS; QL (60 EA per 365 days)
anastrozole	T2	
AUGTYRO ORAL CAPSULE 160 MG	T5	PA-NS; QL (62 EA per 31 days)
AUGTYRO ORAL CAPSULE 40 MG	T5	PA-NS; QL (186 EA per 31 days)
AVMAPKI-FAKZYNJA	T5	PA-NS; QL (66 EA per 28 days)
AYVAKIT	T5	PA-NS; QL (31 EA per 31 days)
azathioprine oral tablet 100 mg, 75 mg	T4	PA-BvD
azathioprine oral tablet 50 mg	T2	PA-BvD
BALVERSA	T5	PA-NS
bexarotene oral	T5	PA-NS
bexarotene topical	T5	PA-NS; QL (60 GM per 28 days)
bicalutamide	T2	
BOSULIF ORAL CAPSULE 100 MG	T5	PA-NS; QL (186 EA per 31 days)
BOSULIF ORAL CAPSULE 50 MG	T5	PA-NS; QL (341 EA per 31 days)
BOSULIF ORAL TABLET 100 MG	T5	PA-NS; QL (93 EA per 31 days)
BOSULIF ORAL TABLET 400 MG, 500 MG	T5	PA-NS; QL (31 EA per 31 days)
bosutinib oral tablet 400 mg	T5	PA-NS; QL (31 EA per 31 days)
BRAFTOVI	T5	PA-NS; QL (186 EA per 31 days)
BRUKINSA ORAL TABLET	T5	PA-NS; QL (62 EA per 31 days)
CABOMETYX	T5	PA-NS; QL (31 EA per 31 days)
CALQUENCE (ACALABRUTINIB MAL)	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
CAPRELSA ORAL TABLET 300 MG	T5	PA-NS; QL (31 EA per 31 days)
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
COMETRIQ ORAL CAPSULE 140 MG/DAY(80 MG X1-20 MG X3)	T5	PA-NS; QL (112 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
COMETRIQ ORAL CAPSULE 60 MG/DAY (20 MG X 3/DAY)	T5	PA-NS; QL (84 EA per 28 days)
COPIKTRA	T5	PA-NS; QL (62 EA per 31 days)
COTELLIC	T5	PA-NS; LA; QL (63 EA per 28 days)
cyclophosphamide oral capsule	T3	PA-BvD
cyclophosphamide oral tablet 50 mg	T3	PA-BvD
cyclosporine modified oral capsule	T2	PA-BvD
cyclosporine modified oral solution	T5	PA-BvD
cyclosporine oral capsule	T2	PA-BvD
DANZITEN	T5	PA-NS; QL (124 EA per 31 days)
dasatinib oral tablet 100 mg, 140 mg, 50 mg, 70 mg, 80 mg	T5	PA-NS; QL (31 EA per 31 days)
dasatinib oral tablet 20 mg	T5	PA-NS; QL (93 EA per 31 days)
DAURISMO ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
DAURISMO ORAL TABLET 25 MG	T5	PA-NS; QL (62 EA per 31 days)
ELIGARD	T4	QL (1 EA per 30 days)
ELIGARD (3 MONTH)	T4	QL (1 EA per 90 days)
ELIGARD (4 MONTH)	T4	QL (1 EA per 120 days)
ELIGARD (6 MONTH)	T4	QL (1 EA per 180 days)
ENSACOVE	T5	PA-NS; QL (62 EA per 31 days)
ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HR 0.75 MG, 1 MG	T4	PA-BvD
ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HR 4 MG	T5	PA-BvD
ERIVEDGE	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 240 MG	T5	PA-NS; QL (31 EA per 31 days)
ERLEADA ORAL TABLET 60 MG	T5	PA-NS; QL (93 EA per 31 days)
erlotinib	T5	PA-NS; QL (31 EA per 31 days)
EULEXIN	T4	
everolimus (antineoplastic) oral tablet	T5	PA-NS; QL (31 EA per 31 days)
everolimus (antineoplastic) oral tablet for suspension 2 mg, 5 mg	T5	PA-NS; QL (62 EA per 31 days)
everolimus (antineoplastic) oral tablet for suspension 3 mg	T5	PA-NS; QL (93 EA per 31 days)
everolimus (immunosuppressive) oral tablet 0.25 mg	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
everolimus (immunosuppressive) oral tablet 0.5 mg, 0.75 mg, 1 mg	T5	PA-BvD
exemestane	T2	
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 120 MG	T5	PA-NS
FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS RECON SOLN 80 MG	T4	PA-NS
FOTIVDA	T5	PA-NS; QL (21 EA per 28 days)
FRUZAQLA ORAL CAPSULE 1 MG	T5	PA-NS; QL (84 EA per 28 days)
FRUZAQLA ORAL CAPSULE 5 MG	T5	PA-NS; QL (21 EA per 28 days)
GAVRETO	T5	PA-NS; QL (124 EA per 31 days)
gefitinib	T5	PA-NS; QL (31 EA per 31 days)
GENGRAF ORAL CAPSULE	T2	PA-BvD
GILOTRIF	T5	PA-NS; QL (31 EA per 31 days)
GOMEKLI ORAL CAPSULE 1 MG	T5	PA-NS; QL (126 EA per 28 days)
GOMEKLI ORAL CAPSULE 2 MG	T5	PA-NS; QL (84 EA per 28 days)
GOMEKLI ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (168 EA per 28 days)
HERNEXEOS	T5	PA-NS; QL (93 EA per 31 days)
hydroxyurea	T2	
HYRNUO	T5	PA-NS; QL (124 EA per 31 days)
IBRANCE ORAL CAPSULE 125 MG	T5	PA-NS; QL (21 EA per 28 days)
IBRANCE ORAL TABLET	T5	PA-NS; QL (21 EA per 28 days)
IBTROZI	T5	PA-NS; QL (93 EA per 31 days)
ICLUSIG	T5	PA-NS; QL (31 EA per 31 days)
IDHIFA	T5	PA-NS; QL (31 EA per 31 days)
imatinib oral tablet 100 mg	T5	PA-NS; QL (93 EA per 31 days)
imatinib oral tablet 400 mg	T5	PA-NS; QL (62 EA per 31 days)
IMBRUVICA ORAL CAPSULE 140 MG	T5	PA-NS; QL (124 EA per 31 days)
IMBRUVICA ORAL CAPSULE 70 MG	T5	PA-NS; QL (31 EA per 31 days)
IMBRUVICA ORAL SUSPENSION	T5	PA-NS; QL (216 ML per 25 days)
IMBRUVICA ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
IMKELDI	T5	PA-NS; QL (280 ML per 28 days)
INLURIYO	T5	PA-NS; QL (62 EA per 31 days)
INLYTA	T5	PA-NS; QL (124 EA per 31 days)
INQOVI	T5	PA-NS; QL (5 EA per 28 days)
INREBIC	T5	PA-NS; QL (124 EA per 31 days)
ITOVEBI ORAL TABLET 3 MG	T5	PA-NS; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ITOVEBI ORAL TABLET 9 MG	T5	PA-NS; QL (31 EA per 31 days)
IWILFIN	T5	PA-NS; QL (248 EA per 31 days)
JAKAFI	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 100 MG	T5	PA-NS; QL (62 EA per 31 days)
JAYPIRCA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
JIDEYTRO ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
JIDEYTRO ORAL TABLET 25 MG	T5	PA-NS; QL (93 EA per 31 days)
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (21 EA per 28 days)
KISQALI ORAL TABLET 400 MG/DAY (200 MG X 2)	T5	PA-NS; QL (42 EA per 28 days)
KISQALI ORAL TABLET 600 MG/DAY (200 MG X 3)	T5	PA-NS; QL (63 EA per 28 days)
KLISYRI (250 MG)	T5	PA
KOMZIFTI	T5	PA-NS; QL (90 EA per 30 days)
KOSELUGO ORAL CAPSULE 10 MG	T5	PA-NS; QL (279 EA per 31 days)
KOSELUGO ORAL CAPSULE 25 MG	T5	PA-NS; QL (124 EA per 31 days)
KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG	T5	PA-NS; QL (600 EA per 30 days)
KOSELUGO ORAL CAPSULE, SPRINKLE 7.5 MG	T5	PA-NS; QL (360 EA per 30 days)
KRAZATI	T5	PA-NS; QL (186 EA per 31 days)
lapatinib	T5	PA-NS; QL (186 EA per 31 days)
LAZCLUZE ORAL TABLET 240 MG	T5	PA-NS; QL (30 EA per 30 days)
LAZCLUZE ORAL TABLET 80 MG	T5	PA-NS; QL (60 EA per 30 days)
lenalidomide	T5	PA-NS; QL (21 EA per 28 days)
LENVIMA	T5	PA-NS
letrozole	T2	
leucovorin calcium oral	T2	
LEUKERAN	T5	
leuprolide subcutaneous kit	T4	QL (2 EA per 28 days)
LIFYORLI ORAL CAPSULE 125 MG/DAY(100 MG X1-25MG X1), 125 MG/DAY(100 MG X1-25MG X3), 125 MG/DAY(100 MG X1-25MG X9)	T5	PA-NS; QL (18 EA per 28 days)
LIFYORLI ORAL CAPSULE 150 MG/DAY(100 MG X1-25MG X2), 150 MG/DAY(100 MG X1-25MG X3), 150 MG/DAY(100 MG X1-25MG X9)	T5	PA-NS; QL (27 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
lomustine oral capsule 10 mg, 40 mg	T4	PA-NS
lomustine oral capsule 100 mg	T5	PA-NS
LONSURF	T5	PA-NS
LORBRENA ORAL TABLET 100 MG	T5	PA-NS; QL (31 EA per 31 days)
LORBRENA ORAL TABLET 25 MG	T5	PA-NS; QL (93 EA per 31 days)
LUMAKRAS ORAL TABLET 120 MG, 240 MG	T5	PA-NS; QL (62 EA per 31 days)
LUMAKRAS ORAL TABLET 320 MG	T5	PA-NS; QL (93 EA per 31 days)
LUPRON DEPOT (3 MONTH) INTRAMUSCULAR SYRINGE KIT 11.25 MG	T5	QL (1 EA per 90 days)
LUPRON DEPOT INTRAMUSCULAR SYRINGE KIT 3.75 MG	T5	QL (1 EA per 30 days)
LYNPARZA	T5	PA-NS; QL (124 EA per 31 days)
LYSODREN	T5	
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3)	T5	PA-NS; QL (93 EA per 31 days)
LYTGOBI ORAL TABLET 16 MG/DAY (4 MG X 4)	T5	PA-NS; QL (124 EA per 31 days)
LYTGOBI ORAL TABLET 20 MG/DAY (4 MG X 5)	T5	PA-NS; QL (155 EA per 31 days)
MATULANE	T5	
megestrol oral suspension 400 mg/10 ml (40 mg/ml)	T2	PA
megestrol oral suspension 625 mg/5 ml (125 mg/ml)	T4	PA
megestrol oral tablet	T2	PA-NS
MEKINIST ORAL RECON SOLN	T5	PA-NS; QL (1260 ML per 31 days)
MEKINIST ORAL TABLET 0.5 MG	T5	PA-NS; QL (93 EA per 31 days)
MEKINIST ORAL TABLET 2 MG	T5	PA-NS; QL (31 EA per 31 days)
MEKTOVI	T5	PA-NS; QL (186 EA per 31 days)
mercaptopurine oral suspension	T5	
mercaptopurine oral tablet	T2	
mesna oral	T3	
methotrexate sodium (pf) injection solution	T2	PA-BvD
methotrexate sodium injection	T2	PA-BvD
methotrexate sodium oral	T1	PA-BvD
MODEYSO	T5	PA-NS; QL (20 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
mycophenolate mofetil oral capsule	T2	PA-BvD
mycophenolate mofetil oral suspension for reconstitution	T5	PA-BvD
mycophenolate mofetil oral tablet	T2	PA-BvD
mycophenolate sodium	T2	PA-BvD
NEMLUVIO	T5	PA; QL (2 EA per 28 days)
NERLYNX	T5	PA-NS; QL (186 EA per 31 days)
nilotinib hcl	T5	PA-NS; QL (124 EA per 31 days)
nilutamide	T5	
NINLARO	T5	PA-NS; QL (3 EA per 28 days)
NUBEQA	T5	PA-NS; QL (124 EA per 31 days)
octreotide acetate injection solution 1,000 mcg/ml	T3	PA
octreotide acetate injection solution 100 mcg/ml, 50 mcg/ml	T2	PA
octreotide acetate injection solution 200 mcg/ml, 500 mcg/ml	T4	PA
ODOMZO	T5	PA-NS; LA; QL (31 EA per 31 days)
OGSIVEO ORAL TABLET 100 MG, 150 MG	T5	PA-NS; QL (62 EA per 31 days)
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION	T5	PA-NS; QL (96 ML per 28 days)
OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4)	T5	PA-NS; QL (16 EA per 28 days)
OJEMDA ORAL TABLET 500 MG/WEEK (100 MG X 5)	T5	PA-NS; QL (20 EA per 28 days)
OJEMDA ORAL TABLET 600 MG/WEEK (100 MG X 6)	T5	PA-NS; QL (24 EA per 28 days)
OJJAARA	T5	PA-NS; QL (31 EA per 31 days)
ONUREG	T5	PA-NS; QL (14 EA per 28 days)
ORGOVYX	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 345 MG	T5	PA-NS; QL (31 EA per 31 days)
ORSERDU ORAL TABLET 86 MG	T5	PA-NS; QL (93 EA per 31 days)
pazopanib oral tablet 200 mg	T5	PA-NS; QL (124 EA per 31 days)
PEMAZYRE	T5	PA-NS; QL (14 EA per 21 days)
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1)	T5	PA-NS; QL (28 EA per 28 days)
PIQRAY ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)	T5	PA-NS; QL (56 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
pomalidomide	T5	PA-NS; QL (21 EA per 28 days)
PROGRAF ORAL GRANULES IN PACKET	T4	PA-BvD
QINLOCK	T5	PA-NS; QL (93 EA per 31 days)
RETEVMO ORAL TABLET 120 MG, 160 MG, 80 MG	T5	PA-NS; QL (62 EA per 31 days)
RETEVMO ORAL TABLET 40 MG	T5	PA-NS; QL (93 EA per 31 days)
REVUFORJ ORAL TABLET 110 MG	T5	PA-NS; QL (124 EA per 31 days)
REVUFORJ ORAL TABLET 160 MG	T5	PA-NS; QL (62 EA per 31 days)
REVUFORJ ORAL TABLET 25 MG	T5	PA-NS; QL (248 EA per 31 days)
REZLIDHIA	T5	PA-NS; QL (62 EA per 31 days)
ROMVIMZA	T5	PA-NS; QL (8 EA per 28 days)
ROZLYTREK ORAL CAPSULE 100 MG	T5	PA-NS; QL (155 EA per 31 days)
ROZLYTREK ORAL CAPSULE 200 MG	T5	PA-NS; QL (93 EA per 31 days)
ROZLYTREK ORAL PELLETS IN PACKET	T5	PA-NS; QL (372 EA per 31 days)
RUBRACA	T5	PA-NS; QL (124 EA per 31 days)
RYDAPT	T5	PA-NS; QL (248 EA per 31 days)
SCEMBLIX ORAL TABLET 100 MG	T5	PA-NS; QL (124 EA per 31 days)
SCEMBLIX ORAL TABLET 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SCEMBLIX ORAL TABLET 40 MG	T5	PA-NS; QL (248 EA per 31 days)
SIGNIFOR	T5	PA
sirolimus oral solution	T4	PA-BvD
sirolimus oral tablet	T2	PA-BvD
SOLTAMOX	T4	
sorafenib	T5	PA-NS; QL (124 EA per 31 days)
STIVARGA	T5	PA-NS; QL (84 EA per 28 days)
sunitinib malate	T5	PA-NS; QL (31 EA per 31 days)
TABLOID	T4	
TABRECTA	T5	PA-NS; QL (124 EA per 31 days)
tacrolimus oral capsule	T2	PA-BvD
tacrolimus oral capsule,extended release 24hr 0.5 mg, 1 mg	T3	PA-BvD
tacrolimus oral capsule,extended release 24hr 5 mg	T5	PA-BvD
TAFINLAR ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
TAFINLAR ORAL TABLET FOR SUSPENSION	T5	PA-NS; QL (930 EA per 31 days)
TAGRISSO	T5	PA-NS; LA; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TALZENNA	T5	PA-NS; QL (31 EA per 31 days)
tamoxifen	T1	
TEPMETKO	T5	PA-NS; QL (62 EA per 31 days)
THALOMID ORAL CAPSULE 100 MG	T5	PA-NS; QL (112 EA per 28 days)
THALOMID ORAL CAPSULE 50 MG	T5	PA-NS; QL (28 EA per 28 days)
TIBSOVO	T5	PA-NS; QL (62 EA per 31 days)
toremifene	T4	
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 11.25 MG	T4	QL (1 EA per 84 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 22.5 MG	T4	QL (1 EA per 168 days)
TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 3.75 MG	T4	QL (1 EA per 28 days)
tretinoin (antineoplastic)	T5	
TREXALL	T3	PA-BvD
TRUQAP	T5	PA-NS; QL (64 EA per 28 days)
TUKYSA ORAL TABLET 150 MG	T5	PA-NS; QL (124 EA per 31 days)
TUKYSA ORAL TABLET 50 MG	T5	PA-NS; QL (248 EA per 31 days)
TURALIO	T5	PA-NS; QL (124 EA per 31 days)
VANFLYTA	T5	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 10 MG	T3	PA-NS; QL (62 EA per 31 days)
VENCLEXTA ORAL TABLET 100 MG	T5	PA-NS; QL (186 EA per 31 days)
VENCLEXTA ORAL TABLET 50 MG	T5	PA-NS; QL (31 EA per 31 days)
VENCLEXTA STARTING PACK	T5	PA-NS; QL (84 EA per 365 days)
VERZENIO	T5	PA-NS; QL (62 EA per 31 days)
VIJOICE ORAL GRANULES IN PACKET	T5	PA-NS; QL (31 EA per 31 days)
VIJOICE ORAL TABLET 125 MG, 50 MG	T5	PA-NS; QL (28 EA per 28 days)
VIJOICE ORAL TABLET 250 MG/DAY (200 MG X1-50 MG X1)	T5	PA-NS; QL (56 EA per 28 days)
VITRAKVI ORAL CAPSULE 100 MG	T5	PA-NS; QL (62 EA per 31 days)
VITRAKVI ORAL CAPSULE 25 MG	T5	PA-NS; QL (186 EA per 31 days)
VITRAKVI ORAL SOLUTION	T5	PA-NS; QL (310 ML per 31 days)
VONJO	T5	PA-NS; QL (124 EA per 31 days)
VORANIGO ORAL TABLET 10 MG	T5	PA-NS; QL (62 EA per 31 days)
VORANIGO ORAL TABLET 40 MG	T5	PA-NS; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
WELIREG	T5	PA-NS; QL (93 EA per 31 days)
WYOST	T3	PA-NS
XALKORI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XALKORI ORAL PELLET 150 MG	T5	PA-NS; QL (186 EA per 31 days)
XALKORI ORAL PELLET 20 MG, 50 MG	T5	PA-NS; QL (124 EA per 31 days)
XATMEP	T4	PA-BvD
XERMELO	T5	PA; QL (93 EA per 31 days)
XGEVA	T5	PA-NS
XOSPATA	T5	PA-NS; QL (124 EA per 31 days)
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40MG TWICE WEEK (40 MG X 2), 80 MG/WEEK (40 MG X 2)	T5	PA-NS; QL (8 EA per 28 days)
XPOVIO ORAL TABLET 40 MG/WEEK (10 MG X 4)	T5	PA-NS; QL (16 EA per 28 days)
XPOVIO ORAL TABLET 60 MG/WEEK (60 MG X 1), 80 MG/WEEK (80 MG X 1)	T5	PA-NS; QL (4 EA per 28 days)
XPOVIO ORAL TABLET 60MG TWICE WEEK (120 MG/WEEK)	T5	PA-NS; QL (24 EA per 28 days)
XPOVIO ORAL TABLET 80MG TWICE WEEK (160 MG/WEEK)	T5	PA-NS; QL (32 EA per 28 days)
XTANDI ORAL CAPSULE	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 40 MG	T5	PA-NS; QL (124 EA per 31 days)
XTANDI ORAL TABLET 80 MG	T5	PA-NS; QL (62 EA per 31 days)
XTRENBO	T3	PA-NS
YONSA	T5	PA-NS; QL (124 EA per 31 days)
YULITHIRA	T3	PA-NS; QL (31 EA per 31 days)
ZEJULA	T5	PA-NS; QL (31 EA per 31 days)
ZELBORAF	T5	PA-NS; QL (248 EA per 31 days)
ZENBEXUS	T5	PA-NS; QL (21 EA per 28 days)
ZOLINZA	T5	PA-NS
ZYKADIA	T5	PA-NS; QL (93 EA per 31 days)
Autonomic / Cns Drugs, Neurology / Psych		
ABILIFY MAINTENA	T5	QL (1 EA per 28 days)
acetaminophen-codeine oral solution 120-12 mg/5 ml	T2	PA; QL (5167 ML per 31 days)
acetaminophen-codeine oral tablet	T2	PA; QL (403 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML	T3	PA; QL (1 ML per 28 days)
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 70 MG/ML	T3	PA; QL (2 ML per 28 days)
almotriptan malate oral tablet 12.5 mg	T2	QL (8 EA per 28 days)
almotriptan malate oral tablet 6.25 mg	T2	QL (16 EA per 28 days)
ALPRAZOLAM INTENSOL	T4	PA
alprazolam oral tablet 0.25 mg, 0.5 mg	T2	PA; QL (93 EA per 31 days)
alprazolam oral tablet 1 mg, 2 mg	T2	PA; QL (155 EA per 31 days)
alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg	T2	PA; QL (31 EA per 31 days)
alprazolam oral tablet extended release 24 hr 2 mg	T2	PA; QL (155 EA per 31 days)
alprazolam oral tablet extended release 24 hr 3 mg	T2	PA; QL (93 EA per 31 days)
amitriptyline	T2	PA-NS
amitriptyline-chlordiazepoxide	T4	PA-NS
amoxapine	T1	
aripiprazole oral solution	T4	PA-NS
aripiprazole oral tablet	T3	
aripiprazole oral tablet,disintegrating 10 mg	T4	PA-NS
aripiprazole oral tablet,disintegrating 15 mg	T5	PA-NS
ARISTADA INITIO	T5	QL (4.8 ML per 365 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 1,064 MG/3.9 ML	T5	QL (3.9 ML per 28 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 441 MG/1.6 ML	T5	QL (1.6 ML per 28 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 662 MG/2.4 ML	T5	QL (2.4 ML per 28 days)
ARISTADA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 882 MG/3.2 ML	T5	QL (3.2 ML per 28 days)
armodafinil	T4	PA; QL (31 EA per 31 days)
asenapine maleate	T4	PA-NS; QL (62 EA per 31 days)
atomoxetine oral capsule 10 mg, 25 mg, 40 mg	T3	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
atomoxetine oral capsule 100 mg, 60 mg, 80 mg	T3	QL (31 EA per 31 days)
atomoxetine oral capsule 18 mg	T3	QL (124 EA per 31 days)
AUSTEDO ORAL TABLET 12 MG, 6 MG	T5	PA; QL (124 EA per 31 days)
AUSTEDO ORAL TABLET 9 MG	T5	PA; QL (155 EA per 31 days)
AUSTEDO XR	T5	PA; QL (31 EA per 31 days)
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG	T5	PA; QL (56 EA per 365 days)
AUVELITY ORAL TABLET, IR AND ER, BIPHASIC	T5	PA-NS; QL (62 EA per 31 days)
baclofen oral tablet 10 mg	T1	
baclofen oral tablet 20 mg	T2	
BAFIERTAM	T5	PA; QL (124 EA per 31 days)
benztropine oral	T2	PA
brivaracetam oral solution	T5	QL (620 ML per 31 days)
brivaracetam oral tablet	T5	QL (62 EA per 31 days)
bromocriptine	T4	
buprenorphine	T4	PA; QL (4 EA per 28 days)
buprenorphine hcl sublingual	T3	
buprenorphine-naloxone sublingual film	T2	
buprenorphine-naloxone sublingual tablet	T4	
bupropion hcl oral tablet	T2	
bupropion hcl oral tablet extended release 24 hr 150 mg	T2	QL (93 EA per 31 days)
bupropion hcl oral tablet extended release 24 hr 300 mg	T2	QL (31 EA per 31 days)
bupropion hcl oral tablet sustained-release 12 hr	T2	QL (62 EA per 31 days)
buspirone oral tablet 10 mg, 15 mg, 30 mg, 5 mg, 7.5 mg	T2	
butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg	T4	PA; QL (403 EA per 31 days)
butalbital-acetaminop-caf-cod oral capsule 50-325-40-30 mg	T4	PA; QL (372 EA per 31 days)
butalbital-acetaminophen oral tablet 50-325 mg	T2	QL (372 EA per 31 days)
butalbital-acetaminophen-caff oral capsule 50-300-40 mg	T4	QL (403 EA per 31 days)
butalbital-acetaminophen-caff oral capsule 50-325-40 mg	T4	QL (372 EA per 31 days)
butalbital-acetaminophen-caff oral tablet	T4	QL (372 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
butalbital-aspirin-caffeine oral capsule	T4	
butorphanol nasal	T2	QL (5 ML per 28 days)
CAPLYTA	T5	PA-NS; QL (31 EA per 31 days)
carbamazepine oral capsule, er multiphase 12 hr	T2	
carbamazepine oral suspension 100 mg/5 ml	T1	
carbamazepine oral tablet	T1	
carbamazepine oral tablet extended release 12 hr	T2	
carbamazepine oral tablet,chewable 100 mg	T1	
carbamazepine oral tablet,chewable 200 mg	T4	
carbidopa-levodopa oral tablet	T2	
carbidopa-levodopa oral tablet extended release	T2	
carbidopa-levodopa oral tablet,disintegrating	T2	
carbidopa-levodopa-entacapone	T2	
celecoxib	T2	QL (62 EA per 31 days)
chlorpromazine oral	T4	
citalopram oral solution	T3	
citalopram oral tablet	T1	
clobazam oral suspension	T4	PA-NS; QL (496 ML per 31 days)
clobazam oral tablet	T3	PA-NS; QL (62 EA per 31 days)
clomipramine	T4	PA-NS
clonazepam oral tablet 0.5 mg	T2	PA-NS; QL (93 EA per 31 days)
clonazepam oral tablet 1 mg	T2	PA-NS; QL (124 EA per 31 days)
clonazepam oral tablet 2 mg	T2	PA-NS; QL (310 EA per 31 days)
clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg	T2	PA-NS; QL (93 EA per 31 days)
clonazepam oral tablet,disintegrating 1 mg	T2	PA-NS; QL (124 EA per 31 days)
clonazepam oral tablet,disintegrating 2 mg	T2	PA-NS; QL (310 EA per 31 days)
clonidine hcl oral tablet extended release 12 hr	T4	PA
clorazepate dipotassium oral tablet 15 mg	T2	PA-NS; QL (186 EA per 31 days)
clorazepate dipotassium oral tablet 3.75 mg, 7.5 mg	T2	PA-NS; QL (93 EA per 31 days)
clozapine oral tablet 100 mg, 25 mg	T2	QL (279 EA per 31 days)
clozapine oral tablet 200 mg	T2	QL (124 EA per 31 days)
clozapine oral tablet 50 mg	T2	QL (93 EA per 31 days)
clozapine oral tablet,disintegrating 100 mg, 25 mg	T2	QL (279 EA per 31 days)
clozapine oral tablet,disintegrating 12.5 mg	T2	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
clozapine oral tablet,disintegrating 150 mg	T4	QL (186 EA per 31 days)
clozapine oral tablet,disintegrating 200 mg	T4	QL (124 EA per 31 days)
COBENFY	T5	PA-NS; QL (62 EA per 31 days)
COBENFY STARTER PACK	T5	PA-NS; QL (112 EA per 365 days)
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
COPAXONE SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
cyclobenzaprine oral tablet 10 mg, 7.5 mg	T2	QL (93 EA per 31 days)
cyclobenzaprine oral tablet 5 mg	T2	QL (155 EA per 31 days)
dalfampridine	T5	PA; QL (62 EA per 31 days)
dantrolene oral	T4	
desipramine oral tablet 10 mg, 25 mg, 50 mg	T3	
desipramine oral tablet 100 mg, 150 mg, 75 mg	T4	
desvenlafaxine oral tablet extended release 24 hr 100 mg	T4	QL (124 EA per 31 days)
desvenlafaxine oral tablet extended release 24 hr 50 mg	T4	QL (31 EA per 31 days)
desvenlafaxine succinate	T3	QL (31 EA per 31 days)
dexmethylphenidate oral capsule,er biphasic 50-50	T4	QL (31 EA per 31 days)
dexmethylphenidate oral tablet 10 mg	T2	QL (62 EA per 31 days)
dexmethylphenidate oral tablet 2.5 mg, 5 mg	T2	QL (93 EA per 31 days)
dextroamphetamine sulfate oral tablet 10 mg	T4	QL (186 EA per 31 days)
dextroamphetamine sulfate oral tablet 5 mg	T4	QL (341 EA per 31 days)
dextroamphetamine-amphetamine oral capsule,extended release 24hr	T3	QL (31 EA per 31 days)
dextroamphetamine-amphetamine oral tablet 10 mg, 15 mg, 30 mg, 5 mg, 7.5 mg	T1	QL (62 EA per 31 days)
dextroamphetamine-amphetamine oral tablet 12.5 mg	T2	QL (62 EA per 31 days)
dextroamphetamine-amphetamine oral tablet 20 mg	T1	QL (93 EA per 31 days)
dextromethorphan-quinidine	T5	PA; QL (62 EA per 31 days)
DIACOMIT ORAL CAPSULE 250 MG	T5	PA-NS; QL (341 EA per 31 days)
DIACOMIT ORAL CAPSULE 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIACOMIT ORAL POWDER IN PACKET 250 MG	T5	PA-NS; QL (341 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
DIACOMIT ORAL POWDER IN PACKET 500 MG	T5	PA-NS; QL (186 EA per 31 days)
DIAZEPAM INTENSOL	T2	PA-NS; QL (248 ML per 31 days)
diazepam oral solution 5 mg/5 ml (1 mg/ml)	T2	PA-NS; QL (1500 ML per 31 days)
diazepam oral tablet	T2	PA-NS; QL (124 EA per 31 days)
diazepam rectal	T4	
diclofenac epolamine	T4	PA; QL (62 EA per 31 days)
diclofenac potassium oral tablet 50 mg	T1	
diclofenac sodium oral	T1	
diclofenac sodium topical drops	T2	QL (450 ML per 28 days)
diclofenac-misoprostol	T3	
diflunisal	T2	
dihydroergotamine nasal	T5	PA; QL (8 ML per 28 days)
DILANTIN	T3	
DILANTIN EXTENDED	T4	
dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg	T4	PA; QL (62 EA per 31 days)
dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg (14)- 240 mg (46)	T5	PA; QL (120 EA per 365 days)
dimethyl fumarate oral capsule,delayed release(dr/ec) 240 mg	T5	PA; QL (62 EA per 31 days)
divalproex oral capsule, delayed rel sprinkle	T2	
divalproex oral tablet extended release 24 hr 250 mg	T2	
divalproex oral tablet extended release 24 hr 500 mg	T3	
divalproex oral tablet,delayed release (dr/ec)	T2	
donepezil oral tablet 10 mg, 5 mg	T1	
donepezil oral tablet 23 mg	T3	QL (31 EA per 31 days)
donepezil oral tablet,disintegrating	T2	
doxepin oral capsule	T2	PA-NS
doxepin oral concentrate	T2	PA-NS
doxepin oral tablet	T2	PA
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG	T3	PA-NS; QL (93 EA per 31 days)
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 30 MG, 60 MG	T3	PA-NS; QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 40 MG	T3	PA-NS; QL (31 EA per 31 days)
duloxetine oral capsule,delayed release(dr/ec) 20 mg, 60 mg	T2	QL (62 EA per 31 days)
duloxetine oral capsule,delayed release(dr/ec) 30 mg	T2	QL (31 EA per 31 days)
duloxetine oral capsule,delayed release(dr/ec) 40 mg	T3	QL (31 EA per 31 days)
DUOPA	T5	PA-BvD
eletriptan oral tablet 20 mg	T4	QL (12 EA per 28 days)
eletriptan oral tablet 40 mg	T4	QL (6 EA per 28 days)
EMGALITY PEN	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML	T3	PA; QL (1 ML per 28 days)
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 300 MG/3 ML (100 MG/ML X 3)	T5	PA; QL (3 ML per 28 days)
EMSAM	T5	QL (30 EA per 30 days)
entacapone	T2	
EPIDIOLEX	T5	PA-NS
EQUETRO	T4	
ergotamine-caffeine	T3	PA
escitalopram oxalate oral capsule	T4	PA-NS; QL (31 EA per 31 days)
escitalopram oxalate oral solution	T4	QL (620 ML per 31 days)
escitalopram oxalate oral tablet 10 mg	T2	QL (45 EA per 30 days)
escitalopram oxalate oral tablet 20 mg, 5 mg	T2	QL (30 EA per 30 days)
eslicarbazepine oral tablet 200 mg	T4	QL (186 EA per 31 days)
eslicarbazepine oral tablet 400 mg	T5	QL (93 EA per 31 days)
eslicarbazepine oral tablet 600 mg, 800 mg	T5	QL (62 EA per 31 days)
estazolam	T2	PA
eszopiclone	T2	PA; QL (31 EA per 31 days)
ethosuximide oral capsule	T3	
ethosuximide oral solution	T2	
etodolac oral capsule	T3	
etodolac oral tablet	T2	
etodolac oral tablet extended release 24 hr	T3	
EXXUA ORAL TABLET EXTENDED RELEASE 24 HR	T4	PA-NS; QL (31 EA per 31 days)
EXXUA ORAL TABLET, EXT REL 24HR DOSE PACK	T4	PA-NS; QL (64 EA per 365 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
FANAPT ORAL TABLET 1 MG	T4	PA-NS; QL (62 EA per 31 days)
FANAPT ORAL TABLET 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG	T5	PA-NS; QL (62 EA per 31 days)
FANAPT TITRATION PACK A	T4	PA-NS; QL (16 EA per 365 days)
FANAPT TITRATION PACK B	T4	PA-NS; QL (24 EA per 365 days)
FANAPT TITRATION PACK C	T4	PA-NS; QL (16 EA per 365 days)
felbamate	T4	
fenoprofen oral capsule 400 mg	T4	
fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 75 mcg/hr	T4	PA; QL (10 EA per 30 days)
fentanyl transdermal patch 72 hour 25 mcg/hr, 50 mcg/hr	T2	PA; QL (10 EA per 30 days)
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)	T4	PA-NS; QL (56 EA per 365 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 40 MG, 80 MG	T4	PA-NS; QL (31 EA per 31 days)
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 20 MG	T4	PA-NS; QL (93 EA per 31 days)
fingolimod	T5	PA; QL (31 EA per 31 days)
FINTEPLA	T5	PA-NS; QL (360 ML per 30 days)
fluoxetine (pmdd)	T1	
fluoxetine oral capsule	T1	
fluoxetine oral solution	T3	
fluoxetine oral tablet 10 mg, 20 mg	T1	
fluphenazine decanoate	T2	
fluphenazine hcl injection	T2	
fluphenazine hcl oral concentrate	T2	
fluphenazine hcl oral elixir	T4	
fluphenazine hcl oral tablet	T1	
flurazepam	T2	PA
flurbiprofen oral tablet 100 mg	T2	
fluvoxamine oral capsule,extended release 24hr	T4	
fluvoxamine oral tablet	T2	
frovatriptan	T4	QL (12 EA per 28 days)
gabapentin oral capsule 100 mg, 400 mg	T2	QL (270 EA per 30 days)
gabapentin oral capsule 300 mg	T2	QL (360 EA per 30 days)
gabapentin oral solution 250 mg/5 ml	T3	QL (2160 ML per 30 days)
gabapentin oral tablet 600 mg	T2	QL (180 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
gabapentin oral tablet 800 mg	T2	QL (120 EA per 30 days)
galantamine oral capsule,ext rel. pellets 24 hr	T3	
galantamine oral solution	T4	
galantamine oral tablet	T2	
glatiramer subcutaneous syringe 20 mg/ml	T5	PA; QL (31 ML per 31 days)
glatiramer subcutaneous syringe 40 mg/ml	T5	PA; QL (12 ML per 28 days)
GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (31 ML per 31 days)
GLATOPA SUBCUTANEOUS SYRINGE 40 MG/ML	T5	PA; QL (12 ML per 28 days)
guanfacine oral tablet extended release 24 hr	T2	PA
haloperidol decanoate	T2	
haloperidol lactate injection	T1	
haloperidol lactate oral	T2	
haloperidol oral tablet 0.5 mg, 1 mg, 10 mg, 2 mg, 5 mg	T1	
haloperidol oral tablet 20 mg	T2	
hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg	T2	PA; QL (372 EA per 31 days)
hydromorphone oral liquid	T4	PA; QL (1240 ML per 31 days)
hydromorphone oral tablet 2 mg, 4 mg	T2	PA; QL (186 EA per 31 days)
hydromorphone oral tablet 8 mg	T3	PA; QL (155 EA per 31 days)
IBU ORAL TABLET 600 MG, 800 MG	T1	
ibuprofen oral suspension	T1	
ibuprofen oral tablet 400 mg, 600 mg, 800 mg	T1	
imipramine hcl	T2	PA-NS
imipramine pamoate	T4	PA-NS
INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE	T5	PA; QL (300 EA per 30 days)
indomethacin oral capsule	T1	
indomethacin oral capsule, extended release	T1	
indomethacin oral suspension	T4	
INGREZZA INITIATION PK(TARDIV)	T5	PA; QL (56 EA per 365 days)
INGREZZA ORAL CAPSULE 40 MG	T5	PA; QL (62 EA per 31 days)
INGREZZA ORAL CAPSULE 60 MG, 80 MG	T5	PA; QL (31 EA per 31 days)
INGREZZA SPRINKLE	T5	PA; QL (31 EA per 31 days)
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,092 MG/3.5 ML	T5	QL (3.5 ML per 180 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
INVEGA HAFYERA INTRAMUSCULAR SYRINGE 1,560 MG/5 ML	T5	QL (5 ML per 180 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 117 MG/0.75 ML	T5	QL (0.75 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 156 MG/ML	T5	QL (1 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 234 MG/1.5 ML	T5	QL (1.5 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 39 MG/0.25 ML	T4	QL (0.25 ML per 28 days)
INVEGA SUSTENNA INTRAMUSCULAR SYRINGE 78 MG/0.5 ML	T5	QL (0.5 ML per 28 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 273 MG/0.88 ML	T5	QL (0.88 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 410 MG/1.32 ML	T5	QL (1.32 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 546 MG/1.75 ML	T5	QL (1.75 ML per 84 days)
INVEGA TRINZA INTRAMUSCULAR SYRINGE 819 MG/2.63 ML	T5	QL (2.63 ML per 84 days)
KESIMPTA PEN	T5	PA; QL (0.4 ML per 28 days)
ketoprofen oral capsule 25 mg, 50 mg	T2	
ketoprofen oral capsule,ext rel. pellets 24 hr 200 mg	T2	
ketorolac oral	T2	
KLOXXADO	T3	
lacosamide oral	T4	
lamotrigine oral tablet	T2	
lamotrigine oral tablet extended release 24hr	T4	
lamotrigine oral tablet, chewable dispersible	T2	
lamotrigine oral tablet,disintegrating	T4	
levetiracetam oral solution 100 mg/ml	T2	
levetiracetam oral tablet	T2	
levetiracetam oral tablet extended release 24 hr	T2	
lisdexamfetamine oral capsule	T4	ST; QL (31 EA per 31 days)
lithium carbonate	T1	
lithium citrate oral solution	T1	
lofexidine	T5	
LORAZEPAM INTENSOL	T2	PA; QL (155 ML per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
lorazepam oral tablet 0.5 mg	T2	PA; QL (124 EA per 31 days)
lorazepam oral tablet 1 mg	T2	PA; QL (186 EA per 31 days)
lorazepam oral tablet 2 mg	T2	PA; QL (155 EA per 31 days)
loxapine succinate	T2	
lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg	T4	PA-NS; QL (31 EA per 31 days)
lurasidone oral tablet 80 mg	T4	PA-NS; QL (62 EA per 31 days)
LYBALVI	T5	PA-NS; QL (31 EA per 31 days)
MARPLAN	T3	
mefenamic acid	T4	
meloxicam oral tablet	T1	
memantine oral capsule,sprinkle,er 24hr	T3	
memantine oral solution	T3	
memantine oral tablet	T2	
memantine-donepezil	T4	PA
metaxalone oral tablet 400 mg	T2	PA
metaxalone oral tablet 800 mg	T3	PA
methadone oral solution 10 mg/5 ml	T2	PA; QL (620 ML per 31 days)
methadone oral solution 5 mg/5 ml	T2	PA; QL (1240 ML per 31 days)
methadone oral tablet 10 mg	T2	PA; QL (124 EA per 31 days)
methadone oral tablet 5 mg	T2	PA; QL (248 EA per 31 days)
methsuximide	T4	
methylphenidate hcl oral capsule, er biphasic 30-70	T4	QL (31 EA per 31 days)
methylphenidate hcl oral capsule,er biphasic 50-50 10 mg	T4	QL (186 EA per 31 days)
methylphenidate hcl oral capsule,er biphasic 50-50 20 mg	T4	QL (93 EA per 31 days)
methylphenidate hcl oral capsule,er biphasic 50-50 30 mg, 40 mg	T4	QL (62 EA per 31 days)
methylphenidate hcl oral capsule,er biphasic 50-50 60 mg	T4	QL (31 EA per 31 days)
methylphenidate hcl oral tablet	T2	QL (93 EA per 31 days)
methylphenidate hcl oral tablet extended release	T2	QL (93 EA per 31 days)
methylphenidate hcl oral tablet extended release 24hr 18 mg, 18 mg (bx rating), 27 mg, 36 mg, 54 mg	T4	QL (31 EA per 31 days)
methylphenidate transdermal	T4	PA; QL (30 EA per 30 days)
mirtazapine oral tablet 15 mg, 30 mg, 45 mg	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
mirtazapine oral tablet 7.5 mg	T3	
mirtazapine oral tablet,disintegrating	T3	
modafinil	T2	PA; QL (31 EA per 31 days)
molindone	T4	
morphine concentrate oral solution	T2	PA; QL (310 ML per 31 days)
morphine oral solution 10 mg/5 ml (2 mg/ml)	T2	PA; QL (2800 ML per 31 days)
morphine oral solution 20 mg/5 ml (4 mg/ml)	T2	PA; QL (1400 ML per 31 days)
morphine oral tablet	T2	PA; QL (186 EA per 31 days)
morphine oral tablet extended release 100 mg	T3	PA; QL (62 EA per 31 days)
morphine oral tablet extended release 15 mg, 30 mg, 60 mg	T3	PA; QL (100 EA per 31 days)
morphine oral tablet extended release 200 mg	T3	PA; QL (31 EA per 31 days)
nabumetone	T1	
naloxone injection solution	T2	
naloxone injection syringe	T2	
naltrexone	T2	
naproxen oral tablet	T1	
naproxen oral tablet,delayed release (dr/ec)	T2	
naproxen sodium oral tablet 275 mg, 550 mg	T1	
naproxen sodium oral tablet, er multiphase 24 hr 750 mg	T4	
naratriptan oral tablet 1 mg	T2	QL (20 EA per 28 days)
naratriptan oral tablet 2.5 mg	T2	QL (9 EA per 28 days)
NAYZILAM	T4	PA-NS; QL (10 EA per 30 days)
nefazodone	T2	
NEUPRO	T4	
nortriptyline	T2	
NOURIANZ	T5	PA; QL (31 EA per 31 days)
NUCYNTA ORAL TABLET 100 MG, 75 MG	T5	PA; QL (186 EA per 31 days)
NUCYNTA ORAL TABLET 50 MG	T4	PA; QL (186 EA per 31 days)
NUEDEXTA	T5	PA; QL (62 EA per 31 days)
NUPLAZID	T5	PA-NS; QL (31 EA per 31 days)
NURTEC ODT	T5	PA; QL (18 EA per 28 days)
olanzapine intramuscular	T2	
olanzapine oral	T2	QL (31 EA per 31 days)
olanzapine-fluoxetine	T2	
ONGENTYS	T4	PA; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ONZETRA XSAIL	T4	QL (16 EA per 28 days)
OPIPZA	T5	PA-NS
oxaprozin oral tablet	T2	
oxazepam	T4	PA
oxcarbazepine oral suspension	T2	
oxcarbazepine oral tablet	T2	
oxcarbazepine oral tablet extended release 24 hr 150 mg, 300 mg	T4	
oxcarbazepine oral tablet extended release 24 hr 600 mg	T5	
oxycodone oral capsule	T2	PA; QL (186 EA per 31 days)
oxycodone oral concentrate	T4	PA; QL (180 ML per 31 days)
oxycodone oral tablet 10 mg, 15 mg, 20 mg, 5 mg	T2	PA; QL (186 EA per 31 days)
oxycodone oral tablet 30 mg	T3	PA; QL (138 EA per 31 days)
oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg	T3	PA; QL (372 EA per 31 days)
oxycodone-acetaminophen oral tablet 5-325 mg, 7.5-325 mg	T2	PA; QL (372 EA per 31 days)
paliperidone oral tablet extended release 24hr 1.5 mg, 3 mg, 9 mg	T4	QL (31 EA per 31 days)
paliperidone oral tablet extended release 24hr 6 mg	T4	QL (62 EA per 31 days)
paroxetine hcl oral suspension	T4	PA-NS
paroxetine hcl oral tablet	T1	PA-NS
paroxetine hcl oral tablet extended release 24 hr	T3	PA-NS
perampanel oral suspension	T5	QL (744 ML per 31 days)
perampanel oral tablet 10 mg, 12 mg, 4 mg, 6 mg, 8 mg	T5	QL (31 EA per 31 days)
perampanel oral tablet 2 mg	T4	QL (31 EA per 31 days)
perphenazine	T2	
perphenazine-amitriptyline	T2	PA-NS
phenelzine	T2	
phenobarbital	T2	PA-NS
PHENYTEK	T3	
phenytoin oral suspension 125 mg/5 ml	T2	
phenytoin oral tablet,chewable	T2	
phenytoin sodium extended oral capsule 100 mg	T2	
pimozide	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
piroxicam	T2	
pramipexole oral tablet	T2	
pregabalin oral capsule 100 mg, 150 mg, 200 mg, 25 mg, 50 mg, 75 mg	T2	PA-NS; QL (93 EA per 31 days)
pregabalin oral capsule 225 mg, 300 mg	T2	PA-NS; QL (62 EA per 31 days)
pregabalin oral solution	T2	PA-NS; QL (930 ML per 31 days)
primidone oral tablet 125 mg	T3	
primidone oral tablet 250 mg, 50 mg	T2	
protriptyline	T4	
pyridostigmine bromide oral syrup	T5	
pyridostigmine bromide oral tablet 30 mg	T3	
pyridostigmine bromide oral tablet 60 mg	T2	
pyridostigmine bromide oral tablet extended release 180 mg	T2	
quetiapine oral tablet 100 mg, 200 mg, 300 mg, 400 mg, 50 mg	T2	QL (62 EA per 31 days)
quetiapine oral tablet 150 mg	T3	QL (62 EA per 31 days)
quetiapine oral tablet 25 mg	T1	QL (62 EA per 31 days)
quetiapine oral tablet extended release 24 hr	T3	QL (62 EA per 31 days)
QULIPTA	T3	PA; QL (31 EA per 31 days)
RADICAVA ORS STARTER KIT SUSP	T5	PA; QL (70 ML per 28 days)
RALDESY	T5	
ramelteon	T4	QL (31 EA per 31 days)
rasagiline	T3	
RELGAABI ORAL CAPSULE 300 MG	T2	QL (360 EA per 30 days)
RELGAABI ORAL CAPSULE 400 MG	T2	QL (270 EA per 30 days)
REXULTI ORAL TABLET	T5	PA-NS; QL (31 EA per 31 days)
REZENOPY	T3	
risperidone microspheres intramuscular suspension,extended rel recon 12.5 mg/2 ml	T4	QL (2 EA per 28 days)
risperidone microspheres intramuscular suspension,extended rel recon 25 mg/2 ml, 37.5 mg/2 ml, 50 mg/2 ml	T5	QL (2 EA per 28 days)
risperidone oral solution	T2	QL (496 ML per 31 days)
risperidone oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg	T1	QL (31 EA per 31 days)
risperidone oral tablet 3 mg	T1	QL (93 EA per 31 days)
risperidone oral tablet 4 mg	T1	QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
risperidone oral tablet,disintegrating 0.25 mg, 0.5 mg, 1 mg, 2 mg	T4	QL (31 EA per 31 days)
risperidone oral tablet,disintegrating 3 mg	T4	QL (93 EA per 31 days)
risperidone oral tablet,disintegrating 4 mg	T4	QL (124 EA per 31 days)
rivastigmine	T2	QL (30 EA per 30 days)
rivastigmine tartrate	T2	
rizatriptan oral tablet 10 mg	T2	QL (12 EA per 28 days)
rizatriptan oral tablet 5 mg	T2	QL (24 EA per 28 days)
rizatriptan oral tablet,disintegrating 10 mg	T2	QL (12 EA per 28 days)
rizatriptan oral tablet,disintegrating 5 mg	T2	QL (24 EA per 28 days)
ropinirole oral tablet	T2	
ropinirole oral tablet extended release 24 hr	T4	
ROWEEPRA	T2	
rufinamide oral suspension	T5	PA-NS
rufinamide oral tablet 200 mg	T4	PA-NS
rufinamide oral tablet 400 mg	T5	PA-NS
RYTARY	T3	ST
SECUADO	T5	PA-NS; QL (31 EA per 31 days)
selegiline hcl	T2	
sertraline oral concentrate	T2	
sertraline oral tablet	T1	
SKYCLARYS	T5	PA; QL (93 EA per 31 days)
sodium oxybate	T5	PA; QL (540 ML per 30 days)
SPRITAM ORAL TABLET FOR SUSPENSION 250 MG, 500 MG	T4	
SUBVENITE ORAL SUSPENSION	T5	
SUBVENITE ORAL TABLET	T2	
sulindac	T2	
sumatriptan nasal spray,non-aerosol 20 mg/actuation	T2	QL (8 EA per 28 days)
sumatriptan nasal spray,non-aerosol 5 mg/actuation	T2	QL (32 EA per 28 days)
sumatriptan succinate oral tablet 100 mg	T2	QL (9 EA per 28 days)
sumatriptan succinate oral tablet 25 mg	T2	QL (36 EA per 28 days)
sumatriptan succinate oral tablet 50 mg	T2	QL (18 EA per 28 days)
sumatriptan succinate subcutaneous pen injector 6 mg/0.5 ml	T4	QL (4 ML per 28 days)
sumatriptan succinate subcutaneous solution	T4	QL (4 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
sumatriptan-naproxen	T4	QL (9 EA per 28 days)
SUNOSI	T4	PA; QL (31 EA per 31 days)
SYMPAZAN ORAL FILM 10 MG, 20 MG	T5	PA-NS; QL (62 EA per 31 days)
SYMPAZAN ORAL FILM 5 MG	T4	PA-NS; QL (62 EA per 31 days)
TASCENSO ODT	T5	PA; QL (31 EA per 31 days)
tasimelteon	T5	PA; QL (31 EA per 31 days)
temazepam	T2	PA; QL (31 EA per 31 days)
TENCON	T2	QL (372 EA per 31 days)
teriflunomide	T5	PA; QL (31 EA per 31 days)
tetrabenazine oral tablet 12.5 mg	T5	PA; QL (93 EA per 31 days)
tetrabenazine oral tablet 25 mg	T5	PA; QL (124 EA per 31 days)
thioridazine	T2	
thiothixene	T1	
tiagabine oral tablet 12 mg, 16 mg	T3	
tiagabine oral tablet 2 mg, 4 mg	T4	
tizanidine oral capsule 2 mg, 4 mg, 6 mg	T2	
tizanidine oral tablet	T2	
tolcapone	T5	
topiramate oral capsule, sprinkle 15 mg, 25 mg	T2	
topiramate oral capsule,extended release 24hr 100 mg, 25 mg, 50 mg	T4	
topiramate oral capsule,extended release 24hr 200 mg	T5	
topiramate oral solution	T4	PA-NS; QL (496 ML per 31 days)
topiramate oral tablet	T2	
tramadol oral tablet 100 mg	T4	PA; QL (124 EA per 31 days)
tramadol oral tablet 25 mg	T4	PA; QL (496 EA per 31 days)
tramadol oral tablet 50 mg	T2	PA; QL (240 EA per 30 days)
tramadol oral tablet 75 mg	T4	PA; QL (155 EA per 31 days)
tramadol-acetaminophen	T2	PA; QL (372 EA per 31 days)
tranylcypromine	T2	
trazodone oral tablet 100 mg, 150 mg, 50 mg	T1	
trazodone oral tablet 300 mg	T2	
triazolam	T4	PA
trifluoperazine	T2	
trihexyphenidyl	T2	
trimipramine	T3	PA-NS

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TRINTELLIX	T3	PA-NS
UBRELVY ORAL TABLET 100 MG	T5	PA; QL (17 EA per 28 days)
UBRELVY ORAL TABLET 50 MG	T5	PA; QL (34 EA per 28 days)
valproic acid	T2	
valproic acid (as sodium salt) oral solution 250 mg/5 ml	T2	
VALTOCO	T5	PA-NS; QL (10 EA per 30 days)
venlafaxine oral capsule,extended release 24hr 150 mg, 37.5 mg	T2	QL (31 EA per 31 days)
venlafaxine oral capsule,extended release 24hr 75 mg	T2	QL (93 EA per 31 days)
venlafaxine oral tablet	T2	
venlafaxine oral tablet extended release 24hr	T4	QL (31 EA per 31 days)
VERSACLOZ	T5	QL (558 ML per 31 days)
vigabatrin	T5	PA-NS
VIGADRONE	T5	PA-NS
vilazodone	T3	PA-NS; QL (31 EA per 31 days)
VIVITROL	T5	
VRAYLAR	T5	PA-NS; QL (31 EA per 31 days)
VUMERITY	T5	PA; QL (124 EA per 31 days)
XCOPRI	T5	PA-NS
XCOPRI MAINTENANCE PACK	T5	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 12.5 MG (14)- 25 MG (14)	T4	PA-NS
XCOPRI TITRATION PACK ORAL TABLETS,DOSE PACK 150 MG (14)- 200 MG (14), 50 MG (14)- 100 MG (14)	T5	PA-NS
zaleplon oral capsule 10 mg	T2	PA; QL (62 EA per 31 days)
zaleplon oral capsule 5 mg	T2	PA; QL (93 EA per 31 days)
ZAVZPRET	T5	PA; QL (8 EA per 30 days)
ZELAPAR	T5	
ZEPOSIA	T5	PA; QL (31 EA per 31 days)
ZEPOSIA STARTER KIT (28-DAY)	T5	PA; QL (56 EA per 365 days)
ZEPOSIA STARTER PACK (7-DAY)	T5	PA; QL (14 EA per 365 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML	T5	PA; QL (11.648 ML per 28 days)
ZILBRYSQ SUBCUTANEOUS SYRINGE 23 MG/0.574 ML	T5	PA; QL (16.072 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ZILBRYSQ SUBCUTANEOUS SYRINGE 32.4 MG/0.81 ML	T5	PA; QL (22.68 ML per 28 days)
ziprasidone hcl	T3	QL (62 EA per 31 days)
ziprasidone mesylate	T3	
zolmitriptan nasal spray,non-aerosol 5 mg	T4	QL (8 EA per 28 days)
zolmitriptan oral tablet 2.5 mg	T2	QL (16 EA per 28 days)
zolmitriptan oral tablet 5 mg	T2	QL (8 EA per 28 days)
zolmitriptan oral tablet,disintegrating 2.5 mg	T2	QL (16 EA per 28 days)
zolmitriptan oral tablet,disintegrating 5 mg	T2	QL (8 EA per 28 days)
zolpidem oral tablet	T2	PA; QL (31 EA per 31 days)
zolpidem oral tablet,ext release multiphase	T2	PA; QL (31 EA per 31 days)
zolpidem sublingual	T3	PA; QL (31 EA per 31 days)
ZONISADE	T5	PA-NS; QL (930 ML per 31 days)
zonisamide	T2	
ZTALMY	T5	PA-NS; QL (1100 ML per 30 days)
ZUBSOLV	T3	
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG	T5	PA-NS; QL (28 EA per 180 days)
ZURZUVAE ORAL CAPSULE 30 MG	T5	PA-NS; QL (14 EA per 180 days)
Cardiovascular, Hypertension / Lipids		
acebutolol	T1	
aliskiren	T4	
amiloride	T1	
amiloride-hydrochlorothiazide	T1	
amiodarone oral	T2	
amlodipine	T1	
amlodipine-atorvastatin	T2	
amlodipine-benazepril	T1	
amlodipine-olmesartan	T2	QL (31 EA per 31 days)
amlodipine-valsartan	T2	
aspirin-dipyridamole	T2	
atenolol	T1	
atenolol-chlorthalidone	T1	
atorvastatin	T1	
ATTRUBY	T5	PA; QL (124 EA per 31 days)
benazepril	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
benazepril-hydrochlorothiazide	T1	
betaxolol oral	T1	
bisoprolol fumarate oral tablet 10 mg, 5 mg	T1	
bisoprolol-hydrochlorothiazide	T1	
bumetanide	T2	
candesartan	T2	
candesartan-hydrochlorothiazid	T2	
captopril	T1	
CARTIA XT	T1	
carvedilol	T1	
carvedilol phosphate	T4	
chlorthalidone oral tablet 25 mg, 50 mg	T1	
cholestyramine (with sugar) oral powder in packet	T2	
CHOLESTYRAMINE LIGHT ORAL POWDER IN PACKET	T2	
cilostazol	T2	
clonidine	T4	
clonidine hcl oral tablet 0.1 mg, 0.2 mg, 0.3 mg	T1	
clopidogrel	T1	
colesevelam	T4	
colestipol oral packet	T2	
colestipol oral tablet	T2	
dabigatran etexilate oral capsule 110 mg	T4	QL (124 EA per 31 days)
dabigatran etexilate oral capsule 150 mg, 75 mg	T4	QL (62 EA per 31 days)
digoxin oral solution	T3	QL (155 ML per 31 days)
digoxin oral tablet 125 mcg (0.125 mg)	T1	QL (62 EA per 31 days)
digoxin oral tablet 250 mcg (0.25 mg)	T2	QL (31 EA per 31 days)
digoxin oral tablet 62.5 mcg (0.0625 mg)	T4	QL (124 EA per 31 days)
diltiazem hcl oral capsule,extended release 12 hr	T1	
diltiazem hcl oral capsule,extended release 24 hr 360 mg, 420 mg	T1	
diltiazem hcl oral capsule,extended release 24hr 120 mg, 180 mg, 240 mg, 300 mg	T1	
diltiazem hcl oral tablet	T1	
diltiazem hcl oral tablet extended release 24 hr	T1	
DILT-XR	T1	
dofetilide	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
doxazosin	T1	
EDARBYCLOR	T4	
ELIQUIS DVT-PE TREAT 30D START	T3	QL (74 EA per 30 days)
ELIQUIS ORAL TABLET 2.5 MG	T3	QL (60 EA per 30 days)
ELIQUIS ORAL TABLET 5 MG	T3	QL (74 EA per 30 days)
ELIQUIS ORAL TABLET FOR SUSPENSION 0.5 MG	T3	QL (112 EA per 28 days)
ELIQUIS ORAL TABLET FOR SUSPENSION 1.5 MG (0.5 MG X 3)	T3	QL (336 EA per 28 days)
ELIQUIS ORAL TABLET FOR SUSPENSION 2 MG (0.5 MG X 4)	T3	QL (560 EA per 28 days)
ELIQUIS SPRINKLE	T3	QL (70 EA per 28 days)
eltrombopag olamine oral powder in packet 12.5 mg	T5	PA; QL (372 EA per 31 days)
eltrombopag olamine oral powder in packet 25 mg	T5	PA; QL (31 EA per 31 days)
eltrombopag olamine oral tablet 12.5 mg, 25 mg	T5	PA; QL (31 EA per 31 days)
eltrombopag olamine oral tablet 50 mg, 75 mg	T5	PA; QL (62 EA per 31 days)
enalapril maleate oral solution	T4	
enalapril maleate oral tablet	T1	
enalapril-hydrochlorothiazide	T1	
enoxaparin subcutaneous syringe 100 mg/ml, 120 mg/0.8 ml, 150 mg/ml	T4	
enoxaparin subcutaneous syringe 30 mg/0.3 ml, 40 mg/0.4 ml, 60 mg/0.6 ml, 80 mg/0.8 ml	T2	
ENTRESTO ORAL TABLET 24-26 MG	T4	ST; QL (186 EA per 31 days)
ENTRESTO ORAL TABLET 49-51 MG	T4	ST; QL (93 EA per 31 days)
ENTRESTO ORAL TABLET 97-103 MG	T4	ST; QL (62 EA per 31 days)
ENTRESTO SPRINKLE	T3	PA; QL (248 EA per 31 days)
eplerenone	T2	
ethacrynic acid	T4	
ezetimibe	T2	
ezetimibe-simvastatin	T3	QL (31 EA per 31 days)
felodipine	T2	
fenofibrate micronized oral capsule 134 mg, 200 mg, 43 mg, 67 mg	T2	
fenofibrate nanocrystallized	T2	
fenofibrate oral tablet 120 mg, 40 mg	T4	
fenofibrate oral tablet 160 mg, 54 mg	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
fenofibric acid (choline)	T3	
flecainide	T2	
fluvastatin oral capsule	T1	
fluvastatin oral tablet extended release 24 hr	T3	
fondaparinux subcutaneous syringe 10 mg/0.8 ml, 5 mg/0.4 ml, 7.5 mg/0.6 ml	T5	
fondaparinux subcutaneous syringe 2.5 mg/0.5 ml	T4	
fosinopril	T1	
fosinopril-hydrochlorothiazide	T1	
furosemide injection	T2	
furosemide oral solution	T2	
furosemide oral tablet	T1	
gemfibrozil	T1	
heparin (porcine) injection solution	T2	PA-BvD
hydralazine oral	T1	
hydrochlorothiazide	T1	
icosapent ethyl oral capsule 0.5 gram	T4	QL (248 EA per 31 days)
icosapent ethyl oral capsule 1 gram	T4	QL (124 EA per 31 days)
indapamide	T1	
irbesartan	T1	QL (31 EA per 31 days)
irbesartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
isosorbide dinitrate oral tablet 10 mg, 20 mg, 30 mg, 5 mg	T2	
isosorbide dinitrate oral tablet 40 mg	T4	
isosorbide mononitrate	T1	
isosorbide-hydralazine	T4	
isradipine	T2	
ivabradine oral tablet 5 mg	T4	PA; QL (93 EA per 31 days)
ivabradine oral tablet 7.5 mg	T4	PA; QL (62 EA per 31 days)
JANTOVEN	T1	
KERENDIA	T4	PA; QL (31 EA per 31 days)
labetalol oral tablet 100 mg, 200 mg, 300 mg	T1	
LIPOFEN	T4	
lisinopril	T1	
lisinopril-hydrochlorothiazide	T1	
losartan oral tablet 100 mg	T1	QL (31 EA per 31 days)
losartan oral tablet 25 mg	T1	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
losartan oral tablet 50 mg	T1	QL (62 EA per 31 days)
losartan-hydrochlorothiazide	T1	
lovastatin	T1	
MATZIM LA	T2	
metolazone	T2	
metoprolol succinate	T1	
metoprolol ta-hydrochlorothiaz	T1	
metoprolol tartrate oral tablet 100 mg, 25 mg, 37.5 mg, 50 mg, 75 mg	T1	
metyrosine	T5	PA
mexiletine	T3	
minoxidil oral	T2	
moexipril	T1	
MULPLETA	T5	PA
MULTAQ	T4	
nadolol	T1	
nebivolol oral tablet 10 mg, 2.5 mg	T2	QL (93 EA per 31 days)
nebivolol oral tablet 20 mg	T2	QL (62 EA per 31 days)
nebivolol oral tablet 5 mg	T2	QL (217 EA per 31 days)
NEXLETOL	T4	PA; QL (31 EA per 31 days)
NEXLIZET	T4	PA; QL (31 EA per 31 days)
niacin oral tablet extended release 24 hr 1,000 mg, 750 mg	T3	
niacin oral tablet extended release 24 hr 500 mg	T3	QL (31 EA per 31 days)
nicardipine oral capsule 20 mg	T2	
nicardipine oral capsule 30 mg	T4	
nifedipine oral tablet extended release	T2	
nifedipine oral tablet extended release 24hr	T2	
nimodipine oral capsule	T2	
nisoldipine oral tablet extended release 24 hr 17 mg, 34 mg, 8.5 mg	T2	
NITRO-BID	T2	
nitroglycerin sublingual	T2	
nitroglycerin transdermal patch 24 hour	T2	
nitroglycerin translingual	T3	
olmesartan oral tablet 20 mg, 40 mg	T1	QL (31 EA per 31 days)
olmesartan oral tablet 5 mg	T1	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
olmesartan-amlodipin-hcthiazid	T3	
olmesartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
omega-3 acid ethyl esters	T3	QL (124 EA per 31 days)
PACERONE ORAL TABLET 100 MG, 200 MG	T2	
pentoxifylline	T2	
perindopril erbumine	T1	
pindolol	T1	
pitavastatin calcium	T3	
prasugrel hcl	T3	
pravastatin	T1	
prazosin	T1	
PREVALITE ORAL POWDER IN PACKET	T2	
propafenone	T2	
propranolol oral capsule,extended release 24 hr	T2	
propranolol oral solution	T1	
propranolol oral tablet	T1	
quinapril	T1	
quinapril-hydrochlorothiazide	T1	
quinidine gluconate oral	T4	
quinidine sulfate oral tablet	T2	
ramipril	T1	
ranolazine	T3	QL (62 EA per 31 days)
REPATHA SURECLICK	T3	PA; QL (3 ML per 28 days)
REPATHA SYRINGE	T3	PA; QL (3 ML per 28 days)
rosuvastatin	T1	
sacubitril-valsartan oral tablet 24-26 mg	T2	QL (186 EA per 31 days)
sacubitril-valsartan oral tablet 49-51 mg	T2	QL (93 EA per 31 days)
sacubitril-valsartan oral tablet 97-103 mg	T2	QL (62 EA per 31 days)
simvastatin	T1	
SOTALOL AF	T1	
sotalol oral	T1	
spironolactone oral tablet	T1	
spironolacton-hydrochlorothiaz	T1	
TAVALISSE	T5	PA; QL (62 EA per 31 days)
telmisartan	T1	
telmisartan-amlodipine	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
telmisartan-hydrochlorothiazid	T1	
terazosin	T1	
TIADYLT ER	T2	
ticagrelor	T2	
timolol maleate oral	T1	
torsemide oral	T1	
trandolapril	T1	
triamterene	T4	
triamterene-hydrochlorothiazid	T1	
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 400 MCG, 600 MCG, 800 MCG	T5	PA; QL (62 EA per 31 days)
UPTRAVI ORAL TABLET 200 MCG	T5	PA; QL (224 EA per 28 days)
UPTRAVI ORAL TABLETS,DOSE PACK	T5	PA; QL (400 EA per 365 days)
valsartan oral tablet 160 mg, 40 mg, 80 mg	T1	QL (62 EA per 31 days)
valsartan oral tablet 320 mg	T1	QL (31 EA per 31 days)
valsartan-hydrochlorothiazide	T1	QL (31 EA per 31 days)
verapamil oral	T2	
VERQUVO	T4	PA; QL (31 EA per 31 days)
warfarin	T1	
XARELTO DVT-PE TREAT 30D START	T3	QL (51 EA per 30 days)
XARELTO ORAL SUSPENSION FOR RECONSTITUTION	T3	QL (930 ML per 31 days)
XARELTO ORAL TABLET 10 MG, 20 MG	T3	QL (31 EA per 31 days)
XARELTO ORAL TABLET 15 MG	T3	QL (52 EA per 31 days)
XARELTO ORAL TABLET 2.5 MG	T3	QL (62 EA per 31 days)
Dermatologicals/Topical Therapy		
ACCUTANE ORAL CAPSULE 10 MG, 20 MG, 40 MG	T2	
acitretin	T4	PA
acyclovir topical cream	T3	QL (5 GM per 28 days)
acyclovir topical ointment	T1	QL (30 GM per 30 days)
adapalene topical gel 0.3 %	T2	PA
adapalene topical swab	T2	PA
adapalene-benzoyl peroxide	T4	PA
ALA-CORT TOPICAL CREAM	T1	
alclometasone	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
amcinonide topical cream	T2	
amcinonide topical ointment	T2	
ammonium lactate	T2	
AMNESTEEM	T2	
ANZUPGO	T5	PA; QL (60 GM per 30 days)
azelaic acid	T4	ST; QL (50 GM per 28 days)
AZELEX	T4	QL (50 GM per 28 days)
betamethasone dipropionate	T1	
betamethasone valerate topical cream	T2	
betamethasone valerate topical foam	T4	
betamethasone valerate topical lotion	T2	
betamethasone valerate topical ointment	T2	
betamethasone, augmented	T2	
calcipotriene scalp	T2	QL (60 ML per 28 days)
calcipotriene topical cream	T2	QL (60 GM per 28 days)
calcipotriene topical ointment	T2	QL (60 GM per 28 days)
calcipotriene-betamethasone topical ointment	T4	ST; QL (400 GM per 28 days)
calcipotriene-betamethasone topical suspension	T5	ST; QL (400 GM per 28 days)
calcitriol topical	T4	ST
ciclopirox topical cream	T2	QL (90 GM per 28 days)
ciclopirox topical gel	T2	QL (45 GM per 28 days)
ciclopirox topical shampoo	T2	QL (120 ML per 28 days)
ciclopirox topical solution	T2	
ciclopirox topical suspension	T3	QL (60 ML per 28 days)
CLARAVIS	T4	
CLINDACIN	T2	QL (100 GM per 28 days)
clindamycin phosphate topical foam	T4	QL (100 GM per 28 days)
clindamycin phosphate topical gel	T2	QL (60 GM per 28 days)
clindamycin phosphate topical lotion	T2	QL (60 ML per 28 days)
clindamycin phosphate topical solution	T2	QL (60 ML per 28 days)
clindamycin phosphate topical swab	T2	
clindamycin-benzoyl peroxide topical gel	T2	
clindamycin-benzoyl peroxide topical gel with pump 1.2-2.5 %	T2	
clindamycin-tretinoin	T2	PA; QL (60 GM per 28 days)
clobetasol scalp	T2	QL (50 ML per 28 days)
clobetasol topical cream 0.05 %	T3	QL (60 GM per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
clobetasol topical foam	T2	QL (100 GM per 28 days)
clobetasol topical gel	T2	QL (60 GM per 28 days)
clobetasol topical lotion	T2	QL (118 ML per 28 days)
clobetasol topical ointment	T3	QL (60 GM per 28 days)
clobetasol topical shampoo	T4	QL (118 ML per 28 days)
clobetasol topical spray,non-aerosol	T2	QL (125 ML per 28 days)
clobetasol-emollient topical cream	T3	QL (60 GM per 28 days)
clobetasol-emollient topical foam	T3	QL (100 GM per 28 days)
clocortolone pivalate	T4	
CLODAN	T2	QL (118 ML per 28 days)
clotrimazole topical cream	T2	QL (45 GM per 28 days)
clotrimazole topical solution	T2	QL (30 ML per 28 days)
clotrimazole-betamethasone topical cream	T2	QL (45 GM per 28 days)
clotrimazole-betamethasone topical lotion	T2	QL (60 ML per 28 days)
COSENTYX (2 SYRINGES)	T5	PA; QL (2 ML per 28 days)
COSENTYX PEN (2 PENS)	T5	PA; QL (2 ML per 28 days)
COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML	T5	PA; QL (0.5 ML per 28 days)
COSENTYX UNOREADY PEN	T5	PA; QL (2 ML per 28 days)
dapsone topical gel 5 %	T4	QL (90 GM per 28 days)
dapsone topical gel with pump	T4	QL (90 GM per 28 days)
desonide topical cream	T3	QL (60 GM per 28 days)
desonide topical gel	T3	QL (60 GM per 28 days)
desonide topical lotion	T3	QL (118 ML per 28 days)
desonide topical ointment	T3	QL (60 GM per 28 days)
desoximetasone topical cream	T2	QL (100 GM per 28 days)
desoximetasone topical gel	T2	QL (60 GM per 28 days)
desoximetasone topical ointment 0.25 %	T2	QL (100 GM per 28 days)
desoximetasone topical spray,non-aerosol	T2	QL (100 ML per 28 days)
diclofenac sodium topical gel 3 %	T4	PA; QL (100 GM per 28 days)
doxepin topical	T4	PA; QL (45 GM per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML	T5	PA; QL (2.28 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 300 MG/2 ML	T5	PA; QL (8 ML per 28 days)
econazole nitrate topical cream	T2	QL (85 GM per 28 days)
ERY PADS	T2	
erythromycin with ethanol topical gel	T2	QL (60 GM per 28 days)
erythromycin with ethanol topical solution	T2	QL (60 ML per 28 days)
erythromycin-benzoyl peroxide	T2	
FILSUVEZ	T5	PA
FINACEA TOPICAL FOAM	T4	ST; QL (50 GM per 28 days)
fluocinolone and shower cap	T2	QL (118.28 ML per 28 days)
fluocinolone topical cream 0.01 %	T2	QL (60 GM per 28 days)
fluocinolone topical cream 0.025 %	T2	QL (120 GM per 28 days)
fluocinolone topical ointment	T2	QL (120 GM per 28 days)
fluocinolone topical solution	T2	QL (90 ML per 28 days)
fluocinonide topical cream 0.05 %	T2	QL (60 GM per 28 days)
fluocinonide topical gel	T2	QL (60 GM per 28 days)
fluocinonide topical ointment	T2	QL (60 GM per 28 days)
fluocinonide topical solution	T2	QL (60 ML per 28 days)
fluocinonide-emollient	T2	QL (60 GM per 28 days)
fluorouracil topical cream 5 %	T2	
fluorouracil topical solution	T2	
flurandrenolide topical lotion	T4	QL (120 ML per 28 days)
fluticasone propionate topical cream	T2	
gentamicin topical	T2	QL (60 GM per 28 days)
halcinonide topical cream	T4	QL (60 GM per 28 days)
halobetasol propionate topical cream	T2	QL (50 GM per 28 days)
halobetasol propionate topical ointment	T2	QL (50 GM per 28 days)
hydrocortisone butyrate topical cream	T2	QL (45 GM per 28 days)
hydrocortisone butyrate topical lotion	T4	QL (118 ML per 28 days)
hydrocortisone butyrate topical ointment	T2	QL (45 GM per 28 days)
hydrocortisone butyrate topical solution	T2	QL (60 ML per 28 days)
hydrocortisone topical cream 1 %	T1	
hydrocortisone topical lotion 2.5 %	T1	QL (118 ML per 28 days)
hydrocortisone topical ointment 1 %, 2.5 %	T1	
hydrocortisone valerate	T2	QL (60 GM per 28 days)
imiquimod topical cream in metered-dose pump	T4	
imiquimod topical cream in packet 5 %	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
isotretinoin	T2	
ketoconazole topical cream	T2	QL (60 GM per 28 days)
ketoconazole topical shampoo	T2	QL (120 ML per 28 days)
lidocaine hcl mucous membrane solution 2 %	T2	
lidocaine hcl mucous membrane solution 4 % (40 mg/ml)	T2	PA; QL (50 ML per 28 days)
lidocaine topical adhesive patch,medicated 5 %	T3	PA; QL (93 EA per 31 days)
lidocaine topical ointment	T2	PA; QL (50 GM per 28 days)
lidocaine-prilocaine topical cream	T2	PA; QL (30 GM per 28 days)
malathion	T4	
metronidazole topical cream	T2	
metronidazole topical gel 0.75 %	T2	
metronidazole topical gel 1 %	T1	
metronidazole topical lotion	T2	
mometasone topical	T2	
mupirocin	T2	
naftifine topical cream 1 %	T4	ST; QL (90 GM per 28 days)
naftifine topical cream 2 %	T4	ST; QL (60 GM per 28 days)
naftifine topical gel	T4	ST; QL (60 GM per 28 days)
NEO-SYNALAR	T4	
NEUAC	T2	
NYAMYC	T2	QL (60 GM per 28 days)
nystatin topical cream	T2	QL (30 GM per 28 days)
nystatin topical ointment	T2	QL (30 GM per 28 days)
nystatin topical powder	T2	QL (60 GM per 28 days)
nystatin-triamcinolone	T3	QL (60 GM per 28 days)
NYSTOP	T2	QL (60 GM per 28 days)
PANRETIN	T5	PA-NS
penciclovir	T3	QL (5 GM per 28 days)
permethrin	T2	
pimecrolimus	T3	QL (100 GM per 28 days)
podofilox topical solution	T2	
selenium sulfide topical lotion	T1	
SILIQ	T5	PA; QL (6 ML per 28 days)
silver sulfadiazine	T1	
SKYRIZI SUBCUTANEOUS PEN INJECTOR	T5	PA; QL (1 ML per 84 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML	T5	PA; QL (1 ML per 84 days)
SKYRIZI SUBCUTANEOUS SYRINGE 55 MG/0.37 ML	T5	PA; QL (0.37 ML per 84 days)
SSD	T4	
STELARA SUBCUTANEOUS SOLUTION	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T5	PA; QL (0.5 ML per 84 days)
STELARA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
STEQEYMA SUBCUTANEOUS SOLUTION	T3	PA; QL (0.5 ML per 84 days)
STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T3	PA; QL (0.5 ML per 84 days)
STEQEYMA SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
sulfacetamide sodium (acne)	T1	
SULFAMYLON TOPICAL CREAM	T3	
tacrolimus topical	T2	QL (100 GM per 28 days)
tazarotene topical cream	T4	PA; QL (60 GM per 28 days)
tazarotene topical gel 0.1 %	T4	PA; QL (100 GM per 28 days)
TOVET EMOLLIENT	T3	QL (100 GM per 28 days)
tretinoin topical cream	T2	PA; QL (45 GM per 28 days)
tretinoin topical gel 0.01 %, 0.025 %	T2	PA; QL (45 GM per 28 days)
tretinoin topical gel 0.05 %	T3	PA; QL (45 GM per 28 days)
triamcinolone acetonide topical cream	T1	
triamcinolone acetonide topical lotion	T1	
triamcinolone acetonide topical ointment 0.025 %, 0.1 %, 0.5 %	T1	
triamcinolone acetonide topical ointment 0.05 %	T4	
TRIDERM TOPICAL CREAM 0.5 %	T1	
ustekinumab subcutaneous solution	T5	PA; QL (0.5 ML per 84 days)
ustekinumab subcutaneous syringe 45 mg/0.5 ml	T5	PA; QL (0.5 ML per 84 days)
ustekinumab subcutaneous syringe 90 mg/ml	T5	PA; QL (1 ML per 56 days)
VALCHLOR	T5	PA-NS
VEREGEN	T5	QL (30 GM per 28 days)
YESINTEK SUBCUTANEOUS SOLUTION	T3	PA; QL (0.5 ML per 84 days)
YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML	T3	PA; QL (0.5 ML per 84 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
YESINTEK SUBCUTANEOUS SYRINGE 90 MG/ML	T5	PA; QL (1 ML per 56 days)
ZELSUVMI	T5	PA; QL (31 GM per 28 days)
ZORYVE TOPICAL CREAM 0.15 %	T4	PA; QL (60 GM per 28 days)
Diagnostics / Miscellaneous Agents		
acamprosate	T2	
anagrelide	T3	
ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG	T5	PA
bupropion hcl (smoking deter)	T2	QL (62 EA per 31 days)
carglumic acid	T5	PA
cevimeline	T4	
CHEMET	T4	
CLINIMIX 4.25%/D5W SULFIT FREE	T4	PA-BvD
CLINIMIX E 2.75%/D5W SULF FREE	T4	PA-BvD
d10 %-0.45 % sodium chloride	T2	
d2.5 %-0.45 % sodium chloride	T2	
d5 % and 0.9 % sodium chloride	T2	
d5 %-0.45 % sodium chloride	T2	
deferasirox oral tablet, dispersible 125 mg	T4	PA
deferasirox oral tablet, dispersible 250 mg, 500 mg	T5	PA
dextrose 10 % in water (d10w)	T2	
dextrose 5 % in water (d5w) intravenous parenteral solution	T2	
dextrose 5%-0.2 % sod chloride	T2	
disulfiram	T2	
droxidopa oral capsule 100 mg	T5	PA; QL (465 EA per 31 days)
droxidopa oral capsule 200 mg, 300 mg	T5	PA; QL (186 EA per 31 days)
DUVYZAT	T5	PA; QL (420 ML per 35 days)
FABHALTA	T5	PA; QL (62 EA per 31 days)
GLASSIA	T5	PA
glycerol phenylbutyrate	T5	PA
INCRELEX	T5	PA
JOENJA ORAL TABLET 70 MG	T5	PA; QL (60 EA per 30 days)
KIONEX ORAL SUSPENSION	T2	
KYGEVVI	T5	PA; QL (496 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
lanthanum	T4	PA-BvD
levocarnitine (with sugar)	T2	PA-BvD
levocarnitine oral tablet	T2	PA-BvD
LITFULO	T5	PA; QL (28 EA per 28 days)
LITHOSTAT	T4	
LOARGYS	T5	PA
LOKELMA	T3	PA; QL (93 EA per 31 days)
midodrine	T2	
NICOTROL NS	T4	
nitisinone	T5	PA
PHEBURANE	T5	PA; QL (620 GM per 31 days)
pilocarpine hcl oral	T2	
PROLASTIN-C INTRAVENOUS SOLUTION	T5	PA
REVCOVI	T5	
REZDIFFRA	T5	PA; QL (31 EA per 31 days)
riluzole	T4	
risedronate oral tablet 30 mg	T5	
sevelamer carbonate	T3	PA-BvD
sevelamer hcl	T4	PA-BvD
sodium chloride 0.9 % intravenous parenteral solution	T2	
sodium chloride irrigation	T2	
sodium phenylbutyrate	T5	PA
sodium polystyrene sulfonate	T2	
SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 5 MG	T5	PA; QL (31 EA per 31 days)
SOHONOS ORAL CAPSULE 10 MG	T5	PA; QL (62 EA per 31 days)
SPS (WITH SORBITOL) ORAL	T2	
trientine oral capsule 250 mg	T5	QL (248 EA per 31 days)
varenicline tartrate oral tablet	T4	QL (60 EA per 30 days)
varenicline tartrate oral tablets,dose pack	T4	QL (106 EA per 365 days)
ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG	T5	PA
ZOKINVY	T5	PA
Ear, Nose / Throat Medications		
acetic acid otic (ear)	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
azelastine nasal spray,non-aerosol 137 mcg (0.1 %)	T2	QL (30 ML per 25 days)
chlorhexidine gluconate mucous membrane	T1	
ciprofloxacin-dexamethasone	T3	
fluocinolone acetonide oil	T2	
hydrocortisone-acetic acid	T2	
ipratropium bromide nasal spray,non-aerosol 21 mcg (0.03 %)	T1	QL (30 ML per 28 days)
ipratropium bromide nasal spray,non-aerosol 42 mcg (0.06 %)	T1	QL (15 ML per 28 days)
KOURZEQ	T2	
neomycin-polymyxin-hc otic (ear)	T2	
ofloxacin otic (ear)	T2	
olopatadine nasal	T2	QL (30.5 GM per 30 days)
PERIOGARD	T1	
triamcinolone acetonide dental	T2	
Endocrine/Diabetes		
acarbose	T1	QL (93 EA per 31 days)
ALCOHOL PADS	T2	PA
AVEED	T4	PA
cabergoline	T2	
calcitonin (salmon) nasal	T2	PA-BvD
calcitriol oral capsule	T2	PA-BvD
calcitriol oral solution	T3	PA-BvD
CERDELGA	T5	PA; QL (62 EA per 31 days)
cinacalcet oral tablet 30 mg, 60 mg	T3	PA-BvD; QL (62 EA per 31 days)
cinacalcet oral tablet 90 mg	T3	PA-BvD; QL (124 EA per 31 days)
danazol	T2	
dapagliflozin	T3	QL (31 EA per 31 days)
DEPO-TESTOSTERONE INTRAMUSCULAR OIL 100 MG/ML	T4	PA
desmopressin nasal spray,non-aerosol 10 mcg/spray (0.1 ml)	T4	
desmopressin oral	T2	
dexamethasone oral solution	T1	
dexamethasone oral tablet	T1	
diazoxide	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
doxercalciferol oral capsule 0.5 mcg, 2.5 mcg	T2	PA-BvD
doxercalciferol oral capsule 1 mcg	T4	PA-BvD
FIASP FLEXTOUCH U-100 INSULIN	T3	
FIASP PENFILL U-100 INSULIN	T3	
FIASP U-100 INSULIN	T3	
fludrocortisone	T2	
glimepiride oral tablet 1 mg, 2 mg, 4 mg	T1	
glipizide oral tablet 10 mg, 5 mg	T1	
glipizide oral tablet extended release 24hr	T1	
glipizide-metformin	T1	
GLUCAGON EMERGENCY KIT (HUMAN)	T3	
glyburide	T2	
glyburide-metformin	T2	
GLYXAMBI	T3	QL (31 EA per 31 days)
GVOKE	T3	
GVOKE HYPOPEN 2-PACK	T3	
GVOKE PFS 1-PACK SYRINGE SUBCUTANEOUS SYRINGE 1 MG/0.2 ML	T3	
HUMALOG JUNIOR KWIKPEN U-100	T3	
HUMALOG KWIKPEN INSULIN	T3	
HUMALOG MIX 50-50 KWIKPEN	T3	
HUMALOG MIX 75-25 KWIKPEN	T3	
HUMALOG MIX 75-25(U-100)INSULN	T3	
HUMALOG U-100 INSULIN	T3	
HUMULIN 70/30 U-100 INSULIN	T3	
HUMULIN 70/30 U-100 KWIKPEN	T3	
HUMULIN N NPH INSULIN KWIKPEN	T3	
HUMULIN N NPH U-100 INSULIN	T3	
HUMULIN R REGULAR U-100 INSULN	T3	
HUMULIN R U-500 (CONC) KWIKPEN	T3	
hydrocortisone oral	T1	
insulin lispro	T3	
insulin lispro protamin-lispro	T3	
ISTURISA ORAL TABLET 1 MG	T5	PA; QL (558 EA per 31 days)
ISTURISA ORAL TABLET 5 MG	T5	PA; QL (372 EA per 31 days)
JANUMET	T3	QL (62 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 100-1,000 MG, 50-500 MG	T3	QL (31 EA per 31 days)
JANUMET XR ORAL TABLET, ER MULTIPHASE 24 HR 50-1,000 MG	T3	QL (62 EA per 31 days)
JANUVIA ORAL TABLET 100 MG, 50 MG	T3	QL (31 EA per 31 days)
JANUVIA ORAL TABLET 25 MG	T3	QL (93 EA per 31 days)
JARDIANCE ORAL TABLET 10 MG	T3	QL (62 EA per 31 days)
JARDIANCE ORAL TABLET 25 MG	T3	QL (31 EA per 31 days)
JAVYGTOR	T5	PA
JENTADUETO ORAL TABLET 2.5-1,000 MG, 2.5-500 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG	T3	QL (62 EA per 31 days)
JENTADUETO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 5-1,000 MG	T3	QL (31 EA per 31 days)
LANTUS SOLOSTAR U-100 INSULIN	T3	
LANTUS U-100 INSULIN	T3	
levothyroxine oral capsule	T4	
levothyroxine oral tablet	T1	
LEVOXYL ORAL TABLET 100 MCG, 112 MCG, 125 MCG, 137 MCG, 150 MCG, 175 MCG, 200 MCG, 25 MCG, 50 MCG, 75 MCG, 88 MCG	T3	
LIOMNY	T2	
liothyronine oral	T2	
liraglutide	T4	PA; QL (9 ML per 30 days)
MEDROL ORAL TABLET 2 MG	T4	
metformin oral solution	T4	ST; QL (791 ML per 31 days)
metformin oral tablet 1,000 mg, 500 mg, 850 mg	T1	
metformin oral tablet extended release 24 hr	T1	
methimazole oral tablet 10 mg, 5 mg	T2	
methylprednisolone	T2	
mifepristone oral tablet 300 mg	T5	PA; QL (124 EA per 31 days)
miglitol	T2	
miglustat	T5	PA; QL (93 EA per 31 days)
MOUNJARO	T3	PA; QL (2 ML per 28 days)
nateglinide	T1	QL (93 EA per 31 days)
NOVOLIN 70/30 U-100 INSULIN	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
NOVOLIN 70-30 FLEXPEN U-100	T3	
NOVOLIN N FLEXPEN	T3	
NOVOLIN N NPH U-100 INSULIN	T3	
NOVOLIN R FLEXPEN	T3	
NOVOLIN R REGULAR U100 INSULIN	T3	
NOVOLOG FLEXPEN U-100 INSULIN	T3	
NOVOLOG MIX 70-30 U-100 INSULN	T3	
NOVOLOG MIX 70-30FLEXPEN U-100	T3	
NOVOLOG PENFILL U-100 INSULIN	T3	
NOVOLOG U-100 INSULIN ASPART	T3	
OZEMPIC ORAL	T3	PA; QL (31 EA per 31 days)
OZEMPIC SUBCUTANEOUS	T3	PA; QL (3 ML per 28 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (15 ML per 30 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 2.5 MG/0.5 ML	T5	PA; QL (4 ML per 30 days)
PALYNZIQ SUBCUTANEOUS SYRINGE 20 MG/ML	T5	PA; QL (90 ML per 30 days)
paricalcitol oral	T2	PA-BvD
pioglitazone	T1	QL (31 EA per 31 days)
pioglitazone-metformin	T2	QL (93 EA per 31 days)
prednisolone oral solution	T2	
prednisolone sodium phosphate oral solution 10 mg/5 ml, 20 mg/5 ml (4 mg/ml), 25 mg/5 ml (5 mg/ml), 5 mg base/5 ml (6.7 mg/5 ml)	T2	
prednisone oral solution	T2	
prednisone oral tablet	T1	
prednisone oral tablets,dose pack	T2	
propylthiouracil	T2	
repaglinide oral tablet 0.5 mg	T1	QL (124 EA per 31 days)
repaglinide oral tablet 1 mg	T2	QL (124 EA per 31 days)
repaglinide oral tablet 2 mg	T2	QL (248 EA per 31 days)
sapropterin	T5	PA
sitagliptin oral tablet 100 mg, 50 mg	T3	QL (31 EA per 31 days)
sitagliptin oral tablet 25 mg	T3	QL (93 EA per 31 days)
sitagliptin phos-metformin oral tablet	T3	QL (62 EA per 31 days)
sitagliptin phos-metformin oral tablet, er multiphase 24 hr 100-1,000 mg, 50-500 mg	T3	QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
sitagliptin phos-metformin oral tablet, er multiphase 24 hr 50-1,000 mg	T3	QL (62 EA per 31 days)
sitagliptin phosphate oral tablet 100 mg, 50 mg	T3	QL (31 EA per 31 days)
sitagliptin phosphate oral tablet 25 mg	T3	QL (93 EA per 31 days)
sitagliptin-metformin oral tablet	T3	QL (62 EA per 31 days)
sitagliptin-metformin oral tablet, er multiphase 24 hr 100-1,000 mg, 50-500 mg	T3	QL (31 EA per 31 days)
sitagliptin-metformin oral tablet, er multiphase 24 hr 50-1,000 mg	T3	QL (62 EA per 31 days)
SOLIQUA 100/33	T3	QL (18 ML per 30 days)
SOMAVERT	T5	PA
SYNAREL	T5	PA
SYNJARDY	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 12.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
SYNJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 25-1,000 MG	T3	QL (31 EA per 31 days)
SYNTHROID	T3	
testosterone cypionate	T2	PA
testosterone enanthate	T2	PA
testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)	T4	PA
testosterone transdermal solution in metered pump w/app	T4	PA
TIROSINT	T4	
TIROSINT-SOL	T4	
tolvaptan	T5	PA
tolvaptan (polycys kidney dis) oral tablet	T5	PA; QL (112 EA per 28 days)
tolvaptan (polycys kidney dis) oral tablets, sequential	T5	PA; QL (56 EA per 28 days)
TOUJEO MAX U-300 SOLOSTAR	T3	
TOUJEO SOLOSTAR U-300 INSULIN	T3	
TRADJENTA	T3	QL (31 EA per 31 days)
TRESIBA FLEXTOUCH U-100	T3	
TRESIBA FLEXTOUCH U-200	T3	
TRESIBA U-100 INSULIN	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-5-1,000 MG, 25-5-1,000 MG	T3	QL (31 EA per 31 days)
TRIJARDY XR ORAL TABLET, IR - ER, BIPHASIC 24HR 12.5-2.5-1,000 MG, 5-2.5-1,000 MG	T3	QL (62 EA per 31 days)
TRULICITY	T3	PA; QL (2 ML per 28 days)
UNITHROID	T3	
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 10-1,000 MG, 10-500 MG, 5-500 MG	T3	QL (31 EA per 31 days)
XIGDUO XR ORAL TABLET, IR - ER, BIPHASIC 24HR 2.5-1,000 MG, 5-1,000 MG	T3	QL (62 EA per 31 days)
XULTOPHY 100/3.6	T3	QL (15 ML per 30 days)
YARGESA	T5	PA; QL (93 EA per 31 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML	T5	PA; QL (1.12 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 294 MCG/0.98 ML	T5	PA; QL (1.96 ML per 28 days)
YORVIPATH SUBCUTANEOUS PEN INJECTOR 420 MCG/1.4 ML	T5	PA; QL (2.8 ML per 28 days)
ZELVYSIA	T5	PA
Gastroenterology		
alosetron oral tablet 0.5 mg	T4	PA; QL (93 EA per 31 days)
alosetron oral tablet 1 mg	T5	PA; QL (62 EA per 31 days)
amoxicil-clarithromy-lansopraz	T4	
aprepitant oral capsule 125 mg	T5	PA-BvD
aprepitant oral capsule 40 mg, 80 mg	T4	PA-BvD
aprepitant oral capsule,dose pack	T4	PA-BvD
balsalazide	T2	
betaine	T5	
budesonide oral capsule,delayed,extend.release	T4	
budesonide oral tablet,delayed and ext.release	T5	
cimetidine	T2	
cimetidine hcl oral	T2	
COMPRO	T3	
CONSTULOSE	T2	
CREON	T3	
cromolyn oral	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
dicyclomine oral capsule	T2	
dicyclomine oral solution	T2	
dicyclomine oral tablet 20 mg	T2	
diphenoxylate-atropine oral liquid	T3	
diphenoxylate-atropine oral tablet	T2	
dronabinol oral capsule	T4	PA-BvD
EMEND ORAL SUSPENSION FOR RECONSTITUTION	T4	PA-BvD
ENULOSE	T2	
EOHILIA	T5	PA; QL (1800 ML per 365 days)
esomeprazole magnesium oral capsule,delayed release(dr/ec)	T2	QL (31 EA per 31 days)
famotidine oral suspension for reconstitution	T1	
famotidine oral tablet 20 mg, 40 mg	T1	
GATTEX 30-VIAL	T5	PA
GAVILYTE-C	T1	
GAVILYTE-G	T1	
GAVILYTE-N	T1	
GENERLAC	T2	
glycopyrrolate oral tablet 1 mg, 2 mg	T2	
granisetron hcl oral	T4	PA-BvD
hydrocortisone rectal	T3	
hydrocortisone topical cream with perineal applicator 2.5 %	T1	
hydrocortisone-pramoxine rectal cream 1-1 %	T4	
lactulose oral solution	T1	
lansoprazole oral capsule,delayed release(dr/ec) 15 mg	T3	QL (31 EA per 31 days)
lansoprazole oral capsule,delayed release(dr/ec) 30 mg	T3	QL (62 EA per 31 days)
LINZESS	T3	QL (31 EA per 31 days)
loperamide oral capsule	T2	
lubiprostone	T4	QL (62 EA per 31 days)
meclizine oral tablet 12.5 mg, 25 mg	T2	
mesalamine oral capsule (with del rel tablets)	T3	QL (186 EA per 31 days)
mesalamine oral capsule, extended release 500 mg	T4	PA; QL (248 EA per 31 days)
mesalamine oral capsule,extended release 24hr	T4	QL (124 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
mesalamine oral tablet,delayed release (dr/ec) 1.2 gram	T3	QL (124 EA per 31 days)
mesalamine rectal enema	T4	QL (1860 ML per 31 days)
methscopolamine	T2	
metoclopramide hcl oral solution	T2	
metoclopramide hcl oral tablet	T2	
misoprostol	T2	
MOVANTIK	T3	QL (31 EA per 31 days)
MYTESI	T5	QL (62 EA per 31 days)
nitroglycerin rectal	T4	
nizatidine oral capsule	T2	
omeprazole oral capsule,delayed release(dr/ec)	T1	
ondansetron hcl oral solution	T2	PA-BvD
ondansetron hcl oral tablet 4 mg, 8 mg	T2	PA-BvD
ondansetron oral tablet,disintegrating 4 mg, 8 mg	T2	PA-BvD
pantoprazole oral tablet,delayed release (dr/ec)	T1	
peg 3350-electrolytes	T1	
peg3350-sod sul-nacl-kcl-asb-c	T4	
peg-electrolyte soln	T1	
prochlorperazine	T3	
prochlorperazine maleate	T2	
PROCTO-MED HC	T2	
PROCTOSOL HC TOPICAL	T2	
PROCTOZONE-HC	T2	
rabeprazole oral tablet,delayed release (dr/ec)	T2	QL (62 EA per 31 days)
RELISTOR ORAL	T5	PA; QL (93 EA per 31 days)
RELISTOR SUBCUTANEOUS SOLUTION	T5	PA; QL (18.6 ML per 31 days)
RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML	T5	PA; QL (18.6 ML per 31 days)
RELISTOR SUBCUTANEOUS SYRINGE 8 MG/0.4 ML	T5	PA; QL (12.4 ML per 31 days)
SANCUSO	T5	
scopolamine base	T3	QL (10 EA per 30 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML)	T5	PA; QL (1.2 ML per 56 days)
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 360 MG/2.4 ML (150 MG/ML)	T5	PA; QL (2.4 ML per 56 days)
sodium,potassium,mag sulfates	T4	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
sucralfate	T2	
sulfasalazine	T2	
SYMPROIC	T4	PA; QL (31 EA per 31 days)
ursodiol oral capsule 300 mg	T3	
ursodiol oral tablet	T3	
VIBERZI	T5	PA; QL (62 EA per 31 days)
VOWST	T5	PA; QL (12 EA per 14 days)
ZENPEP ORAL CAPSULE,DELAYED RELEASE(DR/EC) 10,000-32,000 -42,000 UNIT, 15,000-47,000 -63,000 UNIT, 20,000-63,000- 84,000 UNIT, 25,000-79,000- 105,000 UNIT, 3,000-10,000 -14,000-UNIT, 40,000-126,000- 168,000 UNIT, 5,000-17,000- 24,000 UNIT, 60,000-189,600- 252,600 UNIT	T3	
Immunology, Vaccines / Biotechnology		
ABRYSVO (PF)	T3	QL (1 EA per 365 days)
ACTHIB (PF)	T3	
ACTIMMUNE	T5	PA
ADACEL(TDAP ADOLESN/ADULT)(PF)	T3	
AREXVY (PF)	T3	QL (1 EA per 365 days)
AVONEX INTRAMUSCULAR PEN INJECTOR KIT	T5	PA; QL (1 EA per 28 days)
AVONEX INTRAMUSCULAR SYRINGE KIT	T5	PA; QL (1 EA per 28 days)
bcg vaccine, live (pf)	T3	
BESREMI	T5	PA-NS; QL (2 ML per 28 days)
BESREMI PEN	T5	PA-NS; QL (1 ML per 28 days)
BETASERON SUBCUTANEOUS KIT	T5	PA; QL (14 EA per 28 days)
BEXSERO	T3	
BOOSTRIX TDAP INTRAMUSCULAR SYRINGE	T3	
DAPTACEL (DTAP PEDIATRIC) (PF)	T3	
ENGERIX-B (PF)	T3	PA-BvD
ENGERIX-B PEDIATRIC (PF)	T3	PA-BvD
GAMMAGARD LIQUID	T5	PA
GAMMAGARD LIQUID ERC	T5	PA
GAMMAGARD S-D (IGA < 1 MCG/ML)	T5	PA
GARDASIL 9 (PF)	T3	
GENOTROPIN	T5	PA

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
GENOTROPIN MINIQUICK SUBCUTANEOUS SYRINGE 0.2 MG/0.25 ML	T4	PA
GENOTROPIN MINIQUICK SUBCUTANEOUS SYRINGE 0.4 MG/0.25 ML, 0.6 MG/0.25 ML, 0.8 MG/0.25 ML, 1 MG/0.25 ML, 1.2 MG/0.25 ML, 1.4 MG/0.25 ML, 1.6 MG/0.25 ML, 1.8 MG/0.25 ML, 2 MG/0.25 ML	T5	PA
GRASTEK	T4	PA
HAVRIX (PF)	T3	
HEPLISAV-B (PF)	T3	PA-BvD
HIBERIX (PF)	T3	
HUMATROPE INJECTION CARTRIDGE	T5	PA
IMOVAX RABIES VACCINE (PF)	T3	PA-BvD
INFANRIX (DTAP) (PF)	T3	
IPOL	T3	
IXIARO (PF)	T3	
JYNNEOS (PF)	T3	PA-BvD
KINRIX (PF)	T3	
LEUKINE INJECTION RECON SOLN	T5	PA
MENQUADFI (PF)	T4	
MENVEO A-C-Y-W-135-DIP (PF) INTRAMUSCULAR KIT	T3	
M-M-R II (PF)	T3	
MRESVIA (PF)	T3	QL (1 ML per 365 days)
NIVESTYM	T5	
NORDITROPIN FLEXPRO SUBCUTANEOUS PEN INJECTOR 10 MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML (10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML)	T5	PA
ODACTRA	T4	PA
OMNITROPE SUBCUTANEOUS CARTRIDGE 10 MG/1.5 ML (6.7 MG/ML)	T5	PA
OMNITROPE SUBCUTANEOUS CARTRIDGE 5 MG/1.5 ML (3.3 MG/ML)	T4	PA
OMNITROPE SUBCUTANEOUS RECON SOLN	T5	PA
PEDIARIX (PF)	T3	
PEDVAX HIB (PF)	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
PEGASYS	T5	PA
PENBRAYA (PF)	T3	
PENMENVY MEN A-B-C-W-Y (PF)	T3	
PENTACEL (PF)	T3	
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML	T5	PA; QL (1 ML per 28 days)
PRIORIX (PF)	T3	
PRIVIGEN	T5	PA
PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML	T3	PA-BvD
PROCRIT INJECTION SOLUTION 20,000 UNIT/ML, 40,000 UNIT/ML	T5	PA-BvD
PROQUAD (PF)	T3	
QUADRACEL (PF)	T3	
RABAVERT (PF)	T3	PA-BvD
RECOMBIVAX HB (PF)	T3	PA-BvD
RETACRIT	T3	PA-BvD
ROTARIX	T3	
ROTATEQ VACCINE	T3	
SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG	T5	PA
SHINGRIX (PF)	T3	QL (2 EA per 999 days)
TENIVAC (PF)	T3	
TICOVAC	T3	
TRUMENBA	T3	
TWINRIX (PF)	T3	
TYPHIM VI INTRAMUSCULAR SYRINGE	T3	
VAQTA (PF)	T3	
VARIVAX (PF)	T3	
VAXCHORA VACCINE	T3	QL (200 ML per 365 days)
VIMKUNYA (PF)	T3	
VIVOTIF	T3	
XOLREMDI	T5	PA; QL (124 EA per 31 days)
YF-VAX (PF)	T3	
ZARXIO	T5	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
Miscellaneous Supplies		
ASSURE ID INSULIN SAFETY SYRINGE 1 ML 29 GAUGE X 1/2"	T4	
GAUZE PAD TOPICAL BANDAGE 2 X 2 "	T2	PA
insulin syringe-needle u-100 syringe 0.3 ml 29 gauge, 1 ml 29 gauge x 1/2", 1/2 ml 28 gauge	T3	
pen needle, diabetic needle 29 gauge x 1/2"	T4	
Musculoskeletal / Rheumatology		
ACTEMRA ACTPEN	T5	PA; QL (3.6 ML per 28 days)
ACTEMRA SUBCUTANEOUS	T5	PA; QL (3.6 ML per 28 days)
alendronate oral solution	T1	
alendronate oral tablet 10 mg, 35 mg, 70 mg	T1	
allopurinol oral tablet 100 mg, 300 mg	T1	
BENLYSTA SUBCUTANEOUS	T5	PA; QL (4 ML per 28 days)
colchicine oral tablet	T2	QL (62 EA per 31 days)
ENBREL MINI	T5	PA; QL (8 ML per 28 days)
ENBREL SUBCUTANEOUS SOLUTION	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5)	T5	PA; QL (4 ML per 28 days)
ENBREL SUBCUTANEOUS SYRINGE 50 MG/ML (1 ML)	T5	PA; QL (8 ML per 28 days)
ENBREL SURECLICK	T5	PA; QL (8 ML per 28 days)
ENOBY	T3	PA; QL (1 ML per 180 days)
HADLIMA	T5	PA; QL (1.6 ML per 28 days)
HADLIMA PUSHTOUCH	T5	PA; QL (1.6 ML per 28 days)
HADLIMA(CF)	T5	PA; QL (0.8 ML per 28 days)
HADLIMA(CF) PUSHTOUCH	T5	PA; QL (0.8 ML per 28 days)
ibandronate oral	T2	
JUBBONTI	T3	PA; QL (1 ML per 180 days)
KINERET	T5	PA; QL (18.76 ML per 28 days)
leflunomide	T2	
milnacipran oral tablet	T4	PA
ORENCIA CLICKJECT	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML	T5	PA; QL (4 ML per 28 days)
ORENCIA SUBCUTANEOUS SYRINGE 50 MG/0.4 ML	T5	PA; QL (1.6 ML per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
ORENCIA SUBCUTANEOUS SYRINGE 87.5 MG/0.7 ML	T5	PA; QL (2.8 ML per 28 days)
OTEZLA	T5	PA; QL (62 EA per 31 days)
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47)	T5	PA; QL (110 EA per 365 days)
OTEZLA XR	T5	PA; QL (30 EA per 30 days)
OTEZLA XR INITIATION	T5	PA; QL (82 EA per 365 days)
penicillamine oral tablet	T5	
probenecid	T2	
probenecid-colchicine	T2	
PROLIA	T4	PA; QL (1 ML per 180 days)
raloxifene	T3	
RASUVO (PF)	T4	PA
RINVOQ LQ	T5	PA; QL (372 ML per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG	T5	PA; QL (31 EA per 31 days)
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 45 MG	T5	PA; QL (168 EA per 365 days)
risedronate oral tablet 150 mg, 35 mg, 35 mg (12 pack), 35 mg (4 pack), 5 mg	T3	
risedronate oral tablet,delayed release (dr/ec)	T3	
SAVELLA ORAL TABLETS,DOSE PACK	T4	PA
SIMLANDI(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
SIMLANDI(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML	T5	PA; QL (2 EA per 28 days)
teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)	T5	PA; QL (2.48 ML per 28 days)
tofacitinib oral solution	T5	PA; QL (310 ML per 31 days)
tofacitinib oral tablet	T4	PA; QL (62 EA per 31 days)
tofacitinib oral tablet extended release 24 hr 11 mg	T4	PA; QL (31 EA per 31 days)
tofacitinib oral tablet extended release 24 hr 22 mg	T5	PA; QL (31 EA per 31 days)
TYMLOS	T5	PA; QL (1.56 ML per 30 days)
YUFLYMA(CF) AI CROHN'S-UC-HS	T5	PA; QL (6 EA per 365 days)
YUFLYMA(CF) AUTOINJECTOR	T5	PA; QL (2 EA per 28 days)
YUFLYMA(CF) SUBCUTANEOUS SYRINGE KIT 40 MG/0.4 ML	T5	PA; QL (2 EA per 28 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
Obstetrics / Gynecology		
ABIGALE	T2	
ALTAVERA (28)	T2	
ALYACEN 1/35 (28)	T2	
ANGELIQ ORAL TABLET 0.5-1 MG	T4	
APRI	T2	
ARANELLE (28)	T2	
AVIANE	T2	
AZURETTE (28)	T2	
BALZIVA (28)	T2	
BLISOVI FE 1.5/30 (28)	T2	
BRIELLYN	T2	
CAMILA	T2	
CLEOCIN VAGINAL SUPPOSITORY	T4	
CLIMARA PRO	T4	
clindamycin phosphate vaginal	T2	
CLINDESSE	T4	
CRYSELLE (28)	T2	
CYRED EQ	T2	
DEPO-ESTRADIOL	T4	
DEPO-SUBQ PROVERA 104	T3	
DOLISHALE	T4	
DOTTI	T2	
drospirenone-ethinyl estradiol	T2	
ELURYNG	T3	
ENSKYCE	T2	
ERRIN	T2	
ESTARYLLA	T2	
estradiol oral	T1	
estradiol transdermal patch semiweekly	T2	
estradiol transdermal patch weekly	T2	
estradiol vaginal cream	T2	
estradiol vaginal tablet	T4	
estradiol-norethindrone acet	T2	
ESTRING	T4	
etonogestrel-ethinyl estradiol	T3	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
EVAMIST	T4	
FEMRING	T4	
FYAVOLV	T4	
GALLIFREY	T2	
HAILEY 24 FE	T2	
HAILEY FE 1/20 (28)	T2	
HEATHER	T2	
ICLEVIA	T2	
IMVEXXY MAINTENANCE PACK	T3	
IMVEXXY STARTER PACK	T3	
INCASSIA	T2	
INTROVALE	T2	
ISIBLOOM	T2	
JASMIEL (28)	T2	
JINTELI	T2	
JULEBER	T2	
JUNEL 1.5/30 (21)	T2	
JUNEL 1/20 (21)	T2	
JUNEL FE 1.5/30 (28)	T2	
JUNEL FE 1/20 (28)	T2	
JUNEL FE 24	T2	
KAITLIB FE	T2	
KARIVA (28)	T2	
KELNOR 1/35 (28)	T2	
KURVELO (28)	T2	
LESSINA	T2	
LEVONEST (28)	T2	
levonorgestrel-ethinyl estrad oral tablet 0.1-20 mg-mcg	T2	
levonorgestrel-ethinyl estrad oral tablet 90-20 mcg (28)	T4	
levonorgestrel-ethinyl estrad oral tablets,dose pack,3 month	T2	
levonorg-eth estrad triphasic	T2	
LILETTA	T3	
LORYNA (28)	T2	
LOW-OGESTREL (28)	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
LUIZZA	T2	
LYLEQ	T2	
LYLLANA	T2	
LYZA	T2	
MARLISSA (28)	T2	
medroxyprogesterone	T2	
metronidazole vaginal gel 0.75 % (37.5mg/5 gram)	T3	
MICONAZOLE-3 VAGINAL SUPPOSITORY	T2	
MICROGESTIN 1.5/30 (21)	T2	
MICROGESTIN 1/20 (21)	T2	
MICROGESTIN FE 1.5/30 (28)	T2	
MICROGESTIN FE 1/20 (28)	T2	
MILI	T2	
NECON 0.5/35 (28)	T2	
NEXPLANON	T3	
norethindrone (contraceptive)	T2	
norethindrone acetate	T2	
norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-20 mg-mcg, 1-5 mg-mcg	T2	
norgestimate-ethinyl estradiol oral tablet 0.18/0.215/0.25 mg-0.035mg (28), 0.25-0.035 mg	T2	
NORTREL 0.5/35 (28)	T2	
NORTREL 1/35 (21)	T2	
NORTREL 1/35 (28)	T2	
NORTREL 7/7/7 (28)	T2	
NYLIA 1/35 (28)	T2	
NYLIA 7/7/7 (28)	T2	
PIMTREA (28)	T2	
PORTIA 28	T2	
PREMARIN VAGINAL	T3	
progesterone micronized oral	T3	
RECLIPSEN (28)	T2	
SETLAKIN	T2	
SPRINTEC (28)	T2	
SYEDA	T2	
terconazole	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
tranexamic acid oral	T2	
TRI-ESTARYLLA	T2	
TRI-MILI	T2	
TRI-SPRINTEC (28)	T2	
TRI-VYLIBRA	T2	
TURQOZ (28)	T2	
VALTYA	T2	
VANDAZOLE	T3	
VELIVET TRIPHASIC REGIMEN (28)	T2	
VESTURA (28)	T2	
VIENVA	T2	
VIORELE (28)	T2	
VYFEMLA (28)	T2	
VYLIBRA	T2	
XELRIA FE	T2	
YUVAFEM	T4	
ZAFEMY	T3	
ZOVIA 1-35 (28)	T2	
Ophthalmology		
acetazolamide	T2	
ALPHAGAN P OPHTHALMIC (EYE) DROPS 0.1 %	T3	
apraclonidine	T2	
atropine ophthalmic (eye) drops 1 %	T2	
AZASITE	T4	
azelastine ophthalmic (eye)	T2	
bacitracin-polymyxin b	T2	
BESIVANCE	T4	
betaxolol ophthalmic (eye)	T2	
bimatoprost ophthalmic (eye) drops 0.03 %	T2	
brimonidine ophthalmic (eye) drops 0.1 %	T3	
brimonidine ophthalmic (eye) drops 0.15 %, 0.2 %	T2	
brimonidine-timolol	T3	
brinzolamide	T4	
bromfenac ophthalmic (eye) drops 0.09 %	T2	
carteolol	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
CILOXAN OPHTHALMIC (EYE) OINTMENT	T3	
ciprofloxacin hcl ophthalmic (eye)	T1	
COMBIGAN	T3	
cromolyn ophthalmic (eye)	T2	
cyclosporine ophthalmic (eye)	T3	QL (60 EA per 30 days)
CYSTADROPS	T5	PA; QL (20 ML per 28 days)
CYSTARAN	T5	PA; QL (60 ML per 28 days)
dexamethasone sodium phosphate ophthalmic (eye)	T2	
diclofenac sodium ophthalmic (eye)	T1	
difluprednate	T4	
dorzolamide	T2	
dorzolamide-timolol	T2	
epinastine	T2	
erythromycin ophthalmic (eye)	T2	
fluorometholone	T2	
flurbiprofen sodium	T2	
gatifloxacin	T3	
gentamicin ophthalmic (eye) drops	T1	
ketorolac ophthalmic (eye)	T2	
latanoprost	T1	
levobunolol ophthalmic (eye) drops 0.5 %	T1	
levofloxacin ophthalmic (eye) drops 0.5 %	T3	
LUMIGAN OPHTHALMIC (EYE) DROPS 0.01 %	T3	QL (5 ML per 31 days)
methazolamide	T2	
moxifloxacin ophthalmic (eye) drops	T3	
neomycin-bacitracin-poly-hc	T2	
neomycin-bacitracin-polymyxin	T2	
neomycin-polymyxin b-dexameth	T2	
neomycin-polymyxin-gramicidin	T2	
neomycin-polymyxin-hc ophthalmic (eye)	T2	
ofloxacin ophthalmic (eye)	T2	
pilocarpine hcl ophthalmic (eye) drops 1 %, 2 %, 4 %	T2	
polymyxin b sulf-trimethoprim	T1	
prednisolone acetate	T2	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
prednisolone sodium phosphate ophthalmic (eye)	T2	
RESTASIS	T3	QL (60 EA per 30 days)
RESTASIS MULTIDOSE	T3	QL (5.5 ML per 27 days)
RHOPRESSA	T4	ST
ROCKLATAN	T4	ST
SIMBRINZA	T4	
sulfacetamide sodium ophthalmic (eye) drops	T2	
sulfacetamide-prednisolone	T2	
tafluprost (pf)	T4	
timolol maleate ophthalmic (eye) drops	T1	
timolol maleate ophthalmic (eye) drops, once daily	T4	
timolol maleate ophthalmic (eye) gel forming solution	T2	
TOBRADEX	T3	
TOBRADEX ST	T3	
tobramycin ophthalmic (eye)	T1	
tobramycin-dexamethasone	T2	
tobramycin-lotepred	T4	
TOBREX OPHTHALMIC (EYE) OINTMENT	T3	
travoprost	T3	
trifluridine	T2	
XDEMVY	T5	PA; QL (10 ML per 42 days)
XIIDRA	T3	QL (60 EA per 30 days)
ZIRGAN	T4	ST
Respiratory And Allergy		
acetylcysteine	T2	PA-BvD
ADEMPAS	T5	PA; QL (93 EA per 31 days)
ADVAIR HFA	T3	QL (12 GM per 30 days)
albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation	T3	QL (17 GM per 30 days)
albuterol sulfate inhalation hfa aerosol inhaler 90 mcg/actuation (nda020503)	T3	QL (13.4 GM per 30 days)
albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg /3 ml (0.083 %), 2.5 mg/0.5 ml	T2	PA-BvD
albuterol sulfate oral syrup	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
albuterol sulfate oral tablet	T1	
ambrisentan	T5	PA; QL (31 EA per 31 days)
ANORO ELLIPTA	T3	QL (60 EA per 30 days)
arformoterol	T4	PA-BvD
ASMANEX HFA	T3	QL (13 GM per 30 days)
ASMANEX TWISTHALER INHALATION AEROSOL POWDR BREATH ACTIVATED 110 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (120), 220 MCG/ ACTUATION (30), 220 MCG/ ACTUATION (60)	T3	QL (1 EA per 30 days)
ATROVENT HFA	T3	QL (25.8 GM per 30 days)
azelastine-fluticasone	T4	QL (23 GM per 30 days)
bosentan oral tablet	T5	PA; QL (62 EA per 31 days)
BREO ELLIPTA	T3	QL (60 EA per 30 days)
BREYNA	T3	QL (10.3 GM per 30 days)
BREZTRI AEROSPHERE	T3	QL (10.7 GM per 30 days)
BRINSUPRI	T5	PA; QL (31 EA per 31 days)
budesonide inhalation	T4	PA-BvD
budesonide-formoterol	T3	QL (10.2 GM per 30 days)
cetirizine oral solution 1 mg/ml	T1	QL (310 ML per 31 days)
CINRYZE	T5	PA; QL (20 EA per 28 days)
CLARINEX-D 12 HOUR	T4	
COMBIVENT RESPIMAT	T3	QL (4 GM per 30 days)
cromolyn inhalation	T4	PA-BvD
cyproheptadine oral syrup	T2	PA
cyproheptadine oral tablet	T2	PA
desloratadine oral tablet	T2	QL (31 EA per 31 days)
DULERA	T3	QL (13 GM per 30 days)
epinephrine injection auto-injector	T3	
FASENRA PEN	T5	PA; QL (1 ML per 28 days)
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML	T5	PA; QL (0.5 ML per 56 days)
FASENRA SUBCUTANEOUS SYRINGE 30 MG/ML	T5	PA; QL (1 ML per 28 days)
flunisolide	T2	QL (50 ML per 25 days)
fluticasone furoate-vilanterol	T3	QL (60 EA per 30 days)
fluticasone propionate inhalation blister with device 100 mcg/actuation, 50 mcg/actuation	T4	ST; QL (60 EA per 30 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
fluticasone propionate inhalation blister with device 250 mcg/actuation	T4	ST; QL (240 EA per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 110 mcg/actuation	T4	ST; QL (12 GM per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 220 mcg/actuation	T4	ST; QL (24 GM per 30 days)
fluticasone propionate inhalation hfa aerosol inhaler 44 mcg/actuation	T4	ST; QL (10.6 GM per 30 days)
fluticasone propionate nasal	T1	QL (16 GM per 30 days)
fluticasone propion-salmeterol inhalation blister with device	T3	QL (60 EA per 30 days)
HAEGARDA	T5	PA
hydroxyzine hcl oral solution 10 mg/5 ml	T2	PA
hydroxyzine hcl oral tablet	T2	PA
icatibant	T5	PA; QL (18 ML per 30 days)
ipratropium bromide inhalation solution	T1	PA-BvD
ipratropium-albuterol	T2	PA-BvD
JASCAYD	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 5.8 MG, 50 MG, 75 MG	T5	PA; QL (56 EA per 28 days)
KALYDECO ORAL GRANULES IN PACKET 25 MG	T5	PA; QL (62 EA per 31 days)
KALYDECO ORAL TABLET	T5	PA; QL (62 EA per 31 days)
levalbuterol hcl inhalation solution for nebulization 0.31 mg/3 ml, 0.63 mg/3 ml, 1.25 mg/3 ml	T2	PA-BvD
levalbuterol tartrate	T3	QL (30 GM per 30 days)
levocetirizine oral solution	T4	QL (310 ML per 31 days)
levocetirizine oral tablet	T1	QL (31 EA per 31 days)
macitentan	T4	PA; QL (31 EA per 31 days)
mometasone nasal	T3	QL (34 GM per 30 days)
montelukast oral tablet	T2	QL (31 EA per 31 days)
montelukast oral tablet,chewable	T2	QL (31 EA per 31 days)
nintedanib	T5	PA; QL (62 EA per 31 days)
OPSUMIT	T5	PA; QL (31 EA per 31 days)
OPSYNVI	T5	PA; QL (31 EA per 31 days)
ORKAMBI ORAL GRANULES IN PACKET	T5	PA; QL (62 EA per 31 days)
ORKAMBI ORAL TABLET	T5	PA; QL (124 EA per 31 days)
pirfenidone oral capsule	T5	PA; QL (279 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
pirfenidone oral tablet	T5	PA; QL (93 EA per 31 days)
promethazine oral syrup	T2	PA
promethazine oral tablet	T4	PA
promethazine rectal	T2	
PULMOZYME	T5	PA
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 40 MCG/ACTUATION	T3	QL (10.6 GM per 30 days)
QVAR REDIHALER INHALATION HFA AEROSOL BREATH ACTIVATED 80 MCG/ACTUATION	T3	QL (21.2 GM per 30 days)
roflumilast oral tablet 250 mcg	T4	QL (31 EA per 31 days)
roflumilast oral tablet 500 mcg	T3	QL (31 EA per 31 days)
SAJAZIR	T5	PA; QL (18 ML per 30 days)
SEREVENT DISKUS	T3	QL (60 EA per 30 days)
sildenafil (pulm.hypertension) oral tablet	T3	PA; QL (372 EA per 31 days)
SPIRIVA RESPIMAT	T3	QL (4 GM per 30 days)
STIOLTO RESPIMAT	T3	QL (4 GM per 30 days)
STRIVERDI RESPIMAT	T4	QL (4 GM per 30 days)
tadalafil (pulm. hypertension)	T5	PA; QL (62 EA per 31 days)
terbutaline oral	T4	
TEZSPIRE SUBCUTANEOUS PEN INJECTOR	T5	PA; QL (1.91 ML per 28 days)
THEO-24	T3	
theophylline oral solution	T2	
theophylline oral tablet extended release 12 hr	T2	
theophylline oral tablet extended release 24 hr	T2	
TRELEGY ELLIPTA	T3	QL (60 EA per 30 days)
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL	T5	PA; QL (56 EA per 28 days)
TRIKAFTA ORAL TABLETS, SEQUENTIAL	T5	PA; QL (84 EA per 28 days)
VENTOLIN HFA	T3	QL (36 GM per 30 days)
WINREVAIR	T5	PA
WIXELA INHUB	T3	QL (60 EA per 30 days)
XOLAIR	T5	PA
YUPELRI	T5	PA-BvD
zafirlukast oral tablet 10 mg	T4	QL (93 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
zafirlukast oral tablet 20 mg	T4	QL (62 EA per 31 days)
zileuton	T5	PA; QL (124 EA per 31 days)
Urologicals		
alfuzosin	T2	QL (31 EA per 31 days)
bethanechol chloride oral tablet 10 mg, 25 mg, 5 mg	T2	
bethanechol chloride oral tablet 50 mg	T3	
CYSTAGON	T3	
darifenacin	T3	QL (31 EA per 31 days)
dutasteride	T2	QL (31 EA per 31 days)
dutasteride-tamsulosin	T3	QL (31 EA per 31 days)
ELMIRON	T4	
finasteride oral tablet 5 mg	T2	
flavoxate	T2	
GEMTESA	T4	QL (31 EA per 31 days)
MYRBETRIQ ORAL SUSPENSION,EXTENDED REL RECON	T3	QL (300 ML per 30 days)
MYRBETRIQ ORAL TABLET EXTENDED RELEASE 24 HR	T3	QL (31 EA per 31 days)
oxybutynin chloride oral syrup	T2	
oxybutynin chloride oral tablet	T2	
oxybutynin chloride oral tablet extended release 24hr 10 mg, 5 mg	T2	QL (31 EA per 31 days)
oxybutynin chloride oral tablet extended release 24hr 15 mg	T3	QL (62 EA per 31 days)
OXYTROL	T4	QL (8 EA per 28 days)
potassium citrate oral tablet extended release	T2	
PROCYSBI ORAL GRANULES DEL RELEASE IN PACKET	T5	PA
RIVFLOZA SUBCUTANEOUS SOLUTION	T5	PA; QL (1 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML	T5	PA; QL (0.8 ML per 28 days)
RIVFLOZA SUBCUTANEOUS SYRINGE 160 MG/ML	T5	PA; QL (1 ML per 28 days)
silodosin	T4	
solifenacin	T4	QL (31 EA per 31 days)
tadalafil oral tablet 2.5 mg	T4	PA; QL (62 EA per 31 days)
tadalafil oral tablet 5 mg	T4	PA; QL (31 EA per 31 days)

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
tamsulosin	T1	
tolterodine oral capsule,extended release 24hr	T3	QL (31 EA per 31 days)
tolterodine oral tablet	T3	QL (62 EA per 31 days)
trospium oral capsule,extended release 24hr	T4	QL (31 EA per 31 days)
trospium oral tablet	T2	QL (93 EA per 31 days)
Vitamins, Hematinics / Electrolytes		
calcium acetate(phosphat bind)	T2	PA-BvD
CLINIMIX 5%/D15W SULFITE FREE	T4	PA-BvD
CLINIMIX 4.25%/D10W SULF FREE	T4	PA-BvD
CLINIMIX 5%-D20W(SULFITE-FREE)	T4	PA-BvD
CLINIMIX E 4.25%/D10W SUL FREE	T4	PA-BvD
CLINIMIX E 4.25%/D5W SULF FREE	T4	PA-BvD
CLINIMIX E 5%/D15W SULFIT FREE	T4	PA-BvD
CLINIMIX E 5%/D20W SULFIT FREE	T4	PA-BvD
fluoride (sodium) oral tablet	T2	
INTRALIPID INTRAVENOUS EMULSION 20 %	T4	PA-BvD
ISOLYTE S PH 7.4	T3	PA-BvD
ISOLYTE-P IN 5 % DEXTROSE	T4	PA-BvD
KLOR-CON	T4	
KLOR-CON 10	T3	
KLOR-CON 8	T3	
KLOR-CON M10	T1	
KLOR-CON M15	T1	
KLOR-CON M20	T1	
magnesium sulfate injection	T2	
PLENAMINE	T3	PA-BvD
potassium chlorid-d5-0.45%nacl	T2	
potassium chloride in 0.9%nacl intravenous parenteral solution 20 meq/l, 40 meq/l	T2	
potassium chloride in 5 % dex intravenous parenteral solution 20 meq/l	T2	
potassium chloride in lr-d5	T2	
potassium chloride in water intravenous piggyback 10 meq/100 ml, 20 meq/100 ml, 40 meq/100 ml	T2	
potassium chloride intravenous	T2	
potassium chloride oral capsule, extended release	T1	

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Drug Name	Drug Tier	Requirements/Limits
potassium chloride oral liquid	T2	
potassium chloride oral tablet extended release 10 meq, 20 meq, 8 meq	T1	
potassium chloride oral tablet,er particles/crystals	T1	
potassium chloride-0.45 % nacl	T2	
potassium chloride-d5-0.2%nacl	T2	
potassium chloride-d5-0.9%nacl	T2	
PRENATAL VITAMIN PLUS LOW IRON	T2	PA
PROSOL 20 %	T4	PA-BvD
sodium chloride 0.45 % intravenous	T2	
sodium chloride 3 % hypertonic	T2	
sodium chloride 5 % hypertonic	T2	
TRAVASOL 10 %	T4	PA-BvD
TROPHAMINE 10 %	T4	PA-BvD

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

You can find information on what the symbols and abbreviations on this table mean by going to the beginning of this table.

Index of Drugs

abacavir 3
abacavir-lamivudine 3
ABIGALE 64
ABILIFY MAINTENA 20
abiraterone 12
ABIRTEGA 12
ABRYSVO (PF) 59
acamprosate 49
acarbose 51
ACCUTANE 43
acebutolol 37
acetaminophen-codeine 20
acetazolamide 67
acetic acid 50
acetylcysteine 69
acitretin 43
ACTEMRA 62
ACTEMRA ACTPEN 62
ACTHIB (PF) 59
ACTIMMUNE 59
acyclovir 3, 43
acyclovir sodium 3
ADACEL(TDAP ADOLESN/ADULT)(PF) 59
adapalene 43
adapalene-benzoyl peroxide 43
adefovir 3
ADEMPAS 69
ADVAIR HFA 69
AIMOVIG AUTOINJECTOR 21
AKEEGA 12
ALA-CORT 43
albendazole 3
albuterol sulfate 69, 70
alclometasone 43
ALCOHOL PADS 51
ALECENSA 12
alendronate 62
alfuzosin 73
aliskiren 37
allopurinol 62
almotriptan malate 21
alosetron 56
ALPHAGAN P 67
alprazolam 21
ALPRAZOLAM INTENSOL ..21
ALTAVERA (28) 64
ALUNBRIG 12
ALYACEN 1/35 (28) 64
amantadine hcl 3
ambrisentan 70
amcinonide 44
amikacin 3
amiloride 37
amiloride-hydrochlorothiazide ... 37
amiodarone 37
amitriptyline 21
amitriptyline-chlordiazepoxide ... 21
amlodipine 37
amlodipine-atorvastatin 37
amlodipine-benazepril 37
amlodipine-olmesartan 37
amlodipine-valsartan 37
ammonium lactate 44
AMNESTEEM 44
amoxapine 21
amoxicil-clarithromy-lansopraz . 56
amoxicillin 3
amoxicillin-pot clavulanate 3, 4
amphotericin b 4
amphotericin b liposome 4
ampicillin 4
ampicillin sodium 4
ampicillin-sulbactam 4
anagrelide 49
anastrozole 12
ANGELIQ 64
ANORO ELLIPTA 70
ANZUPGO 44
apraclonidine 67
aprepitant 56
APRI 64
APTIVUS 4
ARALAST NP 49
ARANELLE (28) 64
AREXVY (PF) 59
arformoterol 70
ARIKAYCE 4
aripiprazole 21
ARISTADA 21
ARISTADA INITIO 21
armodafinil 21
asenapine maleate 21
ASMANEX HFA 70
ASMANEX TWISTHALER 70
aspirin-dipyridamole 37
ASSURE ID INSULIN SAFETY 62
atazanavir 4
atenolol 37
atenolol-chlorthalidone 37
atomoxetine 21, 22
atorvastatin 37
atovaquone 4
atovaquone-proguanil 4
atropine 67
ATROVENT HFA 70
ATTRUBY 37
AUGTYRO 12
AUSTEDO 22
AUSTEDO XR 22
AUSTEDO XR TITRATION KT(WK1-4) 22
AUVELITY 22
AVEED 51
AVIANE 64
AVMAPKI-FAKZYNJA 12
AVONEX 59
AVYCAZ 4
AYVAKIT 12
AZASITE 67
azathioprine 12
azelaic acid 44
azelastine 51, 67
azelastine-fluticasone 70
AZELEX 44
azithromycin 4
aztreonam 4
AZURETTE (28) 64
bacitracin-polymyxin b 67
baclofen 22
BAFIERTAM 22
balsalazide 56
BALVERSA 12
BALZIVA (28) 64
bcg vaccine, live (pf) 59
benazepril 37
benazepril-hydrochlorothiazide .. 38
BENLYSTA 62
benztropine 22
BESIVANCE 67
BESREMI 59
BESREMI PEN 59
betaine 56
betamethasone dipropionate 44
betamethasone valerate 44
betamethasone, augmented 44
BETASERON 59
betaxolol 38, 67
bethanechol chloride 73
bexarotene 12
BEXSERO 59
bicalutamide 12

BICILLIN C-R 4
BICILLIN L-A 4
BIKTARVY 4
bimatoprost 67
bisoprolol fumarate 38
bisoprolol-hydrochlorothiazide .. 38
BLISOVI FE 1.5/30 (28) 64
BLUJEPA 4
BOOSTRIX TDAP 59
bosentan 70
BOSULIF 12
bosutinib 12
BRAFTOVI 12
BREO ELLIPTA 70
BREYNA 70
BREZTRI AEROSPHERE 70
BRIELLYN 64
brimonidine 67
brimonidine-timolol 67
BRINSUPRI 70
brinzolamide 67
brivaracetam 22
bromfenac 67
bromocriptine 22
BRUKINSA 12
budesonide 56, 70
budesonide-formoterol 70
bumetanide 38
buprenorphine 22
buprenorphine hcl 22
buprenorphine-naloxone 22
bupropion hcl 22
bupropion hcl (smoking deter) 49
buspirone 22
butalbital-acetaminop-caf-cod 22
butalbital-acetaminophen 22
butalbital-acetaminophen-caff 22
butalbital-aspirin-caffeine 23
butorphanol 23
cabergoline 51
CABOMETYX 12
calcipotriene 44
calcipotriene-betamethasone 44
calcitonin (salmon) 51
calcitriol 44, 51
calcium acetate(phosphat bind) .. 74
CALQUENCE (ACALABRUTINIB MAL) 12
CAMILA 64
candesartan 38
candesartan-hydrochlorothiazid .. 38
CAPLYTA 23
CAPRELSA 12
captopril 38
carbamazepine 23
carbidopa-levodopa 23
carbidopa-levodopa-entacapone 23
carglumic acid 49
carteolol 67
CARTIA XT 38
carvedilol 38
carvedilol phosphate 38
caspofungin 4
CAYSTON 4
cefaclor 4
cefadroxil 4
cefazolin 4
cefdinir 4
cefepime 4
cefixime 4
cefotetan 4
cefoxitin 4
cefpodoxime 5
cefprozil 5
ceftaroline fosamil 5
ceftazidime 5
ceftriaxone 5
cefuroxime axetil 5
cefuroxime sodium 5
celecoxib 23
cephalexin 5
CERDELGA 51
cetirizine 70
cevimeline 49
CHEMET 49
chlorhexidine gluconate 51
chloroquine phosphate 5
chlorpromazine 23
chlorthalidone 38
cholestyramine (with sugar) 38
CHOLESTYRAMINE LIGHT 38
ciclopirox 44
cilostazol 38
CILOXAN 68
CIMDUO 5
cimetidine 56
cimetidine hcl 56
cinacalcet 51
CINRYZE 70
ciprofloxacin hcl 5, 68
ciprofloxacin in 5 % dextrose 5
ciprofloxacin-dexamethasone 51
citalopram 23
CLARAVIS 44
CLARINEX-D 12 HOUR 70
clarithromycin 5
CLEOCIN 64
CLIMARA PRO 64
CLINDACIN 44
clindamycin hcl 5
clindamycin in 5 % dextrose 5
CLINDAMYCIN PEDIATRIC 5
clindamycin phosphate 5, 44, 64
clindamycin-benzoyl peroxide 44
clindamycin-tretinoin 44
CLINDESSE 64
CLINIMIX 5%/D15W SULFITE FREE 74
CLINIMIX 4.25%/D10W SULF FREE 74
CLINIMIX 4.25%/D5W SULFIT FREE 49
CLINIMIX 5%-D20W(SULFITE-FREE) 74
CLINIMIX E 2.75%/D5W SULF FREE 49
CLINIMIX E 4.25%/D10W SUL FREE 74
CLINIMIX E 4.25%/D5W SULF FREE 74
CLINIMIX E 5%/D15W SULFIT FREE 74
CLINIMIX E 5%/D20W SULFIT FREE 74
clobazam 23
clobetasol 44, 45
clobetasol-emollient 45
clocortolone pivalate 45
CLODAN 45
clomipramine 23
clonazepam 23
clonidine 38
clonidine hcl 23, 38
clopidogrel 38
clorazepate dipotassium 23
clotrimazole 5, 45
clotrimazole-betamethasone 45
clozapine 23, 24
COARTEM 5
COBENFY 24
COBENFY STARTER PACK .24
colchicine 62
colesevelam 38
colestipol 38

colistin (colistimethate na) 5
COMBIGAN 68
COMBIVENT RESPIMAT 70
COMETRIQ 12, 13
COMPRO 56
CONSTULOSE 56
COPAXONE 24
COPIKTRA 13
COSENTYX 45
COSENTYX (2 SYRINGES) 45
COSENTYX PEN (2 PENS) 45
COSENTYX UNOREADY PEN 45
COTELLIC 13
CREON 56
CRESEMBA 5
cromolyn 56, 68, 70
CRYSELLE (28) 64
cyclobenzaprine 24
cyclophosphamide 13
cycloserine 5
cyclosporine 13, 68
cyclosporine modified 13
cyproheptadine 70
CYRED EQ 64
CYSTADROPS 68
CYSTAGON 73
CYSTARAN 68
d10 %-0.45 % sodium chloride 49
d2.5 %-0.45 % sodium chloride 49
d5 % and 0.9 % sodium chloride 49
d5 %-0.45 % sodium chloride 49
dabigatran etexilate 38
dalbavancin 5
dalfampridine 24
danazol 51
dantrolene 24
DANZITEN 13
dapagliflozin 51
dapsone 5, 45
DAPTACEL (DTAP PEDIATRIC) (PF) 59
daptomycin 5
darifenacin 73
darunavir 5, 6
dasatinib 13
DAURISMO 13
deferasirox 49
DELSTRIGO 6
demeclocycline 6
DEPO-ESTRADIOL 64
DEPO-SUBQ PROVERA 104 64
DEPO-TESTOSTERONE 51
DESCOVY 6
desipramine 24
desloratadine 70
desmopressin 51
desonide 45
desoximetasone 45
desvenlafaxine 24
desvenlafaxine succinate 24
dexamethasone 51
dexamethasone sodium phosphate 68
dexmethylphenidate 24
dextroamphetamine sulfate 24
dextroamphetamine-amphetamine 24
dextromethorphan-quinidine 24
dextrose 10 % in water (d10w) 49
dextrose 5 % in water (d5w) 49
dextrose 5%-0.2 % sod chloride 49
DIACOMIT 24, 25
diazepam 25
DIAZEPAM INTENSOL 25
diazoxide 51
diclofenac epolamine 25
diclofenac potassium 25
diclofenac sodium 25, 45, 68
diclofenac-misoprostol 25
dicloxacillin 6
dicyclomine 57
DIFICID 6
diflunisal 25
difluprednate 68
digoxin 38
dihydroergotamine 25
DILANTIN 25
DILANTIN EXTENDED 25
diltiazem hcl 38
DILT-XR 38
dimethyl fumarate 25
diphenoxylate-atropine 57
disulfiram 49
divalproex 25
dofetilide 38
DOLISHALE 64
donepezil 25
dorzolamide 68
dorzolamide-timolol 68
DOTTI 64
DOVATO 6
doxazosin 39
doxepin 25, 45
doxercalciferol 52
DOXY-100 6
doxycycline hyclate 6
doxycycline monohydrate 6
DRIZALMA SPRINKLE 25, 26
dronabinol 57
drospirenone-ethinyl estradiol 64
droxidopa 49
DULERA 70
duloxetine 26
DUOPA 26
DUPIXENT PEN 45
DUPIXENT SYRINGE 45, 46
dutasteride 73
dutasteride-tamsulosin 73
DUVYZAT 49
econazole nitrate 46
EDARBYCLOR 39
EDURANT PED 6
efavirenz 6
efavirenz-emtricitabin-tenofov 6
efavirenz-lamivu-tenofov disop 6
eletriptan 26
ELIGARD 13
ELIGARD (3 MONTH) 13
ELIGARD (4 MONTH) 13
ELIGARD (6 MONTH) 13
ELIQUIS 39
ELIQUIS DVT-PE TREAT 30D START 39
ELIQUIS SPRINKLE 39
ELMIRON 73
eltrombopag olamine 39
ELURYNG 64
EMEND 57
EMGALITY PEN 26
EMGALITY SYRINGE 26
EMSAM 26
emtricitabine 6
emtricitabine-tenofovir (tdf) 6
emtricita-rilpivirine-tenof df 6
EMTRIVA 6
enalapril maleate 39
enalapril-hydrochlorothiazide 39
ENBREL 62
ENBREL MINI 62
ENBREL SURECLICK 62
ENGERIX-B (PF) 59
ENGERIX-B PEDIATRIC (PF) 59
ENOBY 62

enoxaparin 39
ENSACOVE 13
ENSKYCE 64
entacapone 26
entecavir 6
ENTRESTO 39
ENTRESTO SPRINKLE 39
ENULOSE 57
ENVARSUS XR 13
EOHILIA 57
EPIDIOLEX 26
epinastine 68
epinephrine 70
eplerenone 39
EQUETRO 26
ERAXIS(WATER DILUENT) .. 7
ergotamine-caffeine 26
ERIVEDGE 13
ERLEADA 13
erlotinib 13
ERRIN 64
ertapenem 7
ERY PADS 46
erythromycin 7, 68
erythromycin with ethanol 46
erythromycin-benzoyl peroxide ... 46
escitalopram oxalate 26
eslicarbazepine 26
esomeprazole magnesium 57
ESTARYLLA 64
estazolam 26
estradiol 64
estradiol-norethindrone acet 64
ESTRING 64
eszopiclone 26
ethacrynic acid 39
ethambutol 7
ethosuximide 26
etodolac 26
etonogestrel-ethinyl estradiol 64
etravirine 7
EULEXIN 13
EVAMIST 65
everolimus (antineoplastic) 13
everolimus (immunosuppressive) 13, 14
EVOTAZ 7
exemestane 14
EXXUA 26
ezetimibe 39
ezetimibe-simvastatin 39
FABHALTA 49
famciclovir 7
famotidine 57
FANAPT 27
FANAPT TITRATION PACK A 27
FANAPT TITRATION PACK B 27
FANAPT TITRATION PACK C 27
FASENRA 70
FASENRA PEN 70
felbamate 27
felodipine 39
FEMRING 65
fenofibrate 39
fenofibrate micronized 39
fenofibrate nanocrystallized 39
fenofibric acid (choline) 40
fenoprofen 27
fentanyl 27
FETZIMA 27
FIASP FLEXTOUCH U-100 INSULIN 52
FIASP PENFILL U-100 INSULIN 52
FIASP U-100 INSULIN 52
fidaxomicin 7
FILSUVEZ 46
FINACEA 46
finasteride 73
fingolimod 27
FINTEPLA 27
FIRMAGON KIT W DILUENT SYRINGE 14
flavoxate 73
flecainide 40
fluconazole 7
fluconazole in nacl (iso-osm) 7
flucytosine 7
fludrocortisone 52
flunisolide 70
fluocinolone 46
fluocinolone acetonide oil 51
fluocinolone and shower cap 46
fluocinonide 46
fluocinonide-emollient 46
fluoride (sodium) 74
fluorometholone 68
fluorouracil 46
fluoxetine 27
fluoxetine (pmdd) 27
fluphenazine decanoate 27
fluphenazine hcl 27
flurandrenolide 46
flurazepam 27
flurbiprofen 27
flurbiprofen sodium 68
fluticasone furoate-vilanterol 70
fluticasone propionate 46, 70, 71
fluticasone propion-salmeterol ... 71
fluvastatin 40
fluvoxamine 27
fondaparinux 40
fosamprenavir 7
fosfomycin tromethamine 7
fosinopril 40
fosinopril-hydrochlorothiazide ... 40
FOTIVDA 14
frovatriptan 27
FRUZAQLA 14
furosemide 40
FYAVOLV 65
gabapentin 27, 28
galantamine 28
GALLIFREY 65
GAMMAGARD LIQUID 59
GAMMAGARD LIQUID ERC 59
GAMMAGARD S-D (IGA < 1 MCG/ML) 59
GARDASIL 9 (PF) 59
gatifloxacin 68
GATTEX 30-VIAL 57
GAUZE PAD 62
GAVILYTE-C 57
GAVILYTE-G 57
GAVILYTE-N 57
GAVRETO 14
gefitinib 14
gemfibrozil 40
GEMTESA 73
GENERLAC 57
GENGRAF 14
GENOTROPIN 59
GENOTROPIN MINIQUICK 60
gentamicin 7, 46, 68
gentamicin in nacl (iso-osm) 7
GENVOYA 7
GILOTRIF 14
GLASSIA 49
glatiramer 28
GLATOPA 28
glimepiride 52
glipizide 52

glipizide-metformin 52
GLUCAGON EMERGENCY KIT (HUMAN) 52
glyburide 52
glyburide-metformin 52
glycerol phenylbutyrate 49
glycopyrrolate 57
GLYXAMBI 52
GOMEKLI 14
granisetron hcl 57
GRASTEK 60
griseofulvin microsize 7
griseofulvin ultramicrosize 7
guanfacine 28
GVOKE 52
GVOKE HYPOPEN 2-PACK 52
GVOKE PFS 1-PACK SYRINGE 52
HADLIMA 62
HADLIMA PUSHTOUCH 62
HADLIMA(CF) 62
HADLIMA(CF) PUSHTOUCH 62
HAEGARDA 71
HAILEY 24 FE 65
HAILEY FE 1/20 (28) 65
halcinonide 46
halobetasol propionate 46
haloperidol 28
haloperidol decanoate 28
haloperidol lactate 28
HAVRIX (PF) 60
HEATHER 65
heparin (porcine) 40
HEPCLUDEX 7
HEPLISAV-B (PF) 60
HERNEXEOS 14
HIBERIX (PF) 60
HUMALOG JUNIOR KWIKPEN U-100 52
HUMALOG KWIKPEN INSULIN 52
HUMALOG MIX 50-50 KWIKPEN 52
HUMALOG MIX 75-25 KWIKPEN 52
HUMALOG MIX 75-25(U-100)INSULN 52
HUMALOG U-100 INSULIN 52
HUMATROPE 60
HUMULIN 70/30 U-100 INSULIN 52
HUMULIN 70/30 U-100 KWIKPEN 52
HUMULIN N NPH INSULIN KWIKPEN 52
HUMULIN N NPH U-100 INSULIN 52
HUMULIN R REGULAR U-100 INSULN 52
HUMULIN R U-500 (CONC) KWIKPEN 52
hydralazine 40
hydrochlorothiazide 40
hydrocodone-acetaminophen 28
hydrocortisone 46, 52, 57
hydrocortisone butyrate 46
hydrocortisone valerate 46
hydrocortisone-acetic acid 51
hydrocortisone-pramoxine 57
hydromorphone 28
hydroxychloroquine 7
hydroxyurea 14
hydroxyzine hcl 71
HYRNUO 14
ibandronate 62
IBRANCE 14
IBTROZI 14
IBU 28
ibuprofen 28
icatibant 71
ICLEVIA 65
ICLUSIG 14
icosapent ethyl 40
IDHIFA 14
IDVYNSO 7
imatinib 14
IMBRUVICA 14
imipenem-cilastatin 7
imipramine hcl 28
imipramine pamoate 28
imiquimod 46
IMKELDI 14
IMOVAX RABIES VACCINE (PF) 60
IMVEXXY MAINTENANCE PACK 65
IMVEXXY STARTER PACK 65
INBRIJA 28
INCASSIA 65
INCRELEX 49
indapamide 40
indomethacin 28
INFANRIX (DTAP) (PF) 60
INGREZZA 28
INGREZZA INITIATION PK(TARDIV) 28
INGREZZA SPRINKLE 28
INLURIYO 14
INLYTA 14
INQOVI 14
INREBIC 14
insulin lispro 52
insulin lispro protamin-lispro 52
insulin syringe-needle u-100 62
INTELENCE 7
INTRALIPID 74
INTROVALE 65
INVEGA HAFYERA 28, 29
INVEGA SUSTENNA 29
INVEGA TRINZA 29
IPOL 60
ipratropium bromide 51, 71
ipratropium-albuterol 71
irbesartan 40
irbesartan-hydrochlorothiazide 40
ISENTRESS 8
ISENTRESS HD 7
ISIBLOOM 65
ISOLYTE S PH 7.4 74
ISOLYTE-P IN 5 % DEXTROSE 74
isoniazid 8
isosorbide dinitrate 40
isosorbide mononitrate 40
isosorbide-hydralazine 40
isotretinoin 47
isradipine 40
ISTURISA 52
ITOVEBI 14, 15
itraconazole 8
ivabradine 40
ivermectin 8
IWILFIN 15
IXIARO (PF) 60
JAKAFI 15
JANTOVEN 40
JANUMET 52
JANUMET XR 53
JANUVIA 53
JARDIANCE 53
JASCAYD 71
JASMIEL (28) 65
JAVYGTOR 53
JAYPIRCA 15
JENTADUETO 53

JENTADUETO XR 53
JIDEYTRO 15
JINTELI 65
JOENJA 49
JUBBONTI 62
JULEBER 65
JULUCA 8
JUNEL 1.5/30 (21) 65
JUNEL 1/20 (21) 65
JUNEL FE 1.5/30 (28) 65
JUNEL FE 1/20 (28) 65
JUNEL FE 24 65
JYNNEOS (PF) 60
KAITLIB FE 65
KALYDECO 71
KARIVA (28) 65
KELNOR 1/35 (28) 65
KERENDIA 40
KESIMPTA PEN 29
ketoconazole 8, 47
ketoprofen 29
ketorolac 29, 68
KINERET 62
KINRIX (PF) 60
KIONEX 49
KISQALI 15
KLISYRI (250 MG) 15
KLOR-CON 74
KLOR-CON 10 74
KLOR-CON 8 74
KLOR-CON M10 74
KLOR-CON M15 74
KLOR-CON M20 74
KLOXXADO 29
KOMZIFTI 15
KOSELUGO 15
KOURZEQ 51
KRAZATI 15
KURVELO (28) 65
KYGEVVI 49
labetalol 40
lacosamide 29
lactulose 57
LAGEVRIO (EUA) 8
lamivudine 8
lamivudine-zidovudine 8
lamotrigine 29
lansoprazole 57
lanthanum 50
LANTUS SOLOSTAR U-100 INSULIN 53
LANTUS U-100 INSULIN 53
lapatinib 15
latanoprost 68
LAZCLUZE 15
leflunomide 62
lenalidomide 15
LENVIMA 15
LESSINA 65
letrozole 15
leucovorin calcium 15
LEUKERAN 15
LEUKINE 60
leuprolide 15
levalbuterol hcl 71
levalbuterol tartrate 71
levetiracetam 29
levobunolol 68
levocarnitine 50
levocarnitine (with sugar) 50
levocetirizine 71
levofloxacin 8, 68
levofloxacin in d5w 8
LEVONEST (28) 65
levonorgestrel-ethinyl estrad 65
levonorg-eth estrad triphasic 65
levothyroxine 53
LEVOXYL 53
lidocaine 47
lidocaine hcl 47
lidocaine-prilocaine 47
LIFYORLI 15
LILETTA 65
linezolid 8
linezolid in dextrose 5% 8
LINZESS 57
LIOMNY 53
liothyronine 53
LIPOFEN 40
liraglutide 53
lisdexamfetamine 29
lisinopril 40
lisinopril-hydrochlorothiazide 40
LITFULO 50
lithium carbonate 29
lithium citrate 29
LITHOSTAT 50
LIVTENCITY 8
LOARGYS 50
lofexidine 29
LOKELMA 50
lomustine 16
LONSURF 16
loperamide 57
lopinavir-ritonavir 8
lorazepam 30
LORAZEPAM INTENSOL 29
LORBRENA 16
LORYNA (28) 65
losartan 40, 41
losartan-hydrochlorothiazide 41
lovastatin 41
LOW-OGESTREL (28) 65
loxapine succinate 30
lubiprostone 57
LUIZZA 66
LUMAKRAS 16
LUMIGAN 68
LUPRON DEPOT 16
LUPRON DEPOT (3 MONTH) 16
lurasidone 30
LYBALVI 30
LYLEQ 66
LYLLANA 66
LYNPARZA 16
LYSODREN 16
LYTGOBI 16
LYZA 66
macitentan 71
magnesium sulfate 74
malathion 47
maraviroc 8
MARLISSA (28) 66
MARPLAN 30
MATULANE 16
MATZIM LA 41
MAVYRET 8
meclizine 57
MEDROL 53
medroxyprogesterone 66
mefenamic acid 30
mefloquine 8
megestrol 16
MEKINIST 16
MEKTOVI 16
meloxicam 30
memantine 30
memantine-donepezil 30
MENQUADFI (PF) 60
MENVEO A-C-Y-W-135-DIP (PF) 60
mercaptopurine 16
meropenem 8
mesalamine 57, 58
mesna 16

metaxalone 30
metformin 53
methadone 30
methazolamide 68
methenamine hippurate 8
methimazole 53
methotrexate sodium 16
methotrexate sodium (pf) 16
methscopolamine 58
methsuximide 30
methylphenidate 30
methylphenidate hcl 30
methylprednisolone 53
metoclopramide hcl 58
metolazone 41
metoprolol succinate 41
metoprolol ta-hydrochlorothiaz .. 41
metoprolol tartrate 41
metronidazole 8, 47, 66
metronidazole in nacl (iso-os) 8
metyrosine 41
mexiletine 41
micafungin 8
MICONAZOLE-3 66
MICROGESTIN 1.5/30 (21) 66
MICROGESTIN 1/20 (21) 66
MICROGESTIN FE 1.5/30 (28) 66
MICROGESTIN FE 1/20 (28) . 66
midodrine 50
mifepristone 53
miglitol 53
miglustat 53
MILI 66
milnacipran 62
minocycline 8
minoxidil 41
mirtazapine 30, 31
misoprostol 58
M-M-R II (PF) 60
modafinil 31
MODEYSO 16
moexipril 41
molindone 31
mometasone 47, 71
montelukast 71
morphine 31
morphine concentrate 31
MOUNJARO 53
MOVANTIK 58
moxifloxacin 8, 68
moxifloxacin-sod.chloride(iso) 8
MRESVIA (PF) 60
MULPLETA 41
MULTAQ 41
mupirocin 47
mycophenolate mofetil 17
mycophenolate sodium 17
MYRBETRIQ 73
MYTESI 58
nabumetone 31
nadolol 41
nafcillin 9
naftifine 47
naloxone 31
naltrexone 31
naproxen 31
naproxen sodium 31
naratriptan 31
nateglinide 53
NAYZILAM 31
nebivolol 41
NECON 0.5/35 (28) 66
nefazodone 31
NEMLUVIO 17
neomycin 9
neomycin-bacitracin-poly-hc 68
neomycin-bacitracin-polymyxin .. 68
neomycin-polymyxin b-dexameth 68
neomycin-polymyxin-gramicidin 68
neomycin-polymyxin-hc 51, 68
NEO-SYNALAR 47
NERLYNX 17
NEUAC 47
NEUPRO 31
nevirapine 9
NEXLETOL 41
NEXLIZET 41
NEXPLANON 66
niacin 41
nicardipine 41
NICOTROL NS 50
nifedipine 41
nilotinib hcl 17
nilutamide 17
nimodipine 41
NINLARO 17
nintedanib 71
nisoldipine 41
nitazoxanide 9
nitisinone 50
NITRO-BID 41
nitrofurantoin macrocrystal 9
nitrofurantoin monohyd/m-cryst ... 9
nitroglycerin 41, 58
NIVESTYM 60
nizatidine 58
NORDITROPIN FLEXPRO ... 60
norethindrone (contraceptive) 66
norethindrone acetate 66
norethindrone ac-eth estradiol ... 66
norgestimate-ethinyl estradiol 66
NORTREL 0.5/35 (28) 66
NORTREL 1/35 (21) 66
NORTREL 1/35 (28) 66
NORTREL 7/7/7 (28) 66
nortriptyline 31
NOURIANZ 31
NOVOLIN 70/30 U-100 INSULIN 53
NOVOLIN 70-30 FLEXPEN U-100 54
NOVOLIN N FLEXPEN 54
NOVOLIN N NPH U-100 INSULIN 54
NOVOLIN R FLEXPEN 54
NOVOLIN R REGULAR U100 INSULIN 54
NOVOLOG FLEXPEN U-100 INSULIN 54
NOVOLOG MIX 70-30 U-100 INSULN 54
NOVOLOG MIX 70-30FLEXPEN U-100 54
NOVOLOG PENFILL U-100 INSULIN 54
NOVOLOG U-100 INSULIN ASPART 54
NUBEQA 17
NUCYNTA 31
NUEDEXTA 31
NUPLAZID 31
NURTEC ODT 31
NUZYRA 9
NYAMYC 47
NYLIA 1/35 (28) 66
NYLIA 7/7/7 (28) 66
nystatin 9, 47
nystatin-triamcinolone 47
NYSTOP 47
octreotide acetate 17
ODACTRA 60
ODEFSEY 9
ODOMZO 17
ofloxacin 9, 51, 68

OGSIVEO 17
OJEMDA 17
OJJAARA 17
olanzapine 31
olanzapine-fluoxetine 31
olmesartan 41
olmesartan-amlodipin-hcthiazid . 42
olmesartan-hydrochlorothiazide . 42
olopatadine 51
omega-3 acid ethyl esters 42
omeprazole 58
OMNITROPE 60
ondansetron 58
ondansetron hcl 58
ONGENTYS 31
ONUREG 17
ONZETRA XSAIL 32
OPIPZA 32
OPSUMIT 71
OPSYNVI 71
ORENCIA 62, 63
ORENCIA CLICKJECT 62
ORGOVYX 17
ORKAMBI 71
ORLYNVAH 9
ORSERDU 17
oseltamivir 9
OTEZLA 63
OTEZLA STARTER 63
OTEZLA XR 63
OTEZLA XR INITIATION 63
oxacillin 9
oxaprozin 32
oxazepam 32
oxcarbazepine 32
oxybutynin chloride 73
oxycodone 32
oxycodone-acetaminophen 32
OXYTROL 73
OZEMPIC 54
PACERONE 42
paliperidone 32
PALYNZIQ 54
PANRETIN 47
pantoprazole 58
paricalcitol 54
paroxetine hcl 32
PAXLOVID 9
pazopanib 17
PEDIARIX (PF) 60
PEDVAX HIB (PF) 60
peg 3350-electrolytes 58
peg3350-sod sul-nacl-kcl-asb-c .. 58
PEGASYS 61
peg-electrolyte soln 58
PEMAZYRE 17
pen needle, diabetic 62
PENBRAYA (PF) 61
penciclovir 47
penicillamine 63
penicillin g pot in dextrose 9
penicillin g potassium 9
penicillin g sodium 9
penicillin v potassium 9
PENMENVY MEN A-B-C-W-Y (PF) 61
PENTACEL (PF) 61
pentamidine 9
pentoxifylline 42
perampanel 32
perindopril erbumine 42
PERIOGARD 51
permethrin 47
perphenazine 32
perphenazine-amitriptyline 32
PHEBURANE 50
phenelzine 32
phenobarbital 32
PHENYTEK 32
phenytoin 32
phenytoin sodium extended 32
PIFELTRO 9
pilocarpine hcl 50, 68
pimecrolimus 47
pimozide 32
PIMTREA (28) 66
pindolol 42
pioglitazone 54
pioglitazone-metformin 54
piperacillin-tazobactam 10
PIQRAY 17
pirfenidone 71, 72
piroxicam 33
pitavastatin calcium 42
PLEGRIDY 61
PLENAMINE 74
podofilox 47
polymyxin b sulfate 10
polymyxin b sulf-trimethoprim 68
pomalidomide 18
PORTIA 28 66
posaconazole 10
potassium chlorid-d5-0.45%nacl 74
potassium chloride 74, 75
potassium chloride in 0.9%nacl . 74
potassium chloride in 5 % dex 74
potassium chloride in lr-d5 74
potassium chloride in water 74
potassium chloride-0.45 % nacl . 75
potassium chloride-d5-0.2%nacl 75
potassium chloride-d5-0.9%nacl 75
potassium citrate 73
pramipexole 33
prasugrel hcl 42
pravastatin 42
praziquantel 10
prazosin 42
prednisolone 54
prednisolone acetate 68
prednisolone sodium phosphate 54, 69
prednisone 54
pregabalin 33
PREMARIN 66
PRENATAL VITAMIN PLUS LOW IRON 75
PREVALITE 42
PREVYMIS 10
PREZCOBIX 10
PREZISTA 10
primaquine 10
primidone 33
PRIORIX (PF) 61
PRIVIGEN 61
probenecid 63
probenecid-colchicine 63
prochlorperazine 58
prochlorperazine maleate 58
PROCRIT 61
PROCTO-MED HC 58
PROCTOSOL HC 58
PROCTOZONE-HC 58
PROCYSBI 73
progesterone micronized 66
PROGRAF 18
PROLASTIN-C 50
PROLIA 63
promethazine 72
propafenone 42
propranolol 42
propylthiouracil 54
PROQUAD (PF) 61
PROSOL 20 % 75

protriptyline 33
PULMOZYME 72
pyrazinamide 10
pyridostigmine bromide 33
pyrimethamine 10
QINLOCK 18
QUADRACEL (PF) 61
quetiapine 33
quinapril 42
quinapril-hydrochlorothiazide 42
quinidine gluconate 42
quinidine sulfate 42
quinine sulfate 10
QULIPTA 33
QVAR REDIHALER 72
RABAVERT (PF) 61
rabeprazole 58
RADICAVA ORS STARTER KIT SUSP 33
RALDESY 33
raloxifene 63
ramelteon 33
ramipril 42
ranolazine 42
rasagiline 33
RASUVO (PF) 63
RECLIPSEN (28) 66
RECOMBIVAX HB (PF) 61
RELENZA DISKHALER 10
RELGAABI 33
RELISTOR 58
repaglinide 54
REPATHA SURECLICK 42
REPATHA SYRINGE 42
RESTASIS 69
RESTASIS MULTIDOSE 69
RETACRIT 61
RETEVMO 18
REVCOVI 50
REVUFORJ 18
REXULTI 33
REYATAZ 10
REZDIFFRA 50
REZENOPY 33
REZLIDHIA 18
RHOPRESSA 69
ribavirin 10
rifabutin 10
rifampin 10
rifapentine 10
rilpivirine hcl 10
riluzole 50
rimantadine 10
RINVOQ 63
RINVOQ LQ 63
risedronate 50, 63
risperidone 33, 34
risperidone microspheres 33
ritonavir 10
rivastigmine 34
rivastigmine tartrate 34
RIVFLOZA 73
rizatriptan 34
ROCKLATAN 69
roflumilast 72
ROMVIMZA 18
ropinirole 34
rosuvastatin 42
ROTARIX 61
ROTATEQ VACCINE 61
ROWEEPRA 34
ROZLYTREK 18
RUBRACA 18
rufinamide 34
RUKOBIA 10
RYDAPT 18
RYTARY 34
sacubitril-valsartan 42
SAJAZIR 72
SANCUSO 58
sapropterin 54
SAVELLA 63
SCEMBLIX 18
scopolamine base 58
SECUADO 34
selegiline hcl 34
selenium sulfide 47
SEREVENT DISKUS 72
SEROSTIM 61
sertraline 34
SETLAKIN 66
sevelamer carbonate 50
sevelamer hcl 50
SHINGRIX (PF) 61
SIGNIFOR 18
sildenafil (pulm.hypertension) 72
SILIQ 47
silodosin 73
silver sulfadiazine 47
SIMBRINZA 69
SIMLANDI(CF) 63
SIMLANDI(CF) AUTOINJECTOR 63
simvastatin 42
sirolimus 18
SIRTURO 10
sitagliptin 54
sitagliptin phos-metformin 54, 55
sitagliptin phosphate 55
sitagliptin-metformin 55
SKYCLARYS 34
SKYRIZI 47, 48, 58
sodium chloride 50
sodium chloride 0.45 % 75
sodium chloride 0.9 % 50
sodium chloride 3 % hypertonic 75
sodium chloride 5 % hypertonic 75
sodium oxybate 34
sodium phenylbutyrate 50
sodium polystyrene sulfonate 50
sodium,potassium,mag sulfates 58
sofosbuvir-velpatasvir 10
SOHONOS 50
solifenacin 73
SOLIQUA 100/33 55
SOLTAMOX 18
SOMAVERT 55
sorafenib 18
sotalol 42
SOTALOL AF 42
SPIRIVA RESPIMAT 72
spironolactone 42
spironolacton-hydrochlorothiaz 42
SPRINTEC (28) 66
SPRITAM 34
SPS (WITH SORBITOL) 50
SSD 48
STELARA 48
STEQEYMA 48
STIOLTO RESPIMAT 72
STIVARGA 18
streptomycin 10
STRIBILD 10
STRIVERDI RESPIMAT 72
SUBVENITE 34
sucralfate 59
sulfacetamide sodium 69
sulfacetamide sodium (acne) 48
sulfacetamide-prednisolone 69
sulfadiazine 10
sulfamethoxazole-trimethoprim 10
SULFAMYLON 48
sulfasalazine 59
sulindac 34
sumatriptan 34
sumatriptan succinate 34

sumatriptan-naproxen 35
sunitinib malate 18
SUNLENCA 10
SUNOSI 35
SYEDA 66
SYMPAZAN 35
SYMPROIC 59
SYMTUZA 10
SYNAREL 55
SYNJARDY 55
SYNJARDY XR 55
SYNTHROID 55
TABLOID 18
TABRECTA 18
tacrolimus 18, 48
tadalafil 73
tadalafil (pulm. hypertension) 72
TAFINLAR 18
tafluprost (pf) 69
TAGRISSO 18
TALZENNA 19
tamoxifen 19
tamsulosin 74
TASCENSO ODT 35
tasimelteon 35
TAVALISSE 42
tazarotene 48
telmisartan 42
telmisartan-amlodipine 42
telmisartan-hydrochlorothiazid 43
temazepam 35
TENCON 35
TENIVAC (PF) 61
tenofovir disoproxil fumarate 10
TEPMETKO 19
terazosin 43
terbinafine hcl 10
terbutaline 72
terconazole 66
teriflunomide 35
teriparatide 63
testosterone 55
testosterone cypionate 55
testosterone enanthate 55
tetrabenazine 35
tetracycline 11
TEZSPIRE 72
THALOMID 19
THEO-24 72
theophylline 72
thioridazine 35
thiothixene 35
TIADYLT ER 43
tiagabine 35
TIBSOVO 19
ticagrelor 43
TICOVAC 61
tigecycline 11
timolol maleate 43, 69
tinidazole 11
TIROSINT 55
TIROSINT-SOL 55
TIVICAY 11
TIVICAY PD 11
tizanidine 35
TOBI PODHALER 11
TOBRADEX 69
TOBRADEX ST 69
tobramycin 11, 69
tobramycin in 0.225 % nacl 11
tobramycin sulfate 11
tobramycin-dexamethasone 69
tobramycin-lotepred 69
TOBREX 69
tofacitinib 63
tolcapone 35
tolterodine 74
tolvaptan 55
tolvaptan (polycys kidney dis) 55
topiramate 35
toremifene 19
torsemide 43
TOUJEO MAX U-300 SOLOSTAR 55
TOUJEO SOLOSTAR U-300 INSULIN 55
TOVET EMOLLIENT 48
TRADJENTA 55
tramadol 35
tramadol-acetaminophen 35
trandolapril 43
tranexamic acid 67
tranylcypromine 35
TRAVASOL 10 % 75
travoprost 69
trazodone 35
TRELEGY ELLIPTA 72
TRELSTAR 19
TRESIBA FLEXTOUCH U-100 55
TRESIBA FLEXTOUCH U-200 55
TRESIBA U-100 INSULIN 55
tretinoin 48
tretinoin (antineoplastic) 19
TREXALL 19
triamcinolone acetonide 48, 51
triamterene 43
triamterene-hydrochlorothiazid 43
triazolam 35
TRIDERM 48
trientine 50
TRI-ESTARYLLA 67
trifluoperazine 35
trifluridine 69
trihexyphenidyl 35
TRIJARDY XR 56
TRIKAFTA 72
trimethoprim 11
TRI-MILI 67
trimipramine 35
TRINTELLIX 36
TRI-SPRINTEC (28) 67
TRIUMEQ 11
TRIUMEQ PD 11
TRI-VYLIBRA 67
TROPHAMINE 10 % 75
trospium 74
TRULICITY 56
TRUMENBA 61
TRUQAP 19
TUKYSA 19
TURALIO 19
TURQOZ (28) 67
TWINRIX (PF) 61
TYMLOS 63
TYPHIM VI 61
UBRELVY 36
UNITHROID 56
UPTRAVI 43
ursodiol 59
ustekinumab 48
valacyclovir 11
VALCHLOR 48
valganciclovir 11
valproic acid 36
valproic acid (as sodium salt) 36
valsartan 43
valsartan-hydrochlorothiazide 43
VALTOCO 36
VALTYA 67
vancomycin 11
VANDAZOLE 67
VANFLYTA 19
VAQTA (PF) 61
varenicline tartrate 50

VARIVAX (PF) 61
VAXCHORA VACCINE 61
VELIVET TRIPHASIC REGIMEN (28) 67
VEMLIDY 11
VENCLEXTA 19
VENCLEXTA STARTING PACK 19
venlafaxine 36
VENTOLIN HFA 72
verapamil 43
VEREGEN 48
VERQUVO 43
VERSACLOZ 36
VERZENIO 19
VESTURA (28) 67
VIBERZI 59
VIENVA 67
vigabatrin 36
VIGADRONE 36
VIJOICE 19
vilazodone 36
VIMKUNYA (PF) 61
VIORELE (28) 67
VIRACEPT 11
VIREAD 11
VITRAKVI 19
VIVITROL 36
VIVJOA 11
VIVOTIF 61
VONJO 19
VORANIGO 19
voriconazole 11
VOSEVI 11
VOWST 59
VRAYLAR 36
VUMERITY 36
VYFEMLA (28) 67
VYLIBRA 67
warfarin 43
WELIREG 20
WINREVAIR 72
WIXELA INHUB 72
WYOST 20
XALKORI 20
XARELTO 43
XARELTO DVT-PE TREAT 30D START 43
XATMEP 20
XCOPRI 36
XCOPRI MAINTENANCE PACK 36
XCOPRI TITRATION PACK 36
XDEMVY 69
XELRIA FE 67
XERMELO 20
XGEVA 20
XIFAXAN 11
XIGDUO XR 56
XIIDRA 69
XOCOVA 11
XOFLUZA 11
XOLAIR 72
XOLREMDI 61
XOSPATA 20
XPOVIO 20
XTANDI 20
XTRENBO 20
XULTOPHY 100/3.6 56
YARGESA 56
YESINTEK 48, 49
YF-VAX (PF) 61
YONSA 20
YORVIPATH 56
YUFLYMA(CF) 63
YUFLYMA(CF) AI CROHN'S-UC-HS 63
YUFLYMA(CF) AUTOINJECTOR 63
YULITHIRA 20
YUPELRI 72
YUVAFEM 67
ZAFEMY 67
zafirlukast 72, 73
zaleplon 36
ZARXIO 61
ZAVZPRET 36
ZEJULA 20
ZELAPAR 36
ZELBORAF 20
ZELSUVMI 49
ZELVYSIA 56
ZEMAIRA 50
ZENBEXUS 20
ZENPEP 59
ZEPOSIA 36
ZEPOSIA STARTER KIT (28-DAY) 36
ZEPOSIA STARTER PACK (7-DAY) 36
zidovudine 11
ZILBRYSQ 36, 37
zileuton 73
ziprasidone hcl 37
ziprasidone mesylate 37
ZIRGAN 69
ZOKINVY 50
ZOLINZA 20
zolmitriptan 37
zolpidem 37
ZONISADE 37
zonisamide 37
ZORYVE 49
ZOSYN IN DEXTROSE (ISO-OSM) 11
ZOVIA 1-35 (28) 67
ZTALMY 37
ZUBSOLV 37
ZURZUVAE 37
ZYKADIA 20

Section:
Prior Authorization

Acitretin

Products Affected

- *acitretin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Actimmune

Products Affected

- **ACTIMMUNE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ADHD Drugs

Products Affected

- *clonidine hcl oral tablet extended release 12 hr*
- *guanfacine oral tablet extended release 24 hr*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ADHD -AND- trial/failure, intolerance or contraindication to a stimulant
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Afinitor

Products Affected

- *everolimus (antineoplastic) oral tablet*
- *everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg*
- **YULITHIRA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For everolimus only, documentation of advanced, hormone receptor-positive, HER2-negative breast cancer -AND- all of the following (1-3): 1) member is a postmenopausal woman 2) using in combination with exemestane 3) therapeutic failure or intolerance to prior treatment with letrozole or anastrozole. For everolimus only, documentation of non-functional neuroendocrine tumors -AND- all of the following (1-2): 1) disease is classified as progressive, well-differentiated, non-functional 2) disease is of gastrointestinal or lung origin that are unresectable, locally advanced or metastatic. For everolimus only, documentation of advanced renal cell carcinoma -AND- therapeutic failure or intolerance to prior treatment with sunitinib or sorafenib. For everolimus only, documentation of renal angiomyolipoma and tuberous sclerosis complex (TSC), member does not require immediate surgery. For everolimus and everolimus tablets for oral suspension, documentation of TSC with Subependymal Giant Cell Astrocytoma -AND- member is not a candidate for curative surgical resection. For everolimus only, documentation of progressive neuroendocrine tumors of pancreatic origin -AND- disease is unresectable, locally advanced or metastatic. For everolimus tablets for oral suspension only, documentation of use for adjunctive treatment of TSC-associated partial-onset seizures.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Aimovig

Products Affected

- **AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML, 70 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-2): 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Akeega

Products Affected

- **AKEEGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. -AND- Concomitant therapy with prednisone. -AND- Concomitant therapy with a gonadotropin-releasing hormone analog or a bilateral orchiectomy, -AND- For metastatic castration-sensitive prostate cancer, disease harbors a deleterious or suspected deleterious BRCA2 mutation based on an FDA-approved test. -AND- For metastatic castration-resistant prostate cancer, BRCA mutations.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alecensa

Products Affected

- **ALECENSA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test -AND- One of the following (1-2): 1) metastatic disease, 2) will be used as adjuvant treatment following tumor resection of node positive or greater than or equal to 4 cm tumor(s).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alpha1-Proteinase Inhibitors

Products Affected

- **ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG**
- **GLASSIA**
- **PROLASTIN-C INTRAVENOUS SOLUTION**
- **ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of panacinar emphysema AND documentation of a decline in forced expiratory volume in 1 second (fev1) despite medical therapy (bronchodilators, corticosteroids) AND documentation of phenotype (pi*zz, pi*znull or pi*nullnull) associated with causing serum alpha 1-antitrypsin of less than 80 mg/dl AND documentation of an alpha 1-antitrypsin serum level below the value of 35% of normal (less than 80 mg/dl).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Covered under Part B when furnished incident to a physician service and is not self-administered.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Alunbrig

Products Affected

- **ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG**
- **ALUNBRIG ORAL TABLETS,DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ampyra

Products Affected

- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure disorder, Cr Cl less than 50ml/min
Required Medical Information	Documentation of diagnosis -AND- documentation that the patient is ambulatory and has walking impairment as evidenced by one of the following. 1. Functional status score (EDSS score). 2. Timed 25-foot Walk Test (T25W).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, documentation supporting improvement in walking impairment from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Anticholinergic Meds

Products Affected

- *paroxetine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- Attestation of counseling on risk-benefit profile and potential anticholinergic adverse effects -AND- Therapeutic failure, contraindication, or intolerance to 2 other antidepressants (i.e., SSRI, SNRI).
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Anzupgo

Products Affected

- **ANZUPGO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic hand eczema (CHE) -AND- symptoms persisting for more than 3 months or recurring at least twice within a 12-month timeframe -AND- therapeutic failure, contraindication, or intolerance to 1 generic, on-formulary, medium/high/super-high potency topical corticosteroid.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Arikayce

Products Affected

- **ARIKAYCE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Mycobacterium avium complex lung disease -AND- Attestation of not achieving negative sputum cultures despite at least 6 months with a multidrug background regimen containing 2 of the following: 1) macrolide 2) rifamycin or 3) ethambutal -AND- Arikayce will be used in conjunction with a background multidrug regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive sputum cultures -OR- negative sputum cultures for insufficient period of time (e.g. less than 12 months).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ATTR-CM drugs

Products Affected

- **ATTRUBY**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitantly with transthyretin-lowering agents
Required Medical Information	Documentation of cardiomyopathy of transthyretin-mediated amyloidosis (ATTR-CM) with amyloid deposits on cardiac biopsy or scintigraphy with heart contralateral greater than 1.5 (Grade III) or visual grade 2 to 3 - AND- Cardiac involvement supported by cardiac magnetic resonance, echocardiography or serum cardiac biomarker (e.g. B-type natriuretic peptide, cardiac troponin) -AND- ATTR-CM type is wild-type or variant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For reauthorization, attestation of improvement or delayed disease progression from baseline demonstrated by 6-minute walk test, cardiac function (e.g. LVEF, NYHA class), Kansas City Cardiomyopathy Questionnaire-Overall Summary, number of cardiovascular-related hospitalizations or serum cardiac biomarkers (e.g. B-type natriuretic peptide, cardiac troponin)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Atypical Antipsychotics

Products Affected

- *aripiprazole oral solution*
- *aripiprazole oral tablet,disintegrating*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Aubagio

Products Affected

- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as fingolimod, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Augtyro

Products Affected

- **AUGTYRO ORAL CAPSULE 160 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive. Documentation of solid tumors -AND- disease harbors a NTRK gene fusion -AND- one of the following (1-2): 1) disease is locally advanced or metastatic, or 2) surgical resection is likely to result in severe morbidity -AND- one of the following (3-4): 3) disease has progressed following treatment, or 4) the member has no satisfactory alternative therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Austedo

Products Affected

- **AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG**
- **AUSTEDO XR**
- **AUSTEDO XR TITRATION KT(WK1-4)**
- **ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3) 1) Chorea associated with Huntington's disease 2) In patients with comorbid depression, attestation of adequate treatment for depression is required. 3) Attestation that patient is not actively suicidal. -OR- 4) Tardive Dyskinesia
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Auvelity

Products Affected

- **AUVELITY ORAL TABLET, IR AND ER, BIPHASIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI). Documentation of Alzheimer's disease (AD) -AND- Experiencing agitation associated with dementia due to AD.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Avmapki-Fakzynja

Products Affected

- **AVMAPKI-FAKZYNJA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent low-grade serous ovarian cancer (LGSOC), disease is KRAS-mutated -AND- member has received at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ayvakit

Products Affected

- **AYVAKIT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable or metastatic gastrointestinal stromal tumor -AND- tumors harbor a PDGFRA exon 18 mutation. Documentation of agressive systemic mastocytosis, systemic mastocytosis with an associated hematological neoplasm, mast cell leukemia, or indolent systemic mastocytosis -AND- platelet count greater than or equal to 50 x 10*9/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Bafiertam

Products Affected

- **BAFIERTAM**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Balversa

Products Affected

- **BALVERSA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) FGFR3 mutation status as detected by an FDA approved test 2) disease progression on or after at least on prior line of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Banzel

Products Affected

- *rufinamide*

PA Criteria	Criteria Details
Exclusion Criteria	Patients with familial short QT syndrome
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Benlysta

Products Affected

- **BENLYSTA SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For diagnosis of active systemic lupus erythematosus (SLE), both of the following (1-2): 1) Documentation of positive anti-nuclear antibody (ANA) titer (greater than or equal to 1:80) or anti-double-stranded DNA antibody (anti-dsDNA) greater than or equal to 30IU/mL, 2) member will continue to receive concomitant standard of care treatment with use of at least one of the following (alone or in combination) (a-c): a) corticosteroids (e.g. prednisone), b) antimalarials (e.g. hydroxychloroquine), c) immunosuppressants (e.g. azathioprine, mycophenolate mofetil, or methotrexate). For diagnosis of active lupus nephritis (LN), one of the following (3 or 4): 3) Diagnosis confirmed by renal biopsy, 4) Contraindication to renal biopsy, laboratory findings specific to LN (e.g., elevated serum creatinine, abnormal urine analysis, decreased eGFR). -AND- Member will continue to receive concomitant standard of care treatment which includes corticosteroids (e.g. prednisone) with at least one of the following (d or e): d) mycophenolate for induction followed by mycophenolate for maintenance, e) cyclophosphamide for induction followed by azathioprine for maintenance.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For SLE reauthorization, attestation of disease stability or improvement -AND- attestation the member will continue to receive standard of care therapy with corticosteroids, antimalarials, or immunosuppressives. For active LN reauthorization, attestation of disease stability or improvement -AND- attestation the member will continue to receive standard of care therapy with mycophenolate or azathioprine. For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Besremi

Products Affected

- **BESREMI**
- **BESREMI PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Bosulif

Products Affected

- **BOSULIF ORAL CAPSULE 100 MG, 50 MG**
- **BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy. For pediatric patients 1 year of age and older, one of the following (3, 4): 3) newly diagnosed PH-positive CML in the chronic phase 4) diagnosis of Ph-positive CML in the chronic phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For capsules and 18 years of age or older, intolerance to generic bosutinib tablets or inability to swallow tablets is required. For tablets and 18 years of age or older, intolerance to generic bosutinib tablets is required. For Bosutinib 100mg capsules and pediatric 1 year of age or older, inability to swallow tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Bosutinib

Products Affected

- *bosutinib oral tablet 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For members 18 years of age and older, one of the following (1, 2): 1) newly diagnosed Philadelphia chromosome (PH) -positive CML in the chronic phase 2) diagnosis of Ph-positive CML in the chronic, accelerated, or blast phase and no longer responding to or intolerant to at least 1 prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Braftovi

Products Affected

- **BRAFTOVI**

PA Criteria	Criteria Details
Exclusion Criteria	Use in wild-type BRAF melanoma, wild-type BRAF CRC, or wild-type BRAF NSCLC.
Required Medical Information	Documentation of diagnosis. For metastatic colorectal cancer (mCRC) and using in combination with cetuximab, all of the following (1-3): 1) BRAF V600E mutation status as detected by an FDA-authorized test 2) using in combination with cetuximab 3) member has received prior therapy for CRC. For mCRC and using in combination with cetuximab and fluorouracil-based chemotherapy, all of the following (4-5): 4) BRAF V600E mutation status, as detected by an FDA-authorized test 5) using in combination with cetuximab and fluorouracil-based chemotherapy. For unresectable or metastatic melanoma, all of the following (6-7): 6) BRAF V600E or V600K mutation status, as detected by an FDA-authorized test 7) using in combination with binimetinib. For metastatic non-small cell lung cancer, all of the following (8-9): 8) BRAF V600E mutation status, as detected by an FDA-authorized test 9) using in combination with binimetinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Brinsupri

Products Affected

- **BRINSUPRI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of non-cystic fibrosis bronchiectasis (NCFB) confirmed by computed tomography (CT) -AND- at least 1 of the following symptoms consistent with bronchiectasis (1-8): 1) cough on most days of the week, 2) chronic sputum production, 3) history of recurrent respiratory infections, 4) dyspnea, 5) wheezing, 6) rhinosinusitis, 7) hemoptysis, 8) recurrent pleurisy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Brukinsa

Products Affected

- **BRUKINSA ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma (MCL), previous treatment with at least 1 prior therapy. For marginal zone lymphoma (MZL), previous treatment with at least 1 anti-CD20-based regimen. For follicular lymphoma (FL), using in combination with obinutuzumab and previous treatment with at least 2 prior lines of systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Buphenyl

Products Affected

- *sodium phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of urea cycle disorders involving deficiencies of carbamylphosphate synthetase, argininosuccinic acid synthetase, or ornithine transcarbamylase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cabometyx

Products Affected

- **CABOMETYX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-6): 1) diagnosis of advanced renal cell carcinoma (RCC), 2) diagnosis of advanced RCC and using as a first-line treatment in combination with nivolumab, 3) member has previously been treated with sorafenib for hepatocellular carcinoma, 4) member has experienced disease progression following prior VEGFR-targeted therapy and is either radioactive iodine-refractory or is ineligible for radioactive iodine therapy for locally advanced or metastatic differentiated thyroid cancer, 5) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated pancreatic neuroendocrine tumors (pNET), 6) diagnosis of previously treated, unresectable, locally advanced or metastatic, well-differentiated extra-pancreatic neuroendocrine tumors (epNET)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Calquence

Products Affected

- **CALQUENCE (ACALABRUTINIB MAL)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For mantle cell lymphoma, member has received at least one prior therapy -OR- all of the following (1-3): 1) member has not received prior therapy for MCL, 2) member is ineligible for autologous hematopoietic stem cell transplantation (HSCT), and 3) using in combination with bendamustine and rituximab.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Caplyta

Products Affected

- **CAPLYTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For schizophrenia, bipolar I disorder, and bipolar II disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Caprelsa

Products Affected

- **CAPRELSA ORAL TABLET 100 MG, 300 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic or progressive medullary thyroid cancer in patients with unresectable locally advanced or metastatic disease.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Carbaglu

Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use as an adjunct therapy for acute hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA) -OR- maintenance therapy for chronic hyperammonemia due to hepatic enzyme N-acetylglutamate synthase (NAGS) deficiency
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cayston

Products Affected

- **CAYSTON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cerdelga

Products Affected

- **CERDELGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of type 1 Gaucher disease confirmed by the following A. or B. A) With one of the following symptoms (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm*3. 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B) Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles -AND- Documentation of CYP2D6 metabolizer status (e.g. intermediate metabolizer).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CF drugs

Products Affected

- **TOBI PODHALER**
- *tobramycin in 0.225 % nacl*
- *tobramycin inhalation*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization of tobramycin products, attestation of decrease in sputum density of pseudomonas aeruginosa, increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cialis

Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of benign prostatic hyperplasia (BPH) and trial/failure of at least two alternative medications in the following classes (alpha-1 adrenergic blockers and/or 5-alpha reductase inhibitors)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cinryze

Products Affected

- **CINRYZE**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) due to deficiency of C1INH (type I & II) with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) with normal C1INH (type III) with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE, FXII mutation, angiopoietin-1 mutation, plasminogen mutation, kininogen-1 mutation, myoferlin mutation, or heparin sulfate-glucosamine 3-O-sulfotransferas 6 mutation. 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cobenfy

Products Affected

- **COBENFY**
- **COBENFY STARTER PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cometriq

Products Affected

- **COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressive, metastatic medullary thyroid cancer
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Copiktra

Products Affected

- **COPIKTRA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is no longer responding or is intolerant to at least 2 prior therapies for chronic lymphocytic leukemia and small lymphocytic leukemia.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Corlanor

Products Affected

- *ivabradine oral tablet 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- All of the following: 1) Normal sinus rhythm, 2) Resting heart rate greater than or equal to 70 beats per minute, 3) For adults, left ventricular ejection fraction less than or equal to 35 percent, 4) For adults, concurrent use of 1 beta-blocker used for treatment of heart failure (i.e. bisoprolol, carvedilol, metoprolol succinate) or therapeutic failure, intolerance, or contraindication to the maximum tolerated dose of 2 beta-blockers used for treatment of heart failure.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Cosentyx

Products Affected

- **COSENTYX (2 SYRINGES)**
- **COSENTYX PEN (2 PENS)**
- **COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML**
- **COSENTYX UNOREADY PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) or all are contraindicated. For non-radiographic axial spondyloarthritis, inadequate response or intolerance to 2 NSAIDs or all are contraindicated.
Age Restrictions	Deny if less than 6 years of age for moderate to severe plaque psoriasis -OR- less than 2 years of age for psoriatic arthritis -OR- less than 4 years of age for enthesitis-related arthritis -OR- less than 12 years of age for hidradenitis suppurativa and ankylosing spondylitis -OR- less than 18 years of age for non-radiographic axial spondyloarthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication. For hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Cotellic

Products Affected

- **COTELLIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following for unresectable or metastatic melanoma (1-2): 1) BRAF V600E or V600K mutation status 2) Concomitant therapy with vemurafenib. For cobimetinib monotherapy, documentation of histiocytic neoplasms.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Cysteamine Ophthalmic Drops

Products Affected

- **CYSTADROPS**
- **CYSTARAN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystinosis -AND- Attestation of accumulation of corneal cystine crystals
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Danziten

Products Affected

- **DANZITEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib -OR- member is newly diagnosed in the chronic phase.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Daraprim

Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For primary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 100 cells/mm*3 -AND- Toxoplasma IgG positive -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For secondary prophylaxis of toxoplasmosis gondii infection, CD4 count less than 200 cells/mm*3. For secondary prophylaxis of cystoisosporiasis with CD4 count less than 200 cells/mm*3 or acute cystoisosporiasis infection: failure, intolerance or contraindication to trimethoprim-sulfamethoxazole. For primary prophylaxis of Pneumocystis jirovecii pneumonia: diagnosis of HIV -AND- CD4 count less than 200 cells/mm*3 -AND- failure, intolerance or contraindication to trimethoprim-sulfamethoxazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Daurismo

Products Affected

- **DAURISMO ORAL TABLET 100 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of newly diagnosed Acute Myeloid Leukemia -AND- Used in combination with cytarabine -AND- At least one comorbidity that preclude use of intensive induction chemotherapy defined as one of the following: 1) Age greater than or equal to 75 2) Severe cardiac or pulmonary comorbidity 3) Reduced renal function 4) Hepatic impairment 5.) Physician attests patient is not a candidate for intensive induction therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Daytrana

Products Affected

- *methylphenidate transdermal*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial/failure or intolerance to 2 of the following generic medications: methylphenidate, atomoxetine, or dextroamphetamine/amphetamine.
Age Restrictions	Deny if less than 6 years of age or greater than 17 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Deferasirox

Products Affected

- *deferasirox oral tablet, dispersible*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic iron overload due to blood transfusions, transfusion history of greater than or equal to 100 mL/kg of packed red blood cells (i.e. at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg) -And- history of serum ferritin consistently greater than 1,000 mcg/L or liver iron concentration (LIC) greater than or equal to 7 iron per gram of liver dry weight (mg Fe/g dw). For Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia (NTDT) Syndrome, LIC of at least 5 mg Fe/g dw -AND- serum ferritin greater than 300 mcg/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of chronic iron overload due to blood transfusion, continued requirement for regular blood transfusions -AND- serum ferritin level greater than or equal to 500mcg/L or LIC greater than or equal to 3 mg Fe/g dw. For reauthorization of chronic iron overload in NTDT syndrome, LIC greater than or equal to 3 mg Fe/g dw.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Diacomit

Products Affected

- **DIACOMIT ORAL CAPSULE 250 MG, 500 MG**
- **DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravets syndrome - AND- Used in combination with clobazam
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting reduction in seizure frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Dihydroergotamine

Products Affected

- *dihydroergotamine nasal*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- requires non-oral route of administration -AND- therapeutic failure or intolerance to generic sumatriptan nasal spray.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Doxepin Cream

Products Affected

- *doxepin topical*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- course of therapy will not exceed 8 days -AND- experienced therapeutic failure, intolerance, or contraindication to one of the following (1 or 2): 1) at least 2 generic formulary topical corticosteroids. 2) generic topical tacrolimus or pimecrolimus for topic dermatitis with the facial or anogenital involvement.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Drizalma

Products Affected

- **DRIZALMA SPRINKLE ORAL**
  CAPSULE, DELAYED REL SPRINKLE
  20 MG, 30 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow tablets/capsules. For fibromyalgia, members must also have widespread bilateral pain above and below the waist for greater than 3 months duration -AND- At least 1 fibromyalgia-related symptom (e.g., cognitive impairment, fatigue, sleep disturbance, neurologic symptoms, exercise intolerance).
Age Restrictions	Deny if less than 18 years of age in the treatment of fibromyalgia, major depressive disorder, diabetic peripheral neuropathy and chronic musculoskeletal pain -OR- if less than 7 years of age in generalized anxiety disorder
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Dupixent

Products Affected

- **DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML**
- **DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Mod/Sev atopic dermatitis (AD): failure/intolerance to 1 topical corticosteroid (TCS) or, if 2+ yrs, topical calcineurin inhibitor (e.g. tacrolimus, etc) OR, if sev AD, topical tx are precluded d/t body surface area involvement or severely damaged skin. TCS in combo w/Dupixent for mod/sev AD is permitted & allows coverage. Mod/sev asthma: hx of 2+ exacerbations req oral/inj corticosteroids in past 12mos or 1+ exacerbation req hospitalization in past 12mos AND blood eosinophils of at least 150cells/uL or current daily/alternate-day oral corticosteroid (OCS) AND inadequate control despite regular med/high-dose inhaled corticosteroids (ICS) & 1 or more add'l asthma controller (e.g. LABA, leukotriene receptor antagonist [LTRA], theophylline) +/- OCS, unless intolerant or contraindicated (c/i) to all AND will continue med/high dose ICS & 1 or more add'l asthma controller +/- OCS. Chronic rhinosinusitis w/ nasal polyposis (CRSwNP): failure, intolerance, or c/i to intranasal corticosteroid. Eosinophilic esophagitis: esophageal eosinophils of 15eos/hpf or higher on esophageal biopsy AND clinical esophageal dysfxn symptoms. Prurigo nodularis. COPD: blood eosinophils of at least 300cells/uL or current daily/alternate-day OCS AND inadequate control despite 3+ mos of regular LAMA, LABA, & ICS tx, unless intolerant or c/i to all. Chronic Spontaneous Urticaria: failure, intolerance, or c/i to 1 2nd-gen H1 antihistamine at max rec dose (e.g. cetirizine, etc). Bullous pemphigoid: failure, intolerance, or c/i on high/super high potency TCS or OCS. Allergic fungal rhinosinusitis: hx of sinonasal surgery AND diagnosis confirmed by at least 2 of the following: Type 1 hypersensitivity to fungus (history, in vitro IgE, or skin testing), positive fungal stain from surgical sinus contents, presence of eosinophilic mucin w/o sinus tissue invasion, presence of CRSwNP, or characteristic CT signs
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For atopic dermatitis reauthorization, attestation of positive clinical response to therapy. For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For EoE reauthorization, attestation of histological remission (less than 15 eos/hpf) on esophageal biopsy or reduced severity or frequency of clinical symptoms of esophageal dysfunction. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline. For COPD reauthorization, attestation of one of the following is required (1-4): 1) reduction in COPD symptoms, 2) improvement in exercise tolerance, 3) delayed disease progression, or 4) reduction in the number of COPD exacerbations. For CSU reauthorization, improved CSU symptoms. For BP reauthorization, 1 of the following is required (1-4): 1) attestation of disease control, 2) reduction in number of relapses, 3) improvement in BP symptoms, or 4) reduction in oral corticosteroid use. For allergic fungal rhinosinusitis reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Duvyzat

Products Affected

- **DUVYZAT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Duchenne muscular dystrophy with pathogenic mutation in the dystrophin gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

EGFR Tyrosine Kinase Inhibitors

Products Affected

- *erlotinib*
- **GILOTRIF**

PA Criteria	Criteria Details
Exclusion Criteria	Afatinib products: tumors with resistant EGFR mutations. Erlotinib products: use in NSCLC tumors with mutations other than those in FDA-approved indications. Use in combination with platinum based chemotherapy.
Required Medical Information	For afatinib, documentation of metastatic non-small cell lung cancer (NSCLC) -AND- one of the following, as detected by an FDA-approved test (1-3): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutation 3) disease harbors non-resistant EGFR mutation (i.e., S768I, L861Q, G719X) -OR- documentation of squamous metastatic NSCLC and member has experienced progression on platinum-based chemotherapy. For erlotinib, documentation of metastatic NSCLC -AND- one of the following, as detected by an FDA-approved test (1-2): 1) disease harbors EGFR exon 19 deletions 2) disease harbors EGFR exon 21 (L858R) substitution mutations -OR- documentation of locally advanced, unresectable or metastatic pancreatic cancer -AND- all of the following (1-2): 1) using erlotinib as first-line therapy 2) using in combination with gemcitabine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Emgality

Products Affected

- **EMGALITY PEN**
- **EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, the following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). For episodic cluster headaches, characterized by severe or very severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes when left untreated -AND- Attack frequency of at least one attack every other day during the cluster period.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency is required. For reauthorization of cluster headache, attestation of reduction in the number of mean weekly cluster headaches from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Ensacove

Products Affected

- **ENSACOVE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally advanced or metastatic non-small cell lung cancer (NSCLC) -AND- Disease is ALK-positive, as detected by an FDA-approved test -AND- Not previously received an ALK inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Entresto Sprinkle

Products Affected

- **ENTRESTO SPRINKLE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of symptomatic heart failure with left ventricular systolic dysfunction -AND- weight less than 40kg, or weight greater than or equal to 40kg and less than 50kg and one of the following (1 or 2): 1) therapeutic failure or intolerance to Entresto (sacubitril/valsartan) tablets, or 2) inability to swallow tablets.
Age Restrictions	Deny if greater than 17 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, all of the following are required (1-3): 1) attestation of positive clinical response to therapy, 2) member is younger than 18 years of age, 3) weight less than 40kg, or weight greater than or equal to 40kg and less than 50kg and one of the following (4 or 5): 5) intolerance to Entresto (sacubitril/valsartan) tablets, or 5) inability to swallow tablets.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Eohilia

Products Affected

- **EOHILIA**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 12 weeks
Required Medical Information	Diagnosis of eosinophilic esophagitis -AND- esophageal eosinophil count greater than or equal to 15 eos/hpf on esophageal biopsy -AND- clinical symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, gastroesophageal reflux)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Epclusa

Products Affected

- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance. For Brand Epclusa, the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Epidiolex

Products Affected

- **EPIDIOLEX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Lennox-Gastaut, Dravet syndromes or Tuberous Sclerosis Complex. For Lennox-Gastaut, trial and failure or intolerance of at least two standard of care treatments (e.g. lamotrigine, clobazam). For Lennox-Gastaut and Dravet syndromes, treatment is in combination with other conventional agents.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Eprontia

Products Affected

- *topiramate oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- inability to swallow pills or food. For migraine, therapeutic failure, intolerance, or contraindication to two generic preventive migraine therapies. For partial onset seizures, primary generalized tonic-clonic seizures, or adjunctive treatment of Lennox-Gastaut Syndrome, therapeutic failure, contraindication, or intolerance to two generic anti-epileptic drugs.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ergotamine

Products Affected

- *ergotamine-caffeine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use to abort a vascular headache -AND- therapeutic failure, intolerance, or contraindication to a generic triptan -OR- documentation of use to prevent a vascular headache -AND- therapeutic failure, intolerance, or contraindication to generic prophylactic migraine medication (e.g., topiramate, propanolol, timolol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Erivedge

Products Affected

- **ERIVEDGE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- if disease is locally advanced all of the following: 1) disease has recurred following surgery, or is not a candidate for surgery, 2) is not a candidate for radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Erleada

Products Affected

- **ERLEADA ORAL TABLET 240 MG, 60 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- the member meets one of the following (1 or 2) 1. Documentation of use in combination with a GnRH analog -OR- 2. The member has had a bilateral orchiectomy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Escitalopram Capsule

Products Affected

- *escitalopram oxalate oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of major depressive disorder or generalized anxiety -AND- Therapy has been initiated with escitalopram 10mg or member has been receiving and is experiencing unfavorable tolerability to escitalopram 20mg -AND- Intolerance to generic escitalopram tablets -AND- Therapeutic failure, intolerance or contraindication to at least one other antidepressant (e.g., SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Etanercept

Products Affected

- **ENBREL MINI**
- **ENBREL SUBCUTANEOUS SOLUTION**
- **ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)**
- **ENBREL SURECLICK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis and Ankylosing Spondylitis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Psoriatic Arthritis or Less than 4 years of age for Plaque Psoriasis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	Therapeutic failure or intolerance to 1 of the following preferred adalimumab products is required, when being utilized for the same medically accepted indication and age: Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Exxua

Products Affected

- **EXXUA ORAL TABLET EXTENDED RELEASE 24 HR**
- **EXXUA ORAL TABLET, EXT REL 24HR DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder (MDD) -AND- Therapeutic failure or intolerance to generic bupropion hydrochloride tablets -AND- Therapeutic failure, intolerance or contraindication to one other generic antidepressant (e.g. SNRI, SSRI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fabhalta

Products Affected

- **FABHALTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For paroxysmal nocturnal hemoglobinuria (PNH), meets one of the following (1 or 2): 1) PNH mutant clones confirmed by flow cytometry, or 2) glycosylphosphatidylinositol-anchored proteins (GPI-AP)-deficient polymorphonuclear cells (PMNs) confirmed by flow cytometry -AND- meets one of the following (3-6): 3) anemia secondary to PNH (e.g. hemoglobin less than 10.5 g/dL with symptoms of anemia), 4) elevated lactate dehydrogenase (LDH) greater than or equal to 1.5 times the upper limit of normal, 5) history of a thromboembolic event, or 6) clinical findings of systemic complications (e.g. fatigue, hemoglobinuria, abdominal pain, dyspnea, dysphagia, erectile dysfunction, history of blood cell transfusion due to PNH) -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)) unless initially cross-titrating. For diagnosis of primary immunoglobulin A nephrology (IgAN) confirmed by biopsy, member is at risk for rapid disease progression evidenced by one of the following (7 or 8): 7) Urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g or 8) Proteinuria greater than or equal to 1 g/day -AND- has experienced therapeutic failure, contraindication, or intolerance to a maximally tolerated dose of one of the following (a or b): a) Angiotensin converting enzyme (ACE) inhibitor, b) Angiotensin receptor blocker (ARB) -AND- experienced therapeutic failure, contraindication, or intolerance to one of the following (c or d): c) Filspari (sparsentan) or d) Tarpeyo (budesonide). For Complement 3 Glomerulopathy (C3G) confirmed by biopsy, meets all of the following (13-14): 13) UPCR greater than or equal to 1.0 g/g, 14) currently therapy of the member should be on the maximally tolerated dose of one of the following (e or f): e) ACE-inhibitor, f) ARB.
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of PNH, attestation of positive clinical response defined as one of the following (1-3): 1) hemoglobin stabilization or increase from baseline, 2) decrease in transfusions from baseline, or 3) decrease in LDH levels from baseline or reduction of hemolysis -AND- will not be used in combination with another complement inhibitor for PNH (e.g. Soliris (eculizumab), Ultomiris (ravulizumab), Empaveli (pegcetacoplan)). For reauthorization of IgAN, reduction in urine protein-to-creatinine ratio (UPCR) or proteinuria from baseline. For reauthorization of C3G, reduction in UPCR from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fanapt

Products Affected

- **FANAPT**
- **FANAPT TITRATION PACK A**
- **FANAPT TITRATION PACK B**
- **FANAPT TITRATION PACK C**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fasenra

Products Affected

- **FASENRA PEN**
- **FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe asthma and all of the following (1-4): 1) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12 mos or at least 1 asthma exacerbation requiring hospitalization in past 12 mos 2) blood eosinophils greater than or equal to 150cells/uL within the past 6 weeks or greater than or equal to 300cells/uL within the past 12 mos in without other potential causes of eosinophilia (e.g. hypereosinophilic syndromes, neoplastic disease, known suspected parasitic infection) 3) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline), with or without oral corticosteroids (OCS), unless intolerant or contraindicated to all 4) will continue treatment with medium or high dose ICS and at least 1 add'l asthma controller medication, with or without OCS -OR- Documentation of eosinophilic granulomatosis with polyangiitis (EGPA) and all of the following (5-6): 5) history of relapsing or refractory disease 6) will be receiving standard of care while on Fasenra therapy with glucocorticoid treatment (e.g. prednisone or prednisolone), with or without immunosuppressive therapy (e.g. cyclosporine, leflunomide, azathioprine). Documentation of Hypereosinophilic syndrome (HES) and all of the following (7-8): 7) no identifiable non-hematologic secondary cause, 8) Treatment with HES therapy (chronic or episodic OCS, immunosuppressive or cytotoxic therapy) for at least 4wks, unless therapeutic failure, intolerant, or contraindicated to all.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For severe asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbation, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For EPGA reauthorization, attestation of one of the following is required (5-8): 5) reduction in the frequency and/or severity of relapses, 6) reduction or discontinuation of doses of corticosteroids and/or immunosuppressant, 7) disease remission, or 8) reduction in severity or frequency of EGPA-related symptoms. For HES reauthorization, 1 of the following is required (9-11): 9) reduction in frequency of HES flares, 10) maintenance or reduction in background HES therapy requirements, 11) reduction in or stabilization of eosinophil-mediated organ damage (e.g., cardiac, pulmonary, gastrointestinal, or neurological manifestations).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fecal Microbiota Products

Products Affected

- **VOWST**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a recent diagnosis of Clostridioides difficile infection (CDI) confirmed by a positive stool test -AND- Classified as recurrent (e.g., at least 2 CDI episodes) -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	For reauthorization, attestation of recurrent CDI episodes after administration of the initial fecal microbiota product -AND- Will be used for prophylaxis and not treatment of recurrent CDI -AND- Attestation that antibiotic treatment for the most recent recurrent CDI is complete or will be completed.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Fetzima

Products Affected

- **FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)**
- **FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder and trial and failure of two other generic antidepressants.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Filsuvez

Products Affected

- **FILSUVEZ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB) -AND- at least one open wound.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of improvement in targeted wound(s) -AND- member requires additional courses of treatment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Fintepla

Products Affected

- **FINTEPLA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Dravet syndrome or Lennox-Gastaut syndrome. For Lennox-Gastaut syndrome, therapeutic failure, contraindication, or intolerance to at least 2 standard of care treatments (e.g. lamotrigine, clobazam).
Age Restrictions	Deny if less than 2 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Firazyr

Products Affected

- *icatibant*
- **SAJAZIR**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two acute therapies simultaneously and acute therapy should not be used as prophylactic therapy
Required Medical Information	For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) due to deficiency of C1INH (type I & II) with the following (1-3): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. For the treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) with normal C1INH (type III) with the following (4-7): 4) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 5) Documentation of family history of HAE, FXII mutation, angiopoietin-1 mutation, plasminogen mutation, kininogen-1 mutation, myoferlin mutation, or heparin sulfate-glucosamine 3-O-sulfotransferas 6 mutation. 6) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 7) Medications known to cause angioedema have been evaluated and discontinued.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Firmagon

Products Affected

- **FIRMAGON KIT W DILUENT SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Flector

Products Affected

- *diclofenac epolamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis AND one of the following (1,2 or 3): 1) trial/failure, intolerance, or contraindication to 2 oral generic NSAIDs one of which must be diclofenac 2) hypersensitivity to oral NSAIDs 3) history or high risk for adverse gastrointestinal effects associated with oral NSAID use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Forteo

Products Affected

- *teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of teriparatide will be limited to a coverage duration of 24 months in the absence of provider attestation that the member remains at or has returned to having a high risk for fracture
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Fotivda

Products Affected

- **FOTIVDA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received at least two prior systemic therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Fruzaqla

Products Affected

- **FRUZAQLA ORAL CAPSULE 1 MG, 5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member has received previous threatment with a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy -AND- member has received previous treatment with an anti-VEGF therapy -AND- one of the following, if member is RAS wild-type (1-2): 1) member has received previous therapy with an anti-EGFR therapy 2) prescriber attests that treatment with an anti-EGFR therapy would not be medically appropriate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gattex

Products Affected

- **GATTEX 30-VIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of short bowel syndrome (SBS) having less than 200 cm of functional small bowel -AND- Dependence on parenteral/intravenous nutrition -AND- weight of at least 10 kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline -AND- 1 of the following (1-2): 1) continued dependence on parenteral nutrition/intravenous nutritional support, 2) not receiving parenteral nutrition/intravenous support and experienced therapeutic failure upon trial of discontinuation of parenteral nutrition/intravenous support.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gavreto

Products Affected

- **GAVRETO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, disease is RET fusion-positive as detected by an FDA approved test. For advanced or metastatic thyroid cancer, all of the following (1-2): 1) disease is RET fusion-positive 2) if radioactive iodine is appropriate, the member is radioactive iodine-refractory.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gilenya

Products Affected

- *fingolimod*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Glatiramer

Products Affected

- **COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- **GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gleevec

Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult agressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvent treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Gleostine

Products Affected

- *lomustine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or metastatic brain tumor(s) -AND- member has previously received surgical and/or radiotherapeutic procedure(s). Documentation of Hodgkin's lymphoma -AND- all of the following (1-2): 1) using lomustine in combination with other chemotherapies 2) member has experienced disease progression with initial chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GLP1

Products Affected

- *liraglutide*
- **MOUNJARO**
- **OZEMPIC SUBCUTANEOUS**
- **TRULICITY**

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Gomekli

Products Affected

- **GOMEKLI ORAL CAPSULE 1 MG, 2 MG**
- **GOMEKLI ORAL TABLET FOR SUSPENSION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) -AND- presence of symptomatic plexiform neurofibromas (PN) not amenable to complete resection.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Grastek

Products Affected

- **GRASTEK**

PA Criteria	Criteria Details
Exclusion Criteria	Asthma (severe, unstable or uncontrolled), concomitant sublingual or subcutaneous immunotherapy, therapy initiation during active allergy season
Required Medical Information	Documentation of allergic rhinitis and use for Timothy grass pollen or cross reactive grass pollens (Sweet Vernal, Orchard, Perennial Rye, Timothy, Kentucky Blue Grass pollen, Redtop, or meadow fescue) -AND- allergic rhinitis with or without conjunctivitis has been confirmed by a pollen specific positive skin test or in vitro testing for pollen-specific IgE antibodies -AND- trial and failure or intolerance to an intranasal steroid and an oral non-sedating antihistamine, intranasal antihistamine or intranasal anticholinergic agent
Age Restrictions	Deny if less than 5 years of age or greater than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Member must also be prescribed an epinephrine auto injector. For reauthorization, attestation of improved allergy symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Growth Hormone

Products Affected

- **GENOTROPIN**
- **GENOTROPIN MINIQUICK**
- **HUMATROPE INJECTION CARTRIDGE**
- **NORDITROPIN FLEXPRO SUBCUTANEOUS PEN INJECTOR 10 MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML (10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML)**
- **OMNITROPE**
- **SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis, growth chart, member has open epiphyses, growth velocity, response to stimulation test, when applicable to meet standard diagnostic criteria. Additionally for growth failure due to chronic kidney disease, glomerular filtration rate is less than 89ml/min per 1.73m*2. For HIV wasting and cachexia, Concurrent use of antiretroviral therapy -AND- weight loss from baseline of at least 10% in the past 12 months or at least 5% in the past 6 months. For short bowel syndrome, receiving management for short bowel syndrome, including specialized nutritional support -AND- less than 200 cm of functional small bowel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of growth velocity and member has open epiphyses, as applicable to meet standard continuation of therapy guidelines. For reauthorization of HIV wasting and cachexia, attestation of increase in weight from start of therapy. For reauthorization of short bowel syndrome, continued dependence on parenteral nutrition/intravenous nutritional support -AND- attestation of increase in weight from baseline or decrease in intravenous parenteral nutrition requirements from baseline.
Indications	All Medically-accepted Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	No

Haegarda

Products Affected

- **HAEGARDA**

PA Criteria	Criteria Details
Exclusion Criteria	Member should not be on two prophylactic therapies simultaneously.
Required Medical Information	For the prophylactic treatment of abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) due to deficiency of C1INH (type I & II) with the following (1-4): 1) Low C4 level of less than or equal to 14mg/dL or C4 below lower limit of laboratory reference range and 1 of the following (A or B). A) C1 inhibitor (C1INH) antigen level less than or equal to 19mg/dL or below lower limit of laboratory reference range. B) Normal C1INH antigen level and a low C1INH functional level below laboratory reference range. 2) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 3) Medications known to cause angioedema have been evaluated and discontinued. 4) Documentation of member's weight. For the prophylactic treatment of acute abdominal, facial, or laryngeal attacks of hereditary angioedema (HAE) with normal C1INH (type III) with the following (5-9): 5) Documentation of clinical laboratory performance C4, C1INH antigen, or C1INH functional level are within normal limits of laboratory reference ranges. 6) Documentation of family history of HAE, FXII mutation, angiopoietin-1 mutation, plasminogen mutation, kininogen-1 mutation, myoferlin mutation, or heparin sulfate-glucosamine 3-O-sulfotransferas 6 mutation. 7) Past medical history of at least 1 symptom of moderate or severe angioedema attack (e.g. airway swelling, painful facial distortion) in absence of concomitant hives. 8) Medications known to cause angioedema have been evaluated and discontinued. 9) Documentation of member's weight.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Hepcludex

Products Affected

- **HEPCLUDEX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of chronic hepatitis delta virus (HDV) and all of the following (1-3): 1) diagnosis supported by detectable serum HDV RNA and positive serum anti-HDV antibodies, 2) attestation that HDV infection is classified as chronic, 3) liver status of no cirrhosis or compensated cirrhosis, 4) attestation that the chronic HBV infection will be managed as clinically appropriate per the FDA-approved labeling.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, all of the following are required (1-3): 1) liver status of no cirrhosis or compensated cirrhosis, 2) attestation that the chronic HBV infection continues to be managed as clinically appropriate per the FDA-approved labeling, 3) attestation that response to treatment warrants continued therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Hernexeos

Products Affected

- **HERNEXEOS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For non-squamous non-small cell lung cancer (NSCLC), all of the following (1-2): 1) disease is unresectable or metastatic, 2) disease harbors HER2 (ERBB2) tyrosine kinase domain activating mutations, as detected by an FDA-authorized test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

High-risk meds

Products Affected

- *amitriptyline*
- *amitriptyline-chlordiazepoxide*
- *benztropine oral*
- *clomipramine*
- *cyproheptadine oral syrup*
- *cyproheptadine oral tablet*
- *doxepin oral capsule*
- *doxepin oral concentrate*
- *doxepin oral tablet*
- *hydroxyzine hcl oral solution 10 mg/5 ml*
- *hydroxyzine hcl oral tablet*
- *imipramine hcl*
- *imipramine pamoate*
- *metaxalone oral tablet 400 mg, 800 mg*
- *perphenazine-amitriptyline*
- *promethazine oral*
- *trimipramine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For all medications subject to this PA group, the following information (1 through 3) is required: 1. Documentation of diagnosis 2. Explanation of risk-benefit profile favoring use of the high-risk medication 3. Attestation of an intent to monitor and address treatment-related adverse events. For target tricyclic antidepressants (TCAs), in addition to criteria 1 through 3 above, trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs for the same indication, if available, is required (e.g. SSRIs and SNRIs). If using a TCA for a medically-accepted indication not shared by the safer alternatives listed, then no trial of alternatives is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Doxepin doses less than or equal to 6 mg per day will receive automatic approval.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

High-risk meds phenobarbital

Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for use in sedation/insomnia
Required Medical Information	For use in seizures the following are required: 1. Explanation of risk-benefit profile favoring use of the high-risk medication 2. Attestation of an intent to monitor and address treatment-related adverse events. 3. For new starts, the trial and failure or documentation of intolerance or contraindication to at least 2 non-high risk alternative drugs used for seizures (e.g. carbamazepine, lamotrigine) is required.
Age Restrictions	Automatic approval if less than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Seizure disorders
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Hyrnuo

Products Affected

- **HYRNUO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For non-squamous non-small cell lung cancer (NSCLC), all of the following (1-2): 1) disease is locally advanced or metastatic, 2) disease harbors HER2 (ERBB2) tyrosine kinase domain activating mutations, as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ibrance

Products Affected

- **IBRANCE ORAL CAPSULE 125 MG**
- **IBRANCE ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of HR-positive, HER2-negative advanced or metastatic breast cancer -AND- meets one of the following (1. or 2.): 1) documentation of use with an aromatase inhibitor as initial endocrine-based therapy -OR- 2) documentation of use with fulvestrant in patients with disease progression following endocrine therapy. Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (3-6): 3) disease is HR-positive, HER2-negative 4) disease is PIK3CA-mutated, as detected by an FDA-authorized test 5) the member is using in combination with inavolisib and fulvestrant 6) the member has experienced recurrence on or after completing adjuvant endocrine therapy. Documentation of locally advanced or metastatic breast cancer -AND- all of the following (7-9): 7) disease is HR-positive, HER2-positive, 8) will be used following induction treatment, 9) will be used in combination with trastuzumab (with or without pertuzumab) and endocrine therapy as maintenance treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Ibtrozi

Products Affected

- **IBTROZI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iclusig

Products Affected

- **ICLUSIG**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of newly-diagnosed chronic phase CML
Required Medical Information	Documentation of T3151+ chronic myeloid leukemia (CML) -OR- documentation of chronic phase CML and member has experienced resistance or intolerance to at least two prior kinase inhibitors -OR- documentation of accelerated phase or blast phase CML and no other kinase inhibitor is indicated -OR- member is using Iclusig as monotherapy and meets one of the following (1-2): 1) documentation of T3151+ acute lymphoblastic leukemia (ALL) 2) documentation of Ph+ ALL and no other tyrosine kinase inhibitor therapy is indicated -OR- member has newly diagnosed Ph+ ALL and is using Iclusig in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Idhifa

Products Affected

- **IDHIFA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IG

Products Affected

- **GAMMAGARD LIQUID**
- **GAMMAGARD LIQUID ERC**
- **GAMMAGARD S-D (IGA**
- **PRIVIGEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For select diagnoses the following apply 1) For Myasthenia Gravis Syndrome, documentation that the patient is refractory to at least one other standard therapy (e.g., cholinesterase inhibitors, corticosteroids, azathioprine) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 2) For Multiple Sclerosis, patient is refractory to at least one other standard therapy (e.g., interferons) given in therapeutic doses over at least 3 months OR is intolerant/contraindication to those standard therapies. 3) For B-cell CLL, associated with recurrent bacterial infections OR with Associated Hypogammaglobulinemia defined as IgG level less than 600mg/dL or evidence of a specific antibody deficiency. 4) For Bone Marrow Transplantation, the member is 20 years of age or older and within the first 100 days after transplantation. 5) For Dermatomyositis/Polymyositis, trial and failure, intolerance, or contraindication to at least one standard first line therapy (i.e. corticosteroids or immunosuppressants). 6) For Pediatric HIV, the patient is less than 13 y.o. who have entry CD4 lymphocyte count greater than or equal to 200/mcl and IgG less than 400mg/dL OR recurrent bacterial infections. 7) For Guillain-Barre syndrome, impaired function by objective assessment and/or objective findings on physical exam at the time of initial therapy and IVIG therapy must be initiated within 2 weeks of symptom onset. 8) For Autoimmune Mucocutaneous Blistering Diseases (e.g. Stevens-Johnson Syndrome), trial and failure, intolerance, or contraindication to at least one conventional therapy (e.g. corticosteroids) or the patient has rapidly progressive disease in which a clinical response could not be affected quickly enough using conventional agents.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	Covered under Part B when administered in the home to a member with a diagnosis of primary immunodeficiency disease.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Myasthenia Gravis syndrome, Multiple Sclerosis, Polymyositis, Bone Marrow Transplant, Pediatric HIV, Guillain-Barre syndrome, Autoimmune Mucocutaneous Blistering Diseases
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Imbruvica

Products Affected

- **IMBRUVICA ORAL CAPSULE 140 MG, 70 MG**
- **IMBRUVICA ORAL SUSPENSION**
- **IMBRUVICA ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For chronic graft versus host disease, previous treatment with at least 1 prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For suspension, inability to swallow oral tablets or oral capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Imkeldi

Products Affected

- **IMKELDI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Ph+ chronic myeloid leukemia -AND- one of the following: 1) member is newly diagnosed in chronic phase 2) member is in blast crisis, the accelerated phase or the chronic phase after failure of interferon-alpha therapy. Documentation of Ph+ acute lymphocytic leukemia -AND- one of the following: 1) for adults, member has relapsed or refractory disease 2) for pediatric patients, member is newly diagnosed and will be using imatinib in combination with a chemotherapy regimen. Documentation of adult agressive systemic mastocytosis -AND- one of the following: 1) documentation that the member does not have D816V c-KIT status 2) member's c-KIT status is unknown. For gastrointestinal stromal tumors (GIST) one of the following: 1) member has diagnosis of KIT (CD117)-positive unresectable and/or metastatic malignant GIST 2) imatinib will be used as adjuvent treatment following resection of KIT (CD117)-positive GIST. Documentation of unresectable, recurrent or metastatic dermatofibrosarcoma protuberans. Documentation of Hypereosinophilic Syndrome/Chronic Eosinophilic Leukemia. Documentation of myeloproliferative disease -AND- disease is associated with PDGFR gene rearrangements. Documentation of myelodysplastic syndrome -AND- disease is associated with PDGFR gene rearrangements.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Documentation of an inability to swallow oral tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inbrija

Products Affected

- **INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use for the treatment of intermittent off episodes of Parkinson's disease while on carbidopa/levodopa
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of positive clinical response
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Increlex

Products Affected

- **INCRELEX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of severe primary IGF-1 deficiency and all of the following: 1) Normal or elevated response (greater than 10 ng/ml) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 2) Serum IGF-1 concentration that is less than or equal to three (3) standard deviations below the normal value based on laboratory reference range. 3) Height less than or equal to three (3) standard deviations below normal (at or below the third percentile for gender and age). 4) Member has open epiphyses. -OR- Documentation of diagnosis of growth hormone deficiency caused by gene deletion and all of the following: 1) Growth velocity at least 2 standard deviations below the age-appropriate mean or height at least 2.25 standard deviations below the age-appropriate mean. 2) Subnormal response (less than 10 ng/mL) to two (2) of the following standard growth hormone stimulation tests: arginine, clonidine, glucagon, insulin, levodopa, propranolol. 3) Development of neutralizing antibodies to growth hormone product(s). 4) Member has open epiphyses
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For pediatric patients, reauthorization requires open epiphyses.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Ingrezza

Products Affected

- **INGREZZA INITIATION PK(TARDIV)**
- **INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG**
- **INGREZZA SPRINKLE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of tardive dyskinesia -OR- documentation of chorea associated with Huntington's disease.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Inluriyo

Products Affected

- **INLURIYO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) disease is ER-positive, HER2-negative, 2) disease harbors an ESR1 mutation, as detected by an FDA-approved test, 3) member has experienced disease progression following at least one line of endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inlyta

Products Affected

- **INLYTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced renal cell carcinoma -AND- one of the following (1-2): 1) if using axitinib as first line therapy, member is using axitinib in combination with avelumab or pembrolizumab 2) if using axitinib as a single-agent, member has been treated with at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inqovi

Products Affected

- **INQOVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For myelodysplastic syndrome, one of the following is required (1-2): 1) French American-British MDS subtypes of refractory anemia, refractory anemia with ringed sideroblasts or refractory anemia with excess blasts, 2) International Prognostic Scoring System group of intermediate-1, intermediate-2 or high-risk. For newly-diagnosed acute myeloid leukemia, use in combination with venetoclax -AND- one of the following is required (3-4): 3) age greater than or equal to 75 years of age, 2) 1 or more comorbidity that precludes the use of intensive induction chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Inrebic

Products Affected

- **INREBIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate-2 or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) -AND- If a new start, baseline platelet count of greater than or equal to 50x10*9/L
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Insulin Supplies

Products Affected

- **ALCOHOL PADS**
- **GAUZE PAD TOPICAL BANDAGE 2 X 2 "**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation the product is being used for the delivery of insulin into the body.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Interferon Alfa

Products Affected

- **PEGASYS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Interferon Beta

Products Affected

- **AVONEX INTRAMUSCULAR PEN INJECTOR KIT**
- **AVONEX INTRAMUSCULAR SYRINGE KIT**
- **BETASERON SUBCUTANEOUS KIT**
- **PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML**
- **PLEGRIDY SUBCUTANEOUS**

SYRINGE 125 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IPF Agents

Products Affected

- *nintedanib*
- *pirfenidone oral capsule*
- *pirfenidone oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 50% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30%. For nintedanib only, documentation of systemic sclerosis-associated interstitial lung disease -AND- baseline forced vital capacity (FVC) of at least 40% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- documentation of a chest computed tomography (CT) scan demonstrating greater than or equal to 10% pulmonary fibrosis. For nintedanib only, documentation of chronic fibrosing interstitial lung disease with progressive phenotype -AND- chest computing tomography (CT) scan demonstrating greater than 10% fibrosing disease -AND- baseline forced vital capacity (FVC) of at least 45% and a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 30% -AND- disease progression in previous 24 months shown by one of the following : 1. Relative decline in FVC greater than or equal to 10% predicted 2. Relative decline in FVC greater than or equal to 5% but less than 10% predicted and either worsening of respiratory symptoms or increased extent of fibrotic changes on CT 3. Worsening of respiratory symptoms and increasing extent of fibrotic changes on CT
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iressa

Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	Use in tumors with EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.
Required Medical Information	Documentation of diagnosis -AND- the following: 1) EGFR exon 19 deletion mutations or exon 21 (L858R) mutations as detected by an FDA-approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Itovebi

Products Affected

- **ITOVEBI ORAL TABLET 3 MG, 9 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endocrine-resistant, locally advanced or metastatic breast cancer -AND- all of the following (1-4): 1) disease is HR-positive, HER2-negative 2) disease is PIK3CA-mutated, as detected by an FDA-approved test 3) the member is using in combination with palbociclib and fulvestrant 4) the member has experienced recurrence on or after completing adjuvant endocrine therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Itraconazole

Products Affected

- *itraconazole oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If using for diagnosis of onychomycosis, confirmation through positive laboratory testing (e.g. KOH preparation, fungal culture, or nail biopsy) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Onychomycosis: 3 months. All other indications: 3 months initial, 12 months reauth
Other Criteria	Documentation of trial/failure or intolerance of amphotericin b must be provided for approval in patients with aspergillosis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ivermectin Oral

Products Affected

- *ivermectin oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of strongyloidiasis of the intestinal tract (non-disseminated disease) or onchocerciasis -AND- Member weighs greater than or equal to 15kg.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Iwilfin

Products Affected

- **IWILFIN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of high-risk neuroblastoma (HRNB) -AND- partial response to anti-glycolipid disialoganglioside (GD2) immunotherapy (e.g., dinutuximab, naxitamab).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jakafi

Products Affected

- **JAKAFI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis per an accepted risk stratification tool for myelofibrosis (e.g., International Prognostic Scoring System [IPSS]) and if a new start, baseline platelet count of greater than or equal to 50x10*9/L -OR- documentation of polycythemia vera and inadequate response or intolerance to hydroxyurea -OR- Documentation of steroid refractory acute graft-versus-host disease and prior therapy with at least one systemic corticosteroid -OR- Documentation of chronic graft-versus-host disease with prior failure of at least one systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Baseline platelet count to be provided.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jascayd

Products Affected

- **JASCAYD**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis -AND- baseline forced vital capacity (FVC) of at least 45% -AND- a percent predicted diffusing capacity of the lungs of carbon monoxide (DLCO) of at least 25% -AND- 1 of the following (1-3): 1) Concomitant use of a stable dose of Esbriet (pirfenidone) or Ofev (nintedanib). For concomitant Esbriet (pirfenidone) therapy, Jascayd dose will remain 18 mg orally twice daily. 2) Receiving Jascayd as monotherapy and intolerance to Esbriet (pirfenidone) and Ofev (nintedanib), 3) Receiving Jascayd as monotherapy and contraindication to Esbriet (pirfenidone) and Ofev (nintedanib). Diagnosis of progressive pulmonary fibrosis (PPF) -AND- documentation of a chest computed tomography (CT) scan demonstrating at least 10% pulmonary fibrosis -AND- a baseline FVC of at least 45% -AND- a percent predicted DLCO of at least 25% -AND- at least 2 of the following (4-6): 4) Worsening of respiratory symptoms 5) Physiological evidence of disease progression defined as 1 of the following (a or b): a) Decline in FVC of at least 5% (relative from baseline). b) Decline in percent predicted DLCO of at least 10% (relative from baseline). 6) Radiological evidence of disease progression defined as increased extent of fibrotic changes on imaging (e.g., chest computed tomography scan) -AND- One of the following (7-9): 7) Concomitant use of a stable dose of Ofev (nintedanib), 8) Treatment naive to antifibrotic treatment, or 9) Previously discontinued Ofev (nintedanib) with no plans to initiate or resume background PPF treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jaypirca

Products Affected

- **JAYPIRCA ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory mantle cell lymphoma, member has received at least two (2) previous lines of systemic therapy, at least one (1) of which was a BTK inhibitor. For relapsed or refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia, member has previously been treated with a covalent BTK inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Jideytro

Products Affected

- **JIDEYTRO ORAL TABLET 100 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer (NSCLC) that is ROS1-positive and member has received a prior ROS1 kinase inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Joenja

Products Affected

- **JOENJA ORAL TABLET 70 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of activated phosphoinositide 3-kinase delta syndrome (APDS) with genetic confirmation of variant in PIK3CD or PIK3R1 gene.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Jynarque

Products Affected

- *tolvaptan (polycys kidney dis) oral tablet*
- *tolvaptan (polycys kidney dis) oral tablets, sequential*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of rapidly progressing autosomal dominant polycystic kidney disease defined by one of the following: 1.) Historical decline in eGFR greater than or equal to 3mL/min/1.73 m*2 within a 12 month period, or 2.) Mayo imaging classification of 1C, 1D, or 1E.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, prescriber attestation of slowed decline of kidney function
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kalydeco

Products Affected

- **KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG**
- **KALYDECO ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to ivacaftor based on clinical and or in vitro assay (e.g. G551D, G1244E, G1349D)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kerendia

Products Affected

- **KERENDIA**

PA Criteria	Criteria Details
Exclusion Criteria	eGFR less than 25 mL/min/1.73 m*2, serum potassium greater than 5.5 mEq/L
Required Medical Information	Documentation of type 2 diabetes mellitus with chronic kidney disease (T2DM+CKD) or heart failure (HF) with left ventricular ejection fraction (LVEF) of greater than or equal to 40% -AND- one of the following (1. 2. or 3.): 1) concomitant use of a sodium-glucose Cotransporter-2 (SGLT2) inhibitor 2) therapeutic failure to at least one SGLT2 inhibitor or 3) contraindication or intolerance to at least one SGLT2 inhibitor.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that signs or symptoms of hyperkalemia are not present.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kesimpta

Products Affected

- **KESIMPTA PEN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kineret

Products Affected

- **KINERET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For Deficiency of Interleukin-1 Receptor Antagonist (DIRA), therapeutic failure or intolerance to at least one (1) corticosteroid, or all corticosteroids are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kisqali

Products Affected

- **KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative -AND- one of the following (1-3): 1) member is using ribociclib in combination with an aromatase inhibitor as initial endocrine-based therapy 2) member is using ribociclib in combination with fulvestrant and member is using fulvestrant as initial endocrine-based therapy or member has experienced disease progression on endocrine therapy. 3) disease is classified as stage II or stage III early breast cancer at high risk of recurrence, ribociclib is being used in combination with an aromatase inhibitor as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Klisyri

Products Affected

- **KLISYRI (250 MG)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of actinic keratosis of the face or scalp -AND- Therapeutic failure or intolerance to 1 of the following 1) generic topical imiquimod 5% cream 2) generic fluorouracil 5% topical cream 3) generic fluorouracil topical solution
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation that the member has previously experienced complete or partial clearance of actinic keratosis lesions with Klisyri -AND- additional course of therapy is required for recurrence of actinic keratosis -AND- member is restarting therapy at least 60 days after cessation of an initial Klisyri 5-day course.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Komzifti

Products Affected

- **KOMZIFTI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory acute myeloid leukemia, the member has a susceptible NPM1 mutation -AND- the member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Korlym

Products Affected

- *mifepristone oral tablet 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of endogenous Cushing's syndrome -AND- Patient is not a candidate for surgery or where surgery has failed -AND- one of the following (1 or 2): 1) Diagnosis of diabetes, 2) Glucose intolerance
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Koselugo

Products Affected

- **KOSELUGO ORAL CAPSULE 10 MG, 25 MG**
- **KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- for neurofibromatosis type 1 (NF1), documentation of symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Koselugo granules, 1 of the following is required (1-2): 1) inability to swallow oral capsules, 2) body surface area of less than 0.55 m*2
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Krazati

Products Affected

- **KRAZATI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Kuvan

Products Affected

- **JAVYGTOR**
- *sapropterin*
- **ZELVYSIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of PKU -AND- documented baseline Phe level greater than 6 mg/dL -AND- clinical documentation of current weight -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, initial therapy has resulted in decrease in phenylalanine levels from baseline or current phenylalanine levels within the range of 120-360 micromol/L -AND- clinical documentation of current weight is required -AND- sapropterin dihydrochloride dose does not exceed 20 mg/kg/day.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Kygevvi

Products Affected

- **KYGEVVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Thymidine Kinase 2 Deficiency (TK2d) -AND- Pathogenic mutation in TK2 gene, as confirmed by genetic testing -AND- symptom onset on or before 12 years of age
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Latuda

Products Affected

- *lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. If medication is being used for bipolar 1 disorder, documentation of trial and failure or intolerance to one other formulary medication indicated in bipolar 1 disorder (e.g. quetiapine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lazcluze

Products Affected

- **LAZCLUZE ORAL TABLET 240 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) -AND- all of the following (1-4): 1) disease harbors EGFR exon 19 deletions or EGFR exon 21 (L858R) substitution mutation, as detected by an FDA-approved test 2) disease is locally advanced or metastatic, 3) member is treatment naive for advanced disease, 4) using in combination with amivantinib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lenvima

Products Affected

- **LENVIMA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of differentiated thyroid cancer -AND- meets all of the following (1-3): 1) disease is locally recurrent or metastatic 2) disease is progressive 3) disease is radioactive iodine refractory. Documentation of advanced renal cell carcinoma -AND- one of the following (4-5): 4) member is using lenvatinib in combination with pembrolizumab and is using lenvatinib and pembrolizumab as first-line treatment 5) member is using lenvatinib in combination with everolimus and has experienced therapeutic failure or intolerance to one prior anti-angiogenic therapy. Documentation of unresectable hepatocellular carcinoma -AND- member is using lenvatinib as first-line treatment. Documentation of endometrial cancer -AND- meets all of the following (6-10): 6) member is using lenvatinib in combination with pembrolizumab 7) disease is advanced 8) disease is not classified as microsatellite instability-high or disease is classified as mismatch repair proficient as determined by an FDA-approved test 9) member has experienced disease progression following prior systemic therapy 10) member is not a candidate for curative surgery or radiation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Leukine

Products Affected

- **LEUKINE INJECTION RECON SOLN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis: following induction chemotherapy in patients who are 55 years or older with acute myelogenous leukemia (AML) -OR- mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation -OR- acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation -OR- acceleration of myeloid reconstitution following allogeneic BMT -OR- treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic BMT -OR- following acute exposure to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Leukotriene Modifiers

Products Affected

- *zileuton*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of asthma -AND- trial/failure of generic montelukast and generic zafirlukast
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lidoderm

Products Affected

- *lidocaine topical adhesive patch,medicated 5 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of postherpetic neuralgia (PHN) or diabetic peripheral neuropathy (DPN) -AND- One of the following (1-3): 1) trial and failure of 1 other agent used to treat diagnosis (e.g. gabapentin for PHN, duloxetine for DPN), 2) inability to swallow oral medication, 3) unable to take an oral medication due to potential adverse events (e.g. sedation).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Diabetic peripheral neuropathy
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lifyorli

Products Affected

- **LIFYORLI ORAL CAPSULE 125 MG/DAY(100 MG X1-25MG X1), 125 MG/DAY(100 MG X1-25MG X3), 125 MG/DAY(100 MG X1-25MG X9), 150 MG/DAY(100 MG X1-25MG X2), 150 MG/DAY(100 MG X1-25MG X3), 150 MG/DAY(100 MG X1-25MG X9)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For epithelial ovarian, fallopian tube, or primary peritoneal cancer, all of the following (1-3): 1) disease is platinum-resistant, 2) member has received 1-3 prior systemic treatment regimens, at least 1 of which included bevacizumab, 3) using in combination with nab-paclitaxel.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Litfulo

Products Affected

- **LITFULO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For severe alopecia areata, therapeutic failure/intolerance to an intralesional corticosteroid or high potency topical corticosteroid, or contraindication to all.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Livtencity

Products Affected

- **LIVTENCITY**

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of CMV prophylaxis.
Required Medical Information	Documentation of refractory cytomegalovirus infection or disease as evidenced by antigenemia or polymerase chain reaction (PCR) test -AND- all of the following (1-3): 1) member weighs at least 35 kg, 2) member is a recipient of hematopoietic stem cell transplant -OR- solid organ transplant. 3) member has experienced therapeutic failure to one of the following: ganciclovir, valganciclovir, cidofovir, or foscarnet.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of a previous reduction in CMV DNA level -AND- documentation of one of the following (1-3): 1) new onset symptomatic CMV infection, 2) virologic relapse with treatment-emergent maribavir resistance or 3) continued antiviral treatment is required to achieve virologic clearance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Loargys

Products Affected

- **LOARGYS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Arginase 1 Deficiency (ARG1-D) -AND- Used in conjunction with dietary protein restriction -AND- Diagnosis is confirmed by 1 of the following (1-3): 1) genetic testing, 2) newborn screening, or 3) elevated arginine levels in plasma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lokelma

Products Affected

- **LOKELMA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hyperkalemia as defined by serum potassium level between 5.1 and 7.4 mmol/L on at least two (2) screenings -AND- Modification of medications to reduce serum potassium levels were not successful, when applicable
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in serum potassium levels following Lokelma administration and continued treatment for hyperkalemia is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lonsurf

Products Affected

- **LONSURF**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer as a single agent or in combination with bevacizumab in patients who have previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy -OR- documentation of metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan, and if appropriate, HER2/neu-targeted therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lorbrena

Products Affected

- **LORBRENA ORAL TABLET 100 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of ALK-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lotronex

Products Affected

- *alosetron oral tablet 0.5 mg, 1 mg*

PA Criteria	Criteria Details
Exclusion Criteria	For irritable bowel syndrome (IBS): Exclude if male gender
Required Medical Information	Documentation of chronic severe diarrhea-predominant IBS -AND- trial and failure or intolerance to one anti-diarrheal (e.g. loperamide), anti-spasmodic, or tricyclic antidepressant, or contraindication to all
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 weeks initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation that symptoms of IBS continue to persist AND positive clinical response.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lumakras

Products Affected

- **LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- member has received at least one prior systemic therapy -AND- using as a single agent. Documentation of metastatic colorectal cancer -AND- disease is KRAS G12C-mutated as determined by an FDA approved test -AND- using in combination with panitumumab -AND- prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lybalvi

Products Affected

- **LYBALVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of schizophrenia or bipolar I disorder -AND- One of the following (1 or 2): 1) Therapeutic failure, intolerance or contraindication to generic risperidone and generic quetiapine, 2) Currently stable and responding to olanzapine but experiencing weight gain from the medication. If using for maintenance monotherapy treatment of bipolar 1 disorder, only criterion 2 is required.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lynparza

Products Affected

- **LYNPARZA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) in complete or partial response to first-line platinum-based chemotherapy 2) diagnosis of deleterious or suspected deleterious BRCA mutated disease or disease is associated with homologous recombination deficiency (HRD) positive status with a deleterious or suspected deleterious BRCA mutation or genomic instability and will be using in combination with bevacizumab. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) deleterious or suspected deleterious germline or somatic BRCA mutation 2) in complete or partial response to platinum-based chemotherapy. For deleterious or suspected deleterious gBRCAm, HER2-negative breast cancer, 1 of the following (1-2): 1) classified as high-risk, early breast cancer and has been treated with neoadjuvant or adjuvant chemotherapy 2) has been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting and if hormone receptor (HR)-positive, has been previously treated with or considered inappropriate for treatment with endocrine therapy. For metastatic pancreatic adenocarcinoma, all of the following (1-2): 1) a deleterious or suspected deleterious gBRCA mutation 2) did not progress on at least 16 weeks of a first-line platinum-based chemotherapy regimen. For deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene mutation metastatic castration-resistant prostate cancer, all of the following (1-2): 1) progressed following prior treatment with enzalutamide or abiraterone 2) concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or has had a bilateral orchiectomy. For deleterious or suspected deleterious BRCA mutation metastatic castration-resistant prostate cancer, using in combination with all of the following (1-2): 1) abiraterone 2) prednisone or prednisolone.
Age Restrictions	

PA Criteria	Criteria Details
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Lyrica

Products Affected

- *pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg*
- *pregabalin oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. When using pregabalin products concomitantly with an opiate agonist, attestation of an intent to monitor and address concomitant drug-drug interaction adverse events for opiate potentiators.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Lytgobi

Products Affected

- **LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-2): 1) disease harbors FGFR2 fusions or other rearrangements 2) member has experienced therapeutic failure or intolerance to at least one prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Mavyret

Products Affected

- **MAVYRET ORAL PELLETS IN PACKET**
- **MAVYRET ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance
Age Restrictions	Deny if less than 3 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Megace

Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Mekinist

Products Affected

- **MEKINIST ORAL RECON SOLN**
- **MEKINIST ORAL TABLET 0.5 MG, 2 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For trametinib monotherapy, documentation of unresectable or metastatic melanoma -AND- meets all of the following (1-2): 1) member has a BRAF V600E or V600K mutation 2) member is BRAF inhibitor treatment naive. For use in combination with dabrafenib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with dabrafenib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA approved test when FDA indicated. For use in combination with dabrafenib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Mekinist powder for oral solution, attestation of inability to swallow Mekinist (trametinib) tablets is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Mektovi

Products Affected

- **MEKTOVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, all of the following (1-2): 1) BRAF V600E or V600K mutation status, as detected by an FDA-authorized test 2) using in combination with encorafenib. For metastatic non-small cell lung cancer, all of the following (1-2): 1) BRAF V600E mutation status, as detected by an FDA-authorized test 2) using in combination with encorafenib.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Metyrosine

Products Affected

- *metyrosine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of pheochromocytoma defined by 1 of the following (1 or 2): 1) Elevated metanephrines in plasma or urine, or 2) Tumor evidence from CT scan or MRI -AND- Documentation of 1 of the following (3., 4., or 5.): 3) Planned resection surgery, 4) Resection surgery is contraindicated, or 5) malignant pheochromocytoma. -AND- therapeutic failure, contraindication, or intolerance to a selective alpha-blocker (e.g., doxazosin, prazosin, terazosin).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Modeyso

Products Affected

- **MODEYSO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of diffuse midline glioma -AND- disease harbors an H3 K27M mutation -AND- member has experienced progressive disease following prior therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

MTX Injection

Products Affected

- **RASUVO (PF)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis AND intolerance or contraindication to oral generic methotrexate tablets. For rheumatoid arthritis and juvenile idiopathic arthritis, disease is classified as severe and active -AND- therapeutic failure or intolerance to first-line therapy (e.g., nonsteroidal anti-inflammatory drug [NSAID]). For psoriasis, disease is classified as severe, recalcitrant, and disabling -AND- therapeutic failure or intolerance to at least one (1) other standard psoriasis therapy (e.g., cyclosporine)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Mulpleta

Products Affected

- **MULPLETA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of thrombocytopenia and chronic liver disease - AND- beneficiary is scheduled to undergo a procedure -AND- trial and failure or intolerance to Doptelet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Myasthenia Gravis

Products Affected

- **ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML, 23 MG/0.574 ML, 32.4 MG/0.81 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of generalized myasthenia gravis (gMG) -AND- Anti-acetylcholine receptor (AChR) antibody-positive -AND- Therapeutic failure, contraindication, or intolerance to generic pyridostigmine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of gMG signs/symptoms improvement (e.g., speech, swallowing, mobility, and/or respiratory function) -OR- decreased gMG exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Namzaric

Products Affected

- *memantine-donepezil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis and intolerance to generic memantine and generic donepezil
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nayzilam

Products Affected

- **NAYZILAM**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nemluvio

Products Affected

- **NEMLUVIO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of prurigo nodularis -OR- Documentation of moderate to severe atopic dermatitis and one of the following (1-3): 1) trial and failure or intolerance to at least one topical corticosteroid or one topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus), 2) severe atopic dermatitis and the member is incapable of applying topical therapies due to the extent of body surface area involvement, or 3) severe atopic dermatitis and topical therapies are contraindicated due to severely damaged skin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For prurigo nodularis, must follow recommended dosing guidelines based upon weight. For atopic dermatitis, must follow recommended dosing based on FDA approved dosing guidelines. For induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For prurigo nodularis reauthorization, attestation of reduction in itch or number of nodules or lesions from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Nerlynx

Products Affected

- **NERLYNX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of early-stage, HER2-positive breast cancer -AND- meets all of the following: 1)member has received adjuvant trastuzumab-based therapy 2) member is using neratinib as a single agent. Documentation of advanced HER-2 positive, or metastatic HER2-positive breast cancer and meets all of the following 1) using neratinib in combination with capecitabine 2) member has received two or more prior anti-HER2 based regimens in the metastatic setting.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nexavar

Products Affected

- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally recurrent or metastatic, progressive, differentiated thyroid carcinoma, refractory to radioactive iodine treatment
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nexletol

Products Affected

- **NEXLETOL**
- **NEXLIZET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	1.HeFH supported by 1 of the following (a-d): a) causal mutation of FH by genetic testing, b) untreated LDL-C at least 190mg/dL w/physical signs of FH (e.g. xanthomas, xanthelasma), c) untreated LDL-C at least 160mg/dL before 20 yrs old w/physical signs of FH, d) diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria w/score over 8 points, definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C, or if very high risk, LDL-C at least 55mg/dL or non-HDL-C at least 85mg/dL -AND- Attestation of combination use with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. 2.ASCVD or attestation of high risk for CVD -AND- Therapeutic failure/intolerance/contraindication to ezetimibe. 3. Primary Hyperlipidemia/Hypercholesterolemia (w/o ASCVD or HeFH) AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C -AND- Attestation of combination use with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. -AND- Therapeutic failure/intolerance/contraindication to ezetimibe.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For reauthorization of ASCVD or High ASCVD Risk, attestation of need for continued therapy is required. For reauthorization associated with all other diagnoses, documentation showing an LDL-C reduction from baseline -AND- attestation of continued use in combination with other LDL-C lowering therapies or concomitant LDL-C lowering therapy is not possible. Therapeutic failure to a max tolerated statin or documented statin intolerance is required. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statin which resolved upon discontinuation of statin or documentation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ninlaro

Products Affected

- **NINLARO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- previous treatment with at least 1 prior therapy -AND- used in combination with lenalidomide and dexamethasone
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nitisinone

Products Affected

- *nitisinone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hereditary tyrosinemia type 1 (HT-1) confirmed by biochemical or genetic testing -AND- dietary restriction of tyrosine and phenylalanine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Northera

Products Affected

- *droxidopa oral capsule 100 mg, 200 mg, 300 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of neurogenic orthostatic hypotension caused by primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, or pure autonomic failure), dopamine beta-hydroxylase deficiency or non-diabetic autonomic neuropathy -AND- documentation of inadequate response, intolerance or contraindication to preferred generic alternative midodrine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of increase from baseline of systolic or diastolic blood pressure upon standing -OR- attestation of decrease from baseline of neurogenic orthostatic hypotension symptoms upon standing (e.g., dizziness, feeling faint, etc.).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nourianz

Products Affected

- **NOURIANZ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease experiencing off episodes -AND- Used as adjunct to levodopa/carbidopa -AND- trial/failure, contraindication or intolerance to one of the following generic products: ropinirole, pramipexole, entacapone, selegiline, or rasagiline
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nubeqa

Products Affected

- **NUBEQA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of non-metastatic castration-resistant prostate cancer or metastatic castration-sensitive prostate cancer -AND- one of the following (1-2): 1) using in combination with a GnRH analog 2) member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nuedexta

Products Affected

- *dextromethorphan-quinidine*
- **NUEDEXTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation supporting improvement in symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nuplazid

Products Affected

- **NUPLAZID**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of hallucinations and delusions associated with Parkinson's disease psychosis
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Nurtec

Products Affected

- **NURTEC ODT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For Episodic Migraine Prevention, defined as 4-14 migraine days per month, the following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound or overutilization (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use). For acute treatment of migraine with or without aura, trial and failure of one generic triptan or all triptans are contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization of episodic migraine prevention, attestation of reduction in migraine frequency. For reauthorization of acute treatment of migraine, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Nuvigil

Products Affected

- *armodafinil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography or home sleep apnea test as established in accordance with ICSD or DSM V criteria acceptable for all indications
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Octreotide

Products Affected

- *octreotide acetate injection solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- therapeutic failure or cannot be treated with surgical resection, pituitary irradiation or bromocriptine mesylate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Odactra

Products Affected

- **ODACTRA**

PA Criteria	Criteria Details
Exclusion Criteria	Asthma (severe, unstable or uncontrolled), concomitant sublingual or subcutaneous immunotherapy
Required Medical Information	Documentation of allergic rhinitis due to house dust mites -AND- allergic rhinitis with or without conjunctivitis has been confirmed by skin testing for licensed house dust mite allergen extracts or in vitro testing for IgE antibodies to D. pteronyssinus or D. Farina -AND- trial and failure or intolerance to an intranasal steroid and an oral non-sedating antihistamine, intranasal antihistamine or intranasal anticholinergic agent
Age Restrictions	Deny if less than 5 years of age or greater than 65 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Member must also be prescribed an epinephrine auto injector. For reauthorization, attestation of improved allergy symptoms is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Odomzo

Products Affected

- **ODOMZO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced basal cell carcinoma (laBCC) that has recurred following surgery or radiation therapy or for use in patients who are not candidates for surgery or radiation therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ogsiveo

Products Affected

- **OGSIVEO ORAL TABLET 100 MG, 150 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of progressing desmoid tumor(s) requiring systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ojemda

Products Affected

- **OJEMDA ORAL SUSPENSION FOR RECONSTITUTION**
- **OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500 MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory pediatric low-grade glioma -AND- one of the following (1-2): 1) BRAF fusion or rearrangement, or 2) BRAF V600K mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Ojjaara

Products Affected

- **OJJAARA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of anemia.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Onfi

Products Affected

- *clobazam oral suspension*
- *clobazam oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ongentys

Products Affected

- **ONGENTYS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Parkinson's disease experiencing off episodes -AND- Used as adjunct to levodopa/carbidopa -AND- trial/failure, contraindication or intolerance to the following (1. and 2.): 1) entacapone 2) one of the following generic products: rasagiline, pramipexole, ropinirole, selegiline
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Onureg

Products Affected

- **ONUREG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute myeloid leukemia that has achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy -AND- Inability to complete intensive curative therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Opipza

Products Affected

- **OPIPZA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- 1 of the following (1-2): 1) intolerance to generic aripiprazole tablets, 2) inability to swallow oral tablets. If medication is being used for major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Orencia

Products Affected

- **ORENCIA CLICKJECT**
- **ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Psoriatic Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. For psoriatic arthritis, patients over 18 years of age must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Cosentyx, Otezla/Otezla XR, Skyrizi SC, Rinvoq, and Xeljanz/Xeljanz XR. For polyarticular juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq/Rinvoq LQ, and Xeljanx/Xeljanz solution. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Orgovyx

Products Affected

- **ORGOVYX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For induction therapy dosing, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimens per indication.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Orkambi

Products Affected

- **ORKAMBI ORAL GRANULES IN PACKET**
- **ORKAMBI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis and homozygous F508del mutation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Orserdu

Products Affected

- **ORSERDU ORAL TABLET 345 MG, 86 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-4): 1) member is male or a postmenopausal female 2) disease is ER-positive, HER2-negative 3) disease harbors an ESR1 mutation, as detected by an FDA-approved test 4) member has experienced disease progression on or after an endocrine based regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Otezla

Products Affected

- **OTEZLA**
- **OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47)**
- **OTEZLA XR**
- **OTEZLA XR INITIATION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For plaque psoriasis, all of the following (1-2): 1) member is greater than or equal to 18 years of age with mild-to-severe disease -OR- if the member is greater than or equal to 6 years of age and less than 18 years, weight is greater than or equal to 20 kg for Otezla or 50 kg for Otelza XR, and member has moderate-to-severe disease -AND- 2) inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For oral ulcers associated with Behcet's Disease, inadequate response or intolerance to one systemic therapy for prevention of recurrent oral ulcers. For psoriatic arthritis, if the member is greater than or equal to 6 years of age and less than 18 years and weight is greater than or equal to 20 kg for Otezla or 50 kg for Otelza XR.
Age Restrictions	Deny if less than 6 years of age for plaque psoriasis or psoriatic arthritis. Deny if less than 18 years of age for Behcet's Disease.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Ozempic Oral

Products Affected

- **OZEMPIC ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	Obesity or use for weight loss
Required Medical Information	Documentation of diabetes mellitus type 2
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Palynziq

Products Affected

- **PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of phenylketonuria -AND- all of the following criteria (1-3): 1) Baseline Phe level greater than 600 micromoles/L, 2) Failure or intolerance to existing management (i.e. sapropterin dihydrochloride)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in baseline pretreatment Phe levels -OR- blood Phe levels are less than or equal to 600 micromoles/L -OR- attestation that additional therapy with Palynziq is needed to allow adequate trial of maximum dose of 60mg per day for 16 weeks or member is not currently on 60mg per day dose.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Panretin

Products Affected

- **PANRETIN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions in patients with AIDS-related Kaposi Sarcoma (KS) who are not receiving systemic therapy for KS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pemazyre

Products Affected

- **PEMAZYRE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of unresectable locally advanced cholangiocarcinoma or metastatic cholangiocarcinoma -AND- all of the following (1-2): 1) disease harbors FGFR2 fusion or other rearrangement as detected by an FDA-approved test 2) member has experienced therapeutic failure or intolerance to at least on prior therapy. Documentation of relapsed or refractory myeloid/lymphoid neoplasms -AND- disease harbors an FGFR1 rearrangement.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Pentasa

Products Affected

- *mesalamine oral capsule, extended release 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Ulcerative Colitis (UC) -AND- Intolerance to 1 of the following (1-3): 1) generic Mesalamine 4 g/60 mL enema, 2) generic Mesalamine 400 mg DR capsule, 3) generic Mesalamine 1.2 g DR tablet
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of disease stability or beneficial response to therapy -AND- additional courses for UC treatment are necessary.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Pheburane

Products Affected

- **PHEBURANE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- intolerence to generic sodium phenylbutyrate.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Piqray

Products Affected

- **PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following: 1) HR-positive, HER2-negative tumor status, 2) PIK3CA mutation positive as detected by an FDA-approved test, 3) disease progression on or after an endocrine-based regimen, 4) Concomitant therapy with fulvestrant
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pomalyst

Products Affected

- *pomalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma, and combination use with dexamethasone, and previous trial of at least 2 therapies including lenalidomide and a proteasome inhibitor, and disease progression on or within 60 days of completion of the last therapy -OR- Documentation of AIDS-related Kaposi sarcoma (KS) after failure of highly active antiretroviral therapy (HAART) or in patients with KS who are HIV negative
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Posaconazole Tablet

Products Affected

- *posaconazole oral tablet,delayed release (dr/ec)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For aspergillus or candida infection prophylaxis, high risk of developing invasive Aspergillosis or Candidiasis infection due to being severely immunocompromised (e.g., hematopoietic stem cell transplant recipients with graft versus host disease, those with hematologic malignancies with prolonged neutropenia from chemotherapy) -AND- weight greater than 40 kg. For invasive aspergillosis infection, trial/failure or contraindication to voriconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Preferred Adalimumab

Products Affected

- **HADLIMA**
- **HADLIMA PUSHTOUCH**
- **HADLIMA(CF)**
- **HADLIMA(CF) PUSHTOUCH**
- **SIMLANDI(CF) AUTOINJECTOR**
- **SIMLANDI(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML**
- **YUFLYMA(CF) AI CROHN'S-UC-HS**
- **YUFLYMA(CF) AUTOINJECTOR**
- **YUFLYMA(CF) SUBCUTANEOUS SYRINGE KIT 40 MG/0.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe rheumatoid arthritis, inadequate response or intolerance to at least one DMARD (e.g. methotrexate, leflunomide). For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID). For moderate to severe polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For uveitis, inadequate response or intolerance to 1 immunosuppressant or corticosteroid, or all are contraindicated.
Age Restrictions	Deny if less than 18 years of age for Rheumatoid Arthritis, Psoriatic Arthritis, Plaque Psoriasis, and Ankylosing Spondylitis or less than 12 years of age for Hidradenitis Suppurative or Less than 6 years of age for Crohn's disease or Less than 5 years of age for Ulcerative Colitis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Uveitis
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	For plaque psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For rheumatoid arthritis therapy without concomitant methotrexate, doses above plan quantity limit will be approved aligned with recommended weekly dosing regimen. For pediatric ulcerative colitis and hidradenitis suppurativa, doses above plan quantity limit will be approved to align with recommended dosing regimen. Induction therapy or treatment regimens for other indications are aligned with plan quantity limit on Humira starter kit. These criteria are applicable to Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Preferred Ustekinumab

Products Affected

- **STEQEYMA SUBCUTANEOUS SOLUTION**
- **STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**
- **YESINTEK SUBCUTANEOUS SOLUTION**
- **YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease or Ulcerative Colitis, attestation of receiving or currently undergoing a single induction dose of IV ustekinumab within 2 months of initiating therapy with ustekinumab SC.
Age Restrictions	Deny if less than 18 years of age for Ulcerative Colitis. Deny if less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis. Deny if less then 2 years of age for Crohn's Disease.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	Must follow recommended dosing guidelines based upon weight. Induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For pediatric Crohn's disease, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Prenatal Vitamins

Products Affected

- **PRENATAL VITAMIN PLUS LOW IRON**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of nutritional supplementation required in a female of child-bearing potential during pre-conception, pregnancy, or lactation
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Prescription Drug Combo

Products Affected

- *acetaminophen-codeine oral solution 120-12 mg/5 ml*
- *acetaminophen-codeine oral tablet*
- **ALPRAZOLAM INTENSOL**
- *alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg, 2 mg, 3 mg*
- *buprenorphine*
- *butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg, 50-325-40-30 mg*
- *clonazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- *clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg*
- *clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg*
- **DIAZEPAM INTENSOL**
- *diazepam oral solution 5 mg/5 ml (1 mg/ml)*
- *diazepam oral tablet*
- *estazolam*
- *eszopiclone*
- *fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr*
- *flurazepam*
- *hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg*
- *hydromorphone oral liquid*
- *hydromorphone oral tablet 2 mg, 4 mg, 8 mg*
- **LORAZEPAM INTENSOL**
- *lorazepam oral tablet 0.5 mg, 1 mg, 2 mg*
- *methadone oral solution 10 mg/5 ml, 5 mg/5 ml*
- *methadone oral tablet 10 mg, 5 mg*
- *morphine concentrate oral solution*
- *morphine oral solution 10 mg/5 ml (2 mg/ml), 20 mg/5 ml (4 mg/ml)*
- *morphine oral tablet*
- *morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg*
- **NUCYNTA**
- *oxazepam*
- *oxycodone oral capsule*
- *oxycodone oral concentrate*
- *oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg*
- *oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg*
- *temazepam*
- *tramadol oral tablet 100 mg, 25 mg, 50 mg, 75 mg*
- *tramadol-acetaminophen*
- *triazolam*
- *zaleplon oral capsule 10 mg, 5 mg*
- *zolpidem oral tablet*
- *zolpidem oral tablet,ext release multiphase*
- *zolpidem sublingual*

PA Criteria	Criteria Details
Exclusion Criteria	

PA Criteria	Criteria Details
Required Medical Information	For concomitant use of opiate agonist and substance abuse therapy, documentation of acute pain condition (eg, acute traumatic injury) in which treatment with other agents would cause insufficient pain control or treatment for terminal illness related pain. For concomitant use of an opiate agonist, benzodiazepine, and centrally acting skeletal muscle relaxant, trial/failure of at least 2 other skeletal muscle relaxants (eg, methocarbamol, metaxalone), understanding these skeletal muscle relaxants are high-risk medications in geriatric patients, attestations that non-opiate (eg, NSAIDs) and non-benzodiazepine therapies (eg, SSRI, SNRI) have been considered, and attestation of intent to monitor & address concomitant drug-drug interaction adverse events. For concomitant use of opiate agonist and other opiate potentiators (eg, gabapentinoids, sedative-hypnotics), attestation of intent to monitor & address concomitant drug-drug interaction adverse events. For long acting opioid medications, the following apply (1-3): 1)Severe pain requiring daily, around-the-clock, long-term opioid treatment. 2)Not opioid naive. 3)Attestations that non-opiate alternative therapies have been explored (eg, NSAIDs), controlled substance history was reviewed in the state Prescription Drug Monitoring Program (PDMP), counseling on potential adverse effects of opioid analgesics, including risk of misuse, abuse, addiction.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Opiate tx for pain+subs. abuse, approve opiate x 1mo. All other combos and dx approve x 12mo.
Other Criteria	Opiate agonists will receive automatic approval if no recent claims for a substance abuse therapy (e.g. buprenorphine-naloxone) OR a benzodiazepine with a centrally acting skeletal muscle relaxant (e.g., carisoprodol) OR a gabapentinoid OR a sedative-hypnotic. Benzodiazepines (e.g. triazolam, alprazolam) will receive automatic approval if no recent claims for an opiate agonist with a centrally acting skeletal muscle relaxant (e.g. carisoprodol). Sedative-hypnotics (e.g. zolpidem) will receive automatic approval if no recent claims for an opiate agonist.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Prevymis Oral Pellets

Products Affected

- **PREVYMIS ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Deny if less than 6 months of age with HSCT. Deny if less than 12 years of age with kidney transplant.
Prescriber Restrictions	
Coverage Duration	7 months
Other Criteria	One of the following is required (1-2): 1) inability to swallow tablets, 2) unable to use Prevymis (letermovir) tablets due to body weight dosing limitations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Procysbi

Products Affected

- **PROCYSBI ORAL GRANULES DEL RELEASE IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- previous trial and failure, intolerance, or contraindication to Cystagon (cysteamine bitartrate immediate-release)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Prolia

Products Affected

- **ENOBY**
- **JUBBONTI**
- **PROLIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -And- For osteoporosis at high risk for fracture, meeting one of the following (1. thru 4.) 1) History of previous fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 (i.e. osteopenia) -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent. 4) Age 40 years or older with T-score between -1.0 and -2.5 -AND- History of glucocorticoid use for at least 3 months at a dose of 5mg per day or more of prednisone (or equivalent).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For osteoporosis and osteopenia, documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated -OR- very high risk for a fracture, defined by one of the following: fracture within past 12 months, fracture while on osteoporosis therapy, history of multiple fractures, fractures while on drugs causing skeletal harm, fall risk, history of injurious falls, T-score less than -3, FRAX 10 year major osteo fracture risk of greater than 30%, or FRAX 10 year hip fracture risk of greater than 4.5%. Covered under Part B for patients eligible for home health services when provider certifies that patient sustained bone fracture related to post-menopausal osteoporosis and is unable to learn the skills needed to self-administer the drug or is otherwise physically or mentally incapable of administering the drug or family/caregivers are unable or unwilling to administer the drug.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Provigil

Products Affected

- *modafinil*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of shift work sleep disorder (SWSD) substantiated by excessive sleepiness or insomnia that is temporarily associated with a recurring work schedule that overlaps the usual time for sleep -AND- Symptoms are accompanied by a reduction of total sleep time -AND- Symptoms experienced for at least 3 months -AND- Sleep log or actigraphy monitoring for at least 14 days including both work and free days -AND- Sleep disturbance is not better explained by another current sleep disorder, medical or neurological disorder, mental disorder, medication use, or substance use disorder. Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography or home sleep apnea test as established in accordance with ICSD or DSM V criteria acceptable for all indications. Diagnosis of fatigue associated with Multiple Sclerosis (MS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months

PA Criteria	Criteria Details
Other Criteria	For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Fatigue associated with Multiple Sclerosis (MS)
Part B Prerequisite	No
Prerequisite Therapy Required	No

Pulmonary Arterial Hypertension

Products Affected

- **ADEMPAS**
- *ambrisentan*
- *bosentan oral tablet*
- *macitentan*
- **OPSUMIT**
- **OPSYNVI**
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group. For bosentan and sildenafil in pediatric individuals, an exception to RHC will be allowed when the risk of RHC outweighs the benefit -AND- prescriber attests alternative studies have been completed (i.e. CT, MRI or specified test ruling out other causes of pulmonary hypertension). For Adempas, additional diagnosis of CTEPH as documented by right heart catheterization and V/Q scan substantiating mPAP greater than 20 mmHg at rest and PWP less than or equal to 15 mmHg and documented presence of occlusive thrombi within the pulmonary arteries will be approved.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Pulmozyme

Products Affected

- **PULMOZYME**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of cystic fibrosis -AND- Used in conjunction with standard therapies for management of cystic fibrosis to improve pulmonary function.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Inhalation solutions covered under Part B when administered in the home setting using a covered nebulizer (i.e. DME). For reauthorization, attestation of increase in FEV1 or decrease in number of hospitalizations or pulmonary exacerbations.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Qinlock

Products Affected

- **QINLOCK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of advanced gastrointestinal stromal tumor -AND- Prior treatment with imatinib and 2 other kinase inhibitors.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Quinine

Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	Treatment or prevention of leg cramps
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	Doses for duration greater than 10 days will not be approved
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Qulipta

Products Affected

- **QULIPTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of Episodic Migraine, defined as 4-14 migraine days per month OR Chronic Migraine, defined as 15 or more headaches per month, of which 8 or more are migraine days. The following criteria will apply (1-2). 1) Documentation of average monthly migraine days. 2) Attestation that headaches are not caused by medication rebound (e.g. not taking triptans exceeding more than 18 doses per month) or lifestyle factors (e.g. sleep patterns, caffeine use).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine frequency
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Radicava ORS

Products Affected

- **RADICAVA ORS STARTER KIT SUSP**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of amyotrophic lateral sclerosis (ALS) -AND- Therapeutic failure, intolerance, or contraindication to riluzole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of experiencing stability or improvement of symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ravicti

Products Affected

- *glycerol phenylbutyrate*

PA Criteria	Criteria Details
Exclusion Criteria	Urea cycle disorders due to N-acetylglutamate synthetase deficiency, Treatment of acute hyperammonemia in urea cycle disorders
Required Medical Information	Documentation of chronic management of a urea cycle disorders (UCDs)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Relistor

Products Affected

- **RELISTOR ORAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-2): 1) chronic non-cancer pain, or 2) chronic pain related to prior cancer or its treatment who do not require frequent (e.g. weekly) opioid dosage escalation -AND- documentation of opioid medication use -AND- trial and failure, contraindication, or intolerance to 2 of the following (3-5): 3) Laxatives, 4) lubiprostone, 5) Movantik
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Relistor SC

Products Affected

- **RELISTOR SUBCUTANEOUS SOLUTION**
- **RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-3): 1) chronic non-cancer pain, 2) advanced illness or active cancer in palliative care or 3) chronic pain related to prior cancer or its treatments who do not require frequent (e.g., weekly) opioid dosage escalation -AND- documentation of opioid medication use -AND- trial and failure, contraindication, or intolerance to 2 of the following (4 to 6): 4) Laxatives 5) lubiprostone 6) Movantik.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Repatha

Products Affected

- **REPATHA SURECLICK**
- **REPATHA SYRINGE**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with another PCSK9 inhibitor or siRNA directed to PCSK9.
Required Medical Information	1.HoFH supported by 1 of the following (a-d): a) bi-allelic causal mutations of FH by genetic testing, b) LDL-C over 560mg/dL, c) LDL-C over 400mg/dL w/aortic valve disease or xanthomata at less than 20 yrs old, d) LDL-C over 400mg/dL and 1 or both parents w/clinically diagnosed FH, positive genetic testing for a known LDL-C-raising gene (e.g., LDLR, ApoB, PCSK9) defect, or autosomal-recessive FH AND current LDL-C over 115mg/dL (if 17yrs or younger) or over 70mg/dL (18yrs or older) or over 55mg/dL w/ASCVD or major ASCVD risk factors (18yrs or older). 2.HeFH supported by 1 of the following (a-d): a) causal mutation of FH by genetic testing, b) untreated LDL-C at least 190mg/dL or untreated LDL-C at least 160mg/dL before 20 yrs old, c) physical signs of FH (e.g. xanthomas, xanthelasma), d) diagnosis based on WHO criteria/Dutch Lipid Clinical Network criteria w/score over 8 points, definite on Simon Broome register, or definite on the Make Early Diagnosis to Prevent Early Deaths tool AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C, or if very high risk, LDL-C at least 55mg/dL or non-HDL-C at least 85mg/dL. 3.Hypercholesterolemia w/ASCVD or attestation of major CV risk AND LDL-C at least 55mg/dL or non-HDL-C at least 85mg/dL or Lp(a) at least 125nmol/L or, if ASCVD, less than 50% decrease in baseline LDL-C. 4.Primary Hyperlipidemia/Hypercholesterolemia (w/o ASCVD, HeFH, or HoFH) AND LDL-C at least 70mg/dL or non-HDL-C at least 100mg/dL or less than 50% decrease in baseline LDL-C.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization

PA Criteria	Criteria Details
Other Criteria	For reauthorization associated with Major Cardiovascular Risk, attestation of need for continued Repatha therapy is required. For reauthorization associated with all other diagnoses, documentation showing an LDL-C reduction on Repatha therapy from baseline must be provided. Therapeutic failure to a max tolerated statin or documented statin intolerance is required. Statin intolerance defined as follows: statin related rhabdomyolysis or skeletal muscle symptoms while receiving at least 2 separate trials of different statins which resolved upon discontinuation of statin or attestation of one of the following during any course of statin therapy: 1. CK increase to 10x upper limit of normal 2. LFTs increase to 3x upper limit of normal 3. Hospitalization due to severe statin-related AEs such as rhabdomyolysis.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Retevmo

Products Affected

- **RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of locally advanced or metastatic non-small cell lung cancer -AND- disease is classified as RET gene fusion as detected by an FDA approved test. Documentation of advanced or metastatic medullary thyroid cancer -AND- disease is classified as RET mutation as detected by an FDA approved test. Documentation of advanced or metastatic thyroid cancer -AND- all of the following (1-2): 1) disease is classified as RET gene fusion as detected by an FDA approved test 2) if radioactive iodine is appropriate for the member, the member is radioactive iodine-refractory. Documentation of locally advanced or metastatic solid tumor(s) -AND- disease harbors a RET gene fusion, as detected by an FDA-approved test -AND- one of the following (1-2): 1) the member has no satisfactory alternative treatments 2) the member's tumors have progressed following prior systemic treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Revlimid

Products Affected

- *lenalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of chronic lymphocytic leukemia outside of a controlled clinical trial
Required Medical Information	Diagnosis of multiple myeloma in combination with dexamethasone -OR- diagnosis of multiple myeloma, as maintenance following autologous hematopoietic stem cell transplant (auto-HSCT) -OR- diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities -OR- diagnosis of mantle cell lymphoma (MCL) in which disease has relapsed or progressed after two prior therapies, one of which included bortezomib -OR- diagnosis of follicular lymphoma in combination with a rituximab product after previous treatment -OR- diagnosis of marginal zone lymphoma in combination with a rituximab product after previous treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Revuforj

Products Affected

- **REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For relapsed or refractory acute leukemia, member has a lysine methyltransferase 2A gene (KMT2A) translocation, as determined by an FDA-authorized test. For relapsed or refractory acute myeloid leukemia, the member has a susceptible NPM1 mutation -AND- the member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rexulti

Products Affected

- **REXULTI ORAL TABLET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI). For schizophrenia, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rezdiffra

Products Affected

- **REZDIFFRA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of noncirrhotic nonalcoholic steatohepatitis (NASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in conjunction with diet and exercise. Liver biopsy (within the past 3 years) or noninvasive tests (NITs) (performed within the past 6 months) indicating stage F2 or F3 fibrosis -AND- attestation of 1 cardiometabolic risk factor (e.g. obesity, hyperglycemia, hypertriglyceridemia, hyperlipidemia, hypertension) -AND- attestation that member does not have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma -AND- attestation that member's alcohol consumption is not defined as excess alcohol use (greater than 20g/day in females or greater than 30g/day in males)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of continued use in conjunction with diet and exercise -AND- member has experienced improvement or stabilization of fibrosis as demonstrated by NIT -AND- member has not progressed to stage F4 fibrosis -AND- attestation that member does not have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma -AND- attestation that member's alcohol consumption is not defined as excess alcohol use (greater than 20g/day in females or greater than 30g/day in males)
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rezlidhia

Products Affected

- **REZLIDHIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA approved test
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rinvoq

Products Affected

- **RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate) or all non-biologic DMARDs are contraindicated. For moderate to severe refractory atopic dermatitis whose disease is not adequately controlled with other systemic drug products, documentation of one of the following (1 or 2): 1) trial & failure, or intolerance to at least one topical corticosteroid -OR- topical calcineurin inhibitor (e.g., tacrolimus, pimecrolimus) 2) The member has severe atopic dermatitis and is incapable of applying topical therapies due to the extent of body surface area involvement or topical therapies are contraindicated due to severely damaged skin. For ankylosing spondylitis, inadequate response or intolerance to one nonsteroidal anti-inflammatory drug (NSAID). For Non-radiographic Axial Spondyloarthritis, trial & failure or intolerance to two nonsteroidal anti-inflammatory drugs (NSAIDs) or contraindication to all. For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Plan Year

PA Criteria	Criteria Details
Other Criteria	For diagnoses in which tumor necrosis factor (TNF) blockers are also indicated (e.g., Rheumatoid Arthritis, Psoriatic Arthritis), the member has experienced therapeutic failure or intolerance to at least 1 TNF blocker. For ulcerative colitis or Crohns disease, the member has experienced therapeutic failure or intolerance to at least 1 systemic therapy if TNF blockers are not advisable.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rinvoq LQ

Products Affected

- **RINVOQ LQ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or all non-biologic DMARDs are contraindicated or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	For PsA, deny if 18 years of age or older
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rivfloza

Products Affected

- **RIVFLOZA SUBCUTANEOUS SOLUTION**
- **RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-4): 1) diagnosis of primary hyperoxaluria type 1 (PH1) confirmed by one of the following (a or b): a) Genetic testing (i.e., AGXT gene mutation), b) Liver biopsy (i.e., absence or significantly reduced AGT activity), 2) relatively preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2), 3) at least two elevated urinary oxalate levels greater than 1.5 times the upper reference limit, and 4) The member has not received a liver transplant.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, continued preserved kidney function (e.g., eGFR greater than or equal to 30 mL/min/1.73 m2) -AND- reduction in urinary oxalate levels from baseline -AND- has not received a liver transplant.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Romvimza

Products Affected

- **ROMVIMZA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor -AND- attestation that surgical resection may cause one of the following (1-2): 1) worsening functional limitation, 2) severe morbidity
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rozlytrek

Products Affected

- **ROZLYTREK ORAL CAPSULE 100 MG, 200 MG**
- **ROZLYTREK ORAL PELLETS IN PACKET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer, the tumor status is ROS1-positive. For solid tumors with NTRK gene fusion without a known acquired resistance mutation, the tumors are metastatic or surgical resection is likely to result in severe morbidity - AND- There are no satisfactory alternative treatments or the tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Rubraca

Products Affected

- **RUBRACA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, all of the following (1-2): 1) Disease harbors a deleterious BRCA mutation (germline or somatic) 2) member is in complete or partial response to platinum-based chemotherapy. For metastatic castration-resistant prostate cancer, all of the following (1-3): 1) disease harbors a deleterious BRCA mutation (germline and/or somatic) 2) member has been treated with androgen receptor-directed therapy 3) member is concurrently receiving a gonadotropin-releasing hormone (GnRH) analog or member has had a bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Rydapt

Products Affected

- **RYDAPT**

PA Criteria	Criteria Details
Exclusion Criteria	Use as single agent induction therapy for AML
Required Medical Information	Documentation of diagnosis. For a new diagnosis of acute myeloid leukemia, member is using in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation chemotherapy regimens and member is FLT3 mutation positive as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Sabril

Products Affected

- *vigabatrin*
- **VIGADRONE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of refractory complex partial seizures -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance to at least two alternative treatments (e.g. carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, tiagabine) -OR- documentation of use as monotherapy in treatment of infantile spasms
Age Restrictions	Deny if less than 2 years of age in treatment of refractory complex partial seizures -OR- if less than 1 month old and greater than 2 years of age in treatment of infantile spasms
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Samsca

Products Affected

- *tolvaptan*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic hypervolemic or euvolemic hyponatremia evidenced by (1. or 2.): 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	Doses must be initiated in the hospital setting to closely monitor serum sodium. Treatment should be limited to 30 days to minimize risk of liver injury. For reauthorization, treatment is for a new episode of a clinically significant euvolemic or hypervolemic hyponatremia -AND- on of the following (1. or 2.) 1.) Serum Na less than 125 mEq/L -OR- 2.) Serum NA less than 135mEq/L with symptoms (e.g. nausea, malaise, lethargy, headache, seizures)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Saphris

Products Affected

- *asenapine maleate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis - AND - trial and failure of one of the following: olanzapine, quetiapine, or risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Savella

Products Affected

- *milnacipran oral tablet*
- **SAVELLA ORAL TABLETS,DOSE PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation to support a diagnosis of fibromyalgia and trial/failure or intolerance to both duloxetine and pregabalin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Scemblix

Products Affected

- **SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- one of the following (1-3): 1) the member is newly-diagnosed, 2) member has been previously treated for Ph+ CML in chronic phase, 3) disease is positive for the T3151 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Secuado

Products Affected

- **SECUADO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- intolerance to generic asenapine sublingual tablets -AND- trial/failure or intolerance to 1 of the following or all are contraindicated (1-3): 1) olanzapine, 2) quetiapine, 3) risperidone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Siliq

Products Affected

- **SILIQ**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For psoriasis induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For psoriasis, patients must have therapeutic failure or intolerance to 2 preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Cosentyx, Otezla/Otezla XR, and Skyrizi SC. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

Sirturo

Products Affected

- **SIRTURO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Criteria will be applied consistent with the current ATS/CDC/ERS/IDSA Clinical Practice Guideline for the Treatment of Drug-Susceptible and Drug-Resistant Tuberculosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of disease improvement -AND- member requires additional antimicrobial therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Skyclarys

Products Affected

- **SKYCLARYS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Friedreichs ataxia confirmed by genetic testing (i.e., FXN gene mutation).
Age Restrictions	Deny if less than 16 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Skyrizi

Products Affected

- **SKYRIZI SUBCUTANEOUS PEN INJECTOR**
- **SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML, 55 MG/0.37 ML**
- **SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohns disease, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC. For moderate to severe ulcerative colitis, attestation of receiving or currently undergoing IV administration of Skyrizi within 3 months of initiating therapy with Skyrizi SC.
Age Restrictions	Deny if less than 18 years of age for Crohn's disease and ulcerative colitis. Deny if less than 6 years of age for plaque psoriasis and psoriatic arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For induction therapy, doses above plan quantity limit will be approved when aligned with recommended induction therapy dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sohonos

Products Affected

- **SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 5 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	Deny if female and less than 8 years of age -OR- if male and less than 10 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in volume of new heterotopic ossification from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Solaraze

Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- trial and failure, intolerance, or contraindication to one of the following (1 or 2): 1) topical fluorouracil solution or fluorouracil 5% cream 2) topical imiquimod 5% cream
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of 30 day washout period since optimal therapeutic effect may not be evident until 30 days following cessation of therapy AND attestation of previous response to diclofenac sodium 3% topical gel therapy
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Somavert

Products Affected

- **SOMAVERT**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For acromegaly, high pretreatment insulin-like growth factor-1 (IGF-1) based on laboratory reference range -AND- inadequate or partial response to surgery or radiotherapy or not a candidate for surgery or radiotherapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization of acromegaly, decreased or normalized IGF-1 from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Sprycel

Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adults with Ph+ chronic myeloid leukemia, the member is newly diagnosed in the chronic phase -OR- the member is in chronic, accelerated, or myeloid or lymphoid blast phase and has resistance or intolerance to prior therapy including imatinib. For adults with Ph+ acute lymphocytic leukemia, member has had resistance or intolerance to prior therapy. For pediatric patients with Ph+ CML, the member is in the chronic phase. For pediatric patients with Ph+ acute lymphoblastic leukemia, the member is newly diagnosed and will be using in combination with chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Steroidogenesis Inhibitors

Products Affected

- **ISTURISA ORAL TABLET 1 MG, 5 MG**
- **SIGNIFOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- For Signifor, patient is not a candidate for pituitary surgery or pituitary surgery has not been curative. For Isturisa, patient is not a candidate for surgery or surgery has not been curative.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of decrease in urinary free cortisol levels from baseline
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Stivarga

Products Affected

- **STIVARGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic colorectal cancer and trial of a fluoropyrimidine-, oxaliplatin-, and irinotecan-containing chemotherapy, AND an anti-VEGF therapy AND if RAS wild-type, an anti-EGFR therapy -OR- documentation of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with both imatinib and sunitinib -OR- documentation of hepatocellular cancer AND previous treatment with sorafenib
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sunosi

Products Affected

- **SUNOSI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) -AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B) A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL, or B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range, 2) Multiple sleep latency test (MSLT) documenting MSL less than or equal to 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP), or 3) MSLT documenting MSL less than or equal to 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. Diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) documented by objective polysomnography or home sleep apnea study as established in accordance with ICSD or DSM V criteria acceptable for all indications.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For narcolepsy and OSAHS, documentation of trial and failure, contraindication or intolerance to modafinil and armodafinil. For reauthorization, provider attestation of improvement in daytime sleepiness is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sutent

Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For gastrointestinal stromal tumor (GIST), the member has experienced therapeutic failure, intolerance, or contraindication to imatinib. For a high risk of recurrent renal cell carcinoma, member has had a nephrectomy and sunitinib is to be used as adjuvant treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Sympazan

Products Affected

- **SYMPAZAN**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures due to Lennox-Gastaut Syndrome -AND- documentation of adjunctive therapy -AND- therapeutic failure or intolerance of a previous antiepileptic therapy -AND- unable to tolerate generic clobazam
Age Restrictions	Deny if less than 2 years old
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Symproic

Products Affected

- **SYMPROIC**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of opioid induced constipation (OIC) due to any of the following (1-2): 1) chronic non-cancer pain, or 2) chronic pain related to prior cancer or its treatments who do not require frequent (e.g., weekly) opioid dosage escalation -AND- documentation of opioid medication use -AND- trial and failure, contraindication, or intolerance to at least 2 of the following (3-5): 3) Laxatives, 4) lubiprostone, 5) Movantik
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Synarel

Products Affected

- **SYNAREL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For central precocious puberty (CPP), advancement of bone age is beyond chronological age -AND- Basal luteinizing hormone (LH) level greater than 0.2-0.3IU/L or leuprolide-stimulating LH level greater than 3.3-5 IU/L. For female with endometriosis, attestation of not pregnant if of childbearing age -AND- Therapeutic failure, contraindication or intolerance to 2 of the following standard of care treatments: NSAIDs, combination hormonal contraceptive, progestin (i.e. medroxyprogesterone injection), GnRH agonist (i.e. Leuprolide) or danazol
Age Restrictions	Deny if greater than 8 years of age for females or greater than 9 years of age for males unless there is medical necessity for treatment of central precocious puberty. Deny if less than 18 years of age for endometriosis.
Prescriber Restrictions	
Coverage Duration	Endometriosis: 6 months, CPP: 6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization for CPP, attestation of pubertal development slowing from baseline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tabrecta

Products Affected

- **TABRECTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with MET exon 14 skipping mutation as detected by an FDA approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tafinlar

Products Affected

- **TAFINLAR ORAL CAPSULE**
- **TAFINLAR ORAL TABLET FOR SUSPENSION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For dabrafenib monotherapy, documentation of unresectable or metastatic melanoma -AND- documentation of a BRAF V600E mutation. For use in combination with trametinib, documentation of unresectable or metastatic melanoma or melanoma with lymph node(s) involvement following complete resection and member is using trametinib and dabrafenib as adjuvant therapy -AND- member has a BRAF V600E or V600K mutation. For use in combination with trametinib, documentation of metastatic non-small cell lung cancer, locally advanced or metastatic anaplastic thyroid cancer, or low-grade glioma -AND- member has a BRAF V600E mutation, as detected by an FDA-approved test when FDA indicated. For use in combination with trametinib, documentation of unresectable or metastatic solid tumors -AND- all of the following (1-3): 1) BRAF V600E mutation 2) disease has progressed following prior treatment 3) member has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For Tafinlar tablets for oral suspension, attestation of inability to swallow Tafinlar (dabrafenib) capsules is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Tagrisso

Products Affected

- **TAGRISSO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of non-small cell lung cancer (NSCLC) and one of the following (1-5): 1) Adjuvant therapy after tumor resection -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test 2) Locally advanced disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used in combination with pemetrexed and platinum-based chemotherapy, 3) Metastatic disease -AND- using as first-line therapy -AND- disease harbors EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test -AND- used with or without combination therapy of pemetrexed and platinum-based chemotherapy, 4) Metastatic disease -AND- disease harbors EGFR T790M mutations, as detected by an FDA-approved test -AND- has progressed on or after EGFR TKI therapy, 5) Locally advanced, unresectable (stage III) disease -AND- disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy -AND- disease harbors EGFR exon 19 deletions or EGFR exon 21 L858R mutations, as detected by an FDA-approved test.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Talzenna

Products Affected

- **TALZENNA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of deleterious or suspected deleterious gBRCAm, HER2-negative locally advanced or metastatic breast cancer as a single agent -OR- Documentation of HRR gene-mutated metastatic castration-resistant prostate cancer in combination with enzalutamide
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Targretin

Products Affected

- *bexarotene oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations of cutaneous T-cell lymphoma in patients who are refractory to at least one prior systemic therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Targretin Gel

Products Affected

- *bexarotene topical*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous lesions associated with one (1) of the following stages of cutaneous T-cell lymphoma (CTCL) (1. or 2.): 1) Stage IA or 2) Stage IB - AND- the member has experience therapeutic failure, contraindication, or intolerance to at least one other therapy for CTCL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tascenso ODT

Products Affected

- **TASCENSO ODT**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as teriflunomide, interferons, Copaxone, Tysabri
Required Medical Information	Documentation of a relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Inability to swallow capsules
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tasigna

Products Affected

- *nilotinib hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For adult patients with Ph+ chronic myeloid leukemia (CML), member's CML is in the chronic or accelerated phase and the member is no longer responding to or is intolerant to imatinib -OR- member is newly diagnosed in the chronic phase. For pediatric patients, one of the following (1-2): 1) member has chronic phase or accelerated phase Ph+ CML and is is resistant or intolerant to prior tyrosine kinase inhibitor therapy 2) member is newly diagnosed with Ph+ CML in the chronic phase.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tasimelteon

Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented diagnosis of Non-24 Sleep-Wake disorder in patient that is totally blind -AND- evidenced by all the following (1 through 4): 1) history of insomnia, excessive daytime sleepiness, or both alternating with asymptomatic episodes 2) symptoms persistent for at least 3 months 3) daily sleep logs for at least 1 month demonstrating a sleep/wake pattern that delays each day 4) sleep disturbances are not better explained by another current disorder or medication/substance use
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of increased total nighttime sleep or decreased daytime nap duration for Non-24 Sleep-Wake disorder
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tavalisse

Products Affected

- **TAVALISSE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For diagnosis of ITP, the following criteria apply (1 and 2): 1) insufficient response to a corticosteroid, immunoglobulin, or splenectomy. 2) One of the following (A or B): A) Platelet count less than or equal to 50 x 10*9/L and has significant mucous member bleeding or at least one risk factor for bleeding (e.g. hypertension, peptic ulcer disease). B) Platelet count of less than or equal to 30 x 10*9/L.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tazorac

Products Affected

- *tazarotene topical cream*
- *tazarotene topical gel 0.1 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 of the following (A or B). A) Documentation of plaque psoriasis -AND- trial and failure or intolerance to at least one topical corticosteroid (e.g. fluocinonide, mometasone, triamcinolone, betamethasone). B) Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two topical acne medications (e.g. adapalene, clindamycin, sulfacetamide, erythromycin) one of which must be generic topical tretinoin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tecfidera

Products Affected

- *dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tepmetko

Products Affected

- **TEPMETKO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of metastatic non-small cell lung cancer with a MET exon 14 skipping alteration
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Testosterone (androgens)

Products Affected

- **AVEED**
- **DEPO-TESTOSTERONE INTRAMUSCULAR OIL 100 MG/ML**
- *testosterone cypionate*
- *testosterone enanthate*
- *testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %)*
- *testosterone transdermal solution in metered*

pump w/app

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of primary or secondary hypogonadism in males -OR- documentation of chronic steroid treatment in men. In all previously noted indications, members must also have documented low total testosterone level below the normal range for the laboratory -OR- a total testosterone level near the lower limit of the normal range with a low free testosterone level which is less than normal based upon the laboratory reference range -OR- the member is not producing any testosterone. Additional approvable indications include female patients with metastatic breast cancer (testosterone enanthate only), primary or secondary hypogonadism in males with testicular failure due to double orchidectomy, and delayed puberty in males (testosterone enanthate only).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications, Some Medically-accepted Indications.
Off Label Uses	Gender Dysphoria, Gender Identity Disorder
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tezspire

Products Affected

- **TEZSPIRE SUBCUTANEOUS PEN INJECTOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of severe asthma AND all of the following (1-3): 1) one of the following (a. or b.): a) history of at least two asthma exacerbations requiring oral or injectable corticosteroid treatment in the past 12 months, b) history of at least one asthma exacerbation requiring hospitalization in the past 12 months, 2) inadequate symptom control despite regular treatment with medium or high dose inhaled corticosteroid (ICS) and at least one additional asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline) with or without oral corticosteroids (OCS), 3) member will continue treatment with medium or high dose ICS and at least one additional asthma controller medication, with or without OCS. Diagnosis of chronic rhinosinusitis with nasal polyposis (CRwNP) and therapeutic failure, contraindication, or intolerance to 1 generic intra-nasal corticosteroid.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For asthma, therapeutic failure, intolerance, or contraindication to 1 of the following: Cinqair, Dupixent, Fasenra, Nucala, or Xolair. For CRwNP, therapeutic failure, intolerance, or contraindication to 1 of the following: Dupixent, Nucala, or Xolair. For asthma reauthorization, attestation of 1 of the following (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of severe asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CRwNP reauthorization, 1 of the following (5-6): 5) decrease in nasal polyp score, or 6) reduction in nasal congestion/obstruction severity score.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Thalomid

Products Affected

- **THALOMID ORAL CAPSULE 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	Use as monotherapy for ENL treatment in the presence of moderate to severe neuritis
Required Medical Information	Documentation of multiple myeloma in combination with dexamethasone -OR- documentation for use in the treatment of cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) -OR- documentation of therapy for prevention and suppression of the cutaneous manifestations of ENL recurrence
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Thrombopoiesis Stimulating Agents

Products Affected

- *eltrombopag olamine oral powder in packet 12.5 mg, 25 mg*
- *eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of persistent or chronic immune idiopathic thrombocytopenia purpura and trial and failure of corticosteroid or immunoglobulin therapy or splenectomy -OR- documentation of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy -OR- severe aplastic anemia who have had an insufficient response to immunosuppressive therapy -OR- documentation of first line treatment for severe aplastic anemia and used in combination with at least two immunosuppressive therapies.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Platelet count to be provided
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tibsovo

Products Affected

- **TIBSOVO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is IDH1 mutation positive as detected by an FDA-approved test. For newly-diagnosed acute myeloid leukemia, member is using as monotherapy or in combination with azacitidine -AND- member meets one of the following (1-5): 1) age is greater than or equal to 75 years of age 2) severe cardiac or pulmonary comorbidity 3) reduced renal function 4) hepatic impairment 5) or prescriber attestation that member is not a candidate for intensive induction therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tocilizumab

Products Affected

- **ACTEMRA ACTPEN**
- **ACTEMRA SUBCUTANEOUS**

PA Criteria	Criteria Details
Exclusion Criteria	Part A covered for Covid-19 in hospitalized patients
Required Medical Information	Documentation of diagnosis. For rheumatoid arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide). For polyarticular juvenile idiopathic arthritis, patients must have an inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For systemic sclerosis-associated interstitial lung disease (SSc-ILD), patients must have therapeutic failure or intolerance to one immunosuppressant (e.g., mycophenolate mofetil, corticosteroids, cyclophosphamide). Documentation of systemic juvenile idiopathic arthritis.
Age Restrictions	Deny if less than 18 years of age for systemic sclerosis-associated interstitial lung disease (SSc-ILD), Rheumatoid Arthritis, and Giant Cell Arteritis or less than 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For rheumatoid arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq and Xeljanz/Xeljanz XR. For polyarticular juvenile idiopathic arthritis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, Rinvoq/Rinvoq LQ, and Xeljanz/Xeljanz solution. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Topical Lidocaine

Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical ointment*
- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tretinoin

Products Affected

- *adapalene topical gel 0.3 %*
- *adapalene topical swab*
- *adapalene-benzoyl peroxide*
- *clindamycin-tretinoin*
- *tretinoin*

PA Criteria	Criteria Details
Exclusion Criteria	Cosmetic use
Required Medical Information	Documentation of acne vulgaris -AND- trial and failure or intolerance of at least two generic topical non-retinoid acne medications (e.g. clindamycin-benzoyl peroxide)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Trikafta

Products Affected

- **TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL**
- **TRIKAFTA ORAL TABLETS, SEQUENTIAL**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cystic fibrosis (CF) in patients who have at least one F508del mutation or another mutation in the cystic fibrosis transmembrane conductance regulator (CFTR gene) that is responsive to elexacaftor/tezacaftor/ivacaftor based on in vitro assay (e.g. E56K, R117C, A455E)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, documentation supporting improvement or stabilization of FEV1 compared to baseline FEV1 -or- increase in body mass index -or- decreased pulmonary exacerbations -or- improved quality of life as demonstrated by CF Questionnaire is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Trintellix

Products Affected

- **TRINTELLIX**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Truqap

Products Affected

- **TRUQAP**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For locally advanced or metastatic breast cancer, all of the following (1-3): 1) HR mutation status, HER2 mutation status, and PIK3CA/AKT1/PTEN status as detected by an FDA-authorized test 2) concomitant therapy with fulvestrant 3) disease progression on at least one endocrine-based regimen in the metastatic setting -OR- recurrence on or within 12 months of completing adjuvant therapy. For metastatic prostate cancer, all of of the following (5-8): 5) disease is androgen pathway modulation-naive or -sensitive 6) disease is PTEN-deficient, as detected by an FDA-authorized test, 7) will be used in combination with abiraterone and prednisone, 8) concurrently receiving a gonadotropin-releasing hormone analog or has had bilateral orchiectomy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tukysa

Products Affected

- **TUKYSA ORAL TABLET 150 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced unresectable HER2-positive breast cancer or metastatic HER2-positive breast cancer, member will be using in combination with trastuzumab and capecitabine -AND- member has received one or more prior anti-HER2 based regimens in the metastatic setting. For RAS wild-type HER2-positive unresectable or metastatic colorectal cancer, member will be using in combination with trastuzumab -AND- member has experienced disease progression following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Turalio

Products Affected

- **TURALIO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of symptomatic tenosynovial giant cell tumor associated with severe morbidity and functional limitations -AND- patient is not amenable to improvement with surgery or not a candidate for surgery
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Tykerb

Products Affected

- *lapatinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced HER2-positive, or metastatic HER2-positive breast cancer, the member has received prior therapy with an anthracycline, a taxane, and trastuzumab -AND- will be using in combination with capecitabine. For HR+, metastatic breast cancer, the member is post-menopausal -AND- the member's cancer over expresses the HER2 receptor -AND- the member will be using lapatinib in combination with letrozole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Tymlos

Products Affected

- **TYMLOS**

PA Criteria	Criteria Details
Exclusion Criteria	Treatment duration greater than 24 months.
Required Medical Information	Documentation of diagnosis -AND- at high risk for fracture, meeting one of the following (1. thru 3.) 1) History of previous hip or vertebral fracture. 2) T-score less than or equal to -2.5. 3) T-score between -1.0 and -2.5 -AND- meets FRAX calculation (A. or B.) A) 10-year risk of major osteoporotic fracture is greater than or equal to 20 percent or B) 10-year risk of hip fracture is greater than or equal to 3 percent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 months
Other Criteria	Documentation of trial/failure or intolerance to at least one oral bisphosphonate or all are contraindicated. Additional documentation of trial/failure, intolerance or contraindication to preferred parathyroid hormone analog teriparatide. Coverage of human parathyroid hormone related peptide analogs beyond 24 months will not be approved. A cumulative lifetime approval of Tymlos will be limited to a coverage duration of 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ubrelvy

Products Affected

- **UBRELVY ORAL TABLET 100 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of acute treatment of migraine with or without aura -AND- trial and failure of one generic triptan or all triptans are contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	Patients must have therapeutic failure, intolerance, or contraindication to Nurtec ODT. For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Uptravi

Products Affected

- **UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG** **PACK**
- **UPTRAVI ORAL TABLETS,DOSE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Trial and failure of Opsumit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ustekinumab

Products Affected

- **STELARA SUBCUTANEOUS SOLUTION**
- **STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML**
- *ustekinumab subcutaneous solution*
- *ustekinumab subcutaneous syringe 45 mg/0.5 ml, 90 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- documentation of member weight and prescribed dose. For moderate to severe plaque psoriasis, inadequate response or intolerance to one systemic therapy (e.g. methotrexate) -OR- inadequate response to phototherapy -OR- contraindication to phototherapy and systemic therapy. For Crohn's Disease or Ulcerative Colitis, attestation of receiving or currently undergoing a single induction dose of IV ustekinumab within 2 months of initiating therapy with ustekinumab SC.
Age Restrictions	Deny if less than 18 years of age for Ulcerative Colitis. Deny if less than 6 years of age for Plaque Psoriasis and Psoriatic Arthritis. Deny if less then 2 years of age for Crohn's Disease.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	For ustekinumab products except Steqeyma and Yesintek, intolerance to both preferred ustekinumab products is required: Steqeyma with NDC starting with 72606 and Yesintek with NDC starting with 83257. Must follow recommended dosing guidelines based upon weight. Subcutaneous induction therapy, doses above plan quantity limit will be approved aligned with recommended induction therapy dosing regimen. For pediatric Crohn's disease, doses above plan quantity limit will be approved to align with recommended dosing regimen.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Valchlor

Products Affected

- **VALCHLOR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of Stage IA or IB mycosis fungoides-type cutaneous T-cell lymphoma in patients who have received at least one prior skin-directed therapy (e.g. topical corticosteroids, topical chemotherapy, local radiation and topical retinoids).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Valtoco

Products Affected

- **VALTOCO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizure clusters or acute repetitive seizures -AND- the member is currently receiving antiepileptic maintenance therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vanflyta

Products Affected

- **VANFLYTA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is FLT3-ITD-positive as detected by an FDA-approved test -AND- one of the following (1-3): 1) member is receiving induction therapy and is using Vanflyta in combination with standard cytarabine and anthracycline induction therapy 2) member is receiving consolidation therapy and is using Vanflyta in combination with standard cytarabine consolidation therapy 3) member is receiving maintenance therapy and is using Vanflyta as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Venclexta

Products Affected

- **VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG**
- **VENCLEXTA STARTING PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For newly-diagnosed AML, member is using in combination with azacitidine, decitabine, or cytarabine -AND- age greater than or equal to 75 years or presence of at least one comorbidity that precludes use of intensive induction chemotherapy (i.e. severe cardiac or pulmonary comorbidity, reduced renal function, hepatic impairment, or physician attestation) is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Verquvo

Products Affected

- **VERQUVO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of heart failure (NYHA Class II to IV) -AND- Left ventricular ejection fraction less than 45% -AND- Hospitalization for heart failure or received outpatient IV diuretics for heart failure -AND- Used in combination with a angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker or Entresto -AND- Used in combination with bisoprolol, carvedilol IR/ER or metoprolol succinate ER.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Verzenio

Products Affected

- **VERZENIO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- disease is classified as HR-positive, HER2-negative. For early breast cancer that is at high risk of recurrence and is node-positive, all of the following (1-2): 1) used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) 2) used as adjuvant treatment. For advanced or metastatic breast cancer, used as initial endocrine-based therapy and used in combination with an aromatase inhibitor -OR- used after documented disease progression following endocrine therapy and used in combination with fulvestrant -OR- used after documented disease progression and used following endocrine therapy and prior chemotherapy in the metastatic setting and will be used as monotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Viberzi

Products Affected

- **VIBERZI**

PA Criteria	Criteria Details
Exclusion Criteria	Severe (Child-Pugh C) hepatic impairment
Required Medical Information	Documentation of diarrhea predominant, irritable bowel syndrome (IBS-D) -AND- trial/failure or intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Viibryd

Products Affected

- *vilazodone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of major depressive disorder -AND- Therapeutic failure, intolerance or contraindication to one generic antidepressant (e.g. SSRI, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vijoice

Products Affected

- **VIJOICE ORAL GRANULES IN MG PACKET**
- **VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of severe manifestations of PIK3CA Related Overgrowth Spectrum (PROS)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vitrakvi

Products Affected

- **VITRAKVI ORAL CAPSULE 100 MG, 25 MG**
- **VITRAKVI ORAL SOLUTION**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation -AND- Tumors are metastatic or surgical resection is likely to result in severe morbidity -AND- There are no satisfactory alternative treatments or tumors have progressed following treatment.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vivjoa

Products Affected

- **VIVJOA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of recurrent vulvovaginal candidiasis defined as at least 3 episodes of vulvovaginal candidiasis in less than one year -AND Documentation the member is NOT of reproductive potential defined as postmenopausal or another reason for permanent infertility (e.g., tubal ligation, hysterectomy, salpingo-oophorectomy) -AND- the member has experienced therapeutic failure, contraindication, or intolerance to a six-month maintenance course of oral fluconazole.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	14 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vonjo

Products Affected

- **VONJO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of intermediate or high-risk myelofibrosis -AND- attestation of a platelet count of less than 50 x 10*9/L.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Voranigo

Products Affected

- **VORANIGO ORAL TABLET 10 MG, 40 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- all of the following (1-3): 1) grade 2 astrocytoma or grade 2 oligodendroglioma, 2) disease harbors a susceptible isocitrate dehydrogenase (IDH)-1 or IDH-2 mutation, as detected by an FDA-approved test, 3) will be used following surgery including biopsy, sub-total resection, or gross total resection.
Age Restrictions	Deny if less than 12 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Voriconazole

Products Affected

- *voriconazole intravenous recon soln*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary cannot take oral voriconazole
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For reauthorization, attestation of continued indicators of active disease (e.g. histopathology, positive cultures) is required
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Vosevi

Products Affected

- **VOSEVI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guidance - AND- the member has a contraindication to or is otherwise not a candidate for one of the following regimens recommended by the AASLD/IDSA guidelines containing the following agents: sofosbuvir/velpatasvir (i.e. Epclusa authorized generic), Mavyret.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Criteria/duration applied consistent with current AASLD-IDSA guidance
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Votrient

Products Affected

- *pazopanib oral tablet 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Documentation of adipocytic soft tissue sarcoma or gastrointestinal stromal tumor
Required Medical Information	Documentation of diagnosis. For advanced soft-tissue sarcoma, trial/failure of at least one prior chemotherapy regimen.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vraylar

Products Affected

- **VRAYLAR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For a diagnosis of schizophrenia and bipolar I disorder, therapeutic failure, intolerance, or contraindication to one other generic atypical antipsychotic (e.g. quetiapine). For major depressive disorder, documentation of adjunctive therapy and therapeutic failure, contraindication or intolerance to one other generic antidepressant in addition to the antidepressant currently being used for the treatment of MDD (e.g. SSRI, SNRI, NDRIs, TCA, MAOI).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Vumerity

Products Affected

- **VUMERITY**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other disease modifying agents such as interferons, Copaxone, Tysabri, Aubagio, Gilenya
Required Medical Information	Documentation of relapsing form of multiple sclerosis (e.g. relapsing-remitting, clinically isolated syndrome, or active secondary progressive disease) -AND- Therapeutic failure or intolerance to generic dimethyl fumarate
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	5 years
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Welireg

Products Affected

- **WELIREG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For von Hippel Lindau (VHL) syndrome, one of the following diagnoses not requiring immediate surgery (1, 2, or 3): 1) Renal cell carcinoma (RCC) 2) CNS hemangioblastoma 3) Pancreatic neuroendocrine tumor. For advanced RCC with a clear cell component, all of the following are required (4-6): 4) prior treatment with a vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI), 5) prior treatment with a programmed death receptor-1 (PD-1) or a programmed death-ligand (PD-L1) inhibitor, 6) will be used as monotherapy. For RCC with a clear cell component, all of the following are required (7-8): 7) at intermediate-high or high risk of recurrence following nephrectomy -OR- has undergone nephrectomy and resection of metastatic lesions, 8) will be used as adjuvant treatment in combination with pembrolizumab or pembrolizumab berahyaluronidase alfa-pmph.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

Winrevair

Products Affected

- **WINREVAIR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of pulmonary arterial hypertension, substantiated by results from right heart catheterization (RHC), defined as a mean pulmonary arterial pressure (mPAP) of greater than 20 mmHg at rest, with a pulmonary capillary wedge pressure (PWP) of less than or equal to 15 mmHg, and a PVR greater than or equal to 3 Wood units -AND- WHO Group -AND- Concomitant use of at least one of the following (1-4): 1) generic endothelin-1 receptor antagonist, 2) phosphodiesterase type 5 inhibitor, 3) soluble guanylate cyclase stimulator, 4) generic prostacyclin agent -AND- Background therapy with PAH-specific therapies will be continued while being treated with Winrevair.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xalkori

Products Affected

- **XALKORI ORAL CAPSULE**
- **XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For metastatic non-small cell lung cancer (NSCLC), disease is ALK-positive as detected by an FDA-approved test or ROS1-positive as detected by an FDA-approved test. For relapsed or refractory anaplastic large cell lymphoma (ALCL), disease is ALK-positive. For unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT), disease is ALK-positive.
Age Restrictions	Deny if less than 1 year of age or greater than 21 years of age for ALCL.
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For crizotinib oral pellets and NSCLC, inability to swallow capsules is required. For crizotinib oral pellets and ALCL / IMT, inability to swallow oral capsules -OR- body surface area less than 1.34 m2 is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xcopri

Products Affected

- **XCOPRI**
- **XCOPRI MAINTENANCE PACK**
- **XCOPRI TITRATION PACK**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Therapeutic failure, intolerance or contraindication to 1 other anti-epileptic drug (e.g. carbamazepine, levetiracetam)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xdemvy

Products Affected

- **XDEMVY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis confirmed by identification of Demodex infection via microscopic examination of pulled eyelashes -OR- identification of collarettes via slit-lamp evaluation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xeljanz

Products Affected

- *tofacitinib oral tablet*
- *tofacitinib oral tablet extended release 24 hr*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For moderate to severe active rheumatoid arthritis, an inadequate response or intolerance to at least one non-biologic DMARD (e.g., leflunomide, methotrexate). For immediate release formulation for polyarticular juvenile idiopathic arthritis, inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) -OR- requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage. For ankylosing spondylitis, inadequate response or intolerance to at least one nonsteroidal anti-inflammatory drug (NSAID).
Age Restrictions	Deny if less than 18 years of age for rheumatoid arthritis, ulcerative colitis, ankylosing spondylitis. For extended release, deny if less than 18 years of age for psoriatic arthritis. For immediate release, deny if less than 2 years of age for polyarticular juvenile idiopathic arthritis and psoriatic arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced inadequate response or intolerance to at least 1 tumor necrosis factor (TNF) blocker. Doses greater than 10 mg per day (immediate release) and 11 mg per day (extended release) will not be approved for rheumatoid arthritis and psoriatic arthritis. Doses greater than 20mg per day (immediate release) and 22 mg per day (extended release) will not be approved for ulcerative colitis.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xeljanz Solution

Products Affected

- *tofacitinib oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. Documentation of polyarticular juvenile idiopathic arthritis -AND- Inadequate response or intolerance to at least one DMARD (e.g., methotrexate, leflunomide) or requires initial biologic therapy due to involvement of high-risk joints, high disease activity or at high risk of disabling joint damage.
Age Restrictions	Deny if less than 2 years of age for polyarticular juvenile idiopathic arthritis and psoriatic arthritis.
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	The member has experienced therapeutic failure or intolerance to at least 1 tumor necrosis factor (TNF) blocker.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xenazine

Products Affected

- *tetrabenazine oral tablet 12.5 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- attestation that the beneficiary is not actively suicidal
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	In patients with comorbid depression, attestation of adequate treatment for depression is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xermelo

Products Affected

- **XERMELO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of carcinoid syndrome diarrhea AND used in combination with a somatostatin analog AND trial and failure of somatostatin analog monotherapy
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	3 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in average number of daily bowel movements -AND- the member will continue to use in combination with a somatostatin analog.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xgeva

Products Affected

- **WYOST**
- **XGEVA**
- **XTRENBO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For hypercalcemia of malignancy, refractory to bisphosphonates. For giant cell tumor of bone, unresectable or surgical resection is likely to result in severe morbidity -AND- one of the following (1. or 2.)- 1.) the member is 18 years old or older -OR- 2.) the member is a skeletally mature adolescent (e.g. has at least one mature long bone)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Plan Year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xifaxan

Products Affected

- **XIFAXAN ORAL TABLET 550 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of 1 or 2. 1) Diagnosis of hepatic encephalopathy AND trial/failure, intolerance, or contraindication to lactulose. 2) Diagnosis of Irritable Bowel Syndrome with Diarrhea (IBS-D) AND trial/failure, intolerance to one of the following medications for IBS-D or documentation of contraindication to all: antidiarrheal (e.g., loperamide), antispasmodic (e.g., dicyclomine, hyoscyamine), tricyclic antidepressant (e.g., amitriptyline, nortriptyline).
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	Hepatic encephalopathy: 1 year. IBS-D: 14 days.
Other Criteria	No more than three courses of rifaximin for the treatment of IBS-D will be approved per lifetime.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xolair

Products Affected

- **XOLAIR**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of the following (1-2): 1) Chronic Spontaneous Urticaria, 2) trial/failure or intolerance of a second-generation non-sedating H1 antihistamine at the maximum recommended doses (e.g. cetirizine, fexofenadine, loratadine, desloratadine, levocetirizine). -OR- Documentation of the following (3-9): 3) moderate to severe persistent asthma, 4) a positive skin test or in vitro reactivity to a perennial aeroallergen, 5) Baseline IgE titer greater than or equal to 30 IU/mL, 6) documented pretreatment FEV1 less than 80% predicted in adults or less than 90% predicted in children and adolescents or FEV1 reversibility of at least 12% and 200mL after albuterol administration, 7) history of at least 2 asthma exacerbations requiring oral or injectable corticosteroid treatment in past 12mos or at least 1 asthma exacerbation requiring hospitalization in past 12mos, 8) inadequate symptom control despite regular treatment w/ medium- or high-dose inhaled corticosteroids (ICS) and at least 1 add'l asthma controller medication (e.g. long-acting beta2-agonist [LABA], leukotriene receptor antagonist [LTRA], theophylline) w/ or w/o OCS, unless intolerant or contraindicated to all, 9) will continue treatment with a medium- or high-dose ICS and at least 1 add'l asthma controller medication w/ or w/o OCS -OR- Documentation of the following (10-11): 10) chronic rhinosinusitis with nasal polyps (CRSwNP), 11) will use concomitantly with nasal corticosteroid maintenance treatment, -OR- Documentation of the following (12-17): 12) IgE mediated food allergy, 13) diagnosis confirmed by skin prick test or food-specific antibodies, 14) previous allergic reaction to food, 15) using for the reduction of allergic reactions (type 1), including anaphylaxis, 16) will be used in conjunction with food allergen avoidance, 17) member has a documented prescription for epinephrine.
Age Restrictions	
Prescriber Restrictions	

PA Criteria	Criteria Details
Coverage Duration	12 months
Other Criteria	For asthma reauthorization, attestation of one of the following is required (1-4): 1) decreased rescue medication or oral corticosteroid use, 2) decreased frequency of asthma exacerbations, 3) increased pulmonary function from baseline (e.g. FEV1), or 4) reduction in reported asthma related symptoms. For CSU reauthorization, improved CSU symptoms. For CRSwNP reauthorization, attestation of decrease in nasal polyp score or reduction in nasal congestion/obstruction severity score. For IgE-mediated food allergy reauthorization, member requires continuation of therapy and will continue food allergen avoidance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xolremdi

Products Affected

- **XOLREMDI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of WHIM syndrome (warts, hypogammaglobulinemia, infections, myelokathexis).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of reduction in incidence of infections is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xospata

Products Affected

- **XOSPATA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- member is FLT3 mutation-positive.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xpovio

Products Affected

- **XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80 MG/WEEK (80 MG X 1), 80MG TWICE WEEK (160 MG/WEEK)**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of use in combination with dexamethasone for relapse or refractory multiple myeloma with failure, intolerance or contraindication to 5 therapies (e.g. bortezomib, carfilzomib, lenalidomide, pomalidomide and daratumumab) -OR- Documentation of use in combination with both bortezomib and dexamethasone for multiple myeloma after receiving 1 prior multiple myeloma therapy -OR- Documentation of relapsing or refractory diffuse large B-cell lymphoma with failure, intolerance or contraindication to at least 2 lines of systemic therapy
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Xtandi

Products Affected

- **XTANDI ORAL CAPSULE**
- **XTANDI ORAL TABLET 40 MG, 80 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For castration-resistant prostate cancer and metastatic castration sensitive-prostate cancer, the member is using in combination with a GnRH analog or the member has had a bilateral orchiectomy. For non-metastatic castration-sensitive prostate cancer, the member has biochemical recurrence at high risk for metastasis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Xyrem

Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of narcolepsy -AND- Documentation of baseline data of excessive daytime sleepiness (EDS) via the Epworth Sleepiness Scale (ESS) or Maintenance of Wakefulness Test (MWT) - AND- Documentation of the following (1, 2, or 3): 1) Hypocretin-1 deficiency defined by (A or B), A) Cerebrospinal fluid hypocretin-1 less than 110 pg/mL. B) Cerebrospinal fluid hypocretin-1 less than 1/3 of the normal value based on laboratory reference range -OR- 2) Multiple sleep latency test (MSLT) documenting MSL less than 8 minutes and 2 sleep-onset rapid eye movement periods (SOREMP) -OR- 3) MSLT documenting MSL less than 8 minutes and 1 SOREMP and Polysomnography substantiating 1 SOREMP. If the member has a diagnosis of cataplexy provision of baseline number of cataplexy episodes is required.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	If 18 years or older and no diagnosis of cataplexy, trial and failure, intolerance, or contraindication to generic modafinil -AND- a generic CNS stimulant indicated for use in narcolepsy (e.g. methylphenidate, amphetamine salts) is required. For reauthorization, attestation supporting improvement in symptoms of narcolepsy and cataplexy (if applicable) is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Yonsa

Products Affected

- **YONSA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with methylprednisolone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Yorvipath

Products Affected

- **YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML, 294 MCG/0.98 ML, 420 MCG/1.4 ML**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hypoparathyroidism -AND- albumin-corrected serum calcium greater than or equal to 7.8 mg/dL -AND- trial and failure of an active form of vitamin D (e.g. calcitriol, alfacalcidol).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	For reauthorization, attestation of normalization or improvement in total serum calcium from baseline is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zavesca

Products Affected

- *miglustat*
- **YARGESA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of mild to moderate type 1 Gaucher disease confirmed by the following (A. or B.) A. (1, 2, 3, 4, or, 5): 1)Hepatomegaly. 2)Splenomegaly. 3)Bone disease (i.e. osteonecrosis, osteopenia, secondary pathologic fractures, bone infarct). 4)Bone marrow complications as defined by anemia with hemoglobin less than or equal to 11.5 g/dL for females or 12.5 g/dL for males or thrombocytopenia with platelet count less than or equal to 120,000/mm*3 -OR- 5)Symptomatic disease (e.g. bone pain, exertional limitation, cachexia). -OR- B. Attestation of deficiency in glucocerebrosidase activity in peripheral leukocytes or genetic testing confirms mutant alleles.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	Attestation enzyme replacement therapy (e.g. Cerezyme, Elelyso, or VPRIV) is not a therapeutic option
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zavzpret

Products Affected

- **ZAVZPRET**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of acute migraine headaches with or without aura -AND- Therapeutic failure, contraindication or intolerance to one generic triptan -AND- Inability to swallow capsules/tablets
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months initial authorization, 12 months reauthorization
Other Criteria	For reauthorization, attestation of reduction in migraine symptoms.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zejula

Products Affected

- **ZEJULA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis. For advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer, member is in complete or partial response to first-line platinum-based therapy -AND- disease is associated with homologous recombination deficiency-positive status, defined by at least one of the following (1 or 2): 1) a deleterious or suspected deleterious BRCA mutation or 2) genomic instability. For recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, all of the following (1-2): 1) disease harbors a deleterious or suspected deleterious germline BRCA mutation 2) member is in a complete or partial response to platinum-based chemotherapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zelboraf

Products Affected

- **ZELBORAF**

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Documentation of diagnosis. For unresectable or metastatic melanoma, member has a BRAF V600E mutation. For Erdheim-Chester Disease, member has a BRAF V600 mutation.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zelsuvmi

Products Affected

- **ZELSUVMI**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of molluscum contagiosum
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	For reauthorization, attestation that the member has previously experienced complete or partial clearance of molluscum contagiosum lesions with Zelsuvmi -AND- additional course of therapy is required for recurrence of molluscum contagiosum
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zenbexus

Products Affected

- **ZENBEXUS**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of multiple myeloma -AND- combination use with daratumumab/hyaluronidase-fihj and dexamethasone -AND- received at least 1 prior line of therapy that included a proteasome inhibitor and an immunomodulatory agent.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zeposia

Products Affected

- **ZEPOSIA**
- **ZEPOSIA STARTER KIT (28-DAY)**
- **ZEPOSIA STARTER PACK (7-DAY)**

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of Zeposia and other disease modifying agents such as interferons, Copaxone, Tysabri.
Required Medical Information	Documentation of diagnosis.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Ulcerative Colitis: Plan Year, Multiple Sclerosis: 24 months
Other Criteria	For moderate to severe active ulcerative colitis, patients must have therapeutic failure or intolerance to 2 of the following preferred products: a preferred adalimumab product, a preferred ustekinumab SC product, Skyrizi SC, Rinvoq, and Xeljanz/Xeljanz XR. Please Note: Preferred adalimumab products include Hadlima with NDC starting with 78206, Simlandi with NDC starting with 51759, and Yuflyma with NDC starting 72606. Preferred ustekinumab products include Steqema with NDC starting with 72606 and Yesintek with NDC starting with 83257.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zokinvy

Products Affected

- **ZOKINVY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- BSA of at least 0.39 m*2. For Hutchinson-Gilford Progeria Syndrome (HGPS), a mutation in the LMNA gene is required. For Processing-Deficient Progeroid Lamionpathy (PL), 1 of the following is required (1-3): 1) heterozygous LMNA mutation and progerin-like protein accumulation, 2) homozygous ZMPSTE24 mutations, or 3) compound heterozygous ZMPSTE24 mutations.
Age Restrictions	Deny if less than 12 months of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zolinza

Products Affected

- **ZOLINZA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of cutaneous manifestations in patients with cutaneous T-cell lymphoma (CTCL) who have progressive, persistent, or recurrent disease on or following 2 systemic therapies. Systemic therapies include bexarotene, interferon alpha, extracorporeal photochemotherapy, PUVA, single agent or combination chemotherapies (e.g. cyclophosphamide, vinblastine, romidepsin)
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zonisade

Products Affected

- **ZONISADE**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of partial-onset seizures -AND- Documentation of adjunctive therapy -AND- Inability to swallow capsules -AND- Therapeutic failure/intolerance to 2 or contraindication to all of the following (1-6): 1) generic carbamazepine suspension/chewable tablet/extended-release capsule, 2) generic gabapentin capsules/solution, 3) generic lacosamide solution, 4) generic levetiracetam solution, 5) generic oxcarbazepine suspension, 6) generic pregabalin capsules/solution.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Zoryve 0.15 Cream

Products Affected

- **ZORYVE TOPICAL CREAM 0.15 %**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of all of the following (1-3). 1) Documentation of mild to moderate atopic dermatitis -AND- 2) therapeutic failure, contraindication or intolerance to one generic formulary topical corticosteroid -OR- documentation of facial or anogenital involvement -AND- 3) therapeutic failure, contraindication or intolerance to topical generic tacrolimus or topical generic pimecrolimus.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

Ztalmy

Products Affected

- **ZTALMY**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of seizures associated with CDKL5 deficiency confirmed by genetic testing.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zurzuvae

Products Affected

- **ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis of postpartum depression -AND- attestation of symptom onset began during the third trimester of pregnancy or up to 4 weeks post-delivery -AND- less than or equal to 12 months postpartum.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 days
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zykadia

Products Affected

- **ZYKADIA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- ALK mutations, as detected by an FDA approved test.
Age Restrictions	Deny if less than 18 years of age
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Zytiga

Products Affected

- *abiraterone oral tablet 250 mg, 500 mg*
- **ABIRTEGA**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of diagnosis -AND- using in combination with prednisone.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

Index of Drugs

abiraterone oral tablet 250 mg, 500 mg 402
ABIRTEGA 402
acetaminophen-codeine oral solution 120-12 mg/5 ml 250
acetaminophen-codeine oral tablet 250
acitretin 1
ACTEMRA ACTPEN 332
ACTEMRA SUBCUTANEOUS 332
ACTIMMUNE 2
adapalene topical gel 0.3 % 335
adapalene topical swab 335
adapalene-benzoyl peroxide 335
ADEMPAS 259
AIMOVIG AUTOINJECTOR SUBCUTANEOUS AUTO-INJECTOR 140 MG/ML, 70 MG/ML 6
AKEEGA 7
ALCOHOL PADS 140
ALECENSA 8
alosetron oral tablet 0.5 mg, 1 mg 184
ALPRAZOLAM INTENSOL 250
alprazolam oral tablet 0.25 mg, 0.5 mg, 1 mg, 2 mg 250
alprazolam oral tablet extended release 24 hr 0.5 mg, 1 mg, 2 mg, 3 mg 250
ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG 10
ALUNBRIG ORAL TABLETS,DOSE PACK 10
ambrisentan 259
amitriptyline 118
amitriptyline-chlordiazepoxide 118
ANZUPGO 13
ARALAST NP INTRAVENOUS RECON SOLN 1,000 MG 9
ARIKAYCE 14
aripiprazole oral solution 16
aripiprazole oral tablet,disintegrating 16
armodafinil 217
asenapine maleate 289
ATTRUBY 15
AUGTYRO ORAL CAPSULE 160 MG, 40 MG 18
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG 19
AUSTEDO XR 19
AUSTEDO XR TITRATION KT(WK1-4) ORAL TABLET, EXT REL 24HR DOSE PACK 12-18-24-30 MG 19
AUVELITY ORAL TABLET, IR AND ER, BIPHASIC 20
AVEED 325
AVMAPKI-FAKZYNJA 21
AVONEX INTRAMUSCULAR PEN INJECTOR KIT 142
AVONEX INTRAMUSCULAR SYRINGE KIT 142
AYVAKIT 22
BAFIERTAM 23
BALVERSA 24
BENLYSTA SUBCUTANEOUS 26
benztropine oral 118
BESREMI 28
BESREMI PEN 28
BETASERON SUBCUTANEOUS KIT 142
bexarotene oral 316
bexarotene topical 317
bosentan oral tablet 259
BOSULIF ORAL CAPSULE 100 MG, 50 MG 29
BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG 29
bosutinib oral tablet 400 mg 30
BRAFTOVI 31
BRINSUPRI 33
BRUKINSA ORAL TABLET 34
buprenorphine 250
butalbital-acetaminop-caf-cod oral capsule 50-300-40-30 mg, 50-325-40-30 mg 250
CABOMETYX 36
CALQUENCE (ACALABRUTINIB MAL) 37
CAPLYTA 38
CAPRELSA ORAL TABLET 100 MG, 300 MG 39
carglumic acid 40
CAYSTON 41
CERDELGA 42
CINRYZE 45
clindamycin-tretinoin 335

clobazam oral suspension.....225
clobazam oral tablet.....225
clomipramine.....118
*clonazepam oral tablet 0.5 mg, 1 mg, 2 mg*250
clonazepam oral tablet,disintegrating 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg.....250
clonidine hcl oral tablet extended release 12 hr.....3
clorazepate dipotassium oral tablet 15 mg, 3.75 mg, 7.5 mg..... 250
COBENFY.....47
COBENFY STARTER PACK..... 47
COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)..... 48
COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML..... 105
COPIKTRA.....49
COSENTYX (2 SYRINGES).....51
COSENTYX PEN (2 PENS).....51
COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML..... 51
COSENTYX UNOREADY PEN.....51
COTELLIC.....53
cyproheptadine oral syrup.....118
cyproheptadine oral tablet.....118
CYSTADROPS.....54
CYSTARAN.....54
dalfampridine..... 11
DANZITEN.....55
dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg.....301
DAURISMO ORAL TABLET 100 MG, 25 MG..... 57
deferasirox oral tablet, dispersible.....59
DEPO-TESTOSTERONE INTRAMUSCULAR OIL 100 MG/ML. 325
dextromethorphan-quinidine..... 214
DIACOMIT ORAL CAPSULE 250 MG, 500 MG..... 61
DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG.....61
DIAZEPAM INTENSOL.....250
diazepam oral solution 5 mg/5 ml (1 mg/ml).....250
diazepam oral tablet..... 250
diclofenac epolamine.....97
diclofenac sodium topical gel 3 %.....299
dihydroergotamine nasal.....62
dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg.....323
doxepin oral capsule.....118
doxepin oral concentrate.....118
doxepin oral tablet.....118
doxepin topical..... 63
DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG, 30 MG, 40 MG, 60 MG..... 64
droxidopa oral capsule 100 mg, 200 mg, 300 mg..... 211
DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML.....65
DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 200 MG/1.14 ML, 300 MG/2 ML..... 65
DUVYZAT..... 67
eltrombopag olamine oral powder in packet 12.5 mg, 25 mg.....330
eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg.....330
EMGALITY PEN.....70
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3).....70
ENBREL MINI.....82
ENBREL SUBCUTANEOUS SOLUTION.....82
ENBREL SUBCUTANEOUS SYRINGE 25 MG/0.5 ML (0.5), 50 MG/ML (1 ML)..82
ENBREL SURECLICK.....82
ENOBY.....255
ENSACOVE.....72
ENTRESTO SPRINKLE.....73
EOHILIA.....74
EPIDIOLEX.....76
ergotamine-caffeine.....78
ERIVEDGE.....79

ERLEADA ORAL TABLET 240 MG, 60 MG 80
erlotinib 68
escitalopram oxalate oral capsule 81
estazolam 250
eszopiclone 250
everolimus (antineoplastic) oral tablet 4
everolimus (antineoplastic) oral tablet for suspension 2 mg, 3 mg, 5 mg 4
EXXUA ORAL TABLET EXTENDED RELEASE 24 HR 84
EXXUA ORAL TABLET, EXT REL 24HR DOSE PACK 84
FABHALTA 85
FANAPT 87
FANAPT TITRATION PACK A 87
FANAPT TITRATION PACK B 87
FANAPT TITRATION PACK C 87
FASENRA PEN 88
FASENRA SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 30 MG/ML 88
fentanyl transdermal patch 72 hour 100 mcg/hr, 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr 250
FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26) 91
FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR 120 MG, 20 MG, 40 MG, 80 MG 91
FILSUVEZ 92
fingolimod 104
FINTEPLA 93
FIRMAGON KIT W DILUENT SYRINGE 96
flurazepam 250
FOTIVDA 100
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG 101
GAMMAGARD LIQUID 127
GAMMAGARD LIQUID ERC 127
GAMMAGARD S-D (IGA < 1 MCG/ML) 127
GATTEX 30-VIAL 102
GAUZE PAD TOPICAL BANDAGE 2 X 2 " 140
GAVRETO 103
gefitinib 146
GENOTROPIN 112
GENOTROPIN MINIQUICK 112
GILOTRIF 68
GLASSIA 9
glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml 105
GLATOPA SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML 105
glycerol phenylbutyrate 266
GOMEKLI ORAL CAPSULE 1 MG, 2 MG 110
GOMEKLI ORAL TABLET FOR SUSPENSION 110
GRASTEK 111
guanfacine oral tablet extended release 24 hr 3
HADLIMA 245
HADLIMA PUSHTOUCH 245
HADLIMA(CF) 245
HADLIMA(CF) PUSHTOUCH 245
HAEGARDA 114
HEPCLUDEX 116
HERNEXEOS 117
HUMATROPE INJECTION CARTRIDGE 112
hydrocodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg 250
hydromorphone oral liquid 250
hydromorphone oral tablet 2 mg, 4 mg, 8 mg 250
hydroxyzine hcl oral solution 10 mg/5 ml 118
hydroxyzine hcl oral tablet 118
HYRNUO 121
IBRANCE ORAL CAPSULE 125 MG 122
IBRANCE ORAL TABLET 122
IBTROZI 124
icatibant 94
ICLUSIG 125
IDHIFA 126
imatinib oral tablet 100 mg, 400 mg 106
IMBRUVICA ORAL CAPSULE 140 MG, 70 MG 129
IMBRUVICA ORAL SUSPENSION 129

IMBRUVICA ORAL TABLET 129
imipramine hcl 118
imipramine pamoate 118
IMKELDI 130
INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE 132
INCRELEX 133
INGREZZA INITIATION PK(TARDIV) 135
INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG 135
INGREZZA SPRINKLE 135
INLURIYO 136
INLYTA 137
INQOVI 138
INREBIC 139
ISTURISA ORAL TABLET 1 MG, 5 MG 302
ITOVEBI ORAL TABLET 3 MG, 9 MG 147
itraconazole oral capsule 148
ivabradine oral tablet 5 mg, 7.5 mg 50
ivermectin oral 149
IWILFIN 150
JAKAFI 151
JASCAYD 152
JAVYGTOR 168
JAYPIRCA ORAL TABLET 100 MG, 50 MG 154
JIDEYTRO ORAL TABLET 100 MG, 25 MG 155
JOENJA ORAL TABLET 70 MG 156
JUBBONTI 255
KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG 158
KALYDECO ORAL TABLET 158
KERENDIA 159
KESIMPTA PEN 160
KINERET 161
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3) 162
KLISYRI (250 MG) 163
KOMZIFTI 164
KOSELUGO ORAL CAPSULE 10 MG, 25 MG 166
KOSELUGO ORAL CAPSULE, SPRINKLE 5 MG, 7.5 MG 166
KRAZATI 167
KYGEVVI 169
lapatinib 341
LAZCLUZE ORAL TABLET 240 MG, 80 MG 171
lenalidomide 273
LENVIMA 172
LEUKINE INJECTION RECON SOLN 174
lidocaine hcl mucous membrane solution 4 % (40 mg/ml) 334
lidocaine topical adhesive patch,medicated 5 % 176
lidocaine topical ointment 334
lidocaine-prilocaine topical cream 334
LIFYORLI ORAL CAPSULE 125 MG/DAY(100 MG X1-25MG X1), 125 MG/DAY(100 MG X1-25MG X3), 125 MG/DAY(100 MG X1-25MG X9), 150 MG/DAY(100 MG X1-25MG X2), 150 MG/DAY(100 MG X1-25MG X3), 150 MG/DAY(100 MG X1-25MG X9) 177
liraglutide 109
LITFULO 178
LIVTENCITY 179
LOARGYS 180
LOKELMA 181
lomustine 108
LONSURF 182
LORAZEPAM INTENSOL 250
lorazepam oral tablet 0.5 mg, 1 mg, 2 mg 250
LORBRENA ORAL TABLET 100 MG, 25 MG 183
LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG 185
lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg 170
LYBALVI 186
LYNPARZA 187
LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5) 190

macitentan 259
MAVYRET ORAL PELLETS IN PACKET 191
MAVYRET ORAL TABLET 191
megestrol oral suspension 400 mg/10 ml (40 mg/ml), 625 mg/5 ml (125 mg/ml) 192
megestrol oral tablet 192
MEKINIST ORAL RECON SOLN 193
MEKINIST ORAL TABLET 0.5 MG, 2 MG 193
MEKTOVI 195
memantine-donepezil 201
mesalamine oral capsule, extended release 500 mg 240
metaxalone oral tablet 400 mg, 800 mg 118
methadone oral solution 10 mg/5 ml, 5 mg/5 ml 250
methadone oral tablet 10 mg, 5 mg 250
methylphenidate transdermal 58
metyrosine 196
mifepristone oral tablet 300 mg 165
miglustat 388
milnacipran oral tablet 290
modafinil 257
MODEYSO 197
morphine concentrate oral solution 250
morphine oral solution 10 mg/5 ml (2 mg/ml), 20 mg/5 ml (4 mg/ml) 250
morphine oral tablet 250
morphine oral tablet extended release 100 mg, 15 mg, 200 mg, 30 mg, 60 mg 250
MOUNJARO 109
MULPLETA 199
NAYZILAM 202
NEMLUVIO 203
NERLYNX 205
NEXLETOL 207
NEXLIZET 207
nilotinib hcl 319
NINLARO 209
nintedanib 144
nitisinone 210
NORDITROPIN FLEXPRO SUBCUTANEOUS PEN INJECTOR 10 MG/1.5 ML (6.7 MG/ML), 15 MG/1.5 ML (10 MG/ML), 5 MG/1.5 ML (3.3 MG/ML) 112
NOURIANZ 212
NUBEQA 213
NUCYNTA 250
NUEDEXTA 214
NUPLAZID 215
NURTEC ODT 216
octreotide acetate injection solution 219
ODACTRA 220
ODOMZO 221
OGSIVEO ORAL TABLET 100 MG, 150 MG 222
OJEMDA ORAL SUSPENSION FOR RECONSTITUTION 223
OJEMDA ORAL TABLET 400 MG/WEEK (100 MG X 4), 500 MG/WEEK (100 MG X 5), 600 MG/WEEK (100 MG X 6) 223
OJJAARA 224
OMNITROPE 112
ONGENTYS 226
ONUREG 227
OPIPZA 228
OPSUMIT 259
OPSYNVI 259
ORENCIA CLICKJECT 229
ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML 229
ORGOVYX 231
ORKAMBI ORAL GRANULES IN PACKET 232
ORKAMBI ORAL TABLET 232
ORSERDU ORAL TABLET 345 MG, 86 MG 233
OTEZLA 234
OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)- 20 MG (51), 10 MG (4)-20 MG (4)-30 MG (47) 234
OTEZLA XR 234
OTEZLA XR INITIATION 234

oxazepam 250
oxycodone oral capsule 250
oxycodone oral concentrate 250
oxycodone oral tablet 10 mg, 15 mg, 20 mg, 30 mg, 5 mg 250
oxycodone-acetaminophen oral tablet 10-325 mg, 2.5-325 mg, 5-325 mg, 7.5-325 mg 250
OZEMPIC ORAL 236
OZEMPIC SUBCUTANEOUS 109
PALYNZIQ SUBCUTANEOUS SYRINGE 10 MG/0.5 ML, 2.5 MG/0.5 ML, 20 MG/ML 237
PANRETIN 238
paroxetine hcl 12
pazopanib oral tablet 200 mg 362
PEGASYS 141
PEMAZYRE 239
perphenazine-amitriptyline 118
PHEBURANE 241
phenobarbital 120
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2) 242
pirfenidone oral capsule 144
pirfenidone oral tablet 144
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML 142
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML 142
pomalidomide 243
posaconazole oral tablet,delayed release (dr/ec) 244
pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg 189
pregabalin oral solution 189
PRENATAL VITAMIN PLUS LOW IRON 249
PREVYMIS ORAL PELLETS IN PACKET 253
PRIVIGEN 127
PROCYSBI ORAL GRANULES DEL RELEASE IN PACKET 254
PROLASTIN-C INTRAVENOUS SOLUTION 9
PROLIA 255
promethazine oral 118
PULMOZYME 261
pyrimethamine 56
QINLOCK 262
quinine sulfate 263
QULIPTA 264
RADICAVA ORS STARTER KIT SUSP 265
RASUVO (PF) 198
RELISTOR ORAL 267
RELISTOR SUBCUTANEOUS SOLUTION 268
RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML 268
REPATHA SURECLICK 269
REPATHA SYRINGE 269
RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG 271
REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG 274
REXULTI ORAL TABLET 275
REZDIFFRA 276
REZLIDHIA 278
RINVOQ LQ 281
RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG 279
RIVFLOZA SUBCUTANEOUS SOLUTION 282
RIVFLOZA SUBCUTANEOUS SYRINGE 128 MG/0.8 ML, 160 MG/ML 282
ROMVIMZA 283
ROZLYTREK ORAL CAPSULE 100 MG, 200 MG 284
ROZLYTREK ORAL PELLETS IN PACKET 284
RUBRACA 285
rufinamide 25
RYDAPT 286
SAJAZIR 94
sapropterin 168

SAVELLA ORAL TABLETS,DOSE PACK 290
SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG 291
SECUADO 292
SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG 112
SIGNIFOR 302
sildenafil (pulm.hypertension) oral tablet 259
SILIQ 293
SIMLANDI(CF) AUTOINJECTOR 245
SIMLANDI(CF) SUBCUTANEOUS SYRINGE KIT 20 MG/0.2 ML, 40 MG/0.4 ML 245
SIRTURO 295
SKYCLARYS 296
SKYRIZI SUBCUTANEOUS PEN INJECTOR 297
SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML, 55 MG/0.37 ML 297
SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML) 297
sodium oxybate 384
sodium phenylbutyrate 35
sofosbuvir-velpatasvir 75
SOHONOS ORAL CAPSULE 1 MG, 1.5 MG, 10 MG, 5 MG 298
SOMAVERT 300
sorafenib 206
STELARA SUBCUTANEOUS SOLUTION 345
STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML 345
STEQEYMA SUBCUTANEOUS SOLUTION 247
STEQEYMA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML 247
STIVARGA 303
sunitinib malate 306
SUNOSI 304
SYMPAZAN 307
SYMPROIC 308
SYNAREL 309
TABRECTA 310
tadalafil (pulm. hypertension) 259
tadalafil oral tablet 2.5 mg, 5 mg 44
TAFINLAR ORAL CAPSULE 311
TAFINLAR ORAL TABLET FOR SUSPENSION 311
TAGRISSO 313
TALZENNA 315
TASCENSO ODT 318
tasimelteon 320
TAVALISSE 321
tazarotene topical cream 322
tazarotene topical gel 0.1 % 322
temazepam 250
TEPMETKO 324
teriflunomide 17
teriparatide subcutaneous pen injector 20 mcg/dose (560mcg/2.24ml) 98
testosterone cypionate 325
testosterone enanthate 325
testosterone transdermal gel in metered-dose pump 20.25 mg/1.25 gram (1.62 %) 325
testosterone transdermal solution in metered pump w/app 325
tetrabenazine oral tablet 12.5 mg, 25 mg 374
TEZSPIRE SUBCUTANEOUS PEN INJECTOR 327
THALOMID ORAL CAPSULE 100 MG, 50 MG 329
TIBSOVO 331
TOBI PODHALER 43
tobramycin in 0.225 % nacl 43
tobramycin inhalation 43
tofacitinib oral solution 373
tofacitinib oral tablet 371
tofacitinib oral tablet extended release 24 hr 371
tolvaptan 288
tolvaptan (polycys kidney dis) oral tablet 157
tolvaptan (polycys kidney dis) oral tablets, sequential 157
topiramate oral solution 77
tramadol oral tablet 100 mg, 25 mg, 50 mg, 75 mg 250
tramadol-acetaminophen 250
tretinoin 335

triazolam 250
TRIKAFTA ORAL GRANULES IN PACKET, SEQUENTIAL 336
TRIKAFTA ORAL TABLETS, SEQUENTIAL 336
trimipramine 118
TRINTELLIX 337
TRULICITY 109
TRUQAP 338
TUKYSA ORAL TABLET 150 MG, 50 MG 339
TURALIO 340
TYMLOS 342
UBRELVY ORAL TABLET 100 MG, 50 MG 343
UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG 344
UPTRAVI ORAL TABLETS,DOSE PACK 344
ustekinumab subcutaneous solution 345
ustekinumab subcutaneous syringe 45 mg/0.5 ml, 90 mg/ml 345
VALCHLOR 347
VALTOCO 348
VANFLYTA 349
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG 350
VENCLEXTA STARTING PACK 350
VERQUVO 351
VERZENIO 352
VIBERZI 353
vigabatrin 287
VIGADRONE 287
VIJOICE ORAL GRANULES IN PACKET 355
VIJOICE ORAL TABLET 125 MG, 250 MG/DAY (200 MG X1-50 MG X1), 50 MG 355
vilazodone 354
VITRAKVI ORAL CAPSULE 100 MG, 25 MG 356
VITRAKVI ORAL SOLUTION 356
VIVJOA 357
VONJO 358
VORANIGO ORAL TABLET 10 MG, 40 MG 359
voriconazole intravenous recon soln 360
VOSEVI 361
VOWST 90
VRAYLAR 363
VUMERITY 364
WELIREG 365
WINREVAIR 367
WYOST 376
XALKORI ORAL CAPSULE 368
XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG 368
XCOPRI 369
XCOPRI MAINTENANCE PACK 369
XCOPRI TITRATION PACK 369
XDEMVY 370
XERMELO 375
XGEVA 376
XIFAXAN ORAL TABLET 550 MG 377
XOLAIR 378
XOLREMDI 380
XOSPATA 381
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80 MG/WEEK (80 MG X 1), 80MG TWICE WEEK (160 MG/WEEK) 382
XTANDI ORAL CAPSULE 383
XTANDI ORAL TABLET 40 MG, 80 MG 383
XTRENBO 376
YARGESA 388
YESINTEK SUBCUTANEOUS SOLUTION 247
YESINTEK SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML 247
YONSA 386
YORVIPATH SUBCUTANEOUS PEN INJECTOR 168 MCG/0.56 ML, 294 MCG/0.98 ML, 420 MCG/1.4 ML 387
YUFLYMA(CF) AI CROHN'S-UC-HS 245

YUFLYMA(CF) AUTOINJECTOR.....245
YUFLYMA(CF) SUBCUTANEOUS SYRINGE KIT 40 MG/0.4 ML..... 245
YULITHIRA..... 4
zaleplon oral capsule 10 mg, 5 mg.....250
ZAVZPRET..... 389
ZEJULA..... 390
ZELBORAF..... 391
ZELSUVMI.....392
ZELVYSIA.....168
ZEMAIRA INTRAVENOUS RECON SOLN 1,000 MG..... 9
ZENBEXUS.....393
ZEPOSIA.....394
ZEPOSIA STARTER KIT (28-DAY).....394
ZEPOSIA STARTER PACK (7-DAY)...394
ZILBRYSQ SUBCUTANEOUS SYRINGE 16.6 MG/0.416 ML, 23 MG/0.574 ML, 32.4 MG/0.81 ML..... 200
zileuton..... 175
ZOKINVY.....395
ZOLINZA.....396
zolpidem oral tablet.....250
*zolpidem oral tablet,ext release multiphase*250
zolpidem sublingual.....250
ZONISADE..... 397
ZORYVE TOPICAL CREAM 0.15 %...398
ZTALMY.....399
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG..... 400
ZYKADIA..... 401

Section: Step Therapy

Brand ADHD

Products Affected

- *lisdexamfetamine 10 mg capsule*
- *lisdexamfetamine 20 mg capsule*
- *lisdexamfetamine 30 mg capsule*
- *lisdexamfetamine 40 mg capsule*
- *lisdexamfetamine 50 mg capsule*
- *lisdexamfetamine 60 mg capsule*
- *lisdexamfetamine 70 mg capsule*

Details

Criteria	Require a 1 month trial of 2 of the following generic ADHD medications (Step 1 drug) when being utilized for the same medically accepted indication: methylphenidate (i.e. generics of Concerta, Metadate CD, Methylin, Relexxii, Ritalin/Ritalin LA), atomoxetine, dextroamphetamine/amphetamine (i.e. generics of Adderall/Adderal XR), or dexmethylphenidate in the last 180 days.

Entresto Tablet

Products Affected

- **ENTRESTO 24 MG-26 MG TABLET**
- **ENTRESTO 49 MG-51 MG TABLET**
- **ENTRESTO 97 MG-103 MG TABLET**

Details

Criteria	Require a 1 month trial of generic sacubitril/valsartan tablets (Step 1 drug) in the last 90 days

Herpetic Keratitis

Products Affected

- **ZIRGAN 0.15 % EYE GEL**

Details

Criteria	Require a 1 month trial of generic trifluridine eye drops (Step 1 drug) in the last 90 days when utilized for the same medically accepted indication and age

Pulmonary Antiinflammatory

Products Affected

- *fluticasone propionate 100 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 110 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 220 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 250 mcg/actuation blister powder for inhalation*
- *fluticasone propionate 44 mcg/actuation hfa aerosol inhaler*
- *fluticasone propionate 50 mcg/actuation blister powder for inhalation*

Details

Criteria	Require a 1 month trial of Qvar and Asmanex/Asmanex HFA (Step 1 drugs) in the last 180 days

Rho Kinase Inhibitors

Products Affected

- **RHOPRESSA 0.02 % EYE DROPS DROPS**
- **ROCKLATAN 0.02 %-0.005 % EYE**

Details

Criteria	Require a 1 month trial of one preferred glaucoma drug (Step 1 drug) in the last 120 days

Riomet

Products Affected

- *metformin 500 mg/5 ml oral solution*

Details

Criteria	Require a 1 month trial of generic metformin IR tablets in the last 90 days -OR- documentation supporting the inability to swallow or difficulty swallowing tablets containing metformin

Rytary

Products Affected

- **RYTARY 23.75 MG-95 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 36.25 MG-145 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 48.75 MG-195 MG CAPSULE,EXTENDED RELEASE**
- **RYTARY 61.25 MG-245 MG CAPSULE,EXTENDED RELEASE**

Details

Criteria	Require a trial of generic carbidopa/levodopa product (Step 1 drug) in the last 90 days

Topical Antifungals

Products Affected

- *naftifine 1 % topical cream*
- *naftifine 2 % topical cream*
- *naftifine 2 % topical gel*

Details

Criteria	Require a 1 month trial of generic econazole cream and one of the following: generic ketoconazole cream or ketoconazole shampoo (Step 1 drugs),when being utilized for the same medically accepted indication, in the last 180 days

Topical Psoriasis

Products Affected

- *calcipotriene-betamethasone 0.005 %-0.064 % topical ointment*
- *calcipotriene-betamethasone 0.005 %-0.064 % topical suspension*
- *calcitriol 3 mcg/gram topical ointment*

Details

Criteria	Require a 1 month trial of generic calcipotriene cream, ointment or solution (Step 1 drug) in the last 90 days when utilized for the same medically accepted indication and age

Topical Rosacea

Products Affected

- *azelaic acid 15 % topical gel*
- **FINACEA 15 % TOPICAL FOAM**

Details

Criteria	Require a 1 month trial of generic metronidazole cream, lotion, or gel (Step 1 drug) in the last 90 days.

Index of Drugs

azelaic acid 15 % topical gel.....10
calcipotriene-betamethasone 0.005 %-0.064 % topical ointment.....9
calcipotriene-betamethasone 0.005 %-0.064 % topical suspension.....9
calcitriol 3 mcg/gram topical ointment.....9
ENTRESTO 24 MG-26 MG TABLET.....2
ENTRESTO 49 MG-51 MG TABLET.....2
ENTRESTO 97 MG-103 MG TABLET.....2
FINACEA 15 % TOPICAL FOAM.....10
fluticasone propionate 100 mcg/actuation blister powder for inhalation.....4
fluticasone propionate 110 mcg/actuation hfa aerosol inhaler.....4
fluticasone propionate 220 mcg/actuation hfa aerosol inhaler.....4
fluticasone propionate 250 mcg/actuation blister powder for inhalation.....4
fluticasone propionate 44 mcg/actuation hfa aerosol inhaler.....4
fluticasone propionate 50 mcg/actuation blister powder for inhalation.....4
lisdexamfetamine 10 mg capsule.....1
lisdexamfetamine 20 mg capsule.....1
lisdexamfetamine 30 mg capsule.....1
lisdexamfetamine 40 mg capsule.....1
lisdexamfetamine 50 mg capsule.....1
lisdexamfetamine 60 mg capsule.....1
lisdexamfetamine 70 mg capsule.....1
metformin 500 mg/5 ml oral solution.....6
naftifine 1 % topical cream.....8
naftifine 2 % topical cream.....8
naftifine 2 % topical gel.....8
RHOPRESSA 0.02 % EYE DROPS.....5
ROCKLATAN 0.02 %-0.005 % EYE DROPS.....5
RYTARY 23.75 MG-95 MG CAPSULE,EXTENDED RELEASE.....7
RYTARY 36.25 MG-145 MG CAPSULE,EXTENDED RELEASE.....7
RYTARY 48.75 MG-195 MG CAPSULE,EXTENDED RELEASE.....7
RYTARY 61.25 MG-245 MG CAPSULE,EXTENDED RELEASE.....7
ZIRGAN 0.15 % EYE GEL.....3